Sensing, Feeling, and Regulating: Investigating the Association of Focal Brain Damage with
Voluntary Respiratory and Motor Control

Henrik Bischoff[1], Christopher Kovach[2,3], Sukbhinder Kumar[2], Joel Bruss[4], Daniel Tranel[5], Sahib S Khalsa[6]

1 Department of Psychology, University of Stockholm, Stockholm, Sweden; Department of Psychology,
Carl-von-Ossietzky University Oldenburg, Oldenburg, Germany

2 Department of Neurosurgery, University of Iowa, Iowa City, USA.

3 Department of Neurosurgery, University of Nebraska Medical Center, Omaha, NE, USA

4 Departments of Pediatrics, Neurology, and Psychiatry, University of Iowa, Iowa City, USA.

5 Departments of Neurology and Psychological and Brain Sciences, University of Iowa, Iowa City, USA.

6 Laureate Institute for Brain Research, Tulsa, OK, USA; Oxley College of Health Sciences, University of
Tulsa, Tulsa, OK, USA.

Correspondence:

Sahib S. Khalsa MD, PhD
Laureate Institute for Brain Research
6655 S Yale Ave, Tulsa, OK 74136-3326
Phone: 918 502 5743
Fax: 918 502 5135
Email: skhalsa@laureateinstitute.org
26

33 Abstract: 200 words

34 Word count: About 6000 words

Abstract

Breathing is a complex, vital function that can be modulated to influence physical and mental well-being. However, the role of cortical and subcortical brain regions in voluntary control of human respiration is underexplored. Here we investigated the influence of damage to human frontal, temporal, or limbic regions on the sensation and regulation of breathing patterns. Participants performed a respiratory regulation task across regular and irregular frequencies ranging from 6 to 60 breaths per minute (bpm), with a counterbalanced hand motor control task. Interoceptive and affective states induced by each condition were assessed via questionnaire and autonomic signals were indexed via skin conductance. Participants with focal lesions to the bilateral frontal lobe, right insula/basal ganglia, and left medial temporal lobe demonstrated significantly lower performance than individually matched healthy comparisons during the breathing and motor tasks. They also reported significantly higher anxiety during the 60-bpm regular and irregular breathing trials than healthy participants. Correlations between task performance, anxiety, and difficulty varied across experimental conditions and groups. This study demonstrates that damage to frontal, temporal, or limbic regions is associated with abnormal voluntary respiratory and motor regulation and tachypnea-related anxiety, highlighting the role of the forebrain in affective and motor responses during breathing.

Keywords: respiration, interoception, brain damage, emotion, affect, amygdala, motor control

59 **Highlights**

60 • Impaired human respiratory regulation is associated with cortical/subcortical brain lesions
61 • Frontolimbic/temporal regions contribute to rhythmic breathing and hand motor control
62 • Frontolimbic/temporal damage is associated with anxiety during tachypnea/irregular breathing
63 • The human forebrain is vital for affective and interoceptive experiences during breathing

Introduction

Breathing is a vital yet complex function performed continuously by vertebrates from birth until death. Crucial for cellular metabolism and gas exchange and often automatically performed, human breathing can be consciously controlled (Shweta, Pramod & Chen, 2021; Lumb, 2016), impacting physical (Hernandez, Manning & Zhang, 2019; Kim, Shin & Cho, 2022; Russo, Santarelli & O'Rourke, 2017) and mental health outcomes (Banushi et al., 2023; Balban et al., 2023) as well as central (Karjalainen, Kujala & Parviainen, 2023) and peripheral autonomic arousal (Laborde et al., 2022). Voluntary breathing refers to the conscious control of breathing, such as when we hold our breath, take a deep breath, or exhale forcefully. Voluntary breathing has been shown to modulate neural (McKay et al., 2003; Herrero et al., 2018), cognitive (Zaccaro et al., 2018) and emotional states (Jerath & Beveridge, 2020), demonstrating the relevance of this function to numerous health-related processes. While automatic breathing is presumptively generated by brainstem-related mechanisms operating outside conscious awareness, voluntary breathing is predicated on the individual's conscious awareness and control of respiration and is a core component of yoga and other meditative practices (Anālayo, 2019; Peng et al., 2004). Some forms of voluntary respiration, such as slow breathing, are widely considered an important means of quickly reducing anxiety or stress (Balban et al., 2023; Birdee et al., 2023), whereas other forms such as rapid breathing (so called "breath of fire") are sometimes considered to quickly increase energy levels (Peng et al., 2004). Despite increasing studies on the effects of voluntary breathing on the body and mind (Herrero et al., 2018; Zaccaro et al., 2018), the neural mechanisms underlying voluntary breathing control remain unclear.

Automatic breathing, controlled by the brainstem, maintains oxygen and carbon dioxide levels reflexively and independent of conscious control mechanisms. It is initiated by neurons in the formatio reticularis (FR), with the rhythm regulated by the dorsal respiratory group (DRG), the ventral respiratory group (VRG), and the pre-Bötzinger complex (preBötC) (Anderson & Ramirez, 2017; Shweta, Pramod & Chen, 2021; Ikeda et al., 2017; Yang & Feldman, 2018) (Figure 1b). The pneumotaxic and apneustic centers function to prevent lung over-distension. Automatic breathing can be interrupted or overridden by voluntary control mechanisms, and while there are separate anatomic pathways for automatic and voluntary breathing, these pathways demonstrate significant functional overlap (Prasad, Pal & Chen, 2021; Betka et al., 2022). Automatic breathing, controlled by FR's respiratory networks, chemoreceptors, and respiratory mechanoreceptors, can be influenced by descending inputs from emotional and cognitive neural networks. Voluntary breathing thus appears to shift neural control towards supratentorial brain centers (Del Negro, Funk & Feldman, 2018; Homma & Masaoka, 2008; McKay et al., 2003; Herrero et al., 2018).

Functional neuroimaging and intracranial recording studies have identified frontal, parietal, and temporal lobe involvement in voluntary breathing phases, suggesting a potential regulatory influence of widespread cortical and subcortical regions (McKay et al., 2003; Herrero et al., 2018) (Figure 1a). The prefrontal cortex (PFC), encompassing the dorsolateral, ventrolateral, and dorsomedial sections, connects to key subcortical respiratory regions (Jodo et al., 1998; Gabbott et al., 2005; Terreberry & Neafsey, 1987; Dutschmann & Dick, 2012), leaving it well positioned for a role in voluntary respiration in addition to its well-recognized roles in emotion regulation (Graham & Milad, 2013; Dixon et al, 2017; Silvers et al., 2015), motor control (Fine & Hayden, 2021; Watanabe et al., 2002), regulation of executive functions (Yuan & Raz, 2014; Friedman & Robbins, 2022), response inhibition (Aron, Robbins & Poldrack, 2014; Anderson, Bunce & Barbas, 2016), working memory (Curtis & Esposito, 2003; Lara & Wallis, 2015) and decision making (Bechara et al., 1998; Gläscher et al., 2012). The anterior cingulate cortex (ACC), another frontal region contributing to emotion (Etkin, Egner & Kalisch, 2011; Etkin, Büchel & Gross, 2015; García-Cabezas & Barbas, 2017), motivation (Holroyd & Yeung, 2012; Rolls, 2023), visceromotor regulation (Burns & Wyss, 1985) and cognitive control (Botvinick et al., 2001;

García et al., 2022), has also been implicated in voluntary control of breathing. For example, using intracranial recordings in humans, a recent study linked dorsal ACC (dACC) activity with voluntary breath holding and increased blood carbon dioxide levels (Holton et al., 2021). The insular cortex (IC), a key region for processing and integrating interoceptive signals (Craig, 2002; Cameron, 2009; Khalsa et al., 2009a; Hassanpour et al., 2018) and emotion (Damasio et al., 2000), is engaged by voluntary breathing (Pattinson et al., 2009), and right IC damage can lead to reduced dyspnea perception (Schön et al., 2008) and sleep-disordered breathing (Harper et al., 2021). The subcortical basal ganglia (BG) are also found to play a role in voluntary respiratory regulation (McKay et al., 2008) in addition to roles in initiating desired skeletomotor movement and inhibiting competing movements (Young, Reddy & Sonne, 2021; Martinu et al., 2012), as are the primary motor/sensory, premotor and supplementary motor areas. Temporal lobe damage can impact breathing regulation (Homma & Masaoka, 2008; Nobis et al., 2019), and patients with temporal lobe epilepsy may experience oxygen desaturation and central apnea during seizure-induced interference with amygdala activity (Dlouhy et al., 2015; Nobis et al., 2019; Rhone et al., 2020). In sudden unexplained death from epilepsy (SUDEP), studies have pointed toward a deficient arousal sensor to increased levels of CO2, which commonly occur during states of apnea (Buchanan, 2019), and a recent study even identified a subregion within the amygdala responsible for suppressing breathing and air hunger following stimulation (Harmata et al., 2023). Studies showing a role for the amygdala in interoceptive anxiety and cardiorespiratory interoception (Feinstein et al., 2013, Khalsa et al., 2016) have led to the suggestion that a lack of awareness of amygdala-mediated apnea may be a key mechanism provoking interoceptive arousal, fear, and anxiety (Feinstein, Gould & Khalsa, 2022). In summary, the intracranial and functional neuroimaging evidence to date suggests that the regulation of voluntary respiration is associated with neural processes spanning the PFC, ACC, insula, BG, amygdala, and adjacent temporal structures. However, there is a paucity of studies evaluating the impact of damage to these regions on voluntary breathing or associated affective, interoceptive, and autonomic processes.

The current study investigated the association of damage to human frontal, temporal, or limbic (Heimer & Van Hoesen, 2005) brain regions with the ability to sense and voluntarily regulate breathing patterns using a multilevel repeated measures design incorporating interoceptive, affective, and autonomic signals. We compared individuals with focal acquired brain lesions to healthy individuals, aiming to identify associations between cortical and subcortical damage and voluntary respiratory regulation, as well as to measures of physiological (i.e., autonomic) and subjective (i.e., affective and interoceptive) experience. To rule out general motor skill impairment, we included motor control tasks. We selected a number of symmetric respiratory rates, reflecting an interest in evaluating respiratory performance across the full physiological range of rates used in relaxation-, yoga-, and exercise-related behaviors, and to examine the impact of both slow and fast breathing on affective, interoceptive, and autonomic indices. We also selected an irregular breathing task to evaluate autonomic dynamics via respiratory sinus arrhythmia-induced heart rate variability and its impact on affective and interoceptive states. We hypothesized that individuals with damage to the aforementioned regions would show a reduced ability to regulate their respiratory performance during voluntary breathing, as indicated by decreased ability to follow a visually presented breathing trace in comparison to age-, sex-, and body mass index-matched healthy individuals without brain injury. Performance was evaluated with the peak cross-correlation between the instructed trace and chest wall excursion, measured with a chest belt. We hypothesized that any difference of performance would be specific to breathing, i.e., there would be no significant group differences in motor regulation ability on a hand movement task, due to the lesion locations in the selected group. With respect to affective and interoceptive indices, we hypothesized that the lesion group would report more anxiety across both slow and rapid breathing conditions versus resting rates, reflecting a U-shaped pattern of anxiety and task difficulty during extreme perturbations of breathing rate. Although slow breathing

159 techniques are often recommended for reducing anxiety and stress, we hypothesized that increased
160 anxiety would occur during slow breathing in the lesion group due to a difficulty in voluntary breathing
161 regulation. Finally, we explored multilevel associations between respiratory regulation parameters and
162 autonomic (i.e., skin conductance responses), affective (i.e., anxiety ratings), and metacognitive
163 perception (i.e., task difficulty rating) aspects.

Methods

Participants

Healthy participants were recruited through public advertisements at the University of Iowa. Lesion
participants with acquired focal brain injury were recruited from the Iowa Neurological Patient Registry
at the University of Iowa Hospitals and Clinics. Inclusion criteria for the lesion participants included age
18 to 85 years, presence of a focal brain lesion at least 3 months after onset, no history of learning
disabilities, psychiatric disorders, substance abuse, personality disorders, developmental epilepsy, or
other unrelated neurological conditions. The lesions among the participants with focal brain injury
were caused by ischemic or haemorrhagic strokes, meningioma resection, or surgical intervention for
epilepsy. Exclusion criteria for the lesion participants included cardiovascular disease, chronic
obstructive pulmonary disease, active substance use disorder, peripheral/central neurological
disorders other than the defined brain lesion. Healthy participants were required to have no brain
lesions and to meet the remainder of the inclusion/exclusion criteria.

Experimental Design

After obtaining written informed consent participants completed a screening interview to verify
eligibility. Inclusion and exclusion criteria for the lesion group were examined using medical history
data from the Iowa Neurological Patient Registry. When applicable, new demographic information was
also gathered.

Prior to completing the regulation tasks participants with brain lesions were asked to perform a
neuropsychological motor coordination task, called the grooved pegboard test (Trites, 1989). The task
involves fitting small irregularly shaped rods into openings on a pegboard using only one hand,
providing a measure of the motor skills of the fingers, fingertips, hands, and forearms. This test of
motor skill helped to identify fine hand-motor dysfunctions associated with acquired brain lesions,
which could potentially confound the measurement of dial rotation in the hand motor regulation task.
Since this measure was collected only in the lesion participants, their performance was compared
against an age-matched normative sample (Lafayette Instrument Company, 2002). We employed
individual scoring criteria, using Trites' (1989) age-based norms and considered scores greater than 2
standard deviations (SD) above the age-based mean to reflect motor impairment (performance was
measured via completion time, with higher durations indicating worse performance).

The main experimental condition included two regulation paradigms: respiratory and hand motor,
each with two distinct blocks of tests (regular and irregular). For each paradigm, participants were
seated and connected to measurement devices (see Measurements and Recording Devices section
below). Each regular respiratory condition consisted of four 2-minute symmetric breathing trials at
different instructed frequencies (6, 10, 16, and 60 nasal breaths per minute, bpm) and one irregular
trial consisting of a pseudo-random frequency and amplitude modulation of the nasal inspiration and
expiration phases. During the respiratory regulation trials, participants followed a visual cue with their
breath, inhaling nasally with their mouth closed when a red dot moved up and exhaling nasally when
it moved down a sinusoidal curve (Figure 2a). Prior to the respiratory task participants underwent an

amplitude calibration test where they were instructed to demonstrate a maximum amplitude (i.e., full vital capacity) inhalation followed by exhalation. They were then told that the top and bottom of the screen represented their full vital capacity inhalation/exhalation, shown that the amplitude of the instructed sinusoid did not reach this boundary, and were told to ensure that the amplitudes of their inhalation/exhalations should be not exceed their comfort level. The hand motor task, which served as a control, involved the same visual cue, with participants instead making rotational dial movements with their dominant hand (or unaffected hemisphere/hand pair) to track the movement of the dot along the line. It included five trials with hand dial rotational movement cues that were matched to the respiratory task. Participants were also told not to twist their hands during the dial task to the point of discomfort. The presentation order of the experimental trials within each task was randomized, and the sequence of regulation tasks (breathing or motor first) was counterbalanced. Practice rounds were performed at the start of each block to familiarize participants with the experimental procedures.

After each trial, participants were asked to rate the physical sensations and emotional states they had experienced via a questionnaire involving multiple questions using a Likert response scale ranging from 1 (very little) to 5 (very much). This also included their perception of the difficulty and enjoyment of the task. Post-trial ratings of task difficulty were also evaluated to ascertain any differences in perceived difficulty of the respiratory and motor regulation tasks between the groups. They were also asked to rate other feelings and sensations experienced during each trial, including anxiety, boredom, focus, calm, excitement, and others. This study focused on evaluating task-related affect, specifically, anxiety and difficulty ratings.

Measurements and Recording Devices

Demographic data (age, gender, handedness, ethnicity, and height) were gathered through a questionnaire. Body weight was physically measured to calculate body mass index (BMI).

Physiological data were gathered via a MP100 acquisition unit (Biopac Inc., Santa Barbara, CA). Breathing activity was monitored using an elastic thoracic breathing belt and amplifier (RSP100C) at the mid-sternal level. Cardiac activity was measured using two electrodes paired with an amplifier (ECG100C) designed for electrocardiogram (ECG) monitoring and recorded at 200 Hz with a high pass filter to stabilize the ECG baseline. The positive electrode was positioned on the left lower anterior abdomen and the negative electrode on the right anterior chest at the mid-clavicular line (lead II configuration). Galvanic skin response electrodes, placed on the thenar and hypothenar eminences of the hand contralateral to the dial hand, were collected using a single-channel amplifier (EDA100C) at 200 Hz.

A custom-built dial was employed to measure hand movement regulation, consisting of a rotating potentiometer with a continuous rating scale ranging from 0.000 to 5.000 Volts at a sampling rate of 200 Hz, as utilized in our previous studies (Khalsa et al., 2009b; 2020). Participants were asked to rotate the dial using their dominant hand following a visual cue, while breathing normally during each trial.

Statistical Analysis

Descriptive statistics for demographic data were compared for the lesion and matched-healthy comparison group. T-tests were used for metric variables (age, BMI), with normality checked using the Shapiro-Wilk test and histogram distributions.

To evaluate respiratory and motor regulation performance, maximum cross-correlation values were calculated between the instructed and measured signals (respiratory trace for breathing trials and dial trace for motor trials) using MatLab R2021a software (Mathworks Inc., Natick, MA) and transformed using Fisher's Z-transformation to address skewed sampling distributions of correlation coefficients

(Silver & Dunlap, 1987). Cross-correlations were calculated using the built-in xcorr function. Linear
mixed effects models were used to compare performance across the respiration and motor regulation
tasks via separate models for the regular and irregular tasks using the lme4 package in R. The lmerTest
package (Bates et al., 2015) was used to calculate statistical significance and to perform Bonferroni-
corrected post-hoc tests for group comparisons, employing the emmeans R-package (Lenth, 2018) for
post-hoc analysis. The analysis excluded participants missing respiratory or motor regulation data for
more than two trials per task. For the multilevel correlation analysis, Pearson correlation coefficients
were calculated in R.

For skin conductance response (SCR) analysis, we utilized the open-source software Ledalab (version
3.x; Benedek & Kaernbach, 2010) for Matlab. Data were first preprocessed using a 1st order low-pass
Butterworth filter with a cutoff frequency of 5 Hz. Adaptive smoothing was performed using Gaussian
windows with a width of 16 samples. CDA.ISCR (Cumulative Driver Amplitude Integrated with Skin
Conductance Response) was the dependent variable calculated for in-depth analysis of the SCR signals.
CDA.ISCR represents the area covered by the temporal integration of the phasic driver of the SCR with
respect to the duration of the SCR. This is achieved by multiplying the SCR by the size of the response
window. CDA.ISCR is expressed in microsiemens-seconds ($\mu S*s$) and accounts for both the intensity
and duration of the SCR over time. This allows a more detailed study of the temporal evolution and
intensity of responses to different stimuli.

Ethics and Data Source

266 This study was approved by the University of Iowa Institutional Review Board. All volunteers provided
267 written informed consent prior to study participation and received compensation for their
268 involvement.

Results

Participant Demographics

A total of 76 participants were consented, consisting of 56 healthy individuals (mean (*M)* = 43.2,
standard deviation (*SD)* = 12.5 years, range from 20 to 63 years) and 20 participants with acquired focal
brain injury (*M* = 53.7, *SD* = 10.05 years, range from 36 to 84 years). Due to substantial variations in
age and sex ratio between groups, the sample of 20 lesion participants (14 female and 6 male) was
individually matched to an equal-sized sample of healthy participants (*n* = 20; 14 female and 6 male
participants) based on age, sex, and BMI (Table 1). The mean age of lesion (*M* = 53.8; *SD* = 7.05 years)
and healthy (*M* = 53.6; *SD* = 10.6 years) participants did not significantly differ (*t* (33) = -0.07, *p* = 0.94),
nor did the mean BMI of lesion (*M* = 27.6; *SD* = 5.0) and healthy (*M* = 30.7; *SD* = 5.9) participants
significantly differ (*t* (35) = 1.81, *p* = 0.079). Shapiro-Wilk testing showed no evidence of non-normality
of matching variables within experimental groups (Age: Lesion, *W* = 0.97, *p* = 0.73; Healthy, *W* = 0.92,
p = 0.098; BMI: Lesion, *W* = 0.91, *p* = 0.071; Healthy, *W* = 0.97, *p* = 0.73).

Brain Lesion Characteristics

284 The lesion locations for the 20 study participants with focal brain injury included a clustering of regions
285 in the left and right frontal lobe, right hemispheric lesions in the insular cortex and basal ganglia, and
286 left hemispheric lesions in the temporal lobe (Figure 1c).

Grooved Pegboard Task

288 For the nondominant hand, the lesion group mean completion time was M = 87.9 seconds (SD = 27.5);
289 for the dominant hand, the mean score was M = 82.8 seconds (SD = 29.8). According to the scoring
290 criteria, 29.4% (5 out of 17 participants) were classified as impaired for the non-dominant hand; for
291 the dominant hand (i.e., the hand used to rotate the dial), 11.1% (2 of 18 participants) were classified
292 as impaired.

Respiratory and Motor Regulation Task Performance

294 Due to the putative widespread involvement of frontal, parietal, and temporal lobe structures in
295 voluntary breathing, we conducted omnibus tests comparing performance of the entire lesion sample
296 against the healthy sample for both the respiratory and motor regulation tasks. Due to the substantial
297 dissimilarity of the irregular trial stimuli, we divided the evaluation of performance across the regular
298 and irregular tasks.

Regular Task

On the regular task there were significant main effects of group (F(1, 39) = 26.29, p < .001) and task
(F(1, 270) = 33.51, p < 0.001), and a significant interaction of group and task (F(1, 270) = 6.58, p =
0.011). Post-hoc tests showed that the healthy comparison group significantly outperformed the lesion
group across both breathing (p = 0.046, mean difference (MDiff) = 0.24, 95%-CI [0.00, 0.48]) and motor
control tasks (p < 0.001, Mdiff = 0.51, 95%-CI [0.26, 0.76]). Both groups performed significantly better
on the hand motor control task as compared to the breathing task (healthy, p < 0.001, Mdiff = -0.44,
95%-CI [-0.63, -0.24]; lesion, p = 0.155, Mdiff = -0.17, 95%-CI [-0.37, -0.03]). See Figure 2c and
Supplemental Figure 1 for performance at the individual task frequencies. In summary, while the lesion
group performed significantly worse than the healthy comparison group on both regular breathing and
regular motor tasks, the disparity in performance between the tasks was smaller in the lesion group as
evidenced by the significant group by task interaction.

Irregular Task

On the irregular task there was a significant main effect of group (F(1,38) = 14.50, p < 0.001) and task
(F(1,37) = 17.04, p < 0.001). However, no significant interaction was observed between group and task
(F(1,37) = 0.0031, p = 0.956). Post-hoc comparisons revealed that the healthy participants significantly
outperformed those with lesions for both the breathing (p = 0.015, Mdiff = 0.16, 95%-CI [0.02, 0.3])
and motor tasks (p = 0.017, Mdiff = 0.16, 95%-CI [0.02, 0.3]). However, contrary to the pattern
observed on the regular task, for the irregular task both groups performed significantly better on the
breathing component of the task as opposed to the hand motor control (healthy, p = 0.037, Mdiff =
0.12, 95%-CI [0.00, 0.23]; lesion, p = 0.035, Mdiff = 0.12, 95%-CI [0.00, 0.24]). See figure 2d and
Supplemental Figure 1. In summary, participants with lesions demonstrated lower performance on
both irregular tasks compared to healthy participants, with performance in the breathing condition
being superior for both groups.

Affective Ratings

For anxiety experienced during the regular breathing condition significant group differences were
observed (F(1, 37) = 7.81, p = 0.008), with the lesion participants reporting higher anxiety (M = 2.02
out of 5, SD = 1.39) than the healthy comparisons (M = 1.39 out of 5, SD = 0.70) across all breathing
tasks. There was also a significant effect of breathing condition (F(4, 147) = 14.48, p < 0.001), and an
interaction between group and breathing condition (F(4, 147) = 3.16, p = 0.016). Post-hoc tests
revealed significant within-group differences in anxiety only at the 60 bpm condition relative to
baseline for participants with lesions (p < 0.001, Mdiff = -1.91, 95%-CI[-2.77, -1.04]). This was also

reflected by a significant difference in anxiety between the lesion versus healthy comparison group at
60 bpm (p = 0.001, Mdiff = 1.42, 95%-CI [0.32, 2.52]) (Figure 3a).

For anxiety ratings during the regular motor task conditions, there was a significant main effect for
task condition ($F_{(4, 134)}$ = 12.50, p < 0.001) but no significant main effect for the experimental groups
($F_{(1, 35)}$ = 1.44, p = 0.238). Post-hoc tests showed that both groups reported significant anxiety
increases for the fastest motor regulation trial (60 rpm) compared to baseline. Lesion participants also
reported significant anxiety increases in slow and normal regulation trials (6 & 10 rpm), in the absence
of significant group differences (Figure 3b).

For anxiety experienced during the irregular breathing condition, there was an interaction between
group and breathing condition ($F_{(1, 38)}$ = 6.65, p = 0.013). Post-hoc tests showed significantly increased
anxiety scores for the lesion versus healthy comparison group (p = 0.002, Mdiff = 1.08, 95%-CI [0.32,
1.84]) (Figure 3c).

For anxiety experienced during the irregular motor task, there was a significant main effect of task ($F_{(1,}$
$_{39)}$ = 33.13, p < 0.001) but no significant main effect for the experimental groups ($F_{(1, 39)}$ = 1.85, p =
0.182) or interactions (Figure 3d).

Difficulty Ratings

For the difficulty rating during the regular tasks, there was a main effect for the groups ($F_{(1, 38)}$ = 8.20,
p = 0.007) such that participants with lesions perceived the regular tasks as more challenging overall.
Post hoc analyses clarified that these participants rated the regular breathing tasks significantly more
difficult than healthy controls (p = 0.054, Mdiff = 0.73, 95% CI[0.008, 1.48]). However, there were no
significant group differences in perceived difficulty between the respiratory and motor tasks for
participants with lesions (p > 0.999, Mdiff = -0.08, 95%-CI[-0.58, 0.41]) or for the healthy comparisons
(p > 0.999, Mdiff = - 0.15, 95%-CI[-0.62, 0.31]).

For the irregular task types, no statistically significant differences in perceived difficulty were found
between groups, either within one group (between breathing and motor) or between groups for both
task types.

SCR

For SCR during the regular breathing tasks, there was a main effect for the task ($F_{(4, 136)}$ = 5.70, p <
0.001). Post hoc analyses clarified that healthy participants in the 60 bpm showed significantly higher
physiological arousal, than in the 6 bpm (p < 0.001, Mdiff = -6044.1, 95% CI[-10372, -1717]) and 10
bpm condition (p = 0.024, MDiff = -4586.2, 95% CI[-8892, -280]). For lesion participants, there were no
significant differences in SCR between the different respiratory tasks.

For SCR during the regular motor tasks, there was also a main effect for the task ($F_{(1, 122)}$ = 2.95, p =
0.023). Post hoc analyses clarified that healthy participants in the 60 bpm showed significantly higher
physiological arousal, than in the 6 bpm (p = 0.035, MDiff = -5500, 95% CI[-10834, -168]). For lesion
participants, there were no significant differences in SCR between the different motor tasks.

For the irregular breathing and motor tasks, there were no significant SCR effects for both
experimental groups compared with baseline.

Multilevel Correlation Analysis

370 This exploratory analysis examined the correlations between respiratory regulatory performance
371 (behavioral/physiological measure), SCR (physiological measure), anxiety rating (subjective-affective
372 measure), and rating of the respective perceived task difficulty (a metacognitive interoceptive

373   measure). Based on our interest in the neural regulation of respiration, associations were evaluated
374   for each group separately.

Correlations with Breathing

For healthy participants, significant correlations were observed across different conditions of the
regular breathing task: At 10 bpm, a positive correlation was found between anxiety and perceived
difficulty ($r(20) = 0.73$, $p = 0.002$) as well as between anxiety and SCR ($r(20) = 0.65$, $p = 0.008$). At 16
bpm, anxiety was negatively correlated with performance accuracy ($r(20) = -0.60$, $p = 0.028$), and
positively correlated with perceived difficulty ($r(20) = 0.51$, $p = 0.049$). In addition, a significant positive
correlation was also found between perceived difficulty and SCR ($r(20) = 0.58$, $p = 0.027$). There were
no significant correlations found between variables in the slow breathing (6 bpm), fast breathing (60
bpm) or the irregular breathing task. To summarize the correlational breathing trial results for healthy
participants, increased perceived difficulty was linked both to higher SCR values and heightened
anxiety during normal breathing (10 and 16 bpm). Interestingly, no significant correlations between
these variables were identified in the slow, fast, and irregular breathing tasks.

For lesion participants, significant correlations were observed across different conditions of the regular
breathing task: At 16 bpm, anxiety was negatively correlated with performance accuracy ($r(20) = -0.53$,
$p = 0.016$). Additionally, a significant positive correlation was found between perceived difficulty and
skin conductance ($r(20) = 0.60$, $p = 0.040$). At 60 bpm, a significant negative correlation was found
between performance accuracy and SCR ($r(20) = -0.59$, $p = 0.040$). Another significant positive
correlation was found between anxiety and perceived difficulty ($r(20) = 0.61$, $p = 0.021$). To summarize
the correlational breathing trial results for lesion participants in the regular breathing tasks, at specific
bpm rates, performance accuracy was negatively correlated with anxiety, with no significant
correlations observed at 6 bpm, 10 bpm, or the irregular breathing task. Task difficulty was consistently
associated with heightened skin conductance, particularly at 16 and 60 bpm.

Correlations with Hand Movement

For healthy participants, significant correlations were observed across different experimental
conditions of the motor dial task: At 6 rpm, anxiety was positively correlated with perceived difficulty
($r(20) = 0.75$, $p = 0.009$). No significant correlations were identified for the experimental trials at the
10, 16 and 60 rpm frequencies, nor were any found for the irregular motor task.

For lesion participants, the only significant correlation were observed at 60 rpm, between anxiety and
perceived difficulty ($r(20) = 0.73$, $p = 0.007$). No significant correlations were identified for the
experimental conditions for the experimental trials at the 6, 10 and 16 rpm frequencies, nor were any
found for the irregular motor task.

Exploratory Analysis

407   Given previous studies demonstrating breathing frequency-specific changes in heart rate variability
408   (HRV), we conducted an exploratory analysis of HRV across all tasks. This analysis revealed no group
409   differences across the different instructed respiratory frequencies but found that the lesion group
410   (relative to healthy comparisons) exhibited slightly higher heart rates as well as lower low-frequency
411   and higher high-frequency power of the HRV during the hand movement task (see Supplement).

Discussion

The current study investigated the association of damage to cortical and subcortical brain areas with
human respiratory and motor regulation, characterizing the interoceptive and affective responses
occurring during with these tasks. As hypothesized, participants with frontal, temporal, and limbic
brain lesions exhibited a significantly reduced ability to regulate their breathing across regular and
irregular tasks, in addition to reductions in hand motor control. While both groups performed better
on motor control tasks compared to breathing tasks during regular instruction conditions, this
discrepancy was less pronounced among the lesion group. Moreover, the lesion participants reported
heightened anxiety levels, particularly during performance of rapid symmetric or irregular breathing
frequencies, as well as greater task-related difficulty with the voluntary top-down modulation of their
breathing.

A fundamental, yet under-researched, aspect of human brain-body interactions is the connection
between the voluntary control of breathing and cortical and subcortical regions of the central nervous
system. Here, we examined an omnibus scenario by testing a group of lesion patients with damage to
a heterogeneous set of cortical and subcortical areas and comparing them to healthy participants. Our
hypothesis that participants with damage to the frontal, temporal or limbic regions would show a
reduced ability to control their breathing performance during a voluntary breathing task was partially
supported. Indeed, the participant group with lesions was found to have, on average, a lower
respiratory regulatory capacity than the healthy participant group across both regular and irregular
instruction conditions. However, against our expectations, this reduction in performance ability was
not restricted solely to breathing, suggesting that damage to the observed regions was also associated
with other forms of rhythmic motor control. This inference is supported by the fact that the lesion
group did not show substantial impairment in other forms of non-rhythmic hand movement, as
measured by the grooved pegboard test. One implication for this finding is that the neural circuits for
production of complex skeletomotor movements are recruited, and may even be necessary, for the
optimization of voluntary breathing control patterns, such as deep breathing and rapid breathing, that
are practiced during yoga, meditation, and stress management routines. While there is evidence from
structural and functional brain imaging approaches supporting the engagement of temporal and
frontal brain regions by yoga (Gothe et al., 2019), this is a sparsely studied topic and more data are
needed.

Although both groups performed better on motor control tasks as compared to breathing tasks during
regular instruction conditions, this discrepancy was less pronounced among the lesion group. The exact
mechanisms underlying this phenomenon warrant further investigation, but the finding may suggest
a specialized impairment of respiratory (versus motor) control following injury to the involved frontal,
limbic, and temporal brain regions. At the same time, the non-significant interaction between group
and task in the irregular instruction condition suggests a lower impact on breathing control
mechanisms when respiratory frequency and amplitude is randomly fluctuating. Respiratory rhythms
vary considerably depending on the internal and external environmental context, with symmetric
changes more common during homeostatic perturbations in the sympathetic (e.g., hyperventilation
during fight or flight response) and parasympathetic (e.g., slow breathing during relaxation) domains.
Accordingly, it seems plausible that the lesion group's difficulty in symmetric respiratory regulation
could manifest via tachypnea-associated anxiety during the 60-bpm condition; a possibility that is
further supported by their greater respiratory-specific difficulty ratings.

The lack of respiratory/hand-motor discrepancy for irregular breathing conditions raises several
considerations worth exploring. First, respiratory rhythms are increasingly acknowledged to influence
the balance of excitatory and inhibitory activity within the brain, with a particular influence of shifts in
excitability that are inducible by respiratory phase (Kluger & Gross, 2021; Kluger et al., 2023). This
suggests that aperiodic changes in respiratory rhythm, such as those produced during the irregular
breathing condition, could alter the brain's excitatory/inhibitory coupling in a manner different from
the regular breathing condition. Why these "respiration-modulated brain oscillations" (RMBOs) would
lead to a relative reduction in breathing performance in the regular vs. irregular breathing condition is

uncertain, but future studies might evaluate the impact of excitatory vs. inhibitory input from cortical
and subcortical areas to respiratory brainstem nuclei. Such an approach would require an animal
model of respiratory control, although neurosurgical patients could be another means of gaining some
insight into the role of specific subregions (e.g., medial temporal regions in epilepsy patients awaiting
a corrective temporal resection procedure). Second, respiratory rhythms are well established to
influence cognition, affect, and perception through influences on olfactory, insular, and
somatosensory pathways (reviewed in Allen, Varga & Heck, 2023) that are closely anatomically
connected to the frontal, limbic, and temporal regions damaged in the present lesion sample. It seems
plausible that the volatility inherent to the irregular respiration condition could have altered the
excitability of cortical circuits in a manner different from the regular respiration conditions, but the
precise impact of this volatility (e.g., selective influence on interoceptive prediction error processing,
or global influence on exteroceptive or affective representations; Brændholt et al., 2023) is unknown
and only the subject of speculation. Third, the present task relied primarily on the voluntary top-down
control of motor circuitry. Adapting this approach to incorporate spontaneous breathing perturbations
(e.g., via transient inspiratory restrictions (Herzog et al., 2021; Burt et al., 2023)) would allow for the
discernment of bottom-up processes contributing to the interoceptive and affective differences
reported during specific forms of breathing. Adapting such a protocol to the concurrent measurement
of neural oscillations would facilitate the identification interactions between top-down and bottom-
up processing streams, which could be used to develop fine-grained explanatory models of respiratory
sensation and regulation.

The affective responses observed during different task conditions provide intriguing insights into the
interplay between brain lesions, task performance, and affective responses. The fact that the lesion
participants reported significantly higher anxiety levels during specific respiratory conditions, notably
during fast breathing, highlights a potential heightened respiratory-affective sensitivity or alternatively
a decreased capacity to emotionally regulate tachynpeic states following brain injury. Although there
is no pre-lesion data from which to make firm claims of causality, the fact that the anxiety increase was
specific to one breathing frequency band increases the likelihood that reduced neural resources post-
injury might render these individuals more susceptible to interpreting tachypnea as an anxiety-
inducing stimulus. Moreover, the lack of anxious responses to slow breathing suggests that individuals
with brain lesions in frontal, limbic, and temporal areas may be able to utilize slow breathing
techniques as taught for reducing anxiety and stress without concern for a paradoxical effect.

In general, the correlations between regular task performance, anxiety, perceived difficulty, and
physiological measures (such as SCR) showed similar patterns for both groups. For example, in both
groups breathing performance accuracy was associated with lower difficulty ratings, greater difficulty
was associated with greater SCRs, and greater difficulty was associated with greater anxiety. During
the hand motor task there were fewer significant correlations for both groups. Collectively, these data
suggest that increased anxiety is detrimentally associated with breathing regulation ability irrespective
of brain injury status. Future studies could examine whether these associations are altered in clinically
anxious patients. The correlations between affective states and task performance observed during the
hand movement task also illustrate the challenging nature of the task instruction. In healthy
participants, anxiety was strongly correlated with perceived difficulty at the slowest rotation (6 rpm),
while in lesioned participants, a negative correlation between the two was observed at the highest
speed (60 rpm). Such rate-related variations could imply different coping or regulatory mechanisms
were activated depending on the neural integrity and the nature of the task. The lack of significant
correlational patterns for either group on the irregular breathing condition is unclear.

This study has several limitations and future directions. These include a small sample size, which was
limited by the rarity of the lesion sample, and the heterogeneity of lesion location, which was
determined by random sampling from available participants visiting the laboratory during the data
collection period. As study measurements were taken from patients during the chronic injury phase, it
is possible that there were undetected impacts of injury on respiration or motor ability during the

acute injury phase that had been compensated for by neural reorganization by the time of the study.
Respiration was only measured using a thoracic chest belt, which can yield inaccurate measurements,
thus future studies would benefit from more sophisticated measurement approaches capable of
assessing variables such as tidal/vital capacity and nasal flow rates. The groups also differed with
respect to educational level; although there are no clear suggestions this difference may have
influenced the current results future studies might consider controlling for this variable. Future
research directions include mechanistically investigating the influence of activity within specific brain
regions on respiratory and motor regulation with larger, more homogeneous sample groups; including
more cases of bilateral lesions; and possibly employing animal models and human intracranial
recording studies to further understand the neural basis of respiratory and motor control.
531

Conclusion

533
534 This study demonstrates that damage to frontal, temporal, or limbic brain regions is associated with
535 abnormal voluntary respiratory and motor regulation and tachypnea-related anxiety, highlighting the
536 role of the forebrain in affective and motor responses during breathing.

Author Contributions

SSK, CK, and DT: conceived and designed the experiments; SSK, CK, HB: carried out the experiments and/or processed the data. SSK, SK, CK, DT and HB: contributed to the analysis of the results. JB and HB: contributed the illustrations. Figure 1A and B was created with Biorender by HB and Figure 1C was created by JB, with input from SSK. All other figures were created by HB with input from SSK. SSK and HB: wrote the paper, with input from all authors.

Funding

This project was supported by the National Center for Complementary and Integrative Health Grant/Award Number F31AT003061 and the Kiwanis International Neuroscience Research Foundation.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

Allen, M., Varga, S., & Heck, D. H. (2023). Respiratory rhythms of the predictive mind. *Psychological review, 130*(4), 1066–1080. https://doi.org/10.1037/rev0000391

Anālayo, B. (2019). Meditation on the Breath: Mindfulness and Focused Attention. *Mindfulness, 10*(8), 1684–1691. https://doi.org/10.1007/s12671-019-01169-9

Anderson, M. C., Bunce, J. G., & Barbas, H. (2016). Prefrontal-hippocampal pathways underlying inhibitory control over memory. *Neurobiology of learning and memory, 134 Pt A*(Pt A), 145–161. https://doi.org/10.1016/j.nlm.2015.11.008

Anderson, T. M., & Ramirez, J. M. (2017). Respiratory rhythm generation: triple oscillator hypothesis. *F1000Research*, *6*, 139. https://doi.org/10.12688/f1000research.10193.1.

Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: one decade on. *Trends in cognitive sciences, 18*(4), 177–185. https://doi.org/10.1016/j.tics.2013.12.003

Balban, M. Y., Neri, E., Kogon, M. M., Weed, L., Nouriani, B., Jo, B., Holl, G., Zeitzer, J. M., Spiegel, D., & Huberman, A. D. (2023). Brief structured respiration practices enhance mood and reduce physiological arousal. *Cell reports. Medicine, 4*(1), 100895. https://doi.org/10.1016/j.xcrm.2022.100895

Banushi, B., Brendle, M., Ragnhildstveit, A., Murphy, T., Moore, C. E., Egberts, J., & Robison, R. (2023). Breathwork Interventions for Adults with Clinically Diagnosed Anxiety Disorders: A Scoping Review. *Brain Sciences, 13*(2), 256. https://doi.org/10.3390/brainsci13020256

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i0.

Bechara, A., Damasio, H., Tranel, D., & Anderson, S. W. (1998). Dissociation Of working memory from decision making within the human prefrontal cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 18*(1), 428–437. https://doi.org/10.1523/JNEUROSCI.18-01-00428.1998

Benedek, M., & Kaernbach, C. (2010). A continuous measure of phasic electrodermal activity. *Journal of Neuroscience Methods, 190*(1), 80–91. https://doi.org/10.1016/j.jneumeth.2010.04.028

Betka, S., Adler, D., Similowski, T., & Blanke, O. (2022). Breathing control, brain, and bodily self-consciousness: Toward immersive digiceuticals to alleviate respiratory suffering. *Biological Psychology, 171*, 108329. https://doi.org/10.1016/j.biopsycho.2022.108329

Birdee, G., Nelson, K., Wallston, K., Nian, H., Diedrich, A., Paranjape, S., Abraham, R., & Gamboa, A. (2023). Slow breathing for reducing stress: The effect of extending exhale. *Complementary therapies in medicine, 73*, 102937. https://doi.org/10.1016/j.ctim.2023.102937

Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological review*, *108*(3), 624–652. https://doi.org/10.1037/0033-295x.108.3.624

Brændholt, M., Kluger, D. S., Varga, S., Heck, D. H., Gross, J., & Allen, M. G. (2023). Breathing in waves: Understanding respiratory-brain coupling as a gradient of predictive oscillations. *Neuroscience and biobehavioral reviews, 152*, 105262. https://doi.org/10.1016/j.neubiorev.2023.105262

Buchanan, G.F. (2019). Impaired CO_2-Induced Arousal in SIDS and SUDEP. *Trends Neurosci*. 42(4):242-250. https://doi.org/10.1016/j.tins.2019.02.002.

Burns, S. M., & Wyss, J. M. (1985). The involvement of the anterior cingulate cortex in blood pressure control. *Brain research, 340*(1), 71–77. https://doi.org/10.1016/0006-8993(85)90774-7

Burt, J. S., Davenport, M. P., Welch, J. F., & Davenport, P. W. (2023). fNIRS analysis of rostral prefrontal cortex activity and perception of inspiratory loads. *Respiratory physiology & neurobiology, 316*, 104113. Advance online publication. https://doi.org/10.1016/j.resp.2023.104113

Cameron O. G. (2009). Visceral brain-body information transfer. NeuroImage, 47(3), 787–794. https://doi.org/10.1016/j.neuroimage.2009.05.010.

Craig A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nature reviews. Neuroscience*, *3*(8), 655–666. https://doi.org/10.1038/nrn894.

Curtis, C. E., & D'Esposito, M. (2003). Persistent activity in the prefrontal cortex during working memory. *Trends in cognitive sciences*, *7*(9), 415–423. https://doi.org/10.1016/s1364-6613(03)00197-9.

Damasio, A. R., Grabowski, T. J., Bechara, A., Damasio, H., Ponto, L. L., Parvizi, J., & Hichwa, R. D. (2000). Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nature neuroscience, 3*(10), 1049–1056. https://doi.org/10.1038/79871

Del Negro, C.A., Funk, G.D., Feldman, J.L.. Breathing matters. Nat Rev *Neurosci*. 2018 Jun, *19*(6), 351-367. doi: 10.1038/s41583-018-0003-6.

Dixon, M. L., Thiruchselvam, R., Todd, R., & Christoff, K. (2017). Emotion and the prefrontal cortex: An integrative review. *Psychological Bulletin, 143*(10), 1033–1081. https://doi.org/10.1037/bul0000096.

Dlouhy, B. J., Gehlbach, B. K., Kreple, C. J., Kawasaki, H., Oya, H., Buzza, C., Granner, M. A., Welsh, M. J., Howard, M. A., Wemmie, J. A., & Richerson, G. B. (2015). Breathing Inhibited When Seizures Spread to the Amygdala and upon Amygdala Stimulation. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *35*(28), 10281–10289. https://doi.org/10.1523/JNEUROSCI.0888-15.2015.

Dutschmann, M., & Dick, T. E. (2012). Pontine mechanisms of respiratory control. *Comprehensive Physiology*, *2*(4), 2443–2469. https://doi.org/10.1002/cphy.c100015.

Etkin, A., Büchel, C., & Gross, J. J. (2015). The neural bases of emotion regulation. *Nature reviews. Neuroscience*, *16*(11), 693–700. https://doi.org/10.1038/nrn4044.

Etkin, A., Egner, T., & Kalisch, R. (2011). Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends in cognitive sciences, 15*(2), 85–93. https://doi.org/10.1016/j.tics.2010.11.004

Feinstein, J. S., Buzza, C., Hurlemann, R., Follmer, R. L., Dahdaleh, N. S., Coryell, W. H., Welsh, M. J., Tranel, D., & Wemmie, J. A. (2013). Fear and panic in humans with bilateral amygdala damage. Nature neuroscience, 16(3), 270–272. https://doi.org/10.1038/nn.3323

Feinstein, J. S., Gould, D., & Khalsa, S. S. (2022). Amygdala-driven apnea and the chemoreceptive
origin of anxiety. *Biological psychology, 170*, 108305.
https://doi.org/10.1016/j.biopsycho.2022.108305

Fine, J. M., & Hayden, B. Y. (2021). The whole prefrontal cortex is premotor cortex. *Philosophical
Transactions of the Royal Society B, 377*(1844). https://doi.org/10.1098/rstb.2020.0524

Friedman, N. P., & Robbins, T. W. (2022). The role of prefrontal cortex in cognitive control and
executive function. *Neuropsychopharmacology, 47*(1), 72–89. https://doi.org/10.1038/s41386-021-
01132-0

Gabbott, P. L., Warner, T. A., Jays, P. R., Salway, P., & Busby, S. J. (2005). Prefrontal cortex in the rat:
projections to subcortical autonomic, motor, and limbic centers. *The Journal of comparative
neurology*, *492*(2), 145–177. https://doi.org/10.1002/cne.20738.

García, M. C., Wang, Y., Jiang, M., Anderson, M. C., & Lei, X. (2022). Anterior Cingulate Cortex Signals
the Need to Control Intrusive Thoughts during Motivated Forgetting. *The Journal of Neuroscience,
42*(21), 4342–4359. https://doi.org/10.1523/jneurosci.1711-21.2022

García-Cabezas, M. Á., & Barbas, H. (2017). Anterior Cingulate Pathways May Affect Emotions
Through Orbitofrontal Cortex. *Cerebral Cortex, 27*(10), 4891–4910.
https://doi.org/10.1093/cercor/bhw284

Gläscher, J., Adolphs, R., Damasio, H., Bechara, A., Rudrauf, D., Calamia, M., Paul, L. K., & Tranel, D.
(2012). Lesion mapping of cognitive control and value-based decision making in the prefrontal cortex.
Proceedings of the National Academy of Sciences of the United States of America, 109(36), 14681–
14686. https://doi.org/10.1073/pnas.1206608109

Gothe, N. P., Khan, I., Hayes, J., Erlenbach, E., & Damoiseaux, J. S. (2019). Yoga Effects on Brain
Health: A Systematic Review of the Current Literature. *Brain plasticity (Amsterdam, Netherlands),
5*(1), 105–122. https://doi.org/10.3233/BPL-190084

Graham, B. M., & Milad, M. R. (2013). Prefrontal cortex regulation of emotion and anxiety. In D. S.
Charney, J. D. Buxbaum, P. Sklar, & E. J. Nestler (Eds.), *Neurobiology of mental illness* (pp. 580–592).
Oxford University Press. https://doi.org/10.1093/med/9780199934959.003.0043

Harmata, G.I., Rhone, A.E., Kovach, C.K., Kumar, S., Mowla, M.R., Sainju, R.K., Nagahama, Y., Oya, H.,
Gehlbach, B.K., Ciliberto, M.A., Mueller, R.N., Kawasaki, H., Pattinson, K.T., Simonyan, K., Davenport,
P.W., Howard III, M.A., Steinschneider, M., Chan, A.C., Richerson, G.B., Wemmie, J.A., Dlouhy, B.J.
Failure to breathe persists without air hunger or alarm following amygdala seizures. (2023). *JCI
Insight*. e172423. https://doi.org/10.1172/jci.insight.172423.

Harper, R. M., Kumar, R., Macey, P. M., Woo, M. A., & Ogren, J. A. (2014). Affective brain areas and
sleep-disordered breathing. *Progress in brain research, 209*, 275–293. https://doi.org/10.1016/B978-
0-444-63274-6.00014-X

Hassanpour, M. S., Simmons, W. K., Feinstein, J. S., Luo, Q., Lapidus, R. C., Bodurka, J., Paulus, M. P.,
& Khalsa, S. S. (2018). The IC Cortex Dynamically Maps Changes in Cardiorespiratory
Interoception. *Neuropsychopharmacology*, *43*(2), 426–434. https://doi.org/10.1038/npp.2017.154.

Heimer, L., & Van Hoesen, G.W. (2006). The limbic lobe and its output channels: implications for
emotional functions and adaptive behavior. *Neurosci Biobehav Rev* 30(2):126-47.
https://doi.org/10.1016/j.neubiorev.2005.06.006.

Hernandez, L., Manning, J., & Zhang, S. (2019). Voluntary control of breathing affects center of pressure complexity during static standing in healthy older adults. *Gait & Posture, 68*, 488–493. https://doi.org/10.1016/j.gaitpost.2018.12.032

Herrero, J. L., Khuvis, S., Yeagle, E., Cerf, M., & Mehta, A. D. (2018). Breathing above the brain stem: volitional control and attentional modulation in humans. *Journal of neurophysiology*, *119*(1), 145–159. https://doi.org/10.1152/jn.00551.2017.

Herzog, M., Sucec, J., Jelinčić, V., Van Diest, I., Van den Bergh, O., Chan, P. S., Davenport, P., & von Leupoldt, A. (2021). The test-retest reliability of the respiratory-related evoked potential. *Biological psychology, 163*, 108133. https://doi.org/10.1016/j.biopsycho.2021.108133

Holroyd, C. B., & Yeung, N. (2012). Motivation of extended behaviors by anterior cingulate cortex. *Trends in cognitive sciences, 16*(2), 122–128. https://doi.org/10.1016/j.tics.2011.12.008

Holton, P., Huang, Y., Bahuri, N., Boccard, S., Hyam, J. A., Paterson, D. J., Dorrington, K. L., Aziz, T. Z., Moosavi, S. H., & Green, A. L. (2021). Differential responses to breath-holding, voluntary deep breathing and hypercapnia in left and right dorsal anterior cingulate. *Experimental physiology*, *106*(3), 726–735. https://doi.org/10.1113/EP088961.

Homma, I., & Masaoka, Y. (2004). Amygdala and emotional breathing in humans. *Advances in experimental medicine and biology*, *551*, 9–14. https://doi.org/10.1007/0-387-27023-x_2.

Homma, I., & Masaoka, Y. (2008). Breathing rhythms and emotions. *Experimental physiology*, *93*(9), 1011–1021. https://doi.org/10.1113/expphysiol.2008.042424.

Ikeda, K., Kawakami, K., Onimaru, H., Okada, Y., Yokota, S., Koshiya, N., Oku, Y., Iizuka, M., & Koizumi, H. (2017). The respiratory control mechanisms in the brainstem and spinal cord: integrative views of the neuroanatomy and neurophysiology. *The journal of physiological sciences: JPS*, *67*(1), 45–62. https://doi.org/10.1007/s12576-016-0475-y.

Jerath, R., & Beveridge, C. (2020). Respiratory Rhythm, Autonomic Modulation, and the Spectrum of Emotions: The Future of Emotion Recognition and Modulation. *Frontiers in psychology*, *11*, 1980. https://doi.org/10.3389/fpsyg.2020.01980.

Jodo, E., Chiang, C., & Aston-Jones, G. (1998). Potent excitatory influence of prefrontal cortex activity on noradrenergic locus coeruleus neurons. *Neuroscience*, *83*(1), 63–79. https://doi.org/10.1016/s0306-4522(97)00372-2.

Karjalainen, S., Kujala, J., & Parviainen, T. (2023). Neural activity is modulated by spontaneous and volitionally controlled breathing. *bioRxiv*. https://doi.org/10.1101/2023.08.30.555484

Khalsa, S. S., Rudrauf, D., Feinstein, J. S., & Tranel, D. (2009a). The pathways of interoceptive awareness. *Nature neuroscience*, *12*(12), 1494–1496. https://doi.org/10.1038/nn.2411.

Khalsa, S. S., Rudrauf, D., Sandesara, C., Olshansky, B., & Tranel, D. (2009b). Bolus isoproterenol infusions provide a reliable method for assessing interoceptive awareness. *International journal of psychophysiology : official journal of the International Organization of Psychophysiology, 72*(1), 34–45. https://doi.org/10.1016/j.ijpsycho.2008.08.010

Khalsa, S. S., Feinstein, J. S., Li, W., Feusner, J. D., Adolphs, R., & Hurlemann, R. (2016). Panic Anxiety in Humans with Bilateral Amygdala Lesions: Pharmacological Induction via Cardiorespiratory Interoceptive Pathways. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, *36*(12), 3559–3566. https://doi.org/10.1523/JNEUROSCI.4109-15.2016.

Khalsa, S. S., Rudrauf, D., Hassanpour, M. S., Davidson, R. J., & Tranel, D. (2020). The practice of
meditation is not associated with improved interoceptive awareness of the heartbeat.
Psychophysiology, 57(2), e13479. https://doi.org/10.1111/psyp.13479

Kim, S. H., Shin, H. J., & Cho, H. Y. (2022). Impact of Types of Breathing on Static Balance Ability in
Healthy Adults. *International journal of environmental research and public health, 19*(3), 1205.
https://doi.org/10.3390/ijerph19031205

Kluger, D. S., & Gross, J. (2021). Respiration modulates oscillatory neural network activity at rest.
PLoS biology, 19(11), e3001457. https://doi.org/10.1371/journal.pbio.3001457

Kluger, D.S., Forster, C., Abbasi, O. et al. (2023). Modulatory dynamics of periodic and aperiodic
activity in respiration-brain coupling. *Nat Commun 14*, 4699. https://doi.org/10.1038/s41467-023-
40250-9

Laborde, S., Allen, M. S., Borges, U., Dosseville, F., Hosang, T. J., Iskra, M., Mosley, E., Salvotti, C.,
Spolverato, L., Zammit, N., & Javelle, F. (2022). Effects of voluntary slow breathing on heart rate and
heart rate variability: A systematic review and a meta-analysis. Neuroscience and biobehavioral
reviews, 138, 104711. https://doi.org/10.1016/j.neubiorev.2022.104711.
729
730   Lafayette Instrument Company. (2002). Grooved Pegboard Test User instructions. Fabication
731   Enterprises   Inc.   Retrieved   September   8,   2023,   from   https://www.fab-
732   ent.com/MEDIA/41_INSTRUCTIONS/12-3010_MANUAL.PDF
733

Lara, A. H., & Wallis, J. D. (2015). The Role of Prefrontal Cortex in Working Memory: A Mini Review.
Frontiers in systems neuroscience, 9, 173. https://doi.org/10.3389/fnsys.2015.00173

Lenth, R., 2018. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version
1.1.3. https://CRAN.R-project.org/package=emmeans.

Lumb, A. (2016). Nunn's Applied Respiratory Physiology (8th ed., Vol. 8). Elsevier.

Martinu, K., Degroot, C., Madjar, C., Strafella, A. P., & Monchi, O. (2012). Levodopa influences striatal
activity but does not affect cortical hyper-activity in Parkinson's disease. *European Journal of*
Neuroscience, 35(3-4), 572–583. https://doi.org/10.1111/j.1460-9568.2011.07979.x.

McKay, L. C., Evans, K. C., Frackowiak, R. S., & Corfield, D. R. (2003). Neural correlates of voluntary
breathing in humans. *Journal of applied physiology (Bethesda, Md.: 1985), 95*(3), 1170–1178.
https://doi.org/10.1152/japplphysiol.00641.2002.

Nobis, W. P., González Otárula, K. A., Templer, J. W., Gerard, E. E., VanHaerents, S., Lane, G., Zhou,
G., Rosenow, J. M., Zelano, C., & Schuele, S. (2019). The effect of seizure spread to the amygdala on
respiration and onset of ictal central apnea. *Journal of neurosurgery, 132*(5), 1313–1323.
https://doi.org/10.3171/2019.1.JNS183157.

Pattinson, K. T., Governo, R. J., MacIntosh, B. J., Russell, E. C., Corfield, D. R., Tracey, I., & Wise, R. G.
(2009). Opioids depress cortical centers responsible for the volitional control of respiration. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience, 29(25), 8177–8186.
https://doi.org/10.1523/JNEUROSCI.1375-09.2009

Peng, C. K., Henry, I. C., Mietus, J. E., Hausdorff, J. M., Khalsa, G., Benson, H., & Goldberger, A. L.
(2004). Heart rate dynamics during three forms of meditation. *International journal of*
cardiology, 95(1), 19–27. https://doi.org/10.1016/j.ijcard.2003.02.006.

Prasad, S., Pal, P. K., & Chen, R. (2021). Chapter 1 - Breathing and the Nervous System. In M. Aminoff & S. A. Josephson (Eds.), *Aminoff's Neurology and General Medicine* (6th ed., pp. 3–19). essay, Elsevier Gezondheidszorg

Rhone, A. E., Kovach, C. K., Harmata, G. I., Sullivan, A. W., Tranel, D., Ciliberto, M. A., Howard, M. A., Richerson, G. B., Steinschneider, M., Wemmie, J. A., & Dlouhy, B. J. (2020). A human amygdala site that inhibits respiration and elicits apnea in pediatric epilepsy. *JCI insight*, *5*(6), e134852. https://doi.org/10.1172/jci.insight.134852.

Rolls E. T. (2023). Emotion, motivation, decision-making, the orbitofrontal cortex, anterior cingulate cortex, and the amygdala. *Brain structure & function, 228*(5), 1201–1257. https://doi.org/10.1007/s00429-023-02644-9

Russo, M. A., Santarelli, D. M., & O'Rourke, D. (2017). The physiological effects of slow breathing in the healthy human. *Breathe (Sheffield, England), 13*(4), 298–309. https://doi.org/10.1183/20734735.009817

Schön, D., Rosenkranz, M., Regelsberger, J., Dahme, B., Büchel, C., & von Leupoldt, A. (2008). Reduced perception of dyspnea and pain after right IC cortex lesions. *American journal of respiratory and critical care medicine*, *178*(11), 1173–1179. https://doi.org/10.1164/rccm.200805-731OC.

Shweta, P., Pramod, K., & Chen, R. (2021). Chapter 1 - Breathing and the Nervous System. In *Aminoff's neurology and general medicine* (6th ed., pp. 3–19). essay, Academic Press.

Silver, N. C., & Dunlap, W. P. (1987). Averaging correlation coefficients: Should Fisher's z transformation be used? *Journal of Applied Psychology, 72*(1), 146–148. https://doi.org/10.1037/0021-9010.72.1.146.

Silvers, J. A., Weber, J., Wager, T. D., & Ochsner, K. N. (2015). Bad and worse: Neural systems underlying reappraisal of high- and low-intensity negative emotions. *Social Cognitive and Affective Neuroscience, 10*(2), 172–179. https://doi.org/10.1093/scan/nsu043.

Terreberry, R. R., & Neafsey, E. J. (1987). The rat medial frontal cortex projects directly to autonomic regions of the brainstem. *Brain research bulletin*, *19*(6), 639–649. https://doi.org/10.1016/0361-9230(87)90050-5.

Trites, R. (1989). Grooved Pegboard Test. Lafayette Instruments.

Watanabe, J., Sugiura, M., Sato, K., Sato, Y., Maeda, Y., Matsue, Y., Fukuda, H., & Kawashima, R. (2002). The human prefrontal and parietal association cortices are involved in NO-GO performances: an Event-Related FMRI study. *NeuroImage, 17*(3), 1207–1216. https://doi.org/10.1006/nimg.2002.1198.

Yang, C. F., & Feldman, J. L. (2018). Efferent projections of excitatory and inhibitory preBötzinger Complex neurons. *The Journal of comparative neurology*, *526*(8), 1389–1402. https://doi.org/10.1002/cne.24415.

Young, C. B., Reddy, V., & Sonne, J. (2021). Neuroanatomy, Basal Ganglia. In *StatPearls*. StatPearls Publishing.

Yuan, P., & Raz, N. (2014). Prefrontal cortex and executive functions in healthy adults: a meta-analysis of structural neuroimaging studies. Neuroscience and biobehavioral reviews, 42, 180–192. https://doi.org/10.1016/j.neubiorev.2014.02.005.

796   Zaccaro, A., Piarulli, A., Laurino, M., Garbella, E., Menicucci, D., Neri, B., & Gemignani, A. (2018). How

797   Breath-Control Can Change Your Life: A Systematic Review on Psycho-Physiological Correlates of

798   Slow Breathing. *Frontiers in human neuroscience*, *12*, 353.

799   https://doi.org/10.3389/fnhum.2018.00353.

     Table 1. Demographic characteristics of the lesion and healthy samples.

Variables	Lesion (n = 20)	Healthy (n = 20)	χ2	t-ratio (df)	p-value
Age (years)				-0.07 (33)	0.94
Mean ± SD	53.8 ± 7.1	53.6 ± 10.6			
Sex (%)					
Female	70	70			
Male	30	30			
BMI (kg / m²)				1.81 (35)	0.08
Mean ± SD	27.6 ± 5.0	30.7 ± 5.9			
Median (IQR)	26.7 (23.5 - 29.6)	32.9 (27.1 - 34.5)			
Handedness (n)			2.11		0.15
Right	18	20			
Left	2	0			
Education					
Mean ± SD	14.1 ± 1.9	16.6 ± 2.1		-3.8 (38)	<0.001
Race (number)			1.02		0.31
White / Caucasian	20	19			
Asian	0	1			
Black	0	0			
Hispanic	0	0			
Mixed	0	0			

Note. Data shown compared by *t*-test or χ²- test. BMI: Body mass index; IQR: Interquartile range; kg: Kilogram; m²: Square meter; n: Sample size; SD: Standard deviation;

Figure 1

Overview of Brain Areas Involved in Voluntary and Automatic Respiration and Lesion Overlap Map for Participants with Brain Lesions.

A   **Voluntary Respiration Brain Regions (putative)**

C   **Lesion Overlap Map**

lateral   horizontal   saggital

B   **Automatic Respiratory Network**

Note. **(A)** displays the location of cortical areas putatively involved in the regulation of voluntary breathing projected onto lateral, horizontal, and mid-sagittal surface views of the brain. In addition to the respiratory centers of automatic breathing in the FR in the brainstem, several higher-order subcortical and cortical areas are putatively involved in voluntary breathing. The precise manner in which these regions control voluntary respiration is unknown. Abbreviations: Amyg – Amygdala, CB – Cerebellum, HC – Hippocampus, ILC - Infralimbic cortex, LC - Locus coeruleus, M1 - Primary motor cortex (1), OC - Olfactory cortex, OFC - Orbitofrontal cortex, PAG - Periaqueductal gray, PMA - Pre-motor cortex, SMA - Supplementary motor area, S1 - Sensory motor cortex (S1), TH – Thalamus. **(B)** shows the location of the major brainstem respiratory groups projected onto the planes of a lateral sagittal section (2) and a ventral view (3). The respiratory groups (right inset (2) and (3)) form a set of interacting structures. The Pontine Respiratory Group (PRG) in the superior part of the pons; sends inhibitory signals to the Dorsal respiratory group (DRG) and controls the filling phase of respiration. DRG receives impulses from peripheral chemoreceptors and other receptors via vagus and glossopharyngeal nerves. These impulses produce inspiratory movements and are responsible for the basic rhythm of breathing. The Ventral Respiratory Group (VRG) causes either inhalation or exhalation. It is inactive during quiet breathing but plays an important role in stimulating abdominal expiratory muscles during high respiratory demand. Abbreviations: DRG - Dorsal Respiratory Group, PRG - Pontine Respiratory Group, VRG - Ventral Respiratory Group. **(C)** This figure shows the cortical and subcortical overlap of lesions for the 20 lesion participants via lateral (1) and axial (2) view. Lesions were concentrated in the left temporal lobe, bilateral frontal lobe, the cingulate and insular cortices, and in subcortical structures such as the basal ganglia.

Figure 2

831 *Overview of the Experimental Setup and Overall Performance for the Regular and Irregular Breathing*

832 *and Motor Control Tasks.*

A   Experimental Setting

B   Visual Examples of the Instructed Waveforms

C   Group by Task Interaction Plot for Regular Tasks

D   Group by Task Interaction Plot for Irregular Task

Note. **(A)** Participants were seated upright in front of a screen showing the corresponding instruction signal as a wave
movement with a red dot moving up and down. **(B)** Exemplary 15-second illustration of instructed waveforms for regular and
irregular experimental task conditions. For the regular version of the respiratory and motor regulation tasks, there were four
experimental conditions ranging from 6 to 60 breaths (or repetitions) per minute. For the irregular task, both the amplitude
and rate were randomly varied, with a frequency range from 6 to 60 breaths (or repetitions) per minute. Each experimental
condition in the study lasted 2 minutes. Mean group performance across all conditions of the **(C)** regular task and **(D)** irregular
task; performance was indexed via Fisher`s Z-transformed maximum cross correlation values (range: –2.65 to 2.65). P-values
indicate the group by task interaction. Error bars indicate standard error.

855 **Figure 3**

856 *Mean Anxiety Ratings for All Task Types and Conditions*

858
859
860 *Note.* P-values indicate post-hoc between-group differences in anxiety for the 60-bpm condition of the regular breathing task
861 and for the irregular breathing condition, calculated following observation of a significant group by condition interaction.
862 Error bars indicate standard error.