# Electrical brain activations in young children during a probabilistic reward-learning task are associated with behavioral strategy.

2  **Yu Sun Chung[1*], Berry van den Berg[2], Kenneth C. Roberts[3], Marty G. Woldorff     [1,3], Michael**
3  **S. Gaffrey[1,4,5]**

4  [1] Department of Psychology and Neuroscience, Duke University, Reuben-Cooke Building, 417
5  Chapel Drive, Durham, NC 27708, USA
6
7  [2] Experimental Psychology, University of Groningen, The Netherlands
8
9  [3] Center for Cognitive Neuroscience, Department of Psychiatry, Psychology & Neuroscience and
10  Neurobiology, Duke University, Durham, NC, 27708
11  USA.
12
13  [4] Children's Wisconsin, 9000 W. Wisconsin Avenue, Milwaukee, WI, 53226
14
15  [5] Medical College of Wisconsin, Division of Pediatric Psychology and Developmental Medicine,
16  Department of Pediatrics, 8701 Watertown Plank Road, Milwaukee, WI, 53226
17
**\* Correspondence:**
Corresponding Author
Yu Sun Chung, Ph.D. [1]
Reuben-Cooke Building
417 Chapel Drive
Campus Box 90086
Duke University
Department of Psychology and Neuroscience
Durham, NC 27708
Email: yusun08@gmail.com
28

**Keywords**: reinforcement learning, reward-related positivity, attention, N2pc, P300, win-stay-lose-
switch strategy

**Abstract**

Both adults and children learn through feedback which events and choices in the environment are
associated with higher probability of reward. This probability reward-learning ability is thought to be
supported by the development of fronto-striatal reward circuits. Recent developmental studies have
applied computational models of reward learning to investigate such learning in children. However,
there has been limited development of task tools capable of measuring the cascade of neural reward-
learning processes in children. Using a child-version of a probabilistic reward-learning task while
recording event-related-potential (ERP) measures of electrical brain activity, this study examined key
processes of reward learning in preadolescents (n=30), namely: (1) reward-feedback sensitivity, as
measured by the early reward-related frontal ERP positivity, (2) rapid attentional shifting of
processing toward favored visual stimuli, as measured by the N2pc component, and (3) longer-
latency attention-related responses to reward feedback as a function of behavior strategies (i.e., Win-
Stay-Lose-Shift), as measured by the central-parietal P300. Consistent with our prior work in adults,
the behavioral findings indicate that preadolescents could learn stimulus-reward outcome
associations, but at varying levels of performance. Neurally, poor preadolescent learners (those with
slower learning rates) showed greater reward-related positivity amplitudes relative to good learners,
suggesting greater reward sensitivity. We also found attention shifting towards to-be-chosen stimuli,
as evidenced by the N2pc, but not to more highly rewarded stimuli. Lastly, we found an effect of
behavioral learning strategies (i.e., Win-Stay-Lose-Shift) on the feedback-locked P300 over the
parietal cortex. These findings provide novel insights into the key neural processes underlying
reinforcement learning in preadolescents.

1. **Introduction**

From an early age, children learn about the world around them through trial and error. That is, through repeated experience, children progressively learn what actions are likely to result in relatively positive or negative outcomes given a specific environment and/or set of choices. The ability to learn associations between choice behavior and outcomes and change patterns of subsequent choice behavior in light of this information, has been called probabilistic reinforcement learning (PRL). Computational models of PRL separate such learning into computationally-driven components associated with both behaviorally and neurobiologically distinguishable processes. Specifically, PRL model involves a sequence of neurocognitive processes, including evaluation of the relative value associated with available choices, updating of stimulus-reward outcome associations based on feedback, and shaping of selective attention toward a more-likely-winning stimulus to guide future choice (Anderson, 2017). However, while the neural correlates of PRL have been well studied in adults, relatively few studies have investigated this question in adolescents or children (reviewed by (DePasque & Galván, 2017). This is a critical knowledge of gap because the study of PRL in children is highly promising for identifying specific neurocognitive processes that may account for differences in  these important cognitive functions across development (Nussenbaum & Hartley, 2019b) and/or may predict risk for depression, which is heightened during adolescence (Chen et al., 2015; Keren et al., 2018a).

At the behavioral level, PRL is evident during childhood, adolescence, and adulthood (Raab & Hartley, 2018). However, findings from studies using computational models of PRL to examine how value-based learning ability changes with age have been mixed. As reviewed by (Nussenbaum & Hartley, 2019a), some developmental studies report that learning rates increase from childhood to adulthood (Davidow et al., 2016; Master et al., 2020) while others report that they do not change with age (Palminteri et al., 2016). Such mixed behavior findings may reflect individual differences in learning rates that may come from neurobiological alterations in one or all of the processes underlying PRL. For example, learning rate captures individual differences in the value-updating process, reflecting the degree to which reward feedback and reward prediction errors are incorporated into the updated value estimates during learning. A number of empirical studies using functional Magnetic Resonance Imaging (fMRI) have suggested that reward prediction errors are signaled by mid-brain dopaminergic, fronto-striatal reward brain circuits (Averbeck & O'Doherty, 2022; Nasser et al., 2017; Schultz, 2016), including in adolescents  (Christakou et al., 2013; Cohen et al., 2010; Hauser et al., 2015).  However, the sluggishness of fMRI signals limits the ability of this method to the different subprocesses underlying learning on a trial-by-trial basis, which requires more precise temporal resolution.

The high temporal resolution of electroencephalogram (EEG) data makes it particularly well-suited for studying the neural subprocesses associated with PRL.  More specifically, the ability to measure event-related potentials (ERPs) time-locked to stimulus and reward-feedback events provides an exceptional opportunity to delineate the temporal cascade of cognitive processes related to reward processing within a trial (Amodio et al., 2014). One ERP component extensively examined in reward studies is the reward positivity (RewP), a frontal, positive-polarity, brain wave measured in response to gain minus loss feedback, occurring approximately 250-350ms after that feedback. With respect to RL theory (Holroyd & Coles, 2002; Sutton & Barto, 2018), the RewP is thought to reflect a reward prediction error signal that is sensitive to outcome valence and is larger for unexpected positive events relative to unexpected negative events (Holroyd & Coles, 2002). Although the RewP (or its inverse, the Feedback-related Negativity, FN**,** calculated as loss-minus gain in some earlier research (Krigolson, 2018)) as a measure of response to reward has been investigated in children, its role in PRL during this age has not.

As one learns which stimuli are more likely to generate reward outcomes, one tends to pay
more attention to the stimuli associated with higher probability of leading to reward. Such attention
shifting toward reward-associated stimuli has been indexed by the lateralized attention-shifting-
sensitive N2pc component in prior work with adults (San Martin et al., 2016; Hickey et al., 2010). In
contrast, there have been only a handful of ERP studies that have examined the N2pc in children,
either in the context of a reward task or more generally during visual search (Couperus & Quirk,
2015; Li et al., 2022; Shimi et al., 2014; Sun et al., 2018; Turoman et al., 2021) and/or working
memory tasks (Rodríguez-Martínez et al., 2021; Shimi et al., 2015). Consistent with prior work with
adults, existing studies with children indicate the elicitation of the N2pc component during non-PRL,
attentional tasks, indicating that this component can be used as a marker of attentional section in
children like in adults (Couperus & Quirk, 2015; Li et al., 2022; Shimi et al., 2014; Sun et al., 2018;
Turoman et al., 2021). However, the presence of this attentional neural process has not been directly
tested within the framework of computational PRL model.  Accordingly, we know little about the
functionality of the N2pc ERP component in children within a PRL framework – that is, whether it
may reflect an index of early attentional shifting as a function of reward learning in children, as
shown previously in adults.

Lastly, the P300 has been thought more generally to reflect longer-latency attentional
processes involved in information updating of salient outcomes during evaluative processing (for
reviews, see Glazer et al., 2018, San Martín, 2012). The P300 is a centro-parietally distributed,
positive-going wave that peaks between 400 and 600 ms following a stimulus (San Martín et al.,
2013; Wu & Zhou, 2009). It has been successfully used to investigate PRL with adults (e.g.,
(Donaldson et al., 2016). More specifically, prior work has shown that P300 is sensitive to a later,
top-down-control process of outcome evaluation, suggesting that outcome-related feedback
information in a probability choice task (e.g., probability and magnitude) is used to guide future
decisions for the goal of reward maximization (San Martín et al., 2013, Wu and Zhou, 2009, Zhang et
al., 2013). However, like the N2pc, the functionality of the feedback locked P300 (reward gain vs.
loss feedback) during PRL in children has been rarely examined.

The purpose of the current study was to establish the feasibility and initial validity of a
developmentally appropriate probabilistic reward learning task, which we have termed the C-PLearn
task (for Child Probability Learning task).  Based on our results, we believe that this task would be a
promising tool for studying both typical and atypical development of key neural processes underlying
PRL. For example, alterations in PRL abilities have been known to be critical components of hedonic
functioning that predicts depression risk for children and adolescents (Keren et al., 2018b; Morris et
al., 2015; Saulnier et al., 2023). The design of the C-PLearn task was based on our prior work
investigating neural correlates of PRL processes in adults (van den Berg et al., 2019b). In the present
study, we examined key processes of PRL in a group of preadolescent children, namely (1) reward-
feedback sensitivity, as measured by the RewP, (2) early attentional processing toward stimuli with
higher reward value or likelihood as a shaped by reward learning, as measured by the N2pc, (3)
longer-latency attentional processing in response to the feedback as a function of behavioral learning
strategies (i.e., Win-Stay-Lose-Shift, WSLS (Hernstein et al., 2000)), as measured by the central-
parietal P300. More specifically, first we wanted to replicate the presence of the RewP (reflecting
reward-feedback sensitivity and reward prediction errors), the N2pc (reflecting the shifting of
attention to certain stimuli versus others), and the P300 (reflecting longer-latency attention-related
cognitive processes) as a function of reward learning for preadolescents within the PRL framework.
Second, we predicted that preadolescents who didn't learn the stimulus-outcome associations very
well (i.e., "bad" learners) would show smaller RewP amplitudes, consistent with the RL framework,
compared to "good" learners, and/or may adopt less effective behavior strategy.     Finally, we

hypothesized that the central-parietal, feedback-locked P300 may reflect a modulation of later
cognitive processing to update reward value as a function of behavior strategy for PRL learning (e.g.,
WSLS) in preadolescents, an idea that has been supported by the literature in adults (Donaldson et
al., 2016; von Borries et al., 2013).
153
2. **Methods**
2.1. Participants
Thirty children between 8 to 12 years old (mean age = 9.94 years ±1.26 (SD); 15 female;
30 right-handed; 20% Hispanic or Latino ethnicity, 80% non-Hispanic white) were recruited from a
Research Participation Database at Duke University maintained by the Human Subjects Coordinator
for the Department of Psychology and Neuroscience. In addition, methods included posting flyers on
social media sites and distributing flyers to local schools were used to recruit participants.
Children who were color blind, reported a current and/or historical diagnosis of any
psychiatric disorder, had a history of neurological disorder or insult (e.g., seizures, hydrocephalus,
cerebral palsy, brain tumor, extended loss of consciousness, head trauma), took psychotropic
mediation for a mood or behavioral difficulty, or left-handed were excluded from study participation.
At the end of EEG sessions, a $30 Amazon e-gift card was sent to parents via email for their
participation in the study. Children also received candy and/or small toys after each completed
session. Additional compensation was provided ($10 Amazon e-gift card per hour) to parents if
testing sessions exceeded 2.5 hours. This study was conducted in accordance with protocols
approved by the Duke Institutional Review Board.
170
2.2. The Child Probabilistic Reward Learning Task (C-PLearn)
The C-PLearn task used in the current study for children was based on our adult PLearn task
previously used successfully in college students (van den Berg et al., 2019b). Participants completed
the C-PLearn, followed by a face-vs-house neural localizer task (for localizing the face-selective
neural activity on the scalp), while EEG was recorded. The task was programmed and run on the
OpenSesame platform for behavioral research (Mathôt et al., 2012), and event codes were sent using
the Python 'egi' package.
The C-PLearn consisted of 20 blocks of 18 trials each,    slightly different than the 20 trials
per block in the original adult version to accommodate reduced attention spans and lower tolerance
for prolonged experimental tasks in children. Also, the probability of reward was set to 0.77 (instead
of ranging randomly from 0.5 to 0.75 as in the original adult version) to mitigate child frustration and
potential confusion noted during early piloting sessions using the original adult reward probability
levels. At the beginning of each trial, participants were presented with two images, one of a face and
one of a house, and were instructed to choose between the two. They were also instructed to try to
learn during each block whether faces or houses were more likely to result in winning points, and that
winning more points would allow them to choose small toys with differing levels of cost. Each block
was followed by feedback as to how many points they had accrued up until that point. After a
practice period in which the subject needed to demonstrate understanding of how to do the task, the
test session started with the C-PLearn task runs.
190
191
192
2.3. Computational modeling of reward learning behavior
To calculate each subject's reward learning behavior within the PRL framework, we defined a
win-stay (WS) trial as those in which the previous trial ($n − 1$) was rewarded (won) and the choice on
the current trial $n$ is the same as the choice on trial $n − 1$ (i.e., "stayed" with the same choice of face

or house made on the previous trial). We then calculated WS probabilities as the proportion of trials
with WS behavior given a previously rewarded trial (i.e., after receiving gain feedback). Also, we
defined a lose shift (LS) trial as those in which the previous trial ($n − 1$) was not rewarded (i.e.,
receiving loss feedback) and the choice on the current trial $n$ differs from the choice on trial $n − 1$.
We then calculated LS probabilities as a proportion of trials with LS behavior given that the previous
trial was unrewarded (i.e., after receiving loss feedback). Similarly, WS probabilities were calculated
as a proportion of trials with WS behavior given that the previous trial was rewarded (i.e., after
receiving gain feedback).
205
2.4. EEG Data Collection and Processing
Continuous EEG was recorded using EGI's Net Station acquisition software version 5.4.2 and
Net Amps 400 series amplifiers with the associated 128-channel Geodesic Sensor Net system, at a
sampling rate of a 1000 Hertz (Hz per channel). Data was referenced online to the vertex (Cz) during
acquisition, and impedances were maintained below 50 kilohms throughout the paradigm.
Offline EEG data preprocessing was performed using EEGLAB version 2021 (Delorme &
Makeig, 2004) in MATLAB R2019b (The MathWorks Inc, Natick, MA) using custom scripts (codes
are available at:https://github.com/DEEDLabEEG). According to the Maryland analysis of
developmental EEG pipeline recommended for pediatric populations (Debnath et al., 2020), we
followed several preprocessing steps as follows. First, twenty-four channels located on the outer ring
of the sensor net were removed from the analyses. As participants were required to wear masks
during the EEG sessions according to Duke COVID-related policy, these channels contained many
artifacts during the recordings in these children due to poor connections. The data were then
downsampled to 250Hz and segments of data without relevant events (i.e., breaks) were removed.
Using the ERPLAB v8.2 plugin (Lopez-Calderon & Luck, 2014), a 0.1 to 30 Hz, 4th order
Butterworth, bandpass filter was applied. CleanLine (Kappenman & Luck, 2012) was used to remove
any remaining 60 Hz line noise (Mitra & Bokil, 2007). All data were re-referenced to the average of
the mastoids. Bad channels were removed by running the Clean Rawdata plugin; a channel was
considered bad if 1) it was flat for more than five seconds, 2) contained more than four standard
deviations of line noise relative to its signal, or 3) correlated at less than .8 to nearby channels.
According to several recommendations (Debener et al., 2010; Debnath et al., 2020), a "copy"
of the data was made and then, the following steps were applied (Delorme & Makeig, 2004): 1) the
high-pass filter at a 1 Hz (4th order Butterworth) , 2) Artifact Subspace Reconstruction (ASR;(Mullen
et al., 2015) with the burst criterion set to 20 (Chang et al., 2018) to remove large artifacts with Clean
Rawdata, and 3) extended infomax Independent Component Analysis (ICA; (Lee et al., 1999) with
PCA dimension reduction (50 components). The resulting ICA matrix was then copied over to the
original full-length data (i.e., the data just before the copy was made and the 1 Hz high-pass filter was
applied). The ICLabel plugin (Pion-Tonachini et al., 2019) automatically removed independent
components with a probability greater than .7 of being associated with eye movements and blinks.
Application of the ICA matrix to the full-length data and subsequent removal of eye-related
components allowed for the preservation of data that would have otherwise been removed by ASR or
other artifact removal methods. Bad channels that were previously removed from the analyses were
interpolated back into the data set using spherical splines (Perrin, 1989).
For both the C-PLearn task and the localizer task, epochs were extracted from 400 ms before
until 800 ms after the onset of the stimulus (either the cue-pairs or feedback screens in the main task
or face/house images in the localizer task). All epochs were baseline corrected from -200 ms to the
onset of the stimuli. Artifact rejection using the TBT plugin (Trial-By-Trial basis; (Ben-Shachar,
2018)) removed epochs with at least 10 channels meeting the following criteria: 1) peak-to-peak
amplitudes exceeding 100 µV within 200 ms windows sliding across the epoch by 20 ms increments,
2) voltages below -150 or greater than +150 µV, or 3) joint probabilities above three standard

deviations for local and global thresholds. If less than 10 channels in an epoch met criteria for
rejection, the epoch was not removed, but the identified channels were interpolated for that epoch
only. But if there were more than 10 channels in the epoch that met criteria for rejection, the epoch
was not included. Lastly, the averaged ERPs for each bin (i.e., Gain, Loss) were computed. Mean of
the number of accepted Gain and Loss trials per subject were 184.3 and 128.2, respectively.
251
2.5. Data binning and averaging
The epoched EEG data were binned according to feedback (gains and losses) and choice (face
and house), which resulted in an average number of trials for each subject (Brown et al.) for face gain
[90.53 (± 24.97)], face loss [62.13(±13.73)], house gain [93.76(±19.60)], and house loss [66.06
(±14.76)], after rejection of noisy epochs. As expected, due to learning, there were significantly more
gain trials than loss trials [$F(1,29) = 43.52$, $p < .001$]. On the other hand, there were no significant
differences in the number of gain or loss trials for choosing a face vs a house ([$F(1,29) = 0.46$, $p$
$=.50$].
260
2.6. Analysis of cue-evoked ERP data
We calculated the cue-evoked N2pc response to the cue-pair at the beginning of each trial, to
assess attentional orientation to the two stimulus types as a function of what they will later choose
that trial, following a similar procedure used in our prior work in adults (van den Berg et al., 2019b).
More specifically, the N2pc neural responses were assayed by the standard N2pc contralateral-vs-
ipsilateral analysis this component (Luck & Kappenman, 2012), that is by subtracting the activity in
the contralateral channels (relative to the chosen side) minus the ipsilateral channels and then
collapsing over the left and right sides.
269
2.7. Statistical Analysis
For statistical analysis, we first calculated the time-locked ERPs as a function of the various
event types and conditions. Based on previous literature, the reward-related Positivity (RewP) was
measured from 275 to 375 ms in a fronto-central ROI (E10, E11, E16). The P300 was measured from
400 to 600 ms in a central-parietal ROI (Left/Right: E37/E87, E31/E80, E53/E86, E60/E85, E67/E77,
E61/E78, E54/E79; middle line: E129, E55, E62, E71; (Fischer & Ullsperger, 2013; San Martín,
2012; San Martín et al., 2013). Subsequently, the extracted values were analyzed with a      repeated-
measures analyses of variance (ANOVAs) to test for statistical significance ($p < .05$) using the SPSS
2021 version.
The EEG data collected during the independent localizer task were used to confirm that the
stimuli that we used in this study elicited face-vs-house activity differences as has been observed in
the existing literature and to delineate potential influences of reward learning on stimulus-specific
regions. To do so, we first visually inspected all posterior electrodes (i.e., over visual cortex) for the
ERP differences between the neural responses to face and house images to make sure these effects
were representative of previously reported findings with adults, including the hallmark, face-
selective, lateral inferior occipital N170 responses. The N170 was measured from 170 to 220 ms after
onset of the cue image, from corresponding left and right occipital ROIs (E66/E84, E67/E91,
E61/E92).
The cue-related attentional bias (N2pc) was measured from 175 to 225 ms after onset of the
cue image pair, from corresponding left and right occipital ROIs (E50/E101, E58/96; (Hickey et al.,
2010; Kappenman & Luck, 2012; San Martín et al., 2016)). Subsequently, we also calculated the
difference in voltage contralateral vs ipsilateral relative to the side on which the set-winner was
presented.
To calculate each subject's individual behavioral learning rate, we used the same formula (see
(van den Berg et al., 2019b) for detailed formula) and calculated on a trial-by-trial number basis

using a mixed-modeling approach using the *lme4* statistical package (Bates et al., 2015). A varying
slope of condition per subject (the random effect) was included in the model if the Akaike
Information Criterion (a measure of the quality of the model) improved. Statistical significance was
set at $p < 0.05$, with the Satterthwaite's degrees of freedom method  as given by the R package *l*
*merTest* (Kuznetsova et al., 2017).
300
301
302
303

**3.  Results**

3.1. Behavioral Results
Consistent with our prior work in adults (van den Berg et al., 2019b), and as predicted,
children demonstrated behaviorally reinforcement learning ability based on trial-by-trial reward-
related feedback. The probability learning behavior across blocks in children for choosing the
probabilistic set-winner ranged from 0.40 to 0.73 successful, indicating that, as expected, the children
were able to integrate reward information throughout the block through reinforcement learning. As
presented in Figure 2.a, at the beginning of each 18-trial set, the participants chose the most likely
winner for that set at chance level (0.45-0.50), as would be expected. But by the end of the set, this
proportion of choosing set-winner increased up to ~0.75 on average, indicating that the children had
used the feedback across the trials of that set to learn to choose the more likely probability-based
winner (Fig2a). That said, the individual learning curves revealed very large individual difference in
reward learning, as the final proportion of choosing set winners in a particular block ranged from
0.26 to 1.0 across the subjects and blocks. Due to this large range, we wanted to explore whether
'good' vs. 'bad' learners might show different patterns of reward-related ERP components, in
particular by dividing the subjects using a median split based on their learning performance (Fig1).
320
As presented in Figure 2, good vs. bad learners adopted different behavior strategies during
the C-PLearn task. Good learners were more likely to choose the same stimulus type (i.e., Win-Stay,
WS behavior strategy) on the trial after one with a gain feedback ($t(28) = 3.92$, $p < .001$, Cohen's $d$
$=1.43$). In sharp contrast, bad learners were more likely to change their response choice (i.e., Lose-
Shift, LS behavior strategy) on the next trial after loss feedbacks ($t(28) = -2.89$, $p = 0.007$, Cohen's $d$
$=1.05$).
327
328         ##### figure 1 here #####.
329
330          ##### figure 2 here #####.
331
332    3.2. Electrophysiological Measures
333
334    3.2.1. The Reward-related Positivity (RewP)
335         Consistent with prior work, we confirmed the presence of the RewP around 275-375 ms after
336    gain vs. loss feedback in our frontal ROI (E10, E11, E16) for young children (see Figure 3.1).
337    According to the independent $t$-tests, bad learners had greater RewP amplitudes in the frontal ROIs
338    compared to good learners [$t(27) = -2.14$, $p = .04$, Cohen's $d = 0.79$] (see Figure 3.2     )
339
340         ##### figures 3.1 and 3.2 here #####.
341
3.2.2. P300 during reward gain vs. loss feedback as a function of behavior strategy

We examined updating reward values as a function of reward learning, as reflected by
behaviorally dependent modulation of the P300. Figure 5.1 shows the P300 during gain vs. loss
feedback as a function of behavior strategy (i.e., Switch vs. Stay) between good and bad learners.
Using P300 amplitudes in the central ROIs between 400 and 600 ms for good and bad learners, we
conducted repeated measures ANOVAs with two within-subject factors (i.e., behavior strategy:
switch vs stay, feedback: gain vs. loss) and one between-subject factor (i.e., good vs. bad learners).
The ANOVAs were conducted to examine whether bad learners showed greater P300 amplitudes

than did good learners during gain vs. loss feedback as a function of switch behavior strategy
compared to good learners. As shown in Figure 4.2(a), bad learners (i.e., those with low learning
rates), relative to good learners, showed higher P300 amplitude differences in central, parietal ROIs
when they switched their responses on the next trial after gain feedback vs. after loss feedback. The
repeated ANOVA analysis revealed no main effect on the P300 of learner type [$F(1, 27) = 1.45$, $p$
$=.23$, $\eta_p^2$: 0.05]. We did observe, however, main effects of behavior strategy [$F(1, 27) = 9.11$, $p$
$=.005$, $\eta_p^2$: 0.25] and feedback [$F(1, 27) = 7.00$, $p =.01$, $\eta_p^2$: 0.21], as presented in Figure 4.2.(b) and
(c), respectively, but no interaction of feedback by behavior strategy [$F(1, 27) = 0.34$, $p =.56$, $\eta_p^2$:
0.01] or of behavior strategy by learner type [$F(1, 27) = 0.38$, $p =.54$, $\eta_p^2$: 0.01] or feedback by
behavior strategy by learner type [$F(1, 27) = 0.09$, $p =.76$, $\eta_p^2$: 0.003] (see Figure 4.2). According to
the *post-hoc* paired *t*-tests, after receiving gain vs. loss feedback, children showed greater P300
amplitude differences in the central, parietal ROIs when they switched their choice on the subsequent
trial compared to when they chose the same responses [$t(28)= 2.68$, $p = .012$, Cohen's $d =0.32$ ] (see
Figure 4.2.b). Also, P300 amplitudes in the same central and parietal ROIs were higher when
children received loss feedback compared to gain feedback, collapsed across both good and bad
learners [$t(28)= 3.03$, $p = .005$, Cohen's $d = 0.39$] (see Figure 4.2.c)

##### figure 4.1  and 4.2 here #####.
367

3.2.4. N2pc reflects attentional bias towards choice behavior

In addition, in order to examine attention shift towards behavior choice regardless of whether
or not reward learning had been consolidated, we analyzed the effect of choice on the N2pc over the
course of the 18-trial set. We collapsed over whether the chosen image was that of a face or a house
to increase signal-to-noise ratio. We analyzed the relationship between the N2pc elicited by the cue-
pair stimulus relative to the choice made later in the trial, regardless of what the set-winner was. As
expected, we found the presence of N2pc around 175-225 ms after the onset of each cue-pair
stimulus in the ROIs (E50/E58) [$t(29)=$ , $p <.008$]. These results indicate that children's reward
learning behavior was reflected by the modulation of the N2pc, indexing early attentional bias
towards their choice behavior.  Like our prior work with college students (van den Berg et al., 2019),
we did not analyze choice behavior in the analysis of the neural N2pc. Because if one calculates the
N2pc amplitudes and its changes with learning across a block separately for when faces vs. houses
were the set-winner, the lateralized attention-related N2pc activity could be potentially conflated with
lateralized face-selective N170 activity. Independent *t*-test on the N2pc amplitudes measured in the
ROIs from 175 to 275 ms between good and bad learners revealed no statistical differences ($t(28) = -$
1.76, $p=.09$), even though the behavior indicated that they had learned the reward association.

##### figure 5 here #####.

**4.  Discussion**

Consistent with our prior work in adults (van den Berg et al., 2019b), our behavioral C-PL
earn findings indicate that at a group level, preadolescents are able to learn which stimulus type in a
pair (a face vs. a house) is more likely to result in a higher probability of reward over a series of
trials. As we observed large individual differences in individual learning rates, however, we divided
all participants into good (i.e., with high learning rates) vs. bad learners (i.e., with low learning rates).
Further, our behavior strategy analysis revealed that bad learners used a different strategy than did
good learners, namely a "Lose-Shift" behavior strategy rather than a "Gain-Stay" strategy.  In line
with prior work with adults (Worthy & Maddox, 2014), choosing the same choice on the next trial
after receiving reward feedback, namely, "Gain-Stay" strategy appears to be the better heuristic
learning strategy for preadolescent children in binary decision-making situations.

At the neural level, the current study's findings support C-PLearn as a developmentally
appropriate task for identifying and measuring key ERPs associated with PRL processes and
behavior in preadolescents. In contrast to our predictions, we found that poor preadolescent learners
with low learning rates showed greater RewP ERP amplitudes than good learners, indicating greater
reward-feedback sensitivity and/or greater prediction errors. However, our behavior results revealed
that poor learners used a different strategy than did good learners, namely a "Lose-Shift" behavior
strategy, in line with prior work with adults (Worthy & Maddox, 2014), rather than a "Gain-Stay"
strategy. Furthermore, we found early attention modulation toward the stimulus to be chosen, as
evidenced by the N2pc, although not toward the stimulus of higher likelihood of reward, which has
been shown in adult subjects (van den Berg et al. 2019a). Lastly, we demonstrated the contextual
modulation of feedback-locked P300 over the parietal cortex as a function of behavior strategy in
preadolescents. These results provide novel insights into the associations between RL-related ERP
components and computationally-driven reward learning behavior (i.e., Win-Stay-Lose-Shift, WSLS:
(Hernstein et al., 2000). As a result, the current study supports the use of C-PLearn to measure key
behavioral and neural correlates of PRL processing in preadolescent children.

To our knowledge, this is the first study, not just to examine the key neural processes
underlying PRL learning (i.e., reward-related feedback processing and attention shaping as a function
of reward learning) in preadolescents, but also with a combination of PRL and WSLS models. Most
existing ERP studies with children have focused on the RewP effect on general reward feedback
processing (e.g., (Burani et al., 2019; Ethridge et al., 2017; Ferdinand et al., 2016; Kessel et al., 2016;
Kessel et al., 2019; Lukie et al., 2014). However, little is known about the RewP in the context of
reinforcement learning in preadolescents. This is critical gap of knowledge in the field because poor
PRL learning may be a key process for underlying adolescent-onset of depression (Chen et al., 2015;
O'Callaghan & Stringaris, 2019). Although the exact underlying mechanisms are not yet clear, a
meta-analytic review provides strong evidence that depression is associated with impaired brain
signals of reward prediction error and with poor reinforcement learning behavior (Halahakoon et al.,
2020). We extended the current literature by examining the RewP effect in the domain of RL
processing. Different from our predictions, we found the presence of a small frontal RewP ERP
following reward feedback for preadolescents, which was mostly driven by _poor_ learners, as
described more below. Given that we observed large individual differences in computationally-driven
behavior learning rates in the present study, we compared the RewP effect between "good" and "bad"
learners (i.e., those with lower learning rates), as defined by their computationally-derived learning
rates. This adaptation of WSLS model in current study revealed that poor preadolescent learners
tended to "switch" their choice following losing feedback. These findings are consistent with prior
work with college students(Worthy & Maddox, 2014). WSLS is a rule-based strategy that has been

shown to be commonly used in binary outcome choice tasks (e.g., Otto et al., 2011). According to
(Worthy & Maddox, 2014), college students with higher probabilities of shifting following a loss trial
(i.e., "Lose-Shift" behavior strategy) showed poorer learning performance. Along with prior work
with adults, current findings add converging evidence that combining the WSLS and PRL models
may provide a better account of binary decision-making behavior even in preadolescents compared to
a single PRL model.
442
One key component of PRL processing is attentional shifts towards stimuli associated with
higher probability of getting rewards. Previously learned reward values can have a pronounced effect
on the allocation of selective attention, which has been indexed by an increasing amplitude of the
attention-sensitive, lateralized negative deflection (the N2pc) contralateral to the set-winner in the
cue-pair presentation. The prior ERP studies in adults showed that once a stimulus-reward
association has been learned, it tends to induce a larger attentional shift when presented and also
predicts the likelihood of subsequent choice behavior ((Hickey et al., 2010; Hickey & van Zoest,
2012; San Martín et al., 2016). Such pronounced attentional shift in the context of reward learning, as
indexed by the N2pc has been little investigated in children. Only a handful of ERP studies with
children reported the role of N2pc during visual search (Couperus & Quirk, 2015; Li et al., 2022;
Shimi et al., 2014; Sun et al., 2018; Turoman et al., 2021) and/or in working memory
tasks(Rodríguez-Martínez et al., 2021; Shimi et al., 2015). In addition to the RewP effect, the current
study revealed the modulation of N2pc ERP amplitude indexing early attentional processing towards
the choice behavior (which stimulus would be chosen on that trial). Our prior work in adults (van den
Berg et al., 2019a) revealed the presence of significant N2pc to the to-be-chosen stimulus _and_ to the
more rewarded stimulus in a set, particularly on the later trials of the set, indicating early attention
modulation as a function of reward learning. However, we were not able to replicate such reward-
learning findings in the attention-shifting in the present study with preadolescents. Rather, we found
a modulation of attention bias only towards the chosen side, which was elicited over the whole
course of learning. We speculate that bad learners' limited attention towards rewarded stimuli might
have diluted the presence of small N2pc amplitudes during later vs. early trials of learning in the
good learners. This speculation should be tested in future studies with large samples of both good and
bad child learners.
466
Although we found that, like college students, preadolescents learned the stimulus-reward
outcome association at a group level, our data indicated large individual differences in behavioral
learning rates. With the computationally-driven PRL framework, our behavior data indicated major
differences in strategy between good and bad learners, with good learners using win-stay strategy and
bad learners using more of a lose-switch strategy. PRL theory generally assumes that the most recent
outcomes exert the most influence on the current choice, such that behavior on a given trial, $n$,
depends particularly upon the choice and outcome on the preceding trial, $n - 1$. This relationship has
been formalized as a simple, heuristic learning strategy called Win-Stay-Lose-Shift (WSLS:
(Hernstein et al., 2000).  In our prior work in college students (San Martín et al., 2013), the amplitude
of the feedback-locked P300 over central and parietal cortex, but not the FN (RewP), predicted
behavioral adjustments (i.e., lose-switching strategy) on the next trial. These results are in line with
the context-updating P300 hypothesis (Nieuwenhuis et al., 2005): decisions on each trial are
informed by an internal model of the symbol/probability(win) contingencies, and the P300 amplitude
reflects the extent of the feedback-triggered revision of such a model. Thus, the feedback-locked
P300 distributed over central and parietal cortex may reflect adjustment processes to maximize gain
and/or minimize loss. In line with this, the present study also found a modulation of P300 over
central and parietal cortex as a function of behavioral strategy and feedback type. The feedback-

locked P300 amplitudes were greater on lose-switching trials compared to win-staying trials, and
during loss vs. gain reward feedback.
In contrast to our expectation, we did not find differences in the feedback-locked P300
amplitudes located over central and parietal cortex between good and bad learners even though our
data indicated a modulation of the P300 amplitudes as a function of behavioral strategy and feedback
types more generally. We cannot rule out a possibility that there was not enough power to detect the
interaction effect of P300 amplitudes and subgroup types due to limited number of participants. But
we speculate that better reward learning in children may be more driven by reward sensitivity, rather
than later attention processing related to the updating of salient reward outcomes, as evidenced by
significant differences in the RewP amplitude patterns between the two subtypes of learners.
The main limitation of the current study is a relatively small sample size for examining
individual level differences, which may have resulted in our missing more nuanced distinctions in
our behavioral measures of RL behavior and brain function (e.g., no significant N2pc to the stimulus
with a higher likelihood of reward). In addition, our sample was limited to preadolescent children. As
a result, future work will be needed to replicate the current findings and test their generalizability in
other age ranges and in larger groups reflective of the general US population in terms of race and
ethnicity.
In summary, results of the current study provide novel insights into how a cascade of core
neurocognitive processes underlying RL are associated with preadolescents' behavior learning
strategy: 1) poor preadolescent learners (i.e., with low learning rates) showed greater reward-
feedback sensitivity, which might reflect greater reward-prediction error compared to good learners,
as evidenced by the RewP, 2) there was an early attention modulation towards the stimulus that
would be chosen on a trial, as evidenced by the N2pc, and 3) updating reward values as a function of
behavior strategy, as evidenced by the modulation of P300 amplitudes depending on lose-switch vs.
win-stay strategy in the different learning groups. The alterations in these cognitive processes have
been known to be a key mechanism underlying adolescent-onset depression (Chen et al., 2015).
Adolescents have showed heightened vulnerability for depression with the onset of puberty when
there is a significant reorganization of frontal-striatal reward brain circuitry, as implicated by
differences in RL behavior (Kessler et al., 2001; Walker et al., 2017). The current study shows that
the CPLearn task with the use of EEG measures of brain activity may be a promising tool to further
investigate the relationship between attention and reward sensitivity in preadolescents, which is turn
may advance our understanding of the factors underpinning the risk for depression and other
cognition-development issues in this key demographic group.

**Conflict of Interest**

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest

**Author Contributions**

Chung Y.S. led all aspects of the study, including the development of the task paradigm, the
collection of the data, the analyses of the data, the interpretation of the results, and the writing of the
manuscript. Van den Berg B. and Roberts K.C. contributed to the development of the C-PLearn task,
the preprocessing pipelines, and the data analyses. Woldorff M.G. contributed to the development of
the task paradigm, provided guidance on the EEG data analyses, and contributed to the interpretation
of the results and feedback on the manuscript. Gaffrey MS. contributed to the development of the
task paradigm, guidance on the analyses, interpretation of the results, and feedback on the
manuscript, and provided the research funding to support the study.
531
532

533 **Funding**

534 This study was supported by the National Institute of Mental Health, USA (R01MH110488 to MSG).
535

**Acknowledgments**

We would like to thank to children and their parents who participated and completed this study
despite challenges due to the COVID pandemic (i.e., wearing face masks throughout all sessions).
Also, we would like to specially thank to research staff, Amanda Neal and Summer Lawrence for
their effort and support of data collection. Without their help, we could not have successfully
completed this project.
542

**References**

Amodio, D. M., Bartholow, B. D., & Ito, T. A. (2014). Tracking the dynamics of the social brain: ERP approaches for social cognitive and affective neuroscience. *Soc Cogn Affect Neurosci*, *9*(3), 385-393. https://doi.org/10.1093/scan/nst177

Anderson, B. A. (2017). Going for It: The Economics of Automaticity in Perception and Action. *Curr Dir Psychol Sci*, *26*, 140-145.

Averbeck, B. A.-O., & O'Doherty, J. P. (2022). Reinforcement-learning in fronto-striatal circuits. (1740-634X (Electronic)).

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Package lme4. *Journal of Statistical Software*, *67*, 1-91.

Ben-Shachar, M. (2018). TBT: Reject and Interpolate channels on a trial-by-trial basis. Zenodo. *https://doi.org/10.5281/zenodo.1241518*.

Brown, V. M., Zhu, L., Solway, A., Wang, J. M., McCurry, K. L., King-Casas, B., & Chiu, P. H. (2021). Reinforcement Learning Disruptions in Individuals With Depression and Sensitivity to Symptom Change Following Cognitive Behavioral Therapy. *JAMA Psychiatry*, *78*(10), 1113-1122. https://doi.org/10.1001/jamapsychiatry.2021.1844

Burani, K., Mulligan, E. M., Klawohn, J., Luking, K. R., Nelson, B. D., & Hajcak, G. (2019). Longitudinal increases in reward-related neural activity in early adolescence: Evidence from event-related potentials (ERPs). *Dev Cogn Neurosci*, *36*, 100620. https://doi.org/10.1016/j.dcn.2019.100620

Chang, C. Y., Hsu, S. H., Pion-Tonachini, L., & Jung, T. P. (2018). Evaluation of Artifact Subspace Reconstruction for Automatic EEG Artifact Removal. *Annu Int Conf IEEE Eng Med Biol Soc*, *2018*, 1242-1245. https://doi.org/10.1109/embc.2018.8512547

Chen, C., Takahashi, T., Nakagawa, S., Inoue, T., & Kusumi, I. (2015). Reinforcement learning in depression: A review of computational research. *Neurosci Biobehav Rev*, *55*, 247-267. https://doi.org/10.1016/j.neubiorev.2015.05.005

Christakou, A., Gershman, S. J., Niv, Y., Simmons, A., Brammer, M., & Rubia, K. (2013). Neural and psychological maturation of decision-making in adolescence and young adulthood. *J Cogn Neurosci*, *25*(11), 1807-1823. https://doi.org/10.1162/jocn_a_00447

Cohen, J. R., Asarnow, R. F., Sabb, F. W., Bilder, R. M., Bookheimer, S. Y., Knowlton, B. J., & Poldrack, R. A. (2010). A unique adolescent response to reward prediction errors. *Nat Neurosci*, *13*(6), 669-671. https://doi.org/10.1038/nn.2558

Couperus, J. W., & Quirk, C. (2015). Visual search and the N2pc in children. *Atten Percept Psychophys*, *77*(3), 768-776. https://doi.org/10.3758/s13414-015-0833-5

Davidow, J. Y., Foerde, K., Galván, A., & Shohamy, D. (2016). An Upside to Reward Sensitivity: The Hippocampus Supports Enhanced Reinforcement Learning in Adolescence. (1097-4199 (Electronic)).

Debener, S., Thorne, J. D., Schneider, T. R., & Viola, F. C. (2010). Using ICA for the Analysis of Multi-Channel EEG Data.

Debnath, R. A.-O., Buzzell, G. A., Morales, S. A.-O. X., Bowers, M. E., Leach, S. C., & Fox, N. A. (2020). The Maryland analysis of developmental EEG (MADE) pipeline. (1540-5958 (Electronic)).

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial
EEG dynamics including independent component analysis. *J Neurosci Methods*, *134*(1), 9-21.
https://doi.org/10.1016/j.jneumeth.2003.10.009

DePasque, S., & Galván, A. (2017). Frontostriatal development and probabilistic reinforcement
learning during adolescence. (1095-9564 (Electronic)).

Donaldson, K. R., Ait Oumeziane, B., Hélie, S., & Foti, D. (2016). The temporal dynamics of
reversal learning: P3 amplitude predicts valence-specific behavioral adjustment. *Physiol*
*Behav*, *161*, 24-32. https://doi.org/10.1016/j.physbeh.2016.03.034

Ethridge, P., Kujawa, A., Dirks, M. A., Arfer, K. B., Kessel, E. M., Klein, D. N., & Weinberg, A.
(2017). Neural responses to social and monetary reward in early adolescence and emerging
adulthood. *Psychophysiology*, *54*(12), 1786-1799. https://doi.org/10.1111/psyp.12957

Ferdinand, N. K., Becker, A. M. W., Kray, J., & Gehring, W. J. (2016). Feedback processing in
children and adolescents: Is there a sensitivity for processing rewarding feedback?
*Neuropsychologia*, *82*, 31-38. https://doi.org/10.1016/j.neuropsychologia.2016.01.007

Fischer, A. G., & Ullsperger, M. (2013). Real and fictive outcomes are processed differently but
converge on a common adaptive mechanism. *Neuron*, *79*(6), 1243-1255.
https://doi.org/10.1016/j.neuron.2013.07.006

Halahakoon, D. C., Kieslich, K., O'Driscoll, C., Nair, A., Lewis, G., & Roiser, J. P. (2020). Reward-
Processing Behavior in Depressed Participants Relative to Healthy Volunteers: A Systematic
Review and Meta-analysis. *JAMA Psychiatry*, *77*(12), 1286-1295.
https://doi.org/10.1001/jamapsychiatry.2020.2139

Hauser, T. U., Iannaccone, R., Walitza, S., Brandeis, D., & Brem, S. (2015). Cognitive flexibility in
adolescence: neural and behavioral mechanisms of reward prediction error processing in
adaptive decision making during development. *Neuroimage*, *104*, 347-354.
https://doi.org/10.1016/j.neuroimage.2014.09.018

Hernstein, R., Rachlin, H., & DI, L. (2000). The matching law. *Cambridge, MA: Harvard University*
*Press*.

Hickey, C., Chelazzi, L., & Theeuwes, J. (2010). Reward changes salience in human vision via the
anterior cingulate. *J Neurosci*, *30*(33), 11096-11103. https://doi.org/10.1523/jneurosci.1026-
10.2010

Hickey, C., & van Zoest, W. (2012). Reward creates oculomotor salience. *Curr Biol*, *22*(7), R219-
220. https://doi.org/10.1016/j.cub.2012.02.007

Holroyd, C. B., & Coles, M. G. H. (2002). The neural basis of human error processing: reinforcement
learning, dopamine, and the error-related negativity. *Psychol Rev*, *109*(4), 679-709.
https://doi.org/10.1037/0033-295x.109.4.679

Kappenman, E. S., & Luck, S. J. (2012). The Oxford Handbook of Event-Related Potential
Components. *New York, NY: Oxford University Press*.

Keren, H., O'Callaghan, G., Vidal-Ribas, P., Buzzell, G. A., Brotman, M. A., Leibenluft, E., Pan, P.
M., Meffert, L., Kaiser, A., Wolke, S., Pine, D. S., & Stringaris, A. (2018a). Reward
Processing in Depression: A Conceptual and Meta-Analytic Review Across fMRI and EEG
Studies. *Am J Psychiatry*, *175*(11), 1111-1120.
https://doi.org/10.1176/appi.ajp.2018.17101124

Keren, H., O'Callaghan, G., Vidal-Ribas, P., Buzzell, G. A., Brotman, M. A., Leibenluft, E., Pan, P.
M., Meffert, L., Kaiser, A., Wolke, S., Pine, D. S., & Stringaris, A. (2018b). Reward
Processing in Depression: A Conceptual and Meta-Analytic Review Across fMRI and EEG
Studies. (1535-7228 (Electronic)).

Kessel, E. M., Dougherty, L. R., Kujawa, A., Hajcak, G., Carlson, G. A., & Klein, D. N. (2016).
Longitudinal Associations Between Preschool Disruptive Mood Dysregulation Disorder
Symptoms and Neural Reactivity to Monetary Reward During Preadolescence. *J Child*
*Adolesc Psychopharmacol*, *26*(2), 131-137. https://doi.org/10.1089/cap.2015.0071

Kessel, E. M., Nelson, B. D., Finsaas, M., Kujawa, A., Meyer, A., Bromet, E., Carlson, G. A.,
Hajcak, G., Kotov, R., & Klein, D. N. (2019). Parenting style moderates the effects of
exposure to natural disaster-related stress on the neural development of reactivity to threat and
reward in children. *Dev Psychopathol*, *31*(4), 1589-1598.
https://doi.org/10.1017/s0954579418001347

Kessler, R. C., Avenevoli, S., & Ries Merikangas, K. (2001). Mood disorders in children and
adolescents: an epidemiologic perspective. *Biol Psychiatry*, *49*(12), 1002-1014.
https://doi.org/10.1016/s0006-3223(01)01129-5

Krigolson, O. E. (2018). Event-related brain potentials and the study of reward processing:
Methodological considerations. (1872-7697 (Electronic)).

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). ImerTest Package: tests in linear
mixed effects models. *Journal of Statistical Software*, *82*.

Lee, T. W., Girolami, M., & Sejnowski, T. J. (1999). Independent component analysis using an
extended infomax algorithm for mixed subgaussian and supergaussian sources. *Neural*
*Comput*, *11*(2), 417-441. https://doi.org/10.1162/089976699300016719

Li, B., Guo, J., Zhao, C., Luo, X., Kong, Y., Chen, Y., Liu, H., Sun, L., & Song, Y. (2022). Lack of
an association between anticipatory alpha oscillations and attentional selection in children
with attention-deficit/hyperactivity disorder. *Clin Neurophysiol*, *138*, 25-37.
https://doi.org/10.1016/j.clinph.2022.02.026

Lopez-Calderon, J., & Luck, S. J. (2014). ERPLAB: an open-source toolbox for the analysis of
event-related potentials. *Front Hum Neurosci*, *8*, 213.
https://doi.org/10.3389/fnhum.2014.00213

Luck, S., & Kappenman, E. (2012). ERP components and selective attention In:

The Oxford Handbook of Event-Related Potential Components. *New York: Oxford University Press.*

Lukie, C. N., Montazer-Hojat, S., & Holroyd, C. B. (2014). Developmental changes in the reward
positivity: an electrophysiological trajectory of reward processing. *Dev Cogn Neurosci*, *9*,
191-199. https://doi.org/10.1016/j.dcn.2014.04.003

Master, S. L., Eckstein, M. K., Gotlieb, N., Dahl, R., Wilbrecht, L., & Collins, A. G. E. (2020).
Distentangling the systems contributing to changes in learning during adolescence. (1878-
9307 (Electronic)).

Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: An open-source, graphical experiment
builder for the social sciences. *Behav Res*, *44*(314-324).
https://doi.org/https://doi.org/10.3758/s13428-011-0168-7

Mitra, P., & Bokil, H. (2007). Observed Brain Dynamics. *Oxford University Press*, Chapter 7.3.4.

Morris, B. H., Bylsma Lm Fau - Yaroslavsky, I., Yaroslavsky I Fau - Kovacs, M., Kovacs M Fau - Rottenberg, J., & Rottenberg, J. (2015). Reward learning in pediatric depression and anxiety: preliminary findings in a high-risk sample. (1520-6394 (Electronic)).

Mullen, T. R., Kothe, C. A., Chi, Y. M., Ojeda, A., Kerth, T., Makeig, S., Jung, T. P., & Cauwenberghs, G. (2015). Real-Time Neuroimaging and Cognitive Monitoring Using Wearable Dry EEG. *IEEE Trans Biomed Eng*, *62*(11), 2553-2567. https://doi.org/10.1109/tbme.2015.2481482

Nasser, H. M., Calu, D. J., Schoenbaum, G., & Sharpe, M. J. (2017). The Dopamine Prediction Error: Contributions to Associative Models of Reward Learning. (1664-1078 (Print)).

Nieuwenhuis, S., Aston-Jones, G., & Cohen, J. D. (2005). Decision making, the P3, and the locus coeruleus-norepinephrine system. *Psychol Bull*, *131*(4), 510-532. https://doi.org/10.1037/0033-2909.131.4.510

Nussenbaum, K., & Hartley, C. A. (2019a). Reinforcement learning across development: What insights can we draw from a decade of research? (1878-9307 (Electronic)).

Nussenbaum, K., & Hartley, C. A. (2019b). Reinforcement learning across development: What insights can we draw from a decade of research? *Dev Cogn Neurosci*, *40*, 100733. https://doi.org/10.1016/j.dcn.2019.100733

O'Callaghan, G., & Stringaris, A. (2019). Reward Processing in Adolescent Depression Across Neuroimaging Modalities. *Z Kinder Jugendpsychiatr Psychother*, *47*(6), 535-541. https://doi.org/10.1024/1422-4917/a000663

Palminteri, S., Kilford, E. A.-O., Coricelli, G., & Blakemore, S. J. (2016). The Computational Development of Reinforcement Learning during Adolescence. (1553-7358 (Electronic)).

Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. . (1989). Spherical splines for scalp potential and current density mapping. *Electroencephalography and clinical neurophysiology*, *72(2)*, 184-187.

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated electroencephalographic independent component classifier, dataset, and website. *Neuroimage*, *198*, 181-197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Raab, H. A., & Hartley, C. A. (2018). Chapter 13 - The Development of Goal-Directed Decision-Making. In R. Morris, A. Bornstein, & A. Shenhav (Eds.), *Goal-Directed Decision Making* (pp. 279-308). Academic Press. https://doi.org/https://doi.org/10.1016/B978-0-12-812098-9.00013-9

Rodríguez-Martínez, E. I., Arjona Valladares, A., Gómez-González, J., Diaz-Sanchez, J. A., & Gómez, C. M. (2021). Neurophysiological differences between ADHD and control children and adolescents during the recognition phase of a working memory task. *Neurosci Res*, *164*, 46-54. https://doi.org/10.1016/j.neures.2020.03.011

San Martín, R. (2012). Event-related potential studies of outcome processing and feedback-guided learning. *Front Hum Neurosci*, *6*, 304. https://doi.org/10.3389/fnhum.2012.00304

San Martín, R., Appelbaum, L. G., Huettel, S. A., & Woldorff, M. G. (2016). Cortical Brain Activity Reflecting Attentional Biasing Toward Reward-Predicting Cues Covaries with Economic Decision-Making Performance. *Cereb Cortex*, *26*(1), 1-11. https://doi.org/10.1093/cercor/bhu160

San Martín, R., Appelbaum, L. G., Pearson, J. M., Huettel, S. A., & Woldorff, M. G. (2013). Rapid brain responses independently predict gain maximization and loss minimization during economic decision making. *J Neurosci*, *33*(16), 7011-7019. https://doi.org/10.1523/jneurosci.4242-12.2013

Saulnier, K. G., Marr, N. S., van Geen, C., Babinski, D. E., & Mukherjee, D. (2023). Reinforcement-based responsiveness, depression, and anhedonia: A multi-method investigation of intergenerational risk. (1879-1379 (Electronic)).

Schultz, W. (2016). Dopamine reward prediction error coding. (1958-5969 (Electronic)).

Shimi, A., Kuo, B. C., Astle, D. E., Nobre, A. C., & Scerif, G. (2014). Age group and individual differences in attentional orienting dissociate neural mechanisms of encoding and maintenance in visual STM. *J Cogn Neurosci*, *26*(4), 864-877. https://doi.org/10.1162/jocn_a_00526

Shimi, A., Nobre, A. C., & Scerif, G. (2015). ERP markers of target selection discriminate children with high vs. low working memory capacity. *Front Syst Neurosci*, *9*, 153. https://doi.org/10.3389/fnsys.2015.00153

Sun, M., Wang, E., Huang, J., Zhao, C., Guo, J., Li, D., Sun, L., Du, B., Ding, Y., & Song, Y. (2018). Attentional selection and suppression in children and adults. *Dev Sci*, *21*(6), e12684. https://doi.org/10.1111/desc.12684

Sutton, R. S., & Barto, A. G. (2018). Reinforcement Learning: An Introduction. *MIT Press, Cambridge, MA*, *Second Edition*.

Turoman, N., Tivadar, R. I., Retsa, C., Maillard, A. M., Scerif, G., & Matusz, P. J. (2021). The development of attentional control mechanisms in multisensory environments. *Dev Cogn Neurosci*, *48*, 100930. https://doi.org/10.1016/j.dcn.2021.100930

van den Berg, B., Geib, B. R., San Martín, R., & Woldorff, M. G. (2019a). A key role for stimulus-specific updating of the sensory cortices in the learning of stimulus-reward associations. (1749-5024 (Electronic)).

van den Berg, B., Geib, B. R., San Martín, R., & Woldorff, M. G. (2019b). A key role for stimulus-specific updating of the sensory cortices in the learning of stimulus-reward associations. *Soc Cogn Affect Neurosci*, *14*(2), 173-187. https://doi.org/10.1093/scan/nsy116

von Borries, A. K., Verkes, R. J., Bulten, B. H., Cools, R., & de Bruijn, E. R. (2013). Feedback-related negativity codes outcome valence, but not outcome expectancy, during reversal learning. *Cogn Affect Behav Neurosci*, *13*(4), 737-746. https://doi.org/10.3758/s13415-013-0150-1

Walker, D. M., Bell, M. R., Flores, C., Gulley, J. M., Willing, J., & Paul, M. J. (2017). Adolescence and Reward: Making Sense of Neural and Behavioral Changes Amid the Chaos. *J Neurosci*, *37*(45), 10855-10866. https://doi.org/10.1523/jneurosci.1834-17.2017

Worthy, D. A., & Maddox, W. T. (2014). A Comparison Model of Reinforcement-Learning and Win-Stay-Lose-Shift Decision-Making Processes: A Tribute to W.K. Estes. *J Math Psychol*, *59*, 41-49. https://doi.org/10.1016/j.jmp.2013.10.001

Wu, Y., & Zhou, X. (2009). The P300 and reward valence, magnitude, and expectancy in outcome evaluation. *Brain Res*, *1286*, 114-122. https://doi.org/10.1016/j.brainres.2009.06.032

Figure 1. Individual differences in behavioral probabilistic reward learning

*Note.* Behavioral reward learning: During 18 blocks, the proportion of choosing set-winner for all participants ranged from 0.40 to 0.73. The individual learning curves revealed very large individual difference in reward learning, as the final proportion of choosing set winners in a particular block ranged from 0.26 to 1.0. Due to this large individual difference observed, we divided participants into good vs. bad learners with a median split method.

Figure 2. Behavioral strategies during the Child-version of Probabilistic Reward Learning (C-PLearn)

*Note.* Bad learners with low learning rates are more likely to change their choice on the next trial following loss feedback (i.e., Lose-Shift strategy). On the other hand, good learners with high learning rates are more like to make the same choice on the next trial following gain feedback (i.e., Win-Stay strategy). The error bars indicate standard error. **$p$ = .007, ***$p$ <.001.

Figure 3.1. The Frontal reward-related potential (gain vs. loss feedback) in preadolescents (n=30):

*Note.* (a) Grand average feedback-cue evoked responses for Gain and Loss reward feedback from the fronto-central ROI (E10, E11, E16) (see red dots in the topography for the location of the ROI).
(b) The reward-related potential difference waveform during Gain minus Loss reward feedback. The reward-related potential was measured from 275 to 375 ms in a fronto-central ROI.

Figure 3.2. The reward-related positivity between good and bad learners

*Note.* The comparison of the reward-related potential ERP marker measured from 275 to 375ms in the fronto-central ROI between good and bad learners. Dark red colored electrode sites in the frontal ROI shows significantly greater reward-related positivity amplitudes for bad learners (i.e., those with low learning rates) compared to good learners (i.e., those with high learning rates).

Figure 4.1. Topography of P300 during gain vs. loss feedback as a function of behavioral strategy

Central-parietal
(top view)

*Note.* P300 amplitude differences in central and parietal ROIs when participants switched their responses on the next trial after gain feedback vs. after loss feedback. The red dot in the topography at the bottom shows the location of the central-parietal ROI.
The top row: warm colored electrode sites indicate bigger P300 amplitudes on stay trials for good learners with high learning rates.
The bottom row: warm colored electrode sites indicate bigger P300 on stay trials for bad learners with low learning rates.

Figure 4.2 P300 amplitudes in central, parietal ROIs as a function of feedback type and behavioral strategy

*Note.* The bar indicates standard error; **$p = 0.01$, ***$p = .005$.
(a) Using P300 amplitudes in the central-parietal ROIs between 400 and 600 ms for good and bad learners, we found that bad learners (i.e., those with low learning rates), relative to good learners, showed higher P300 amplitude differences in central, parietal ROIs when they switched their responses on the next trial after gain feedback vs. after loss feedback.
(b) For all children, we found greater P300 amplitudes in the central-parietal ROI when participants switched their choice on the next trial after loss feedback (i.e., lose-switch strategy) compared to when they chose the same choice after gain feedback (i.e., win-stay strategy).
(c) The P300 amplitudes in the same central-parietal ROIs were higher when participants received loss feedback compared to gain feedback.

Figure 5. The cue-evoked attentional bias of N2pc towards the later chosen side

Note. To increase signal-to-noise ratio, we collapsed over whether the chosen image was that of a face or a house and analyzed the effect of choice behavior on the N2pc over the course of the 18-trial set. The cue-evoked N2pc amplitudes measured after onset of the cue image pair, from corresponding left and right occipital ROIs was found from 175 to 225ms (see red dots in the topography for the location of the ROI electrodes).