CaBP1 and 2 enable sustained Ca$_V$1.3 calcium currents and synaptic transmission in inner hair cells

[1,2]David Oestreicher, [3,4]Shashank Chepurwar, [5,6]Kathrin Kusch, [6,7]Vladan Rankovic, [6,#]Sangyong Jung, [3,4]Nicola Strenzke, [1,2,4,8,*]Tina Pangrsic

[1] Experimental Otology Group, InnerEarLab, Department of Otolaryngology, University Medical Center Göttingen, 37075 Göttingen, Germany.
[2] Auditory Neuroscience Group, Max Planck Institute for Multidisciplinary Sciences, 37075 Göttingen, Germany.
[3] Auditory Systems Physiology Group, Institute for Auditory Neuroscience, InnerEarLab, University Medical Center Göttingen, 37075 Göttingen, Germany
[4] Collaborative Research Center 889, University of Göttingen, 37075 Göttingen, Germany
[5] Functional Auditory Genomics, Institute for Auditory Neuroscience, University Medical Center Göttingen, 37075 Göttingen, Germany
[6] Institute for Auditory Neuroscience and InnerEarLab, University Medical Center Göttingen, 37075 Göttingen, Germany.
[7] Restorative Cochlear Genomics Group, Auditory Neuroscience and Optogenetics Laboratory, German Primate Center, 37075 Göttingen, Germany.
[8] Multiscale Bioimaging Cluster of Excellence (MBExC), University of Göttingen, 37075 Göttingen, Germany.
[#] Current address: Department of Medical Science, College of Medicine, CHA University, Seongnam-si, Gyeonggi-do, 13488, South Korea

*Corresponding author: Tina Pangrsic

Email: tpangrs@gwdg.de

Author Contributions: TP and DO designed the study. DO performed *in vitro* cell electrophysiology, immunohistochemistry, and ABR/DPOAE recordings from young and injected animals. SC performed the *in vivo* single unit recordings and the ABR/DPOAE recordings in older animals under the supervision of NS. KK and VR generated the viral vectors. SY performed the initial injection experiments. DO, TP, SC and NS analysed the data. TP and DO wrote the manuscript, with contribution from NS. All authors revised the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Keywords: DFNB93, calcium channel, hearing impairment, ribbon synapse, CACNA1D

Abstract

To encode continuous sound stimuli, the inner hair cell (IHC) ribbon synapses utilize calcium-binding proteins (CaBPs), which reduce the inactivation of their Ca$_V$1.3 calcium channels. Mutations in the *CABP2* gene underlie non-syndromic autosomal recessive hearing loss DFNB93. Besides CaBP2, the structurally related CaBP1 is highly abundant in the IHCs. Here, we investigated how the two CaBPs cooperatively regulate IHC synaptic function. In *Cabp1/2* double-knockout mice, we find strongly enhanced Ca$_V$1.3 inactivation, slowed recovery from inactivation and impaired sustained exocytosis. Already mild IHC activation further reduces the availability of channels to trigger synaptic transmission and may effectively silence synapses. Spontaneous and sound-evoked responses of spiral ganglion neurons *in vivo* are strikingly reduced and strongly depend on stimulation

rates. Transgenic expression of CaBP2 leads to substantial recovery of IHC synaptic function and hearing sensitivity. We conclude that CaBP1 and 2 act together to suppress voltage- and calcium-dependent inactivation of IHC $Ca_V1.3$ channels in order to support sufficient rate of exocytosis and enable fast, temporally precise and indefatigable sound encoding.

Introduction

Calcium influx through voltage-gated calcium channels (Ca_Vs) is a pivotal initiator of neurotransmission. These channels typically undergo considerable Ca^{2+}- and voltage-dependent inactivation (CDI and VDI), which limits the amount of incoming Ca^{2+} ions and thus possible excitotoxicity. In sensory receptor cells such as sensory inner hair cells and photoreceptors, however, the extent of inactivation is significantly reduced to allow encoding of graded and sustained membrane potential fluctuations in response to ongoing sound or visual (t.i. "darkness") stimuli [reviewed in [1]]. The attenuation of channel inactivation may be largely attributed to the action of calcium-binding proteins, CaBPs [2], which primarily abate the calmodulin (CaM)-mediated CDI of Ca_V channels [2–7]. At the inner hair cell (IHC) ribbon synapses, the Ca^{2+} current is carried through the L-type $Ca_V1.3$ channels [8,9]. Their biophysical properties may be regulated by several CaBP family members [10,11], but most prominently by CaBP2 [12]. Mutations in the *CaBP2* gene cause moderate to severe autosomal recessive hearing impairment DFNB93 [see e.g. [12–16]]. Interestingly, the loss of CaBP2 seems to significantly affect VDI, but not CDI of the IHC Ca^{2+} currents [12]. Furthermore, isolated disruption of CaBP1 or CaBP4 in mice leads to at most modest enhancement of IHC $Ca_V1.3$ inactivation [10,11]. Moreover, the automodulatory C-terminal domain of the long $Ca_V1.3$ isoform, abundant at the IHC ribbon synapses [17,18], does not seem critical for CDI suppression either, as its disruption does not lead to hearing impairment [18]. The question of which mechanisms are primarily responsible for suppressing IHC $Ca_V1.3$ CDI thus remains unresolved. Since CaBP2, followed by the structurally similar CaBP1, are the most abundant members of the CaBP family in the cochlea [19], we set out to investigate the phenotype of *Cabp1/2* double-knockout (DKO) animals at the systems and cellular levels. Severe hearing impairment that we observed in these animals reveals that CaBP1 and 2 act together to attenuate the inactivation of the IHC $Ca_V1.3$ channels. Furthermore, we demonstrate that the combined absence of CaBP1 and 2 almost abolishes sustained, and, upon increased hair cell synaptic activity, the fast component of exocytosis. Such phenotype was observed neither in *Cabp1* nor in *Cabp2* single KO animals [11,12]. To obtain a more comprehensive understanding of the roles of the two calcium-binding proteins, we attempted to rescue the auditory function of the CaBP1/2-depleted auditory system by viral delivery of a *Cabp2* coding sequence in the inner ear. These experiments revealed a substantial recovery of hearing, which may be related to the successful restoration of Ca^{2+} currents and IHC exocytosis upon transgenic expression of CaBP2. Our findings indicate that the functions of CaBP1 and 2 partially overlap at the IHC ribbon synapse. Together, they provide sufficient attenuation of channel inactivation to sustain Ca^{2+} currents and exocytosis required for temporally precise, indefatigable neurotransmission at this synapse.

Results

Pronounced VDI and CDI of $Ca_V1.3$ channels in CaBP1/2-deficient IHCs. CaBP1 and 2 show high expression levels in the organ of Corti [19], where they may modulate voltage-gated $Ca_V1.3$ channels at the IHC ribbon synapses (Fig. 1A). Deletion of CaBP2 results in an early-onset, progressive hearing impairment in the KO mouse model and moderate to severe hearing impairment, DFNB93, in affected human patients [12–14]. Loss of CaBP1 in mice, on the other hand, leaves hearing unperturbed in the first few months after birth [11]. Since these proteins show a high degree of homology [2], it is plausible that they possess similar features and are partially redundant, at least initially. Therefore, we were interested in understanding their cumulative impact on the function of hair cells and hearing. For this, we analyzed the phenotype of the *Cabp1/2*-DKO animals and tested

the extent of functional rescue following the transgenic expression of CaBP2 (Fig. S1A) in the ears of the DKO mice.

To investigate possible synergistic functions of the two major IHC CaBPs, we first measured Ca^{2+} currents in the IHCs of 3-4-week-old *Cabp1/2*-DKO and WT animals in the presence of 1.3 mM extracellular $[Ca^{2+}]$. Employing 500 ms-long depolarization steps to maximal Ca^{2+}-current potential (-25 to -15 mV) revealed pronounced inactivation of $Ca_V1.3$ channels in the DKO IHCs (Fig. 1B). This was significantly stronger when compared to the inactivation observed in the IHCs of *Cabp1* or *Cabp2* single KOs in similar conditions (Fig. 1C) [11,12,20], and even when comparing to data from 5-week-old *Cabp2* KO mice obtained at physiological temperature (Fig. S1D), further substantiating the evidence of severity and early onset of the DKO phenotype.

The "non-inactivating" nature of the IHC $Ca_V1.3$ channels was partially restored in the *Cabp1/2*-DKOs by transgenic expression of WT CaBP2 (Student's *t*-test: $p < 0.0001$; Fig. 1B-C; see also Fig. S1). The lack of complete recovery suggests either an inadequate expression of transgenic material or irreversible adaptive changes in response to the lack of major CaBPs that may occur prior to gene delivery. Similarly, Ca^{2+} currents were greatly reduced upon application of short train stimuli and rescued efficiently by the introduction of the transgenic CaBP2 (Fig. 1D). However, it has to be noted that the cells in which train stimuli were applied showed a considerably lower calcium current inactivation upon a 500-ms step depolarization than the average (0.36 (*n*=5) vs 0.47 (*n*=21)). Short stimuli at different potentials revealed maximal Ca^{2+}-current amplitudes in the IHCs of DKO mice that were comparable to controls (Fig. 1E). We observed a slight shift in the voltage-dependence of $Ca_V1.3$ activation towards more hyperpolarized values (Fig. 1F; potential of half-activation – $V_{0.5}$ of -33.6 ± 0.5 vs -30.9 ± 0.5 mV in *Cabp1/2* DKO and WT IHCs, respectively; $p < 0.0001$, Student's *t*-test), and an increased slope of the Boltzmann fit function (7.6 ± 0.1 vs 7.1 ± 0.1, $p < 0.0001$, Student's *t*-test). This finding is somewhat surprising as the opposite would be expected based on data from heterologous expression systems [4,12,21]. At present we can not exclude potential changes in the relative amounts of different $Ca_V1.3$ splice variants, auxiliary subunits or other interaction partners that may in the absence of CaBPs lead to these changes in the IHCs.

To further investigate the nature of the enhanced $Ca_V1.3$ inactivation, we recorded $Ca_V1.3$ currents in the presence of 2 mM Ca^{2+} or Ba^{2+} in the extracellular solution (Fig. 2), and analyzed the two types of inactivation as described previously [22]. The lack of CaBP1 and 2 significantly enhanced both, VDI and CDI of IHC $Ca_V1.3$ channels, which were approx. two- and six-times larger than those in WT IHCs (Fig. 2B; VDI: 0.30 ± 0.02 vs 0.13 ± 0.02 and CDI: 0.65 ± 0.01 vs 0.11 ± 0.02 in *Cabp1/2* DKO and WT IHCs, respectively; $p < 0.0001$, Student's *t*-test).

Impaired synaptic transmission in the IHCs lacking CaBP1 and 2. Since Ca^{2+} influx through $Ca_V1.3$ channels triggers IHC neurotransmission we next investigated how the prominent inactivation affects exocytosis in the CaBP1/2-deficient IHCs. Exocytosis was probed by recordings of membrane capacitance (Fig. 3). In response to short stimuli [primarily recruiting the readily-releasable pool (RRP) of vesicles], increases in capacitance were comparable, while long stimuli (reporting sustained exocytosis) evoked significantly smaller responses in the CaBP1/2-deficient IHCs (Fig. 3A-B). These results are in line with the observation of comparable peak Ca^{2+}-current amplitudes but reduced calcium charge transfer (Q_{Ca}) for stimulus durations above 50 ms. However, calculating exocytosis efficiency revealed that the reduction in sustained exocytosis cannot completely be explained by the diminished Ca^{2+} influx (Fig. 3C). This suggests an additional impairment in exo- or endocytic processes, the underlying mechanism of which is currently unknown. Transgenic expression of CaBP2 largely restored the efficiency of exocytosis, indicating that the impairment as observed in the IHCs of DKOs is related to the absence of the CaBPs. Such exocytosis defect was not observed in *Cabp2*-KOs, suggesting that when IHCs are held at hyperpolarized potential at RT, CaBP1 provides sufficient compensation to obscure the synaptic phenotype[12,20].

Next, we have reasoned that holding a cell at a relatively hyperpolarized potential of -85 mV as typically utilized in patch-clamp recordings may not reveal the synaptic phenotype in its severity.

We thus performed additional sets of experiments (Fig. 3D-E), in which the cells were held at either -65 or -55 mV, closer to the suggested resting potential of IHCs [23-25]. Raising the holding voltage to -55 mV resulted in a significant reduction in the maximal Ca^{2+} currents, which was more pronounced in the CaBP1/2-deficient IHCs as compared to controls (60 ± 4 % vs 25 ± 4 % reduction in DKO and WT IHCs, respectively; $p < 0.005$, Student's t-test). Consequently, further discrepancy between the Q_{Ca} in IHCs of WT and DKO animals was observed ($p < 0.001$), which in DKO animals resulted in a complete loss of sustained exocytosis (Fig. 3D; $p < 0.005$). The latter was well restored upon AAV-mediated transgenic expression of CaBP2 ($p < 0.05$), which also reduced the amount of calcium channel inactivation (38 ± 8 % reduction in the maximal Ca^{2+} current as well as a partial recovery of Q_{Ca} as compared to *Cabp1/2*-DKO IHCs; $p < 0.05$). Rapid voltage oscillations during capacitance recordings may additionally activate calcium channels prior to the depolarizing step. We thus reduced the sine wave amplitude, which resulted in an intermediate phenotype, indicating pronounced activity-dependent deterioration of synaptic function in the absence of CaBPs (Fig. 3E). We propose that *in vivo* in the absence of CaBP1 and 2 a large fraction of IHC $Ca_V1.3$ channels are in a state of prolonged inactivation. This does not support sufficient RRP exocytosis (Fig. 3D) to sustain sensory coding during ongoing stimulation of IHCs.

Lastly, we asked whether the deletion of CaBP1 and 2 affects the speed of recovery from $Ca_V1.3$ inactivation (Fig. 4). We applied an initial 5-ms long pulse followed by a conditioning stimulus and a series of 5-ms long pulses to test the kinetics of recovery of the Ca^{2+}-current amplitude. Upon 500-ms depolarization step (Fig. 4B), the speed of recovery from inactivation was fast both in DKO as well as WT IHCs where a much smaller fraction of channels underwent inactivation (time constants of 0.62 and 0.75 s in DKO and WT IHCs, respectively; Table 1, left). As expected, transgenic expression of CaBP2 reduced the initial inactivation (Ca^{2+}-current amplitude 10 ms after the conditioning pulse, expressed as the fraction of initial current – I_{10ms_post}/I_{pre}, WT: 0.73 ± 0.01; DKO: 0.36 ± 0.03; DKO+PHP.eB-*Cabp2*: 0.64 ± 0.02; $p < 0.001$ for DKO vs DKO+PHP.eB-*Cabp2*). However, the recovery from the step-evoked inactivation was slowest in the *Cabp2*-injected DKO IHCs. We hypothesize that this may be related to the speed at which the intracellular $[Ca^{2+}]$ after depolarization is restored to the basal level allowing $Ca_V1.3$ recovery from CDI (see discussion). A 2-min long sine wave (± 15 mV) from -55 mV to mimic a milder but prolonged hair cell stimulation (Fig. 4C) elicited smaller inactivation of $Ca_V1.3$ channels as compared to a 500-ms depolarization step (I_{10ms_post}/I_{pre} of 0.73 ± 0.01 in WT; 0.58 ± 0.03 in DKO; 0.68 ± 0.03 in DKO+PHP.eB-*Cabp2*; $p < 0.05$ for DKO vs DKO+PHP.eB-*Cabp2*). In these conditions, however, the recovery from inactivation was slowest in the DKO IHCs (Table 1, right). This suggests that in the absence of CaBP1 and 2 even mild but prolonged stimulation of IHCs *in vivo* may drive a significant fraction of $Ca_V1.3$ channels into "deep-state" of inactivation with very slow recovery, as was described for Na_V, K_V and Ca_V channels [26-29], not allowing efficient sound encoding.

Altered synaptic morphology in *Cabp1/2*-DKO mice. Immunolabeling of the organs of Corti indicated normal presynaptic $Ca_V1.3$ clustering with no apparent change in size or shape of the clusters (Fig. S2A), and comparable abundance of ribbon synapses in the apical IHCs of WT and DKO animals (12.9 ± 0.2 vs 12.8 ± 0.8; respectively) at 3 weeks of age (Fig. S2B-C'). However, the intensities of pre- (RibeyeA/CtBP2) and postsynaptic (Homer1) immunofluorescent puncta were increased in the CaBP1/2-deficient IHCs as compared to age-matched controls, processed in parallel (Fig. S2C''). A similar alteration in the synaptic size was observed in some other synaptic mutants as well as upon noise trauma [30,31], but not in *Cabp2* single KOs [12], which show significantly less impaired presynaptic Ca^{2+} signalling. This data is in line with compensatory postsynaptic changes in conditions of no or very little postsynaptic activity, possibly governed through adaptive changes in the postsynaptic elements.

As earlier studies of $Ca_V1.3$ channelopathies demonstrated a lag or stop in IHC development [30,32,33] we next examined the expression of small- (SK) and large-conductance Ca^{2+}-activated potassium channels (BK) in the IHCs at different ages. As expected for normal maturation, the SK2 puncta were mostly absent from DKO IHCs by 3 weeks of age and the BK puncta underwent a typical transition from a spotty appearance throughout the cell membrane to the patches at the IHC necks

(Fig. S2D-G). However, by 3 weeks and more obviously later on, the BK channel clusters appeared smaller in the DKO IHCs as compared to controls, which could be partially reversed by virally-mediated expression of CaBP2 (Fig. S2F-G). We hypothesize that CaBPs may directly (e.g. as transcription factors) regulate BK expression, a notion that needs to be tested in the future. Alternatively, the effects may be more indirect, as they are common in mutants with impaired intracellular Ca^{2+} signaling [30,32,33].

Severe hearing impairment in mice deficient in CaBP1 and CaBP2. Hearing was assessed by measuring auditory brainstem responses (ABRs; Fig. 5, Fig. S3). Already in young (3-4-week-old) *Cabp1/2*-DKO animals, ABR thresholds were highly elevated, with tone bursts up to a maximum intensity of 90 dB barely eliciting any ABR response; click responses were better preserved but greatly reduced in amplitude (Fig. S3A-B). At 7-13 weeks of age, further worsening of the click thresholds was observed. Upon 80-dB click stimulation no sizable ABR wave I was detected in average traces of DKO animals at either age tested (Fig. S3B), suggesting reduced and/or desynchronized spiking of spiral ganglion neurons (SGNs). Like in other mouse mutants with defects in the IHC synapse [34,35], the later ABR peaks were better preserved, indicating partially preserved activation of neurons of the auditory brainstem. To assess cochlear amplification by the outer hair cells (OHCs), we recorded distortion product otoacoustic emissions (DPOAEs). While OAEs were so far only detectable in some of the younger DFNB93 patients [12,36], mouse work corroborates the notion of at least initially intact cochlear amplification in the absence of CaBP2 [11,12,20]. Our results suggest that cochlear amplification is largely preserved also in DKO animals (Fig. S3C-F). In about half of the DKO animals in both age groups, the DPOAEs were normal. In the other half, they were reduced in amplitude but still detectable (Fig. S3E-F), which may suggest partially hampered cochlear amplification in a subset of DKO animals.

Transgenic expression of CaBP2 led to significant improvement of the IHC $Ca_V1.3$ function and synaptic exocytosis (Figs. 1-4), we thus next asked, how this may influence the hearing of the animals. As previously observed [20], unilateral injection of the PHP.eB-*Cabp2-eGFP* resulted in strong eGFP immunofluorescence signal in the vast majority of hair cells and elsewhere (Fig. S4A-B). Transgenic expression of CaBP2 in DKO animals led to significantly improved hearing with lower ABR thresholds, increased click-evoked ABR amplitudes, and shortened wave I latency (Fig. 5, Fig. S4C-D).

Finally, we studied sound encoding by inspecting sound-induced spiking responses of SGNs (Fig. 6). In WT mice, we recorded responses in 39 neurons, which were classified as SGNs based on their location, response patterns, spike waveforms and recording stability. Their spiking characteristics resembled typical WT data [12,37–40]. In response to 50- or 500-ms tone burst at the characteristic frequency 30 dB above threshold (i.e. where the sigmoidal rate-intensity functions have saturated spike rates) and at a stimulus repetition rate of 5 Hz, they showed the SGN-typical very high onset firing rates, gradually adapting to reach steady-state rates between 100 and 400 Hz (Fig. 6A-B). In contrast, only one out of the 20 neurons we recorded and analyzed from DKO animals showed a normal response in our automated tuning curve algorithm with a threshold of 35 dB at 7.8 kHz, while all others displayed thresholds above 80 dB and extremely low spike rates (Fig. 6A-B). Since in the majority of neurons frequency tuning was very broad, we used subjectively chosen best frequency tone burst or white noise burst stimulation of at least 80 dB for subsequent characterization. 16 out of 19 DKO units responded with a relatively high onset rate but strong adaptation to a minimal sustained spiking (Fig. 6A-B), which prevented unit classification based on the response patterns. Judging from other criteria described above, we assume that approximately 13 of these represent SGNs and 3 represent bushy cells or multipolar cells of the anteroventral cochlear nucleus. Three additional units (two resembling a pauser/buildup- and one displaying a clear chopper neuron-type response) were excluded from further analysis. Overall, in DKO neurons, peak and adapted rates were greatly reduced as compared to WT controls (Fig. 6C), which came along with an increased latency (12.6 ± 2.8 ms vs WT 4.8 ± 0.2 ms; $p < 0.001$, Wilcoxon rank-sum test)

and increased jitter (18.2 ± 5.0 vs WT 1.5 ± 0.4; p < 0.00001, Wilcoxon rank-sum test) of the first spike that followed the spike onset.

DKO responses greatly improved when we lowered the stimulus repetition rate to 0.5 Hz, and plateaued at 0.2 Hz (Fig. 6C-D). We hypothesize that in DKO SGNs, recovery of $Ca_V1.3$ channels after a silent interval in the range of 1-2 seconds is the limiting factor for determining the maximal SGN spiking rate at the sound onset. This may be further contributed by potentially slowed re-supply of synaptic vesicles, as suggested by reduced efficiency of sustained exocytosis in the CaBP1/2-deficient IHCs (see Fig. 3). In contrast, in WT SGNs, the maximal spike rates are mostly recovered after a silent interval of 150 ms, which may be mostly limited by vesicle replenishment at the IHC ribbon synapse [35,41]. During a 5-Hz sound stimulation, the strength of adaptation was greatly enhanced (peak to adapted ratio 12.5 ± 3.0 vs WT 3.4 ± 0.1; p < 0.00001, Wilcoxon rank-sum test) whereas the time constant of adaptation was unchanged (7.9 ± 2.0 ms in DKO compared to 7.0 ± 0.4 ms in WT (p = 0.5, Wilcoxon rank-sum test). However, for the longer 500-ms tone bursts used in the 0.5 Hz stimulation dataset, it becomes obvious that despite the strong adaptation, DKO neurons are still able to sustain spiking at a low rate throughout the stimulation (Fig. 6B). DKOs also showed reduced spontaneous spiking activity (Fig. 6E), which supports the hypothesis that already in the absence of sound stimulation over tens of seconds a considerable fraction of calcium channels may be in steady-state inactivation.

The reduction in spike rates and their strong dependence on the strength of stimulation in the preceding seconds introduces experimental bias to our data. In the units of the DKO animals, it was not feasible to faithfully assess sound thresholds or dynamic ranges. Furthermore, the overall experimental yield was generally much lower in DKO animals. On average we hit 0.3 sound-responsive neurons per hour of recording, compared to 2.5 in WT. Very likely many DKO neurons failed to generate spontaneous and evoked action potentials, thus we were not able to identify them. Based on the DPOAE data we assume that cochlear mechanics and active amplification and thus the IHC receptor potential thresholds are mostly normal in the DKOs. We cannot exclude that IHC receptor potentials are decreased in amplitude, which however should rather reduce SGN adaptation. We thus interpret the observed increase in adaptation and delay of recovery of SGN spiking to be the effects of changed synaptic $Ca_V1.3$ channel kinetics as described above.

In summary, sound responses of DKO SGNs and cochlear nucleus neurons showed a drastic reduction in spontaneous and evoked action potential rates, which was associated with increased sound thresholds and reduced temporal precision of action potential generation. Together, these factors explain well the poor ABR thresholds (Fig. 5, Fig. S4A).

Discussion

In this study, we examined the effects of genetic disruption of the two major CaBPs expressed in the cochlea. In the IHCs of CaBP1/2-deficient animals, we found largely inactivated Ca^{2+} currents with prolonged recovery from inactivation upon mild oscillatory stimulation, which did not support sufficient neurotransmission for proper sound encoding. The extent of channel inactivation and impairment of IHC synaptic exocytosis aggravated with the extent of IHC stimulation. Similarly, SGN action potential rates *in vivo* declined at increased sound stimulation rates and displayed very strong adaptation upon stimulation, going along with impaired sustained IHC exocytosis. The impairment was significantly stronger as compared to the phenotype in the absence of CaBP2 alone. This suggests that under *in vivo* conditions the roles of CaBP1 and 2 are partially overlapping.

Function and expression profile of CaBPs in the cochlea. The murine cochlea shows strong expression of CaBP1 and 2, while the levels of CaBP4 and 5 seem negligible [19,42,43]. Among the three putative splice isoforms of CaBP2 differing in the N-terminal aminoacid sequence, the alternative long isoform, termed CaBP2-alt, may be the most abundant splice variant in the IHCs, as suggested by in situ hybridization [19], while SGNs seem to express none. So far our gene-rescue experiments were conducted with the conventional L-isoform only, which however shows strong

inhibition of $Ca_V1.3$ inactivation in the HEK cells as well as IHCs [12,20]. Alternative splicing also generates three CaBP1 variants [19]. The alternative long variant caldendrin is primarily expressed in the SGNs or the surrounding glia cells, while weak expression of CaBP1-S/-L (as judged by the mRNA production) was detected in hair cells [19]. Our DKO model lacks all splice isoforms of both CaBPs. Since *Cabp1* KO animals show normal hearing several weeks after birth, and CaBP2 is absent from SGNs, we interpret the hearing impairment of the *Cabp1/2* DKO animals to be primarily due to defects of hair cell function and judge contribution by SGN dysfunction as rather unlikely, at least in the early postnatal weeks.

The ABR thresholds of *Cabp1* KO animals are normal at 4 weeks of age, but deteriorate with age, starting with high-frequency hearing loss at 9 weeks. Conversely, the suprathreshold ABRs to tone burst stimulation show increased amplitudes and reduced latencies during the first two months after birth. This suggested enhanced SGN activity at first, followed by slowly progressing functional decline, possibly due to overexcitability-related toxicity [11]. Due to early strong impairment of hair cell synaptic function, the latter effects are not expected to occur in the DKO animals. However, ABR thresholds for clicks were further increased and an additional reduction in the summating potential was observed in the DKO animals at two as compared to one month of age despite relatively well-preserved DPOAEs (Fig. S4). It thus remains to be clarified whether the absence of CaBP1 and 2 may affect (IHC) mechanotransduction at more mature stage. Unfortunately, this is difficult to assess due to the increased fragility of hair bundles at mature age. Intriguingly, CaBPs share similarities with calcium and integrin-binding protein 2 (CIB2), associated with hair bundle defects [44–46]. When we overexpressed CaBP2 in the DKO animals, interestingly, suprathreshold responses of ABR wave I were very large (Fig. 5) and the summated wave I-IV responses were not distinguishable from WT responses. However, tone burst and click thresholds did not completely recover to WT levels. The difference in the extent of rescue of threshold and suprathreshold responses may be due to possibly distinct requirements for CaBPs at synapses providing inputs to low- or high-threshold SGN fibers. Apart from a small shift in voltage activation and abundance of the channels [47], not much is known about the possible differences in the $Ca_V1.3$ properties and interaction partners at distinct ribbons. Neither is it known whether the composition of the channel complex [t.i. proportions of alpha subunit splice variants [18,48,49], and different beta subunits [50]] differs systematically between ribbon-AZs partnered by different classes of SGN fibers. Investigation of intracellular Ca^{2+} signals from individual AZs may offer some answers in the future. Different amounts of rescue of threshold vs suprathreshold responses could be further due to confounding effects of CaBP1 and 2, e.g. augmented SGN response in *Cabp2*-injected DKO animals still lacking CaBP1 (reminiscent of the phenotype of young *Cabp1* KOs) in combination with "only" partial rescue of CaBP2 function. Possible adverse effects of exogenous expression of CaBP2, e.g. off-target toxicity and uncontrolled overexpression of the transgene, have been discussed previously [20], and further experiments need to be performed in the future to achieve (close to) complete rescue of hearing function in these mutants.

Our results reveal that the amount of $Ca_V1.3$ inactivation increases dramatically when IHCs lack both major CaBPs. This suggests that the sole loss of either CaBP1 or 2, can be largely compensated by the other of the two CaBP members. Yet, the loss of CaBP2 alone still results in progressive hearing impairment in mice and DFNB93 in humans. The inability of CaBP1 to fully replace CaBP2 in the CaBP2-deficient IHCs may stem from either too low concentration of CaBP1 (and other CaBPs), or from distinct properties of the two proteins (enabling CaBP2 to exert more potent suppression of $Ca_V1.3$ inactivation, e.g. VDI, or potentially performing additional, yet unidentified cellular function). Interestingly, even the overexpression of CaBP2 did not completely rescue the hearing phenotype of DKO animals. Future experiments involving overexpression of CaBP1 (and possibly CaBP1-2 molecular chimeras) in *Cabp2* SKO and/or *Cabp1/2* DKO animals together with further biophysical interaction studies are required to gain a better understanding of the potentially distinct functions performed by the two major CaBPs in the hair cells.

CaBPs as the regulators of IHC synaptic transmission. Previously, patch-clamp recordings surprisingly revealed normal whole-cell exocytosis in CaBP2-deficient IHCs despite enhanced inactivation of calcium channels [12], and no obvious defects in the CaBP1-deficient IHCs [11]. In contrast, IHCs deficient in both, CaBP1 and 2, displayed strongly impaired exocytosis. Transgenic overexpression of CaBP2 in IHCs resulted in a partial recovery of the Ca^{2+} influx and an even better recovery of exocytosis. Together, these results corroborate the hypothesis of the role of CaBPs in preventing extensive steady-state inactivation of $Ca_V1.3$ channels to support sufficient synaptic exocytosis for proper sound encoding [12]. The lack of CaBP1 and 2 seems to impair sustained exocytosis and efficiency of exocytosis beyond the levels that could be attributed to reduced Q_{Ca}. This became particularly apparent when the IHCs were challenged by higher holding potentials in combination with large sine wave amplitudes during capacitance measurements, which we assume to partially deplete the pool of RRP vesicles. At present, it is not clear how CaBPs may directly affect the exocytosis, e.g. by influencing the coupling of the $Ca_V1.3$ channels and synaptic vesicles or potentially affecting the expression levels of other synaptic proteins that mediate synaptic vesicle replenishment. In this respect, it is worth noting that CaBP1 and 2 have been reported to modulate CaMKII activity [2], which in turn may through phosphorylation modify the Ca^{2+} affinity of otoferlin C2 domains [51]. In the future, it is thus essential to investigate if CaBPs regulate the function of otoferlin and thus synaptic processes. Furthermore, other members of the CaBP/CaM family have been suggested to interact with synaptic proteins and/or be involved in the regulation of synaptic vesicle cycle: CaM in endocytosis [52], CaBP5 in priming via interaction with Munc18-1 [53], CaBP1 via inhibition of IP_3 receptors [54,55]. An important challenge for future studies is to identify further CaBP(2)-interaction partners and possible functions beyond modulation of calcium channels.

The recovery of $Ca_V1.3$ channels from inactivation upon prolonged milder stimulation (2 min sine wave) was following a single exponential course in WT IHCs, but additional slower component was detected in the IHCs deficient in CaBP1 and 2. In the IHCs of injected DKO animals, this second component appeared smaller and faster, suggesting that CaBPs speed-up recovery of $Ca_V1.3$ channels from inactivation. Upon a comparably shorter (500 ms) but stronger depolarization, the recovery of $Ca_V1.3$ from inactivation was very fast in DKO IHCs, in fact faster than in (WT or) *Cabp2*-injected DKO IHCs. We hypothesize this to be due to faster Ca^{2+} clearance from the DKO IHC synapses as the intracellular $[Ca^{2+}]$ levels are expected to rise significantly less in CaBP1/2-deficient IHCs due to fast and pronounced $Ca_V1.3$ inactivation. In contrast, higher $[Ca^{2+}]$ may build-up at the synapses of WT and "rescued" DKO IHCs; consequently, CDI may be prolonged. Another plausible explanation for the observation is, if we consider that CaPB2 may act as a slow mobile calcium buffer [12]. In the presence of excessive "overexpressed" CaBP2 in injected IHCs of DKO animals, the depolarization-evoked intracellular $[Ca^{2+}]$ transients may become biexponential as has been described for parvalbumin alpha [56], which may prolong the time constant of Ca^{2+} clearance and determine the kinetics of channel recovery from inactivation. Our results of the recovery upon 500-ms depolarization may thus mostly reflect the kinetics of the Ca^{2+} clearance from the cytosol, while the experiments with prolonged sine wave stimulation (when e.g. the function of calcium pumps and other extrusion systems and intracellular $[Ca^{2+}]$ levels may be at equilibrium), may report the speed of calcium channel recovery from a slow inactivation. Slow inactivation with very slow recovery was described for Na_V, K_V and Ca_V channels [26–29]. It has to be noted that the time constant of recovery of $Ca_V1.2$ from slow VDI was comparably shorter (4 s) as observed in the present study [29]. Possibly different conditions (e.g. stimulus parameters) or channel subtype may underlie this difference. Future experiments combining patch-clamp recordings with calcium imaging should be employed to address this question further.

From our *in vitro* recordings at higher holding potential or upon prolonged stimulation with small voltage oscillations we predict that *in vivo* a noteworthy proportion of $Ca_V1.3$ channels may be effectively inactivated and not available for mediating IHC synaptic transmission. This could explain the low experimental yield, the low "spontaneous" spiking rates of the SGNs and lower sound-evoked spike rates with higher thresholds. Similarly, thresholds may be high for tone bursts in the ABR measurements due to high proportion of inactivated channels upon stimuli at high repetition

rate used for these recordings (20-40 Hz repetition rate for click and tone burst stimulation, respectively). We believe that any sound stimulation and potentially even spontaneous activity may effectively silence DKO synapses. As expected, longer interstimulus intervals may allow a larger proportion of channels to recover between stimuli, increasing the onset rates of SGN spiking while decreasing the first spike latency and jitter. Still, high-intensity stimulation had to be used in the DKO animals to evoke an increase in SGN spiking. It is possible that spontaneous activity and/or our routines used to determine the SGN characteristic frequency inactivate a majority of IHC $Ca_V1.3$ channels thus not allowing reliable assessment of the characteristic frequency. In addition to $Ca_V1.3$ channel inactivation, other factors may contribute to low hearing sensitivity, such as possible impairment of synaptic vesicle replenishment, or IHC mechanotransduction, and, in some cases, impaired cochlear amplification. A complete lack of BK channels was shown to affect the IHC resting potential, receptor potential amplitude and timing, and consequently the timing and frequency of spiking in the SGNs [57]. Lower expression of BK channels is not expected to significantly affect hearing thresholds, at least initially [58–60], and thus likely does not contribute to the hearing defect of the *Cabp1/2* DKO animals. However, at advanced ages, the effects described above may combine with the lack of CaBPs to additionally desynchronize already scarce spikes in the auditory nerve fibers.

Clinical implications for the diagnosis and treatment of DFNB93. The SGN sound encoding phenotype in *Cabp2*-KO mice was surprisingly mild compared to the deficits observed in DFNB93 patients [e.g. 14,12]: SGN thresholds and evoked spike rates were only minimally affected [12]. Given the strong recovery phenotype in the DKO we now consider the possibility that synaptic activity in response to everyday sounds could already inactivate calcium channels and reduce SGN spiking. In this case, communication in daily life would be more strongly impaired than audiometric data obtained in a soundproof chamber, especially when short stimuli separated by silent intervals are used. In mild cases, a gap detection test may be used to assess auditory fatigue. Based on our data we further predict that patients that have a stronger channel inactivation phenotype might not benefit much from sound amplification by hearing aids. Instead, they should use technologies that improve the signal to noise ratio (e.g. noise cancellation, directional microphones, wireless transmitters). An alternative or additional future treatment approach may be gene therapy. In our hands, re-expression of CaBP2 by genetic treatment partially restored $Ca_V1.3$ properties, IHC exocytosis and consequently ABR wave amplitudes and thresholds. This successful viral rescue supports the view that a reversible inactivation of IHC $Ca_V1.3$ currents is the main mechanism of hearing loss in CaBP-related deafness, which can be targeted by gene therapy in the future.

Materials and Methods

Animals
For the generation of the *Cabp1/Cabp2*-double-knockout (*Cabp1/2*-DKO) line, *CaBP2^{LacZ/LacZ}* mice, carrying LacZ trapping cassette in place of the exons 3 and 4 [12], were crossbred with *Cabp1* KO mice [61] and maintained on a C57/Bl6 background. Mice from 3-13 weeks were used for electrophysiological and systems physiology recordings. All experiments complied with the national animal care guidelines and were approved by the University of Göttingen Board for animal welfare and the animal welfare office of the state of Lower Saxony.

Systems physiology recordings: ABR and DPOAE
Prior to the experiments mice were weighed and anesthetized via intraperitoneal injections of ketamine (125 µg/g) and xylazine (2.5 µg/g). Recordings were obtained in a custom-made soundproof box with a Tucker-Davis (TDT) BioSig System II/BioSig software (Tucker-Davis Technologies, Alachua, FL). Body temperature was maintained using a heating pad (37°C) and ECG was monitored continuously. For ABR recordings, subcutaneous needles were placed at the vertex, the ipsilateral pinna and at the lower back. Using a JBL2402 speaker in free-field configuration, tone bursts (10-ms plateau, 1-ms cos^2 rise/fall) or clicks of 0.03 ms were ipsilaterally presented at 40 Hz for tone bursts or 20 Hz for clicks. ABR traces were amplified 50000-fold, band-pass filtered between 400

Hz and 4 kHz (NeuroAmp) and averaged from 1300 repetitions sampled at 50 kHz for 20ms. Thresholds were defined in stacked waveforms as the lowest stimulus intensity (dB SPL) at which a reproducible waveform could be visually detected with a 10 dB precision. Wave I amplitudes were measured peak-to-trough. DPOAEs were measured using a custom-written MATLAB (MathWorks) routine. Two primary tones f1 and f2 (ratio f1/f2 = 1.2, level difference f2 = f1 + 10 dB) were presented via two speakers (MF-1, TDT) and a custom-made ear probe containing an MKE-2 microphone (Sennheiser). Stimulus duration was 16 ms and f2 levels varied from 10 to 70 dB SPL in 10 dB steps. The microphone signal was amplified (DMX 6Fire; Terratec (young animals), or UAC-2; Zoom (older animals)) and digitized (TDT System III). DPOAE amplitude was analyzed at 2×f2-f1 using custom-written MATLAB software (MathWorks) with fast Fourier transformation. DPOAE thresholds were determined as responses that exceeded -18 dB SPL in young and -4 dB SPL in older animals.

Auditory nerve fiber recordings

Mice were anesthetized by injection of xylazine (5 mg/kg), urethane (1.32 mg/kg) and buprenorphine (0.1 mg/kg) i.p., tracheotomized, and placed in a stereotactic apparatus. The left occipital bone and the cerebellum were partially removed to visualize the left superior semicircular canal. The latter served as a landmark to navigate a glass microelectrode through the cochlear nucleus, aiming at the location where the auditory nerve leaves the internal auditory canal and enters the cochlear nucleus. 85 or 90 dB broadband noise bursts (ScanSpeak, Avisoft Bioacoustics) were used as search stimuli to identify neurons that displayed sound-driven spiking activity. They were characterized by spontaneous rate and frequency tuning and the responses to tone bursts at the characteristic frequency, 30 dB above the threshold, as previously described [35]. In the majority of sound-responsive DKO neurons, thresholds were so high and tuning so broad that neither the automated tuning curve algorithm nor systematic or manual changes in sound frequency elicited frequency-specific responses. In these cases, we either chose a subjective "best frequency" or used white noise burst stimulation at 85 dB.

Patch-clamp recordings

For patch-clamp recordings, the apical coils of 3-4-week-old animals were dissected in HEPES-Hanks solution containing (in mM): 10 HEPES, 5.35 KCl, 141.7 NaCl, 0.5 MgSO$_4$, 1 MgCl$_2$, 11.1 D-glucose and 3.42 L-glutamine. pH was adjusted to 7.2. IHCs of the ~6 kHz tonotopical region were investigated using the perforated patch configuration. The pipette solution contained (in mM): 130 Cs-gluconate, 10 TEA-Cl, 10 4-AP, 1 MgCl$_2$, 10 HEPES, 300 mg/mL amphotericin B, pH 7.2 ~290 mOsm. The bath solution was composed of (in mM): 111 NaCl, 35 TEA-Cl, 2.8 KCl, 1 CsCl, 1 MgCl$_2$, 10 NaOH-Hepes, 11.3 D-glucose, and 1.3 or 2 CaCl$_2$ or BaCl$_2$ (pH 7.2, 300–310 mOsm). When the bath solution contained 2 mM BaCl$_2$/CaCl$_2$, NaCl was adjusted to 110 mM. The bath solution was supplemented by the SK2 channel blocker apamin (100 nM; Peptanova, Germany) freshly on the day of the experiment. Recordings were performed at room temperature using an EPC-9 amplifier (Heka-Germany) controlled by Pulse software. Currents were leak-corrected using a p/10 protocol, sampled at 50 kHz and corrected offline for the liquid junction potential (-15 mV). IHCs were patched at a holding potential of -85 mV (in a subset of experiments, -65 and -55 mV) and stimulated with depolarizations to the peak Ca^{2+}-current potential. Changes in membrane capacitance were monitored by applying a 2-kHz-sinewave with an amplitude of 35 mV. In a subset of experiments, the amplitude was reduced to 15 mV to diminish possible stimulation of the cells caused by the sinewave. VDI and CDI were calculated as follows: VDI = 1 - I_{Ba_500}/$I_{Ba\_initial}$ and CDI = 1 − (I_{Ca_500}/$I_{Ca\_initial}$)/(I_{Ba_500}/$I_{Ba\_initial}$), where I_{Ba_500} and I_{Ca_500} represent whole cell Ba^{2+} and Ca^{2+} currents 500 ms after beginning of depolarization, and $I_{Ba\_initial}$ and $I_{Ca\_initial}$ correspond to the initial, peak currents with the respective divalent cations [22].

Immunohistochemistry

Apical coils (6-8 kHz tonotopical region) of the organ of Corti were fixed with 4% (vol/vol) formaldehyde in PBS on ice. After incubation for 1 h at room temperature in goat serum dilution buffer [16% (vol/vol) normal goat serum, 450 mM NaCl, 0.6% Triton X-100, 20 mM phosphate buffer (pH

7.4)], the explants were incubated in primary antibodies over night at 4 °C. The following antibodies were used: chicken anti-Homer1 (1:200, Synaptic Systems), rabbit anti-Myosin6 (1:200, Proteus), guinea pig anti-Synapsin1/2 (1:200, Synaptic Systems), rabbit anti-RibeyeA (1:200, Synaptic Systems), guinea pig anti–parvalbumin-α (1:200; Synaptic Systems), rabbit anti-BK (1:100; Alomone), rabbit anti-SK2 (1:200; Alomone), mouse IgG1 anti-CtBP2 (1:200; BD Biosciences), rabbit anti-GluA3 (1:200; Santa Cruz), rabbit–anti-CaV1.3 (1:100; Alomone), chicken anti-GFP (1:200, Abcam). The explants were then incubated with secondary AlexaFluor-labeled antibodies (1:200; Molecular Probes) for 1 h at room temperature, washed 3 times in PBS and mounted. Confocal images were acquired with an Abberior Instruments Expert Line STED microscope using a 1.4 NA 100x oil immersion objective. ImageJ software (NIH) was employed to analyze 2D confocal stacks and generate maximum projections. Volumetric synapse analysis ran on Imaris (Bitplane, Zurich, Switzerland) paired with a custom-written Matlab software [62].

Molecular biology and AAV vector production
The construct containing the long isoform of mouse CaBP2 (CaBP2-L; NCBI Reference Sequence: NP_038906.2) and eGFP is depicted in Fig. S1A. AAV9-PHP.eB particles were generated using our standard AAV purification procedure previously described in more detail in ref. [63]. In brief, triple transfection of HEK-293T cells was performed using pHelper plasmid (TaKaRa/Clontech), the *trans*-plasmid providing viral capsid PHP.eB (a generous gift from Viviana Gradinaru, Addgene plasmid #103005) and the *cis*-plasmid providing hCMV/HBA_wtCaBP2-P2A-eGFP. PHP.eB viral particles were harvested from the medium 72 h after, and from cells and the medium 120 h after transfection. Precipitation was done with 40% polyethylene glycol 8000 (Acros Organics, Germany) in 500 mM NaCl for 2 h at 4°C. Both, precipitate and cells were lysed in high salt buffer (500 mM NaCl, 2 mM $MgCl_2$, 40 mM Tris-HCl, pH 8,0) and non-viral DNA was degraded using salt-activated nuclease (SAN, Arcticzymes, USA). The cell lysates were clarified by centrifugation at 2000×g for 10 min and then purified over iodixanol (Optiprep, Axis Shield, Norway) step gradients (15, 25, 40, and 60%) at 350000×g for 2.25 h [64,65]. Finally, viral particles were concentrated using Amicon filters (EMD, UFC910024) and formulated in sterile PBS supplemented with 0.001% Pluronic F-68 (Gibco, Germany). Virus titer was $8.2×10^{12}$ genome copies/ml measured by determining the number of DNase I resistant genome copies using qPCR (StepOne, Applied Biosystems) and AAV titration kit (TaKaRa/Clontech). The purity of viruses was routinely checked by silver staining (Pierce, Germany) after gel electrophoresis (NovexTM 4-12% Tris-Glycine, Thermo Fisher Scientific).

Virus injections
Mice were injected at postnatal day 6 using the round window approach as described in earlier studies [66]. In brief, anesthesia was established with isoflurane (5% for induction, 2-3% for maintenance, frequent testing of the absence of hind-limb withdrawal reflex). For analgesia, buprenorphine (0.1 mg/kg, 30 min before surgery) and carprofen (5 mg/kg, during and 1 day after surgery) were applied subcutaneously and Xylocain (10 mg spray) locally. Body temperature was maintained by a custom-built heating blanket. Following a retro-auricular approach, the facial nerve was exposed to determine where to puncture the cartilaginous bulla with the injection pipette and target the scala tympani where virus suspension (1-1.5 µl, corresponding to 0.82-$1.23×10^{10}$ AAV particles) was injected. Following injection, the endogenous tissue was relocated and the surgical situs closed by suturing the skin. 2-3 weeks later, the injected mice were used for hearing tests, *in vitro* electrophysiology and immunohistochemistry of the cochleae.

Data Analysis
Data were analyzed using Igor Pro Software (Wavemetrics), GraphPad PRISM (Graphpad Software, Inc.), Matlab (Mathworks) and Imaris (Bitplane). All data are presented as mean ± SEM. For the analysis of half-max Ca^{2+} current (V_{half}) and activation constant (k_{act}), current-voltage traces were transformed into an activation function ($I_{Ca} = g_{max}(V-V_{rev})$), which was subsequently fitted by a Boltzmann function: $1/(1 + exp((V_{0.5}−V)/k_{act}))$. G_{max} describes the maximum conductance, V the holding potential and V_{rev} the reversal potential. Data were tested for randomness, normality

(Jaque-Bera test), equality of variances (F-test) and finally compared for statistical significance by performing Student's *t*-test for normally distributed data of equal variance or else by the Wilcoxon rank sum test. Systems physiology data were tested for significance using ANOVA with Tukey's or Šidak's multicomparisons tests.

Acknowledgments

The authors would like to thank Christiane Senger-Freitag for the virus injections, Daniela Gerke for help with virus production, and Sandra Gerke, Ina Preuss, and Sina Langer for excellent technical support. The authors would further like to thank Amy Lee (University of Iowa) for providing the *Cabp1* KO animals, and Mohona Mukhopadhyay and Tobias Moser for critical reading of the paper, useful comments and suggestions. This work was supported by the German Research Foundation: DFG Priority Program 1608 "Ultrafast and temporally precise information processing: Normal and dysfunctional hearing" [to TP (PA 2769/1-1)], the Clinician Scientist Program "Cell Dynamics in Disease and Therapy" at the University Medical Center Göttingen (to DO; project number 413501650), CRC889 (subproject A06; to NS, and B09; to TP); and by the Multiscale Bioimaging Cluster of Excellence (EXC 2067/1- 390729940; to TP).

References

1. Pangrsic, T., Singer, J. H. & Koschak, A. Voltage-Gated Calcium Channels: Key Players in Sensory Coding in the Retina and the Inner Ear. *Physiol. Rev.* **98**, 2063–2096 (2018).
2. Haeseleer, F. *et al.* Five members of a novel Ca(2+)-binding protein (CABP) subfamily with similarity to calmodulin. *J. Biol. Chem.* **275**, 1247–1260 (2000).
3. Findeisen, F. & Minor, D. L. Structural basis for the differential effects of CaBP1 and calmodulin on Ca(V)1.2 calcium-dependent inactivation. *Struct. Lond. Engl. 1993* **18**, 1617–1631 (2010).
4. Haeseleer, F. *et al.* Essential role of Ca2+-binding protein 4, a Cav1.4 channel regulator, in photoreceptor synaptic function. *Nat. Neurosci.* **7**, 1079–1087 (2004).
5. Lee, A. *et al.* Differential modulation of Ca(v)2.1 channels by calmodulin and Ca2+-binding protein 1. *Nat. Neurosci.* **5**, 210–217 (2002).
6. Yang, P. S. *et al.* Switching of Ca2+-dependent inactivation of Ca(v)1.3 channels by calcium binding proteins of auditory hair cells. *J. Neurosci.* **26**, 10677–10689 (2006).
7. Yang, P. S., Ben Johny, M. & Yue, D. T. Allostery in Ca2+ channel modulation by calcium binding proteins. *Nat. Chem. Biol.* **10**, 231–238 (2014).
8. Baig, S. M. *et al.* Loss of Ca(v)1.3 (CACNA1D) function in a human channelopathy with bradycardia and congenital deafness. *Nat. Neurosci.* **14**, 77–84 (2011).
9. Platzer, J. *et al.* Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type Ca^{2+} channels. *Cell* **102**, 89–97 (2000).
10. Cui, G. *et al.* Ca2+-binding proteins tune Ca2+-feedback to Cav1. 3 channels in mouse auditory hair cells. *J. Physiol.* **585**, 791–803 (2007).
11. Yang, T. *et al.* Functions of CaBP1 and CaBP2 in the peripheral auditory system. *Hear. Res.* **364**, 48–58 (2018).
12. Picher, M. M. *et al.* Ca2+-binding protein 2 inhibits Ca2+-channel inactivation in mouse inner hair cells. *Proc. Natl. Acad. Sci.* **114**, E1717–E1726 (2017).
13. Koohiyan, M. *et al.* A Novel Pathogenic Variant in the CABP2 Gene Causes Severe Nonsyndromic Hearing Loss in a Consanguineous Iranian Family. *Audiol. Neurotol.* **24**, 258–263 (2019).
14. Schrauwen, I. *et al.* A Mutation in CABP2, Expressed in Cochlear Hair Cells, Causes Autosomal-Recessive Hearing Impairment. *Am. J. Hum. Genet.* **91**, 636–645 (2012).
15. Sheyanth, I. N. *et al.* First reported *CABP2* -related non-syndromic hearing loss in Northern Europe. *Mol. Genet. Genomic Med.* **9**, (2021).
16. Tabatabaiefar, M. A. *et al.* DFNB93, a novel locus for autosomal recessive moderate-to-severe hearing impairment. *Clin. Genet.* **79**, 594–598 (2011).

17. Bock, G. *et al.* Functional Properties of a Newly Identified C-terminal Splice Variant of Cav1.3 L-type Ca2+ Channels. *J. Biol. Chem.* **286**, 42736–42748 (2011).
18. Scharinger, A. *et al.* Cell-type-specific tuning of Cav1.3 Ca(2+)-channels by a C-terminal auto-modulatory domain. *Front. Cell. Neurosci.* **9**, 309 (2015).
19. Yang, T. *et al.* Expression and Localization of CaBP Ca2+ Binding Proteins in the Mouse Cochlea. *PLoS ONE* **11**, (2016).
20. Oestreicher, D., Picher, M. M., Rankovic, V., Moser, T. & Pangrsic, T. Cabp2-Gene Therapy Restores Inner Hair Cell Calcium Currents and Improves Hearing in a DFNB93 Mouse Model. *Front. Mol. Neurosci.* **14**, 689415 (2021).
21. Shaltiel, L. *et al.* Complex regulation of voltage-dependent activation and inactivation properties of retinal voltage-gated Cav1.4 L-type Ca2+ channels by Ca2+-binding protein 4 (CaBP4). *J. Biol. Chem.* **287**, 36312–36321 (2012).
22. Tadross, M. R., Ben Johny, M. & Yue, D. T. Molecular endpoints of Ca2+/calmodulin- and voltage-dependent inactivation of Ca(v)1.3 channels. *J. Gen. Physiol.* **135**, 197–215 (2010).
23. Russell, I. J. & Sellick, P. M. Low-frequency characteristics of intracellularly recorded receptor potentials in guinea-pig cochlear hair cells. *J. Physiol.* **338**, 179–206 (1983).
24. Dierich, M. *et al.* Optimised tuning of auditory inner hair cells to encode complex sound through synergistic activity of six independent K+ current entities. *Mol. Cell* **in press**, (2020).
25. Oliver, D., Knipper, M., Derst, C. & Fakler, B. Resting potential and submembrane calcium concentration of inner hair cells in the isolated mouse cochlea are set by KCNQ-type potassium channels. *J. Neurosci. Off. J. Soc. Neurosci.* **23**, 2141–2149 (2003).
26. Bezanilla, F., Taylor, R. E. & Fernández, J. M. Distribution and kinetics of membrane dielectric polarization. 1. Long-term inactivation of gating currents. *J. Gen. Physiol.* **79**, 21–40 (1982).
27. Payandeh, J., Gamal El-Din, T. M., Scheuer, T., Zheng, N. & Catterall, W. A. Crystal structure of a voltage-gated sodium channel in two potentially inactivated states. *Nature* **486**, 135–139 (2012).
28. Olcese, R., Latorre, R., Toro, L., Bezanilla, F. & Stefani, E. Correlation between Charge Movement and Ionic Current during Slow Inactivation in Shaker K+ Channels. *J. Gen. Physiol.* **110**, 579–589 (1997).
29. Ferreira, G., Ríos, E. & Reyes, N. Two components of voltage-dependent inactivation in Ca(v)1.2 channels revealed by its gating currents. *Biophys. J.* **84**, 3662–3678 (2003).
30. Eckrich, S. *et al.* Cochlea-Specific Deletion of Cav1.3 Calcium Channels Arrests Inner Hair Cell Differentiation and Unravels Pitfalls of Conditional Mouse Models. *Front. Cell. Neurosci.* **13**, (2019).
31. Kim, K. X. *et al.* Vesicular Glutamatergic Transmission in Noise-induced Loss and Repair of Cochlear Ribbon Synapses. *J. Neurosci.* **39**, 2228–18 (2019).
32. Brandt, A., Striessnig, J. & Moser, T. Ca $_V$ 1.3 Channels Are Essential for Development and Presynaptic Activity of Cochlear Inner Hair Cells. *J. Neurosci.* **23**, 10832–10840 (2003).
33. Neef, J. *et al.* The Ca^{2+} Channel Subunit beta2 Regulates Ca^{2+} Channel Abundance and Function in Inner Hair Cells and Is Required for Hearing. *J. Neurosci.* **29**, 10730 (2009).
34. Khimich, D. *et al.* Hair cell synaptic ribbons are essential for synchronous auditory signalling. *Nature* **434**, 889–894 (2005).
35. Strenzke, N. *et al.* Hair cell synaptic dysfunction, auditory fatigue and thermal sensitivity in otoferlin Ile515Thr mutants. *EMBO J.* **35**, e201694564 (2016).
36. Bharadwaj, H. M., Verhulst, S., Shaheen, L., Liberman, M. C. & Shinn-Cunningham, B. G. Cochlear neuropathy and the coding of supra-threshold sound. *Front. Syst. Neurosci.* **8**, 26 (2014).
37. Jean, P. *et al.* The synaptic ribbon is critical for sound encoding at high rates and with temporal precision. *eLife* **7**, e29275 (2018).
38. Jing, Z. *et al.* Disruption of the presynaptic cytomatrix protein bassoon degrades ribbon anchorage, multiquantal release, and sound encoding at the hair cell afferent synapse. *J. Neurosci.* **33**, 4456–4467 (2013).
39. Taberner, A. M. & Liberman, M. C. Response Properties of Single Auditory Nerve Fibers in the Mouse. *J. Neurophysiol.* **93**, 557–569 (2005).

40. Vogl, C. *et al.* The BEACH protein LRBA is required for hair bundle maintenance in cochlear hair cells and for hearing. *EMBO Rep.* **18**, 2015–2029 (2017).
41. Pangrsic, T. *et al.* Hearing requires otoferlin-dependent efficient replenishment of synaptic vesicles in hair cells. *Nat. Neurosci.* **13**, 869–876 (2010).
42. Jean, P. *et al.* Single-cell transcriptomic profiling of the mouse cochlea: An atlas for targeted therapies. *Proc. Natl. Acad. Sci.* **120**, e2221744120 (2023).
43. Xu, Z. *et al.* Profiling mouse cochlear cell maturation using 10× Genomics single-cell transcriptomics. *Front. Cell. Neurosci.* **16**, 962106 (2022).
44. Giese, A. P. J. *et al.* CIB2 interacts with TMC1 and TMC2 and is essential for mechanotransduction in auditory hair cells. *Nat Commun* **8**, 43 (2017).
45. Riazuddin, S. *et al.* Mutations in CIB2, a calcium and integrin binding protein, cause Usher syndrome type 1J and nonsyndromic deafness DFNB48. *Nat. Genet.* **44**, 1265–1271 (2012).
46. Wang, Y. *et al.* Loss of CIB2 Causes Profound Hearing Loss and Abolishes Mechanoelectrical Transduction in Mice. *Front. Mol. Neurosci.* **10**, 401 (2017).
47. Ohn, T.-L. *et al.* Hair cells use active zones with different voltage dependence of Ca^{2+} influx to decompose sounds into complementary neural codes. *Proc. Natl. Acad. Sci.* **113**, E4716–E4725 (2016).
48. Singh, A. *et al.* Modulation of Voltage- and Ca2+-dependent Gating of CaV1.3 L-type Calcium Channels by Alternative Splicing of a C-terminal Regulatory Domain. *J. Biol. Chem.* **283**, 20733–20744 (2008).
49. Vincent, P. F. Y. *et al.* Different CaV1.3 Channel Isoforms Control Distinct Components of the Synaptic Vesicle Cycle in Auditory Inner Hair Cells. *J. Neurosci. Off. J. Soc. Neurosci.* **37**, 2960–2975 (2017).
50. Ortner, N. J. *et al.* RBP2 stabilizes slow Cav1.3 Ca2+ channel inactivation properties of cochlear inner hair cells. *Pflüg. Arch. - Eur. J. Physiol.* **472**, 3–25 (2020).
51. Meese, S. *et al.* Activity-Dependent Phosphorylation by CaMKIIδ Alters the Ca2+Affinity of the Multi-C2-Domain Protein Otoferlin. *Front. Synaptic Neurosci.* **9**, 13 (2017).
52. Wu, X.-S. *et al.* Ca2+ and calmodulin initiate all forms of endocytosis during depolarization at a nerve terminal. *Nat. Neurosci.* **12**, 1003–1010 (2009).
53. Sokal, I. & Haeseleer, F. Insight into the Role of Ca $^{2+}$ -Binding Protein 5 in Vesicle Exocytosis. *Investig. Opthalmology Vis. Sci.* **52**, 9131 (2011).
54. Haynes, L. P., Tepikin, A. V. & Burgoyne, R. D. Calcium-binding Protein 1 Is an Inhibitor of Agonist-evoked, Inositol 1,4,5-Trisphosphate-mediated Calcium Signaling. *J. Biol. Chem.* **279**, 547–555 (2004).
55. Kasri, N. N. *et al.* Regulation of InsP3 receptor activity by neuronal Ca2+-binding proteins. *EMBO J.* **23**, 312–321 (2004).
56. Lee, S.-H., Schwaller, B. & Neher, E. Kinetics of Ca2+ binding to parvalbumin in bovine chromaffin cells: implications for [Ca2+] transients of neuronal dendrites. *J. Physiol.* **525**, 419–432 (2000).
57. Oliver, D. *et al.* The role of BKCa channels in electrical signal encoding in the mammalian auditory periphery. *J. Neurosci. Off. J. Soc. Neurosci.* **26**, 6181–6189 (2006).
58. Rüttiger, L. *et al.* Deletion of the Ca2+-activated potassium (BK) alpha-subunit but not the BKbeta1-subunit leads to progressive hearing loss. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 12922–12927 (2004).
59. Maison, S. F., Pyott, S. J., Meredith, A. L. & Liberman, M. C. Olivocochlear suppression of outer hair cells in vivo: evidence for combined action of BK and SK2 channels throughout the cochlea. *J. Neurophysiol.* **109**, 1525–1534 (2013).
60. Pyott, S. J. *et al.* Cochlear Function in Mice Lacking the BK Channel α, β1, or β4 Subunits. *J. Biol. Chem.* **282**, 3312–3324 (2007).
61. Kim, K. Y. *et al.* Localization and expression of CaBP1/caldendrin in the mouse brain. *Neuroscience* **268**, 33–47 (2014).
62. Neef, J. *et al.* Quantitative optical nanophysiology of Ca^{2+} signaling at inner hair cell active zones. *Nat. Commun.* **9**, 290 (2018).

63. Huet, A. T. & Rankovic, V. Application of Targeting-Optimized Chronos for Stimulation of the Auditory Pathway. in *Methods in Molecular Biology (Clifton, N.J.)* vol. 2191 261–285 (2021).
64. Grieger, J. C., Choi, V. W. & Samulski, R. J. Production and characterization of adeno-associated viral vectors. *Nat. Protoc.* **1**, 1412–1428 (2006).
65. Zolotukhin, S. *et al.* Recombinant adeno-associated virus purification using novel methods improves infectious titer and yield. *Gene Ther.* **6**, 973–985 (1999).
66. Rankovic, V. *et al.* Overloaded Adeno-Associated Virus as a Novel Gene Therapeutic Tool for Otoferlin-Related Deafness. *Front. Mol. Neurosci.* **13**, (2021).

Figures and Tables

Figure 1. Strong inactivation of IHC Ca$_V$1.3 channels upon deletion of *Cabp1* and *Cabp2*.
(A) Schematic of the IHC and one of its several ribbon synapses. CaBPs and CaM modulate voltage-gated calcium channels at the IHC ribbon synapses to shape the presynaptic Ca^{2+} signal. **(B)** Peak normalized Ca^{2+}-currents after a 500-ms depolarization step to the maximum current potential in the apical IHCs of WT and *Cabp1/2*-DKO animals. Note a pronounced inactivation of the Ca^{2+} current in the absence of CaBP1 and 2 and partial recovery upon *Cabp2* gene replacement. **(C)** The fraction of the remaining Ca^{2+}-current after a 500-ms depolarization step is depicted for different mouse models acquired at comparable conditions (room temperature, 2 mM extracellular [Ca^{2+}]). The histogram combines the results of this study (*Cabp1/2*-DKO) with previously published data on CaBP1 [11] and CaBP2 [12]. The WT data combines pooled data from the current study and controls as obtained previously [11]. Note that the recordings from *Cabp2*-KO animals were acquired at the lower apex to mid-cochlear tonotopic positions as the phenotype in the apex (low-frequency positions investigated otherwise) is mild [12]. **(D)** Ca^{2+}-currents upon a train of 10-ms long depolarization steps normalized to the amplitude of the first pulse. **(E)** Ca^{2+}-current-voltage relationships show slightly reduced amplitudes in the IHCs of the *Cabp1/2*-DKO animals transduced with the PHP.eB-*Cabp2*. **(F)** Activation curves were calculated with Boltzmann fits. Note a significant difference in the half-voltage activation (V$_{0.5}$) and slope in the CaBP1/2-deficient IHCs and the persistence of the voltage shift upon re-expression of CaBP2 (*inserts*). See main text for the *p* values. Data in B and D-F was acquired in 1.3 mM, data in C in 2 mM [Ca]$^{2+}$.

Figure 2. Voltage- and Ca²⁺-dependent inactivation are enhanced after deletion of *Cabp1* and *Cabp2*. (A-B) Peak normalized Ca²⁺- and Ba²⁺ currents in WT and CaBP1/2-deficient IHCs recorded in the extracellular solution containing 2 mM of the respective divalent cation. The two types of inactivation were calculated as depicted. **(C)** VDI and CDI were both significantly enhanced in the IHCs of *Cabp1/2*-DKO animals (*asterisks*; Student's *t* test, $p < 0.0001$).

Figure 3. Reduced sustained component of exocytosis and calcium charge transfer (Q_{Ca}) can be partially rescued by *Cabp2*-transgene-expression. (A) Representative Ca^{2+}-current traces and corresponding membrane capacitance changes upon 100-ms long depolarization steps to the peak Ca^{2+}-current potential. **(B)** Capacitance increments with the corresponding Q_{Ca} were probed by different depolarization durations from a holding potential of -85 mV (Student's *t* test or Wilcoxon rank-sum test). CaBP1/2-deficient IHCs showed significant impairment of the sustained exocytosis ($p < 0.05$ for 20-ms, and < 0.001 for 100- and 200-ms step, respectively) and reduced cumulative Ca^{2+} influx ($p < 0.005$ for the longest two pulses) as compared to WT controls (*blue-black asterisks*). AAV-mediated delivery of *Cabp2* improved IHC synaptic function as compared to non-injected *Cabp1/2*-DKO controls ($p < 0.05$ for ΔC_m upon 100- and 200-ms step, and Q_{Ca} upon 200-ms step; *green-blue asterisks*). **(C)** Note a reduced efficiency of Ca^{2+}-dependent exocytosis ($p < 0.001$ for the two longest test pulses; *blue-black asterisks*), which can be efficiently rescued by intracochlear delivery of PHP.eB-*Cabp2* ($p < 0.05$ and 0.005 for 100- and 200-ms pulse, respectively; *green-blue asterisks*). **(D)** Capacitance measurements at a holding potential of -55 mV reveal an aggravation of the phenotype by additional activation of calcium channels between the test pulses ($p < 0.05$ for 10- and 50ms, and $p < 0.005$ for 20-, 100- and 200-ms pulse) and a further discrepancy between the Q_{Ca} of WT and CaBP1/2-deficient IHCs (2-10 ms: $p < 0.05$; 20-200 ms: $p < 0.001$). Also in these recording conditions, the IHCs from *Cabp2*-injected *Cabp1/2*-DKO animals showed increased Q_{Ca} (50-200 ms: $p < 0.005$) and exocytosis (100 ms: $p < 0.005$; 200 ms: $p < 0.001$) as compared to non-injected controls. **(E)** Capacitance increments and the corresponding Q_{Ca} upon 100-ms depolarization steps as recorded in different conditions. Note worsening of the phenotype with increasing IHC activation between the test pulses.

Figure 4. The speed of recovery of Ca²⁺ currents is affected in CaBP1/2-deficient IHCs. (A) Ca²⁺ currents were probed by 5-ms long depolarization steps to the maximum current potential. An initial test pulse was followed by a conditioning stimulus (a 500-ms depolarization step or a 120-s long sine wave). Subsequently, Ca²⁺ currents were tested at various time points to assess the recovery of Ca²⁺-current amplitudes. **(A')** Example Ca²⁺-current recording using a 500-ms long depolarization step as a conditioning stimulus. **(A")** Representative Ca²⁺-current traces in WT and CaBP1/2-deficient IHCs at different time points after a 500-ms long conditioning stimulus. **(B-C)** Ca²⁺-current amplitudes normalized to the amplitude of the pre-conditioning test pulse up to 150 s after a 500-ms depolarization step (B) or a 2-min long sine wave stimulus (C). The inserts show the initial 20 s after the conditioning stimulus. The data were fitted by a single or double exponential function (fit parameters in the Table 1). Note stronger inactivation in the *Capb1/2*-DKO IHCs and delayed recovery of Ca²⁺ currents after the presentation of a long sine wave and a partial recovery upon viral gene transfer of *Cabp2*. Recovery upon a 500-ms pulse was much faster and slowest in the injected animals.

Figure 5. AAV-mediated transgene-expression of *Cabp2* ameliorates hearing loss and recovers ABR amplitude responses in *Cabp1/2*-DKO mice. (A) Significant improvement of ABR thresholds in the injected ears of 3-4-week-old *Cabp1/2*-DKO animals for all tone burst frequencies tested as well as 20-Hz-click stimulus (see Table S1 for the *p* values). **(B)** Average click-evoked responses to an 80-dB 20-Hz-click stimulus. **(C)** A strong recovery of the amplitude of the ABR wave I (Tukey's multicomparison test; $p \leq 0.0001$ for DKO vs WT or vs injected DKO) upon *Cabp2*-transgene expression.

Figure 6. Nearly blocked sound encoding in _Cabp1/2_-DKO mice. Single neuron responses to sound stimulation _in vivo_ in the region of the auditory nerve of _Cabp1/2_-DKO mice were extremely scarce. **(A)** WT poststimulus time histograms of SGNs to 50-ms suprathreshold tone or noise burst stimuli presented at a rate of 5 Hz showed a very high sound onset rate gradually adapting to a sustained rate persisting throughout the stimulus duration. Spike rates were much lower in the mutants. **(B)** Applying a lower stimulus rate of 0.5 Hz partially restored sound onset responses in _Cabp1/2_-DKOs, highlighting the enhanced strength of adaptation. WT and mutant neurons sustained their adapted spiking response throughout the longer 500 ms stimulus used in these recordings. For direct comparison, the average 5-Hz poststimulus time histograms from the same neurons are shown. **(C)** Correlations of peak and adapted rates for different stimulus rates. **(D)** Peak rates for different stimulus rates / interstimulus intervals in WT SGNs strongly increase between 10 and 5 Hz, while in DKOs a longer period of silence is required to observe such difference. **(E)** _Cabp1/2_-DKO neurons had lower spontaneous spike rates than WT SGNs.

Test group	500-ms depolarization					2-min sine wave				
	A_1 (%)	τ_1 (s)	A_2 (%)	τ_2 (s)	y_0 (%)	A_1 (%)	τ_1 (s)	A_2 (%)	τ_2 (s)	y_0 (%)
DKO	65	0.62	-	-	99	19	1.36	21	19.41	98
WT	28	0.75	-	-	100	24	3.83	-	-	97
DKO+PHP.eB-*Cabp2*	22	0.47	14	3.87	99	13	1.88	18	11.56	99

Table 1. Ca^{2+}-current recovery fit parameters. The data was fitted with the single or the double exponential with the following equations:

$y(t) = y_0 - A_1 \times \exp(-(t/\tau_1))$ and $y(t) = y_0 - A_1 \times \exp(-(t/\tau_1)) - A_2 \times \exp(-(t/\tau_2))$, where $y(t)$ represents the peak current amplitude at the time t after the end of conditioning pulse (either 500-ms depolarization or 2-min sine wave) given in % of the initial pre-conditioning peak current amplitude ($100 \times I_{post}/I_{pre}$). While current recovery upon 500-ms depolarization in the IHCs of WT and DKO animals could be well fitted by a single exponential, the data obtained in the injected DKO animals required a double exponential fitting. Upon 2-min sine wave, the current recovery in DKO IHCs (both, injected and non-injected group) also required fitting with a double exponential function.