Mechanical characterization of regenerating *Hydra* tissue spheres

Thomas Perros[1], Anaïs Biquet-Bisquert[1,2], Zacchari Ben Meriem[3], Morgan Delarue[3], Pierre Joseph[3],
Philippe Marcq[4], Olivier Cochet-Escartin[1,*]

1 : Universite Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, Villeurbanne, France.

2 : Centre de Biologie Structurale, Université de Montpellier-CNRS-INSERM, Montpellier, France.

3 : Laboratory for Analysis and Architecture of Systems, Université de Toulouse-CNRS, UPS, INP,
Toulouse, France.

4 : Physique et Mécanique des Milieux Hétérogènes, PMMH, CNRS, ESPCI Paris, Université PSL,
Sorbonne Université, Université Paris Cité, F-75005, Paris, France

* : correspondence should be addressed to olivier.cochet-escartin@univ-lyon1.fr

Abstract

Hydra vulgaris, long known for its remarkable regenerative capabilities, is also a longstanding source of inspiration for models of spontaneous patterning. Recently, it became clear that early patterning during *Hydra* regeneration is an integrated mechano-chemical process where morphogen dynamics is influenced by tissue mechanics. One roadblock to understand *Hydra* self-organization is our lack of knowledge about the mechanical properties of these organisms. In this paper, we combined microfluidic developments to perform parallelized microaspiration rheological experiments and numerical simulations to characterize these mechanical properties. We found three different behaviors depending on the applied stresses: an elastic response, a visco-elastic one and tissue rupture. Using models of deformable shells, we quantify their Young's modulus, shear viscosity as well as the critical stresses required to switch between behaviors. Based on these experimental results, we propose a description of the tissue mechanics during normal regeneration. Our results provide a first step towards the development of original mechano-chemical models of patterning grounded in quantitative, experimental data.

Statement of significance

Hydra vulgaris is a remarkable organism thanks to its regenerative abilities. One can cut this animal into several pieces which will reform a full *Hydra* in a few days. In this process, the pieces have to define a new organizing axis. Recently, researchers have shown that this axis definition is under mechanical control. One roadblock to understand the relationship between tissue mechanics and *Hydra* biology is our lack of knowledge about the mechanical state of this organism. Here, we perform a mechanical characterization using a combination of microaspiration setups and numerical simulations. We finally propose a description of what happens at the mechanical level during *Hydra* regeneration, allowing quantitative approaches questioning the role of mechanical cues in axis definition.

Introduction

Hydra vulgaris has long been a model of choice in developmental biology because of its remarkable regenerative capabilities (1, 2). Almost any excised tissue piece as well as cellular re-aggregates are capable of reforming a fully viable adult in just a few days. In the former example, an excised tissue piece folds back into a closed spherical shape with both its epithelial monolayers, endoderm and ectoderm, engulfing a water-filled lumen. In the latter, the same tissues start by spontaneously sorting into their relative positions (3, 4) before expelling excess cells to reform a hollow sphere. At this point, both regenerative trajectories converge and these tissue spheres start undergoing osmotically-driven oscillations during which they swell because of water flowing from the environment to the lumen up to the point where the tension building up within the tissues becomes too large and leads to a local rupture (5, 6). After rupture, the samples deflate, close the resulting wound and start another swelling-rupture cycle. These osmotic oscillations induce deformations and therefore stresses within the tissues with relative changes in the radii of the spheres up to 30%. These have been referred to as phase I oscillations and are characterized by a high amplitude and a low frequency (6). Indeed, after a few of these cycles, the oscillations clearly change and enter phase II where they have lower amplitude and higher frequency (6).

In parallel to these mechanical oscillations, the samples establish a chemical pattern involving some characterized morphogens to define the oral-aboral axis of the organism. Most notably, a local expression of HyWnt3 has been shown to be an early signal of axial patterning with this activation

defining the future position of the head organizer (7). This local activation is then followed by the
establishment of various chemical gradients within the spheres effectively patterning the whole axis
(8, 9). At that point, the originally symmetrical samples start elongating in an oblong shape at which
ends the adult organs of the head and foot will be regenerated, and the patterning can be considered
complete. The switch between phase I and phase II oscillations was thought to be a signature of the
establishment of axial patterning (6, 10). Since small excised tissue pieces showed both phases, it was
thought that they underwent spontaneous symmetry breaking and retained no memory of axial
patterning just as cellular re-aggregates (5, 6).

The question was then to determine how the spherical symmetry was broken during phase I
oscillations and how the local head organizer was defined. For a long time, the main hypothesis was
that of a purely biochemical spontaneous symmetry breaking in the form of Turing-instabilities of an
unknown reaction-diffusion system. In his seminal work, partially inspired by *Hydra* regeneration,
Turing has shown how a system of two interacting and diffusing chemical species, which he named
morphogens, could become unstable in their homogenous state and spontaneously start forming
structures such as dots or stripes (11). These ideas were critical in developing the field of pattern
formation and have been adapted to a wide variety of systems and organisms (see for example (12,
13) for recent reviews in the context of developmental biology). In this historical development, *Hydra*
has retained its place as a model organism. Notably, Gierer and Meinhardt have expanded on the
seminal ideas of Turing and developed a modified version of his reaction-diffusion system specifically
designed to explain regeneration and grafting experiments in *Hydra* (14). As a result, the so-called
Gierer-Meinhardt model has long been the gold standard in the field. At its inception, it was purely
speculative but, since then, some proteins have been shown to possess many of the characteristics
required by the Gierer-Meinhardt model. Most notably, the protein HyWnt3, involved in the canonical
Wnt pathway is now generally considered to be the activator represented in the Gierer-Meinhardt
model (15) since its expression is restricted to the head organizer, it has self-activating capacities and
it is the first temporal signature of symmetry breaking during regeneration. However, the necessary
long-range inhibitor is notoriously missing. Some promising candidates were put forward, such as
Dickkopf (Dkk) (16), Sp5 (17) or *Hydra* astacin-7 (18) but none of these could reproduce the predictions
of these models, mostly because of differences in expression patterns.

These models also ignore the mechanical aspects of the process, most notably the osmotic oscillations
although it is now well established that they are necessary for proper regeneration (19, 20). This
observation suggested a possible coupling between the mechanical state of the tissues and their
biological response, as has been observed in a variety of contexts and organisms with impacts on cell
division or gene expression including in the canonical Wnt pathway (21–24). Recent results have
demonstrated such a coupling in *Hydra* by which the expression of HyWnt3 is reduced when osmotic
oscillations are blocked (20) providing a potential direct coupling between tissue mechanics and
chemical patterning.

Recent modelling efforts have thus been made in order to incorporate mechano-chemical couplings
(19, 25, 26), for instance by making the diffusion constant of the morphogens within the tissues
dependent on tissue stretch (19). In most cases though, these couplings were not grounded in
experimental evidence. In addition, recent experimental results have started to question the
assumptions described so far and underlying these models. First, it was shown that the shift from
phase I to phase II oscillations was not a direct signature of axial patterning being established. Instead,
the onset of phase II is due to the early apparition of the *Hydra* mouth (27) allowing it to regulate its
osmotic imbalance by mouth opening rather than by tissue rupture. One consequence was that axial
patterning had to be anterior to the shift between oscillation phases. It then became unclear whether

small, excised tissue pieces really went through *de novo* patterning or if they could inherit this
patterning from their original host organism. It now appears that they do retain axial patterning
although the exact mechanism by which they do so remains unclear, whether by the organization of
their ectodermal actin structures called myonemes (28) or by pre-existing biochemical gradients (29).

Although small excised tissue pieces retain axial patterning, it still remains that 1- cellular re-aggregates
which cannot conserve either supracellular actin structures or chemical gradients have to show *de*
novo axial patterning during osmotic oscillations, 2- there is evidence of a direct coupling between
tissue deformations and Hywnt3 expression (20), one of the most important morphogens involved in
Hydra patterning and 3- osmotic oscillations are required for the proper elongation, morphogenesis
and regeneration of both excised tissue pieces and cellular re-aggregates. For all these reasons, the
focus of the field is currently shifting to an integrated view of *Hydra* regeneration as a mechano-
biological process (26, 30). One clear roadblock to the development of these ideas is our lack of
understanding of the rheological properties and mechanical state of regenerating *Hydra* tissue
spheres. This often leads to assumptions as to the rheology of these samples and rough estimates of
their key mechanical parameters. One exception is the development by Veschgini and colleagues of a
two-fingered robotic hand allowing them to apply known, constant deformations on the tissue spheres
and measure the resulting forces (31) giving the first local, quantitative measurements of mechanical
features in *Hydra*. These measurements were not, however, used to deepen our understanding of the
spontaneous osmotic oscillations.

In this work, we try to overcome these limitations and to offer a quantitative characterization of the
mechanics of *Hydra* tissue spheres. To do so, we used the well-established micro-aspiration technique
(32–35) which we adapted to increase its throughput through the use of original microfluidic
constructs following (36). We found three different mechanical behaviors as the aspiration pressure
was increased: first an elastic response, then a viscoelastic one and finally tissue rupture, as observed
in phase I oscillations. Combining our experimental observations and measurements with the
development of a rheological model of elastic shells and numerical simulations, we obtained
quantitative measurements of both the main rheological parameters of *Hydra* tissue spheres and the
critical pressures required to switch between the three regimes. Thanks to these results, we provide a
description of internal tissue mechanics, strains and stresses during osmotic oscillations and reveal
that the tissue spheres behave largely as hyperelastic spherical shells during these oscillations.
Hopefully, this mechanical characterization will serve as a stepping stone for the study of mechano-
biochemical couplings in a quantitative manner.

Material and Methods

<u>*Hydra* maintenance and lines used</u>

3 *Hydra vulgaris* lines were maintained and used for experiments: a watermelon (WM) line
(ectoderm GFP / endoderm DsRed2), a reverse watermelon (RWM) one (ectoderm DsRed2 / endoderm
GFP) and the AEP line from which embryos were obtained for making transgenic animals. *Hydras* were
kept in *Hydra* medium (HM), which consists of 1mM CaCl2, 0.1mM MgCl2, 0.3mM KNO3, 0.5mM
NaHCO3 and 0.08mM MgSO4 at a PH between 7 and 7.3. Cultures were stored at 18°C in the dark in
an incubator (Pol-Ekko-Aparatura). Animals were fed two to three times per week with newly hatched
Artemia (Hobby) and cleaned every two days by changing the medium in which they sit. Animals were
starved for at least 24h before the start of any experiment.

<u>Preparation of tissue spheres</u>

Tissue spheres were made by cutting a whole animal using a sterile scalpel (Holtex, Bistouri
UU n°10), as follows: The head and tail were removed by two transverse cuts and the remaining body
column was first sliced into 3 or 4 pieces. These pieces were cut further to obtain 6 to 8 samples per
adult animal. These samples were left to fold into tissue spheres for 3 to 4 hours in HM.

<u>Design and fabrication of the removable inserts</u>

Sliding elements containing the cylindrical pipettes were fabricated in dry film according to the
protocol detailed in (36). Briefly, photolithography was realized on a 500 μm-thick photosensitive dry
film (DF series), laminated on a silicon wafer. Lateral dimension of the photolithography mask defined
the length of the insert (20 mm) and its height (1 mm), as well as pipette diameter (50 and 100 μm
were used). 100 μm-wide square openings placed at the bottom of the insert were also used for
imaging purposes in Movie S3 and Fig S4. After post-exposure bake on a hot-plate, the inserts were
released from the wafer during overnight development in an acetone bath that both revealed the
patterns and unstick the dry films from the wafer. A single fabrication run allowed to fabricate 100
such sliding elements.

<u>Mounting and preparing of the microfluidic device</u>

We used a microfluidic tool that allows several micro aspirations to be performed in parallel.
This system is composed of a chamber crossed by a removable insert bearing multiple tunnels allowing
suction to happen. Both the chambers and inserts were designed using the CAD software Inventor
2020 (Autodesk). The chamber is composed of a main channel which is 20.5mm long, 4mm wide and
0.5mm high (see Fig S1 for the schematics). Halfway through its length, it is barred by another channel
which will host the removable insert. The perpendicular channel is 13mm long, 0.5mm wide and 1mm
high (Fig S1).

From this design, a mold was manufactured by micromilling using the CNC Mini-Mill/GX (Minitech
machinery corporation) with a two size cutter of 1mm diameter on a brass template. The chamber was
then made of polydimethylsiloxane (PDMS) (Sylgard 184, Dow Corning) supplemented with 10% curing
agent (Dow Corning) from the mold. The PDMS was left to solidify overnight at 70°C and the channel
was manually peeled. Inlets and outlets were created by manual punching using a 1mm wide puncher
(Harris Uni-Core) at each end of the main channel. The insert channel was opened by manual,
transverse cutting to allow insertion of the sliding element. The resulting channel was then bound to
a thin PDMS layer, itself bound to a glass slide (Biosigma VBS655/A) by using an oxygen plasma cleaner
(Harrick plasma PDC002 with the associated PlasmaFlo gaz mixer). The thin PDMS layer was used to
facilitate the introduction of the insert by avoiding direct friction with the glass slide and to avoid
leakage. This introduction is done manually with tweezers and is aided by the fact that the insert
cannot deviate from its associated channel because of the height difference between both channels.
The outlet was connected to a plastic tube of 0.5mm internal diameter and 1.25mm external diameter,
sealed at the other end by a 0.6x30mm needle (BD, microlance 3) attached to a 3mL syringe (Braun,
Omnifix3P). The needle is deliberately slightly wider than the tube to ensure a hermetic connection.
The whole setup was placed in a 100mm Petri dish and immersed in HM. By slowly removing the
syringe's plunger, the channel started filling with HM from the inlet all the way to the syringe while
avoiding to create air bubbles in the main channel and checking for proper sealing of the system. Once
the plunger was fully removed, the height difference ΔH between the water in the syringe and the
water in the Petri dish (see Fig S2 for a schematic of the whole setup) was manually controlled and
created a pressure difference between chamber inlet and outlet according to:

$$\Delta P = \rho g \Delta H$$

Where ρ is the HM density and g the gravitational acceleration.

Micro-aspiration experiments and data analysis

The syringe was first placed 4cm below the water level of the dish to create a light water flow induced
by hydrostatic pressure. Several *Hydra* tissue spheres were manually pipetted into the inlet and were
thus naturally driven to the holes in the inserts by the water flow, at which point the height difference
was set back to 0 for a couple of minutes. A typical experiment could then be started by the application
of a controlled pressure by moving the syringe down again. Images were captured under a microscope
(Zeiss Axio Vert.A1 equipped with an Axiocam 202 camera). In most experiments, timelapses were
recorded in fluorescent microscopy using WM and RWM lines.

In the case of elastic measurements, increasing steps of pressure were applied onto the same samples.
Each step lasted for 10min and images were recorded every 10s to make sure that no flow was
observed on the samples. After these 10min, the samples were released for 5min to allow them to
regain their original shape before applying a new, larger pressure difference.

For visco-elastic measurements, a single pressure step was applied and samples were imaged by
fluorescent microscopy for 30min with a 10s time step and the dynamics of the aspirated length was
monitored.

To automate the measurement of the aspirated length in these experiments, kymographs of the
aspirated tongue were generated in ImageJ (NIH), binarized and analyzed in Python with custom made
code to extract the length of the tongue in each image. For elastic measurements, the average length
over the last 5min of recording was used whereas for visco-elastic measurements, the whole dynamics
was kept.

The acquired images were also used to measure the mean projected area *A* of each analyzed sample
by usual thresholding techniques in ImageJ. This measurement was then turned into an average radius

R as $R = \sqrt{\dfrac{A}{\pi}}$.

Osmotic oscillations measurements

To measure osmotic oscillations shown in Fig S8, *Hydra* tissue spheres were prepared as described
above. In parallel, a 1% agarose gel was prepared at the bottom of a Petri dish, left to solidify and was
manually punched to create 1mm wide wells which then hosted the samples. These wells were used
to limit the samples movement during osmotic oscillations. They were then imaged for 24h at a 15min
interval. The resulting images were thresholded and analyzed in ImageJ to extract the projected area
as a function of time $A(t)$. Finally, the samples' radii $R(t)$ were computed as described above. The
equilibrium radius R_0 of a given sample was defined as the minimum of $R(t)$ over the whole recording
allowing us to estimate the strains on the inflating sphere.

Finite element simulations

Numerical simulations were performed with the finite-element solver Comsol Multiphysics (COMSOL,
Inc). The micropipette was represented as a rigid, motionless cylinder of inner radius $R_p = 50\mu m$. The
reference state of the *Hydra* sphere was a shell of outer radius $R_0 + {h_0}/{2}$ and inner radius $R_0 - {h_0}/{2}$,
whose center O was located on the cylinder's longitudinal symmetry axis Oz. We used contact
boundary conditions ensuring that the (deformed) *Hydra* sphere and the micropipette never overlap.
Hydra tissue was modeled as a hyperelastic Saint-Venant Kirchhoff material. We denote by E and ν
the Young's modulus and Poisson's ratio of the tissue and considered nearly incompressible conditions
($\nu = 0.495$). The external pressure ΔP was applied on the spherical section contained within the
micropipette. Simulations were performed in 2D, enforcing the rotational invariance of the system

(*Hydra* + micropipette) about Oz. The equations of elasticity were treated within a pressure
formulation, adapted to approximately incompressible elastic materials. For each value of the applied
pressure, we recorded the maximal deformation observed on the symmetry axis denoted $\delta z(r = 0)$,
and computed as a spatial average along the shell's thickness. We checked that our results were robust
to smoothing the sharp corner of the micropipette by a fillet over a scale of the order of $5 \mu m$. They
were also insensitive to decreasing the typical mesh size.

Extraction of rheological parameters

To measure Young's moduli in the elastic phase, each sample was treated separately. Based on the
acquired images, we measured their projected area S_0 and turned them into rest radii $R_0 = \sqrt{\frac{S_0}{\pi}}$. Then,
we used our measurements of ΔP as a function of δz (Eq 7) and fitted them by a straight line in Python.
The slope of this straight line was kept and multiplied, according to Eq 7, by $\frac{(1-\nu)R_0^2}{2C_S h_0}$ to extract the
Young's modulus of that specific sample. We then repeated the procedure on many samples and report
the mean and standard deviation of this distribution in the text.

In the visco-elastic phase, we measured the dynamics of the aspirated length as a function of time $L(t)$
and fitted them in Python according to Eq 8. To extract Young's moduli, we used the same formula as
above except that we replaced the slope of a linear fit of several points by the ratio of ΔP over $\delta(1 -$
$\kappa)$, δ and κ stemming from the fit of the full dynamics.

To estimate the order of magnitude of the effective viscosities in the same experiments, we used the
same fit according to Eq 8 to extract the speed of tissue flow U and estimated the viscosity as $\eta = \frac{\Delta P R_p}{U C_v}$
with $C_v = 1$.

Measurement and data analysis of threshold pressure measurements.

For the measurement of the pressure thresholds distinguishing each regime, we used the
fraction of pieces changing their rheological behavior at different applied pressures. To observe the
switch between the elastic and viscous regimes the pressure was gradually increased by lowering the
syringe in 3cm increments, starting low enough that no viscoelastic behavior was observed (usually 10
or 14cm). At each pressure level and after 20 minutes, we noted the fraction of pieces that had
changed regime, and the experiment was stopped when all samples had done so. Kymographs were
used to visually separate both regimes. The length profile over time was constant in the elastic regime,
whereas it evolved linearly with a clear slope in the viscous regime.

The same strategy was used to measure the rupture threshold, but we started at higher applied
pressures. Images were taken in transmission, as it facilitated the observation of briefly detached
pieces of tissue in the holes of the inserts, the first sign of the rupture regime. They were also taken
this time at a rate of ten minutes of suction per threshold. Identifying the switch between the viscous
and the rupture regime was done visually.

In both cases, we thus ended up with a fraction of samples adopting a certain behavior as a function
of applied pressure. Assuming that individual thresholds are variable and normally distributed, this
cumulative frequency followed a sigmoid shape of the form $\frac{1}{1+e^{-(\Delta P - \Delta P_c)/\alpha}}$ where ΔP_c is the threshold
pressure at which 50% of samples have switched behavior and α is a parameter controlling the
steepness of the sigmoid function. We thus fitted our data with that functional form using Matlab's
(Mathworks) curve fitting toolbox. In the manuscript, we report threshold pressures as ΔP_c resulting
from the fits and error bars represent the 95% confidence interval of these fits.

EDTA/blebbistatin experiments

Rheological measurements on *Hydra* whose cell adhesion had been inhibited were carried out by immersing the tissue spheres in solutions of either 2 mM EDTA (Sigma-Aldrich) or 5μm blebbistatin (Tocris Bioscience) in HM. The entire microfluidic system was immersed in the solution, and the *Hydra* pieces were left to bathe in it for around ten minutes before aspiration to give the drug time to take effect before measuring. The rest of the protocol was identical to that of the rheological measurements explained above.

Spinning disk imaging

To estimate the thickness of *Hydra* tissue spheres, several WM samples were imaged underneath an Eclipse Ti2 microscope (Nikon) equipped with a CSU-W1 spinning disk unit (Yokogawa) and an Orca Fusion BT camera (Hamamatsu). Z-stacks were automatically acquired with a step of 1 micron and vertical slices were made using ImageJ (Fig S4). The thickness of six samples were manually measured using the same software and were all found to be in the $15 - 25\mu m$ range.

Mechanical description of *Hydra* tissue spheres

We consider a thin spherical shell of undeformed inner and outer radii $R_0 - \frac{h_0}{2}$ and $R_0 + \frac{h_0}{2}$. Normal components of the Cauchy stress tensor are denoted σ_a ($a = 1,2,3$). Briefly, we re-express the strain energy function, Eq 4, of a Saint-Venant Kirchhoff material as a function of the stretch ratios α_a: $W(\bar{\bar{E}}) = \Psi(\{\alpha_a\})$. Denoting P the shell pressure field, stress components are given by $\sigma_a = -P + \alpha_a \frac{\partial \Psi}{\partial \alpha_a}$. By the assumption of plane stress, $\sigma_3 = 0$, we obtain the expression of P and deduce the circumferential, or hoop stress $\sigma = \sigma_1 = \sigma_2$. For an incompressible material, $4\pi R_0^2 h_0 = 4\pi R^2 h$, where R and h denote the deformed radius and thickness, respectively. We note $\alpha = \alpha_1 = \alpha_2 = \frac{R}{R_0}$ the circumferential stretch ratios, and deduce the normal stretch ratio $\alpha_3 = \frac{h}{h_0} = \frac{1}{\alpha^2}$. The calculation follows (37, 38) and yields the expression of the hoop stress of a Saint-Venant Kirchhoff spherical shell, as given by Eq 11.

Results

Multiplexed micro-aspiration setup

In order to define and measure the rheological properties of *Hydra* tissue spheres, we settled on the use of micro-aspiration experiments. These experiments are now well-established, robust and measure rheological properties at length scales and frequencies relevant for the regenerative process. In contrast with Atomic Force Microscopy (AFM), previously used to locally probe the stiffness of adult *Hydras* (39), our experiments were particularly designed to study the mechanics of tissue spheres during the osmotic oscillations which is why we employed elongations rather than indentations and probed the large deformation regime. Their main drawback is their intrinsically low throughput. In their original form, micro-aspiration experiments can only probe one sample at a time with a single experiment running for around 1h (4, 33). We thus started by developing and adapting a microfluidics device making use of removable inserts (36) to parallelize the experiment, in the same spirit as the technique developed in (40) but with circular pipettes which prevent singularities and leakage. This device was composed of two different objects. First, a microfluidic channel was designed to host the samples, it was 3cm in length and 500μm in height (Fig 1A, Fig S1). Halfway through this channel and perpendicular to it sat another channel meant to host the removable insert. This channel was 500μm wide and 1mm high. This whole construct was then manufactured in polydimethylsiloxane (PDMS).

Most of the experiments presented here were carried out with removable inserts, microfabricated in
dry film, bearing either 6 or 10 circular openings, 500µm or 300µm away from one another, 300µm in
length and 100µm in diameter. Of note, these holes sat, vertically, at the midpoint of the samples'
channel (250µm, Fig 1B). Tissue spheres aspirated in these holes therefore didn't touch either the
bottom or the top of the PDMS channel which had no influence on the aspiration.

Fig 1. Parallelized micro-aspiration experiments. A: Schematics of the mold for the PDMS channel (details on dimensions can
be found in Fig S1). B: Representation of the principle of the experiment, the insert, in orange, effectively creates an array of
cylindrical tunnels, equivalent to model micropipets. The tissue spheres, in yellow, flow towards these openings and, once
sealed, get aspirated within. C: Snapshot of a mounted channel with an insert containing ten tunnels in parallel. D:
Fluorescent imaging of the ectoderm of six samples aspirated within the tunnels.

Using hydrostatic pressure, we applied a difference in pressure between the inlet and outlet of the
main chamber (with $P_{in} > P_{out}$, see Fig S2) which initially created a flow of water from the Petri dish
to the outlet of the chamber. Previously prepared *Hydra* tissue spheres were then manually pipetted
into the entrance of the main channel. Thanks to the flow of water, the samples naturally aligned with
the openings in the insert. Once all openings were blocked by a sample, the flow of water stopped and
aspiration of the tissue spheres was observed (Fig 1D, Movie S1). Of note, in most experiments, the
pressure difference was set to 0 as soon as the samples aligned with the holes so that a controlled step
in pressure could be applied thereafter.

Rheological behavior depends on applied stress.

Using this multiplexed setup, we started by studying the response of newly formed *Hydra* tissue
spheres to micro-aspirations in the kPa range using circular openings with a radius $R_p = 50µm$. We

observed two different rheological responses depending on the applied pressure. Up to around 2.5kPa,
tissue spheres exhibited an elastic response in the sense that for a given applied pressure difference,
a tongue of tissue was aspirated within the holes which length increased in the first tens of seconds
but then saturated and remained constant over much longer time scales. When the pressure was
released, we also observed that the samples retrieved their original shape, and that aspiration and
release dynamics were very similar (Fig S3), consistent with the reversible behavior of an elastic
material. To further make sure that we were not missing a viscous behavior at longer time scales, we
managed to trap tissue spheres for tens of hours in this elastic phase without any indication of tissue
flow (Movie S2).

However, when we increased the applied pressure, we started observing samples flowing inside the
holes, a signature of a viscous behavior. This demonstrates a real change in mechanical behavior of the
tissue spheres akin to the behavior of yield stress fluids and requiring two different experimental and
theoretical approaches.

Finally, at even higher applied stresses, we started observing rupture of the tissue and cell
detachments, reminiscent of the rupture observed under normal phase I osmotic oscillations (Movie
S3). Since all three behaviors seemed relevant to understand the mechanical behavior of normally
regenerating tissue spheres, we decided to characterize them all, as well as the pressure thresholds
between these different phases.

Elastic behavior at low applied stresses.

To characterize the elastic behavior of the tissue spheres, we applied steps of pressure on the same
samples and recorded, for each step and for each sample in an experiment, the length of the aspirated
tongue L (Fig 2A). We monitored the samples closely to make sure they did not start flowing in the
holes, in which case their analysis was stopped. As expected, we found that the higher the pressure,
the longer the aspirated tongue (Fig 2A). To estimate a Young's modulus from these measurements,
we needed a rheological model of an elastic spherical shell of mean radius R_0 and thickness h_0.

As is the case for cell aggregates, we expect that intercellular adhesion and cellular cortical tension will
contribute to an effective surface tension of the *Hydra* tissue spheres, which we denote by γ.

A pressure difference ΔP through the micropipette of radius R_p generates an aspiration force $f_p =
\pi R_p^2 \Delta P$. For an aspirated length L, the free energy F of the aspirated shell reads:

$$F = 2\gamma S - f_p L \qquad (1)$$

Where S denotes the surface area of the sample and the factor 2 takes into account the presence of
two interfaces. Assuming incompressibility, the volume V of the sample is constant. For small enough
deformations during aspiration, we expected that the thickness h_0 of the sample would remain
constant. To validate this hypothesis, we performed spinning disk confocal microscopy on normal
tissue spheres and aspirated ones to reconstruct their 3d structure and estimate their thicknesses. We
found $h_0 = 20 \pm 5\mu m$ in both cases and didn't observe any clear change due to micro aspiration (Fig
S4). Since $V = Sh_0$, the surface area of the sample also remains constant $dS = 0$. In other words, in
the case of the small deformations of an incompressible shell of approximately constant thickness,
tissue surface tension does not contribute to the total force f exerted on the sample:

$$f = -\frac{dF}{dL} = -2\gamma \frac{dS}{dL} + f_p = f_p = \pi R_p^2 \Delta P \qquad (2)$$

Indeed, our experiments do not exhibit a clear half-sphere of radius R_p, typical of the early entry of a
full cell aggregate into a micropipette (41). In addition, (33) elegantly demonstrates that surface
tension manifests itself as a critical pressure below which no aspiration is observed and which needs
to correct all subsequent applied pressures. The fact that we observed aspirations with applied
pressures as low as 200Pa also indicates that surface tension can largely be ignored in our experiments.

The rest state of the system is simply that of the initial spherical shell of radius R_0 and due to that
initial curvature, the displacement of the tip of the tissue tongue inside the micropipette δz reads:

$$\delta z = L - \left(R_0 - \sqrt{R_0^2 - R_p^2}\right) \qquad (3)$$

As expected for an elastic material, we found a linear relationship between this displacement and the
applied pressure ΔP (Fig 2B). We also noticed that for many samples, our linear fits had negative
intercepts at the origin. This could be the signature of a stress-stiffening effect by which the effective
Young's modulus would increase with the applied pressure, *i.e.* the existence of non-linear elastic
behavior, as observed in suspended cell monolayers (42).

To take into account this observation, we turned to a nonlinear, hyperelastic description of the *Hydra*
tissue, using, as previously advocated by (43), the Saint-Venant Kirchhoff model, defined by the strain
energy function:

$$W(E) = \frac{\lambda}{2} Tr(\bar{\bar{E}})^2 + \mu Tr(\bar{\bar{E}}^2) \qquad (4)$$

where λ and μ denote the Lamé coefficients, and $\bar{\bar{E}}$ is the Green-Lagrange strain tensor (37). This
equation reduces to its linear elastic counterpart for small deformations, when $\bar{\bar{E}} \approx \bar{\bar{\varepsilon}}$ with $\bar{\bar{\varepsilon}}$ the
infinitesimal strain tensor. Of note, the Saint-Venant Kirchhoff model is defined by the same
parameters as would be the case for a linear elastic material, with the following classical relationships
between Lamé coefficients and other elastic moduli: E and v, respectively, the Young's modulus and
the Poisson's ratio of the sample.

$$E = \mu \frac{3\lambda + 2\mu}{\lambda + \mu} \qquad (5)$$

$$v = \frac{\lambda}{2(\lambda+\mu)} \qquad (6)$$

430 In analogy with the behavior of linearly elastic thin spherical shells (see for instance (44)), and taking
431 into account the non-zero intercept ΔP_0 of the linear fits of our data (Fig 2C, $\Delta P_0 =$
432 $(-6.4 \pm 1.3).10^2 \, Pa$), we expect that the displacement and the applied pressure should be related
433 according to:

$$\Delta P = C_s \frac{2Eh_0}{(1-v)R_0^2} \delta z + \Delta P_0 \qquad (7)$$

Where C_s is a dimensionless, geometry-dependent pre-factor. Of note, in this description, we ignore
the complex composite structure of the shell which includes two different epithelial layers as well as
the extra-cellular matrix.

In order to estimate C_s, we performed numerical simulations of micro-aspiration of hyperelastic
spherical shells by a single cylindrical tunnel (Fig 3A). In our experimental data, we had $R_0 = 162 \pm$
$28\mu m$ (mean ± standard deviation, n=36) and $h_0 = 20 \pm 5\mu m$. As we had no measurement of the
Poisson's ratio of our samples, we made an assumption of near incompressibility and set $v = 0.495$.
Based on these values, we submitted spherical shells to applied pressure ΔP and measured the

resulting δz. As shown in Fig 3B, these simulations confirmed the linear relationship between ΔP and
δz over a similar range of applied pressures and displacements and allowed to estimate C_s thanks to
the knowledge of the value of E and ΔP_0. Of note, other element-based numerical simulations were
performed with parameters varying in the following ranges: $25 \leq \Delta P \leq 2000 Pa$, $15 \leq h_0 \leq 25 \mu m$
,$130 \leq R_0 \leq 200 \mu m$, $200 \leq E \leq 750 Pa$ and with a fillet size varied between 2 and $10 \mu m$. Overall,
our estimate of the coefficient C_S reads $C_S = 23.4 \pm 4.3$, in the vicinity of 23, suggesting that the order
of magnitude of our results was unaffected by uncertainties on these different parameters. Further,
our estimate of the intercept in these simulations is $\Delta P_0 = (-8.5 \pm 2.1).10^2 \, Pa$, consistent with the
above experimental values.

Fig 3. Numerical simulations. A: Graphical representation of the deformed shell under. The color code represents the z-
component of the displacement field. Black lines delineate the reference state of the system. Parameter values for these
simulations are $R_0 = 160 \, \mu m$, $h_0 = 20 \, \mu m$, $E = 440 \, Pa$, $\nu = 0.495$, $\Delta P = 10^3 Pa$. B: applied pressure Δp versus the
displacement δz for the same parameter values. The slope of the fitted line, in orange, corresponds to a value $C_S = 23.4$.

Thanks to our multiplexed setup, we were able to measure up to six samples in parallel in one sitting.
By then repeating these experiments eight different times, we achieved a total of 36 tissue spheres
(Fig 2C) characterized in that phase. For each of them, we extracted the slope of the linear relationship
between ΔP and δz, its initial radius R_0 and finally obtained a quantitative measurement of its Young's
modulus using Eq 7 (Fig 2D). From the above estimate of the coefficient C_S, we found the Young's
modulus of *Hydra* tissue spheres to be $(4.4 \pm 3.3).10^2$ Pa (mean ± standard deviation, n=36).

We also observed a large variability in our measurements of individual Young's moduli (Fig 2D) which
could hint at subtle effects missed by our averaging approach. The different tissue spheres could differ
in initial size or degree of inflation since we do not precisely control the time they need to fold back
into tissue spheres after being cut. To better understand this variability, we looked at the correlation
between the samples' radii and their extracted elastic moduli. We indeed found a positive correlation,
albeit weak, between these measurements (Fig S5). We then asked whether this correlation could be
due to differences in inflation state. To do so, we ran new experiments measuring the elastic moduli
of tissue spheres whose oscillations were blocked by the addition of 70mM sucrose in the outside
medium. Doing so, we found 1- that the obtained elastic moduli distributions in both cases were not
statistically different (Fig S6) and 2- that addition of sucrose abolished neither the weak positive
correlation between size and elasticity (Fig S5) nor the existence of stiffer samples. We conclude that
some, but not all, of the observed variability comes from differences in initial size rather than inflation
rate between different samples. A better understanding of this dependency would be a natural
continuation of this work.

In addition, we validated the relevance and precision of our experimental procedure by repeating
elastic measurements under two chemical modulations: EDTA and blebbistatin. EDTA is a chemical
well-known for weakening cell-cell adhesions in many epithelial tissues (45) whereas blebbistatin

inhibits acto-myosin contractility (46). We expected that both treatments should lower the samples'
elasticity either by allowing larger tissue-scale deformations in the case of EDTA or cell-scale
deformations in the case of blebbistatin by making the actin cortex of each cell softer. We indeed found
a clear and statistically significant decrease in Young's modulus with both treatments (Fig S6, EDTA:
E=(3.0 ± 2.0).10^2 Pa, blebbistatin: E=(1.9 ± 0.7).10^2 Pa), confirming that the observed elastic behavior
of the tissue spheres stems from a complex interplay of single cell mechanics and tissue structure.

Overall, we characterized in a quantitative way, the response of *Hydra* tissue spheres to applied
stresses and found them to behave hyperelastically, *i.e.* as soft ($E{\sim}500Pa$) elastic materials at small
deformations, in line with AFM measurements on adult *Hydras* (39), while they become increasingly
stiffer as the applied pressure increases.

Visco-elastic behavior at intermediate applied stresses

We then turned our attention to the next observed behavior which involved flowing of the samples
within the tunnels. This visco-elastic behavior has long been observed and studied, both on single cells
(32, 47, 48) and multi-cellular spheroids (4, 33, 41). We followed the usual approach in that situation
where a single, controlled pressure step was applied in the device at t=0 after the samples were already
placed close to the entrance of the tunnels. The dynamics of the aspirated length $L(t)$ was then
recorded by fluorescent microscopy for tens of minutes (Fig 4A). It was important to keep recording
on these longer time scales in order to be certain of the behavior adopted by each sample as a very
slow flow could easily be misinterpreted as an elastic response if the observation time scale was too
short.

Fig 4. Visco-elastic flow at large applied pressure. A: snapshots of a tissue sphere showing flow at a constant applied
pressure. B: four examples of typical dynamics of the flow as a function of time. Dots of different colours represent four
independent samples and solid lines fits by Eq 8. C: distribution of measured viscosities as black dots with box-whisker plot in
orange. We found it to be (2.4 ± 1.2).10^6 Pa.s (mean ± standard deviation, n=22).

Since the behavior is qualitatively similar to that observed for cellular aggregates aspirated in
cylindrical micropipets, we first applied the well-known modified Maxwell model developed in (33).
Briefly, a dashpot representing tissue scale viscosity is combined in series with a standard linear solid
unit incorporating cell scale viscoelasticity as well as tissue scale elasticity. We fitted our data with the
form of $L(t)$ predicted by this model:

$$L(t) = \delta\left(1 - \kappa e^{\frac{-t}{\tau}}\right) + Ut \qquad (8)$$

Where δ is linked to the short-term elastic deformation, κ is a constant linked to the relative
contributions of the two springs in the standard linear solid unit, τ is the cell scale visco-elastic
relaxation-time and U is the long term viscous flow rate. We found that this functional form was also
relevant to describe our data quantitatively (Fig 4B) despite the morphological differences between
our samples (empty shells) and ball-shaped aggregates (full spheres). We could however employ a
scaling argument similar to the one applied in this model to estimate an order of magnitude of the
effective viscosity of our samples. As for cellular aggregates, friction between the tissue and the
tunnels walls can be neglected since the long-term aspirated length scales not as the square-root of
time, but as a linear function of time. Assuming that dissipation is due to cell rearrangements at the
entrance of the tunnels thus acting on a length scale of order R_p, the total dissipative force can
therefore be approximated as $f_{viscous} \approx C_v R_p U \eta$ with η the viscosity of the samples and C_v a
dimensionless, geometrical pre-factor equal to 3π for a ball-shaped aggregate and unknown in our
geometry. This viscous force has to be balanced by the aspiration force which we have shown to be
$f_p = \pi R_p^2 \Delta P$ which leads to:

$$\eta \approx \frac{\Delta P R_p}{U C_v} \qquad (9)$$

To the best of our knowledge the computation of C_v has never been done for spherical viscous shells
and in the same geometry as the one in our experiments. We will see below that a precise
measurement of this effective viscosity has little impact on the description of *Hydra* osmotic
oscillations and we limited ourselves to obtaining an order of magnitude estimation of the viscosity
by taking $C_v = 1$. Note that assuming that tissue scale dissipation acts across the thickness h of this
multilayered tissue does not change our estimate as h and R_p have the same order of magnitude.
Again, our microaspiration setup allowed us to efficiently gather data on multiple samples and
combining them, we found an estimation of the viscosity of $(2.4 \pm 1.2).10^6$ Pa.s (mean $\pm$ standard
deviation, n=22, Fig 4C).

The full dynamics of aspiration in that phase can also yield measurements of the Young's modulus of
the samples. In line with our analysis of the purely elastic response, we expect here that:

$$\Delta P = C_s \frac{2Eh}{(1-\nu)R_0^2}\delta(1-\kappa) \qquad (10)$$

Similar to Eq. 7 with $\delta(1 - \kappa)$ the immediate elastic aspirated length in the notation of this model.
Applying this relationship to the results of our fitted data, we found $E = (3.7 \pm 1.6).10^2 Pa$ (Fig S7),
in accordance with our results on the purely elastic behavior of *Hydra* tissue spheres.

These visco-elastic experiments were made difficult by the fact that in many instances, we observed,
in addition to the flow of the samples, cells detaching at the tip of the aspirated tongue, indicative that
the tissues were already rupturing. These cases were obviously discarded from the analysis, limiting
the amount of available data. Overall, it appeared that at these applied pressures, the visco-elastic
phase and rupture were often occurring together.

Critical pressures between different behaviors

To further study this question, we then turned our attention to the critical pressures required to
transition between the three regimes: elastic, viscoelastic and rupture. We often found cases where,
for a given applied pressure, a fraction of the samples loaded in the same setup would behave as elastic
solids while other behaved as viscoelastic fluids (Fig 5A). To characterize the first critical pressure, we
therefore measured, at different applied pressures, the fraction of samples showing a viscoelastic
versus elastic behavior while discarding rupturing ones. We fitted this fraction as a function of the
applied pressure by a sigmoid (see Materials and Methods, Fig 5B). We defined the critical pressure as
the center of this sigmoid where 50% of samples had switched from one behavior to the other.
Following this method, we found the critical pressure between elastic and viscoelastic behaviors to be
2.22 ± 0.14 kPa (error bars represent the 95% confidence interval of the sigmoid fit). We then employed
the same strategy to characterize the critical pressure leading to tissue rupture (Fig 5C). We found it
to be 2.37 +- 0.08 kPa, very close to the previous one confirming our intuition that both behaviors were
linked together and that tissue flows inside the holes were already a signature of their lack of integrity.

Fig 5. Critical stresses between different mechanical behaviors. A: snapshot of an experiment showing two similar samples
displaying different behaviors at the same applied pressure. B: Quantification of the fraction of viscous behavior at different
applied pressures. C: quantification of the fraction of samples showing tissue rupture as a function of applied pressure. In B
and C, dots are data and solid lines fit by a sigmoid function. In black are control samples and in red samples pretreated with
2mM EDTA. Each point is derived from 18 to 40 samples stemming from 8 different experiments in B and 4 in C. D:
Numerical simulations. The color code represents the hoop stress within the aspirated shell when aspirated at a pressure of
1kPa. Parameter values are the same as in Fig. 3.

The most natural explanation for this lack of integrity is the weakening and rupture of cell-cell contacts
since we often observed detachment of single cells, making a possible deadhesion of the tissues to the

extracellular matrix unlikely. To verify this hypothesis, we repeated our measurement of the critical
pressures required to induce either a viscous behavior or tissue rupture on tissue spheres pre-treated
with 2mM EDTA as we expected this treatment to facilitate the induction of a viscous behavior and
tissue rupture. This is indeed what we found with a shift of the critical pressure from 2.22 kPa to 1.19
± 0.16 kPa for the viscous behavior (Fig 5B) and from 2.37 kPa to 1.29 ± 0.19 kPa for rupture (Fig 5C).

It is important to note that all these values are difference of pressure applied between the two sides
of the holes and not necessarily the stress actually acting in the tissue called the hoop stress. Said
differently, the applied pressure required to rupture the tissue shouldn't be directly taken as a
measurement of the ultimate tensile stress of the whole tissue. To relate the applied pressure and
internal tissue stresses, we turned again to numerical simulations. Within the range of parameters
studied, the maximal values of the hoop stress we found was on the order of a 2-7 kPa (see Fig 5D for
an example).

Since in normal conditions, the threshold for viscous behavior cannot be clearly distinguished from the
threshold for rupture, our measurements revealed that up to rupture, which is also observed in normal
osmotic oscillations, regenerating *Hydra* tissue spheres behave as hyperelastic spherical shells with a
stress stiffening effect.

Mechanical description of oscillating Hydra tissue spheres during regeneration.

After folding back into a spherical shape, regenerating *Hydra* tissue spheres experience a series of
swelling-rupture cycles which have been shown to be driven by the difference in osmolarity between
the inside and outside of the spheres. These oscillations are required for the proper morphogenesis
and regeneration of the samples although their exact role remains unknown. Thanks to previous work
on the topic (5, 49) and our own mechanical characterization, we can provide a good understanding of
the internal tissue mechanics occurring during this phase.

The key quantity is the circumferential or hoop stress, noted here σ, which is known to control tissue
integrity and to be able to alter key biological processes. It is given for an incompressible, thin elastic
spherical shell of a Saint-Venant Kirchhoff material (see (37, 38) and the Methods section) by the
expression:

$$\sigma = (\alpha^4 - \frac{1}{\alpha^2})\left(\left(\frac{\lambda}{2} + \mu\right)\left(2 + \frac{1}{\alpha^6}\right) - \left(\frac{3\lambda}{2} + \mu\right)\frac{1}{\alpha^2} - \mu\right) \qquad (11)$$

599 With $\alpha = \frac{R(t)}{R_0}$ is the stretch ratio, and λ and μ the Lamé coefficients. This gives the nonlinear
600 constitutive relationship between stresses and strains in the spherical shell during the oscillations
601 which allows to directly extract the hoop stress from a measurement of the radius of the sample as a
602 function of time (see Fig S8). One can check that Eq 11 reduces to the relationship expected of a linear
603 elastic thin shell in the limit of small deformations where $|R - R_0| \ll R_0$:

$$\sigma = \frac{E}{1 - \nu} \cdot \frac{R - R_0}{R_0} \qquad (12)$$

At larger deformations, the tissues get stiffer and the experimentally measured values of E imply that,
in this regime, the differential elastic modulus $\frac{\partial \sigma}{\partial \alpha}$ is on the order of 10kPa, in accordance with previous
estimates of the sphere's stiffness during osmotic oscillations (5).

Based on our experimental observations, we will approximate our samples as hyperelastic, symmetric,
spherical shells of time-dependent thickness, $h(t)$ and radius $R(t)$ (Fig S8) submitted to a difference
in osmolite concentrations $(C_{in} - C_{out}) > 0$. The fact that the spheres are constantly swelling
indicates a constant influx of water into their lumen and that the system never reaches osmotic
equilibrium. Furthermore, Kücken et al. (5) showed that the increase of $R(t)$ can be described by a
Darcy-type law of the form:

$$\frac{dV}{dt} = 4\pi R(t)^2 L_w k_b T (C_{in} - C_{out}) \qquad (13)$$

Where $V(t)$ is the volume of the sphere, L_w is the water permeability coefficient, k_b is Boltzmann's
constant. The dynamics of swelling is thus entirely controlled by the influx of water which imposes a
constantly increasing deformation on the tissue sphere with a growth rate of the radius on the order
of $5\mu m/h$. This influx of water effectively inflates the sphere and creates a restoring elastic pressure
P_{el} which our mechanical description can also predict. Following the mechanics of spherical thin shells,
this pressure can be written as:

$$P_{el}(t) = \frac{2h(t)}{R(t)} \sigma(t) \qquad (14)$$

Using the experimental measurement of E we previously obtained and the same value of the Poisson's
ratio, $\nu = 0.495$, we were able to approximate the values of λ and μ ($\mu = 148Pa, \lambda = 100\mu$). Using
these values and experimental measurements of $\alpha(t)$ during the oscillations allowed us to estimate
$\sigma(t)$ and we found it to be on the order of a few kPa (Fig S8). Most notably, this approach allowed us
to estimate the critical stress, within the tissue, to induce rupture and we found it to be on the order
of 4-6kPa. This is the same order of magnitude as the maximum hoop stress obtained by numerical
simulations of the elastic deformation during micropipette aspiration. Overall, we demonstrate that
our results allow to map measurements of the radius of the tissue spheres, which are experimentally
easy to access, into both internal stresses within the tissue and restoring elastic pressure acting on the
fluid inside of the spheres.

Discussion

The effect of stretching on the biology of epithelial tissues has emerged in recent years as an important
topic in developmental biology since developing organisms often undergo substantial deformations
which have an increasingly well recognized impact on their biological response. In the context of *Hydra*
regeneration, Ferenc et al. observed that the levels of expression of Wnt ligands, especially HyWnt3,
are directly linked to the cumulative amount of stretching in the tissue (20). In a similar fashion,
stretching adhering monolayers of epithelial cells has been shown to induce cell division through the
activation of Piezo1 mechanosensitive ion channels (22) to restore proper cell density. Anisotropic
deformations also have a large impact on the tissue as uniaxial cyclic stretch leads to re-orientation of
the cells perpendicular to the direction of application (50, 51), as was observed for the mitotic spindle
of single cells (52). In many of these experiments, however, stresses within the tissues couldn't be
recorded after stretching.

One notable exception was achieved on freely suspended epithelia axially stretched with a measurable
force (42). This experimental setup allowed the authors to characterize the rheology of Madine–Darby

Canine Kidney (MDCK) monolayers. Interestingly, they observed a non-linear response with three distinct phases: a linear one at small deformations with a small equivalent elastic modulus, a second linear one for intermediate values of stresses and strains with much higher stiffness and a third one corresponding to the rupture of the tissue. This behavior is very similar to the one we observed in this work. The authors quantified the second phase by linear fitting of the stress-strain curves with negative intercepts, as we did. Since they approximated the tissue in this phase as linearly elastic, they found a high elastic modulus on the order of 20kPa, similar to our observations of *Hydra* tissue spheres differential moduli at intermediate stresses. Finally, the authors found that the monolayer was able to withstand deformations up to 60% before rupturing, a value slightly higher than what we observed for *Hydra* tissue spheres. Our results are thus well in line with these observations, and both ask the question of the biological origin of the non-linear behavior of epithelial tissues which remains elusive for the moment.

Regarding the specific rheology of *Hydra*, our work is the first to focus specifically on regenerating tissue spheres but measurements exist on full adults. In (39), the authors employed a modified AFM experimental procedure to generate indentations on fixed, full *Hydras* with a 20μm wide sphere. By co-recording the deformation of the tissue and the force exerted, they observed a linear relationship between both which allowed them to estimate the elastic modulus of these samples. One key result was the observation that the elasticity of the body column was inhomogeneous, the tissue being roughly three times stiffer near the head than close to the foot. Although it is a well-known fact that different experimental approaches can lead to different estimations of rheological properties (53) and potential differences between fully mature *Hydras* and tissue spheres, our results are in good agreement with this work. Indeed, in the softest part of the adult, which we would imagine to most resemble tissue spheres, the authors report an elastic modulus on the order of 400Pa. Since the deformations involved are of a few microns, we expect these measurements to probe the small deformation regime where our own measurements also find an elastic modulus on the order of 400Pa. In the large deformation regime, our results also confirm those of (5) which, in order to properly reproduce the osmotic oscillations based on chemical considerations implemented an elastic modulus $E = 11kPa$. Still, AFM measurements on tissue spheres, albeit technically challenging, would be a great complement to our work to obtain a comprehensive understanding of the mechanics of these samples.

Another consequence of our focus on osmotic oscillations and tissue-scale mechanics is that our mechanical description of the tissue spheres assumes a homogenous shell, ignoring the complex three-layer (endoderm-extra-cellular matrix-ectoderm) structure of the samples. It is clearly possible that the deformations and stresses induced by the osmotic oscillations are differently transmitted and have different effects on these different layers, a refinement which will need to be addressed in the future. In particular, it was recently suggested that the mechanical properties of the ECM, which were not specifically measured here, could be linked to morphogenesis in *Hydra* (54).

Since we believe deformations and stresses within the tissues to be central in the morphogenesis of *Hydra*, another natural question is the existence of inhomogeneities of deformations, forces or rheological properties within the same sample. Mechanical fluctuations, if coupled to biochemistry, could well be sufficient to destabilize a reaction-diffusion system into a patterned state. Our own experimental setup could allow to measure fluctuations in rheological properties by repeating measurements on the same samples with small holes but at different locations, the main limitation being the lack of control of the position of the tested zone. In the same logic, it has been proposed that the amount of HyWnt3 protein could locally affect the Young's modulus of the tissue (25) which could couple the chemical patterning of the head organizer and morphogenesis. If the future location

of the head becomes softer, the homogenous osmotic pressure will deform this region more, leading
to a deviation from a spherical shape to an oblong one. Here too, our setup could allow to measure
the Young's modulus of tissue spheres overexpressing HyWnt3 to directly test this hypothesis.

Conclusion

In this work, we have employed an original parallelized microaspiration device to characterize the
mechanical behavior of *Hydra* tissue spheres. Although we found different rheological behaviors
depending on the applied stresses, the main result we obtained is that during normal osmotic
oscillations, these samples behave largely as hyperelastic thin shells, displaying a nonlinear, stress-
stiffening rheology. Doing so, we also obtained quantitative measurements of their relevant
rheological parameters ($E = (4.4 \pm 3.3) . 10^2$ Pa, $\eta = (2.4 \pm 1.2) . 10^6$ Pa.s, critical applied
pressures for viscous behavior and tissue rupture). This allowed us to estimate the stresses acting
within the tissues and, in particular, the critical stress at which they start rupturing. These results, by
allowing a quantification of both stresses and deformations during *Hydra* regeneration, will hopefully
pave the way for quantitative approaches aiming at correlating these mechanical cues with molecular
ones to understand *Hydra* patterning as an integrated mechano-biochemical process.

Author contributions

OCE designed the research. ZBM, MD and PJ manufactured the inserts. TP, ABB, PM and OCE
performed the research and analyzed data. TP, PM and OCE wrote the manuscript which was
reviewed by all authors.

Declaration of interests

The authors have declared no conflict of interest.

Acknowledgments

This work was funded by the Mission pour les Initiatives Transverses et Interdisciplinaires of CNRS
(Project MeChemReg to OCE and PM). We thank H. Delanoë-Ayari for fruitful discussions.

References

1.   Galliot, B., and V. Schmid. 2002. Cnidarians as a model system for understanding evolution
and regeneration. *Int. J. Dev. Biol.* 46:39–48.

2.   Technau, U., and R.E. Steele. 2011. Evolutionary crossroads in developmental biology:
Cnidaria. *Development*. 138:1447–58.

3.   Gierer, A., S. Berking, H. Bode, C.N. David, K. Flick, G. Hansmann, H. Schaller, and E. Trenkner.
1972. Regeneration of hydra from reaggregated cells. *Nat. New Biol.* 239:98–101.

4.   Cochet-Escartin, O., T.T. Locke, W.H. Shi, R.E. Steele, and E.-M.S. Collins. 2017. Physical
Mechanisms Driving Cell Sorting in Hydra. *Biophys. J.* 113:2827–2841.

5.   Kücken, M., J. Soriano, P.A. Pullarkat, A. Ott, and E.M. Nicola. 2008. An osmoregulatory basis
for shape oscillations in regenerating hydra. *Biophys. J.* 95:978–85.

6.   Fütterer, C., C. Colombo, F. Jülicher, and A. Ott. 2003. Morphogenetic oscillations during
symmetry breaking of regenerating Hydra vulgaris cells. *Europhys. Lett.* 64:137.

7.   Broun, M., L. Gee, B. Reinhardt, and H.R. Bode. 2005. Formation of the head organizer in
hydra involves the canonical Wnt pathway. *Development*. 132.

8.   Bode, H.R. 2009. Axial Patterning in Hydra. *Cold Spring Harb. Perspect. Biol.* 1.

9.   Böttger, A., and M. Hassel. 2012. Hydra, a model system to trace the emergence of
boundaries in developing eumetazoans. *Int. J. Dev. Biol.* 56:583–91.

10. Soriano, J., C. Colombo, and A. Ott. 2006. Hydra molecular network reaches criticality at the
symmetry-breaking axis-defining moment. *Phys. Rev. Lett.* 97:258102.

11. Turing, A.M. 1952. The Chemical Basis of Morphogenesis. *Philos. Trans. R. Soc. B Biol. Sci.*
237:37–72.

12. Marcon, L., and J. Sharpe. 2012. Turing patterns in development: What about the horse part?
Curr. Opin. Genet. Dev. 22:578–584.

13. Schweisguth, F., and F. Corson. 2019. Self-Organization in Pattern Formation. *Dev. Cell*.
49:659–677.

14. Gierer, A., and H. Meinhardt. 1972. A theory of biological pattern formation. *Kybernetik*.
12:30–9.

15. Nakamura, Y., C.D. Tsiairis, S. Özbek, and T.W. Holstein. 2011. Autoregulatory and repressive
inputs localize Hydra Wnt3 to the head organizer. *Proc. Natl. Acad. Sci. U. S. A.* 108:9137–42.

16. Augustin, R., A. Franke, K. Khalturin, R. Kiko, S. Siebert, G. Hemmrich, and T.C.G. Bosch. 2006.
Dickkopf related genes are components of the positional value gradient in Hydra. *Dev. Biol.*
296:62–70.

17. Vogg, M.C., L. Beccari, L. Iglesias Ollé, C. Rampon, S. Vriz, C. Perruchoud, Y. Wenger, and B.
Galliot. 2019. An evolutionarily-conserved Wnt3/β-catenin/Sp5 feedback loop restricts head
organizer activity in Hydra. *Nat. Commun.* 10:312.

18. Ziegler, B., I. Yiallouros, B. Trageser, S. Kumar, M. Mercker, S. Kling, M. Fath, U. Warnken, M.
Schnölzer, T.W. Holstein, M. Hartl, A. Marciniak-Czochra, J. Stetefeld, W. Stöcker, and S.
Özbek. 2021. The Wnt-specific astacin proteinase HAS-7 restricts head organizer formation in
Hydra. *BMC Biol.* 19:1–22.

19. Soriano, J., S. Rüdiger, P. Pullarkat, and A. Ott. 2009. Mechanogenetic coupling of Hydra
symmetry breaking and driven Turing instability model. *Biophys. J.* 96:1649–60.

20. Ferenc, J., P. Papasaikas, J. Ferralli, Y. Nakamura, S. Smallwood, and C.D. Tsiairis. 2021.
Mechanical oscillations orchestrate axial patterning through Wnt activation in Hydra. *Sci. Adv.*
7:6897.

21. Desprat, N., W. Supatto, P.-A. Pouille, E. Beaurepaire, and E. Farge. 2008. Tissue deformation
modulates twist expression to determine anterior midgut differentiation in Drosophila
embryos. *Dev. Cell*. 15:470–7.

22. Gudipaty, S.A., J. Lindblom, P.D. Loftus, M.J. Redd, K. Edes, C.F. Davey, V. Krishnegowda, and J.
Rosenblatt. 2017. Mechanical stretch triggers rapid epithelial cell division through Piezo1.
Nature. 543:118–121.

23. Pouille, P.-A., P. Ahmadi, A.-C. Brunet, and E. Farge. 2009. Mechanical signals trigger Myosin II
redistribution and mesoderm invagination in Drosophila embryos. *Sci. Signal.* 2:ra16.

24.   Shinozuka, T., R. Takada, S. Yoshida, S. Yonemura, and S. Takada. 2019. Wnt produced by
772   stretched roof-plate cells is required for the promotion of cell proliferation around the central
773   canal of the spinal cord. *Dev.* 146.

774   25.   Mercker, M., A. Köthe, and A. Marciniak-Czochra. 2015. Mechanochemical symmetry breaking
775   in Hydra aggregates. *Biophys. J.* 108:2396–407.

776   26.   Brinkmann, F., M. Mercker, T. Richter, and A. Marciniak-Czochra. 2018. Post-Turing tissue
777   pattern formation: Advent of mechanochemistry. *PLoS Comput. Biol.* 14:e1006259.

778   27.   Wang, R., T. Goel, K. Khazoyan, Z. Sabry, H.J. Quan, P.H. Diamond, and E.-M.S. Collins. 2019.
779   Mouth Function Determines the Shape Oscillation Pattern in Regenerating Hydra Tissue
780   Spheres. *Biophys. J.* 117:1145–1155.

781   28.   Livshits, A., L. Shani-Zerbib, Y. Maroudas-Sacks, E. Braun, and K. Keren. 2017. Structural
782   Inheritance of the Actin Cytoskeletal Organization Determines the Body Axis in Regenerating
783   Hydra. *Cell Rep.* 18:1410–1421.

784   29.   Wang, R., R.E. Steele, and E.M.S. Collins. 2020. Wnt signaling determines body axis polarity in
785   regenerating Hydra tissue fragments. *Dev. Biol.* 467:88–94.

786   30.   Wang, R., A.L. Bialas, T. Goel, and E.M.S. Collins. 2023. Mechano-Chemical Coupling in Hydra
787   Regeneration and Patterning. *Integr. Comp. Biol.* 63:1422–1441.

788   31.   Veschgini, M., F. Gebert, N. Khangai, H. Ito, R. Suzuki, T.W. Holstein, Y. Mae, T. Arai, and M.
789   Tanaka. 2016. Tracking mechanical and morphological dynamics of regenerating Hydra tissue
790   fragments using a two fingered micro-robotic hand. *Appl. Phys. Lett.* 108.

791   32.   Evans, E.A. 1973. New Membrane Concept Applied to the Analysis of Fluid Shear- and
792   Micropipette-Deformed Red Blood Cells. *Biophys. J.* 13:941–954.

793   33.   Guevorkian, K., M.-J. Colbert, M. Durth, S. Dufour, and F. Brochard-Wyart. 2010. Aspiration of
794   Biological Viscoelastic Drops. *Phys. Rev. Lett.* 104:1–4.

795   34.   Hochmuth, R.M. 2000. Micropipette aspiration of living cells. *J. Biomech.* 33:15–22.

796   35.   Lim, C.T., E.H. Zhou, and S.T. Quek. 2006. Mechanical models for living cells--a review. *J.
797   Biomech.* 39:195–216.

798   36.   Landiech, S., M. Elias, P. Lapèze, H. Ajiyel, M. Plancke, A. Laborde, F. Mesnilgrente, D. Bourrier,
799   D. Berti, C. Montis, L. Mazenq, J. Baldo, C. Roux, M. Delarue, and P. Joseph. 2023. Parallel on-
800   chip micropipettes enabling quantitative multiplexed characterization of vesicle mechanics
801   and cell aggregates rheology. *bioRxiv*. 2023.10.19.562871.

802   37.   Holzapfel, G.A. 2000. Nonlinear solid mechanics: a continuum approach for engineering
803   science. Wiley, Hoboken.

804   38.   Green, A.E., and W. Zerna. 1968. Theoretical Elasticity. Dover Publications, Mineola.

805   39.   Naik, S., M. Unni, D. Sinha, S.S. Rajput, P.C. Reddy, E. Kartvelishvily, I. Solomonov, I. Sagi, A.
806   Chatterji, S. Patil, and S. Galande. 2020. Differential tissue stiffness of body column facilitates
807   locomotion of Hydra on solid substrates. *J. Exp. Biol.* 223.

808   40.   Boot, R.C., A. Roscani, L. van Buren, S. Maity, G.H. Koenderink, and P.E. Boukany. 2023. High-
809   throughput mechanophenotyping of multicellular spheroids using a microfluidic micropipette
810   aspiration chip. *Lab Chip*. 23:1768.

811   41.   Guevorkian, K., F. Brochard-Wyart, and D. Gonzalez-Rodriguez. 2021. Flow dynamics of 3D
812   multicellular systems into capillaries. In: Viscoelasticity and Collective Cell Migration: An

Interdisciplinary Perspective Across Levels of Organization. Academic Press. pp. 193–223.

42.   Harris, A.R., L. Peter, J. Bellis, B. Baum, A.J. Kabla, and G.T. Charras. 2012. Characterizing the
mechanics of cultured cell monolayers. *Proc. Natl. Acad. Sci. U. S. A.* 109:16449–16454.

43.   Brinkmann, F. 2020. Mathematical models and numerical simulation of mechanochemical
pattern formation in biological tissues. *PhD thesis, Ruprecht-Karls-Universität.*

44.   Ugural, A.C. 2009. Stresses in beams, plates, and shells, third edition. CRC Press, Boca Raton.

45.   Takeichi, M. 1977. Functional correlation between cell adhesive properties and some cell
surface proteins. *J. Cell Biol.* 75:464–474.

46.   Straight, A.F., A. Cheung, J. Limouze, I. Chen, N.J. Westwood, J.R. Sellers, and T.J. Mitchison.
2003. Dissecting temporal and spatial control of cytokinesis with a myosin II inhibitor. *Science*
(80-.). 299:1743–1747.

47.   Sato, M., D.P. Theret, L.T. Wheeler, N. Ohshima, and R.M. Nerem. 1990. Application of the
Micropipette Technique to the Measurement of Cultured Porcine Aortic Endothelial Cell
Viscoelastic Properties. *J. Biomech. Eng.* 112:263–268.

48.   Tsai, M.A., R.S. Frank, and R.E. Waugh. 1993. Passive mechanical behavior of human
neutrophils: power-law fluid. *Biophys. J.* 65:2078–2088.

49.   Ruiz-Herrero, T., K. Alessandri, B. V. Gurchenkov, P. Nassoy, and L. Mahadevan. 2017. Organ
size control via hydraulically gated oscillations. *Development*. 144:4422–4427.

50.   Lien, J.C., and Y. li Wang. 2021. Cyclic stretching-induced epithelial cell reorientation is driven
by microtubule-modulated transverse extension during the relaxation phase. *Sci. Reports*
2021 111. 11:1–12.

51.   Gérémie, L., E. Ilker, M. Bernheim-Dennery, C. Cavaniol, J.L. Viovy, D.M. Vignjevic, J.F. Joanny,
and S. Descroix. 2022. Evolution of a confluent gut epithelium under on-chip cyclic stretching.
Phys. Rev. Res. 4:023032.

52.   Fink, J., N. Carpi, T. Betz, A. Bétard, M. Chebah, A. Azioune, M. Bornens, C. Sykes, L. Fetler, D.
Cuvelier, and M. Piel. 2011. External forces control mitotic spindle positioning. *Nat. Cell Biol.*
2011 137. 13:771–778.

53.   Wu, P.H., D.R. Ben Aroush, A. Asnacios, W.C. Chen, M.E. Dokukin, B.L. Doss, P. Durand-Smet,
A. Ekpenyong, J. Guck, N. V. Guz, P.A. Janmey, J.S.H. Lee, N.M. Moore, A. Ott, Y.C. Poh, R. Ros,
M. Sander, I. Sokolov, J.R. Staunton, N. Wang, G. Whyte, and D. Wirtz. 2018. A comparison of
methods to assess cell mechanical properties. *Nat. Methods 2018 157*. 15:491–498.

54.   Veschgini, M., R. Suzuki, S. Kling, H.O. Petersen, B.G. Bergheim, W. Abuillan, P. Linke, S.
Kaufmann, M. Burghammer, U. Engel, F. Stein, S. Özbek, T.W. Holstein, and M. Tanaka. 2023.
Wnt/β-catenin signaling induces axial elasticity patterns of Hydra extracellular matrix.
iScience. 26:106416.