Sibling chimerism among microglia in marmosets

Ricardo C.H. del Rosario[1,2,*], Fenna M. Krienen[1,2,5], Qiangge Zhang[2,3], Melissa Goldman[1,2], Curtis Mello[1,2], Alyssa Lutservitz[1,2], Kiku Ichihara[1,2], Alec Wysoker[1,2], James Nemesh[1,2], Guoping Feng[2,3], Steven A. McCarroll[1,2,4,*]

[1] Department of Genetics, Harvard Medical School, Boston, MA 02115
[2] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA
[3] McGovern Institute for Brain Research, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
[4] Howard Hughes Medical Institute, Boston, MA 02115
[5] current address: Princeton Neuroscience Institute
*Correspondence: rcdelros@broadinstitute.org, smccarro@broadinstitute.org

Abstract

Chimerism happens rarely among most mammals but is common in marmosets and tamarins, a result of fraternal twin or triplet birth patterns in which *in utero* connected circulatory systems (through which stem cells transit) lead to persistent blood chimerism (12-80%) throughout life. The presence of Y-chromosome DNA sequences in organs of female marmosets has long suggested that chimerism might also affect these organs. However, a longstanding question is whether this chimerism is driven by blood-derived cells or involves contributions from other cell types. To address this question, we analyzed single-cell RNA-seq data from blood, liver, kidney and many brain regions across a number of marmosets, using transcribed single nucleotide polymorphisms (SNPs) to identify cells with the sibling's genome in various cell types within these tissues. Sibling-derived chimerism in all tissues arose entirely from cells of hematopoietic origin (i.e., myeloid and lymphoid lineages). In brain tissue this was reflected as sibling-derived chimerism among microglia (20-52%) and macrophages (18-64%) but not among other resident cell types (neurons, glia or ependymal cells). The percentage of microglia that were sibling-derived showed significant variation across brain regions, even within individual animals, likely reflecting distinct responses by genetic-sibling microglia to local recruitment or proliferation cues or, potentially, distinct clonal expansion histories in different brain areas. In the animals and tissues we analyzed, microglial gene expression profiles bore a much stronger relationship to local/host context than to sibling genetic differences. Naturally occurring marmoset chimerism will provide new ways to recognize the effects of genes, mutations and brain contexts on microglial biology and to distinguish between effects of microglia and other cell types on brain phenotypes.

Introduction

Chimerism, in which an organism contains cells from genetically distinct animals, happens rarely among mammals. Chimerism is common, however, in the *Callitrichidae* family that consists of the marmosets (*Callithrix*) and their close relatives the tamarins (*Saguinus*): in these primate species, animals usually give birth to dizygotic twins or trizygotic triplets whose blood contains cells from siblings. During development, the siblings share circulation *in utero*, allowing the exchange of hematopoietic stem cells (Gengozian et al., 1969; Wislocki, 1939). Most marmosets then exhibit blood chimerism throughout life: their blood-derived DNA is a mixture of both twins' genomes, with the twin's genome contributing 12% to 80% of the DNA in the blood (Niblack et al., 1977; The Marmoset Genome Sequencing and Analysis Consortium, 2014). This indicates that the twins' hematopoietic stem cells establish permanent residency in one another's bodies and contribute to blood cell populations throughout life.

A longstanding mystery involves whether other tissues and organs also harbor chimerism. Beyond the blood, Y-chromosome DNA has been detected in the brain and other organs of female marmosets with male twins (Ross et al., 2007; Sweeney et al., 2012), eliciting much speculation about how chimerism might have shaped behavior and natural selection in marmosets. However, it is still not known what cell types harbor this sibling DNA; such observations could in principle be explained by the presence of blood cells within these organs.

Here, we analyze chimerism in the common marmoset (*Callithrix jacchus*) brain, liver, kidney and blood, with single-nucleus RNA-seq (snRNA-seq), using each cell's RNA-expression pattern to identify its type and using combinations of transcribed SNPs (visible in the snRNA-seq sequence reads) to determine which marmoset sibling is the source of each cell. This approach makes it possible to determine whether chimerism arises from blood cells, resident immune cells, or other cell types, and to explore what chimerism can teach us about cellular migrations and cellular population dynamics.

Results

Marmoset chimerism can be characterized at single-cell resolution

To identify which individual cells have the genome of the host marmoset, and which have the genome of the host's birth sibling, we used combinations of many transcribed SNPs that were visible in the snRNA-seq sequencing reads for each nucleus (using the methods described in Wells et al., 2023).

We first determined whether marmosets have sufficient sequence variation to enable the distinction between host and sibling cells. From whole-genome sequences of 123 marmosets, we identified 13 million polymorphic bi-allelic SNPs in the marmoset genome, with individual marmosets harboring 2.3 to 3.8 million (average 3.4 million) heterozygous sites – comparable to levels of heterozygosity in humans. For sibling comparisons, we detected a large number of sites at which any two siblings' genomes differed, ranging from 2.0 to 3.7 million sites (average 2.9 million sites) across 96 sibling comparisons (**Fig. 1A**). To determine how many of these sites were visible in snRNA-seq data, we analyzed snRNA-seq data for several marmoset tissues. The results varied by cell type, reflecting that different cell types' nuclei harbored different quantities of RNA. The hundreds of transcribed variant sites that differed between siblings (median>311 per nucleus) suggested ample power to distinguish between siblings in all cell types (Wells et al., 2023) (**Fig. 1B**).

We next evaluated whether the genome variation visible in snRNA-seq reads was sufficient to distinguish between host and sibling cells. For this we used Dropulation, which identifies the donors of individual cells (from a set of genome-sequenced candidate donors) by using combinations of the transcribed SNPs visible on the snRNA-seq reads of the individual cells (Wells et al., 2023). We first analyzed the blood cells by snRNA-seq of a marmoset (CJ028) born with two birth siblings and used Dropulation (Wells et al., 2023) to assign individual cells to the correct sibling (**Fig. 1C**). The relative likelihoods of the original source of each cell could be strongly differentiated (relative likelihoods of 10^3 to 10^{23}) for >99% of the nuclei (**Fig. 1C**). We found a high level of chimerism: 84% of all nuclei sampled in this marmoset's blood appeared to contain the genome of one (67% - sibling #1) or the other (17% - sibling #2) of its two birth siblings (**Fig. 1C**), consistent with the wide range of chimerism found in previous studies: 4-82% in marmoset T cells and B cells (Niblack et al., 1977); 13-37% in marmoset whole blood ("The Marmoset Genome Sequencing and Analysis Consortium", 2014).

Apparent liver and kidney chimerism arises from infiltrating monocytes

Earlier studies have identified Y-chromosome-derived DNA sequences in the organs of female marmosets with male birth siblings, suggesting that these organs harbor chimerism (Sweeney et al., 2012). However, such observations could also in principle arise from blood or from blood-derived immune cells that are present in those organs (Sweeney et al., 2012).

We performed snRNA-seq analysis of the blood (1,741 nuclei), liver (10,877 nuclei) and kidney (9,262 nuclei) of a marmoset (CJ026) with one birth sibling (**Fig. 1D**). The snRNA-seq profiles clustered into groups that were readily recognized (based on the RNAs expressed) as the principal cell types of each organ; we determined the identity of each cluster using scType, a cell-type identification tool that uses a database of known marker genes (Ianevski et al., 2022).

In kidney and liver, the only clearly twin-derived cells were cells of hematopoietic origin: the resident macrophages in liver (Kupffer cells), lymphocytes in liver, and lymphocytes in kidney (**Fig. 1E,F**). All non-hematopoietic cell types in liver and kidney appeared to contain only the host marmoset's own genome. Chimerism levels for the two chimeric liver immune cell types appeared to diverge, with sibling-derived cells accounting for 15% of Kupffer cells and just 4% of lymphocytes (5/122 vs 57/383; Chi-square test *P*-value=0.003). In the blood, chimerism levels varied across the various cell types: the most abundant cell types, the Naive B cells and Naive CD8+ T cells, were respectively 29% and 32% sibling-derived, while the less-abundant CD8+ NKT-like cells were 15% sibling-derived (**Fig. 1F**, Chi-square test *P*-value=0.01).

These results indicate that, in this animal's liver and kidney, apparent DNA chimerism likely arose from infiltrating immune cells rather than other cell types. These results also indicate that cells with siblings' genomes can differ in their tendency to acquire specific hematopoietic cell fates and in their tendency to infiltrate into organs.

Marmoset brain microglia and macrophages exhibit abundant chimerism

To characterize chimerism in the marmoset brain, we utilized a large snRNA-seq data set that was generated for a marmoset brain cell atlas (Krienen et al., 2020; Krienen et al., 2023). Brain snRNA-seq was performed on 11 animals (6 adults, 3 neonates and 1 six months old; **Table 1**). All were unrelated except for CJ006 and CJ007 which are birth siblings, and CJ025 and CJ026 which are (non-birth) siblings. All animals come from the three main marmoset colonies that comprise the animals in our facilities: New England Primate Research Center (NEPRC), CLEA Japan, and from a non-clinical contract research organization in Massachusetts. All adult

marmosets had no known previous disease and were selected as part of a larger project to create a single-cell atlas of the marmoset brain (Krienen et al., 2020; Krienen et al., 2023). The three neonates died shortly after birth due to unknown reasons and were subsequently selected for snRNA-seq analysis.

We first analyzed 497,000 single-nucleus RNA-expression profiles from the neocortex, thalamus, striatum, hippocampus, basal forebrain, hypothalamus and amygdala of an adult marmoset with two birth siblings (marmoset CJ028). We clustered the cell types using gene expression similarities and identified brain cell types as in earlier work (Krienen et al., 2020), identifying neurons, astrocytes, oligodendrocytes, ependymal cells, endothelial cells, microglia and macrophages (**Fig. 2A**). Microglia (which expressed markers *TREM2*, *LAPTM5* and *C3*) and macrophages (which expressed *LYVE1* and *F13A1*) were a small fraction of all nuclei analyzed (about 3.6%), but due to the large number of nuclei we profiled (53 brain tissue dissections, 497,000 nuclei), we were able to ascertain sufficient numbers of microglia (18,185 nuclei) and to a lesser extent macrophages (172 nuclei) for many downstream analyses. We found microglia and macrophages in snRNA-seq data from 10 additional marmosets with different genetic backgrounds from 3 different colonies. Brain snRNA-seq of all 11 marmosets showed consistently the presence of these two myeloid cell types in the brain (**Supplementary Fig. 1**; number of microglia and macrophages in **Supplementary Table 1**).

Donor-of-origin analysis of snRNA-seq data from 2.2 million nuclei sampled from 137 brain tissue samples from these 11 marmosets showed a clear and consistent pattern: microglia and macrophages, but not neurons, glia or endothelial cells, harbored chimerism (**Fig. 2B**, **Supplementary Fig. 2**). Microglia exhibited abundant chimerism – across the 11 marmosets, the total fraction of cells with the sibling's genome ranged from 20% to 52% (for triplets, sum of two siblings; **Fig. 2C**). Macrophages exhibited a similarly wide range of sibling fractions across marmosets (18% to 64%, **Fig. 2C**).

The quantitative extent of microglial chimerism varied across individuals (**Supplementary Fig. 3A**; test of heterogeneity *P*-value<2.2×10^{-16}), as did that of macrophage chimerism (**Supplementary Fig. 3B**; test of heterogeneity *P*-value=1×10^{-4}). We asked whether microglial and macrophage chimerism were correlated. Intriguingly, only a modest correlation of chimerism levels across 14 host-sibling pairs was observed between the microglia and macrophages (**Fig. 2D**; Pearson correlation 0.31). We investigated further by performing a statistical test that takes into account the uncertainty in the estimates of the chimeric cell proportion using a binomial framework (Methods); in this analysis, microglia chimerism fraction was not a statistically significant predictor of macrophage chimerism fraction (Methods). This suggests that in addition to the cell's genome, other factors such as local host environment play a role in differential recruitment, proliferation or survival of sibling cells. (We note that macrophages often transit the fluid-filled perivascular space, with a substantially different migration history and arrival dynamics than microglia.) Neither of these two myeloid cell types showed consistently higher chimerism than the other cell type did (**Fig. 2C**).

Sibling contributions in blood vs. brain

Though microglia (like macrophages) are myeloid cells that derive from hematopoietic stem cells, the ontogenies of microglia and brain macrophages are distinct from those of bone-marrow-derived peripheral blood mononuclear cells (Perdiguero & Geissmann, 2016). As such, differences in the developmental and migration histories of these cell populations could in principle have caused their chimerism fractions to diverge in a systematic way.

We analyzed three marmosets for which snRNA-seq was performed on both blood and brain tissues. Sibling contributions to microglia and brain macrophages were in general quite different from those in blood (**Fig. 3**).

Marmoset CJ028's chimerism (involving two birth siblings) provided a setting in which cells with three different genomes shared the same environment through development until adulthood (to two years of age). Among CJ028's microglia, the fraction of cells from sibling 1 (35%) was greater than that from sibling 2 (13%) (two-sided test of proportionality P-value<2.2^{-16}), while in blood, the opposite was true (fraction of cells from sibling 1 across all blood cell types was 18%, fraction of cells from sibling 2 across all blood cell types was 67%; two-sided test of proportionality P-value<2.2^{-16}) (**Fig. 3D,E**, **Supplementary Table 2**).

Microglia chimerism fraction varies across brain regions

Sibling contributions to the microglial population could in principle be shaped by effects that are local to specific brain areas, including differential response of sibling microglia to local recruitment or proliferation cues, or population bottlenecks such as clonal expansions. To evaluate whether the sibling contribution to the microglial population varied across brain areas within individual marmosets, we performed chimerism analysis for each of the brain regions profiled in the snRNA-seq datasets: neocortex, thalamus, striatum, hippocampus, basal forebrain, hypothalamus and amygdala (Krienen et al., 2020; Krienen et al., 2023). Within each marmoset, the fraction of microglia with a sibling's genome diverged across a marmoset's brain regions (**Fig. 4**). For example, in marmoset CJ025, sibling contributions to microglial populations ranged from 11% (21/193) in the thalamus to 56% (174/310) in the striatum (P-value=$1.1\times10^{-23,}$ Chi-square test of thalamus vs striatum; P-value=1.5×10^{-40}, Chi-square test across all 4 brain regions). For marmoset brains profiled with at least 300,000 nuclei, CJ027, CJ028 and CJ029, tests of heterogeneity P-values were even more significant: 8.8×10^{-83}, <1×10^{-300}, and 6.1×10^{-47}, respectively (the error bars are very short in the corresponding panels in Fig. 4). We used the binomial generalized linear mixed-model framework and found that all brain regions were statistically significant predictors for microglia chimerism fraction, supporting the conclusion that chimerism varies across brain regions (Methods). Analysis of finer brain substructures showed a similar result **(Supplementary Fig. 4**; the binomial generalized linear mixed-model framework determined that 18 out of 27 brain substructures were statistically significant as predictors for microglia chimerism fraction, Methods). None of the brain regions exhibited consistently higher or lower chimerism levels, suggesting that these divergences did not result from differential physical access of host and sibling microglia to different brain areas (**Fig. 4** and **Supplementary Fig. 4**).

Gene expression comparisons of host- to sibling-derived microglia

Chimerism provides the unusual opportunity to compare cells with different genomes in a shared *in vivo* biological context. We compared RNA expression between host- and sibling-derived microglia of a female marmoset with two birth siblings. Sex differences among the siblings (the host (CJ028) was a female and one of the two siblings was a male) allowed a natural control: the *XIST* gene encodes a non-coding RNA involved in silencing one copy of chromosome X in females and thus exhibits sex-specific expression due to cell-autonomous mechanisms. We found that (as expected) *XIST* transcripts were detected at far higher levels in the snRNA-seq profiles of microglia with the female twin's genome relative to the male twin's (**Fig. 5A,C**). By contrast, *XIST* transcripts were detected at similar levels in two microglial populations with the genomes of female twins (**Fig. 5B**).

Gene-expression differences between host- and sibling-derived microglia in the same brain could in principle arise from asymmetries in their developmental histories (which would be shared across host animals) or from genomic differences (which would vary from host animal to host animal). In all eleven individual marmosets, analysis identified genes whose differential expression distinguished microglia with the two sibling genomes (hundreds of genes in total), documenting a substantial effect of sibling genetic differences on microglial gene expression. However, we did not find any gene whose expression level recurrently distinguished "host" microglia (microglia with the same genome as neural cell types) from "guest" microglia (microglia with the sibling genome), aside from the XIST gene (a proxy for sibling sex differences, which were of course common) (**Supplementary Fig. 5**, **Fig. 5A-C**). In other words, although there were always gene-expression differences between sibling microglia, none of them consistently distinguished between host and guest microglia, suggesting that they were instead due to sibling genetic differences. We note that both analyses are power-limited, as the number of microglia in most animals, especially guest microglia, were modest (**Supplementary Fig. 5**); thus, we cannot rule out the possibility that there may be one or more genes whose expression levels reflect developmental histories (host vs. guest origin), just as there are likely far more genes (than the hundreds we identified) that can have sibling expression differences due e.g. to genetic differences between siblings. We sought to increase power (beyond single-gene analysis) by using latent factor analysis (Ling et al., 2024) to identify and quantify the expression of microglial gene-expression programs; however, even this analysis did not find any gene expression programs that exhibited consistent host-twin differences in expression levels (Methods).

Brain context vs. genetic differences as determinants of microglial gene expression

Chimerism presents interesting opportunities to distinguish between cell-autonomous and contextual effects on a cell's biology, and to compare the magnitudes of such effects.

We first considered the difference in contexts provided by pairs of brain areas by analyzing snRNA-seq data from the neocortex and striatum of marmoset CJ027; the resident microglial populations with different genomes make it possible to compare contextual to genetic effects on microglial gene expression (**Fig. 5D**). Genetic effects appeared to elicit very many small-magnitude gene-expression differences; these differences were shared between cortical and striatal microglia (**Fig. 5E**). Brain-area context elicited much larger-magnitude gene-expression differences, which were experienced in common by microglia with both genotypes (**Fig. 5F**). We obtained similar results for all pairs of brain areas analyzed (52 context vs genetic effect from brain snRNA-seq of 6 marmosets with at least 60 cells available for analysis in each context; **Supplementary Table 3**; **Supplementary Fig. 6**).

We next considered the difference in contexts provided by the same brain area in different marmosets. Two of the marmosets profiled, CJ006 and CJ007, were birth siblings who passed away as neonates (the only birth siblings in our dataset), and thus provided the additional opportunity to distinguish genetic from contextual effects by analyzing the two sibling microglial populations in the cortex, striatum and hippocampus of both marmosets (**Fig. 5G**). The effects of context (host marmoset) in microglia from all three brain areas appeared to be far larger than the cell-autonomous effects of genetic differences (**Fig. 5H,I**).

Discussion

A longstanding debate concerns the extent of chimerism in marmosets and tamarins. Chimerism in these species has been detected in diverse organs but arises from unknown cell

300 types (Ross et al., 2007; Sweeney et al., 2012). Here we found that chimerism in the brain, liver
301 and kidney is present but appears to arise entirely from cells of the myeloid and lymphoid
302 lineages, including infiltrating macrophages, monocytes, and microglia.
303
Cells of the myeloid and lymphoid lineages derive developmentally from hematopoietic stem
cells. We found no strong evidence of chimerism among 2.2 million non-hematopoietic cells in
the liver (from one marmoset), kidney (from one marmoset) or brain (from 11 marmosets).
Thus, while marmosets share a circulation *in utero*, we found no evidence that other kinds of
stem cells or progenitors (beyond those of hematopoietic lineage) had been shared via this
route in any appreciable number. However, we found that in the marmoset brain, the microglia
and macrophages, which also derive from this lineage, routinely harbor abundant chimerism,
with 10-50% of a marmoset's microglia containing the genome(s) of birth sibling(s).
312
Organs in the same marmoset (liver, kidney, brain) differed markedly in the sibling contribution
to resident macrophage and monocyte populations, with microglial chimerism fraction (the
fraction of cells contributed by siblings) varying by as much as 40 percentage points across a
marmoset's brain areas. This phenomenon has more than one potential explanation. First,
cells from the host and sibling could in principle respond differently to recruitment or proliferation
cues that vary spatially; if this is the case, marmoset chimerism could provide a model for
studying the effects of mutations and natural sequence variation on cell migration and
recruitment. (Although we found that genetic effects were smaller than contextual effects in
shaping microglial gene expression at any moment in time, genetic effects were clear (**Fig. 5E**),
and even small effects on proliferation rates would tend to have effects that increase
exponentially over time.) Second, beyond such recruitment effects, it is also possible that these
differences suggest a substantial role of clonal expansions and population bottlenecks in
shaping local microglial and macrophage populations.
326
We found that the cellular contribution of birth siblings to myeloid cell populations was
significantly different in blood than in brain in the modest number of marmosets analyzed (**Fig.
3**). Unlike the differences among brain areas, the blood-brain differences tended to be
directional, with more-modest sibling contributions in the brain than in the blood. Though this
would need to be confirmed in many more marmosets to be definitive on its own, it is plausibly
connected to this aspect of marmoset fetal development: sharing of a blood circulation between
the two fetuses occurs during a window that is more temporally extended than the waves of
colonization of the brain by microglia, potentially allowing for greater exchange in the centers of
blood hematopoiesis (the liver and then the bone marrow). In microglia and macrophages, due
to the defined waves of hematopoiesis in the yolk sac and migration patterns of microglia and
macrophages to the developing brain, the opportunity for a progenitor cell from a twin to
colonize a host's brain may need to occur during a more restricted temporal window.
339
Comparisons of gene expression between microglial cells with host and sibling genomes in a
shared brain context may provide many future opportunities to distinguish the cell-autonomous
from non-cell autonomous genetic effects of genetic differences and engineered mutations.
Such analyses could become especially useful scientifically as genome editing increasingly
enables the utilization of marmosets as a model organism in translational neuroscience (Aida &
Feng, 2020; Feng et al., 2020). Our pilot analysis of host-sibling microglial gene expression
differences in the brains of two co-twins revealed a large role of animal context (relative to
genetic differences) in shaping microglial gene expression. This result points to an important
principle: the ability to isolate the effects of a mutation will be greatly strengthened by the ability
to make within-animal (rather than just between-animal) comparisons of cells with different
genotypes.

351
A long history of innovation in genetics involves elaborate ways to create mosaics in mice, *C.*
elegans and other laboratory organisms in order to distinguish cell-autonomous from non-cell
autonomous genetic effects. Natural chimerism in marmosets may enable many straightforward
ways to pursue such kinds of studies.   Natural chimerism may also make it possible to
determine when microglia or macrophages, as opposed to other cell types, mediate the effect of
a mutation on an animal's phenotype.
358
Chimerism could also enable interesting future analyses of whether there are adaptive benefits
of chimerism in marmoset immune cells, among whom chimerism could in principle allow
presentation of a wider variety of antigens for adaptive immunity. In a recent outbreak of yellow
fever in Brazil in 2016-2018, marmosets were found to be less susceptible than other primates
that lack immune system chimerism, including the howler monkeys (*Alouatta*), robust capuchins
(*Sapajus*), and titi monkeys (*Callicebus*) (de Azebedo Fernandes, et al., 2021). In studying
future outbreaks in marmosets, one could use single-cell RNA-seq and the methods described
here to study how genetically distinct immune cells (in the same animal) have differentially
migrated to affected tissues and/or assumed "activated" immune cell states. Recent innovations
in spatial transcriptomics with sequencing readouts (that detect SNP alleles) may also make it
possible to identify any differential recruitment of genetically distinct immune cells to focal
infection sites.
371
Microglia perform essential roles in the development and regulation of the central nervous
system (CNS), including by sculpting or "pruning" neuronal circuits (Hammond et al., 2018;
Schafer et al., 2012; Schafer & Stevens, 2015), and are implicated in or hypothesized to
contribute to a wide range of brain disorders and diseases, including Alzheimer's disease,
Parkinson's disease, autism spectrum disorder, and schizophrenia.   Marmoset microglial
chimerism will enable many new ways of studying microglia and the effects of genes and alleles
upon brain biology.
379
380 **Methods**
381
382 **Ethical compliance**
383 Marmoset experiments were approved by and in accordance with Massachusetts Institute of
384 Technology IACUC protocol number 051705020.
385
Nucleus Drop-seq library preparation and sequencing
Nucleus suspensions were prepared from frozen tissue and used for nucleus Drop-seq following
the protocol we have described at https://doi.org/10.17504/protocols.io.2srged6. Drop-seq
libraries were prepared as previously described (Macosko et al., 2015), with modifications,
quantification and quality control as described in a previous study (Saunders et al., 2018), as
well as the following modifications optimized for nuclei: in the Drop-seq lysis buffer, 8M
guanidine hydrochloride (pH 8.5) was substituted for water, nuclei were loaded into the syringe
at a concentration of 176 nuclei/µl, and cDNA amplification was performed using around 6,000
beads per reaction, 15 PCR cycles. Raw sequencing reads were aligned to the calJac3
marmoset reference genome assembly and reads that mapped to exons or introns of each
assembly were assigned to annotated genes (https://github.com/broadinstitute/Drop-seq). Drop-
seq libraries are indicated in **Table 1**.
398
Nucleus 10X Chromium library preparation and sequencing
Single-nucleus suspensions from frozen tissue were generated as for Drop-seq; GEM
generation and library preparation followed the manufacturer's protocol (protocol versions

#CG00052 Chromium Single Cell 3' v2 and #CG000183 Chromium Single Cell3′ v3 UG_Rev-A).
Raw sequencing reads were processed and aligned using the same method for aligning Drop-
seq reads. 10X Chromium libraries are indicated in **Table 1**.
405
Clustering of cells using Independent Component Analysis
Nuclei from intact cells were identified and clustered into cell types using a method that we have
previously described (Krienen et al., 2020; Saunders et al., 2018). Briefly, nuclei with less than
400 detected genes were not used in the analysis. A digital gene expression matrix was
created for a set of libraries from the same animal that were to be co-analyzed (**Supplementary
Table 4**), and independent component analysis using the fastICA package in R was used after
normalization and variable gene selection as previously described (Krienen et al., 2020;
Saunders et al., 2018). A Louvain-based clustering algorithm was performed on the top 60
independent components. Due to the large number of nuclei profiled in some marmosets
(CJ027, CJ028, CJ029), memory requirements exceeded machine limits and for these
marmosets, we divided the clustering analysis into two or three batches (**Supplementary
Table4**). The brain of marmoset CJ022 was profiled using both Drop-seq and 10X and a
separate clustering was done for each snRNA-seq method (**Supplementary Table 4**). We ran
the clustering algorithm 12 times using 3 nearest neighbor parameters (10,20,30) and 4
resolution parameters (0.3, 0.5, 0.1, 1.0). Markers for each cluster were identified using
differential gene expression analysis (Krienen et al., 2020; Saunders et al., 2018). We inspected
each clustering result and chose the one which yielded separate clusters for microglia and
macrophages (**Supplementary Table 4**).
424
Identification of cell types
For the brain datasets, the microglia and macrophage clusters were identified by the markers
TREM2, C3, LAPTM5 for microglia and *F13A1, LYVE1* for macrophages. The other brain cell
types were identified using cell type markers for neurons, astrocytes, oligodendrocytes,
polydendrocytes and endothelial cells that we used as before (Krienen et al., 2020). Cell types
in blood, liver and kidney were identified using the ScType method (Ianevski et al., 2022).
431
Donor-of-origin analysis and detection of host-sibling doublets (Dropulation)
We used the Dropulation suite to calculate a donor likelihood for each cell (Wells et al., 2023)
(software available at https://github.com/broadinstitute/Drop-seq). The host and birth sibling
genotypes were provided as input to Dropulation's AssignCellsToSamples tool, together with
the snRNA-seq BAM file and a list of cell barcodes that were identified to be intact cells. To
generate chimerism-free reference genotypes, we cultured fibroblasts and performed whole
genome sequencing (WGS) on the resulting DNA. We found that the difference in likelihoods
between host and sibling increase with the number of UMI of the cell, and hence we imposed a
minimum number of UMI for each marmoset's cells (**Supplementary Table 4**). We also
performed doublet detection using Dropulation's DetectDoublets to obtain a likelihood of a cell
having a mix of transcripts from the host and sibling(s). Doublets lie between the host and
sibling curves (**Supplementary Fig. 7**) and for each marmoset we empirically obtained a
threshold for the Dropulation test statistic to identify them and were discarded in all analyses
(**Supplementary Fig. 7**, **Supplementary Table 4**). The likelihoods plotted in Figures 1C,1E,2C
and Supplementary Fig. 2 are from cells that have been filtered for minimum UMI and doublets.
447
Marmosets CJ006 and CJ007 were born in a triplet litter (tri-zygotic) that all died shortly after
birth. We did not have access to any tissue from the third sibling and were not able to perform
whole genome sequencing on it. For Dropulation analysis of CJ006 and CJ007's brains, we
provided only the genotypes of marmosets CJ006 and CJ007. Nuclei that contain the genome

of the third unknown sibling will mostly be identified as doublets, which were discarded in our
analysis.
454
Additional filtering for microglia and macrophage clusters
We performed additional filtering of microglia and macrophage cells. When we compared the
gene expression of host microglia and sibling microglia using cell-types from first-round
clustering (and with UMI and doublet filtering), we found an abundance of genes that have
higher expression in host than in the sibling (**Supplementary Fig. 8**, see panels B, C, G, I and
K). The asymmetry could arise from neuronal cells mis-classified as microglia or macrophages.
To filter out these mis-classified cells, we sub-clustered the microglia and macrophage cell
types of each marmoset using the same fastICA and Louvain-based clustering used in the first
round of clustering. We found that some sub-clusters were not chimeric, indicating that they
were not cells of hematopoietic origin, and that discarding these cells improved the symmetry
between host and sibling gene expression (**Supplementary Fig. 8**).
466
Whole genome sequencing
Illumina libraries from fibroblast, blood, brain and buccal cells (**Supplementary Table 5**) were
created as follows. An aliquot of genomic DNA (150ng in 50µL) is used as the input into DNA
fragmentation (aka shearing). Shearing is performed acoustically using a Covaris focused-
ultrasonicator, targeting 385bp fragments. Following fragmentation, additional size selection is
performed using a SPRI cleanup. Library preparation is performed using a commercially
available kit provided by KAPA Biosystems (KAPA Hyper Prep with Library Amplification Primer
Mix, product KK8504), and with palindromic forked adapters using unique 8-base index
sequences embedded within the adapter (purchased from Roche). The libraries are then
amplified by 10 cycles of PCR. Following sample preparation, libraries are quantified using
quantitative PCR (kit purchased from KAPA Biosystems) with probes specific to the ends of the
adapters. This assay is automated using Agilent's Bravo liquid handling platform. Based on
qPCR quantification, libraries are normalized to 2.2nM and pooled into 24-plexes. Sample pools
are combined with NovaSeq Cluster Amp Reagents DPX1, DPX2 and DPX3 and loaded into
single lanes of a NovaSeq 6000 S4 flowcell cell using the Hamilton Starlet Liquid Handling
system. Cluster amplification and sequencing occur on NovaSeq 6000 Instruments utilizing
sequencing-by-synthesis kits to produce 151bp paired-end reads. Output from Illumina software
is processed by the Picard data-processing pipeline to yield CRAM or BAM files containing
demultiplexed, aggregated aligned reads. All sample information tracking is performed by
automated LIMS messaging. All samples were sequenced to 30X coverage.
487
Variant site detection and genotyping from whole genome sequencing
Illumina paired-end reads were aligned to the calJac3 reference marmoset genome assembly
using bwa (Li & Durbin, 2010) with command "bwa mem". Duplicate reads were marked using
Picard Markduplicates and for each chromosome, the GATK Haplotype Caller (McKenna et al.,
2010) was run in genotype discovery GVCF mode. For each chromosome, the GVCFs of all
samples analyzed in this study (from fibroblasts, blood, buccal cells, skin, brain and hair), were
combined into a single GVCF file using GATK CombineGVCFs. To obtain the highest sensitivity
in calling SNPs, we included in the GVCF additional fibroblasts whole genome sequences from
the colony, yielding a total of 113 marmosets for multi-sample variant calling. The GVCF of
each chromosome was genotyped using GATK GenotypeGVCFs. Only bi-allelic SNPs were
used in the analysis and the following filters were used: QD<4.0 | FS>60.0 | MQ<40.0 |
MQRankSum<-12.5 | ReadPosRankSum<-8.0 | MAF<0.01 | QUAL<500. SNP calls from all
chromosomes were combined into one VCF file and additional filtering was performed to discard
heterozygous sites that exhibited extreme allelic imbalance, i.e., the fraction of non-reference
allele (from all samples) is less than 0.2 or greater than 0.8, and furthermore, sites in copy

number variant regions were discarded (copy number variant regions were obtained by running
Genome STRiP (Handsaker et al., 2015) on whole genome sequencing data from 113 fibroblast
samples).
506
Dropulation analysis using sibling genotypes from whole genome sequencing of buccal
cells
For four marmosets in our dataset (CJ022, CJ025, CJ026, CJ102; all born with one sibling and
the siblings are CJ106, CJ104, CJ105 and CJ103, respectively), only the buccal cells (from
cheek swabs) of their siblings were available for whole genome sequencing. Using a method
that quantifies chimerism from whole genome sequencing data (Census-seq; software available
at https://github.com/broadinstitute/Drop-seq) (Mitchell et al., 2020), we estimated the
chimerism fraction in buccal cells as follows: CJ106: 10%, CJ104: 24%, CJ105: 24% and
CJ103: 9%. Thus, the genotypes we obtained for these marmosets will include errors and those
genotyping errors could subsequently affect the Dropulation (donor-of-origin) analysis that was
used to estimate chimerism. To empirically estimate how sibling genotypes obtained from a
chimeric tissue affect Dropulation analysis, we selected a host-sibling pair whose genome
sequencing were both obtained from fibroblast cultures: CJ027 and its birth sibling CJ140. To
simulate DNA contamination, we fixed the sequencing coverage of CJ140 to 40X, and replaced
between 1% to 60% of the reads from CJ027's sequencing reads (random subsampling using
'samtools view -s'). We genotyped CJ140's "chimeric" bam files using GATK's "genotype given
alleles" mode and compared the genotypes with CJ140's true genotypes from its pure fibroblast
WGS. We found that the sensitivity at heterozygous sites remains constant with different
contamination levels, while the false positive rate increases. The false positive calls at
heterozygous sites come from homozygous sites incorrectly genotyped as heterozygous
(**Supplementary Fig. 9A-F**). Next, we re-analyzed donor-of-origin on brain snRNA-seq of
CJ027 sibling (CJ140) genotypes from simulated contaminated DNA (300,000 nuclei; CJ027
genotypes from pure fibroblast WGS, CJ140 genotypes from WGS with various contamination
levels). We found that chimerism in CJ140's WGS resulted in doublets being assigned to the
twin (**Supplementary Fig. 9G-L**), which subsequently causes a slight increase in chimerism
estimates (**Supplementary Fig. 9M-U**). The contamination levels in buccal cells were from 9%
to 24%, which we estimate will result in an overestimation in microglia chimerism of up to 3.5
percentage points and 4.5 percentage points in macrophage chimerism. Our results will not be
affected by this over-estimation of chimerism in 4 marmosets since the conclusions were made
from the analysis of all 11 marmosets (including 7 marmosets whose siblings were genotyped
from fibroblast cultures).
538
Gene expression analysis of host and sibling meta cells
For each cell type, the host and sibling "meta cells" were calculated from the sum of UMI counts
per gene across cells and were scaled to counts per 100,000 transcripts. The fold-changes and
P-values of differentially expressed genes were identified using the binomTest method from the
edgeR package (Robinson et al., 2010). The filters we used to identify statistically significant
differentially expressed genes are: FDR Q-value<0.05 and fold-change>1.5 (in either direction)
and the gene must be expressed in at least 10% of at least one of the two sets of microglia
being compared.
547
Binomial generalized linear mixed-effects model analysis
To perform an analysis of **Fig. 2D** that takes into account the uncertainty in the estimate of the
chimeric cell proportion, we performed a binomial generalized linear mixed-effects model
analysis in R using the command **glmer(y~(1|indiv)+ chimerism_micro, family=binomial)**,
where **y** is a vector (of length 1,333) containing the genomic identity of each macrophage (either
host or twin), **1|indiv** models a random effect for the identity of each animal, and

chimerism_micro is the microglia chimerism of the animal's brain. The fixed effects probability of **chimerism_micro** was 0.795 indicating that microglial chimerism fraction was not statistically significant as a predictor for macrophage chimerism fraction. The estimate for the intercept was -0.8115 and estimate for **chimerism_micro** was 0.3106, which indicates that the probability of a cell is a macrophage given the microglia chimerism fraction was only 0.57 (plogis(-0.8115+0.3106)).

We used the same framework to further analyze **Fig. 4**. We included brain region as a covariate in the binomial framework: **glmer(y~(1|indiv) + brain_reg + assay, family=binomial)**, where **y** is a vector of length 48,439 containing the genomic identity of each microglia (either host or twin) and **assay** is either "Drop-seq" or "10X". The brain regions assayed in **Fig. 4** are the cortex, hippocampus, hypothalamus, striatum, thalamus, and basal forebrain. All these brain regions were statistically significant as predictors for microglia chimerism fractions (all *P*-values$<2 \times 10^{-16}$), supporting the conclusion that chimerism varies across brain regions. We also re-analyzed **Supplementary Fig. 4** using the same framework and found that 18 out of 27 brain substructures were statistically significant as predictors for microglia chimerism fraction.

Latent factor analysis
Following the method described in (Ling et al., 2024), we performed latent factor analysis using the probabilistic estimation of expression residuals (PEER, Stegle et al., 2010) on the gene-by-donor matrix expression of microglia. We started by creating a gene-by-cell matrix of microglia gene expression from all animals and we normalized the matrix using SCT transform version 2 (Choudhary and Satija, 2022) with 3000 variable features. We obtained the Pearson residuals from SCT normalization and summed up the residual across cells with the same genome to obtain a gene-by-donor matrix of expression measurements of microglia. We used this matrix as input to PEER and ran the tool with a provided number of factors from 9 to 12. For each gene expression latent factor, to evaluate whether host/sibling identity has a consistent effect on expression levels, we performed a linear regression with host/sibling identity using **glm(peer_factor_k ~ host_or_twin)**. For all factors, the *P*-values for the effect of **host_or_twin** were all insignificant (greater than 0.1), indicating that no PEER factor associated with host-vs-twin identify. Thus our results found no large-scale gene expression program that was consistently expressed differently between hosts and twins.

Software availability
All software used in the analysis are publicly available. Drop-seq (analysis of snRNA-seq data, clustering, marker genes), Census-seq (estimation of chimerism in whole-genome sequencing data) and Dropulation analysis (estimation of chimerism in snRNA-seq data): https://github.com/broadinstitute/Drop-seq; alignment and variant detection of Illumina whole genome sequencing data: bwa (https://github.com/lh3/bwa), GATK (https://gatk.broadinstitute.org), BCFtools (https://github.com/samtools/bcftools), samtools (http://www.htslib.org/download), Picard Tools (https://broadinstitute.github.io/picard); R environment (https://www.rstudio.com/products/rstudio/download and https://www.r-project.org); single-cell analysis in R: Seurat (https://github.com/satijalab/seurat), cell type identification: scType (https://github.com/IanevskiAleksandr/sc-type), SCT transform (https://github.com/satijalab/sctransform/); PEER latent factor analysis (https://github.com/PMBio/peer).

Data availability
Brain snRNA-seq of 6 marmosets (CJ022, CJ023, CJ025, CJ026, CJ027, CJ028) were generated as part of the NIH's Brain Initiative Cell Census Network (BICCN) project, while brain snRNA-seq of 5 marmosets (CJ001, CJ006, CJ007, CJ023, CJ102), and all blood, liver and

kidney snRNA-seq were generated for this project. All snRNA-seq datasets are available in the BICCN NeMO portal (https://assets.nemoarchive.org/dat-hsgdsgu and https://assets.nemoarchive.org/dat-1je0mn3). All whole genome sequencing datasets will be provided in a manuscript (in preparation) that will describe naturally occurring genome variation in captive marmosets.

Acknowledgements

This work was supported by the NIH BRAIN Initiative U01MH114819, the Stanley Center for Psychiatric Research at the Broad Institute of MIT and Harvard, the James and Patricia Poitras Center for Psychiatric Disorders Research at MIT and the Hock E. Tan and K. Lisa Yang Center for Autism Research at MIT.

References

Aida, T., & Feng, G. (2020). The dawn of non-human primate models for neurodevelopmental disorders. *Current Opinion in Genetics & Development*, *65*, 160–168. https://doi.org/10.1016/j.gde.2020.05.040

Choudhary, S. & Satija, R. (2022). Comparison and evaluation of statistical error models for scRNA-seq. *Genome Biology,* 23.1. https://doi.org/10.1186/s13059-021-02584-9

de Azevedo Fernandes NCC, Guerra JM, Díaz-Delgado J, Cunha MS, Saad LD, Iglezias SD, Ressio RA, Dos Santos Cirqueira C, Kanamura CT, Jesus IP, Maeda AY, Vasami FGS, de Carvalho J, de Araújo LJT, de Souza RP, Nogueira JS, Spinola RMF, Catão-Dias JL. (2021) Differential Yellow Fever Susceptibility in New World Nonhuman Primates, Comparison with Humans, and Implications for Surveillance. Emerg Infect Dis, Jan;27(1):47-56. doi: 10.3201/eid2701.191220

Feng, G., Jensen, F. E., Greely, H. T., Okano, H., Treue, S., Roberts, A. C., Fox, J. G., Caddick, S., Poo, M.-M., Newsome, W. T., & Morrison, J. H. (2020). Opportunities and limitations of genetically modified nonhuman primate models for neuroscience research. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(39), 24022–24031. https://doi.org/10.1073/pnas.2006515117

Gengozian, N., Batson, J. S., Greene, C. T., & Gosslee, D. G. (1969). Hemopoietic chimerism in imported and laboratory-bred marmosets. *Transplantation*, *8*(5), 633–652. https://doi.org/10.1097/00007890-196911000-00009

Hammond, T. R., Robinton, D., & Stevens, B. (2018). Microglia and the Brain: Complementary Partners in Development and Disease. *Annual Review of Cell and Developmental Biology*, *34*, 523–544. https://doi.org/10.1146/annurev-cellbio-100616-060509

Handsaker, R. E., Van Doren, V., Berman, J. R., Genovese, G., Kashin, S., Boettger, L. M., & McCarroll, S. A. (2015). Large multiallelic copy number variations in humans. *Nature Genetics*, *47*(3), 296–303. https://doi.org/10.1038/ng.3200

Ianevski, A., Giri, A. K., & Aittokallio, T. (2022). Fully-automated and ultra-fast cell-type identification using specific marker combinations from single-cell transcriptomic data. *Nature Communications*, *13*(1), 1246. https://doi.org/10.1038/s41467-022-28803-w

Krienen, F. M., Goldman, M., Zhang, Q., Del Rosario R, C. H., Florio, M., Machold, R., Saunders, A., Levandowski, K., Zaniewski, H., Schuman, B., Wu, C., Lutservitz, A., Mullally, C. D., Reed, N., Bien, E., Bortolin, L., Fernandez-Otero, M., Lin, J. D., Wysoker, A., … McCarroll, S. A. (2020). Innovations present in the primate interneuron repertoire.

Nature, *586*(7828), 262–269. https://doi.org/10.1038/s41586-020-2781-z

Krienen, F. M., Levandowski, K. M., Zaniewski, H., del Rosario, R. C. H., Schroeder, M. E., Goldman, M., Lutservitz, A., Zhang, Q., Li, K. X., Beja-Glasser, V. F., Sharma, J., Shin, T. W., Mauermann, A., Wysoker, A., Nemesh, J., Kashin, S., Vergara, J., Chelini, G., Dimidschstein, J., … Feng, G. (2023). A marmoset brain cell census reveals persistent influence of developmental origin on neurons. *Sciences Advances*, 9(41) DOI: 10.1126/sciadv.adk3986

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics, 26*(5), 589–595. https://doi.org/10.1093/bioinformatics/btp698

Ling, E., Nemesh, J., Goldman, M., Kamitaki, N., Reed, N., Handsaker, R., Genovese, G., Vogelgsang, J., Gerges, S., Kashin, S., Ghosh, S., … McCarroll S. (2024). A concerted neuron–astrocyte program declines in ageing and schizophrenia. *Nature,* 627, 604–611. https://doi.org/10.1038/s41586-024-07109-5

Macosko, E. Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I., Bialas, A. R., Kamitaki, N., Martersteck, E. M., Trombetta, J. J., Weitz, D. A., Sanes, J. R., Shalek, A. K., Regev, A., & McCarroll, S. A. (2015). Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell*, *161*(5), 1202–1214. https://doi.org/10.1016/j.cell.2015.05.002

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Mitchell, J. M., Nemesh, J., Ghosh, S., Handsaker, R. E., Mello, C. J., Meyer, D., Raghunathan, K., de Rivera, H., Tegtmeyer, M., Hawes, D., Neumann, A., Nehme, R., Eggan, K., & McCarroll, S. A. (2020). Mapping genetic effects on cellular phenotypes with "cell villages." *bioRxiv*, 2020.06.29.174383. https://doi.org/10.1101/2020.06.29.174383

Niblack, G. D., Kateley, J. R., & Gengozian, N. (1977). T-and B-lymphocyte chimerism in the marmoset. *Immunology*, *32*(2), 257–263. https://www.ncbi.nlm.nih.gov/pubmed/139360

Perdiguero, E. G., & Geissmann, F. (2016). The development and maintenance of resident macrophages. *Nature Immunology*, *17*(1), 2–8. https://doi.org/10.1038/ni.3341

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics, 26*(1), 139–140. https://doi.org/10.1093/bioinformatics/btp616

Ross, C. N., French, J. A., & Orti, G. (2007). Germline chimerism and paternal care in marmosets (Callithrix kuhlii). *Proceedings of the National Academy of Sciences of the United States of America*, *104*(15), 6278–6282. https://doi.org/10.1073/pnas.0607426104

Saunders, A., Macosko, E. Z., Wysoker, A., Goldman, M., Krienen, F. M., de Rivera, H., Bien, E., Baum, M., Bortolin, L., Wang, S., Goeva, A., Nemesh, J., Kamitaki, N., Brumbaugh, S., Kulp, D., & McCarroll, S. A. (2018). Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell*, *174*(4), 1015–1030 e16. https://doi.org/10.1016/j.cell.2018.07.028

Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R., Ransohoff, R. M., Greenberg, M. E., Barres, B. A., & Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*, *74*(4),

691–705. https://doi.org/10.1016/j.neuron.2012.03.026

Schafer, D. P., & Stevens, B. (2015). Microglia Function in Central Nervous System

Development and Plasticity. *Cold Spring Harbor Perspectives in Biology*, *7*(10), a020545.

https://doi.org/10.1101/cshperspect.a020545

Stegle O, Parts L, Durbin R, Winn J (2010). A Bayesian Framework to Account for Complex

Non-Genetic Factors in Gene Expression Levels Greatly Increases Power in eQTL Studies.

PLoS Comput Biol, 6(5): e1000770. https://doi.org/10.1371/journal.pcbi.1000770

Sweeney, C. G., Curran, E., Westmoreland, S. V., Mansfield, K. G., & Vallender, E. J. (2012).

Quantitative molecular assessment of chimerism across tissues in marmosets and

tamarins. *BMC Genomics*, *13*, 98. https://doi.org/10.1186/1471-2164-13-98

"The Marmoset Genome Sequencing and Analysis Consortium", (2014). The common

marmoset genome provides insight into primate biology and evolution. *Nature Genetics*,

46(8), 850–857. https://doi.org/10.1038/ng.3042

Wells, M. F., Nemesh, J., Ghosh, S., Mitchell, J. M., Salick, M. R., Mello, C. J., Meyer, D.,

Pietilainen, O., Piccioni, F., Guss, E. J., Raghunathan, K., Tegtmeyer, M., Hawes, D.,

Neumann, A., Worringer, K. A., Ho, D., Kommineni, S., Chan, K., Peterson, B. K., …

McCarroll, S. A. (2023). Natural variation in gene expression and viral susceptibility

revealed by neural progenitor cell villages. *Cell Stem Cell*, *30*(3), 312–332.e13.

https://doi.org/10.1016/j.stem.2023.01.010

Wislocki, G. B. (1939). Observations on twinning in marmosets. *The American Journal of*

Anatomy, *64*(3), 445–483. https://doi.org/10.1002/aja.1000640305

717
Figure 1. Sibling chimerism analysis at single-cell resolution. (A) Numbers of variant sites
at which marmoset sibling genomes differ, for 96 sibling-pairs. Each dot is a sibling-pair; birth
siblings are colored in pink. **(B)** Numbers of transcribed SNPs (per nucleus) visible in various
cell types from blood and brain snRNA-seq datasets of marmoset CJ028. x-axis: snRNA-seq
library; y-axis: number of ascertained, transcribed SNPs for which host and sibling have
different genotypes; black horizontal lines: median values per cell type. **(C)** Donor-of-origin
assignment of each nucleus in blood snRNA-seq of a marmoset. The host marmoset (CJ028)
was born with two siblings; each nucleus was assigned to either the host or to one of the two

birth siblings. x-axis: number of unique molecular identifiers (UMI; a measure of transcript abundance) that contains SNPs for which the host and sibling's genomes differ, in log scale; y-axis: inferred likelihood that the cell has host genome minus likelihood that the cell has sibling genome (log10). (**D**) Two-dimensional visualization (tSNE plot) of snRNA-seq data from a marmoset's (CJ026) blood, kidney and liver. **(E)** Donor-of-origin assignment in marmoset CJ026's blood, kidney and liver. axes: same as in (C). **(F)** Levels of chimerism in each cell type ascertained in blood, kidney and liver snRNA-seq of marmoset CJ026. y-axis: fraction of sibling cells in each cell type; numbers in fraction: number of sibling nuclei over total nuclei in the cell type; percentage in x-axis labels: cell type representation in the tissue; vertical bars: binomial confidence interval (95%); *P*-values: test of heterogeneity (Chi-square) across immune cell types of a tissue (to test for differences in contribution of sibling across immune cell types).

Figure 2. Microglia and macrophages, but not neurons and glia, are chimeric in the marmoset brain. (A) Two-dimensional visualization (tSNE plot) of marmoset CJ028's snRNA-seq profiles from 8 brain regions. **(B)** Donor assignment of each nucleus to one of three possible donors (host, sibling1 or sibling2) of marmoset CJ028's brain snRNA-seq data. x-axis: number of UMI that contains SNPs for which the host and sibling's genomes differ, in log10 scale; y-axis: inferred likelihood that the cell has host genome minus likelihood that the cell has sibling genome (log10). Data from all of CJ028's brain regions. **(C)** Fractions of cells with a sibling's genome among microglia, macrophages, and other brain cell types (neurons, astrocytes, oligodendrocytes, polydendrocytes, ependyma, endothelial) from 11 marmosets. CJ001, CJ023 and CJ028 were part of a triplet litter and the chimerism fraction of each birth sibling is shown in separate panels. y-axis: fraction of sibling-cells in the cell type. vertical bars: binomial confidence interval (95%). *P*-values: Chi-square test from comparison of chimerism fractions in microglia and macrophage; * (*P*-value<0.05). (**D**) Sibling contributions to microglial (x-axis) and macrophage (y-axis) populations in the same animals. The 14 pairs of animals are from 11 host-sibling1 pairs plus 3 host-sibling2 pairs (three of the 11 host animals were born in a triplet litter; see Table 1). Dots and bars are from (D). Pearson correlation *R*=0.32, 95% confidence interval (-0.26 to 0.73), *P*-value=0.27.

763
Figure 3. Sibling contributions to hematopoiesis-derived cells diverge between blood
and brain. (A)-(E) Chimerism fractions in brain and blood of three animals. CJ028 was part of
a triplet litter, and the contribution of each sibling is shown in a separate panel (D,E). Red
horizontal lines: twin contribution to blood cells as ascertained from whole-genome sequencing
of whole-blood-derived genomic DNA (Census-seq). Blue horizontal lines: total twin contribution
to blood cells as estimated from PBMC snRNA-seq (all cells). Vertical bars: binomial confidence
interval (95%). Numbers in fraction: number of sibling cells over total cells in the cell type.

glia+endothelia: astrocytes, oligodendrocytes, polydendrocytes, ependymal cells, endothelial
cells.

773
Figure 4. **Sibling contributions to brain microglial populations vary across an animal's**
brain areas. Contributions of sibling(s) to the microglial populations ascertained in principal
brain areas. CJ001, CJ023 and CJ028 were part of a triplet litter and the chimerism contribution
of each twin is shown in a separate panel. CJ102 was profiled in only one brain region and
hence was not included in the analysis. Brain regions with missing data were not profiled in that
animal. y-axis: fraction of twin cells. Vertical bars: binomial confidence interval (95%); *P*-values:
test of heterogeneity across an animal's brain regions.
781

Figure 5. Utilizing natural chimerism to distinguish cell-autonomous from non-cell-

autonomous effects on gene expression, and to compare the effects context and genetic
variation in shaping gene expression. **(A-C)** Comparisons of RNA expression between
microglial populations within host animal CJ028, who had two birth siblings. Comparisons of
gene expression between microglia with the genomes of **(A)** the female host and male sibling,
(B) the female host and female sibling, and **(C)** the two siblings (male and female). In (A) to (C),
each point represents a gene; its location on the plot represents the level of expression of that
gene among microglia with two different genomes in the same animal. x- and y-axes:
normalized gene expression levels (number of transcripts per 100,000 transcripts). FC: fold-
change of gene expression, female/male for *XIST*. Fold-change and *P*-values were calculated
using the binomTest method from the edgeR package (Robinson et al., 2010). Differentially
expressed genes (black dots) were defined as: FDR *Q*-value<0.05 and fold-change>1.5 (in
either direction) and the gene must be expressed in at least 10% of at least one of the two sets
of microglia being compared. **(D-I)** Higher effect of context than genetic differences in shaping
gene expression. **(D)** In the brain of marmoset CJ027, the neocortex and striatum are two
contexts where two sets of microglia with different genomes reside. **(E)** Effect of genetic
differences. x-axis: log2-fold-change of cortical microglia gene expression between host and
sibling cells; y-axis: log2-fold-change of striatal microglia gene expression between host and
sibling cells. **(F)** Effect of context. x-axis: log2-fold-change of the sibling's gene expression
between the two brain regions; y-axis: log2-fold-change of the host's gene expression between
the two brain regions. **(G-I)** The brains (cortex, striatum and hippocampus) of two birth siblings
provide biological contexts in which populations of microglia with two sibling genomes reside.
The effect of genetic differences (H) and effect of animal context (I) are compared, for the same
brain areas (combined data from cortex, striatum and hippocampus). x- and y-axes: log2-fold-
changes of the gene expression between two sets of microglia (the sets being compared are
indicated in the axis labels). *R*: Spearman correlation.
809
810
811

812 **Table 1.** Marmosets analyzed with snRNA-seq in this study. Colonies: NEPRC - New England
813 Primate Research Colony; CLEA - Central Institute for Experimental Animals, Japan; Company
814 A: marmosets obtained from a non-clinical contract research organization.

ID	Tissue	Sex	Age	No. birth siblings (sex of sibling(s))	Colony	Single-cell platform
CJ001	Brain	F	Neonate	2 (M, M)	NEPRC	Drop-seq
CJ006	Brain	M	Neonate	2 (F, unknown)	CLEA	10X
CJ007	Brain	F	Neonate	2 (M, unknown)	CLEA	10X
CJ022	Brain	F	2 yrs 8 mos	1 (M)	NEPRC	Drop-seq, 10X
CJ023	Brain	F	2 yrs	2 (F, M)	NEPRC	Drop-seq
CJ025	Brain	M	2 yrs	1 (M)	NEPRC	Drop-seq
CJ026	Brain	M	2 yrs 8 mos	1 (F)	NEPRC	Drop-seq
CJ026	Blood					10X
CJ026	Liver					10X
CJ026	Kidney					10X
CJ027	Brain	M	2 years	1 (M)	CompanyA	10X
CJ027	Blood					10X
CJ028	Brain	F	2 years	2 (M, F)	NEPRC	10X
CJ028	Blood					10X
CJ029	Brain	M	2 years	1 (M)	CLEA	10X
CJ102	Brain	F	6 mos	1 (M)	CompanyA	Drop-seq, 10X

816 **Supplementary Table 1.** Number of microglia and macrophage cells identified in brain
817 datasets and number of nuclei profiled in blood, liver and kidney.

ID	Tissue	Microglia	Macrophage	Other Cell Types	Total nuclei	Percent Microglia	Percent Macrophage
CJ001	Brain	1,063	16	89,599	90,678	1.17%	0.02%
CJ006	Brain	632	259	165,272	166,163	0.38%	0.16%
CJ007	Brain	1,155	139	105,407	106,701	1.08%	0.13%
CJ022	Brain	3,659	228	217,312	221,199	1.65%	0.10%
CJ023	Brain	581	38	79692	80,311	0.72%	0.05%
CJ025	Brain	1,366	131	210628	212,125	0.64%	0.06%
CJ026	Brain	693	26	62743	63,462	1.09%	0.04%
CJ027	Brain	8,398	167	342,419	350,984	2.39%	0.05%
CJ028	Brain	18,185	172	478,765	497,333	3.66%	0.03%
CJ029	Brain	11,874	285	320,000	332,159	3.57%	0.09%
CJ102	Brain	1,124	43	56,602	57,769	1.95%	0.07%
CJ026	Blood	NA	NA	1,741	1,741	NA	NA
CJ026	Liver	NA	NA	10,877	10,877	NA	NA
CJ026	Kidney	NA	NA	9,262	9,262	NA	NA
CJ027	Blood	NA	NA	2,529	2,529	NA	NA
CJ028	Blood	NA	NA	10,042	10,042	NA	NA

819 **Supplementary Table 2**. Comparison of chimerism between CJ028's two birth siblings, in blood
820 and in brain myeloid cells (microglia and macrophage). *P*-values are from a two-sided test of
821 proportions between chimerism fractions of sibling 1 and sibling 2 using the prop.test function in
822 R.

Cell type	Total nuclei	Sibling 1 nuclei	Sibling 2 nuclei	Sibling 1 Fraction	Sibling 2 Fraction	Sibling 1 + Sibling 2 Fraction	*P*-value
microglia and macrophage	19701	6922	2689	0.35	0.14	0.49	$<2.2 \times 10^{-16}$
microglia only	19447	6873	2621	0.35	0.13	0.49	$<2.2 \times 10^{-16}$
macrophage only	254	49	68	0.20	0.27	0.46	5.8×10^{-2}
blood	10042	1758	6739	0.18	0.67	0.85	$<2.2 \times 10^{-16}$

823
824

Supplementary Table 3. Summary of context versus genetic effects analysis, with two brain
regions of an animal as two contexts. The analysis described in Fig. 5D-F was repeated across
all animals and brain regions with at least 60 cells that are available for analysis in each context,
and the summary of the correlations are tabulated here. The correlations are plotted in
Supplementary Fig. 5. Abbreviations; STR: striatum; Thal: thalamus; Hippo: hippocampus; Hyp:
hypothalamus; BF: basal forebrain.

Host ID	Sibling ID	Brain Region1	Brain Region2	Genetic Effects Spearman Correlation	Context Effects Spearman Correlation
CJ022	CJ106	Cortex	Thal	0.1	0.73
CJ025	CJ104	Cortex	STR	0.03	0.2
CJ027	CJ140	Cortex	Thal	0.13	0.64
CJ027	CJ140	Cortex	STR	0.14	0.69
CJ027	CJ140	Cortex	Hippo	0.08	0.46
CJ027	CJ140	Cortex	Hyp	0.03	0.68
CJ027	CJ140	Thal	Hippo	0.04	0.56
CJ027	CJ140	Thal	Hyp	0.09	0.66
CJ027	CJ140	STR	Hippo	0.08	0.43
CJ027	CJ140	STR	Hyp	0.08	0.59
CJ027	CJ140	Hippo	Hyp	0.04	0.51
CJ028	CJ141	Cortex	Thal	0.1	0.66
CJ028	CJ142	Cortex	Thal	0.03	0.55
CJ141	CJ142	Cortex	Thal	0.07	0.59
CJ028	CJ141	Cortex	STR	0.13	0.68
CJ028	CJ142	Cortex	STR	0.1	0.58
CJ141	CJ142	Cortex	STR	0.09	0.61
CJ028	CJ141	Cortex	Hippo	0.06	0.39
CJ028	CJ142	Cortex	Hippo	0	0.33
CJ141	CJ142	Cortex	Hippo	0.04	0.38
CJ028	CJ141	Cortex	BF	0.13	0.73
CJ028	CJ142	Cortex	BF	0.07	0.6
CJ141	CJ142	Cortex	BF	0.09	0.6
CJ028	CJ141	Cortex	Hyp	0.06	0.78
CJ028	CJ142	Cortex	Hyp	0.07	0.72
CJ141	CJ142	Cortex	Hyp	0.08	0.75
CJ028	CJ141	Cortex	Amygdala	0.14	0.65
CJ028	CJ142	Cortex	Amygdala	0.1	0.53
CJ141	CJ142	Cortex	Amygdala	0.12	0.57
CJ028	CJ141	STR	Hippo	0.06	0.63
CJ028	CJ142	STR	Hippo	0.01	0.56

CJ141	CJ142	STR	Hippo	0.03	0.61
CJ028	CJ141	STR	BF	0.14	0.63
CJ028	CJ142	STR	BF	0.06	0.5
CJ141	CJ142	STR	BF	0.05	0.52
CJ028	CJ141	Hyp	Amygdala	0.14	0.68
CJ028	CJ142	Hyp	Amygdala	0.21	0.65
CJ141	CJ142	Hyp	Amygdala	0.07	0.68
CJ029	CJ143	Cortex	Thal	0.09	0.61
CJ029	CJ143	Cortex	STR	0.06	0.5
CJ029	CJ143	Cortex	Hippo	0.05	0.62
CJ029	CJ143	Cortex	BF	0.09	0.79
CJ029	CJ143	Cortex	Hyp	0.07	0.78
CJ029	CJ143	Cortex	Amygdala	0.07	0.32
CJ029	CJ143	Thal	STR	0.09	0.48
CJ029	CJ143	Thal	Hippo	0.06	0.71
CJ029	CJ143	Thal	BF	0.14	0.75
CJ029	CJ143	Thal	Hyp	0.13	0.8
CJ029	CJ143	Thal	Amygdala	0.05	0.52
CJ029	CJ143	Hippo	BF	0.06	0.77
CJ029	CJ143	Hippo	Hyp	0.03	0.78
CJ102	CJ103	Cortex	STR	0.07	0.32

831
832
833

Supplementary Table 4. Clustering parameters used to identify microglia and macrophage cell
types, and thresholds for identifying host-sibling doublets. The final number of microglia and
macrophages after the second round of clustering are in Supplementary Table 2. A cell is
assigned as a doublet if the Dropulation tool DetectDoublets assigned the highest likelihood for
the cell as a doublet and if the log10 of the best likelihood minus the log10 of the second-best
likelihood (lrt_test_stat, calculated by DetectDoublets) is greater than the doublet detection
threshold (last column).

Marm oset ID	Tissue	Resolution parameter	Nearest neighbor parameter	UMI threshold (min log-UMI per cell)	Doublet Detection Threshold (lrt_test_stat)
CJ001	Brain	0.8	10	1.2	0.0
CJ006	Brain	0.5	10	1.2	5.0
CJ007	Brain	0.5	20	1.2	10.0
CJ022	Brain Drop-seq	0.5	20	1.2	2.0
CJ022	Brain 10X	0.5	20	1.5	2.0
CJ023	Brain	0.5	20	1.2	3.0
CJ025	Brain	0.5	20	1.0	0.5
CJ026	Brain	0.5	20	1.0	2.0
CJ026	Blood	0.5	20	1.4	4.0
CJ026	Liver	0.5	20	1.9	8.0
CJ026	Kidney	0.5	20	1.9	8.0
CJ027	Brain Cortex A	0.5	20	1.5	20.0
CJ027	Brain Cortex B	0.5	20	1.5	20.0
CJ027	Brain others	0.5	20	1.5	20.0
CJ027	Blood	0.5	20	1.5	1.0
CJ028	Brain Cortex A	0.5	20	1.5	25.0
CJ028	Brain Cortex B	0.8	10	1.5	25.0
CJ028	Brain others	0.5	20	1.5	25.0
CJ028	Blood	0.5	20	1.5	1.0
CJ029	Brain Cortex	0.5	20	1.5	20.0
CJ029	Brain others	0.5	20	1.5	20.0
CJ102	Brain, Drop-seq	0.8	10	1.2	3.0
CJ102	Brain, 10X	0.5	20	1.2	5.0

841
842
843

844 **Supplementary Table 5**. Whole genome sequencing datasets used in (1) donor-of-origin
845 assignment from snRNA-seq (Dropulation), and (2) estimating chimerism from blood whole
846 genome sequencing (Census-seq).

ID	Tissue/DNA source	Sequencing coverage
CJ001	Fibroblast culture	36.4X
CJ119 (CJ001's sibling1)	Fibroblast culture	39.7X
CJ120 (CJ001's sibling2)	Fibroblast culture	36.7X
CJ006 (CJ007's sibling)	Fibroblast culture	45.9X
CJ007 (CJ006's sibling)	Fibroblast culture	55.8X
CJ022	Fibroblast culture	38.4X
CJ106 (CJ022's sibling)	Buccal swab	36.0X
CJ023	Fibroblast culture	34.9X
CJ131 (CJ023's sibling1)	Fibroblast culture	39.5X
CJ116 (CJ023's sibling2)	Fibroblast culture	28.3X
CJ025	Fibroblast culture	72.7X
CJ104 (CJ025's sibling)	Buccal swab	58.0X
CJ026	Fibroblast culture	74.1X
CJ105 (CJ026's sibling)	Buccal swab	43.7X
CJ027	Fibroblast culture	47.5X
CJ027	Blood	46.9X
CJ140 (CJ027's sibling)	Fibroblast culture	40.2X
CJ028	Fibroblast culture	44.6X
CJ028	Blood	52.6X
CJ141 (CJ028's sibling1)	Fibroblast culture	43.6X
CJ142 (CJ028's sibling2)	Fibroblast culture	41.2X
CJ029	Fibroblast culture	49.2X
CJ143 (CJ029's sibling)	Fibroblast culture	40.8X
CJ102	Brain	32.4X
CJ103 (CJ102's sibling)	Buccal swab	38.5X

848
Supplementary Figure 1. Microglia and brain macrophages can be identified in all
animals. Expression of microglia and macrophage markers in microglia (left sub-panels),
macrophages (middle sub-panels) and all other cell types in the brain (right sub-panels;
neurons, glia (astrocytes, oligodendrocytes, polydendrocytes, ependymal cells), and endothelial
cells of each animal. Marmosets CJ022 and CJ102 were profiled using two technologies (DS:
Drop-seq, 10X: 10X Chromium). y-axis: unique molecular identifier (UMI, a measure of
transcript abundance) of each gene across cells, summed and normalized to 100,000

transcripts.   Microglia markers: *TREM2, LAPTM5, C3*; macrophage markers: *F13A1, MSTN,*
LYVE1.

Supplementary Figure 2.   Donor-of-origin assignments from brain snRNA-seq reveals
only microglia and macrophages are chimeric.   (A-L) Donor of origin (Dropulation)
assignments of each nucleus from brain snRNA-seq of 10 animals.   Marmosets CJ022 and
CJ102 were profiled using two technologies (DS: Drop-seq, 10X: 10X Chromium). For each
marmoset, the snRNA-seq data are grouped into microglia, macrophage, and all other cell types

(others: neurons, astrocytes, oligodendrocytes, polydendrocytes, ependymal cells, endothelial
cells). x-axis: number of UMI that contains SNPs for which the host and sibling's genomes differ,
in log scale; y-axis: inferred likelihood that the cell has host genome minus likelihood that the
cell has sibling genome (log10). Nuclei with positive y-values are assigned to the host and
those on the negative y-axes are assigned to the sibling.
870
871

872
873 **Supplementary Figure 3.   Summary of microglia (A) and macrophage (B) chimerism**
874 **across animals.** y-axis: fraction of twin cells. Vertical bars: binomial confidence interval (95%).
875 *P*-values: test of heterogeneity across animals.
876
877

878
879
Supplementary Figure 4. **Sibling contributions to brain microglial populations vary**
across an animal's brain areas. Contributions of sibling(s) to the microglial populations
ascertained in finer-scale brain substructures. CJ001, CJ023 and CJ028 were part of a triplet
litter and the chimerism contribution of each twin is shown in a separate panel. CJ102 was
profiled in only one brain region and hence was not included in the analysis. Brain regions with
missing data were not profiled in that animal. y-axis: fraction of twin cells. Vertical bars:
binomial confidence interval (95%); *P*-values: test of heterogeneity across an animal's brain
regions.

888
Supplementary Figure 5. Comparison of gene expression between microglia with

different genomes in each host animal's brain. Each point represents a gene; its location on
the plot represents the level of expression of that gene among microglia with two different
genomes in the same animal. x- and y-axes: normalized gene expression levels (number of
transcripts per 100,000 transcripts). Fold-change and *P*-values were calculated using the
binomTest method from the edgeR package and differentially expressed genes (black dots)
were defined as: FDR *Q*-value<0.05 and fold-change>1.5 (in either direction) and the gene must
be expressed in at least 10% of at least one of the two sets of microglia being compared.

903
Supplementary Figure 6. Summary of genetic versus context effects. This is a plot of the correlation values from Supplementary Table 5. Abbreviations; STR: striatum; Thal: thalamus; Hippo: hippocampus; Hyp: hypothalamus; BF: basal forebrain

908 **Supplementary Figure 7. Doublet detection using host and sibling genotypes.** The axes
909 are the same as in Supplementary Fig. 2, and each dot is a nucleus. Here, nuclei that were

910   identified as doublets and discarded in analyses were indicated (black dots). Marmosets CJ022
911   and CJ102 were profiled using two technologies (DS: Drop-seq, 10X: 10X Chromium).
912
913
914

915
Supplementary Figure 8. Second round clustering of microglia to discard mis-classified

cells. (A-M) Gene expression comparison between host and sibling cells. Red dots: nuclei
identified as microglia from first-round of clustering, black dots: nuclei that were retained after
second-round clustering. For triplets, only the first sibling is included in the plots. Pearson
correlation as calculated for each set (before and after second round clustering) and shows an
improvement in correlation after discarding mis-classified cells. Marmosets CJ022 and CJ102
were profiled using two technologies (Drop-seq and 10X Chromium). (**N**) Summary (box plot) of
fraction of microglia cells discarded during second round of clustering, for host and birth sibling.
924
925
926
927

930 **Supplementary Figure 9. Analysis of genotyping and chimerism estimates if the whole**
931 **genome sequencing data of the sibling is contaminated by the hosts' DNA. (A)-(F)**
932 Sensitivity and false positives in genotyping; HomRef: homozygous reference, HomAlt:

homozygous alternate allele, Het: heterozygous. **(G)-(L)** Sensitivity and false positives in donor-of-origin assignment. **(M)-(Q)** Microglia chimerism estimates when sibling WGS are contaminated by the hosts' DNA, for 5 brain regions; red horizontal line: chimerism estimates when there are no errors in sibling genotypes. **(R)-(U)** Macrophage chimerism estimates when sibling WGS are contaminated by hosts' DNA; red horizontal line: chimerism estimates when there are no errors in sibling genotypes.