X-Ray Structure and enzymatic study of a Bacterial NADPH oxidase highlight the activation mechanism of eukaryotic NOX

Isabelle Petit-Härtlein[§,1], Annelise Vermot[§,1], Michel Thépaut[§], Anne Sophie Humm[#], Florine Dupeux[§,#], Jerôme Dupuy[§], Vincent Chaptal[£], José A. Marquez[#,∂], Susan M.E. Smith[†*], Franck Fieschi [§,ʄ,*]

[§]Univ. Grenoble Alpes, CNRS, CEA, Institut de Biologie Structurale, Grenoble, France

[†]Department of Molecular and Cellular Biology, Kennesaw State University, GA, USA.

[#] European Molecular Biology Laboratory (EMBL) Grenoble, 71 Avenue des Martyrs, Grenoble, France.

[∂]ALPX S.A.S. 71, Avenue des Martyrs, Grenoble, France

[£] CNRS-Lyon 1 University Laboratory 5086, IBCP, Lyon, France.

[ʄ]Institut Universitaire de France (IUF), Paris, France

1. *These two authors, listed in alphabetical order, contributed equally to the work.*

[*] Corresponding authors: ssmit325@kennesaw.edu, franck.fieschi@ibs.fr

Abstract

NADPH oxidases (NOX) are transmembrane proteins that professionally produce reactive oxygen species (ROS) and are distributed widely in both eukaryotes and prokaryotes. Eukaryotes use the ROS products for innate immune defense and signaling; the seven human isoforms of NOX participate in critical physiological and pathophysiological processes. Recently solved structures of two human NOX isoforms provide much new information, but do not fully elucidate controls on the electron transfer pathway from NAD(P)H substrate through FAD and heme cofactors to the final ROS product. SpNOX, a bacterial NOX homolog from *Streptococcus pneumoniae*, shows robust constitutive activity in detergent solution, making it a good prototype for exploring electron transfer in the NOX family. Here we present crystal structures of wildtype and mutant full-length and dehydrogenase (DH) domain-only constructs of SpNOX. The isolated DH domain acts as a flavin reductase, and both DH and full-length constructs use either NADPH or NADH as substrate. Our data supports hydride transfer from NAD(P)H to FAD as the rate limiting step in electron transfer. Using the DH domain we demonstrate the role of F397 in allowing access of nicotinamide to the flavin isoalloxazine, while in the full length construct we used mutants and flavin analogs to confirm the contribution of both domains to flavin binding observed in the structure. Comparison with homologous enzymes suggests distal heme access may influence the final electron acceptor, while the relative position of DH and TM does not necessarily correlate with activity of a given NOX family member. Based on this comparison, SpNOX appears to be a good model of active NOX2, which allows us to propose an explanation for NOX2's requirement for activation.

Introduction

NADPH oxidases (NOX) are membranous enzymes that 'professionally' produce the reactive oxygen species (ROS) superoxide or H_2O_2 (Vermot et al., 2021); the 'professional' production in NOX contrasts with many other sources of cellular ROS which result from metabolic leaking or byproducts. The first-characterized human enzyme of the NOX family, now called NOX2, has been known since 1978 (Segal and Jones, 1978), and provided a paradigm of its ROS product as a cytotoxic agent in defense mechanisms. However, from 1999 – 2001, six other isoforms of NOX were discovered: NOX1/3/4/5 and DUOX1/2 (Banfi et al., 2001; De Deken et al., 2000; Dupuy et al., 1999; Geiszt et al., 2000; Kikuchi et al., 2000; Suh et al., 1999). These other isoforms of NOX function in many tissues and cell types and the ROS they produce play roles beyond cytotoxic defense, as biochemical and signaling agents. In addition to the identification of NOX2 deficiency as the cause of chronic granulomatous disease (Segal and Jones, 1978, 1979), dysregulation of other NOX family members has since been highlighted in pathophysiologies like hyperthyroidism (De Deken et al., 2000; Dupuy et al., 1999) and inflammatory bowel diseases along with well-documented participation in normal physiological process like cardiovascular tone regulation (Rajagopalan et al., 1996), sense of balance (Banfi et al., 2004), and fertility (Musset et al., 2012) among others. Interestingly, these integrated processes directly link to multicellularity leading to the belief that true NOXes are found only in multicellular eukaryotes (Lalucque and Silar, 2003).

The various isoforms display significant diversity in assembly with accessory membrane proteins (p22[phox] for NOX1-4, DUOXA1/2 for the corresponding DUOX1/2) or not (NOX5); in phosphorylation-promoted association of cytosolic factors required for activation (NOX 1-3) or not (NOX4-5, DUOX1/2); in the presence of EF hands conferring Ca^{2+} dependence (NOX5, DUOX1/2) or not (NOX1-4); and in the species of ROS produced, superoxide for NOX1-3,5 and H_2O_2 for NOX4 and DUOXs enzymes (Vermot et al., 2021).

Despite their variety, all NOX family members share a common catalytic core comprised of two conserved domains. The transmembrane (TM) domain belongs to the Ferric Reductase Domain (FRD) superfamily, which consists of six TM helices containing conserved histidines in TM 3 and 5 that chelate up to two heme b cofactors (Zhang et al., 2013). The cytosolic dehydrogenase (DH) domain belongs to the Ferredoxin NADP+ Reductase (FNR) family which has motifs to bind a flavin cofactor and a pyridine dinucleotide NAD(P)H (Taylor et al., 1993). Thus, the NOX core catalytic domain contains the necessary machinery to move electrons from the electron donor – NADPH – through the flavin and heme cofactors to the ultimate electron acceptor, molecular oxygen, to make the ROS product.

The fusion of the TM (N-terminal) and DH (C-terminal) domains constitutes the conserved architecture for the catalytic subunit of the NOX family members. The widespread appearance of separate FRD and DH domains in prokaryotes motivated a search for prokaryotic NOX homologs. First, separate *E. coli* FRD and FNR representatives MsrQ and Fre were identified and shown to interact, producing a NOX-like flow of electrons from NADPH to flavin and

through the hemes to an electron acceptor (Juillan-Binard et al., 2017). Another bioinformatic search identified a large number of prokaryotic sequences with fused FRD and FNR domains (Hajjar et al., 2017). We demonstrated that one such protein from *Streptococcus pneumoniae*, called SpNOX, is a *bona fide* NOX that produces ROS and shares the spectral signature and biochemical hallmarks (resistance to cyanide, inhibition by DPI) of eukaryotic NOX (Hajjar et al., 2017), although its biological function in *S. pneumoniae* has not yet been determined. These results demonstrate definitively that NOXes also exist in prokaryotes. SpNOX purifies as a monomer and is constitutively active (Hajjar et al., 2017), so it represents the minimal functional core shared by all NOX family members. Furthermore, unlike eukaryotic NOX family members, it overexpresses well in *E. coli* and retains robust activity in detergent solution. These qualities make SpNox well suited to structural determination by several methods.

Published crystal structures of separate TM and DH domains of a NOX5 homolog CsNOX from the cyanobacterium *Cylindrospermum stagnale* (Magnani et al., 2017) provided novel structural information about the NOX family. A low resolution structure of full length SpNOX (Vermot et al., 2020) provided the first look at the fused TM and DH domains in NOX proteins, while cryo-EM structures of mammalian DUOX1 (Sun, 2020; Wu et al., 2021) and NOX2 (Liu et al., 2022; Noreng et al., 2022) furnished important new insights. Still, much remains to be discovered regarding electron transfer, the essential catalytic function of the NOX family. SpNOX thus remains an attractive prototype through which to examine structure/function relationships common to NOX family members.

Here, we obtained X-ray structures of WT and F397W mutant of DH-only, and of a F397W mutant of full-length SpNOX. We combined enzymological and structural characterization to elucidate insights into substrate and cofactor recognition, as well as kinetic and mechanistic aspects of electron transfer for the NOX enzyme family. Comparative analysis of this constitutively active prokaryotic NOX with the previously described human NOX2 and DUOX1 structures provides a molecular explanation NOX2's requirement for an activation step.

Results and Discussion

Production of SpNOX and SpNOX$_{DH}$ domain

SpNOX shares only about 25% identity with the human NOX homologs; nonetheless the NOX family members share several well-conserved motifs identified in eukaryotic NOX by Massari et al (Massari et al., 2022). Figure S1A shows the logos corresponding to the eukaryotic NOX motifs in (Massari et al., 2022), but constructed from our alignment containing about half prokaryotic sequences. Other than the motif identified as NADPH3, for which our joint prokaryotic-eukaryotic NOX alignment lacked consensus, SpNOX (and other prokaryotic NOX) recognizably share all the motifs identified by Massari & al (Massari et al., 2022). The C-terminal motif identified by our alignment includes several residues past the C-terminal end of the motif identified from eukaryotes, indicating the high frequency of additional residues in prokaryotic NOX. A specific acid/base pair ($E_{283}R_{284}$ human NOX2 numbering) highly conserved in eukaryotic NOX appears directly after the predicted end of the 6th TM helix (Sharpe et al., 2010), at the inter-domain hinge (Figure S1B); SpNOX has the similar pair $Q_{180}K_{181}$ at this position which prompted us to define K181 as the start of SpNOX$_{DH}$. A typical SDS-PAGE of purified SpNOX and SpNOX$_{DH}$ is shown in Figure S2A.

Figure 1: Buffer optimization for improvement of SpNOX$_{DH}$ thermostability and activity. (A) Thermal unfolding curves of SpNOX$_{DH}$ (0.2 mg.mL^{-1}) in 50mM B bis TRIS-propane buffer pH 6.5, 300 mM NaCl with various glycerol percentages: 5% (–), 10% (–), 15% (–), 20% (–). (B) Thermal unfolding curves of SpNOX$_{DH}$ (0.2 mg.mL^{-1}) in bis TRIS-propane buffer pH 6.5, 5% glycerol with various NaCl concentrations: 100 mM (–), 200 mM (–), 300 mM (–), 400 mM (–), 500 mM (–), 750 mM (–), 1 M (–). (C) Cytochrome *c* reduction in presence of SpNOX$_{DH}$ (1 μg) NADPH (200 μM) FAD (10 μM) in the standard 50mM bis TRIS-propane buffer at pH 6.5, 300mM NaCl in presence of various glycerol percentages: 0% (–), 5% (–), 10% (–). (D) Cytochrome *c* reductase activity with SpNOX$_{DH}$ (1 μg), NADPH (200 μM), FAD (10 μM), glycerol (5%), in presence of various NaCl concentrations: 0 mM (–), 100 mM (–), 200 mM (–), 400 mM (–), 1 M (–). (E) 3D surface representation of T$_m$ dependence with NaCl concentrations and glycerol percentages. (○) depicts the optimum used for further experiments of this study.

Optimization of the SpNOX$_{DH}$ thermostability and activity

SpNOX$_{DH}$ showed a tendency to aggregate, so we investigated buffer conditions to optimize its stability, solubility and activity (Gekko and Timasheff, 1981; Martins de Oliveira et al., 2018). SpNOX$_{DH}$ thermostability and activity were assessed through wide ranges of salt concentrations (0.05-1M) and glycerol percentage (0-20%). The temperatures of thermal unfolding were determined by nanoDSF and demonstrated a constant improvement of T$_m$ with NaCl concentration and glycerol percentage increases (Figure 1A and 1B). SpNOX$_{DH}$ activity was

assessed by reduction of cytochrome *c*. In reactions without NaCl and glycerol, the DH domain efficiently supported cytochrome *c* reduction (Figure 1Cand 1D). In contrast to thermal stability, however, $SpNOX_{DH}$ exhibits a dramatic loss of activity with increasing NaCl and glycerol. We generated a 3D surface plot of Tm as a function of NaCl and glycerol with specific activity as color scale (Figure 1E) which allowed us to optimize buffer conditions jointly. The region highlighted by (○) in Figure 1E indicates that at 300mM NaCl and 5% glycerol $SpNOX_{DH}$ exhibits both high activity (5 mol reduced cyt. *c*/s/mol.**SpNOX_{DH})** and a good thermostability (melting temperature of 60°C).

We then optimized the pH in order to conjointly maximize the thermostability and the activity of $SpNOX_{DH}$. Direct comparison of the kinetic parameters of $SpNOX_{DH}$ and SpNOX necessitated identifying a pH that would work well for both constructs. The dependences of **SpNOX_{DH}** activity (Figure S3A) and stability (Figure S3B), evaluated along a range of pH (respectively 5-7 and 4-9.5) in the presence of 300 mM NaCl 5% glycerol, exhibit a progressive decrease with increasing alkalinity. Like $SpNOX_{DH}$, the *thermostability* of SpNOX follows a similar and progressive decrease of T_m with alkalinity (Figure S3B), but in contrast the *activity* of SpNOX shows an optimum at higher alkalinity (Figure S3A). The two constructs show similar activity at pH 6.5, so we chose this as the pH. Thus, for all experiments in which we compare activity of the full-length SpNOX to the truncated $SpNOX_{DH}$, we chose a standard buffer of 50mM bis TRIS-propane pH 6.5, 300 mM NaCl, 5% glycerol.

Figure 2: Effect of cyt. *c*, FAD, SOD and DPI supplementation on SpNOX_{DH} NADPH oxidase activity. Assays performed with NADPH (200 µM) and $SpNOX_{DH}$ (1µg) in the initial mix. Monitoring of cytochrome *c* reduction (cyt.*c* reductase activity) (A) or NADPH oxidation (flavin reductase activity) (B) along with successive addition of FAD and SOD (A and B) or DPI (C) with cyt.c (blue trace) or without (grey trace) in the initial mix. (D) Monitoring simultaneously NADPH oxidation (red) and cytochrome c reduction (blue) in presence of $SpNOX_{DH}$, cytochrome *c*, NADPH and FAD, addition of SOD is indicated by an arrow. The table presents the

molecular specific activity measured for the different compositions of the initial mixture and addition of the multiple reactants.

Functional characterization of SpNOX$_{DH}$

The DH domain of NOXes, including SpNOX, as a homolog of the FNR superfamily, should catalyze the transfer of two electrons from the cytosolic NADPH to the FAD coenzyme; in full length NOX the electrons are then transferred singly to the hemes and across the membrane to molecular O$_2$. The functionality of the DH domain of SpNOX was evaluated by monitoring the oxidation of NADPH substrate and also by monitoring ferricytochrome c reduction which in the absence of the TM domain indicates the capacity of FAD to transfer electrons to soluble O$_2$. Superoxide generation was assessed in aerobic conditions using SOD-inhibitable ferricytochrome c reduction.

Like purified or membrane preparations of both the full length and DH constructs of other NOX family members (Han et al., 2001; Nisimoto et al., 2004, 2010), SpNOX$_{DH}$ requires supplemental FAD to sustain superoxide production (Figure 2A) and NADPH oxidation but does not require the addition of Cyt. *c* (Figure 2B). The dependence of SpNOX$_{DH}$ on flavin in the oxidation of NADPH was assessed by addition of diphenyleneiodonium (DPI), a common specific flavoenzyme inhibitor (Nisimoto et al., 2014), which inhibited the enzyme activity (Figure 2C).

In Figure 2D NADPH oxidation and superoxide generation by SpNOX$_{DH}$ was followed simultaneously at both 340 and 550 nm. The addition of superoxide dismutase (SOD) led to a ~40% inhibition of SpNOX$_{DH}$'s Cyt. *c* reductase activity, confirming that the reduction of Cyt. *c* observed in the system is partly due to O$_2^{\bullet-}$ generation (Figure 2D and Table I). However, the remaining ~60% Cyt. *c* reductase activity suggests direct electron transfer from the FADH$_2$ produced by the SpNOX$_{DH}$ to the Cyt. *c*. Thus, SpNOX$_{DH}$, separated from the TM domain, retained the redox properties and sensitivity displayed in the context of the whole SpNOX protein (Hajjar et al., 2017). The strict requirement of FAD addition for SpNOX$_{DH}$ activity suggests that the flavin behaves as a co-substrate rather than a prosthetic group. As an isolated domain, SpNOX$_{DH}$ may work as a flavin reductase enzyme, making an interesting parallel with the *E.coli* flavin reductase *fre* (also in the FNR family) which reduces the MsrQ transmembrane component (in the FRD family, like the NOX TM domain, see Figure 6) (Caux et al., 2021; Juillan-Binard et al., 2017).

Comparative flavin recognition properties and activities of SpNOX$_{DH}$ vs SpNOX

To characterize the flavin reductase activity of the DH domain and the relative contributions to FAD binding of the TM and DH domains of SpNOX, we determined the K_m and k_{cat} of SpNOX and SpNOX$_{DH}$ cytochrome *c* reductase activity using flavins FAD, FMN, riboflavin, and lumiflavin (Table 2). We observed a progressive loss of affinity for SpNOX as flavin size diminishes, with 263-fold higher affinity for FAD than for lumiflavin (Table 2). In contrast, the SpNOX$_{DH}$ retains constant affinity equal to that of lumiflavin (~10µM) regardless of the flavin analog. These results suggest that the isolated DH domain binds to FAD primarily at the isoalloxazine ring of the cofactor, and the addition of ribitol (riboflavin), phosphate groups (FMN), and adenosine nucleotide (FAD) do not further interact significantly with the isolated

DH domain. However, the full-length SpNOX appears to interact along the entire length of the FAD cofactor, indicating significant interactions between the FAD and the TM domain. This FAD binding site shared between the TM and DH domains may contribute to the correct placement of the cofactor for electron transfer to the proximal heme. On the other hand, in full length SpNOX the k_{cat} appears to be independent of the flavin derivative used, suggesting that the binding interactions between TM and FAD play no catalytic role. Finally, the k_{cat} of full length and DH-only constructs are the same order of magnitude, with ratios around 1 (Table 2). This is a strong argument that the initial hydride transfer from the nicotinamide is the rate limiting step in the electron transfer from substrate to O_2.

	K_m (µM)			k_{cat} (s^{-1})		
	SpNOX	SpNOX$_{DH}$	K_{mDH}/K_m	SpNOX	SpNOX$_{DH}$	k_{catDH}/k_{cat}
FAD	0.049 ± 0.01	12.89 ± 0,69	263	4.68 ± 0.74	6.73 ± 0.74	1.44
FMN	0.92 ± 0.01	7.89 ± 0.48	8.6	3.01 ± 0.04	5.14 ± 0.14	1.70
Riboflavin	8.28 ± 0.01	16.76 ± 1.47	2	7.37 ± 0.38	2.77 ± 0.16	0.38
Lumiflavin	16.56 ± 2.77	15.85 ± 1.07	1	4.67 ± 0.45	4.14 ± 0.11	0.89

Table 2: SpNOX and SpNOX$_{DH}$ affinity for flavins. K_m and k_{cat} of SpNOX and SpNOX$_{DH}$ for FAD, FMN, riboflavin and lumiflavin. K_{mDH}/K_m and k_{catDH}/k_{cat} indicate the ratio between K_ms or k_{cat}s measured on the DH domain and on the full-length SpNOX protein. Figure S4 shows the kinetic studies corresponding to the parameters presented here.

SpNOX uses NADPH and NADH with similar affinity and activity

The typical Michaelis Menten type kinetics parameters of SpNOX and SpNOX$_{DH}$ were obtained for increasing NAD(P)H concentrations (Table 3). Full length and DH constructs showed the same order of magnitude affinity to each other for NADPH or NADH as substrate (Table 3). Furthermore, the value of 1 for NADPH/NADH affinity for each construct shows that SpNOX does not discriminate significantly between NADPH and NADH; the agreement of two different assays – cytochrome c reduction and NADPH oxidation -- strengthens this observation. Thus in SpNOX, the NAD(P)H binding energy arises from interactions just in the DH domain with little contribution from the TM domain of SpNOX. The lack of discrimination

of SpNOX between NADH and NADPH represents a significant difference from eukaryotic NOX which strongly select NADPH as substrate. In these experiments that used a range of NAD(P)H concentration, k_{cat}'s of the same order of magnitude for both constructs again supports hydride transfer as the limiting step in NOX electron flow.

Structural characterization of SpNOX and SpNOX$_{DH}$: mutation of F397

We obtained large hexagonal crystals of SpNOX (See Figure S5) but with poor diffraction power beyond 5 Å, and only 6 % between 3 to 5 Å, with clear anisotropy. We therefore tried stabilizing mutations based on the work of Deng *et al* (Deng et al., 1999) in pea FNR, homologous to the NOX DH domain. They showed that the FNR C-terminal Y308 stabilizes the isoalloxazine site of the FAD cofactor and regulates access of the NADP+ substrate to this site. We reasoned that changing the corresponding residue of SpNOX to W would stabilize FAD binding, effectively locking the two subdomains of the DH domain in place and reducing mobility. Supporting this approach, Magnani et al reported the stabilizing effect in CsNOX of the serendipitous insertion of a Trp that stacks with the FAD isoalloxazine ring (Ceccon et al., 2017; Magnani et al., 2017). Similarly, we reasoned that changing the corresponding F residue to S would increase NADPH access to FAD, potentially facilitating crystals with NADPH bound.

Based on sequence alignment (Figure S1), either of two phenylalanine residues in SpNox, F397 and F399, could correspond to the conserved C-terminal residue in FNR. We mutated both positions to tryptophan in the SpNOX$_{DH}$ construct and monitored activity to test their effects. In our standard buffer used for comparing DH with full-length constructs, F399W mutants showed normal activity (Table 3), while F397W mutants showed no activity (Fig 3). We interpreted this to mean that W in position 397 prevents hydride transfer by immobile stacking with the isoalloxazine ring of FAD, while W in position 399 has little effect on hydride transfer. These results indicate that F397 of SpNOX corresponds to the terminal Y of FNR.

Figure 3 : Flavin content and activity measurement of full-length SpNOX WT, SpNox F397S and F397W. A) UV-Visible spectra of SpNOX$_{DH}$ WT, and mutants. Inset) zoom on the 320-500 nm window, a specific spectroscopic feature of the oxidized flavin isoalloxazine ring. B) Reduced cytochrome c was monitored at 550 nm for SpNOX WT (blue), F397W (grey), and F397S (green). A no-protein negative control is shown in black. See also Fig.S6 for spectral characterization of full length SpNOX mutants.

We subsequently explored mutations of F397 in both full length and DH constructs of SpNOX. The flavin content of purified F397W is much higher than that of WT, while the F397S is not able to retain flavin in its binding site (Figure 3A), consistent with our hypothesis about flavin binding. As displayed in Fig 3B, in our standard activity conditions, F397W has little or no activity while F397S shows enhanced activity compared to WT. Increasing the NADPH concentration 10-fold resulted in low but significant activity of the F397W mutant (not shown). These results indicate that the W and S mutations of position 397 do not significantly affect the structural integrity of the SpNOX constructs. We used Michaelis-Menten experiments to test the affinity for both NADPH and NADH of F397S mutants of full-length enzyme and DH constructs; such experiments were not possible with F397W because of its very low activity (Figure 3). F397S mutants show similar results to WT (Table 3) in both constructs indicating that S in this position does not greatly impact affinity for either substrate; this mutation produced an increase in k_{cat} for both constructs, consistent with the hypothesis that S in this position provides increased access of substrate to the catalytic site.

Substrats	Assays	K_m (μM)			k_{cat} (s^{-1})	
		SpNOX	SpNOX$_{DH}$		SpNOX	SpNOX$_{DH}$
		Cyt.C red	Cyt.C red	FAD red	Cyt.C red	Cyt.C red
NADPH	**WT**	**41.5 ± 5**	**106 ± 31**	**87.89 ± 13**	**6.42 ± 0.6**	**5.42 ± 1.15**
	F399W	n.d.	86.7 ± 6.5	-	n.d.	9.07 ± 0.5
	F397S	33.76 ± 5	79.3 ± 19.5	-	10.80 ± 0.5	8.29 ± 1.3
NADH	**WT**	**41.81 ± 4.4**	**133 ± 21**	**80.27 ± 9.1**	**3.5 ± 0.2**	**4.41 ± 0.5**
	F399W	n.d.	148.6 ± 46	-	n.d.	8.65 ± 2
	F397S	25.46 ± 2	66.9 ± 10.3	-	10.3 ± 0.4	6.20 ± 0.5
WT NADPH/NADH		1	0.8	0.96	1.8	1.2

Table 3: K_m and K_{cat} using NADPH or NADH as electron donor for the SpNOX WT, F399W and F397S full length constructs or the corresponding versions of the DH domain (SpNOX$_{DH}$). F399W is in italics to indicate that position 399 is not homologous to the terminal aromatic in FNR; the W mutation at the homologous position, F397W, did not have sufficient activity to perform such analysis and thus does not appear in this table. WT NADPH/NADH is the ratio of the K_m or K_{cat} for NADPH vs NADH. For SpNOX$_{DH}$ WT K_ms were determined by monitoring both Cyt. *c* reductase activity (at 550 nm) or flavin reductase activity (at 340nm) when reported. The kinetic studies corresponding to the parameters presented here are in Figure S7.

High resolution structure of the SpNOX Dehydrogenase domain.

The use of the full-length SpNOX F397W construct resulted in about 20% of the crystals produced displaying diffraction limits close to 4 Å, with some crystals showing an anisotropic diffraction between 3 and 3.3 Å in a privileged direction of the crystals, as often observed for membrane proteins (Martin et al., 2023; Robert et al., 2017). Despite the improvement of the crystal quality, the anisotropy precluded resolution without a good model for molecular replacement. However, the soluble SpNOX$_{DH}$ domain represents more than 50% of the full-length enzyme, making it a suitable model for use in molecular replacement. We obtained some crystals of both SpNOX$_{DH}$ WT and F397W, co-crystallized with FAD. Unfortunately, crystals of the SpNOX F397S mutant with or without NADPH could not be obtained.

Structure resolution at 1.94 Å of SpNOX$_{DH}$ F397W *de novo* was possible thanks to co-crystallization in the presence of sodium bromide and collection of a bromide-SAD dataset. We used this structure as a model to solve SpNOX$_{DH}$ WT by molecular replacement at 2.5 Å. As expected, SpNOX$_{DH}$ presents a typical fold of the FNR superfamily of reductase domain containing two sub-domains, the FAD-binding domain (FBD) and an NADPH-binding domain (NBD) (Figure 4A). Both WT and F397W SpNOX$_{DH}$ display 3 molecules per asymmetric unit and the two structures are identical. In the asymmetric unit of SpNOX$_{DH}$ WT, one of the three molecules contains no flavin (Figure S8A). In the asymmetric unit of SpNOX$_{DH}$ F397W, all molecules of the unit contain a well-defined FAD (Figure S8B). These observations agree with flavin contents observed in solution (Figure 3A) and support the hypothesis that the F397W mutant stabilizes FAD binding.

In strong correlation with the affinity data obtained for the flavins shown in Table 2, the DH domain mainly interacts with the isoalloxazine ring through polar contacts with the protein backbone and a few residues (P234, F235 and S23) belonging to the conserved FAD-1 binding motif (Massari et al., 2022) (see Figure S1). The DH domain shows almost no contact with the ribitol and adenosine parts of the FAD. With the exception of possible polar contact of the first phosphate group with the backbone of the protein, the rest of the FAD molecule points out into the solvent, in the direction of the missing TM domain (Figure 4D).

In the wild type, the isoalloxazine ring is sandwiched between FBD residues P234 and F235 on its *si*-face, and the NBD residue F397 on its *re*-face (Figure 4D). Superposition of the different SpNOX$_{DH}$ WT molecules from the crystal asymmetric unit shows that the F397 aromatic ring imperfectly occupies the position in front of the *re*-face of the isoalloxazine ring (Figure 4B), with orientation 40° with respect to the plane of the isoalloxazine ring, and distance at 5.3 Å. In contrast (and as predicted from the homologous FNR structure), the tryptophan of the F397W mutation makes an extensive and much tighter parallel π-stacking interaction at a distance of 3.5 Å from the isoalloxazine ring (Figure 4C). From these observations, WT SpNOX$_{DH}$ could well behave as a flavin reductase, able to use different flavins as substrate through the sole recognition of the isoalloxazine ring. As Fig.3A shows, the F397W mutant may fix the flavin as a cofactor. Figure 4E shows the sidechains of both the F and W residue overlaid, on which is superimposed a putative binding mode of NADPH based on the pea FNR structure (Deng et al., 1999). Trp at position 397 appears to preclude access of the nicotinamide to the isoalloxazine, preventing direct hydride transfer. Phe in this position, however, leaves more space in this pocket and the sidechain has more degrees of freedom, making it feasible for the nicotinamide to displace it and transfer hydride to the isoalloxazine. These observations most likely explain the drastic difference of activity between WT and F397W; they probably also explain the enhanced activity observed in the F397S mutant which provides even more space near the isoalloxazine (Table 3). The increase of the global k_{cat} in the F397S mutant is an additional argument to support hydride transfer as the limiting step in NOX electron transfer.

Figure 4: Crystal structures of the DH domain of SpNOX WT and F397W. A) Superposition of SpNOX$_{DH}$ WT (pale cyan ribbon, PDB: 8qq5) and SpNOX$_{DH}$ F397W (light blue ribbon, PDB: 8qq1); FAD is shown as sticks colored by atom, side chains of residues in position 397 (respectively F and W), are shown as sticks in the same color as the corresponding ribbon. B) and C) Superposition of the three different molecules of the asymmetric unit of WT SpNOX$_{DH}$ (B) and of SpNOX$_{DH}$ F397W (C), zoomed on the interaction of the isoalloxazine ring with the aromatic residue at position 397. D) Close up of the FAD binding site in SpNOX$_{DH}$ WT; polar contacts are shown with dotted lines. Colors as for A. E) SpNOX$_{DH}$ with the F397 (pale cyan sticks) and W397 (light blue sticks) residues superimposed; FAD as in A; and NADPH (salmon sticks) shown based on a superposition of the pea FNR:NADPH complex (PDB: 1qfz) with SpNOX$_{DH}$.

Structure of full length SpNOX.

The high-resolution model of the DH domain of SpNOX allowed resolution of the anisotropic crystals of full length SpNOX-F397W to 3.6 Å. Small extracellular loops limiting crystal contacts together with intrinsic interdomain flexibility of SpNOX are probably the source of the crystal anisotropy due to subtle variation of packing within the crystals (Fig S9). Like the previously reported structures of the NOX family (Liu et al., 2022; Magnani et al., 2017; Sun, 2020; Wu et al., 2021), the crystallographic model of SpNOX harbors the N-terminal TM domain that encompasses 6 membrane-spanning alpha helices chelating two hemes, and the C-terminal cytosolic DH domain (Figure 5A). The density corresponding to the two heme groups is packed in the channel formed by TM helices 2 to 5. The inner and outer heme are respectively

coordinated by the imidazole rings of pairs of histidines H69 (TM3)/H129 (TM5) and H82 (TM3)/H142 (TM5), holding them perpendicular to the plane of the membrane (Vermot et al., 2020). The metal-to-metal distance of the two hemes is ~21.5 Å, while the shortest interatomic distance is ~9.8 Å, between the vinyl groups.

As observed in previously solved structures of NOX enzymes (Liu et al., 2022; Magnani et al., 2017; Sun, 2020; Wu et al., 2021), a similar electron transfer relay across the membrane was mapped. SpNOX has 9.2 Å edge-to-edge distance between the isoalloxazine ring to the inner heme; a conserved aromatic residue, Y136, at equal distance from both hemes (6.9 Å) could assist the electron transfer between the hemes as previously suggested (Magnani et al., 2017).

Figure 5: Structure of the full length SpNOX. A) overall structure of SpNOX F397W at 3.6 Å with ribbons in the TM colored by helix, hemes (red sticks) and FAD cofactor (colored by atom). B) The FAD binding site; FAD as in A, side chains (sticks) of residues involved are labeled and interactions are indicated by yellow dotted lines. C) Electron pathway from FAD to distal heme; for clarity, the DH domain and TM1 and TM6 are omitted. Two alternative electron pathways are indicated, from FAD to heme (both depicted as in A), with the Y136 relay between hemes (black dotted lines) or without the relay (red dotted line); distances in Å are indicated.

FAD binding site in the full length SpNOX: role of the D-Loop region.

We observed electronic density corresponding to the FAD cofactor at the interface of the DH and TM domain. FAD binding in the full-length construct preserves the π-stacking interaction with W397 and the interactions with F235 and P234 observed in the DH-only structure. In addition, the full-length SpNOX structure shows that the adenine extremity of FAD is packed in a pocket including the amino acid F399 from the DH domain– providing a potential role for this residue – as well as residues from the D-loop of the TM domain. Indeed, Y122 and R126 could stabilize the adenine ring simultaneously. Y122 might have dual action through π-stacking with the adenine ring, as well as H-bonding with the phosphate groups through its hydroxyl group (Figure 5B). R126 is at a reasonable distance to bridge with the adenine ring and simultaneously a carboxylate of the proximal heme. However, the resolution of the structure makes the accurate positioning of side chains difficult, particularly in this region which displays large electron densities.

	FAD		FMN		Riboflavin		NADPH	
	K_m	$\frac{K_m^{mut}}{K_m^{WT}}$	K_m	$\frac{K_m^{mut}}{K_m^{WT}}$	K_m	$\frac{K_m^{mut}}{K_m^{WT}}$	K_m	$\frac{K_m^{mut}}{K_m^{WT}}$
WT	0.06 ± 0.01	-	$0.9 \pm 0,3$	-	6.4 ± 0.2	-	13 ± 0.74	-
Y122A	0.38 ± 0.5	6.4	1.5 ± 0.9	1.7	9.6 ± 0.08	1.5	23.8 ± 0.04	1.8
R126A	3.5 ± 0.33	59,1	2.9 ± 0.48	3.2	6.63 ± 0.38	1.03	24.4 ± 0.45	1.87

Table 4. Determination of K_m for flavins as a function of the mutations in the D-loop region. Data here were obtained in the optimal buffer for SpNOX rather than the co-optimized buffer for comparing full-length and DH constructs. It has no impact on flavin K_m determination (compare WT line with values from Table2) but K_m of full length SpNOX for NAD(P)H are lower by a factor 2 to 3 compared to the co-optimized buffer. For the corresponding curve see figure S10.

To check the validity of our side-chain observations we generated alanine mutants of Y122 and R126 and checked the affinity of several flavins for each mutant (Figure 5B and Table 4). We determined affinity for NADPH for all mutants; similar affinity in all mutants suggests no effect of these mutations on global enzyme structure (Table 4). The results in Table 4 shows that only the affinity for FAD is impacted by Y122A and R126A, confirming the role of these residues in the interaction with the FAD's adenine ring, which is missing in the other flavins tested. Y122A diminishes FAD affinity by a factor of 6 and R126A by a factor of 23. This latter suggests that R126 is crucial for FAD affinity; possibly the simultaneous interaction between R125 and both heme and flavin (proposed above) could hold the adenine ring in a position to be sandwiched between Y122 on one side and F399 on the other. Surprisingly, while the position corresponding to Y122 is very highly conserved as an aromatic residue in the full NOX enzyme family, the position corresponding to SpNOX R126 shows a difference between eukaryotes and prokaryotes. In eukaryotes, the position corresponding to R126 is very highly conserved as an aromatic, while in prokaryotes it is most often a basic residue, although aromatics are not uncommon. The interactions with the TM domain's D-loop (or lack thereof) provide an explanation for the differences among the affinities of the flavins for the full-length protein (Table 2).

Access to the distal heme: diverse situations from prokaryotic to eukaryotic enzymes

The FRD found in NOXs but also in MsrQ and STEAP proteins, transfers electrons from the cytoplasmic to the extracellular side of the cell. Thus, access of the distal heme is essential to the specificity of the electron acceptor and its product. Figure 6 compares the distal heme access of a panel of FRD-containing proteins from prokaryotes and eukaryotes. From the structures available to date, including the present study, some FRDs (MsrQ, SpNOX and CsNOX) exhibit small extracellular loops and free access to the distal heme (Fig 6A), while others (NOX2) have long extracellular loops (Fig 6A) and sometimes another extracellular domain (DUOX) (Fig 6B) create a cap that limits access. Homology models of both NOX4 (O'Neill et al., 2018) and NOX1 (Ward et al., 2023) also predict similar caps. In NOX with large caps, the distal hemes are deeply masked, implying that O_2 must diffuse into the protein. On the basis of structures, several potential tunnels have been proposed for NOX2 and DUOXs (Liu et al., 2022; Noreng et al., 2022; Sun, 2020; Wu et al., 2021) and for the homology model of NOX1 (Ward et al., 2023). For most of the tunnels identified, bottleneck constriction would require protein dynamics to allow diffusion and the exact pathway for O_2 still remains to be defined. However, a potential entry site with direct access to the distal heme from the lateral side can be seen in the NOXs with large caps (Figure 6B). In addition, an internal pocket in close proximity to the

distal heme has been proposed as the site of O_2 reduction through an outer sphere electron transfer mechanism. This pocket is characterized in eukaryotic enzymes by a conserved His and Arg involved in O_2 binding and heme interaction; the residues characteristic of the O_2 reducing center in eukaryotic FRD domains of NOX and DUOX enzymes are not conserved in SpNOX. The importance of the conserved His and Arg in the reducing center pocket of eukaryotes are strongly supported by CGD-induced mutations (Cross et al., 1995; Picciocchi et al., 2011; Rae et al., 1998). A pathological NOX1 variant was used to show that an additional Asn residue, separated from the conserved His by two residues, is necessary for efficient O_2 reduction (Ward et al., 2023); this Asn is highly conserved in animal NOX (see Helix 3 motif in Fig. S1) but not prokaryotes. Entry site, tunnel size and reducing center pocket location ensure the selectivity to O_2 as unique electron acceptor for NOXs with large caps.

Figure 6: FRD containing proteins, represented as surfaces, and accessory proteins (p22[phox] and DUOXA1) represented as ribbons. TM2 to TM5 were structurally aligned; all proteins are depicted in the same orientation based on this structural alignment. A) Top (extracellular) views of the TM domains of MsrQ (AlfaFold model, green), SpNOX (this work, PDB: 8qq7, gray), TM domain of CsNOX (PDB: 5O0t, purple), NOX2/p22[phox] (PDB: 8gz3, blue) and DUOX1 (PDB: 7d3f, pink) B). Each protein's lateral side view (a 90°C rotation from the orientation in A). Black arrow shows potential access to distal heme either from top or from lateral side. Hemes are represented as salmon sticks.

FRD containing proteins with direct access to the distal heme may reflect more versatility with regard to final electron acceptors. MsrQ, to date the only FRD-containing prokaryotic protein with clearly identified physiological function, provides an interesting case. In *E.coli*, MsrQ transfers its electrons to MsrP, a periplasmic methionine sulfoxide reductase, which repairs proteins damaged by methionine oxidation (Gennaris et al., 2015). The absence of long extracellular loops allows MsrP to dock on MsrQ and to receive electrons directly from the

distal heme to its molybdopterin center. Similarly, SpNOX's exposed heme may confer its ability to directly reduce Cyt. c (Hajjar et al., 2017). The physiological functions of neither SpNOX nor CsNOX are known; in addition to their NOX activity, they may use electron acceptors other than O_2.

Domain orientation in SpNOX structure: comparison with other NOX structures.

The previous crystal structure of a prokaryotic NOX was in fact two independent structures of the separate TM and DH domains of CsNOX (Magnani et al., 2017). With the SpNOX structure presented here we provide the first entire prokaryotic NOX structure, affording a view of the orientation of the TM domain relative to the DH. The importance of the relative orientation of the domains in NOX proteins has been underlined as a critical point in the activation and the efficiency of electron transfer (Liu et al., 2022; Wu et al., 2021). Comparison of the recent structures of an inactive "resting state" NOX2 and of the presumably active "high Ca^{2+} state" DUOX1, emphasized the large 49° rotation of the DH domain of one structure relative to the other. This motion leads to a 12.4 Å movement of the FAD, which was suggested as critical for the activation of the electron transfer (Liu et al., 2022). Including SpNOX into this movement analysis is all the more interesting given that SpNOX is constitutively active. Here, Figure 7A shows that in our SpNOX structure, the relative position of TM *vs* DH domain is intermediate between the inactive (NOX2) and active (DUOX) structures. Figure 7B details the potential intermediate motions of SpNOX: the relative position of SpNOX DH with respect to the inactive state NOX2 DH rotates 13° and translates 5 Å towards the plane of membrane, while the relative position of the high Ca^{2+} "activated" state of DUOX1 DH rotates almost an additional 49° and translates an additional 7.6 Å. SpNOX's domain orientation is much closer to that observed in inactive NOX2 than that of DUOX1.

The relative positions of the cofactors involved in the first steps of electron transfer are highlighted in Figure 7C. The distance between hemes in a given FRD domain is invariant; thus, the efficiency of electron transfer depends on two main factors: the distance between the nicotinamide of NAD(P)H and the isoalloxazine ring of FAD, and the distance between the isoalloxazine ring of FAD and the proximal heme. Surprisingly, there is no evident correlation between DH motion from inactive to active state and contraction of the distance between FAD and heme. In fact, this distance actually increases between inactive NOX2 and constitutively active SpNOX, and from SpNOX to 'activated' (high Ca^{2+}) DUOX. Nonetheless, in all cases these distances, ranging from 7.2 to 10.2 from edge to edge (from methyl group of the flavin ring to the carboxylate of the heme), all remain compatible with biologically relevant electron tunneling probability, calculated as ≤ 14 Å. (Page et al., 1999).

Figure 7: Comparison of domain and cofactor organization in NOX2, SpNOX and hDUOX1. A) Proteins were superimposed and are shown in identical orientations based on superposition, resting state NOX2 (PDB: 8gz3, gray ribbon), active SpNOX (this work, PDB: 8qq7, green ribbon) and high Ca^{2+} DUOX1 in (PDB: 7d3f, red ribbon), with DH domains in darker color. B) and C) DH domains only are represented in same orientations and color as in A. In B, angle and translation from NOX2 to SpNOX$_{DH}$ are indicated. In C, angle and translation from SpNOX$_{DH}$ to DUOX1 are indicated. D) Cofactors necessary for electron transfer are shown as sticks, in identical orientations as in A-C, with distances indicated. In SpNOX the sidechain of Trp397 is shown and the distance to FAD isoalloxazine ring indicated. For DUOX1, NADPH is shown and the distance of nicotinamide to the isoalloxazine indicated.

These observations led us to question the paradigm that switching between resting and activated states requires DH motion. The lack of DH motion between the low and the high Ca^{2+} state human DUOX1 structures adds further hesitation. The DH domains of these two structures superimpose with a RMSD of 0.246 Å, and the FAD and NADPH of these structures also superimpose perfectly. The distance between the nicotinamide ring of the NADPH and the isoalloxazine ring of the FAD even in the activated DUOX1 structure is surprisingly large for efficient hydride transfer (Figure 7C). In line with conclusions from DH homologs (Deng et al., 1999; Hermoso et al., 2002), our data (Table 3, Figures 3 and 7) suggests that the SpNOX F397W sidechain represents the position that would be occupied by the nicotinamide ring of NADPH. The distance between the Trp397 sidechain and the isoalloxazine ring of FAD in our structure is 3.3 Å, similar to the corresponding distance of these moieties in homologous DH domains and suitable for hydride transfer. In the case of DUOX1, it has been suggested that the "high Ca^{2+} activated" state, to be competent in hydride transfer, requires tensing from a relaxed conformation to enhances electron transfer efficiency rather than a large domain motion (Wu et al., 2021).

These considerations suggest that differences in relative domain orientations in NOXes might represent isoform specificity rather than intermediate activation states. If so, these differences and those of the cofactor distances could fine tune the electron transfer properties and resulting efficiency of ROS production, linking the isoform with its physiological context. In this context, the impact of the lipid environment on domain orientation is suggested by longstanding observations of lipid effects on NOX activity (Koshkin and Pick, 1993; Shpungin et al., 1989) and by the observation of a lipid in the murine DUOX structure that mediates the interaction between the TM and DH domain through a direct connection with the phospho-ADP-ribose part of the NADPH (Sun, 2020). Structures solved in detergent like ours presented here may be released from interaction with the membrane, increasing flexibility. The question of what happens in NOX2 –whether DH tensing, domain motion, or something else – remains elusive and awaits further structural characterization. In any case the SpNOX domain orientation presented here makes it a rather close model of human NOX2.

Figure 8: Comparison of FNR domains and their putative NADPH binding sites. Top, pea FNR (PDB: 1qfz), DUOX1 domain (PDB: 7d3f) in complex with NADPH. Middle, SpNOX DH domain (PDB: 8qq7) and NOX2 DH domain (PDB: 8gz3). Bottom, SpNOX DH is represented with the NADPH from pea FNR after superposition of those proteins; NOX2 DH is represented with the NADPH from DUOX1 after superposition of those proteins. Arg residues conserved between DUOX1 and NOX2 are highlighted using CPK colors, Side chain of R446 is not defined in NOX2 CryoEM structure. In top and middle line, pale yellow and lime patches represent NADPH motifs 1 and 4, respectively (see figureS1). Assuming that NADPH binding implies movement of F397, residues ^{397}FKF^{399} have been omitted in the SpNOX DH surface to increase clarity and visibility of NAD(P)H binding.

A Close-up view of NOX's NAD(P)H binding domains *vs* the FNR gold standard

The striking difference in distance between NADPH and FAD at the active site (Figure 7C) parallels a structural difference between NOX2 and DUOX1 on one hand, and SpNOX on the other. Here we use the well-characterized FNR structure as a basis of comparison for DH domains. The NADPH binding groove has characteristic motifs universally conserved in DH domains (Piubelli et al., 2000) (motifs NADPH1 and NADPH4 in Figure S1). In FNR, the binding groove for NADPH is separated from the flavin binding site by a narrow crevice (Figure 8 top left), but in DUOX1 this crevice is much wider (Figure 8, top right). Superposition of the four DH domains makes it clear that SpNOX has a short crevice like FNR (Figure 8 middle left), while NOX2 has a long crevice like DUOX1 (Figure 8, middle right). NADPH-binding residues of FNR are mostly conserved in SpNOX (see Figure S11), while the two Arg residues that interact with the 2'-phosphate of NADPH in DUOX 1 are conserved in NOX2, although the side chain of one of those (R446) is not visible in the CryoEM structure of NOX2 (Figure 8, bottom left).

Although the SpNOX and NOX2 structures lack NADPH, superposition produces an excellent picture of the location of the NADPH in these proteins. Figure 8 (bottom) shows that when the proteins are superimposed, the NADPH from pea FNR fits extremely well into the NADPH binding groove of SpNOX and reproduces the short distance between nicotinamide and isoalloxazine seen in FNR. Similarly, the superimposed DUOX1 NADPH fits very well into the corresponding binding groove of NOX2. Even this crude docking with no energy minimization leaves little doubt about the overall NADPH position in SpNOX and NOX2. Our functional characterization of mutants in position 397 strongly support the picture presented here. Of course, new structures for NOX2 and SpNOX in complex with NAD(P)H would be a welcome addition to confirm these observations.

Taking into account the similar interdomain organization and comparable electron pathways (Figure 7D) of SpNOX and NOX2, the different topology at the interface of their respective FBD and NBD subdomains would explain the switch from inactive conformation of NOX2 to constitutively active conformation of SpNOX. This idea is consistent with the idea that 'tensing' of the high Ca^{2+}-state DUOX1 structure may be necessary for electron transfer (Wu et al., 2021). The mechanism by which the inactive state of NOX2 changes to active, and how this situation fits with the activated state of the DUOX1 structure, remains to be elucidated.

Conclusion

The important roles of NOX in human, animal, and plant health and disease explain the field's focus on eukaryotic NOX. However, prokaryotic NOX's advantages in production and handling offer opportunities to decipher structural and mechanistic details relevant to the whole family (eg. (Magnani et al., 2017; Vermot et al., 2020)). We present here extensive enzymological characterization of SpNOX activity and also new structural information on both full-length enzyme and the isolated DH. First, the particular effort to optimize conditions for $SpNOX_{DH}$ allowed the investigation of the first step of NOX electron transport, hydride transfer from NAD(P)H to FAD (Figure 2 and Table I). We showed that NADPH and NADH are equivalent electron donors for SpNOX, constituting a significant difference from eukaryotic NOX. Comparison with the full length SpNOX demonstrates that NAD(P)H binding is largely independent from the TM domain, at least in our conditions in which SpNOX is solubilized in LMNG detergent. Comparison of k_{cat} obtained from both constructs suggests that the hydride

transfer from NAD(P)H to FAD is the rate limiting step in the overall NOX reaction (Table 3). Structural and functional characterization of WT and mutant F397W and F397S highlight the critical role of the aromatic at 397: Phe provides flavin isoalloxazine ring interaction and stabilization and also properties that allow entrance of the nicotinamide ring to the active site, similar to the mechanism long accepted in FNR family domains. Parallel analysis of flavin analogs using both DH and full length SpNOX shows that the DH domain alone has no selectivity for flavin type. SpNOX, on the other hand, displays a marked selectivity for FAD with affinity at the nM level, indicating binding contribution from the TM domain. This is supported by interactions visible in the entire SpNOX structure, and a related mutational analysis, highlighting the role of the D-Loop region in the interaction with the adenine ring of the flavin (Figure 5B).

Structural comparison of NOX structures indicates that accessibility of the distal heme of SpNOX and CsNOX may reflect plasticity in the final electron acceptor, while the inaccessibility of other NOXes may enforce O_2 as a substrate (Figure 6). On the other hand, this comparison also generates a new conundrum, how to explain hydride transfer in activated states of NOX. SpNOX shares similar domain organization with NOX2, and the orientations and distances between cofactors in the electron transfer pathway are similar (Fig S12 for overlay of both structures). Surprisingly, DUOX1 domain organization and cofactor distances are quite different, raising the possibility that domain organization is isoform specific. Considering the constitutive activity of SpNOX, we believe that the electron pathway as organized in NOX2 is poised for electron transfer, and that the difference in position of the NADPH binding groove could explain the difference between active and inactive states. The "high Ca^{2+}" activated state of DUOX1 also displays the distant binding groove suggesting that an additional step might be required to reach an efficient active state. Rather than a global repositioning of the DH, activation of NOX2 may require an internal rearrangement within the DH that brings the NADPH groove closer to the FAD site; a similar rearrangement may be necessary for electron transfer in activated DUOX1. Conformational change within the DH may agree with a previous study concluding that p67[phox], which activates NOX2, regulates hydride transfer from NADPH to the FAD rather than NADPH binding. (Nisimoto et al., 1999). In this view, the assembly of p67[phox] with the DH could promote the required rearrangement within the DH domain. However, the resting state of NOX2 has been reported to resist initial binding of NADPH (Liu et al., 2022). This could be a second level of control for NOX2.

In conclusion, SpNOX seems to be a good model of an activated form of NOX2. The molecular tools and structures provided in this work will allow further investigations to understand additional molecular features at work in those fascinating NOX enzymes.

Materials & Methods

Structure-based sequence alignment

A large set of 549 NOX homolog sequences distributed about evenly between prokaryotes and eukaryotes, was aligned using MAFFT (Katoh et al., 2019). The sequences were split into transmembrane (TM) and dehydrogenase (DH) domains identified by homology and including 30 overlapping residues between the domains. Each domain was realigned separately with PromalS3D (Pei et al., 2008). The TM alignment was constrained to the crystal structure of

CsNOX TM domain (5O0T) while the DH alignment was constrained to the crystal structure of CsNOX DH domain (5O0X) and DH domain homologs 1GJR, 1FNB, 1GVH, 2EIX, 3A1F; all other parameters were at default values. The resulting multiple alignments were inspected in Jalview (Waterhouse et al., 2009); those with missing secondary structural elements or with unusually large insertions were removed and remaining sequences were pruned to 91 % (TM domain) or 95 % (DH domain) identity. The final sequence sets (172 sequences of TM and 207 sequences of DH domain) were realigned in Promals3D with default parameters and the same structural constraints specified above. Logos of subsequences of the full alignment were created in WebLogo (Crooks et al., 2004).

Cloning of SpNOX, SpNOX_DH WT, F397S and F397W.

The synthetic SpNOX gene optimized for expression in *E. coli* including a polyhistidine tag and thrombin cleavage site at its N-terminus was amplified *via* PCR and subcloned into pET-30b (Novagen) between the NdeI/ BamHI sites (Vermot et al., 2020).

The expression vector encoding His-tagged SpNOX_DH - aa 181-417 of full-length SpNOX - was obtained from SpNOX by site-directed mutagenesis according to the manufacturer's protocol (QuikChange® Lightning Site-Directed Mutagenesis Kit, Agilent), using forward primer 5'GTCTGGTCCCGCGTGGCAGTAAAATTAGCTTTCCGTATCTGGG3' and reverse primer 5'CCCAGATACGGAAAGCTAATTTTACTGCCACGCGGGACCAGAC3'. F397S or F397W mutants of SpNOX and SpNOX_DH were created using forward primer 5'ACGGAACTGATCTACGAAGGCTCTAAATTCAAATGAGAATTCGAGC3' and reverse primer 5'GCTCGAATTCTCATTTGAATTTAGAGCCTTCGTAGATCAGTTCCGT3' for F397S or forward primer 5'CGGAACTGATCTAC-GAAGGCTGGAAATTCAAATGAGAATTCGAG3' and reverse primer 5'CTCGAATTCTCATTTGAATTTCCAGCCTTCGTAGATCAGTTCCG3' for F397W. We retain the nomenclature 'F397x' for these mutations in both SpNOX and SpNOX_DH constructs, although in SpNOX_DH construct that residue is actually in position 217.

Production of SpNOX, SpNOX_DH WT, F397S and F397W.

The expression of SpNOX WT, F397S and F397W was performed as described in (Vermot et al., 2020). SpNOX_DH WT, F397S and F397W proteins were over-expressed in *E. coli* BL21(DE3) using TB medium supplemented with 50 μg.ml^{-1} kanamycine. At culture OD_{600} = 1, expression was induced by the addition of 0.5 mM IPTG and the culture was grown overnight at 18 °C. Cells were harvested by centrifugation (5000g, 20 min, 4°C) and stored at -20°C. The purification procedure described below is the final optimized version for SpNOX_DH stabilization. The pellet was resuspended in lysis buffer (50 mM TRIS pH7, 1M NaCl, 10 mM Imidazole and 5% glycerol) supplemented with deoxyribonuclease and a mix of protease inhibitors (chymostatin, leupeptin, antipain, pepstatin at 1μg.ml^{-1}, aprotinin at 5 μg/ml), disrupted by sonication, and cleared by centrifugation at 4°C (18500 rpm, JA-25.50 Beckman, 20 min,). The clarified extract was purified on a 4 ml His60 Ni-IDA Superflow resin (Ozyme)

previously equilibrated in high salt buffer (HS buffer; 50 mM TRIS pH7, 1M NaCl, 10 mM Imidazole and 5% glycerol). After an extensive wash (20 column volumes, HS buffer), the protein was eluted with 50 mM TRIS pH7, 1M NaCl, 300 mM Imidazole and 5% glycerol and the buffer was exchanged against a stabilization buffer (50 mM Bis- TRIS propane pH6.5, 0.3M NaCl, 5% glycerol) *via* gravity using a desalting column (PD10 ref. 17-0851-01, GE Healthcare). The selected fractions were pooled and loaded onto a Superdex75 size exclusion chromatography (GE Healthcare) equilibrated in the same buffer. The protein was eluted as a monomer and concentrated by ultrafiltration (10 kDa centrifugal filter unit, Amicon). The protein purity was checked on SDS-PAGE and the concentration was determined by UV-visible spectroscopy using respectively ε_{280nm} = 24870 $M^{-1}.cm^{-1}$ for SpNOX$_{DH}$ WT and SpNOX$_{DH}$ F397S or ε_{280nm} = 30370 $M^{-1}.cm^{-1}$ for SpNOX$_{DH}$ F397W respectively.

Thermostability assays for SpNOX$_{DH}$ stability improvement.

Thermal stability assays were performed using a Prometheus NT.48 nanoDSF instrument (NanoTemper Inc., Germany). 50 µl of sample per condition were prepared to perform experiments in triplicates using standard-treated glass capillaries. The excitation at 280 nm was optimized to yield emission intensities of intrinsic fluorescence at 330 and 350 nm, in the absence of heat gradient, in the range of 5000 to 16,000 A.U. The temperature gradient was set to an increase of 1 °C/min in a range from 20 °C to 95 °C. Protein unfolding was measured by detecting the temperature-dependent change in the ratio of fluorescence emission at 350 nm vs 330 nm. Samples used for testing pH impact on stability were done by diluting 10 times concentrated SpNOX$_{DH}$ solution (2 mg.ml^{-1}) to a 0.2mg.ml^{-1} protein solution in the following 0.1 M buffers depending on the tested pH: citric acid buffer for pH 4, 5, 5.5 and 6; sodium acetate for pH 4.5; bis TRIS-propane for pH 6.5 and 8, Pipes for pH 7; TRIS for pH 7 and 7.5; Bicine buffer for pH 9 and CAPS buffer for pH 9.5.

Activity Assay on Purified SpNOX and SpNOX$_{DH}$ Domain.

The activity of the purified DH domain was measured under aerobic conditions following simultaneously the oxidation of NAD(P)H (decrease of absorbance at 340 nm, ε = 6.22 $M^{-1}.cm^{-1}$) and the reduction of cytochrome c (increase of absorbance at 550 nm, ε = 21.1 $M^{-1}.cm^{-1}$), as a function of time, using a Cary 50 UV-visible spectrophotometer (Varian).
A typical assay was as follows: 50 mM bis TRIS-propane pH 6.5, 0.003% LMNG supplemented with 200 µM NADPH, 300 mM NaCl, and 5% glycerol, was added to a 2 mm quartz cuvette. Then 0.5 µg.ml^{-1} SpNOX$_{DH}$ (final) and 10 µM FAD (final) were added, mixing after each addition. For Cyt c reductase assay, 100 µM cytochrome c was added prior to the addition of the protein. Final assay volume was 500 µl. When full length SpNOX was used alone, not in parallel study with SpNOX$_{DH}$, we used a buffer optimized for activity of the full length enzyme: Tris pH 7, 300 mM NaCl, and 0.003% LMNG.
The inhibition of superoxide production was verified by addition of DPI at a final concentration of 50 µM or Superoxide dismutase (10U) to the reaction mixture after a sufficient time to allow the slope to be characterized in the presence or absence of the inhibitor.

Data analysis of thermostability and activity assays

The results from thermostability and activity assays were analyzed simultaneously using the JMP® statistical software (Version *13.0*. SAS Institute Inc., Cary, NC, 1989-2019) to plot 2D contour plots of Tm and activity as a function of salt concentrations and glycerol. The median of triplicates was used to compute the graphs, and linear interpolations were done between measurement points to fill the graphs. A 2D linear interpolation was performed on thermostability and activity results as a function of pH and salt concentration using Python® to produce a 3D figure.

Michaelis-Menten kinetic analysis for SpNOX and SpNOX$_{DH}$.

All chemicals and reagents were purchased from Sigma Aldrich. Various NAD(P)H concentrations were obtained by serial dilutions from the saturating NAD(P)H solution in triplicate in a 96-well (flat bottom) Greiner microplate (equilibrated at 25°C). The reaction was initiated (at 25°C) by injection of a mixture of the stabilization buffer (50 mM bis TRIS-propane pH6.5, 300 mM NaCl, 10 µM FAD, 5% glycerol and 0,025 mM LMNG) containing SpNox$_{DH}$ (250 ng) and cytochrome *c* (100 µM, final concentration) or (50 mM Tris pH 7, 300 mM NaCl, 10 µM FAD and 0,025 mM LMNG) containing SpNOX (250 ng) and cytochrome *c* (100 µM, final concentration) and the absorbance at 550 nm was recorded within 3 minutes using a monochromator multimode microplate reader (Clariostar, BMG Labtech) after orbital shaking. Michaelis-Menten saturation curves of SpNOX, initial apparent velocities as a function of NAD(P)H concentration, K_m (µM), and k_{cat} (s^{-1}) were determined with Prism® software. Kinetic parameters for all flavin analogs were determined in the presence of 200 µM of NADH using the same procedure; similarly, kinetic parameters of SpNOX$_{DH}$ were determined for NAD(P)H based on NAD(P)H oxidation by monitoring the absorbance at 340 nm within 3 minutes.

Crystallization of SpNOX DH domains

Crystallization experiments were carried out at the High Throughput Crystallization Laboratory (HTX Lab) at EMBL Grenoble using automated protocols (Cornaciu et al., 2021; Dimasi et al., 2007; Dupeux et al., 2011; Mariaule et al., 2014). SpNOX$_{DH}$ and SpNOX$_{DH}$ F397W were concentrated to 15 mg.ml^{-1}. The crystallization experiments were carried out using the sitting-drop vapor-diffusion method with a crystallization robot (Mosquito, SPTLabtech). 0.1 µL of protein solution and 0.1 µL of reservoir were mixed to equilibrate against 45 µL reservoir solution at 20˚C in 96-well CrystalDirect plates (MiTeGen) and automatically imaged in RockImager robot (Formulatrix). Initially, commercially available screening kits from NeXtal (Classic-suite, PEGs, JCSG) and Molecular Dimension (MemTrans, MemMeso, MemGold, PACT, Morpheus) were used to identify initial crystallization conditions that were optimized in subsequent experiments. Optimal crystals of SpNOX$_{DH}$ grew in crystallization drops with precipitant conditions of 35% PEG-MME500, 0.1M Sodium Citrate pH5. Optimal crystals of SpNOX$_{DH}$F397W grew in crystallization drops with precipitant condition composed of 17% to 20% w/v PEG3350, 0.1M Bis-TRIS propane pH 6.5 and 0.2 M Sodium bromide. Crystals of SpNOX$_{DH}$ WT and SpNOX$_{DH}$ F397W typically appeared after 24h and reached full size in 7 days with dimensions of respectively 100×20×10 µm^3 and 80×30×20 µm^3. Automated high-throughput crystal cryo-cooling and harvesting were performed with CrystalDirect^{TM} Technology (Cipriani et al., 2012; Márquez and Cipriani, 2014; Zander et al., 2016).

Data collection and structural solution of SpNOX DH F397W

A bromide-SAD dataset was collected at the macromolecular crystallography beamline ID30B of the European Synchrotron radiation facility, equipped with Pilatus3 6M detector (McCarthy et al., 2018). The wavelength was set above the Br K-edge at 0.9184 Å in order to maximize the anomalous signal, and a highly redundant data set at a resolution of 1.94 Å, including 7200 images with 0.05 degrees oscillation, was collected.

Data processing was performed by the GrenADES parallelproc pipeline (Monaco et al., 2013) that is based on XDS (Kabsch, 2010) and scaled using aimless (Evans and Murshudov, 2013) from the CCP4-suite (Winn et al., 2011). Attempts to solve the structure with the molecular replacement method using homologous models failed. Therefore, we experimentally solved the structure by the SAD method, using the bromide anomalous dispersion signal from crystals grown in presence of 0.2M sodium bromide. Experimental phasing and initial model building was performed using CRANK2 (Skubák and Pannu, 2013) The final model was obtained by successive rounds of manual and automated refinement using the programs REFMAC (Murshudov et al., 1997), and BUSTER (Bricogne, n.d.). Crystallographic data statistics are presented in Table S1. The PDB deposition code for the SpNOX$_{DH}$ F397W is 8qq1.

Data collection and structural solution of SpNOX DH WT

X-ray diffraction data were collected at ID30A-1/MASSIF-1 beamline at ESRF Grenoble (Bowler et al., 2015). A highly redundant data set at a resolution of 2.4 Å, including 3600 images with 0.1 degrees oscillation, was collected. The structure of SpNOX DH WT was solved at a 2.4 Å resolution by molecular replacement using the coordinates of the chain A of SpNOX DH F397W variant structure (PDB 8qq1) as a search template in Molrep searching for 3 mol/a.u. The initial solution was evaluated using COOT (Emsley et al., 2010) revealing no major clashes and good global packing with contacts between chains and classical size solvent channels. The final model was obtained by successive rounds of refinement using the programs REFMAC (Murshudov et al., 1997) and manual construction using COOT (Emsley et al., 2010). Crystallographic data statistics are presented in Table S1. The PDB deposition code for the SpNOX$_{DH}$ F397W is 8qq5.

Crystallization, Data Collection, and Processing of full length SpNOX

SpNOX F397W at 4.04 mg.ml^{-1} in 50 mM TRIS pH 7, 300 mM NaCl, 0.025 mM LMNG, 0.01 mM FAD was crystallized in 30.5% PEG 300, 0.15 M Li$_2$SO$_4$, 0.15 M NaCl and 0.1 M MES pH 6, using the vapor diffusion method in a 6 µL protein sample plus 6 µL of well hanging drop at 20°C. Hexagonal brick-red crystals were obtained, mounted on mesh litho-loops and flash frozen in native conditions in liquid nitrogen. X-ray diffraction data were collected at ID30A-1/MASSIF-1 beamline at ESRF Grenoble (Bowler et al., 2015). A dataset of 1800 images was collected with an oscillation range of 0.2° per image and an exposure time of 1.699 s and a crystal to detector distance refined to 500.9 mm. As observed for many membrane protein crystals (Martin et al., 2023; Robert et al., 2017), the data collected were very anisotropic going to 3.2 Å resolution in the strongest diffracting direction. After data processing the dataset was finally cut at 3.6 Å to ensure sufficient completeness in the highest resolution shell. Inclusion of further data only resulted in more noise without obvious gain in 2mFo-DFc

electron density. Data were initially processed using the program XDS as spherical to the highest resolution possible (3.2 Å) even though spherical statistics were not usable. Staraniso analysis for diffraction anisotropy (staraniso@globalphasing.org) revealed that completeness was 85.4% in the highest resolution shell, therefore revealing that all the data collectable for this crystal had been collected. Data was cut at the diffraction limits suggested by the Staraniso server using only the first 1000 images.

Phases were solved by molecular replacement using PHASER (McCoy et al., 2007), on amplitudes, with data corrected for anisotropy using Staraniso. The search models were the high-resolution DH domain (PDB 8qq5) and a model of the transmembrane domain obtained using PROMALS3D (Pei et al., 2008) and I-TASSER (Yang and Zhang, 2015) on the basis of multiple sequence alignment of NOX homologs and validated by structural alignment on CsNOX (Vermot et al., 2020). The molecular replacement solution was achieved with placing the DH domain first, followed by the TM domain, yielding LLG=91.8 and TFZ=10.9. The phasing resulted in a model with one molecule of SpNOX in the asymmetric unit and the space group $P6_422$. Crystal packing generation showed a sensible solution (classic type II packing of membrane protein and no clash), showing good density for the DH domain and clearly identifiable transmembrane helices, representing a typical initial anisotropic low-resolution map for a membrane protein solved by molecular replacement.

The resulting model was not biologically possible with an obvious error in the positioning of the transmembrane domain, which was inverted by 180° around the longitudinal axis. A new transmembrane domain model was generated on the basis of a sequence alignment between CsNOX and SpNOX and the use of CHAINSAW (CCP4). A new molecular replacement with PHASER (McCoy et al., 2007) without modification of the DH domain position resulted in a new orientation of transmembrane domain, which was biologically compatible. Fo-Fc map of the model harbored two positive peaks at 6.1 σ and 3.6 σ corresponding to the position of proximal and distal hemes, respectively, sandwiched between two key histidine residues, allowing the sequence assignment of the TM domain. Iterative manual building in COOT (Emsley et al., 2010) and LORESTR pipeline (Kovalevskiy et al., 2016), using TLS (5-183 for TM domain and 184-399 for DH domain) allowed large repositioning of some α-helices of transmembrane domain. A final comparison of a SpNox model generated with AlphaFold (Jumper et al., 2021) drove the repositioning of some loops and some sequence shifts in transmembrane helices not associated with hemes. Final refinement was made with the help of ISOLDE (Croll, 2018) and LORESTR, ramachandran and rotamers outliers were corrected, yielding a final model with R = 26.2% and Rfree = 32.0% that was deposited in the Protein Data Bank under the PDB accession code 8qq7. Crystallographic data statistics are presented in Table S1. The PDB deposition code for the SpNOX$_{DH}$ F397W is 8qq7.

Acknowledgment and funding

Funding for the automated crystallography pipelines at EMBL Grenoble was provided by the grants and iNEXT Discovery (*grant agreement* ID 871037) and Fragment Screen (*grant agreement ID: 101094131)* funded by the European Commission and through Instruct-ERIC. We want to thank the EMBL-ESRF Joint Structural Biology Group (JSBG) and in particular Andrew McCarthy for access and support in the use of MX beam lines at the ESRF. This work used the platforms of the Grenoble Instruct center (UMS 3518 CNRS-CEA-UJF-EMBL) with

support from FRISBI (ANR-10- INSB-0502) and GRAL (ANR-10-LABX-49-01) within the Grenoble Partnership for Structural Biology. This work was supported by the French Agence Nationale de la Recherche Bandit Project (ANR17-CE11-0013) to F.F., A.V., I.P.-H., and M.T. A.V. was supported through the Emergence program from the Univ. Grenoble Alpes; S.M.E.S. was supported by an invited professorship from Univ. Grenoble-Alpes and through Emergence partner Kennesaw State University.

Reference

Banfi B, Malgrange B, Knisz J, Steger K, Dubois-Dauphin M, Krause KH. 2004. NOX3, a superoxide-generating NADPH oxidase of the inner ear. *The Journal of biological chemistry* 279:46065–46072.

Banfi B, Molnar G, Maturana A, Steger K, Hegedus B, Demaurex N, Krause KH. 2001. A Ca(2+)-activated NADPH oxidase in testis, spleen, and lymph nodes. *The Journal of biological chemistry* 276:37594–37601.

Bowler MW, Nurizzo D, Barrett R, Beteva A, Bodin M, Caserotto H, Delagenière S, Dobias F, Flot D, Giraud T, Guichard N, Guijarro M, Lentini M, Leonard GA, McSweeney S, Oskarsson M, Schmidt W, Snigirev A, Von Stetten D, Surr J, Svensson O, Theveneau P, Mueller-Dieckmann C. 2015. MASSIF-1: a beamline dedicated to the fully automatic characterization and data collection from crystals of biological macromolecules. *J Synchrotron Rad* 22:1540–1547. doi:10.1107/S1600577515016604

Bricogne. n.d. BUSTER version X.Y.Z. Cambridge, United Kingdom: Global Phasing Ltd.

Caux C, Guigliarelli B, Vivès C, Biaso F, Horeau M, Hassoune H, Petit-Hartlein I, Juillan-Binard C, Torelli S, Fieschi F, Nivière V. 2021. Membrane-Bound Flavocytochrome MsrQ Is a Substrate of the Flavin Reductase Fre in *Escherichia coli*. *ACS Chem Biol* acschembio.1c00613. doi:10.1021/acschembio.1c00613

Ceccon M, Millana Fananas E, Massari M, Mattevi A, Magnani F. 2017. Engineering stability in NADPH oxidases: A common strategy for enzyme production. *Molecular Membrane Biology* 34:67–76. doi:10.1080/09687688.2018.1535141

Cipriani F, Röwer M, Landret C, Zander U, Felisaz F, Márquez JA. 2012. CrystalDirect: a new method for automated crystal harvesting based on laser-induced photoablation of thin films. *Acta Cryst D* 68:1393–1399. doi:10.1107/S0907444912031459

Cornaciu I, Bourgeas R, Hoffmann G, Dupeux F, Humm A-S, Mariaule V, Pica A, Clavel D, Seroul G, Murphy P, Márquez JA. 2021. The Automated Crystallography Pipelines at the EMBL HTX Facility in Grenoble. *JoVE (Journal of Visualized Experiments)* e62491. doi:10.3791/62491

Croll TI. 2018. *ISOLDE* : a physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallogr D Struct Biol* 74:519–530. doi:10.1107/S2059798318002425

Crooks GE, Hon G, Chandonia J-M, Brenner SE. 2004. WebLogo: A Sequence Logo Generator. *Genome Res* 14:1188–1190. doi:10.1101/gr.849004

Cross AR, Heyworth PG, Rae J, Curnutte JT. 1995. A Variant X-linked Chronic Granulomatous Disease Patient (X91+) with Partially Functional Cytochrome b. *Journal of Biological Chemistry* 270:8194–8200. doi:10.1074/jbc.270.14.8194

De Deken X, Wang D, Many M-C, Costagliola S, Libert F, Vassart G, Dumont JE, Miot F. 2000. Cloning of Two Human Thyroid cDNAs Encoding New Members of the NADPH Oxidase Family. *Journal of Biological Chemistry* 275:23227–23233. doi:10.1074/jbc.M000916200

Deng Z, Aliverti A, Zanetti G, Arakaki AK, Ottado J, Orellano EG, Calcaterra NB, Ceccarelli EA, Carrillo N, Karplus PA. 1999. A productive NADP+ binding mode of ferredoxin-NADP + reductase revealed by protein engineering and crystallographic studies. *Nature structural biology* 6:847–853. doi:10.1038/12307

Dimasi N, Flot D, Dupeux F, Márquez JA. 2007. Expression, crystallization and X-ray data collection from microcrystals of the extracellular domain of the human inhibitory receptor expressed on myeloid cells IREM-1. *Acta Cryst F* 63:204–208. doi:10.1107/S1744309107004903

Dupeux F, Röwer M, Seroul G, Blot D, Márquez JA. 2011. A thermal stability assay can help to estimate the crystallization likelihood of biological samples. *Acta Crystallographica Section D* 67:915–919. doi:10.1107/S0907444911036225

Dupuy C, Ohayon R, Valent A, Noël-Hudson M-S, Dème D, Virion A. 1999. Purification of a Novel Flavoprotein Involved in the Thyroid NADPH Oxidase. *Journal of Biological Chemistry* 274:37265–37269. doi:10.1074/jbc.274.52.37265

Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of ıt Coot. *Acta Crystallographica Section D* 66:486–501. doi:10.1107/S0907444910007493

Evans PR, Murshudov GN. 2013. How good are my data and what is the resolution? *Acta Crystallogr D Biol Crystallogr* 69:1204–1214. doi:10.1107/S0907444913000061

Geiszt M, Kopp JB, Varnai P, Leto TL. 2000. Identification of renox, an NAD(P)H oxidase in kidney. *Proc Natl Acad Sci U S A* 97:8010–4.

Gekko K, Timasheff SN. 1981. Mechanism of protein stabilization by glycerol: preferential hydration in glycerol-water mixtures. *Biochemistry* 20:4667–4676. doi:10.1021/bi00519a023

Gennaris A, Ezraty B, Henry C, Agrebi R, Vergnes A, Oheix E, Bos J, Leverrier P, Espinosa L, Szewczyk J, Vertommen D, Iranzo O, Collet J-F, Barras F. 2015. Repairing oxidized proteins in the bacterial envelope using respiratory chain electrons. *Nature* 528:409–412. doi:10.1038/nature15764

Hajjar C, Cherrier MV, Dias Mirandela G, Petit-Hartlein I, Stasia M-J, Fontecilla-Camps JC, Fieschi F, Dupuy J. 2017. The NOX Family of Proteins Is Also Present in Bacteria. *mBio* 8. doi:10.1128/mBio.01487-17

Han C-H, Nisimoto Y, Lee S-H, Kim ET, Lambeth JD. 2001. Characterization of the Flavoprotein Domain of gp91phox Which Has NADPH Diaphorase Activity. *The Journal of Biochemistry* 129:513–520. doi:10.1093/oxfordjournals.jbchem.a002885

Hermoso JA, Mayoral T, Faro M, Gómez-Moreno C, Sanz-Aparicio J, Medina M. 2002. Mechanism of coenzyme recognition and binding revealed by crystal structure analysis of ferredoxin-NADP+ reductase complexed with NADP+. *Journal of molecular biology* 319:1133–1142. doi:10.1016/S0022-2836(02)00388-1

Juillan-Binard C, Picciocchi A, Andrieu J-P, Dupuy J, Petit-Hartlein I, Caux-Thang C, Vivès C, Nivière V, Fieschi F. 2017. A Two-component NADPH Oxidase (NOX)-like System in Bacteria Is Involved in the Electron Transfer Chain to the Methionine Sulfoxide Reductase MsrP. *Journal of Biological Chemistry* 292:2485–2494. doi:10.1074/jbc.M116.752014

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature*. doi:10.1038/s41586-021-03819-2

Kabsch W. 2010. XDS. *Acta Crystallographica Section D* 66:125–132. doi:10.1107/S0907444909047337

Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20:1160–1166. doi:10.1093/bib/bbx108

Kikuchi H, Hikage M, Miyashita H, Fukumoto M. 2000. NADPH oxidase subunit, gp91phox homologue, preferentially expressed in human colon epithelial cells. *Gene* 254:237–243. doi:10.1016/S0378-1119(00)00258-4

Koshkin V, Pick E. 1993. Generation of superoxide by purified and relipidated cytochrome b559 in the absence of cytosolic activators. *FEBS letters* 327:57–62.

Kovalevskiy O, Nicholls RA, Murshudov GN. 2016. Automated refinement of macromolecular structures at low resolution using prior information. *Acta Crystallogr D Struct Biol* 72:1149–1161. doi:10.1107/S2059798316014534

Lalucque H, Silar P. 2003. NADPH oxidase: an enzyme for multicellularity? *Trends in Microbiology* 11:9–12. doi:10.1016/S0966-842X(02)00007-0

Liu R, Song K, Wu J-X, Geng X-P, Zheng L, Gao X, Peng H, Chen L. 2022. Structure of human phagocyte NADPH oxidase in the resting state. *eLife* 11:e83743. doi:10.7554/eLife.83743

Magnani F, Nenci S, Millana Fananas E, Ceccon M, Romero E, Fraaije MW, Mattevi A. 2017. Crystal structures and atomic model of NADPH oxidase. *Proceedings of the National Academy of Sciences* 114:6764–6769. doi:10.1073/pnas.1702293114

Mariaule V, Dupeux F, Márquez JA. 2014. Estimation of crystallization likelihood through a fluorimetric thermal stability assay. *Methods Mol Biol* 1091:189–195. doi:10.1007/978-1-62703-691-7_13

Márquez JA, Cipriani F. 2014. CrystalDirect^TM: A Novel Approach for Automated Crystal Harvesting Based on Photoablation of Thin Films In: Chen YW, editor. Structural Genomics: General Applications. Totowa, NJ: Humana Press. pp. 197–203. doi:10.1007/978-1-62703-691-7_14

Martin J, Robert X, Gouet P, Falson P, Chaptal V. 2023. Specific Xray diffraction patterns of membrane proteins caused by secondary structure collinearity. *Biochimica et Biophysica Acta (BBA) - Biomembranes* 1865:184065. doi:10.1016/j.bbamem.2022.184065

Martins de Oliveira V, Godoi Contessoto V de, Bruno da Silva F, Zago Caetano DL, Jurado de Carvalho S, Pereira Leite VB. 2018. Effects of pH and Salt Concentration on Stability of a Protein G Variant Using Coarse-Grained Models. *Biophysical Journal* 114:65–75. doi:10.1016/j.bpj.2017.11.012

Massari M, Nicoll CR, Marchese S, Mattevi A, Mascotti ML. 2022. Evolutionary and structural analyses of the NADPH oxidase family in eukaryotes reveal an initial calcium dependency. *Redox Biology* 56:102436. doi:10.1016/j.redox.2022.102436

McCarthy AA, Barrett R, Beteva A, Caserotto H, Dobias F, Felisaz F, Giraud T, Guijarro M, Janocha R, Khadrouche A, Lentini M, Leonard GA, Lopez Marrero M, Malbet-Monaco S, McSweeney S, Nurizzo D, Papp G, Rossi C, Sinoir J, Sorez C, Surr J, Svensson O, Zander U, Cipriani F, Theveneau P, Mueller-Dieckmann C. 2018. ID30B – a versatile beamline for macromolecular crystallography experiments at the ESRF. *J Synchrotron Rad* 25:1249–1260. doi:10.1107/S1600577518007166

McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. 2007. Phaser crystallographic software. *J Appl Crystallogr* 40:658–674. doi:10.1107/S0021889807021206

Monaco S, Gordon E, Bowler MW, Delagenière S, Guijarro M, Spruce D, Svensson O, McSweeney SM, McCarthy AA, Leonard G, Nanao MH. 2013. Automatic processing

of macromolecular crystallography X-ray diffraction data at the ESRF. *J Appl Cryst* 46:804–810. doi:10.1107/S0021889813006195

Murshudov GN, Vagin AA, Dodson EJ. 1997. Refinement of Macromolecular Structures by the Maximum-Likelihood Method. *Acta Crystallographica Section D* 53:240–255. doi:10.1107/S0907444996012255

Musset B, Clark RA, DeCoursey TE, Petheo GL, Geiszt M, Chen Y, Cornell JE, Eddy CA, Brzyski RG, Jamali AE. 2012. NOX5 in Human Spermatozoa: EXPRESSION, FUNCTION, AND REGULATION *. *Journal of Biological Chemistry* 287:9376–9388. doi:10.1074/jbc.M111.314955

Nisimoto Y, Diebold BA, Constentino-Gomes D, Lambeth JD. 2014. Nox4: a hydrogen peroxide-generating oxygen sensor. *Biochemistry* 53:5111–5120. doi:10.1021/bi500331y

Nisimoto Y, Jackson HM, Ogawa H, Kawahara T, Lambeth JD. 2010. Constitutive NADPH-dependent electron transferase activity of the Nox4 dehydrogenase domain. *Biochemistry* 49:2433–2442. doi:10.1021/bi9022285

Nisimoto Y, Motalebi S, Han CH, Lambeth JD. 1999. The p67(phox) activation domain regulates electron flow from NADPH to flavin in flavocytochrome b(558). *The Journal of biological chemistry* 274:22999–23005.

Nisimoto Y, Ogawa H, Miyano K, Tamura M. 2004. Activation of the flavoprotein domain of gp91phox upon interaction with N-terminal p67phox (1-210) and the Rac complex. *Biochemistry* 43:9567–9575. doi:10.1021/bi0400249

Noreng S, Ota N, Sun Y, Ho H, Johnson M, Arthur CP, Schneider K, Lehoux I, Davies CW, Mortara K, Wong K, Seshasayee D, Masureel M, Payandeh J, Yi T, Koerber JT. 2022. Structure of the core human NADPH oxidase NOX2. *Nat Commun* 13:6079. doi:10.1038/s41467-022-33711-0

O'Neill S, Mathis M, Kovačič L, Zhang S, Reinhardt J, Scholz D, Schopfer U, Bouhelal R, Knaus UG. 2018. Quantitative interaction analysis permits molecular insights into functional NOX4 NADPH oxidase heterodimer assembly. *Journal of Biological Chemistry* 293:8750–8760. doi:10.1074/jbc.RA117.001045

Page CC, Moser CC, Chen X, Dutton PL. 1999. Natural engineering principles of electron tunnelling in biological oxidation–reduction. *Nature* 402:47–52. doi:10.1038/46972

Pei J, Kim B-H, Grishin NV. 2008. PROMALS3D: a tool for multiple protein sequence and structure alignments. *Nucleic Acids Research* 36:2295–2300. doi:10.1093/nar/gkn072

Picciocchi A, Debeurme F, Beaumel S, Dagher M-C, Grunwald D, Jesaitis AJ, Stasia M-J. 2011. Role of putative second transmembrane region of Nox2 protein in the structural stability and electron transfer of the phagocytic NADPH oxidase. *Journal of Biological Chemistry* 286:28357–28369. doi:10.1074/jbc.M111.220418

Piubelli L, Aliverti A, Arakaki AK, Carrillo N, Ceccarelli EA, Karplus PA, Zanetti G. 2000. Competition between C-terminal Tyrosine and Nicotinamide Modulates Pyridine Nucleotide Affinity and Specificity in Plant Ferredoxin-NADP+ Reductase*. *Journal of Biological Chemistry* 275:10472–10476. doi:10.1074/jbc.275.14.10472

Rae J, Newburger PE, Dinauer MC, Noack D, Hopkins PJ, Kuruto R, Curnutte JT. 1998. X-Linked Chronic Granulomatous Disease: Mutations in the CYBB Gene Encoding the gp91-phox Component of Respiratory-Burst Oxidase. *The American Journal of Human Genetics* 62:1320–1331. doi:10.1086/301874

Rajagopalan S, Kurz S, Münzel T, Tarpey M, Freeman BA, Griendling KK, Harrison DG. 1996. Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone. *J Clin Invest* 97:1916–1923. doi:10.1172/JCI118623

Robert X, Kassis-Sahyoun J, Ceres N, Martin J, Sawaya MR, Read RJ, Gouet P, Falson P, Chaptal V. 2017. X-ray diffraction reveals the intrinsic difference in the physical properties of membrane and soluble proteins. *Sci Rep* 7:17013. doi:10.1038/s41598-017-17216-1

Segal AW, Jones OT. 1978. Novel cytochrome b system in phagocytic vacuoles of human granulocytes. *Nature* 276:515–517.

Segal AW, Jones OTG. 1979. The subcellular distribution and some properties of the cytochrome b component of the microbicidal oxidase system of human neutrophils. *Biochemical Journal* 182:181–188. doi:10.1042/bj1820181

Sharpe HJ, Stevens TJ, Munro S. 2010. A Comprehensive Comparison of Transmembrane Domains Reveals Organelle-Specific Properties. *Cell* 142:158–169. doi:10.1016/j.cell.2010.05.037

Shpungin S, Dotan I, Abo A, Pick E. 1989. Activation of the Superoxide Forming NADPH Oxidase in a Cell-free System by Sodium Dodecyl Sulfate. *Journal of Biological Chemistry* 264:9195–9203. doi:10.1016/S0021-9258(18)60513-7

Skubák P, Pannu NS. 2013. Automatic protein structure solution from weak X-ray data. *Nat Commun* 4:2777. doi:10.1038/ncomms3777

Suh YA, Arnold RS, Lassegue B, Shi J, Xu X, Sorescu D, Chung AB, Griendling KK, Lambeth JD. 1999. Cell transformation by the superoxide-generating oxidase Mox1. *Nature* 401:79–82.

Sun J. 2020. Structures of mouse DUOX1–DUOXA1 provide mechanistic insights into enzyme activation and regulation. *Nat Struct Mol Biol*. doi:10.1038/s41594-020-0501-x

Taylor WR, Jones DT, Segal AW. 1993. A structural model for the nucleotide binding domains of the flavocytochrome b-245 beta-chain. *Protein science : a publication of the Protein Society* 2:1675–85.

Vermot A, Petit-Härtlein I, Breyton C, Le Roy A, Thépaut M, Vivès C, Moulin M, Härtlein M, Grudinin S, Smith SME, Ebel C, Martel A, Fieschi F. 2020. Interdomain Flexibility within NADPH Oxidase Suggested by SANS Using LMNG Stealth Carrier. *Biophysical Journal* S0006349520305191. doi:10.1016/j.bpj.2020.06.025

Vermot A, Petit-Härtlein I, Smith SME, Fieschi F. 2021. NADPH Oxidases (NOX): An Overview from Discovery, Molecular Mechanisms to Physiology and Pathology. *Antioxidants* 10:890. doi:10.3390/antiox10060890

Ward J, Zhang S, Sikora A, Michalski R, Yin Y, D'Alessio A, McLoughlin RM, Jaquet V, Fieschi F, Knaus UG. 2023. VEO-IBD NOX1 variant highlights a structural region essential for NOX/DUOX catalytic activity. *Redox Biology* 102905. doi:10.1016/j.redox.2023.102905

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25:1189–1191. doi:10.1093/bioinformatics/btp033

Winn MD, Cowtan KD, Dodson EJ, Evans PR, Keegan RM, Leslie AGW, McCoy A. 2011. Overview of the CCP4 suite and current developments.

Wu J-X, Liu R, Song K, Chen L. 2021. Structures of human dual oxidase 1 complex in low-calcium and high-calcium states. *Nat Commun* 12:155. doi:10.1038/s41467-020-20466-9

Yang J, Zhang Y. 2015. I-TASSER server: new development for protein structure and function predictions. *Nucleic Acids Res* 43:W174–W181. doi:10.1093/nar/gkv342

Zander U, Hoffmann G, Cornaciu I, Marquette J-P, Papp G, Landret C, Seroul G, Sinoir J, Röwer M, Felisaz F, Rodriguez-Puente S, Mariaule V, Murphy P, Mathieu M, Cipriani F, Márquez JA. 2016. Automated harvesting and processing of protein

crystals through laser photoablation. *Acta Cryst D* 72:454–466. doi:10.1107/S2059798316000954

Zhang X, Krause K-H, Xenarios I, Soldati T, Boeckmann B. 2013. Evolution of the Ferric Reductase Domain (FRD) Superfamily: Modularity, Functional Diversification, and Signature Motifs. *PloS one* 8:e58126. doi:10.1371/journal.pone.0058126.s005