Full Title: Odorant receptors for floral- and plant-derived volatiles in the yellow fever mosquito, *Aedes*

aegypti (Diptera: Culicidae)

Short Title: *Aedes aegypti* ORs are activated by plant-derived volatiles

Authors: Heidi Pullmann-Lindsley[1], Robert Huff[1], John Boyi[1], R. Jason Pitts[1]*

[1]Department of Biology,

Baylor University

101 Bagby Avenue, Waco, TX 76706

*corresponding author

10
11

Abstract:

Adult mosquitoes require regular sugar meals, usually floral nectar, to survive and flourish in natural habitats. Both males and females locate potential sugar sources using sensory proteins called odorant receptors activated by plant volatiles that facilitate orientation toward flowers or honeydew. The Yellow Fever mosquito, *Aedes aegypti* (Linnaeus, 1762), possesses a large repertoire of odorant receptors, many of which are likely to support floral odor detection and nectar-seeking. In this study, we have employed a heterologous expression system and the two-electrode voltage clamping technique to identify environmentally relevant chemical compounds that activate specific odorant receptors. Importantly, we have uncovered ligand-receptor pairings for a suite of *Aedes aegypti* odorant receptors likely to mediate appetitive or aversive behavioral responses, thus shaping a critical aspect of the life history of a medically important mosquito. Moreover, the high degree of conservation of these receptors in other disease-transmitting species suggests common mechanisms of floral odor detection. This knowledge can be used to further investigate mosquito foraging behavior to either enhance existing, or develop novel, control strategies, especially those that incorporate mosquito bait-and-kill or attractive toxic sugar bait technologies.

Introduction:

Arthropods rely on sensory receptors to interpret and interact with their environments. Arguably, olfaction is the most important sensory modality, conveying the ability to sense volatile chemical cues over large distances [1-3]. Olfaction is responsible for many arthropod behaviors, including host-seeking [4, 5], oviposition [6, 7], mate selection [8, 9], and nectar foraging [10, 11]. Odorant receptors (ORs) comprise the most prominent family of volatile odor detectors encoded in mosquito genomes [12]. Another family of proteins called ionotropic receptors (IRs) also functions in insect olfaction, especially in organic acid detection [13, 14]. ORs are transmembrane proteins expressed by odorant receptor neurons housed in hairlike projections called sensilla [15, 16]. Sensilla sensory appendages such as antennae, maxillary palps, labella, and leg tarsi constitute the primary olfactory appendages in insects [17, 18, 19]. ORs are ligand-

gated ion channels that respond to diverse, environmentally relevant volatile odorants [20]. ORs most likely

form multimers that include co-expression of a single type of tuning OR complexed with an odorant

receptor coreceptor (Orco), which is highly conserved across most insect taxa [21-23]. Chemical volatiles

enter the sensilla through tiny pores on their surfaces and either diffuse through the sensilla lymph or are

shuttled by odorant binding proteins to the neuronal membrane, where they can interact with the OR/Orco

complex [24, 25]. Generally, single ORs respond to one compound most robustly, though they may

demonstrate the responsiveness of a handful of other ligands. Most arthropod species have between 50 and

100 different ORs, which allow for complex and robust interactions with their respective environments

through the perception of various volatile compounds. However, some species have as few as five or as

many as 300 [26, 27]. The diversity of receptor proteins underlies distinct behavioral phenomena across

arthropod species.

OR characterization is a crucial step toward arthropod control and surveillance. By delving into the

molecular intricacies of how these receptors function, we can significantly contribute to our understanding

of insect behavior. Deorphanizing single ORs has yielded valuable insights into many insect species,

directly informing behavioral control strategies. One of the most comprehensively studied relationships lies

in the realm of pheromones [28]. In the silk moth (*Bombyx mori)*, an OR, BmOR1, that responded to the

female-produced pheromone bombykol was identified [29, 30] and characterized using a heterologous

expression system [31]. Notably, disrupting BmOR1 led to the complete loss of sensitivity to bombykol,

subsequently impacting pheromone source detection in male moths[32]. While the intricate relationships

between receptors, ligands, and behavior are best characterized in the context of pheromone reception, ORs

are linked to numerous other behaviors, including nectar foraging. Understanding their function could prove

invaluable to combatting arthropod-borne threats.

The prominent disease vector *Aedes aegypti* (Linnaeus, 1762) has 133 putative odorant receptors

that vary across sex, life stage, and tissue [33-35]. *Ae. aegypti* is a prominent disease vector can transmit

several arboviruses, including Dengue, Zika, and yellow fever. This species inhabits much of Africa, Asia,

South America, Oceania, and parts of North America and Europe [36]. However, as climate change

continues to alter ecosystems and global urbanization accelerates the viable range of *Ae. aegypti* may also

expand, and its threat to human health could increase [37, 38]. Despite the significant impact of this species,

relatively few ORs have been characterized, specifically ones presumed to contribute to female host-

seeking behaviors [39, 40]. Single sensilla recordings (SSR) have revealed that *Ae. aegypti* neurons

responded to a variety of chemical compounds collected from human volunteers, including octanal and

nonanal [41]. We have also previously demonstrated that an OR highly conserved in Culicidae disease

vectors responds to carboxylic acids present on human skin [5]. However, other essential behaviors likely

mediated by ORs have yet to be studied. Remarkably, little is known about the shared behavior of nectar

foraging between males and females, which is crucial for mosquito survival and reproductive fitness [42].

While preference between nectar sources has been documented in numerous mosquito species, the

molecular and olfactory underpinnings are yet to be elucidated [43-45]. This behavior is of particular

importance for disease control as characterizing the ORs that mediate it may provide vital information about

potential nectar-seeking linked attractants that can be used in mosquito bait and kill (MBAK) or attractive

toxic sugar bait (ATSB) systems [46-48]. While MBAKs and ATSBs have demonstrated relative success

in reducing mosquito populations [49, 50], pursuing a more specific attraction to mosquito species could

reduce off-target effects [51].

In this study, we investigated ORs in *Ae. aegypti* that are conserved in several disease vector

mosquito species and are closely related to other ORs that function in floral odor sensing. These odorant

receptors are expressed mainly in male and female antennae [33-35], suggesting they may function in a

shared behavior such as nectar foraging. We initiated this study by completing manual gene annotations

and intron analysis [52]. Following this, we used a heterologous expression system and a technique called

two-electrode voltage clamping (TEVC) to identify essential ligands [53]. After synthesizing cRNA for

odorant receptors of interest, we injected unfertilized *Xenopus laevis* oocytes with the cRNA for the OR of

interest (called the tuning receptor) and cRNA for *Ae. aegypti* Orco. After allowing time for protein

expression, we used TEVC to record changes in membrane potential as a library of 147 potential ligands

was perfused past. Using this method, we identified probable ligands for ten ORs, eight of which were

novel or revealed a previously unknown ligand. Many odorant receptors responded to indole-like

compounds that affect *Ae. aegypti* behavior [54, 55]. These studies highlight the importance of

characterizing ORs to improve our understanding of the molecular basis for *Ae. aegypti* nectar seeking or

other behaviors.

While we have demonstrated that these receptors and ligands interact on a molecular basis, more

research is needed to determine how the ligands might influence a mosquito's behavior. Future studies will

include olfactometry with wild-type and gene-knockout mosquitoes to verify the role of these ORs and

research into the expression patterns of these genes in response to ligand exposure. These studies can help

to identify specific attractants or repellants for use in surveillance systems or bait traps to help control

mosquito populations and combat the spread of disease.

Materials and Methods

Phylogenetic Analysis and Gene Annotations

A conserved clade of genes appears to be activated by floral or nectar odors and utilized by mosquitoes in

foraging. These genes were identified via BLAST via tBLASTn or BLASTp using VectorBase and

confirmed using available RNAseq data (S1 Figure). Closely related ORs from *Aedes albopictus* (Skuse,

1894) and *Culex quinquefasciatus* (Say, 1823) were also identified. DNA sequences from the reference

genomes (*Ae. aegypti* LVP_AGWG, *Ae. albopictus* Foshan FPA, *Cx. quinquefasciatus* JHB 2020) were

downloaded and the genes were manually annotated using SnapGene (Dotmatics, Boston, MA, USA).

Intron analysis was conducted for the *Ae. aegypti* genes and compared with previously identified conserved

introns [52]. A Neighbor-Joining tree based on amino acid alignments was constructed using Geneious

Prime software (Biomatters Ltd., Boston, MA, USA).

Gene cloning and Sequencing

The coding regions of ten genes of interest (S2 Table) and AaegOrco were synthesized by Twist Bioscience

(San Francisco, CA, USA) and cloned into the pENTR plasmid. The Gateway LR directional cloning

system (Invitrogen Corp., Carlsbad, CA, USA) was utilized to subclone AaegOR coding regions into the

pSP64t-RFA oocyte expression plasmid. Plasmids were purified using the GeneJet Plasmid Miniprep Kit

(ThermoFisher Scientific, Waltham, MA, USA). Purified plasmids were sequenced in both directions to

confirm coding regions.

OR Gene Expression Analysis and de novo Transcript Assembly

OR expression values in Transcripts Per Million (TPM) were obtained from VectorBase (vectorbase.org)

based on published studies of *Ae*. *aegypti* chemosensory appendages [34]. *De novo* transcripts were

assembled using non-bloodfed female antennae and maxillary palp RNAseq data sets [34] using the Trinity

software package [35] on the Galaxy server (usegalaxy.org, S3 Figure).

Chemical Reagents

Chemical compounds (S4 Table) used in the deorphanization of AaegORs were purchased from Acros

Organics (Morris, NJ, USA), Alfa Aesar (Ward Hill, MA, USA), and ThermoFisher Scientific (Waltham,

MA, USA) at the highest purity available. Compounds were stored as 1M stocks in 100% DMSO. Serial

dilutions of each compound were prepared in 10% DMSO and blended by chemical class.

Two-electrode voltage clamping

pSP64t-RFA: AaegOR clones were linearized using Xbal, and cRNAs were synthesized using the

mMESSAGE mMACHINE ® SP6 kit (Life Technologies, Carlsbad, CA, USA). Xenopus laevis stage V-

VII oocytes purchased from Xenopus1 (Dexter, MI, USA) were stored at 18°C in Incubation medium

(ND96 96 with 5% dialyzed horse serum,50μg/mL gentamycin,100μg/mL streptomycin, 100μg/mL

penicillin, and 550μg/mL sodium pyruvate). 30nL of cRNA was injected into multiple oocytes using a

microinjection syringe pump (World Precision Instruments, Sarasota, FL, USA) maintained at 18C for 48-

72 hrs. Odorant blends were perfused across individual oocytes for 10-15 seconds, held at a resting

membrane potential of -80 mV using an OC-725C oocyte clamp (Warner Instruments, LLC, Hamden, CT,

USA). Induced currents were measured indirectly using the two-electrode voltage clamp technique

(TEVC). Following each stimulus, currents were allowed to return to baseline before further stimulation.

Data was captured using the Digidata 1550 B digitizer and PClamp10 software (Molecular Devices, San

Jose, CA, USA). Concentration-response curves were generated by perfusing unitary odorants over oocytes

at increasing concentrations from [10^{-3}M] to [10^{-9}M]. Data were analyzed using GraphPad Prism 8

(GraphPad Software Inc., La Jolla, CA, USA). At least two biological replicates were conducted at each

concentration and for each compound. Data for at least seven oocytes responding to each compound was

analyzed (S6 File).

Results:

1. *Aedes aegypti* odorant receptors are expressed in male and female antennae and conserved in other

disease vector mosquitoes.

Analysis of publicly available RNAseq data sets [36] demonstrated that the ORs in our study are

expressed in adult sensory appendages (Table 1). With the exception of AaegOR8, which is expressed in

the maxillary palps [33], ORs were most abundant in the antennae of both sexes (Table 1; [34]).

Phylogenetic analysis of possible floral volatile sensing odorant receptors in three major disease vectors

revealed several different groupings, which may indicate shared or similar ligands (Figure 1A). *Aedes*

aegypti ionotropic receptor 75d (AaegIR75d) served as an outgroup. Available RNAseq analysis [36]

confirmed gene structures (S1 Figure). Our analysis revealed that there are homologs to the *Ae*. *aegypti*

genes of interest in both *Ae*. *albopictus* and *Cx*. *quinquefasciatus* and manual annotations revealed close

homology (S5 Figure) and highly conserved intron structures (Figure 1B). Phylogenetic analysis supports

homologies between ORs from multiple vector species, providing targets for future functional studies, as

the receptors deorphanized in this study are likely to . Through our analysis, we identified and named six

new ORs in *Ae. albopictus*. We located two OR11 homologs, named AalbOR11a and AalbOR11b, as well

as a secondary OR13 homolog (AalbOR13b), and three previously unannotated OR31 homologs

(AalbOR31a, AalbOr31b, AalbOR31c). As with the previously annotated ORs, these clustered in the

cladogram with the homologous genes from *Ae. aegypti* and *Cx. quinquefasciatus*.

Table 1: Abundance of AaegOR transcripts in *Ae. aegypti* sensory appendages.

AaegOR	VectorBase ID	NCBI ID	Transcripts per million (+/- SE) in Ae. aegypti head appendages				
			female ant.	male ant.	female palp	male rost.	female prob.
Orco	AAEL005776	NM_001358471	2545.4 (146.6)	385.1 (9.2)	985.6 (87.3)	80.6 (4.2)	35.7 (2.1)
Or6	AAEL017548	NM_001358467	5.3 (1.0)	6.6 (0.5)	0.1 (0.1)	2.6 (0.4)	5.8 (0.7)
Or8	AAEL012254	XM_001662305.3	0 (0.0)	0 (0.0)	387.7 (37.8)	25.7 (1.3)	0.2 (0.2)
Or10	AAEL006003	NM_001358339	21.0 (3.0)	11.5 (0.6)	0 (0.0)	0.5 (0.0)	0.4 (0.1)
Or11	AAEL011583	NM_001358172	146.1 (8.9)	37.0 (1.1)	0 (0.0)	0.2 (0.1)	2.5 (0.5)
Or13	AAEL008368	NM_001358180	20.9 (1.5)	3.8 (0.6)	0 (0.0)	0.3 (0.0)	0.6 (0.1)
Or15	AAEL008448	NM_001358181	6.7 (1.2)	3.3 (0.2)	0 (0.0)	2.2 (0.2)	4.0 (0.3)
Or28	AAEL027053	NM_001358166	4.0 (0.4)	3.4 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Or31	AAEL013217	NM_001358160	19.0 (1.5)	10.4 (0.7)	0.2 (0.1)	0.5 (0.0)	1.0 (0.3)
Or55	AAEL010415	NM_001358184	4.4 (0.3)	1.4 (0.3)	0 (0.0)	0.5 (0.1)	1.3 (0.5)
Or71	AAEL017564	NM_001358112	43.1 (2.7)	14.6 (0.3)	0.9 (0.3)	2.9 (0.4)	9.4 (0.7)

Data from VectorBase.org (Matthews *et al.* 2016)

TPM scale

0 ----------- 2545

2. Odorant receptors respond to environmentally relevant compounds in a concentration dependent manner.

Ae. aegypti ORs responded to blends of chemical compounds (Figure 2). While there is some

responsiveness to multiple blends by specific ORs, all demonstrated a pronounced specificity for one blend

over the others, as indicated by highly positive kurtosis values (Table 2). The blends that most frequently

activated ORs were indole or alcohol blends. Unitary compounds from the highest activating blend for each

OR were tested individually. We determined the primary ligand each OR responded to from our chemical

library and noted other secondary ligands (Table 2, Figure 3). As with the blends, ORs exhibited

pronounced responses to a single ligand, again supported by highly positive kurtosis values. A defining

feature of OR responsiveness is that it is concentration-based; we also investigated responses to increasing

concentrations with that compound. We determined the half-maximal effective concentration (EC_{50}) for

each primary ligand (Table 2, Figure 4).

Table 2: Functional responses of *Ae. aegypti* odorant receptors

Odorant Receptor	Blend	Kurtosis	Primary Ligand (s)	Kurtosis	EC$_{50}$ (μM)	Secondary Ligand(s)
AaegOr6	Indoles 2	14.86	4-chromanone	10.85	6.84	
AaegOr8	Alcohols 2	15.71	1-octen-3-ol	8.75	2.14	
AaegOr10	Indoles 1	15.61	3-methylindole	5.87	0.16	
AaegOr11	Terpenes 3	5.68	cis-deca-hydro-1-napthol	6.85	68.40	fenchone
AaegOr13	Indoles 1	6.24	4-ethyl-phenol	9.11	0.17	
AaegOr15	Alcohols 2	3.63	1-hexanol	5.23	1.07	phenethyl propionate
AaegOr28	Alcohols 2	15.96	tetrahydrolinalool	7.70	44.13	
AaegOr31	Alcohols 1	14.58	Farnesol	10.70	14.32	
AaegOr55	Indoles 1	14.50	2,3-dimethylindole	0.22	8.65	
AaegOr71	Indoles 2	15.56	4-quinolinecarboxaldehyde	3.08	0.96	

Summarization table of 10 *Ae. aegypti* odorant receptors and the corresponding responsiveness to chemical

and unitary compound blends. Kurtosis values are indicated for both blend and unitary compounds.

EC50 values (μM) are indicated for primary ligands. Secondary ligands are presented if identified during

the functional characterizations.

Three ORs responded most robustly to the Indole 1 blend. OR10, OR13, and OR55 responded to

3-methylindole (0.16 μM), 4-ethylphenol (0.17 μM), and 2,3-dimethylindole (8.65 μM), respectively. OR6

responded to 4-chromanone (6.81 μM) from the Indole 2 blend. The other receptor that responded to this

blend was OR71, which responded to 4-quinoline carboxaldehyde (0.96 μM). From the Alcohols 1 blend,

OR31 responded to farnesol (14.32 μM), while from the Alcohols 2 blend, OR8 and OR15 responded to

1-octen-3-ol (2.14 μM) and 1-hexanol (1.07 μM), respectively. A previously identified ligand of OR15,

phenethyl propionate, demonstrated an EC$_{50}$ of 20.16 μM. OR28 responded to tetrahydrolinalool (44.13

μM) from the Alcohols 3 blend. Finally, the only receptor to respond to a compound not belonging to an

indole or alcohol blend was OR11, which responded to cis-decahydro-1-naphthal (68.40 μM) from the

Terpenes 3 blend. We found that OR11 has an EC$_{50}$ of about 77.10 μM to a previously identified ligand,

fenchone.

Discussion:

For mosquitoes, nectar foraging to obtain sugar meals is essential for survival . Individuals that can

locate and access higher-quality nectar sources exhibit extended lifespans and increased fecundity [43, 57-

58]. Therefore, exploring the molecular mechanisms driving selective preferences is imperative. While past

research has demonstrated the existence of nectar source preferences in various mosquito species

corresponding to nectar qualities [44-45, 59-63], a substantial gap in our comprehension of the sensory

components facilitating these preferences still needs to be addressed. In other insect species, the study of

ORs has proven instrumental in deciphering behavioral patterns and formulating effective control measures

[64-66].

ORs function as sensory gateways, providing critical information about the surrounding

environment and enabling mosquitoes to discern and select nectar sources based on the chemical signals

they receive [3]. In this context, our research centers on the responsiveness of a series of ORs to a library

of ecologically relevant compounds, linking OR-ligand relationships to floral odor sensing. By expanding

our knowledge of the chemoreceptive basis of a critical mosquito behavior, we expect to contribute to

enhanced surveillance and control measures by developing more effective odor-baited trap and kill systems.

In turn, these improvements may help to mitigate the impact of mosquitoes on human health.

We began this study by examining *Ae. aegypti* OR gene structures, expression patterns, and

relatedness to ORs from two other prominent disease vectors, *Cx. quinquefasciatus* and *Ae. albopictus* [67].

We selected a subset of ORs to characterize based on their homologies to ORs or that have been previously

shown to respond to floral odors [68]. While AaegORs are expressed in various chemosensory tissues,

antennae are the primary structures involved in odor sensing [1]. Their expression in the antennae of both

sexes suggests that ORs are used for shared behaviors, such as nectar feeding. Furthermore, by refining and

examining *Ae. aegypti* ORs gene structures, we identified homologs of interest in both *Ae. albopictus* and

Cx. quinquefasciatus, two medically relevant mosquito species. The existence of numerous closely related

genes leads us to propose that the essential act of nectar foraging in vector mosquitoes shares a common

underlying OR mechanism, indeed that OR-ligand pairs may have been conserved through divergence of

these species. Future studies will be required to characterize the response profiles of conserved ORs, and

to establish the requirement of these receptors in nectar foraging in *Ae. aegypti* and other species

222   Using two-electrode voltage clamping in conjunction with the *Xenopus laevis* oocyte expression

system, we assessed the responses of the AaegORs to our chemical library, which contains 147

environmentally significant compounds organized into 14 blends. Many AaegORs responded robustly to

the blends containing indole or indole-like compounds (Supp. Table 1). Indoles are relatively ubiquitous in

natural settings and are present in potential nectar sources. AaegOR71 responded to 4-quinoline

carboxaldehyde, a compound identified in honey samples and endophytic bacteria via gas chromatography

[69-70]. However, indoles are emitted naturally from many sources, including flowers, vertebrate hosts,

and oviposition sites of microbiotic origin, suggesting they influence multiple behaviors across species [71-

72]. AaegOR6 responds best to 4-chromanone, a compound that is found in shrubs of the Asteraceae

family, is structurally related to quinolines and is a key constituent of flavonoid skeletons [73-74].

AaegOR10 responds to 3-methylindole (skatole) found in animal feces and produces behavioral responses

in other Dipteran species [75-78]. 4-ethylphenol, the primary ligand of AaegOR13, is a volatile constituent

in wine and animal waste and has been linked to oviposition in *Culex pipiens* [78-79]. While there is little

evidence about natural sources of 2,3-dimethylindole, the AaegOR55 ligand is closely related structurally

to 3-methylindole. Other ORs responded to compounds from the "Alcohols 1", "Alcohols 2", or "Alcohols

3" blends. AaegOR8 responds to 1-octen-3-ol, a well-characterized fungal volatile and insect attractant

[80]. A green-leaf volatile, 1-hexanol, is another compound with well-characterized physiological effects

in many arthropods [81-84]. Our study discovered that 1-hexanol activates AaegOR15 with a low threshold

(EC_{50} ~1.07 μM). AaegOR28 responds to tetrahydrolinalool, which is structurally related to linalool,

another plant-derived ligand to mosquito ORs [85]. AaegOR31 responds to farnesol, a significant

constituent of the honeybee Nasanov pheromone [86]. Lastly, the AaegOR11 ligand, cis-decahydro-1-

napthol, is structurally related to geosmin, a known oviposition attractant [87]. One limitation of oocyte

studies is the number of compounds that can be screened. More efficacious or potent ligands may exist for

AaegOR11 and AaegOR28, which produced EC_{50} values greater than $40 \mu M$ (Table 2). We will continue to

expand our library by adding compounds structurally related to known ligands.

While four (AaegOR8, AaegOR10, AaegOR11, and AaegOR15) of the ORs tested in this study

already have identified ligands, we assessed them to uncover more potentially behaviorally relevant ligands,

as our chemical library differs from others tested. For AaegOR8 and AaegOR10, our data confirm previous

observations of high selectivity to 1-octen-3-ol and skatole, respectively [56, 88]. For AaegOR11 and

AaegOR15, our data supports and adds to previous studies. For AaegOR11, we found that cis-decahydro-

1-napthol produced a greater magnitude of response than (-)-fenchone or (+)-fenchone, and we identified

1-hexanol as a potent activator of AaegOR15 in addition to phenethyl propionate. This demonstrates that

even for deorphanized ORs, assessing their function with expanded libraries can aid in further

understanding receptor-ligand interactions.

Ae. aegypti and other vector mosquitoes remain persistent global threats to human health and

flourishing. Pursuing receptor-ligand pairings is paramount, as it opens doors to identifying potential

attractants for behavioral control or targets for genetic knockout. In our investigation of floral-odor sensing

receptors, we aim to pinpoint compounds that impart behavioral attraction for both sexes. Such compounds

can be applied in Mosquito Baited Traps (MBAK) and Attractive Toxic Sugar Baits (ATSB) systems. While

these technologies have been successfully deployed in some studies [48-52], identifying compounds that

activate *Ae. aegypti* ORs offers the potential to refine and enhance their effectiveness while minimizing the

impact on non-target species. OR deorphanization is a crucial initial step in unraveling the intricate sensory

systems of arthropods, providing valuable insights for future experiments and control strategies. This

focused endeavor not only aids in addressing the immediate threats posed by vector mosquitoes but also

contributes to our broader understanding of insect behavior and the development of sustainable control

measures.

Literature Cited:

1. Hansson BS, Stensmyr MC. Evolution of Insect Olfaction. Neuron. 2011 Dec 8;72(5):698–711.

2. Ray A. Reception of odors and repellents in mosquitoes. Current Opinion in Neurobiology. 2015 Oct 1;34:158–64.

3. Lutz EK, Lahondère C, Vinauger C, Riffell JA. Olfactory learning and chemical ecology of olfaction in disease vector mosquitoes: a life history perspective. Current Opinion in Insect Science. 2017 Apr 1;20:75–83.

4. Sun H, Liu F, Ye Z, Baker A, Zwiebel LJ. Mutagenesis of the orco odorant receptor coreceptor impairs olfactory function in the malaria vector *Anopheles coluzzii*. Insect Biochem Mol Biol. 2020 Dec;127:103497.

5. Huff RM, Pitts RJ. Carboxylic acid responses by a conserved odorant receptor in culicine vector mosquitoes. Insect Mol Biol. 2020 Dec;29(6):523–30.

6. Hughes DT, Pelletier J, Luetje CW, Leal WS. An odorant receptor from the southern house mosquito narrowly tuned to the oviposition attractant skatole. J Chem Ecol. 2010 Aug;36(8):797–800.

7. Liu H, Liu L, Cheng P, Huang X, Gong M. An odorant receptor from *Anopheles sinensis* in China is sensitive to oviposition attractants. Malar J. 2018 Oct 5;17(1):348.

8. Dahanukar A, Hallem EA, Carlson JR. Insect chemoreception. Curr Opin Neurobiol. 2005 Aug;15(4):423–30.

9. Sun YL, Dong JF, Yang HB, Li DX, Tian CH. Identification and Characterization of

Chemosensory Receptors in the Pheromone Gland-Ovipositor of *Spodoptera frugiperda*

(J. E. Smith). Insects. 2022 May;13(5):481.

10. Huff RM, Pitts RJ. An odorant receptor from *Anopheles gambiae* demonstrates

enantioselectivity to the plant volatile linalool. PLoS One. 2019;14(11):e0225637.

11. Dekel A, Yakir E, Bohbot JD. The sulcatone receptor of the strict nectar-feeding mosquito

Toxorhynchites amboinensis. Insect Biochemistry and Molecular Biology. 2019 Aug

1;111:103174.

12. Robertson HM. Molecular Evolution of the Major Arthropod Chemoreceptor Gene

Families. Annual Review of Entomology. 2019;64(1):227–42.

13. Raji JI, Melo N, Castillo JS, Gonzalez S, Saldana V, Stensmyr MC, *et al*. *Aedes aegypti*

Mosquitoes Detect Acidic Volatiles Found in Human Odor Using the IR8a Pathway. Curr

Biol. 2019 Apr 22;29(8):1253-1262.e7.

14. Stocker RF. Drosophila as a focus in olfactory research: Mapping of olfactory sensilla by

fine structure, odor specificity, odorant receptor expression, and central connectivity.

Microscopy Research and Technique. 2001;55(5):284–96.

15. Jiang X, Breer H, Pregitzer P. Sensilla-Specific Expression of Odorant Receptors in the

Desert Locust *Schistocerca gregaria*. Front Physiol. 2019;10:1052.

16. Sutcliffe JF. Sensory bases of attractancy: morphology of mosquito olfactory sensilla-- a

review. J Am Mosq Control Assoc. 1994 Jun;10(2 Pt 2):309–15.

17. Zahran N, Sawires S, Hamza A. Piercing and sucking mouth parts sensilla of irradiated

mosquito, *Culex pipiens* (Diptera: Culicidae) with gamma radiation. Sci Rep. 2022 Oct

25;12(1):17833.

18. Yamany AS, Abdel-Gaber R. Studies on the sensory sensilla on the tarsi and external genitalia of the Asian tiger mosquito, *Aedes albopictus* (Skuse). Microsc Res Tech. 2023 Feb;86(2):242–51.

19. del Mármol J, Yedlin MA, Ruta V. The structural basis of odorant recognition in insect olfactory receptors. Nature. 2021 Sep;597(7874):126–31.

20. Tiwari V, Sowdhamini R. Structure modelling of odorant receptor from *Aedes aegypti* and identification of potential repellent molecules. Computational and Structural Biotechnology Journal. 2023 Jan 1;21:2204–14.

21. Tiwari V, Sowdhamini R. Structure modelling of odorant receptor from *Aedes aegypti* and identification of potential repellent molecules. Computational and Structural Biotechnology Journal. 2023 Jan 1;21:2204–14.

22. Stengl M, Funk NW. The role of the coreceptor Orco in insect olfactory transduction. J Comp Physiol A Neuroethol Sens Neural Behav Physiol. 2013 Nov;199(11):897–909.

23. Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, *et al*. Cryo-EM structure of the insect olfactory receptor Orco. Nature. 2018 Aug;560(7719):447–52.

24. Zhou JJ. Chapter Ten - Odorant-Binding Proteins in Insects. In: Litwack G, editor. Vitamins & Hormones [Internet]. Academic Press; 2010 [cited 2023 Jul 6]. p. 241–72. (Pheromones; vol. 83). Available from: https://www.sciencedirect.com/science/article/pii/S0083672910830109

25. Li RT, Huang LQ, Dong JF, Wang CZ. A moth odorant receptor highly expressed in the ovipositor is involved in detecting host-plant volatiles. Elife. 2020 May 21;9:e53706.

26. Leal WS. Odorant Reception in Insects: Roles of Receptors, Binding Proteins, and Degrading Enzymes. Annual Review of Entomology. 2013;58(1):373–91.

27. Brand P, Robertson HM, Lin W, Pothula R, Klingeman WE, Jurat-Fuentes JL, *et al*. The

origin of the odorant receptor gene family in insects. Benton R, Tautz D, editors. eLife.

2018 Jul 31;7:e38340.

28. Fleischer J, Krieger J. Insect Pheromone Receptors - Key Elements in Sensing

Intraspecific Chemical Signals. Front Cell Neurosci. 2018;12:425.

29. Sakurai T, Nakagawa T, Mitsuno H, Mori H, Endo Y, Tanoue S, *et al.* Identification and

functional characterization of a sex pheromone receptor in the silkmoth Bombyx mori.

Proceedings of the National Academy of Sciences. 2004 Nov 23;101(47):16653–8.

30. Krieger J, Große-Wilde E, Gohl T, Breer H. Candidate pheromone receptors of the

silkmoth Bombyx mori. European Journal of Neuroscience. 2005;21(8):2167–76

31. Nakagawa T, Sakurai T, Nishioka T, Touhara K. Insect Sex-Pheromone Signals Mediated

by Specific Combinations of Olfactory Receptors. Science. 2005 Mar

11;307(5715):1638–42.

32. Sakurai T, Mitsuno H, Mikami A, Uchino K, Tabuchi M, Zhang F, *et al.* Targeted

disruption of a single sex pheromone receptor gene completely abolishes in vivo

pheromone response in the silkmoth. Sci Rep. 2015 Jun 5;5(1):11001.

33. Bohbot J, Pitts RJ, Kwon HW, Rützler M, Robertson HM, Zwiebel LJ. Molecular

characterization of the *Aedes aegypti* odorant receptor gene family. Insect Mol Biol. 2007

Oct;16(5):525–37.

34. Matthews BJ, McBride CS, DeGennaro M, Despo O, Vosshall LB. The

neurotranscriptome of the *Aedes aegypti* mosquito. BMC Genomics. 2016 Jan 6;17:32.

35. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, *et al*. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.

2011 May 15;29(7):644–52.

36. Matthews BJ, Dudchenko O, Kingan SB, Koren S, Antoshechkin I, Crawford JE, *et al.*

Improved reference genome of *Aedes aegypti* informs arbovirus vector control. Nature.

2018 Nov;563(7732):501–7.

37. Kraemer MUG, Sinka ME, Duda KA, Mylne AQN, Shearer FM, Barker CM, *et al.* The

global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. Elife. 2015

Jun 30;4:e08347.

38. Iwamura T, Guzman-Holst A, Murray KA. Accelerating invasion potential of disease

vector *Aedes aegypti* under climate change. Nat Commun. 2020 May 1;11(1):2130.

39. Khormi HM, Kumar L. Climate change and the potential global distribution of Aedes

aegypti: spatial modelling using GIS and CLIMEX. Geospat Health. 2014 May;8(2):405–

15.

40. Coutinho-Abreu IV, Riffell JA, Akbari OS. Human attractive cues and mosquito host-

seeking behavior. Trends in Parasitology. 2022 Mar 1;38(3):246–64.

41. Wang X, Chen Z, Wang Y, Liu F, Jiang S, Liu N. Heterologous Expression and Functional

Analysis of *Aedes aegypti* Odorant Receptors to Human Odors in Xenopus Oocytes. JoVE

(Journal of Visualized Experiments). 2021 Jun 8;(172):e61813.

42. Ghaninia M, Larsson M, Hansson BS, Ignell R. Natural odor ligands for olfactory receptor

neurons of the female mosquito *Aedes aegypti*: use of gas chromatography-linked single

sensillum recordings. J Exp Biol. 2008 Sep;211(Pt 18):3020–7.

43. Nyasembe VO, Tchouassi DP, Muturi MN, Pirk CWW, Sole CL, Torto B. Plant nutrient

quality impacts survival and reproductive fitness of the dengue vector *Aedes aegypti*.

Parasit Vectors. 2021 Jan 4;14(1):4.

44. Dieng H, Satho T, Binti Arzemi NA, Aliasan NE, Abang F, Wydiamala E, *et al.* Exposure

of a diurnal mosquito vector to floral mimics: Foraging responses, feeding patterns, and

significance for sugar bait technology. Acta Tropica. 2018 Sep 1;185:230–8.

45. Grimstad PR, DeFoliart GR. Nectar Sources of Wisconsin Mosquitoes1. Journal of

Medical Entomology. 1974 Jul 15;11(3):331–41.

46. Nikbakhtzadeh MR, Terbot II JW, Otienoburu PE, Foster WA. Olfactory basis of floral

preference of the malaria vector *Anopheles gambiae* (Diptera: Culicidae) among common

African plants. Journal of Vector Ecology. 2014;39(2):372–83

47. Kashiwagi GA, von Oppen S, Harburguer L, González-Audino P. The main component

of the scent of *Senecio madagascariensis* flowers is an attractant for *Aedes aegypti* (L.)

(Diptera: Culicidae) mosquitoes. Bull Entomol Res. 2022 Dec;112(6):837–46.

48. Sissoko F, Junnila A, Traore MM, Traore SF, Doumbia S, Dembele SM, *et al*. Frequent

sugar feeding behavior by *Aedes aegypti* in Bamako, Mali, makes them ideal candidates

for control with attractive toxic sugar baits (ATSB). PLoS One. 2019;14(6):e0214170.

49. Fikrig K, Johnson BJ, Fish D, Ritchie SA. Assessment of synthetic floral-based attractants

and sugar baits to capture male and female *Aedes aegypti* (Diptera: Culicidae). Parasit

Vectors. 2017 Jan 17;10(1):32.

50. Qualls WA, Müller GC, Revay EE, Allan SA, Arheart KL, Beier JC, *et al.* Evaluation of

attractive toxic sugar bait (ATSB)-Barrier for control of vector and nuisance mosquitoes

and its effect on non-target organisms in sub-tropical environments in Florida. Acta Tropica. 2014;131(1):104–10.

51. Junnila A, Revay EE, Müller GC, Kravchenko V, Qualls WA, Xue R de, *et al.* Efficacy of attractive toxic sugar baits (ATSB) against *Aedes albopictus* with garlic oil encapsulated in beta-cyclodextrin as the active ingredient. Acta Trop. 2015 Dec;152:195–200.

52. Fiorenzano JM, Koehler PG, Xue RD. Attractive Toxic Sugar Bait (ATSB) For Control of Mosquitoes and Its Impact on Non-Target Organisms: A Review. Int J Environ Res Public Health. 2017 Apr 10;14(4):398.

53. Bohbot J, Pitts RJ, Kwon HW, Rützler M, Robertson HM, Zwiebel LJ. Molecular characterization of the *Aedes aegypti* odorant receptor gene family. Insect Mol Biol. 2007 Oct;16(5):525–37.

54. Mowry KL. Using the Xenopus Oocyte Toolbox. Cold Spring Harb Protoc. 2020 Apr 1;2020(4):095844.

55. Baak-Baak CM, Rodríguez-Ramírez AD, García-Rejón JE, Ríos-Delgado S, Torres-Estrada JL. Development and laboratory evaluation of chemically-based baited ovitrap for the monitoring of *Aedes aegypti*. J Vector Ecol. 2013 Jun;38(1):175–81.

56. Ruel DM, Yakir E, Bohbot JD. Supersensitive Odorant Receptor Underscores Pleiotropic Roles of Indoles in Mosquito Ecology. Frontiers in Cellular Neuroscience [Internet]. 2019 [cited 2023 Jul 12];12. Available from: https://www.frontiersin.org/articles/10.3389/fncel.2018.00533

57. Foster WA. Mosquito Sugar Feeding and Reproductive Energetics. Annual Review of Entomology. 1995;40(1):443–74.

58. Stone CM, Taylor RM, Roitberg BD, Foster WA. Sugar Deprivation Reduces

Insemination of *Anopheles gambiae* (Diptera: Culicidae), Despite Daily Recruitment of

Adults, and Predicts Decline in Model Populations. Journal of Medical Entomology. 2009

Nov 1;46(6):1327–37.

59. Barredo E, DeGennaro M. Not Just from Blood: Mosquito Nutrient Acquisition from

Nectar Sources. Trends Parasitol. 2020 May;36(5):473–84.

60. Vargo AM, Foster WA. Responsiveness of Female *Aedes aegypti* (Diptera: Culicidae) to

Flower Extracts. Journal of Medical Entomology. 1982 Nov 30;19(6):710–8.

61. Gouagna LC, Poueme RS, Dabiré KR, Ouédraogo JB, Fontenille D, Simard F. Patterns of

sugar feeding and host plant preferences in adult males of *An. gambiae* (Diptera:

Culicidae). Journal of Vector Ecology. 2010;35(2):267–76.

62. Chen Z, Kearney CM. Nectar protein content and attractiveness to *Aedes aegypti* and

Culex pipiens in plants with nectar/insect associations. Acta Trop. 2015 Jun;146:81–8.

63. Lahondère C, Vinauger C, Okubo RP, Wolff GH, Chan JK, Akbari OS, *et al.* The

olfactory basis of orchid pollination by mosquitoes. Proceedings of the National Academy

of Sciences. 2020 Jan 7;117(1):708–16.

64. Wang Q, Xu P, Andreazza F, Liu Y, Nomura Y, Duran P, *et al.* Identification of multiple

odorant receptors essential for pyrethrum repellency in Drosophila melanogaster. PLoS

Genet. 2021 Jul;17(7):e1009677.

65. Cattaneo AM. Current Status on the Functional Characterization of Chemosensory

Receptors of Cydia pomonella (Lepidoptera: Tortricidae). Frontiers in Behavioral

Neuroscience [Internet]. 2018 [cited 2023 Oct 15];12.

66. Xia Y, Wang G, Buscariollo D, Pitts RJ, Wenger H, Zwiebel LJ. The molecular and
cellular basis of olfactory-driven behavior in *Anopheles gambiae* larvae. Proc Natl Acad
Sci U S A. 2008 Apr 29;105(17):6433–8.

67. Bohbot JD, Jones PL, Wang G, Pitts RJ, Pask GM, Zwiebel LJ. Conservation of indole
responsive odorant receptors in mosquitoes reveals an ancient olfactory trait. Chem
Senses. 2011 Jan;36(2):149–60.

68. Syed Z, Leal WS. Maxillary palps are broad spectrum odorant detectors in *Culex*
quinquefasciatus. Chem Senses. 2007 Oct;32(8):727–38.

69. Rodríguez-Cabo T, Moniruzzaman M, Rodríguez I, Ramil M, Cela R, Gan SH. Time-of-
flight accurate mass spectrometry identification of quinoline alkaloids in honey. Anal
Bioanal Chem. 2015 Aug 1;407(20):6159–70.

70. Moniruzzaman M, Rodríguez I, Ramil M, Cela R, Sulaiman SA, Gan SH. Assessment of
gas chromatography time-of-flight accurate mass spectrometry for identification of
volatile and semi-volatile compounds in honey. Talanta. 2014 Nov 1;129:505–15.

71. Naama-Amar A, Gitman S, Shoshana N, Bahar O, Naor V, Zchori-Fein E, *et al*.
Antimicrobial Activity of Metabolites Secreted by the Endophytic Bacterium *Frateuria*
defendens. Plants. 2020 Jan;9(1):72.

72. Tomberlin JK, Crippen TL, Wu G, Griffin AS, Wood TK, Kilner RM. Indole: An
evolutionarily conserved influencer of behavior across kingdoms. BioEssays.
2017;39(2):1600203.

73. Bohlmann F, Jakupovic J, Schuster A, King RM, Robinson H. Guaianolides and
homoditerpenes from *Lasiolaena morii*. Phytochem. 1982; 21(1):161–65.

74. Liu Y, Qian X, Xing J, Li N, Li J, Su Q, *et al*. Accurate Determination of 12 Lactones and

11 Volatile Phenols in Nongrape Wines through Headspace-Solid-Phase Microextraction

(HS-SPME) Combined with High-Resolution Gas Chromatography-Orbitrap Mass

Spectrometry (GC-Orbitrap-MS). J Agric Food Chem. 2022 Feb 16;70(6):1971–83.

75. Yokoyama M, Carlson J. Microbial metabolites of tryptophan in the intestinal tract with

special reference to skatole1 2. The American Journal of Clinical Nutrition. 1979 Jan

1;32(1):173–8.

76. Cook KL, Rothrock MJ, Loughrin JH, Doerner KC. Characterization of skatole-producing

microbial populations in enriched swine lagoon slurry. FEMS Microbiol Ecol. 2007

May;60(2):329–40.

77. Jeanbourquin P, Guerin PM. Sensory and behavioural responses of the stable fly

Stomoxys calcitrans to rumen volatiles. Med Vet Entomol. 2007 Sep;21(3):217–24.

78. Johnson SD, Sivechurran J, Doarsamy S, Shuttleworth A. Dung mimicry: the function of

volatile emissions and corolla patterning in fly-pollinated Wurmbea flowers. New

Phytologist. 2020;228(5):1662–73.

79. Musonye HA, Njeru EM, Hassanali A, Langata LM, Mijele D, Kaitho T, *et al*. 16S rRNA

gene profiling of bacterial communities mediating production of tsetse attractive phenols

in mammalian urine. Onderstepoort Journal of Veterinary Research. 2019 Jul 17;86(1):12.

80. Zhu F, Xu P, Barbosa RMR, Choo YM, Leal WS. RNAi-based demonstration of direct

link between specific odorant receptors and mosquito oviposition behavior. Insect

Biochemistry and Molecular Biology. 2013 Oct 1;43(10):916–23.

81. Kline DL. Olfactory attractants for mosquito surveillance and control: 1-octen-3-ol. J Am

Mosq Control Assoc. 1994 Jun;10(2 Pt 2):280–7.

82. De Jong R, Visser JH. Integration of olfactory information in the Colorado potato beetle brain. Brain Research. 1988 Apr 26;447(1):10–7.

83. Dickens JC, Prestwich GD, Ng CS, Visser JH. Selectively fluorinated analogs reveal differential olfactory reception and inactivation of green leaf volatiles in insects. J Chem Ecol. 1993 Sep 1;19(9):1981–91.

84. Williams L, Blackmer JL, Rodriguez-Saona C, Zhu S. Plant Volatiles Influence Electrophysiological and Behavioral Responses of *Lygus hesperus*. J Chem Ecol. 2010 May 1;36(5):467–78.

85. von Arx M, Schmidt-Büsser D, Guerin PM. Host plant volatiles induce oriented flight behaviour in male European grapevine moths, Lobesia botrana. J Insect Physiol. 2011 Oct;57(10):1323–31.

86. Huff RM, Pitts RJ. Functional conservation of Anopheline linalool receptors through 100 million years of evolution. Chemical Senses. 2022 Jan 1;47:bjac032.

87. Williams IH, Pickett JA, Martin AP. The Nasonov pheromone of the honeybeeApis mellifera L. (Hymenoptera, Apidae). Part II. Bioassay of the components using foragers. J Chem Ecol. 1981 Mar;7(2):225–37.

88. Melo N, Wolff GH, Costa-da-Silva AL, Arribas R, Triana MF, Gugger M, *et al*. Geosmin Attracts *Aedes aegypti* Mosquitoes to Oviposition Sites. Current Biology. 2020 Jan 6;30(1):127–134.e5.

89. Bohbot JD, Dickens JC. Characterization of an Enantioselective Odorant Receptor in the Yellow Fever Mosquito *Aedes aegypti*. PLOS ONE. 2009 Sep 15;4(9):e7032.

Figure Legends:

Figure 1. Phylogenetic analysis of possible floral volatile sensing ORs in major disease

vectors

(A) Neighbor-joining tree showing relationships of floral volatile sensing OR conceptual peptides

for three major disease vector species of mosquitoes *Ae. aegypti* (black), *Ae. albopictus* (Blue),

and *Cx. quinquefasciatus* (green). Node values signify bootstrap values from 1000

pseudoreplicates. (B) Intron analysis indicates shared exon and intron sizes and structures. Letter

notation indicates intron, and colors vary based on likely designation.

Figure 2. Heat map depicting AaegOR responses to blends of environmentally relevant

odors. Blends of chemical compounds used for receptor screening are listed on the lefthand side

of the figure while receptor identities are listed below. Color gradation is shown on the right and

represents normalized mean oocyte responses from highest (100, yellow) to lowest (0, blue).

Figure 3. *Ae. aegypti* odorant receptor responses to unitary compounds.

Normalized mean current responses of $AaegOR_X$ + AaegOrco expressing oocytes (n=7–10 oocytes

per blend) to the individual components of various blends (A) 4-chromanone (B) 1-octen-3-ol, (C)

3-methylindole (D) cis-decahydro-1-napthol, (E) 4-ethyl-phenol, (F) 1-hexanol, (G)

tetrahydrolinalool, (H) farnesol, (I) 2,3-dimethylindole, and (J) 4-quinolinecarboxyaldehyde. The

structure of each compound is shown in figure inserts.

540   Figure 4. *Ae. aegypti* odorant receptor concentration response curves.

Normalized mean current responses of AaegOR$_X$ + AaegOrco expressing oocytes (n>10 per

concentration) to (A) 4-chromanone, (B) 1-octen-3-ol, (C) 3-methylindole (D) cis-decahydro-1-

napthol, (E) 4-ethyl-phenol, (F) 1-hexanol, (G) tetrahydrolinalool, (H) farnesol, (I) 2,3-

dimethylindole, and (J) 4-quinolinecarboxyaldehyde. Representative traces from oocyte

recordings are shown above each curve. EC50 values are shown in figure insets.

Supporting Information:

S1 Figure. **RNA sequencing validation of ORs**

Pictures showing RNA abundance and gene structure of selected ten (10) Odorant receptors (Or)

present in unique antennal and maxillary palp reads from Matthews *et al.,* 2018 gotten from

Vectorbase. The scale on the left represents the values of transcripts per million. Highlighted in

red and blue are the exons present in each Or. The actual abundance of RNA per exon is

represented in orange. Numbers at the top of the figures represent chromosome numbers and

positions within the AaegL5 assembly. Vectorbase gene ID and descriptions are present in black

and blue text on the pictures.

S2 Table. **Names and peptide sequences for *Ae. aegypti* ORs and homologs**

Table lists names, VectorBase identification, *Ae. aegypti* homolog, and name attribution

S3 Figure. **Sequence alignments of *de novo* transcripts and ORs**

Transcripts were generated using RNA seq data from Matthews *et al.,* 2018 [35] and *Aedes*

aegypti transcripts (AaegL5) of the ten (10) Odorant receptors (OR). Sequences have been aligned

using ClustalW with character counts (Li *et al.,* 2015). Gene IDs of the AaegL5 transcripts and

those of the denovo transcripts are shown on the left-hand side. Identical residues are highlighted

in gray while different residues are highlighted in black. Missing residues are not highlighted

S4 Table. **List of chemical compounds used in TEVC**

S5 Figure. **Homolog alignments and intron locations**

Alignments between *Ae*. *Aegypti*, *Ae*. *Albopictus*, and *Cx*. *quinquefasciatus* odorant receptors

were completed using Clustal Omenga (https://www.ebi.ac.uk/Tools/msa/clustalo/). Gray

shading indicated shared residues, green shading indicated intron position.

Acknowledgements

This work was supported by a grant from the National Institute of Allergy and Infectious Diseases

(award # 1R01AI148300-01A1 to RJP). The authors also acknowledge Drs. Jeffrey Riffell

(University of Washington) and Omar Akbari (University of California San Diego) and members

of their respective laboratories for helpful discussions of results. The authors are grateful to the

Baylor University Molecular Biosciences Center (Dr. Michelle Nemec, Director) for the use of

various equipment and resources.