Full Title: *Aedes aegypti* (Dipetera: Culicidae) odorant receptors respond to floral and plant derived

volatiles: an expansive deorphanization of odorant receptors expressed in male and female mosquitoes.

Short Title: Mosquitos sense plant derived volatiles

Authors: Heidi Pullmann-Lindsley[1], Robert Huff[1], John Boyi[1], R. Jason Pitts[1]*

[1]Department of Biology,

Baylor University

101 Bagby Avenue, Waco, TX 76706

*corresponding author

10
11

Abstract:

Adult mosquitoes require regular sugar meals, usually in the form of floral nectar, to survive in natural habitats. Both males and females locate potential sugar sources using sensory proteins called odorant receptors, that are activated by plant volatiles and facilitate orientation toward flowers or honeydew. The yellow fever mosquito, *Aedes aegypti* (Linnaeus, 1762), possesses a large gene family of odorant receptors, many of which are likely to be involved in floral odor detection. Using a heterologous expression system and a technique called two-electrode voltage clamping, we are able to identify environmentally relevant chemical cues that produce responses in specific odorant receptors. In this study, we have uncovered ligand-receptor pairings for a suite *Aedes aegypti* receptors that are likely to mediate appetitive or aversive behavioral responses. This information can be used to assess mosquito foraging behavior and develop novel control strategies, especially those that incorporate mosquito bait-and-kill technologies.

Introduction:

Arthropods rely on sensory receptors to interpret and interact with their environments. Arguably, olfaction is the most important sensory modality, conveying the ability to sense volatile chemical cues over large distances [1-3]. Olfaction is responsible for many arthropod behaviors, including host-seeking [4, 5], oviposition [6, 7], mate selection [8, 9], and nectar foraging [10, 11]. Odorant receptors (ORs) comprise the largest family of volatile odor detectors in mosquito genomes [12]. Another family of proteins called ionotropic receptors (IRs) also functions in *Aedes aegypti* olfaction, especially in organic acid detection [13, 14]. Odorant receptors are transmembrane proteins expressed by odorant receptor neurons housed in hairlike projections called sensilla [15, 16]. Sensilla on sensory appendages such as antennae, maxillary palps, labella, and leg tarsi constitute the primary olfactory appendages in insects [17, 18, 19]. ORs are ligand-gated ion channels that respond to diverse, environmentally relevant volatile odorants [20]. ORs most likely form multimers that include co-expression of a single type of tuning OR complexed with an odorant receptor coreceptor (Orco), which is highly conserved across most insect taxa [21-23]. Chemical volatiles enter the sensilla through tiny pores on their surfaces and either diffuse through the sensilla lymph

or are shuttled by odorant binding proteins to the neuronal membrane, where they can interact with the

OR/Orco complex [24, 25]. Most arthropod species have between 50 and 100 different ORs, which allow

for complex and robust interactions with their respective environments through the perception of various

volatile compounds, though some species have as few as five or as many as 300 [26, 27]. The diversity of

receptor proteins underlies distinct behavioral phenomena across arthropod species.

The prominent disease vector *Aedes aegypti* (Linnaeus, 1762) has 133 putative odorant receptors

that vary across sex, life stage, and tissue [28-30]. Despite the significant impact of this species on human

health, few ORs have been characterized, particularly ones presumed to function in female host-seeking

behaviors [31-34]. However, significant knowledge gaps remain in our understanding of the molecular

mechanisms that underpin nectar foraging in this species. As both male and female mosquitoes engage in

this behavior, the ORs that mediate this may provide vital information about potential attractants or

repellants that can be used in trapping or control systems [35-37].

In this study, we investigated ORs in *Ae. aegypti* that are conserved in several disease vector

mosquito species and are closely related to other ORs that function in floral odor sensing. These odorant

receptors are expressed mainly in male and female antennae [28-30], suggesting they may function in a

shared behavior such as nectar foraging. To complete this study, we used a heterologous expression system

and a technique called two-electrode voltage clamping (TEVC) to identify essential ligands [38]. After

synthesizing cRNA for odorant receptors of interest, we injected unfertilized *Xenopus laevis* oocytes with

the cRNA for the OR of interest (called the tuning receptor) and cRNA for *Ae. aegypti* Orco. After allowing

time for protein expression, we used TEVC to record changes in membrane potential as a library of 147

potential ligands was perfused past. Using this method, we identified probable ligands for ten ORs, eight

of which were novel or revealed a previously unknown ligand. Many odorant receptors responded to indole-

like compounds that affect *Ae. aegypti* behavior [39, 40]. These studies highlight the importance of

characterizing ORs as a method of improving our understanding of the molecular basis for *Ae. aegypti*

nectar seeking or other behaviors.

While we have demonstrated that these receptors and ligands interact on a molecular basis, more

research is needed to determine how the ligands might influence a mosquito's behavior. Future studies will

include olfactometry with wild-type and gene-knockout mosquitoes to verify the role of these ORs and

research into the expression patterns of these genes in response to ligand exposure. These studies help to

identify specific attractants or repellants that can be leveraged in surveillance systems or bait traps to help

control mosquito populations and combat the spread of disease.

Materials and Methods

Phylogenetic Analysis

A conserved clade of genes was hypothesized to be activated by floral or nectar odors and utilized by

mosquitoes in foraging. These genes were identified via BLAST via tBLASTn or BLASTp using the NCBI

database. These identified genes were sequenced and cloned for experimental purposes.

Gene cloning and Sequencing

The coding regions of ten genes of interest (Supplemental File 1) and AaegOrco were synthesized by Twist

Bioscience (San Francisco, CA, USA) and cloned into the pENTR plasmid. The Gateway LR directional

cloning system (Invitrogen Corp., Carlsbad, CA, USA) was utilized to subclone AaegOR coding regions

into the pSP64t-RFA oocyte expression plasmid. Plasmids were purified using the GeneJet Plasmid

Miniprep Kit (ThermoFisher Scientific, Waltham, MA, USA). Purified plasmids were sequenced in both

directions to confirm coding regions.

OR Gene Expression Analysis and Transcriptome Assembly

RNAseq reads were downloaded from the NCBI database and mapped to the *Aedes aegypti* annotated

transcript (*Aedes aegypti*_63_LVP_AGWG) obtained from VectorBase. We used the Sailfish alignment-

free software in the Galaxy bioinformatics platform (usegalaxy.org) to map the reads and quantify

expression levels in transcripts per million reads (TPM).

Chemical Reagents

Chemical compounds (Supplemental File 2) used in the deorphanization of AaegORs were purchased from

Acros Organics (Morris, NJ, USA), Alfa Aesar (Ward Hill, MA, USA), and ThermoFisher Scientific

(Waltham, MA, USA) at the highest purity available. Compounds were stored as 1M stocks in 100%

DMSO. Serial dilutions of each compound were prepared in 10% DMSO and blended by chemical class.

Two-electrode voltage clamping

pSP64t-RFA:AaegOR clones were linearized using Xbal and cRNAs were synthesized using the

mMESSAGE mMACHINE ® SP6 kit (Life Technologies, Carlsbad, CA, USA). Xenopus laevis stage V-

VII oocytes purchased from Xenopus1 were stored at 18^0C in Incubation medium (ND96 96 with 5%

dialyzed horse serum,50μg/mL gentamycin,100μg/mL streptomycin, 100μg/mL penicillin, and 550μg/mL

sodium pyruvate). 30nL of cRNA was injected into multiple oocytes (Microinjection Syringe Pump, World

Precision Instruments, Sarasota, FL, USA) maintained at 18C for 48-72 hrs. Odorant blends were perfused

across individual oocytes for 10-15 seconds held at a resting membrane potential of -80 mV using an OC-

725C oocyte clamp (Warner Instruments, LLC, Hamden, CT, USA). Induced currents were measured

indirectly using the two-electrode voltage clamp technique (TEVC). Following each stimulus, currents were

allowed to return to baseline before further stimulation. Data was captured and recorded using the Digidata

1550 B digitizer and Pclamp10 software (Molecular Devices, San Jose, CA, USA). Concentration-response

curves were generated by perfusing unitary odorants over oocytes at increasing concentrations from [10^{-3M}]

to [10^{-9}M]. Data were analyzed using GraphPad Prism 8 (GraphPad Software Inc., La Jolla, CA, USA). At

least two biological replicates were conducted at each concentration and for each compound. Data for at

least seven oocytes that responded to each compound was pooled for analysis (Supplemental File 3.

Results:

1. *Aedes aegypti* odorant receptors are expressed in male and female antennae and conserved in other

disease vector mosquitoes.

Analysis of previously published [41] RNAseq data demonstrated that these ORs are expressed in

both male and female sensory appendages (Table 1). Except for AaegOR8, which is expressed in the

maxillary palps [28]. ORs were most highly expressed in the antennae of both sexes (Table 1; [29]).

Phylogenetic analysis of possible floral volatile sensing odorant receptors in three major disease vectors

revealed several different groupings, which may indicate shared or similar ligands (Figure 1). Orco serves

as an outgroup. This analysis revealed previously deorphanized receptors that share ligands cluster together,

as in the case of CquiOR21 and T2, which are skatole receptors [42]. The cladogram also provides insights

for future functional characterization in other species. AaegOR71 clusters with AalbOR71 and CquiOR32,

AaegOR6, AaegOR13, and AaegOR15 with AalbOR6, AalbOR13, and AalbOR15 respectively,

AaegOR28 with CquiOR117, and AaegOR31 with CquiOR96. The phylogenetic analysis reveals that

newly deorphanized genes are closely related to Ors in other species and can provide helpful information

in future studies with *Culex quinquefasciatus* (Say, 1823) and *Aedes albopictus* (Skuse, 1894).

Table 1: *Ae. aegypti* OR Expression in Sensory Appendages proxied via RNAseq

Odorant receptor gene expression profiles in *Ae. aegypti* male and female antennae and proboscis. Darker

shading indicated elevated transcripts per million.

AaegOR	VectorBase ID	NCBI ID	Transcripts per million (+/- SE) in *Ae. aegypti* head appendages				
			female ant.	male ant.	female palp	male rost.	female prob.
Orco	AAEL005776	NM_001358471	2545.4 (146.6)	385.1 (9.2)	985.6 (87.3)	80.6 (4.2)	35.7 (2.1)
Or6	AAEL017548	NM_001358467	5.3 (1.0)	6.6 (0.5)	0.1 (0.1)	2.6 (0.4)	5.8 (0.7)
Or8	AAEL012254	XM_001662305.3	0 (0.0)	0 (0.0)	387.7 (37.8)	25.7 (1.3)	0.2 (0.2)
Or10	AAEL006003	NM_001358339	21.0 (3.0)	11.5 (0.6)	0 (0.0)	0.5 (0.0)	0.4 (0.1)
Or11	AAEL011583	NM_001358172	146.1 (8.9)	37.0 (1.1)	0 (0.0)	0.2 (0.1)	2.5 (0.5)
Or13	AAEL008368	NM_001358180	20.9 (1.5)	3.8 (0.6)	0 (0.0)	0.3 (0.0)	0.6 (0.1)
Or15	AAEL008448	NM_001358181	6.7 (1.2)	3.3 (0.2)	0 (0.0)	2.2 (0.2)	4.0 (0.3)
Or28	AAEL027053	NM_001358166	4.0 (0.4)	3.4 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)
Or31	AAEL013217	NM_001358160	19.0 (1.5)	10.4 (0.7)	0.2 (0.1)	0.5 (0.0)	1.0 (0.3)
Or55	AAEL010415	NM_001358184	4.4 (0.3)	1.4 (0.3)	0 (0.0)	0.5 (0.1)	1.3 (0.5)
Or71	AAEL017564	NM_001358112	43.1 (2.7)	14.6 (0.3)	0.9 (0.3)	2.9 (0.4)	9.4 (0.7)

Data from VectorBase.org (Matthews *et al.* 2016)

TPM scale

0 ----------- 2545

Figure 1: Phylogenetic Analysis of Possible Floral Volatile Sensing ORs in Major Disease Vectors

Neighbor-joining tree showing relationships of floral volatile sensing OR conceptual peptides for three major disease vector species of mosquitoes *Ae. aegypti* (Aaeg, Black), *Ae. albopictus* (Blue), and *Cu. quinquefasciatus* (green). Node values signify bootstrap values from 1000 pseudoreplicates.

2. Odorant receptors respond to environmentally relevant compounds in a concentration dependent manner

Ae. aegypti ORs responded to blends of chemical compounds (Figure 2). While there is some responsiveness to multiple blends by specific ORs, all demonstrated a pronounced specificity for one blend over the others, as indicated by highly positive kurtosis values (Table 2). The blends that were most frequently responded to were indole or alcohol blends. The unitary compounds for the most responsive blend for each OR was tested individually. We determined the primary ligand each OR responded to from our chemical library and noted other secondary ligands (Table 2, Figure 3). As with the blends, ORs had a pronounced responses to a single ligand, again supported by highly positive kurtosis values. Additionally, a defining feature of OR responsiveness is that it is concentration based, we also investigated responses to increasing concentrations with that compound. From this, we were able to determine the half maximal effective concentration (EC_{50}) for each primary ligand (Table 2, Figure 4).

Figure 2: Heat Map of *Ae. aegypti* odorant receptor responses to blends of chemical compounds

Blends of chemical compounds used for receptor screening are listed as rows on the lefthand side of the

figure while receptor

identities are listed as

columns below the

figure. Heat map

gradation is listed on

the right of the figure

and spans from hot

(100, yellow) to cold (0,

blue).

Table 2: Functional

responses of *Ae. aegypti* odorant receptors

Summarization table of 10 *Ae. aegypti* odorant receptors and the corresponding responsiveness to blends

of chemical compounds and unitary compounds. Kurtosis values are indicated for both blend and unitary

compounds. EC50 values (µM) are indicated for primary ligands. Secondary ligands are presented if

identified during the functional characterizations.

Odorant Receptor	Blend	Kurtosis	Primary Ligand (s)	Kurtosis	EC_{50} (µM)	Secondary Ligand(s)
AaegOr6	Indoles 2	14.86	4-chromanone	10.85	6.84	
AaegOr8	Alcohols 2	15.71	1-octen-3-ol	8.75	2.14	
AaegOr10	Indoles 1	15.61	3-methylindole	5.87	0.16	
AaegOr11	Terpenes 3	5.68	cis-deca-hydro-1-napthol	6.85	68.40	fenchone
AaegOr13	Indoles 1	6.24	4-ethyl-phenol	9.11	0.17	
AaegOr15	Alcohols 2	3.63	1-hexanol	5.23	1.07	phenethyl propionate
AaegOr28	Alcohols 2	15.96	tetrahydrolinalool	7.70	44.13	
AaegOr31	Alcohols 1	14.58	Farnesol	10.70	14.32	
AaegOr55	Indoles 1	14.50	2,3-dimethylindole	0.22	8.65	
AaegOr71	Indoles 2	15.56	4-quinolinecarboxaldehyde	3.08	0.96	

170   **Figure 3: *Ae. aegypti* odorant receptor responses to blend components**

171   Normalized mean current responses of AaegORX/Orco expressing oocytes (n>7) to the individual

172   components of various blends (A) 4-chromanone (B) 1-octen-3-ol, (C) 3-methylindole (D) cis-decahydro-

173   1-napthol, (E) 4-ethyl-phenol, (F) 1-hexanol, (G) tetrahydrolinalool, (H) farnesol, (I) 2,3-dimethylindole,

174   and (J) 4-quinolinecarboxyaldehyde. The structure of each compound is shown in figure inserts.

  Figure 4: *Ae. aegypti* **odorant receptor concentration response curves**

Normalized mean current responses of AaegORX/Orco expressing oocytes (n>10) to (A) 4-chromanone

(B) 1-octen-3-ol, (C) 3-methylindole (D) cis-decahydro-1-napthol, (E) 4-ethyl-phenol, (F) 1-hexanol, (G)

tetrahydrolinalool, (H) farnesol, (I) 2,3-dimethylindole, and (J) 4-quinolinecarboxyaldehyde.

Representative

traces from oocyte

recordings are

shown above each

curve. EC50 values

are shown in figure

      insets.

Three ORs

responded most

robustly to the

Indole 1 blend.

OR10, OR13, and

OR55 responded to

3-methylindole

(0.16 µM), 4-

ethylphenol (0.17

µM), and 2,3-

diemthylindole

(8.65 µM),

respectively. OR6

responded to 4-

chromanone (6.81 μM) from the Indole 2 blend. The other receptor that responded to this blend was OR71

which responded to 4-quinoline carboxaldehyde (0.96 μM). From the Alcohol 1 blend, OR31 responded to

farnesol (14.32 μM), while from the Alcohols 2 blend, OR8 and OR15 responded to 1-octen-3-ol (2.14

μM) and 1-hexanol (1.07 μM), respectively. A previously identified ligand of OR15, phenethyl propionate,

demonstrated an EC_{50} of 20.16 μM. OR28 responded to tetrahydrolinalool (44.13 μM) from the Alcohols 3

blend. Finally, the only receptor to respond to a compound not belonging to an indole or alcohol blend was

OR11, which responded to cis-decahydro-1-naphthal (68.40 μM) from the Terpenes 3 blend. We found that

OR11 has an EC_{50} of about 77.10 μM to a previously identified ligand, fenchone.

Discussion:

Nectar foraging is crucial for mosquito species to intake sugar meals for reproduction and survival

[43-46]. Previous studies have shown that several mosquito species, including *Ae. aegypti,* exhibit

preferences for certain nectar sources [47-52]. ORs provide essential details about the surrounding

environment and allow mosquitoes to identify and select nectar sources based on the information provided

by their chemical signals [3]. Here, we assess the responsiveness of various ORs to a library of ecologically

relevant ligands, focusing on receptors that could be related to floral odor sensing.

We began by looking at phylogeny and RNAseq data. This allowed us to select potential targets

based on expression patterns and determine relatedness between ORs within *Ae. aegypti* and between two

other disease vectors, *Cu. quinquefasciatus* and *Ae. albopictus,* assuming that functionally related ORs

would cluster together [53]. Specifically, we selected specific ORs to study based on their relatedness to

other ORs that respond to floral odors [54] or had previously been deorphanized. AaegOR10, AaegOR9,

AaegOR2, and AaegOR8 have been thoroughly investigated previously, and many of the ORs we selected

for study were closely related to them. When looking at RNAseq data, we were interested in ORs expressed

in antennae. While AaegORs are expressed in various chemosensory tissues, antennae are the primary

structures involved in odor sensing [1]. Additionally, OR expression in the antennae of both males and

females indicates that the receptors are used for a shared behavior, such as nectar feeding. These initial

analyses allowed us to predict and select receptors that would respond to our chemical library.

Using two-electrode voltage clamping in conjunction with the heterologous expression system

Xenopus laevis oocytes, we initially assessed the responses of the AaegORs to our chemical library, which

contains 147 environmentally significant compounds organized into 14 blends. This revealed that many of

the AaegORs responded most robustly to the "Indole 1" and "Indole 2" blends, which contain indole or

indole-like compounds (Supp. Table 1). Indoles are relatively ubiquitous in natural settings and are present

in potential nectar sources. AaegOR1 responds to 4-quinoline carboxaldehyde, which has been found in

honey samples and endophytic bacteria via gas chromatography [55-57]. However, indole and its related

compounds have many other sources and influence behaviors across species [58], so it is uncertain if these

ORs are directly related to floral odor sensing. The specific indoles that several ORs respond to are not

typical plants, and their sources may suggest other functions, such as oviposition or host-seeking. AaegOR

responds to 4-chromanone, of which there is little literature regarding its occurrence in natural settings.

However, it is a key element of flavonoid skeletons [59]. AaegOR10 responds to 3-methylindole (skatole),

which is found in animal feces and produces behavioral responses in other Dipteran species [60-63]. 4-

ethylphenol, the primary ligand of AaegOR13, is a volatile constituent in wine and animal waste and has

been linked to oviposition in *Culex pipiens* [64-66]. While there is little evidence about natural sources of

2,3-dimethylindole, the AaegOR55 ligand it is closely related structurally to 3-methylindole. Other ORs

responded to compounds from the "Alcohols 1", "Alcohols 2", or "Alcohols 3" blends. AaegOR8 responds

to 1-octen-3-ol, a well-characterized fungal volatile and insect attractant [67]. A green-leaf volatile, 1-

hexanol, is another compound with well-characterized physiological effects in many arthropods [68-71].

Our study discovered that 1-hexanol activates AaegOR15 with a low threshold (EC_{50} ~1.07 μM).

AaegOR28 responds to tetrahydrolinalool, which is structurally related to linalool, another plant-derived

ligand to mosquito ORs [72]. AaegOR31 responds to farnesol, a significant constituent of the honeybee

Nasanov pheromone [73]. Lastly, the AaegOR11 ligand, cis-decahydro-1-napthol, is structurally related to

geosmin, a known oviposition attractant [74]. However, our compound library needs to be more comprehensive, as more efficacious or potent ligands for AaegOR11 and AaegOR28 may exist, which had the highest EC_{50} values in this set of ORs. We will continue to expand our library by adding compounds structurally related to known ligands.

While four (AaegOR8, AaegOR10, AaegOR11, and AaegOR15) of the ORs tested in this study already have identified ligands, we assessed them to uncover more potentially behaviorally relevant ligands, as our chemical library differs from others tested [40, 75-77]. For AaegOR8 and AaegOR10, our data supports what has previously been demonstrated. For AaegOR11 and AaegOR15, our data both supports and adds to previous studies. For AaegOR11, we found that cis-decahydro-1-napthol produced a greater magnitude of response than (-)-fenchone or (+)-fenchone, and we identified 1-hexanol as a potent activator of AaegOR15 in addition to phenethyl propionate. This demonstrates that even for deorphanized ORs, assessing their function with expanded libraries can aid in further understanding receptor-ligand interactions.

Ae. Aegypti and other vector mosquitoes continue to threaten human health and flourish worldwide. By continuing to elucidate receptor-ligand pairings, we can identify potential attractants for behavioral control or targets for genetic knockout. By investigating floral-odor sensing receptors in particular, we can identify compounds that are potentially behaviorally relevant to both sexes and can be used for attractive toxic sugar bait systems. OR deorphanization is a valuable first step in understanding the sensory systems of arthropods and can inform future experiments and control measures.

Funding Disclosure:

This work was supported by the National Institutes of Allergy and Infectious Diseases at the National Institutes of Health [grant number R01 AI148300-01A1 to RJP].

278 **Literature Cited:**
279

1. Hansson BS, Stensmyr MC. Evolution of Insect Olfaction. Neuron. 2011 Dec 8;72(5):698–
711.

2. Ray A. Reception of odors and repellents in mosquitoes. Current Opinion in
Neurobiology. 2015 Oct 1;34:158–64.

3. Lutz EK, Lahondère C, Vinauger C, Riffell JA. Olfactory learning and chemical ecology
of olfaction in disease vector mosquitoes: a life history perspective. Current Opinion in
Insect Science. 2017 Apr 1;20:75–83.

4. Sun H, Liu F, Ye Z, Baker A, Zwiebel LJ. Mutagenesis of the orco odorant receptor co-
receptor impairs olfactory function in the malaria vector Anopheles coluzzii. Insect
Biochem Mol Biol. 2020 Dec;127:103497.

5. Huff RM, Pitts RJ. Carboxylic acid responses by a conserved odorant receptor in culicine
vector mosquitoes. Insect Mol Biol. 2020 Dec;29(6):523–30.

6. Hughes DT, Pelletier J, Luetje CW, Leal WS. Odorant receptor from the southern house
mosquito narrowly tuned to the oviposition attractant skatole. J Chem Ecol. 2010
Aug;36(8):797–800.

7. Liu H, Liu L, Cheng P, Huang X, Gong M. An odorant receptor from Anopheles sinensis
in China is sensitive to oviposition attractants. Malar J. 2018 Oct 5;17(1):348.

8. Dahanukar A, Hallem EA, Carlson JR. Insect chemoreception. Curr Opin Neurobiol. 2005
Aug;15(4):423–30.

9. Li RT, Huang LQ, Dong JF, Wang CZ. A moth odorant receptor highly expressed in the
ovipositor is involved in detecting host-plant volatiles. Elife. 2020 May 21;9:e53706.

10. Huff RM, Pitts RJ. An odorant receptor from *Anopheles gambiae* that demonstrates
enantioselectivity to the plant volatile, linalool. PLoS One. 2019;14(11):e0225637.

11. Dekel A, Yakir E, Bohbot JD. The sulcatone receptor of the strict nectar-feeding mosquito Toxorhynchites amboinensis. Insect Biochemistry and Molecular Biology. 2019 Aug 1;111:103174.

12. Robertson HM. Molecular Evolution of the Major Arthropod Chemoreceptor Gene Families. Annual Review of Entomology. 2019;64(1):227–42.

13. Raji JI, Melo N, Castillo JS, Gonzalez S, Saldana V, Stensmyr MC, et al. *Aedes aegypti* Mosquitoes Detect Acidic Volatiles Found in Human Odor Using the IR8a Pathway. Curr Biol. 2019 Apr 22;29(8):1253-1262.e7.

14. Stocker RF. Drosophila as a focus in olfactory research: Mapping of olfactory sensilla by fine structure, odor specificity, odorant receptor expression, and central connectivity. Microscopy Research and Technique. 2001;55(5):284–96.

15. Jiang X, Breer H, Pregitzer P. Sensilla-Specific Expression of Odorant Receptors in the Desert Locust Schistocerca gregaria. Front Physiol. 2019;10:1052.

16. Sutcliffe JF. Sensory bases of attractancy: morphology of mosquito olfactory sensilla-- a review. J Am Mosq Control Assoc. 1994 Jun;10(2 Pt 2):309–15.

17. Zahran N, Sawires S, Hamza A. Piercing and sucking mouth parts sensilla of irradiated mosquito, Culex pipiens (Diptera: Culicidae) with gamma radiation. Sci Rep. 2022 Oct 25;12(1):17833.

18. Yamany AS, Abdel-Gaber R. Studies on the sensory sensilla on the tarsi and external genitalia of the Asian tiger mosquito, *Aedes albopictus* (Skuse). Microsc Res Tech. 2023 Feb;86(2):242–51.

19. del Mármol J, Yedlin MA, Ruta V. The structural basis of odorant recognition in insect olfactory receptors. Nature. 2021 Sep;597(7874):126–31.

20. Tiwari V, Sowdhamini R. Structure modelling of odorant receptor from *Aedes aegypti* and identification of potential repellent molecules. Computational and Structural Biotechnology Journal. 2023 Jan 1;21:2204–14.

21. Tiwari V, Sowdhamini R. Structure modelling of odorant receptor from *Aedes aegypti* and identification of potential repellent molecules. Computational and Structural Biotechnology Journal. 2023 Jan 1;21:2204–14.

22. Stengl M, Funk NW. The role of the coreceptor Orco in insect olfactory transduction. J Comp Physiol A Neuroethol Sens Neural Behav Physiol. 2013 Nov;199(11):897–909.

23. Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, et al. Cryo-EM structure of the insect olfactory receptor Orco. Nature. 2018 Aug;560(7719):447–52.

24. Zhou JJ. Chapter Ten - Odorant-Binding Proteins in Insects. In: Litwack G, editor. Vitamins & Hormones [Internet]. Academic Press; 2010 [cited 2023 Jul 6]. p. 241–72. (Pheromones; vol. 83). Available from: https://www.sciencedirect.com/science/article/pii/S0083672910830109

25. Li RT, Huang LQ, Dong JF, Wang CZ. A moth odorant receptor highly expressed in the ovipositor is involved in detecting host-plant volatiles. Elife. 2020 May 21;9:e53706.

26. Leal WS. Odorant Reception in Insects: Roles of Receptors, Binding Proteins, and Degrading Enzymes. Annual Review of Entomology. 2013;58(1):373–91.

27. Brand P, Robertson HM, Lin W, Pothula R, Klingeman WE, Jurat-Fuentes JL, et al. The origin of the odorant receptor gene family in insects. Benton R, Tautz D, editors. eLife. 2018 Jul 31;7:e38340.

28. Bohbot J, Pitts RJ, Kwon HW, Rützler M, Robertson HM, Zwiebel LJ. Molecular

characterization of the *Aedes aegypti* odorant receptor gene family. Insect Mol Biol. 2007

Oct;16(5):525–37.

29. Matthews BJ, McBride CS, DeGennaro M, Despo O, Vosshall LB. The

neurotranscriptome of the *Aedes aegypti* mosquito. BMC Genomics. 2016 Jan 6;17:32.

30. Matthews BJ, Dudchenko O, Kingan SB, Koren S, Antoshechkin I, Crawford JE, et al.

Improved reference genome of *Aedes aegypti* informs arbovirus vector control. Nature.

2018 Nov;563(7732):501–7.

31. Coutinho-Abreu IV, Riffell JA, Akbari OS. Human attractive cues and mosquito host-

seeking behavior. Trends in Parasitology. 2022 Mar 1;38(3):246–64.

32. Wang X, Chen Z, Wang Y, Liu F, Jiang S, Liu N. Heterologous Expression and Functional

Analysis of *Aedes aegypti* Odorant Receptors to Human Odors in Xenopus Oocytes. JoVE

(Journal of Visualized Experiments). 2021 Jun 8;(172):e61813.

33. Ghaninia M, Larsson M, Hansson BS, Ignell R. Natural odor ligands for olfactory receptor

neurons of the female mosquito *Aedes aegypti*: use of gas chromatography-linked single

sensillum recordings. J Exp Biol. 2008 Sep;211(Pt 18):3020–7.

34. Huff RM, Pitts RJ. Carboxylic acid responses by a conserved odorant receptor in culicine

vector mosquitoes. Insect Molecular Biology. 2020;29(6):523–30.

35. Kashiwagi GA, von Oppen S, Harburguer L, González-Audino P. The main component

of the scent of Senecio madagascariensis flowers is an attractant for *Aedes aegypti* (L.)

(Diptera: Culicidae) mosquitoes. Bull Entomol Res. 2022 Dec;112(6):837–46.

36. Sissoko F, Junnila A, Traore MM, Traore SF, Doumbia S, Dembele SM, et al. Frequent

sugar feeding behavior by *Aedes aegypti* in Bamako, Mali makes them ideal candidates

for control with attractive toxic sugar baits (ATSB). PLoS One. 2019;14(6):e0214170.

37. Fikrig K, Johnson BJ, Fish D, Ritchie SA. Assessment of synthetic floral-based attractants

and sugar baits to capture male and female *Aedes aegypti* (Diptera: Culicidae). Parasit

Vectors. 2017 Jan 17;10(1):32.

38. Mowry KL. Using the Xenopus Oocyte Toolbox. Cold Spring Harb Protoc. 2020 Apr

1;2020(4):095844.

39. Baak-Baak CM, Rodríguez-Ramírez AD, García-Rejón JE, Ríos-Delgado S, Torres-

Estrada JL. Development and laboratory evaluation of chemically-based baited ovitrap

for the monitoring of *Aedes aegypti*. J Vector Ecol. 2013 Jun;38(1):175–81.

40. Ruel DM, Yakir E, Bohbot JD. Supersensitive Odorant Receptor Underscores Pleiotropic

Roles of Indoles in Mosquito Ecology. Frontiers in Cellular Neuroscience [Internet]. 2019

[cited   2023   Jul   12];12.   Available   from:

https://www.frontiersin.org/articles/10.3389/fncel.2018.00533

41. Matthews BJ. *Aedes aegypti*. Trends in Genetics. 2019 Jun 1;35(6):470–1.

42. Choo YM, Buss GK, Tan K, Leal WS. Multitasking roles of mosquito labrum in

oviposition and blood feeding. Front Physiol. 2015;6:306.

43. Foster WA. Mosquito Sugar Feeding and Reproductive Energetics. Annual Review of

Entomology. 1995;40(1):443–74.

44. Stone CM, Taylor RM, Roitberg BD, Foster WA. Sugar Deprivation Reduces

Insemination of *Anopheles gambiae* (Diptera: Culicidae), Despite Daily Recruitment of

Adults, and Predicts Decline in Model Populations. Journal of Medical Entomology. 2009
Nov 1;46(6):1327–37.

45. Dieng H, Satho T, Binti Arzemi NA, Aliasan NE, Abang F, Wydiamala E, et al. Exposure
of a diurnal mosquito vector to floral mimics: Foraging responses, feeding patterns, and
significance for sugar bait technology. Acta Trop. 2018 Sep;185:230–8.

46. Barredo E, DeGennaro M. Not Just from Blood: Mosquito Nutrient Acquisition from
Nectar Sources. Trends Parasitol. 2020 May;36(5):473–84.

47. Grimstad PR, DeFoliart GR. Nectar Sources of Wisconsin Mosquitoes1. Journal of
Medical Entomology. 1974 Jul 15;11(3):331–41.

48. Vargo AM, Foster WA. Responsiveness of Female *Aedes aegypti* (Diptera: Culicidae) to
Flower Extracts. Journal of Medical Entomology. 1982 Nov 30;19(6):710–8.

49. Gouagna LC, Poueme RS, Dabiré KR, Ouédraogo JB, Fontenille D, Simard F. Patterns of
sugar feeding and host plant preferences in adult males of *An. gambiae* (Diptera:
Culicidae). Journal of Vector Ecology. 2010;35(2):267–76.

50. Chen Z, Kearney CM. Nectar protein content and attractiveness to *Aedes aegypti* and
Culex pipiens in plants with nectar/insect associations. Acta Trop. 2015 Jun;146:81–8.

51. Lahondère C, Vinauger C, Okubo RP, Wolff GH, Chan JK, Akbari OS, et al. The olfactory
basis of orchid pollination by mosquitoes. Proceedings of the National Academy of
Sciences. 2020 Jan 7;117(1):708–16.

52. Nikbakhtzadeh MR, Terbot II JW, Otienoburu PE, Foster WA. Olfactory basis of floral
preference of the malaria vector *Anopheles gambiae* (Diptera: Culicidae) among common
African plants. Journal of Vector Ecology. 2014;39(2):372–83.

53. Bohbot JD, Jones PL, Wang G, Pitts RJ, Pask GM, Zwiebel LJ. Conservation of indole responsive odorant receptors in mosquitoes reveals an ancient olfactory trait. Chem Senses. 2011 Jan;36(2):149–60.

54. Syed Z, Leal WS. Maxillary palps are broad spectrum odorant detectors in *Culex quinquefasciatus*. Chem Senses. 2007 Oct;32(8):727–38.

55. Rodríguez-Cabo T, Moniruzzaman M, Rodríguez I, Ramil M, Cela R, Gan SH. Time-of-flight accurate mass spectrometry identification of quinoline alkaloids in honey. Anal Bioanal Chem. 2015 Aug 1;407(20):6159–70.

56. Moniruzzaman M, Rodríguez I, Ramil M, Cela R, Sulaiman SA, Gan SH. Assessment of gas chromatography time-of-flight accurate mass spectrometry for identification of volatile and semi-volatile compounds in honey. Talanta. 2014 Nov 1;129:505–15.

57. Naama-Amar A, Gitman S, Shoshana N, Bahar O, Naor V, Zchori-Fein E, et al. Antimicrobial Activity of Metabolites Secreted by the Endophytic Bacterium Frateuria defendens. Plants. 2020 Jan;9(1):72.

58. Tomberlin JK, Crippen TL, Wu G, Griffin AS, Wood TK, Kilner RM. Indole: An evolutionarily conserved influencer of behavior across kingdoms. BioEssays. 2017;39(2):1600203.

59. Liu Y, Qian X, Xing J, Li N, Li J, Su Q, et al. Accurate Determination of 12 Lactones and 11 Volatile Phenols in Nongrape Wines through Headspace-Solid-Phase Microextraction (HS-SPME) Combined with High-Resolution Gas Chromatography-Orbitrap Mass Spectrometry (GC-Orbitrap-MS). J Agric Food Chem. 2022 Feb 16;70(6):1971–83.

60. Yokoyama M, Carlson J. Microbial metabolites of tryptophan in the intestinal tract with
special reference to skatole1 2. The American Journal of Clinical Nutrition. 1979 Jan
1;32(1):173–8.

61. Cook KL, Rothrock MJ, Loughrin JH, Doerner KC. Characterization of skatole-producing
microbial populations in enriched swine lagoon slurry. FEMS Microbiol Ecol. 2007
May;60(2):329–40.

62. Jeanbourquin P, Guerin PM. Sensory and behavioural responses of the stable fly
Stomoxys calcitrans to rumen volatiles. Med Vet Entomol. 2007 Sep;21(3):217–24.

63. Johnson SD, Sivechurran J, Doarsamy S, Shuttleworth A. Dung mimicry: the function of
volatile emissions and corolla patterning in fly-pollinated Wurmbea flowers. New
Phytologist. 2020;228(5):1662–73.

64. Rodríguez-Cabo T, Moniruzzaman M, Rodríguez I, Ramil M, Cela R, Gan SH. Time-of-
flight accurate mass spectrometry identification of quinoline alkaloids in honey. Anal
Bioanal Chem. 2015 Aug 1;407(20):6159–70.

65. Musonye HA, Njeru EM, Hassanali A, Langata LM, Mijele D, Kaitho T, et al. 16S rRNA
gene profiling of bacterial communities mediating production of tsetse attractive phenols
in mammalian urine. Onderstepoort Journal of Veterinary Research. 2019 Jul 17;86(1):12.

66. Zhu F, Xu P, Barbosa RMR, Choo YM, Leal WS. RNAi-based demonstration of direct
link between specific odorant receptors and mosquito oviposition behavior. Insect
Biochemistry and Molecular Biology. 2013 Oct 1;43(10):916–23.

67. Kline DL. Olfactory attractants for mosquito surveillance and control: 1-octen-3-ol. J Am
Mosq Control Assoc. 1994 Jun;10(2 Pt 2):280–7.

68. De Jong R, Visser JH. Integration of olfactory information in the Colorado potato beetle brain. Brain Research. 1988 Apr 26;447(1):10–7.

69. Dickens JC, Prestwich GD, Ng CS, Visser JH. Selectively fluorinated analogs reveal differential olfactory reception and inactivation of green leaf volatiles in insects. J Chem Ecol. 1993 Sep 1;19(9):1981–91.

70. Williams L, Blackmer JL, Rodriguez-Saona C, Zhu S. Plant Volatiles Influence Electrophysiological and Behavioral Responses of Lygus hesperus. J Chem Ecol. 2010 May 1;36(5):467–78.

71. von Arx M, Schmidt-Büsser D, Guerin PM. Host plant volatiles induce oriented flight behaviour in male European grapevine moths, Lobesia botrana. J Insect Physiol. 2011 Oct;57(10):1323–31.

72. Huff RM, Pitts RJ. Functional conservation of Anopheline linalool receptors through 100 million years of evolution. Chemical Senses. 2022 Jan 1;47:bjac032.

73. Williams IH, Pickett JA, Martin AP. The Nasonov pheromone of the honeybeeApis mellifera L. (Hymenoptera, Apidae). Part II. Bioassay of the components using foragers. J Chem Ecol. 1981 Mar;7(2):225–37.

74. Melo N, Wolff GH, Costa-da-Silva AL, Arribas R, Triana MF, Gugger M, et al. Geosmin Attracts *Aedes aegypti* Mosquitoes to Oviposition Sites. Current Biology. 2020 Jan 6;30(1):127-134.e5.

75. Bohbot JD, Dickens JC. Characterization of an Enantioselective Odorant Receptor in the Yellow Fever Mosquito *Aedes aegypti*. PLOS ONE. 2009 Sep 15;4(9):e7032.