Title: Distinguishing mutants that resist drugs via different mechanisms by examining fitness tradeoffs

Keywords: Fitness tradeoffs, drug resistance, DNA barcodes, fitness profiles, gene-by-environment
interactions, molecular mechanisms, genotype-phenotype map, azole resistance

Authors:

Schmidlin[1,2]*, Apodaca[1,2]*, Newell[1,2], Sastokas[1,2], Kinsler[3], Geiler-Samerotte[1,2]

* Co-first authors

Affiliations:

1 Biodesign Center for Mechanisms of Evolution, Arizona State University, Tempe, AZ

2 School of Life Sciences, Arizona State University, Tempe AZ

3 Department of Bioengineering, University of Pennsylvania, Philadelphia, PA

Corresponding Author:

kerry.samerotte@asu.edu

Abstract
There is growing interest in designing multidrug therapies that leverage tradeoffs to combat resistance.
Tradeoffs are common in evolution and occur when, for example, resistance to one drug results in sensitivity
to another. Major questions remain about the extent to which tradeoffs are reliable, specifically, whether
the mutants that provide resistance to a given drug all suffer similar tradeoffs. This question is difficult
because the drug-resistant mutants observed in the clinic, and even those evolved in controlled laboratory
settings, are often biased towards those that provide large fitness benefits. Thus, the mutations (and
mechanisms) that provide drug resistance may be more diverse than current data suggests. Here, we perform
evolution experiments utilizing lineage-tracking to capture a fuller spectrum of mutations that give yeast
cells a fitness advantage in fluconazole, a common antifungal drug. We then quantify fitness tradeoffs for
each of 774 evolved mutants across 12 environments, finding these mutants group into 6 classes with
characteristically different tradeoffs. Their unique tradeoffs may imply that each group of mutants affects
fitness through different underlying mechanisms. Some of the groupings we find are surprising. For
example, we find some mutants that resist single drugs do not resist their combination, while others do.
And some mutants to the same gene have different tradeoffs than others. These findings, on one hand,
demonstrate the difficulty in relying on consistent or intuitive tradeoffs when designing multidrug
treatments. On the other hand, by demonstrating that hundreds of adaptive mutations can be reduced to a
few groups with characteristic tradeoffs, our findings may yet empower multidrug strategies that leverage
tradeoffs to combat resistance. More generally speaking, by grouping mutants that likely affect fitness
through similar underlying mechanisms, our work guides efforts to map the phenotypic effects of mutation.

Introduction

How many different molecular mechanisms can a microbe exploit to adapt to a challenging environment? Answering this question is particularly urgent in the field of drug resistance because infectious populations are adapting to available drugs faster than new drugs are developed (Centers for Disease Control and Prevention (U.S.), 2019; Ventola, 2015). Understanding the mechanistic basis of drug resistance can inform strategies for how to combine existing drugs in a way that prevents the evolution of resistance (Andersson and Hughes, 2010; Melnikov et al., 2020; Pinheiro et al., 2021). For example, one strategy exposes an infectious population to one drug (Drug A) knowing that the mechanism of resistance to Drug A makes cells susceptible to Drug B (Baym et al., 2016; Hall et al., 2009; Pál et al., 2015; Roemhild et al., 2020). Problematically, these multi-drug strategies perform best when all mutants that resist Drug A have the same tradeoff in Drug B (**Figure 1A**). If there are multiple different mechanisms to resist Drug A, some of which lack this tradeoff, treatment strategies could fail (**Figure 1B**), and they sometimes do (Abel zur Wiesch et al., 2014; Grier et al., 2003; Scarborough et al., 2020; Wang et al., 2019).

Figure 1: A multidrug treatment strategy that relies on all mutants having the same tradeoffs. (A) All of the mutants that resist Drug A do so via a similar mechanism such that all are sensitive to Drug B. **(B)** There are multiple different types of mutants that resist Drug A, not all of which are sensitive to Drug B.

Laboratory experiments that have power to search for universal tradeoffs – where all the mutants that perform well in one environment perform poorly in another – often find there are mutants that violate trends or the absence of trends altogether (Ardell and Kryazhimskiy, 2021; Gjini and Wood, 2021; Herren and Baym, 2022; Hill et al., 2015; Kinsler et al., 2020; Nichol et al., 2019). Another way to phrase this observation is to say that adaptive mutations often have collateral effects in environments other than the one in which they originally evolved (Pál et al., 2015). But these effects, referred to in some studies as pleiotropic effects, can be unpredictable and context dependent (Bakerlee et al., 2021; Chen et al., 2023; Geiler-Samerotte et al., 2020; Hinz et al., 2023; Jerison et al., 2020). In simpler terms, some mutants that resist Drug A will suffer a tradeoff in Drug B, but others may suffer a tradeoff in Drug C. To sum, observations from many fields suggest that the mutations that provide a benefit in one environment do not always suffer similar tradeoffs. This begs questions about the extent of diversity among adaptive mutations: does each one suffer a unique set of tradeoffs or can many adaptive mutants be grouped by

75   their common tradeoffs? If there are only a few types of tradeoff present in a collection of adaptive
76   mutations, multidrug treatments that target tradeoffs may be more feasible.
77
The goal of the present study is to count how many different types of adaptive mutation, each type being
defined by its unique tradeoffs, exist in a population of drug-resistant yeast. This simple counting exercise
is surprisingly difficult. One reason why is that the mutations that provide the strongest fitness advantage
often dominate evolution. Thus, in the clinic, and in laboratory experiments, the same drug-resistant
mutations repeatedly emerge (Berkow and Lockhart, 2017; Ksiezopolska et al., 2021; Lupetti et al., 2002;
Melnikov et al., 2020), potentially leading to the false impression that the mechanistic basis of resistance
to a particular drug, and the associated tradeoffs, are less varied than may be true. This problem is
amplified by the limitations of bulk DNA sequencing methods which often miss mutations that are
present in less than 10% of a population's cells (Good et al., 2017). A similar problem results from
strategies to disentangle adaptive from passenger mutations that rely on observing the adaptive ones
multiple times in multiple independent replicates (Martínez and Lang, 2023). In order to design better
multidrug treatment strategies that thwart resistance, or to see if such strategies are feasible, we need
methods to survey a more complete set of mutations and mechanisms that can contribute to resistance.
91
Fortunately, single-cell and single-lineage DNA sequencing technologies are allowing us to more deeply
sample genetic diversity in evolving populations of microbes beyond the mutations that dominate
evolution (Schmidt and Efferth, 2016). Here, we leverage a cutting-edge lineage-tracing method to
perform massively-replicate evolution experiments in yeast (*S. cerevisiae*). This method has been shown
to reveal a fuller spectrum of mutations underlying adaptation to a particular environment (Levy et al.,
2015). One key to its success is that it uses DNA barcodes to track all competing adaptive lineages, not
just the ones that ultimately rise to appreciable frequency. Another key feature is that it captures adaptive
lineages before they accumulate multiple mutations such that it is easy to pinpoint which mutation is
adaptive. We apply this method to investigate mechanisms underlying resistance to a specific antifungal
drug: fluconazole (FLU) (Logan et al., 2022; Wang et al., 2022). Although serious fungal infections are
most common in immunocompromised individuals, their impact on global health is still striking, resulting
in over 1.5 million deaths annually (Iyer et al., 2022; Xie et al., 2014). By focusing on mechanisms of
FLU resistance, we contribute to a growing literature about the tradeoffs that may be leveraged to design
multidrug antifungal treatment strategies (Cowen and Lindquist, 2005; Hill et al., 2015; Iyer et al., 2022;
Ksiezopolska et al., 2021). However, our primary goal is more generic: we seek to explore the utility of a
high-throughput evolutionary approach to enumerate classes of drug resistant mutant and their associated
tradeoffs.
109
To enhance the diversity of drug resistant mutants in our experiment, we performed multiple laboratory
evolutions in a range of FLU concentrations and sometimes in combination with a second drug. We did
so because previous work has shown that different drug concentrations and combinations select for
different azole resistance mechanisms (Cowen and Lindquist, 2005; Hill et al., 2015). Ultimately, we
obtained a large collection of 774 adaptive yeast strains. But how do we know whether we succeeded in
isolating diverse types of FLU-resistant mutants? Typical phenotyping methods, *e.g.,* quantifying
expression levels of drug export pumps (Miyazaki et al., 1998) or of the drug targets themselves (Palmer
and Kishony, 2014), are low throughput and require some a priori knowledge of the phenotypes that may
be involved in drug resistance. Alternatively, many studies focus on identifying the genetic basis of

adaptation in order to glean insights about the underlying mechanisms of resistance (Cowen et al., 2014;
Tenaillon et al., 2012; Venkataram et al., 2016). However, genotyping lineages from barcoded pools is
technically challenging (Venkataram et al., 2016), and further, genotype does not necessarily predict
phenotype (Brettner et al., 2022; Eguchi et al., 2019). For example, previous work using the same
barcoded evolution platform used here discovered that the mutations that provide an advantage under
glucose-limitation are in genes comprising a canonical glucose-sensing pathway (Venkataram et al.,
2016). Yet despite this similarity at the genetic level, follow-up work showed that these mutants did not
experience the same tradeoffs when exposed to new environments (Kinsler et al., 2020; Y. Li et al.,
2018).

Instead of trying to identify the phenotypic or even the genetic basis of adaptation, here we strive to
enumerate different classes of FLU-resistant mutants. We sort evolved FLU-resistant yeast strains into
classes based on whether they share similar tradeoffs across 12 different environments. The intuition here
is as follows. If two groups of drug resistant mutants have different fitness tradeoffs, it could mean that
they provide resistance through different underlying mechanisms. Alternatively, both could provide drug
resistance via the same mechanism, but some mutations might also affect fitness via additional
mechanisms (i.e. they might have unique "side-effects" at the molecular level) resulting in unique fitness
tradeoffs in some environments. Previous work is consistent with the idea that mutants with different
fitness tradeoffs affect fitness through different underlying mechanisms (Li et al., 2019; Pinheiro et al.,
2021; Rodrigues et al., 2016). Our work can be seen as part of a growing push to flip the problem of
mechanism on its head (Kinsler et al., 2020; Li et al., 2019; Petti et al., 2023). Instead of using a
mechanistic understanding to predict a microbe's fitness, here we use how fitness varies across
environments to distinguish mutants that likely affect fitness via different mechanisms. This inverted
approach to investigating the mechanisms by which mutations affect fitness has broad applications; it
could be used to characterize dominant negative mutations (Flynn et al., 2024; Padhy et al., 2023),
mutations with collateral fitness effects (Mehlhoff et al., 2020; Mehlhoff and Ostermeier, 2023), and in
other high-throughput mutational scanning studies (Flynn et al., 2020; Fowler and Fields, 2014; Hinz et
al., 2023; Starr et al., 2017).

The key requirement to being able to implement this approach is having a large collection of barcoded
mutants and the ability to re-measure their fitness, relative to a reference strain, in multiple environments,
such as the 12 different combinations and concentrations of drugs surveyed here. Across our collection of
774 adaptive yeast lineages, we discovered at least 6 distinct groups with characteristic fitness tradeoffs
across these 12 environments. For example, we find some drug resistant mutants are generally
advantageous, while others are advantageous only in specific environments. And we find some mutants
that resist single drugs also resist combinations of those drugs, while others do not. By grouping mutants
with similar tradeoffs, we reduce the number of unique drug-resistant mutants from more than can be
easily phenotyped (774) to a manageable panel of 6 types for investigating the molecular mechanisms by
which mutations impact fitness.

With regard to multidrug regimens that exploit tradeoffs (**Figure 1**), our finding of multiple mutant
classes with different tradeoffs suggests this may not be straightforward. The outlook is further
complicated by our finding that some classes of FLU-resistant mutant primarily emerge from evolution
experiments that did not contain FLU. This, as well as limits on our power to observe mutants with strong

tradeoffs, suggest there may be additional types of FLU-resistant mutant beyond those we sampled. These
observations suggest multidrug strategies that assume resistant mutants suffer consistent or common
tradeoffs will often fail.

On the other hand, nuanced strategies to forestall resistance that allow for multiple mutant types are
emerging (Gjini and Wood, 2021; Maltas and Wood, 2019; Wang et al., 2023). For example, one idea is
to apply a drug regimen that enriches for mutants that suffer a particular tradeoff before exploiting that
tradeoff (Iram et al., 2020). Another idea is to perform single-cell sequencing on infectious populations to
discover which classes of mutants are present (Forsyth et al., 2021; Nagasawa et al., 2021) and design
treatments specific to those (Aissa et al., 2021; Hsieh et al., 2022; Maltas and Wood, 2019). Our findings
support that such ideas may be feasible by demonstrating that there are not as many unique fitness
tradeoffs as there are mutations.

More generally, our work – showing that 774 mutants fall into a much smaller number of groups –
contributes to growing literature suggesting that the phenotypic basis of adaptation is not as diverse as the
genetic basis (Iwasawa et al., 2022; Kinsler et al., 2020; Petti et al., 2023). This winnowing of diversity is
important: it may mean that evolutionary processes, for example, whether an infectious population will
adapt to resist a drug, are sometimes predictable (King et al., 2022; Kinsler et al., 2020; Lässig et al.,
2017; Rodrigues et al., 2016; Wortel et al., 2023; Yoon et al., 2021)**.**

Results
Barcoded evolution experiments uncover hundreds of yeast lineages with adaptive mutations
In order to create a sizable collection of drug-resistant mutants, we performed high-replicate evolution
experiments utilizing barcoded yeast (*S. cerevisiae*) (Boyer et al., 2021; Levy et al., 2015; Y. Li et al.,
2018). This barcoding system allows evolving hundreds of thousands of genetically identical yeast
lineages together in a single flask. Each lineage is tagged with a unique DNA barcode, which is a 26 base
pair sequence of DNA located within an artificial intron. Lineages with unique barcodes can be thought of
as independent replicates of an evolution experiment. This high-replicate system has the potential to
generate many different yeast lineages that differ by single adaptive mutations (Kinsler et al., 2020;
Venkataram et al., 2016).

We performed a total of 12 barcoded evolution experiments, each of which started from the same pool of
approximately 300,000 barcoded yeast lineages (**Figure 2A – B; Figure S1**). These evolutions survey
how yeast cells adapt to different concentrations and combinations of two drugs: fluconazole (FLU) and
radicicol (RAD). FLU is a first line of defense against yeast infections, but over the past two decades
diverse resistant mutations have been identified (Bongomin et al., 2017; Osset-Trénor et al., 2023; Rybak
et al., 2022). Some earlier work suggested that FLU-resistant mutants are sensitive to the second drug we
study, radicicol (RAD) (Cowen et al., 2009; Cowen and Lindquist, 2005), and more generally that RAD
can prevent the emergence of drug resistance in other systems (Whitesell et al., 2014). However, there are
some mutants that are cross resistant to both FLU and RAD (Hill et al., 2015), and the prominent
mechanism of resistance can differ with the intensity of selection and drug concentration (Cowen and
Lindquist, 2005; Yang et al., 2023). We thus chose to evolve yeast to resist different concentrations and
combinations of FLU and RAD to generate a diverse pool of adaptive mutations comprising different
mechanisms of drug resistance.

208
209
Figure 2: An overview of the experimental design. (A) Yeast cells were barcoded to create 300,000 lineages. **(B)**
These lineages were evolved in 12 different conditions (**Table 1**). **(C)** A small sample of evolved isolates were taken
from each evolution experiment and their barcodes were sequenced. These ~21,000 isolates do not represent as many
unique, adaptive lineages because many either have the same barcode or do not possess adaptive mutations. **(D)** These
samples of evolved isolates were all pooled together with control strains representing the ancestral genotype. **(E)**
Barcoded fitness competition experiments were then performed on this pool in each of the 12 evolution conditions.
Fitness was measured by tracking changes in each barcode's frequency over time relative to control strains. Two
replicates per condition were performed. **(F)** The overall goal is to investigate fitness tradeoffs for hundreds of
adaptive lineages. For example, the adaptive lineage depicted in dark blue has higher fitness than the ancestor in some
environments (HR, HF) but lower fitness in others (DMSO, ND). We were able to investigate fitness tradeoffs for
774 adaptive lineages. We excluded lineages when we did not observe their associated barcode at least 500 times in
all 12 environments. In other words, we only included lineages for which we obtained high quality fitness estimates
in all 12 environments.

We evolved yeast to resist three different concentrations of either FLU or RAD for a total of six single-drug conditions (**Table 1**). We also studied four conditions containing combinations of both drugs, as well as two control conditions, for a total of 12 evolution experiments (**Table 1**). We chose to study subclinical drug concentrations with the hope that no drug treatment would be strong enough to reduce the population of yeast cells to only a handful of unique barcodes (**Figure S2**). We needed to maintain barcode diversity in order to evolve a large number of unique lineages that each accumulate different mutations.

Evolution Condition	Abbreviation	SYMBOL
4 µg/ml Fluconazole	Low Flu	Low Flu
6 µg/ml Fluconazole	Med Flu	Med Flu
8 µg/ml Fluconazole	High Flu	High Flu
5 µM Radicicol	Low Rad	Low Rad
15 µM Radicicol	Med Rad	Med Rad
20 µM Radicicol	High Rad	High Rad
5 µM Radicicol + 4 µg/ml Fluconazole	LRLF	LRLF
5 µM Radicicol + 8 µg/ml Fluconazole	LRHF	LRHF
15 µM Radicicol + 4 µg/ml Fluconazole	HRLF	HRLF
15 µM Radicicol +8 µg/ml Fluconazole	HRHF	HRHF
0.5% DMSO	DMSO	DMSO
No Drug	No Drug	No Drug

Table 1: A list of the environments included in this study and the symbol used to represent them in subsequent figures.

With the goal of collecting adaptive lineages from each evolution experiment, we took samples from each one after 3 - 6 growth/transfer cycles (**Figure S1**). This represents roughly 24 to 48 generations of growth assuming 8 generations per growth/transfer cycle (Levy et al., 2015). We sampled early because previous work using this barcoded evolution system demonstrated that the diversity of adaptive lineages peaks after just a few dozen generations (Levy et al., 2015; Venkataram et al., 2016). This happens because the barcoding process is slightly mutagenic, thus there is less need to wait for DNA replication errors to introduce mutations (Levy et al., 2015; Venkataram et al., 2016). We sampled about 2,000 cells from each evolution experiment except those 3 containing high concentrations of FLU from which we sampled only 1,000 cells for a total of ~21,000 isolates (2,000 cells x 9 conditions + 1000 cells x 3 conditions) (**Figure 2C**).

Next, we measured the fitness of each isolate in each of the 12 evolution environments to quantify fitness tradeoffs (e.g., whether mutants that perform well in one environment perform worse in another). This process also indirectly screens isolates for adaptive mutations by comparing the fitness of each evolved

isolate to the ancestor of the evolution experiments (Venkataram et al., 2016). To do so, we pooled these
21,000 isolates and used this pool to initiate fitness competition experiments (**Figure 2D**). We competed
the pool against control strains, i.e., strains of the ancestral genotype that do not possess adaptive
mutations (Kinsler et al., 2020; Venkataram et al., 2016). We performed 24 such competitive fitness
experiments, 2 per each of the original 12 evolution conditions (**Figure 2E**). In each experiment, we
emulated the growth and transfer conditions of the original evolution experiments as precisely as possible,
tracking how barcode frequencies changed over 5 growth/transfer cycles (~40 generations). We used the
log-linear slope of this change, relative to the average slope for the control strains, to quantify relative
fitness.

We found that many barcoded lineages have higher fitness than the control strains in some conditions,
presumably because they possess adaptive mutations that improve their fitness in some conditions
(**Figure S3**). In fact, some of these adaptive lineages outcompeted the other lineages so quickly that it
posed a challenge. Barcodes pertaining to outcompeted lineages were often not present at high enough
coverage to track their fitness. We applied a conservative filter, preserving only 774 lineages with
barcodes that were observed >500 times in at least one replicate experiment per each of the 12 conditions.
The reason we required fitness measurements in all 12 conditions is that our goal is to examine each
lineage's fitness tradeoffs (**Figure 2F**) in order to see if different lineages have different tradeoffs. In
order to compare apples to apples, we need to measure each lineage's fitness in the same set of
environments.

The 774 lineages we focus on are biased towards those that are reproducibly adaptive in multiple
environments we study. This is because lineages that have low fitness in a particular environment are
rarely observed >500 times in that environment (**Figure S4**). By requiring lineages to have high-coverage
fitness measurements in all 12 conditions, we exclude adaptive mutants that have severe tradeoffs in one
or more environments, consequently blinding ourselves to mutants that act via unique underlying
mechanisms. Despite this bias, we will go on to demonstrate that there are different types of mutants with
characteristically different fitness tradeoffs present among the 774 lineages that remain.

To provide additional evidence that these 774 barcoded yeast lineages indeed possess adaptive mutations,
we performed whole genome sequencing on a subset of 53 lineages. Because we sampled these lineages
after only a few dozen generations of evolution, each lineage differs from the ancestor by one or just a
few mutations, making it easy to pinpoint the genetic basis of adaptation. Doing so revealed mutations
that have previously been shown to be adaptive in our evolution conditions (**Table S1**). For example, we
sequenced many FLU-resistant yeast lineages finding 35 with unique single nucleotide mutations in either
PDR1 or PDR3, and a few with mutations in SUR1 or UPC2, genes which have all been shown to
contribute to FLU resistance in previous work (Flowers et al., 2012; Tanaka and Tani, 2018; Uemura and
Moriguchi, 2022; Vasicek et al., 2014; Vu and Moye-Rowley, 2022). Similarly, lineages that have very
high fitness in RAD were found to possess single nucleotide mutations in genes associated with RAD
resistance, such as HDA1 (Robbins et al., 2012) and HSC82, which is the target of RAD (Roe et al.,
1999). We also observed several lineages with similar mutations to those observed in other studies using
this barcoded evolution regime, including mutations to IRA1, IRA2 and GPB2 (Kinsler et al., 2020;
Venkataram et al., 2016). Previous barcoded evolutions also observed that increases in ploidy were
adaptive, with 43% to 60% of cells becoming diploid during the course of evolution (Venkataram et al.,

2016). However, ploidy changes contributed less to adaptation in our experiment, with at most 9.4% of
cells becoming diploid by the time point when we sampled, but often less than 2% (**Table S2**).
303
304   In sum, we have created a diverse pool of 774 barcoded yeast lineages, most of which have a fitness
305   advantage in at least one of the conditions we study and are likely to possess a unique adaptive mutation.
306   The question we address for the rest of this study is to what extent these hundreds of mutant lineages
307   differ from one another in terms of their fitness tradeoffs and the mechanism/s underlying their fitness
308   advantages.
309
A unique mechanisms of FLU resistance emerges among mutants isolated in RAD evolutions
The majority of the 774 adaptive lineages that we study have higher fitness than the ancestral strains in
not one, but often in several drug conditions. This suggests that pleiotropy, and in particular cross-
resistance, is prevalent among the lineages we study. But not all lineages show the same patterns of cross
resistance (**Figure 3**). For example, the 100 most fit lineages in our highest concentration of fluconazole
are also beneficial in our highest concentration of radicicol (**Figure 3A**; leftmost two boxplots). As
expected, these 100 lineages also have high fitness in conditions where high concentrations of FLU and
RAD are combined (**Figure 3A**; third boxplot). And these 100 most-fit lineages in FLU lose their fitness
advantage in conditions where no drug is present (**Figure 3A**; rightmost boxplot).
319
Given their high fitness in conditions containing FLU, it seems likely that these 100 mutants originated
from evolution experiments containing FLU. We can trace every lineage back to the evolution
experiment/s it originated from because we sequenced the lineages we sampled from each evolution
experiment before pooling all 21,000 isolates (**Figure 2C**). As we expected, these 100 best performing
lineages in high FLU largely originate from evolution experiments containing FLU (**Figure 3B**). Given
that these lineages have no fitness advantage in conditions containing no drug, it is also unsurprising that
they are underrepresented in evolution experiments lacking RAD and FLU (**Figure 3B**).
327
It might be tempting to generalize that most mutations that provide drug resistance are not beneficial in
environments without drugs. Afterall, we show this is true for 100 independent lineages (**Figure 3A**).
Further, many previous studies find a similar pattern, whereby drug resistant mutants often do not have
high fitness in the absence of drug (Allen et al., 2019; Andersson and Hughes, 2010; Basra et al., 2018;
Melnikov et al., 2020), such that treatment strategies have emerged that cycle patients between drug and
no drug states, albeit with mixed success (Algazi et al., 2020; Baker et al., 2018; Raymond, 2019; Wang
et al., 2019). However, this type of generalization is not supported by our data. We find that drug
resistance can sometimes come with an advantage, rather than a cost, in the absence of a drug (**Figure
3C**). The top 100 most fit mutants in our highest concentration of RAD provide a fitness advantage in
high RAD, high FLU, as well as in environments with no drug (**Figure 3C**). These observations suggest
that there are at least two different mechanisms by which to resist FLU that result in different tradeoffs in
other environments (**Figure 3A vs. 3C**).
340
341   Intriguingly, these FLU-resistant lineages that maintain their fitness advantage in the absence of drug
342   (**Figure 3C**) mainly originate from evolution experiments performed in conditions lacking FLU (**Figure
343   3D**). This highlights how the potential mechanisms by which a microbe can resist a drug may be more

344 varied than is often believed. Typically, one doesn't search for FLU-resistant mutants by evolving yeast
345 to resist RAD. Thus typical studies might miss this unique class of FLU-resistant mutants.
346
In sum, there appear to be at least two different types of mutants present among our collection of 774
adaptive yeast lineages. One group has almost equally high fitness in RAD and FLU but has no fitness
advantage over the ancestral strain in conditions without either drug (**Figure 3A**). Another group is
defined by very high fitness in RAD, moderately high fitness in FLU and moderately high fitness in
conditions without either drug (**Figure 3C**). When comparing fitness in RAD vs. FLU across all 774
lineages, not only the top 100 best performing in each drug, we see some evidence that they largely fall
into the two main categories highlighted in **figures 3A and 3C** (**Figure 3E**). Thus it might be tempting to
conclude that there are two different types of FLU-resistant mutant in our dataset. However, sorting
mutants into groups using a pairwise correlation plot (**Figure 3E**) excludes data from ten of our twelve
environments.

Figure 3: Two different classes of FLU-resistant mutants with unique tradeoffs. (A) This panel describes the 100 mutant lineages with the highest fitness relative to the control strains in the high FLU environment (8μg/ml FLU). The vertical axis depicts the fitnesses (log-linear slopes relative to control strains) for these 100 strains in four selected environments, including the high FLU environment (boxed). Boxplots summarize the distribution across all 100 lineages for each environment, displaying the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR (whiskers). **(B)** The 100 lineages with highest fitness in high FLU were most often sampled from evolution experiments in which FLU was present. In this pie-chart, colors correspond to the evolution conditions listed in **table 1** and the blue outer ring highlights evolution conditions that contain FLU. The size of each slice of pie represents the relative frequency with which these 100 lineages were found in each evolution experiment. **(C)** Similar to panel A, this panel describes the 100 mutant lineages with the highest fitness relative to the control strains in the high RAD environment (20uM Rad). **(D)** The 100 lineages with highest fitness in high RAD were most often sampled from evolution experiments that did not contain FLU. **(E)** A pairwise correlation plot showing that all 774 mutants, not just the two groups of 100 depicted in panels A and C, to some extent fall into two groups defined by their fitness in high FLU and high RAD. The contours (black curves) were generated using kernel density estimation with bins = 7. These contours describe the density of the underlying data, which is concentrated into two clusters defined by the two smallest black circles. The 100 mutants with highest fitness in high FLU are blue, highest fitness in high RAD are red, and the seven that overlap between the two aforementioned categories are black.

A strategy to differentiate classes of drug-resistant mutants with different tradeoffs

The observation of two distinct types of adaptive mutants (**Figure 3**) made us wonder whether there were additional unique types of FLU-resistant mutants with their own characteristic tradeoffs. This is difficult to tell by using pairwise correlation like that in **Figure 3E** because we are not studying pairs of conditions, as is somewhat common when looking for tradeoffs to leverage in multidrug therapies (Ardell and Kryazhimskiy, 2021; Larkins-Ford et al., 2022; Melnikov et al., 2020; Scarborough et al., 2020). Instead, we have collected fitness data from 12 conditions to yield a more comprehensive set of gene-by-environment interactions for each mutant. This type of data, describing how a particular genotype responds to environmental change, is sometimes called a 'reaction norm' and can inform quantitative genetic models of how selection operates in fluctuating environments (Gomulkiewicz and Kirkpatrick, 1992; Ogbunugafor, 2022) and how much pleiotropy exists in nature (Yadav et al., 2015). More recent

studies refer to the changing performance of a genotype across environments as a 'fitness profile' or in
aggregate, a 'fitness seascape,' and suggest these type of dynamic measurements are the key to designing
effective multi-drug treatments (King et al., 2022) and to predicting evolution (Cairns et al., 2022; Chen
et al., 2023; Iwasawa et al., 2022; Kinsler et al., 2020; Lässig et al., 2017). And when the environments
studied represent different drugs, these types of data are often referred to as "collateral sensitivity
profiles" a term chosen to convey how resistance to one drug can have "collateral" effects on performance
in other drugs (Gjini and Wood, 2021; Maltas and Wood, 2019; Pál et al., 2015). Despite the wide interest
in this type of fitness data, it is technically challenging to generate, thus many previous studies of fitness
profiles focus on a much smaller number of isolates (Imamovic et al., 2018; Maltas and Wood, 2019;
Nichol et al., 2019), sometimes with variation restricted to a single gene (King et al., 2022; Mira et al.,
2015), or evolved in response to a single selection pressure (Kinsler et al., 2020; Y. Li et al., 2018). Here,
we have generated fitness profiles for a large and diverse group of drug-resistant strains using the power
of DNA barcodes. Now we seek to understand whether these mutants fall into distinct classes that each
have characteristic fitness profiles (i.e., characteristic tradeoffs, characteristic reaction norms, or
characteristic gene-by-environment interactions).
410
To address this question, we start by performing dimensional reduction, clustering mutants with fitness
profiles that have a similar shape. It is in theory possible for all mutants to have similar profiles, perhaps
implying they all affect fitness through similar underlying mechanisms (**Figure 4A**). However, the
disparity reported in **Figure 3** suggests otherwise. It's also possible that every mutant will have a different
profile. This could happen if each mutant affects different molecular-level phenotypes that underlie its
drug resistance (**Figure 4B**). But previous work suggests that the phenotypic basis of adaptation is less
diverse than the genotypic basis (Brettner et al., 2024; Iwasawa et al., 2022; Kinsler et al., 2020). A final
possibility, somewhere in between the first two, is that there exist multiple classes of drug-resistant
mutants each with characteristic tradeoffs (**Figure 4C**). This might imply that each class of mutants
provides drug resistance via a different molecular mechanism, or a different set of mechanisms. Overall,
our endeavor to enumerate how many distinct fitness profiles are present across these 774 mutants
(**Figures 4A – C**) informs general questions about the extent of pleiotropy in the genotype-phenotype-
fitness map (Bakerlee et al., 2021; Boyle et al., 2017; Chen et al., 2023; Geiler-Samerotte et al., 2020), the
extent to which fitness tradeoffs are universal (Andersson and Hughes, 2010; Herren and Baym, 2022; Li
et al., 2019), and relatedly, the extent to which evolution is predictable (Iram et al., 2020; King et al.,
2022; Kinsler et al., 2020; Lässig et al., 2017; Petti et al., 2023).
427
To see whether there are distinct fitness profiles present among our drug-resistant yeast lineages, we
applied uniform manifold approximation and projection (UMAP) (McInnes et al., 2018a) to fitness
measurements for 774 yeast strains across all 12 environments. This method places mutants with similar
fitness profiles near each other in two-dimensional space. As might be expected, it largely places mutants
in each of the two categories described in **Figure 3** far apart, with drug-resistant mutants that lose their
benefit in the absence of drug in the top half of the graph, and those that maintain their benefit in the
bottom half (**Figure 4D and Figure S5**).
435
Beyond the obvious divide between the top and bottom clusters of mutants on the UMAP, we used a
gaussian mixture model (GMM) (Fraley and Raftery, 2003) to identify clusters. A common problem in
this type of analysis is the risk of dividing the data into clusters based on variation that represents

measurement noise rather than reproducible differences between mutants (Mirkin, 2011; Zhao et al.,
2008). One way we avoided this was by using a GMM quality control metric (BIC score) to establish how
splitting out additional clusters affected model performance (**Figure S6**). Another factor we considered
were follow-up genotyping and phenotyping studies that demonstrate biologically meaningful differences
between mutants in different clusters (**Figures 5 – 8**). Using this information, we identified seven clusters
of distinct mutants, including one pertaining to the control strains, and six others pertaining to presumed
different classes of adaptive mutant (**Figure 4D**). It is possible that there exist additional clusters, beyond
those we are able to tease apart in this study.

Preliminarily, we investigated whether the clusters we identified capture reproducible differences between
mutants, rather than measurement noise, by reducing the amount of noise in our data and asking if the
same clusters are still present. To do so, we reduced our collection of adaptive lineages from 774 to 617
by requiring 5,000 rather than 500 reads per lineage per experiment in order to infer fitness. This
procedure reduced noise; the Pearson correlation across replicate experiments improved from 0.756% to
0.813%. Despite this reduction in variation, these 617 lineages cluster into the same six groups (plus a
seventh pertaining to the control strains) as do the original 774 (**Figure S7**). The groupings are also
preserved when we perform alternate methods for dimensionality reduction (**Figures S8 and S9**).

Each of the six clusters of adaptive mutants has a characteristic fitness profile (**Figure 4E**). In any given
environment, the fitnesses of the mutants within each cluster are often very similar to one another and
often significantly different from other clusters (**Figure 4E**). Our follow-up investigations (**Figures 5 –**
8), including whole genome sequencing, growth rate measurements, and tracing the evolutionary origins
of the mutants in each cluster, provide additional evidence that the adaptive mutants in each cluster have
characteristically different properties.

466
467
Figure 4: Clustering evolved lineages with similar fitness profiles. (A-C) Simulated data showing potential
fitness profiles when (**A**) all mutants have similar responses to environmental change and thus a similar fitness
profile, (**B**) every mutant has a different profile (five unique profiles are highlighted in color), or (**C**) every mutant
has one of a small number of unique profiles (two unique profiles are depicted). (**D**) Every point in this plot
represents one of the barcoded lineages colored by cluster; clusters were identified using a gaussian mixture model.
The 774 adaptive lineages cluster into 6 groups based on variation in their fitness profiles; the control lineages
cluster separately into the leftmost cluster in light green. (**E**) The fitness profiles of each cluster of adaptive lineages.
Boxplots summarize the distribution across all lineages within each cluster in each environment, displaying the
median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR
(whiskers).
478
A group of mutants with distinct genotypes are primarily resistant to low concentrations of FLU
The upper three clusters of mutants on the UMAP (**Figure 4D**) are all similar in that they have elevated
fitness in at least one FLU-containing environment but ancestor-like fitness in the absence of drug
(**Figure 4E**; upper three profiles). Despite these similarities, there are major differences between these
three groups of mutant lineages, both at the level of genotype and fitness profile (**Figure 5**). For example,

in cluster 1 (depicted in purple in **Figures 4 and 5**), the 3 sequenced lineages have single nucleotide
mutations to either SUR1 or UPC2 (**Figure 5A**). But in clusters 2 and 3 (depicted in blue and orange in
Figures 4 and 5), 35/36 sequenced lineages have unique single nucleotide mutations to one of two genes
associated with 'Pleiotropic Drug Resistance' (PDR1 or PDR3).
488
PDR1 and PDR3 are transcription factors that are well known to contribute to fluconazole resistance
through increased transcription of a drug pump (PDR5) that removes FLU from cells (Fardeau et al.,
2007; Osset-Trénor et al., 2023). However, SUR1 and UPC2 are less commonly mentioned in literature
pertaining to FLU resistance, and have different functions within the cell as compared to PDR1 and PDR3
(Hill et al., 2015; Kapitzky et al., 2010). SUR1 converts inositol phosphorylceramide to mannosylinositol
phosphorylceramide, which is a component of the plasma membrane (Uemura and Moriguchi, 2022).
Similarly, UPC2 is a transcription factor with a key role in activating the ergosterol biosynthesis genes,
which contribute to membrane formation (Tan et al., 2022; Vik and Rine, 2001). The presence of adaptive
mutations in genes involved in membrane synthesis is consistent with fluconazole's disruptive effect on
membranes (Sorgo et al., 2011).
499
Interestingly, the lineages with mutations to UPC2 and SUR1, and the unsequenced lineages in the same
cluster, do not consistently have cross resistance in RAD (**Figure 5B**; cluster 1). Oppositely, lineages
with mutations to PDR1 or PDR3, and the unsequenced lineages in the same clusters, are uniformly cross
resistant to RAD (**Figure 5B**; clusters 2 and 3). Perhaps this cross resistance is reflective of the fact that
the drug efflux pump that PDR1/3 regulates (PDR5) can transport a wide range of drugs and molecules
out of yeast cells (Harris et al., 2021; Kolaczkowski et al., 1996). Overall, the targets of adaptation in
cluster 1 have disparate functions within the cell as compared to the targets of adaptation in clusters 2 and
3. This may suggest that the mutants in cluster 1 confer FLU resistance via a different mechanism than
clusters 2 and 3.
509
The lineages in cluster 1 have additional important differences from clusters 2 and 3. The lineages in
cluster 1 perform best in the lowest concentration of FLU and have decreasing fitness as the concentration
of FLU rises (**Figure 5C**). In fact, about 15% of these mutant lineages perform worse than their ancestor
in the highest concentration of FLU, suggesting the very mutations that provide resistance to low FLU are
costly in higher concentrations of the same drug. The mutants in clusters 2 and 3 show the opposite trend
from those in cluster 1. They perform best in the highest concentration of FLU and have reduced fitness
in lower concentrations (**Figure 5C**). These findings provide additional evidence that a distinct
mechanism of FLU resistance distinguishes cluster 1 from clusters 2 and 3.
518
To confirm that different drug-resistant mutants dominate evolution in just slightly different
concentrations of the same drug, we used each cluster's barcodes to trace mutants back to the evolution
experiment from which they originated. Mutants in cluster 1 predominantly originated in evolutions
containing the lowest concentration of FLU (**Figure 5A**; pie charts), while mutants in clusters 2 and 3
more often originated from evolution experiments containing higher FLU concentrations (see **Figure 6**).
We also confirmed that the mutants in cluster 1 have high fitness in low FLU only by measuring growth
curves for SUR1, UPC2 and PDR mutants in 3 concentrations of fluconazole. Lineages from cluster 1
(SUR1, UPC2) indeed grow faster and reach higher density in low FLU than those from cluster 3 (PDR)
(**Figure 5D**). But lineages from cluster 1 (SUR1, UPC2) grow poorly in higher FLU concentrations,

while lineages from cluster 3 (PDR) do not suffer this tradeoff (**Figure 5E**). The observation that different
mutants acting via different resistance mechanisms dominate evolution in only slightly different
concentrations of the same drug highlights the complexity of adaptation and the potential benefits of more
deeply understanding the diversity of adaptive mechanisms before designing treatment strategies (Berman
and Krysan, 2020; Yang et al., 2023).

533
534

535
536
Figure 5: Evolved lineages comprising cluster 1 have different genotypes and phenotypes from neighboring
clusters. (**A**) The three clusters on the top half of the UMAP differ in their genetic targets of adaptation with cluster
1 being unique in that it does not contain mutations to PDR1 or PDR3. Cluster 1 is also unique in that it contains
lineages that predominantly originated from the low fluconazole evolution condition; the pie chart depicts the
fraction of lineages originating from each of the 12 evolution environments with colors corresponding to **Table 1**.
(**B**) Evolved lineages comprising cluster 1 do not have consistent fitness advantages in conditions containing RAD,
while lineages comprising clusters 2 and 3 are uniformly adaptive in medium and high RAD. Boxplots summarize
the distribution across all lineages within each cluster in each environment, displaying the median (center line),
interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR (whiskers). (**C**) Lineages
comprising cluster 1 are most fit in low concentrations of FLU, and this advantage dwindles as the FLU
concentration increases. Lineages comprising clusters 2 and 3 show the opposite trend. (**D**) In low FLU (4 μg/ml),
Cluster 1 lineages (UPC2 and SUR1) grow faster and achieve higher density than lineages from cluster 3 (PDR).
This is consistent with bar-seq measurements demonstrating that cluster 1 mutants have the highest fitness in low
FLU. (**D**) Cluster 1 lineages are sensitive to increasing FLU concentrations (SUR1 and UPC2). This is apparent in

that the dark blue (8 μg/ml flu) and grey (32 μg/ml flu) growth curves rise more slowly and reach lower density than
the light blue curves (4 μg/ml flu). But this is not the case for the PDR mutants. These observations are consistent
with the bar-seq fitness data (**Fig 4E**).
554
Two groups of mutant lineages possessing similar adaptive mutations differ in sensitivity to RAD
While cluster 1 appears fairly different from its neighbors, it is not immediately obvious why the mutant
lineages in clusters 2 and 3 are placed into separate groups. For one, the mutants in clusters 2 and 3 have
fitness profiles with a very similar shape (**Figure 4E & 6A**). The sequenced lineages in each of these
clusters also possess mutations to the same genes: PDR1 and PDR3 (**Figure 5A**). And finally, the
lineages in each cluster originate from similar evolution experiments, largely those containing FLU
(**Figure 6A**; pie charts). These observations made us wonder whether the difference between cluster 2 and
3 arose entirely because the mutants in cluster 3 have stronger effects than those in cluster 2 (**Figure 6A**;
the solid blue line is above the solid orange line). In other words, we wondered whether the mutant
lineages in clusters 2 and 3 affect fitness via the same mechanism, but to different degrees. To investigate
this idea, we normalized all fitness profiles to have the same height on the vertical axis; this does not
affect their shape (**Figure 6A**; dotted lines). Then we re-clustered and asked whether mutants pertaining
to the original clusters 2 and 3 were now merged into a single cluster. They were not (**Figure 6B**).
568
569

Figure 6: Evolved lineages in clusters 2 and 3 have characteristic differences despite similarities at the genetic level. (**A**) This panel shows the similarities between clusters 2 and 3. The upper right inset displays the same UMAP from **Figure 4D** with only clusters 2 and 3 highlighted and with lineages possessing mutations to the PDR genes depicted as blue diamonds. The line plot displays the same fitness profiles for clusters 2 and 3 as **Figure 4E**, plotting the average fitness for each cluster in each environment and a 95% confidence interval. Dotted lines represent the same data, normalized such that every lineage has an average fitness of 0 across all environments. These line plots show that the fitness profiles for clusters 2 and 3 have a very similar shape. Pie charts display the relative frequency with which lineages in clusters 2 and 3 were sampled from each of the 12 evolution conditions, colors match those in the horizontal axis of the line plot and **table 1**. (**B**) This panel shows the differences between the new clusters 2 and 3 created after all fitness profiles were normalized to eliminate magnitude differences. The upper right inset displays a new UMAP (also see **Figure S10**) that summarizes variation in fitness profiles after each profile was normalized by setting its average fitness to 0. The line plot displays the fitness profiles for the new clusters 2 and 3, which look different from those in panel A because 37% of mutants in the original clusters 2 and 3 switched identity from 2 to 3 or vice versa. The new clusters 2 and 3 are depicted in slightly different shades of blue and orange to reflect that these are not the same groupings as those depicted in **Figure 4**. Pie charts display the relative frequency

586 with which lineages in new clusters 2 and 3 were sampled from each of the 12 evolution conditions, colors match
587 those in the horizontal axis of the line plot and **table 1**.
588
Normalizing in this way did not radically alter the UMAP, which still contains largely the same 6 clusters
of mutants (**Figure S10**). Clusters 2 and 3, containing lineages with mutations to PDR1 or PDR 3,
experienced the largest changes with 37% of mutants switching from one of these two groups to the other.
The new clusters 2 and 3 now differ in the shape of their fitness profiles, whereby slight differences that
existed between the original fitness profiles are exaggerated (**Figure 6B**). For example, mutants in cluster
3 perform better in high and medium concentrations of RAD (**Figure 6B**; line plot). This difference in
fitness is reflected in the evolution experiments, with more mutant lineages in cluster 3 originating from
the evolutions performed in RAD (**Figure 6B**; pie charts). Though cluster 3 mutants tend to have stronger
RAD resistance, they tend to have reduced fitness in conditions containing neither FLU nor RAD as
compared to cluster 2 lineages (**Figure 6B**; line plot). In sum, the differences between lineages in clusters
2 and 3 were not resolved upon normalizing fitness profiles to reduce magnitude differences, instead they
were made more apparent (**Figure 6**). These differences do not appear to be random because they persist
across independent experiments. For example, cluster 3 mutants are more fit in both medium and high
RAD environments (**Figure 6B**; line plot) and were more often isolated from evolutions containing RAD
(**Figure 6B**; pie charts). The observation that PDR mutations fall into two separate clusters begs a
question: how can different mutations to the same gene affect fitness via different molecular mechanisms?
605
Asking this question forces us to consider what we mean by "mechanism." The mechanism by which
mutations to PDR1 and PDR3 affect FLU resistance is well established: they increase transcription of an
efflux pump that removes FLU from cells (Buechel and Pinkett, 2020; Moye-Rowley, 2019; Osset-Trénor
et al., 2023). But if this is the only molecular-level effect of mutations to these genes, it is difficult to
reconcile why PDR mutants fall into two distinct clusters with differently shaped fitness profiles. Others
have also recently observed that mutants to PDR1 do not all behave the same way when exposed to novel
drugs or changes in pH (Chen et al., 2023). This phenomenon is not reserved to PDR mutants, as adaptive
missense mutations to another gene, IRA1, also do not share similarly shaped fitness profiles either
(Kinsler et al., 2020). One explanation may be that, while all adaptive mutations within the same gene
improve fitness via the same mechanism, not all mutants suffer the same costs. For example, perhaps the
adaptive PDR mutations in cluster 2 cause misfolding of the PDR protein, resulting in lower fitness in
RAD because this drug inhibits a chaperone that helps proteins to fold. In this case, it might be more
correct to say that each of our six clusters affects fitness through a different, but potentially overlapping,
suite of mechanisms (Wang et al., 2023). Previous work demonstrating that mutations commonly affect
multiple traits supports this broader view of the mechanistic differences between clusters (Boyle et al.,
2017; Geiler-Samerotte et al., 2020; Kinsler et al., 2020; Paaby and Rockman, 2013).
622
Alternatively, perhaps not all adaptive mutations to PDR improve fitness via the same mechanism. PDR1
and PDR3 regulate transcription of YOR1 and SNQ2 as well as PDR5, and maybe the different clusters
we observe represent mutants that upregulate one of these downstream targets more than the other (Osset-
Trénor et al., 2023). Or, the mutants in each cluster might harbor different aneuploidies or small, difficult
to sequence chromosomal insertions or deletions that affect fitness. We leave identification of the precise
mechanisms that differentiate these clusters for future work. Here, using the example of PDR mutants, we

showcase how genotype may not predict fitness tradeoffs, suggesting there is more to learn about the
mechanisms underlying FLU resistance.
631
One group of RAD resistant mutants does not respond as expected to drug combinations
Though the three clusters of mutants on the bottom half of the UMAP are all advantageous in RAD and in
conditions without any drug (**Figure 4E**; lower three plots), they differ in their fitness in conditions
containing FLU. For example, the cluster of yeast lineages highlighted in green (cluster 4 in **Figures 4**
and 7A) is unique in that it has a slight advantage in the HRLF environment (**Figure 7B**). We found it
especially strange that the neighboring cluster 5 does not also have a fitness advantage in this condition.
Mutants in cluster 5 have a slight advantage in the LF condition, and a big advantage in the HR condition,
thus we expect them to have at least some fitness advantage in the condition where these two drugs are
combined (HRLF), but they do not (**Figure 7B**). The same is true for the combination of LRLF: cluster 5
mutants have an advantage in both single drug conditions which is lost when the drugs are combined
(**Figure S11**). However, the mutants in cluster 4 (green) exhibit no such sensitivity to combined
treatment. They have a slight advantage in all of the aforementioned single drug conditions, which is
preserved in the relevant multidrug conditions (**Figure 7B & Figure S11**). To obtain an independent
measure of the fitness of cluster 4 vs. cluster 5 lineages in these multidrug conditions, we asked from
where the lineages in each cluster originate. About 10% of cluster 4 lineages originated from the HRLF
evolution, while almost none of the lineages in cluster 5 came from this experiment, confirming that
cluster 5 lineages are uniquely sensitive to this multidrug environment (**Figure 7C**).
649
The different fitness profiles of mutants in cluster 4 versus 5 (**Figure 7B & Figure S11**) might imply that
they have different growth phenotypes. We performed a follow-up experiment that supports this
observation. We asked whether there are differences in the growth phenotypes of cluster 4 versus 5
mutants by measuring a growth curve for the lineage we were able to isolate and sequence from cluster 5,
comparing it to a growth curve from a cluster 4 lineage (**Figure 7D**). Indeed, the selected mutants in
cluster 4 and 5 appear to have markedly different growth curves in some conditions (**Figure S11**). The
growth differences echo those we see in the fitness data. For example, the cluster 5 mutant has a lower
maximum growth rate in the HRLF multidrug condition, corresponding with their lower fitness in this
condition (**Figure 7B & 7D**). The cluster 5 mutant also reaches a lower maximum cell density in the
LRLF multidrug condition and also has lower fitness in this condition (**Figure S11**). However, the growth
curves and fitnesses of cluster 4 and 5 mutants are more similar in LF, LR and HR single drug conditions.
The observation of reproducible growth differences between a cluster 4 and cluster 5 mutant provides
some supporting evidence that our clustering approach is effective at separating mutants with different
properties.

Figure 7: Evolved lineages in cluster 4 and 5 differ in response to combined drugs. (A) Adjacent clusters 4 and 5 each contain a small number of sequenced isolates depicted as diamonds; diamond colors correspond to the genes containing adaptive mutations in each sequenced isolate. **(B)** Cluster 5 (red) has an unexpected fitness disadvantage in the HRLF multidrug environment relative to cluster 4 (green), given that cluster 5 lineages do not have a fitness disadvantage in the relevant single drug environments. Boxplots summarize the distribution across all lineages within each cluster in each environment, displaying the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR (whiskers). **(C)** Pie charts display the relative frequency with which lineages in each cluster were sampled from each of the 12 evolution conditions, colors match those in table 1. **(D)** The maximum exponential growth rate for a single lineage isolated from each of clusters 4 (green) and 5 (red), relative to the ancestor. The growth rate of each lineage in each condition was measured twice by measuring changes in optical density over time. Tested lineage from cluster 4 (in green) has a mutation to GBP2 (S317T) while the lineage from cluster 5 (in red) has mutation to HDA1 (S600S).

One group of RAD resistant mutants is exceptionally adaptive in conditions without drug

One group of mutants in the lower half of the UMAP (cluster 6 in **Figure 8A**) appears distinct from the other two in that it has the largest fitness advantage in conditions lacking any drug (**Figure 8B**). This might imply that cluster 6 lineages rose to high frequency during our evolution experiments in environments without either drug, specifically the "no drug" and "DMSO" control conditions. Indeed, this is what we observe: over 50% of the lineages in cluster 6 were sampled from one of these two evolution experiments (**Figure 8A**). On the contrary, the other clusters in the lower half of the UMAP consist mainly of lineages sampled from one of the RAD evolutions (**Figure 7C**). Since our fitness experiments were performed independently of the evolution experiments, this provides two independent pieces of evidence suggesting that lineages in cluster 6 are defined by their superior performance in conditions lacking any drug.

In line with the success of cluster 6 mutants in no drug conditions, the five sequenced mutants in this
cluster include three that have mutations to IRA1, which was the most common target of adaptation in
another evolution experiment in the condition we call "no drug" (**Figure 8A**) (Venkataram et al., 2016).
In that experiment, and in our no drug experiment, mutations to IRA1 result in a greater fitness advantage
than mutations to its paralog, IRA2, or mutations to other negative regulators of the RAS/PKA pathway
such as GPB2 (**Figure 8**). Previous work showed that sometimes IRA1 mutants have very strong
tradeoffs, for example, they become extremely maladaptive in environments containing salt or benomyl
(Kinsler et al., 2020). We do not observe this to be the case for either FLU or RAD. In fact, we observe
that cluster 6 mutants, including those in IRA1, maintain a fitness advantage in our highest concentration
of both drugs (**Figure 4**), being more fit in high FLU than mutants in either of the other clusters in the
lower half of the UMAP (**Figure 8B**). However, cluster 6 mutants are unique in that they lose their fitness
advantage in the lowest concentration of FLU (**Figure 8B**). Being singularly sensitive to a low
concentration of drug seems unusual, so much so that when this was observed previously for IRA1
mutants the authors added a note about the possibility of a technical error (Kinsler et al., 2020). Our
results suggest that there is indeed something uniquely treacherous about the low fluconazole
environment, at least for some genotypes.

Figure 8: Evolved lineages in cluster 6 have higher fitness than other lineages in the absence of FLU and
RAD. (A) Same UMAP as **Figure 4D** with clusters 4, 5, and 6 highlighted and sequenced isolates in these clusters
represented as diamonds. Diamond colors correspond to the targets of adaptation in the sequenced isolates. Pie
charts display the relative frequency with which lineages in cluster 6 were sampled from each of the 12 evolution
conditions; colors match those in **table 1**. Grey outline depicts conditions lacking RAD and FLU. **(B)** Of the three
clusters on the bottom half of the UMAP, cluster 6 lineages perform best in conditions without any drug and in the
highest concentration of FLU. Yet they perform worst in the lowest concentration of FLU. Boxplots summarize the
distribution across all lineages within each cluster in each environment, displaying the median (center line),
interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR (whiskers).

Discussion:

Here, we present a barcoded collection of fluconazole (FLU) resistant yeast strains that is unique in its size, its diversity, and its tractability. One way we were able to isolate diverse types of FLU-resistance was by evolving yeast to resist diverse drug concentrations and combinations. But the more important tool used to increase both the number and type of mutants in our collection was DNA barcodes. These allowed us to sample beyond the drug resistant mutants that rise to appreciable frequency and to collect mutants that would eventually have been outcompeted by others. Our primary goal in collecting these mutants was to get a rough sense of how many different mechanisms of FLU resistance may exist. This question is relevant to evolutionary medicine (because more mechanisms of resistance make it harder to design strategies to avoid resistance), evolutionary theory (because more mechanisms of adaptation make it harder to predict how evolution will proceed), and genotype-phenotype mapping (because more mechanisms makes for a more complex map).

We distinguish mutants that likely act via different mechanisms by identifying those with different fitness tradeoffs across 12 environments, leveraging the mutants' barcodes to track their relative fitness following previous work (Kinsler et al., 2020). The 774 FLU-resistant mutants studied here clustered into a handful of groups (6) with characteristic tradeoffs. We confirmed that each group captures mutants with distinct properties using multiple approaches, including whole genome sequencing, growth curve experiments, tracing the evolutionary origins of the mutants in each cluster, and by using two additional independent clustering methods: hierarchical clustering and PCA (**Figure 5-8, S7-11**). Some groupings are unintuitive in that they segregate mutations within the same gene (**Figure 6**) or are distinguished by unexpectedly low fitness in multidrug conditions (**Figure 7**). These findings are important because they challenge strategies in evolutionary medicine that rely on consistent tradeoffs or intuitive trends when designing sequential drug treatments. On the other hand, the observation that some mutants have very similar tradeoffs such that they cluster together is promising in that it suggests predicting the impact of some mutations by understanding the impacts of others is somewhat feasible.

Problematically, the clusters we present are incomplete and bound to change as additional data presents itself. For one, we have shown that additional FLU-resistant mutants emerge from evolution experiments in conditions lacking FLU (**Figure 3C & 3D**). This begs questions about what other FLU-resistant mutants might emerge in environments we have not studied here. Additionally, previous work has shown that some mutants that group together in our study (e.g., GPB2 and IRA2) have different fitness profiles in conditions that we did not include here (Kinsler et al., 2020). Also of note is that our evolution experiments were conducted for only a few generations and all started from the same genetic background. Additional types of FLU-resistant mutants with unique fitness profiles may emerge from other genetic backgrounds or arise after more mutations are allowed to accumulate (Allen et al., 2021; Bosch et al., 2021; Brandis et al., 2012). Finally, by requiring that all included mutants have sufficient sequencing coverage in all 12 environments, our study is underpowered to detect adaptive lineages that have low fitness in any of the 12 environments. This is bound to exclude large numbers of adaptive mutants. For example, previous work has shown some FLU resistant mutants have strong tradeoffs in RAD (Cowen and Lindquist, 2005). Perhaps we are unable to detect these mutants because their barcodes are at too low a frequency in RAD environments, thus they are excluded from our collection of 774. All of the aforementioned observations combined suggest that there are more unique types of FLU-resistant mutations than those represented by these 6 clusters, and that the molecular mechanisms that can

766 contribute to fitness in FLU are more diverse than we know. This could complicate (or even make
767 impossible) endeavors to design antimicrobial treatment strategies that thwart resistance.
768
On the up side for evolutionary medicine, not every infection harbors all possible types of mutants. This
might explain why strategies that exploit one or two common tradeoffs have some, albeit mixed, success
in delaying or preventing the emergence of resistance (Amin et al., 2015; Imamovic et al., 2018; Kaiser,
2017; Krishna et al., 2022; Nyhoegen and Uecker, 2023; Waller et al., 2023; Wang et al., 2019). Our
results encourage more complex strategies to thwart drug resistance (Iram et al., 2020), such as those that
focus on advance screening to determine the resistance mechanisms that are present (Andersson et al.,
2019a), or on cycling a larger number of drugs to exploit a larger number of tradeoffs (Thomas et al.,
2022; Yoshida et al., 2017). Problematically, these strategies often rely on knowledge about the diversity
of mutants and tradeoffs that exist (or that can emerge) within an infectious population. This type of
information about population heterogeneity, heteroresistance, and substructure is expensive and arduous
to obtain (Andersson et al., 2019b; Bottery et al., 2021). Fortunately, new methods, in addition to the one
presented in this study, are emerging (Aissa et al., 2021; Brettner et al., 2024; Forsyth et al., 2021; Hsieh
et al., 2022; Kuchina et al., 2021; Nagasawa et al., 2021). The richer data provided by these methods
dovetails with emerging population genetic models that predict the likelihood of resistance to a given
drug regimen (Cannataro et al., 2018; Day et al., 2015; Feder et al., 2021; King et al., 2022; Read and
Huijben, 2009; Somarelli et al., 2020; Wilson et al., 2016). In sum, our observation of numerous different
types of drug-resistant mutations suggests that designing resistance-detering therapies is challenging, but
perhaps not impossible.
787
Outside of predicting the evolution of resistance, our findings provide a tool to investigate the phenotypic
impacts of mutation. This task has proven daunting in light of work demonstrating that mutations often
have many phenotypic impacts (Boyle et al., 2017; Paaby and Rockman, 2013) and that these impacts
change with contexts including the environment (Eguchi et al., 2019; Geiler-Samerotte et al., 2020, 2016;
Lee et al., 2019; Paaby et al., 2015). The approach presented in this study provides a way forward by
identifying mutations that cluster together such that the effects of some mutants can be predicted from
others. This clustering strategy can assist high-throughput efforts to identify the phenotypic impacts of a
large panel of mutations (Flynn et al., 2024; Fowler and Fields, 2014; Mehlhoff et al., 2020; Starr et al.,
2017). Further, our approach identifies environments that differentiate one cluster of mutants from
another. This suggests where to look to understand the phenotypes that differentiate each cluster of
mutants. For example, we were able to show that the growth phenotypes of mutants from clusters 4 and 5
are different because we knew to look for these differences in multidrug environments (**Figure 7D**). And
our results suggest radicicol environments may be most helpful in teasing out any phenotypic differences
that set apart some PDR mutations from others (**Figure 6**). Thus, our approach guides efforts to
understand the phenotypic effects of mutation, while also guiding efforts to predict the effects of some
mutations from others as well as efforts to predict the outcomes of adaptive evolution.
804

Acknowledgements:
We are grateful to Sasha Levy for providing plasmids and yeast strains, to Martin Mullis, Xianan Liu,
Sasha Levy and Gavin Sherlock for advice about creating a high diversity library of barcoded yeast,
Jamie Blundell, Sandeep Venkataram and Fangfei Li for discussions about how to infer adaptive lineages,
and to the members of the Geiler-Samerotte lab as well as Michael Lynch for discussions about the

manuscript. The authors acknowledge resources and support from the KE core facilities at Arizona State
University.
812

813 **Funding:**
814 This work was supported by a National Institutes of Health grant R35GM133674 (to KGS), an Alfred P
815 Sloan Research Fellowship in Computational and Molecular Evolutionary Biology grant FG-2021-15705
816 (to KGS), and a National Science Foundation Biological Integration Institution grant 2119963 (to KGS).

817
818
819 **Methods:**
820
821 **Base yeast strains**
822 All of the budding yeast (*Saccharomyces cerevisiae*) lineages studied here originated from the same starting
823 strain referred to as the "landing pad strain" (SHA185) in previous work (Levy et al., 2015). We transformed
824 a barcode library (generously provided by Sasha Levy) into this strain as described below, creating a strain
825 with the following genetic background:
826
827 MATα, ura3Δ0, ybr209w::Gal-Cre-KanMX-1/2URA3-loxP-Barcode-1/2URA3-HygMX-lox66/71.
828
829 **Base media**
830 All experiments were conducted in "M3" media defined in the same study as the landing pad strain (Levy
831 et al., 2015), which is a glucose-limited media lacking uracil. In our study, we supplemented this media
832 with fluconazole, radicicol, or DMSO when appropriate.
833
Selecting drug concentrations
Our goal was to choose concentrations of each drug that would not kill so many yeast cells as to
dramatically decrease barcode diversity. We wanted to maintain a high number of unique barcodes so we
could track a high number of yeast lineages as they independently evolved drug resistance. We measured
the effect of each drug and drug combination on the growth rate of a single barcoded yeast strain using a
plate reader to track changes in optical density (OD) over time. Ultimately we chose a "low"
concentration of each drug that appeared to have no effect on growth rate, and a "high" concentration that
appeared to reduce growth rate by about 15% (**Figure S2**). Though the lowest concentration of radicicol
that we tested on a plate reader was 10 μM, we chose 5 μM as our low RAD concentration because
previous work suggested this concentration had widespread effects on yeast physiology without affecting
growth (Geiler-Samerotte et al., 2016; Jarosz and Lindquist, 2010). To perform our plate reader
experiment, a single colony was grown to saturation. From this culture, 5 μl was added to every well of a
96-well plate, where every well contained 195 μl of M3 media. Some wells also contained either
fluconazole, radicicol, DMSO, or combinations of these drugs. The concentrations that were tested are
listed on the horizontal axis of **figure S2**; each drug condition was replicated six times. The 96-well plate
was incubated at 30°C for 48 hours on a plate reader and OD measurements were taken every 30 minutes.
Raw OD values were exported and maximum exponential growth rates for all tested conditions were
calculated from the log-linear changes in OD over time.
852
Inserting 300,000 unique DNA barcodes into otherwise genetically identical yeast cells
In order to track many yeast lineages as they independently develop drug resistance, we needed to insert
unique DNA barcodes into many yeast cells. Plasmids harboring barcodes (pBar3) were the same as those
used in a previous barcoded evolution experiment (Levy et al., 2015) and were generously provided to us
by Sasha Levy. These barcodes are 25 base pairs in length. They are targeted to an artificial intron within
the Ura3 gene, such that they must be retained in media lacking uracil but are not expressed and thus do

not themselves affect fitness (Levy et al., 2015). We transformed this barcode library (pBar3) into the
landing pad strain (SHA185) as was done previously, activating a Cre-lox recombination system by
growing the cells in YP-galactose, which resulted in genomic integration of the barcode. However, our
efforts to perform extremely high efficiency transformations from which we could isolate hundreds of
thousands of uniquely barcoded yeast were unsuccessful, despite manipulating the levels and timing of
the inducer (galactose). Ultimately we performed 24 separate transformations and pooled many of these
to obtain a large pool of barcoded yeast where every yeast cell was genetically identical except for its
DNA barcode.
867
Examining the frequency of each barcode in the starting pool of cells
We sequenced the barcode region of these 24 transformed yeast populations on the Hiseq X platform
using a dual index system (Kinsler et al., 2023) to discern barcode coverage, i.e., how many total unique
barcodes were successfully inserted into yeast cells and how evenly these barcodes were sampled. We
needed many uniquely barcoded yeast in order to observe many different adaptive lineages within each
evolution experiment. But barcodes with very high frequencies, referred to herein as monster lineages,
were present in 10 of the 24 transformations and present a problem. Monster lineages allow too many
cells to carry the same barcode, giving that barcode more chances to develop an adaptive mutation. This
could allow different cells harboring that same barcode to pick up different adaptive mutations,
destroying our ability to draw conclusions about adaptive mutations by using barcodes. Therefore, our
final library of barcoded lineages was created by pooling 14 individual transformations together, choosing
those 14 that lacked monster lineages, which we defined as lineages representing greater than 1% of all
transformants. Our sequencing results suggest that this library contains about 300,000 unique barcodes.
881
Initiating 12 barcoded evolution experiments
All evolution experiments started from the same pool of roughly 300,000 uniquely barcoded yeast
lineages. To start the evolution experiments, a pea sized amount of the frozen yeast barcode library was
grown up in 4 ml YPD for 4 hours at 30 °C in a shaking incubator at 220 rpm. Then, 300 µl of the grown
barcode library was added to each of 12 pre-prepared 500mL flasks representing the 12 evolution
experiments listed in **Table 1**. To prepare these flasks, first, 1.2 L of M3 media was warmed at 30°C.
Then, 100 ml was added to each of 12 flat bottom flasks. Next, 500 µl of the appropriate drug or drug
combination was added to each flask. Drugs were pre-diluted, aliquoted and frozen such that 500 µl of the
appropriate tube could be added to each flask to achieve the desired concentration as listed in **Table 1**. All
drugs were resuspended in DMSO such that the final concentration of DMSO in all experiments (except
the "no drug" control) was 0.5%.
893
Performing barcoded evolution experiments
Evolution experiments were performed following previous work (Levy et al., 2015). After initiation (see
above), the yeast in every flask were allowed to grow at 30°C with shaking at 200 RPM for 48 hours.
Then, the flasks were removed from the incubator and 400 – 1000 µl of each culture was transferred to a
new pre-prepared flask with identical conditions to the first. The reason we added more volume (1000 µl)
to some flasks than previous work was that the cell counts at the end of the 48 hours were lower for some
of our higher drug conditions. We adjusted the transfer volume to maintain a transfer population of 4×10^7
cells, which was the same as in previous work (Levy et al., 2015). We completed a total of 24
growth/transfer cycles, corresponding to 192 generations of growth assuming 8 generations per 48-hour
cycle (Levy et al., 2015). Following each transfer, the remaining culture from each flask were split into
two 50 ml conical vials, centrifuged for 3 minutes at 4000 rpm, and the supernatant was discarded. The
final pellet was resuspended in 30% glycerol up to a total volume of 6 ml before being split into three 2
ml cryovials and stored at -80°C. These frozen samples were later utilized for barcode sequencing and
isolating adaptive mutants.
908
909 **Isolating a large pool of adaptive mutants**

To generate a large pool of diverse adaptive mutants, our goal was to collect a sample from each evolution experiment at a time point when there were many different adaptive lineages competing. If we sampled too late, the adaptive lineage with the greatest fitness advantage would have already risen to high frequency, thus reducing diversity. But if we sampled too early, adaptive lineages would not yet have risen in frequency above other lineages. Therefore, we chose to sample cells from a time in each evolution experiment when many barcoded lineages appeared to be rising in frequency (**Figure S1**). We sampled either 1 or 2 thousand cells per each evolution experiment by spreading frozen stock from the chosen time point onto agarose plates, scraping 1 or 2 thousand colonies into a 15mL conical tube containing a final concentration of 30% glycerol, and freezing the pool pertaining to each of the 12 evolutions. We sampled 2,000 cells from most evolution experiments, but sampled only 1,000 from those containing a high concentration of FLU as those evolutions appeared to have reduced barcode diversity (**Figure S1**), presumably because high FLU represents a strong selective pressure. We sequenced the barcodes from each of these 12 pools so that we could track which adaptive mutants originated from which evolution experiment (see methods section below entitled, "**Inferring where adaptive lineages originally evolved**").

Initiating barcoded fitness competition experiments

To assess the fitnesses of the 1 or 2 thousand barcoded lineages that we sampled from each evolution experiment, we pooled all sampled lineages together into a larger pool of roughly 21,000 barcoded lineages. We used this larger pool to initiate 24 fitness competition experiments, 2 replicates for each of the 12 conditions listed in **Table 1**. In this type of competition, we measure fitness by tracking changes in each barcode's frequency over time. Barcodes that rise in frequency represent strains that have higher fitness than others.

Our goal was to calculate the fitness effect of adaptive mutations. Therefore, we needed to calculate the fitness of every evolved lineage relative to the unmutated ancestor of the evolution experiments. To do so, we followed previous work by spiking in a large quantity of this unmutated ancestor strain into each fitness competition, with this ancestor making up at least 90% of the final culture (Kinsler et al., 2020; Venkataram et al., 2016). In environments containing a high concentration of FLU which resulted in the ancestral strain having a more severe growth defect, we spiked in the ancestor such that it represented 95% of the final pool.

To avoid wasting 90% or more of our sequencing reads on the ancestor strain's barcode, we created a barcodeless ancestor strain. This strain was created by transforming SHA185 with a linear piece of DNA such that the genetic background was identical to the strains of the barcoded library, but the homology to the primers used to amplify the barcode was missing. Thus the DNA from these cells does not get amplified or sequenced during subsequent steps.

In addition to this barcodeless ancestor, we also spiked in some barcoded ancestral strains at lower frequency (1%) to use as "reference" or "control" strains, following previous work (Kinsler et al., 2023, 2020). These strains have been previously shown to possess no fitness differences from the ancestor. We used these strains as a baseline when calculating relative fitness by setting the fitness of these strains to zero during our fitness inference procedure (see methods section below entitled, "**Inferring fitness**").

All 24 fitness competitions were performed simultaneously in one big batch (Kinsler et al., 2023) and initiated from the same pool of roughly 21,000 barcoded evolved yeast lineages, barcodeless ancestor, and control strains. To initiate the competitions, 7×10^7 cells from this pool were added to 24 pre-prepared 500mL flasks corresponding to the conditions listed in **Table 1**. These flasks were prepared exactly the same way as was done for the evolution experiments (see above in "**Performing barcoded evolution experiments**"). Each flask was allowed to grow for 48 hours at 30°C with shaking at 200 RPM.

Performing barcoded fitness competition experiments

Fitness competitions were performed following previous work (Kinsler et al., 2020). After the initial flasks were allowed to grow for 48 hours, they were removed from the incubator and 400 μl from each culture representing $4x10^7$ cells were transferred to a new flask with identical media. For each of 24 competitions, we completed a total of 4 growth/transfer cycles, corresponding to 40 generations of growth assuming 8 generations per 48-hour cycle (Levy et al., 2015). Following each transfer, the remaining culture from each flask was split into two 50 ml conical vials, centrifuged for 3 minutes at 4000 rpm, and the supernatant was discarded. The final pellet was resuspended in 30% glycerol up to a total volume of 6 ml before being split into three 2 ml cryovials and stored at -80°C. These frozen samples were later utilized for DNA extraction and subsequent barcode sequencing.

Despite the fitness competition experiments being conducted for nearly the same number of generations (40) as were the evolution experiments before isolating adaptive lineages, we do not anticipate that secondary mutations will bias fitness measurements. Previous work has demonstrated that in this evolution platform, most mutations occur during the transformation that introduces the DNA barcodes (Levy et al. 2015). In other words, these mutations are already present and do not accumulate during the 40 generations of evolution. Therefore, the observation that we collect a genetically diverse pool of adaptive mutants after 40 generations of evolution is not evidence that 40 generations is enough time for secondary mutations to bias abundance values. For a detailed treatment of how secondary mutations have a minimal influence on fitness, see Venkataram et al 2016.

Extracting genomic DNA

DNA was extracted from 500 μl of concentrated frozen stocks pertaining to the evolution experiments and fitness competitions. Frozen cells were thawed and pelleted. Cells were treated with 250 μl of 0.1 M Na2EDTA, 1M sorbitol and 5U/μl zymolyase for a minimum of 15 minutes at 37 °C to remove the cell wall. Lysis was completed by adding 250 μl of 1% SDS, 0.2N NaOH and inverting to mix. Proteins and cell debris were removed with 5M KOAc by spinning for 5 min at 15,000 rpm. Supernatant was moved to a new tube and DNA was precipitated with 600 μl isopropanol by spinning for 5 min at 15,000 rpm. The resulting pellet was washed 1 ml of 70% ethanol before being resuspended in 50 μl water plus 10μg/ml RNAse. Extracted DNA was quantified using the NanoDrop spectrophotometer and all samples were diluted to a concentration of 50 ng/μL for barcode amplification and sequencing library preparation.

Preparing barcodes for high-throughput multiplexed sequencing using PCR

Extracted DNA was prepared for sequencing using a two-step PCR that preserves information about the relative frequency of each barcode in each sample (Kinsler et al., 2023, 2020; Venkataram et al., 2016). Briefly, in the first step PCR, the barcode region is amplified from the genomic DNA, labeled with a sample-specific combination of primers, and tagged with a UMI. This step utilizes a short 3 cycle PCR with New England Biolabs OneTaq polymerase. Purification of the first step product to remove excess reagents was performed using Thermo Scientific GeneJET PCR Purification Kit. The second step PCR attached Illumina indices that were used to distinguish samples from different experiments and timepoints. We utilized a dual indexing scheme to prevent index misassignment that is common when sequencing amplicon libraries using patterned flow cell technology (Kinsler et al., 2023). Amplification of this second step of PCR was done with a longer 23 cycle PCR using Q5 polymerase. Final libraries were bead purified using 0.8X Quantabio sparQ Pure Mag beads. Quantification of the final PCR products was done using the Invitrogen Qubit Fluorometer before all samples were pooled at equimolar ratios for sequencing.

Sequencing and clustering barcodes

Next Generation Sequencing was performed at either Psomagen (Rockville, MD) or at the Translation
Genomics Research Institute (Phoenix, AZ) on patterned flow cells (either an Illumina HiSeqX or
NovaSeq) using 2 x 150 base pair paired end reads. Samples were dual indexed to allow multiplexing
while minimizing contamination from index misassignments (Kinsler et al., 2023). The 20 base pairs of
variable sequence referred to as a DNA barcode were identified and clustered to determine the number of
unique barcodes and the frequency of each barcode in each sample. For the evolution experiments, this
was done following our previous work (Kinsler et al., 2020; Venkataram et al., 2016). For the fitness
competition experiments, this was done using updated software (Zhao et al., 2018) with the following
command:
bartender_extractor_com -f ${SAMPLE}_R1_001.fastq -o ${SAMPLE}_extracted -q 0 -p
GTACC[5]AA[5]AA[5]TT[5]ATAAC -m 2 -d f -u 0,8
1021
Inferring fitness
In fitness competition experiments, fitness is often inferred from the log-linear change in a strain's
frequency over time (Bakerlee et al., 2021; Geiler-Samerotte et al., 2011; Kinsler et al., 2023). Recently,
more advanced methods to infer fitness have emerged that take into account nonlinearities in frequency
changes over time, for example, nonlinearities that reflect changes in the mean fitness of the population
(Kinsler et al., 2020; F. Li et al., 2018; Li et al., 2023; Venkataram et al., 2016). We had trouble
implementing these newer methods on our fitness data, perhaps because many of our evolved lineages,
and our control strains, have low fitness in some drugs. This caused their barcodes to rapidly decline in
frequency such that they received low counts only at later time points. Their counts could become so low
that these lineages would seemingly disappear due to sampling error, and then reappear at a subsequent
time point. This dramatic (but false) late increase in frequency was sometimes interpreted as evidence of
very high fitness, especially when we inferred fitness using approaches that account for nonlinearities.
1034
To contend with this issue, we applied strict coverage thresholds to every fitness measurement: we
required at least 500 counts across all timepoints in order to infer fitness for a given lineage in a given
environment. This is stricter than previous work that does not require a minimum number of reads per
each lineage and instead requires a minimum number of reads per time point (Kinsler et al., 2020). We
found that 774 lineages passed our threshold in at least one replicate experiment per all 12 environments.
Of these, 729 passed for both replicates and the final fitness value we report represents the average of
both replicates.
1042
Even with our strict coverage threshold, fitness inference methods that account for nonlinearities still
interpreted minor stochastic fluctuations in fitness at later time points as evidence of a fitness advantage,
even if fitness dramatically declined in earlier time points. Therefore, we calculated fitness via the
traditional method, as the slope of the log-linear change in barcode frequency relative to the average slope
of the control strains. We found that this method is less sensitive to that type of error. Using this method,
we found that our fitness inferences were reproducible between replicates (**Figure S4A**), and between
experiments performed in similar conditions (e.g., medium vs. high concentrations of the same drug)
(**Figure S4B**). When we increased our coverage threshold to require an order of magnitude more reads
per lineage per measurement (from 500 to 5000), we lost 157 lineages (from 774 to 617), saw
reproducibility increase across replicates (from an average Pearson correlation of 0.756 to 0.813) and the
main conclusions of our study were unchanged in that the same 6 clusters were present on a UMAP
(**Figure S7**).
1055
Identifying adaptive mutations using whole-genome sequencing
One downside of barcoded evolution experiments is that all lineages exist together in a pooled culture.
Fishing out adaptive lineages in order to perform whole genome sequencing is a major challenge
(Venkataram et al., 2016). Here, we randomly selected cells from these mixed pools for whole genome

sequencing, sometimes selecting from later time points in the evolution experiments and sometimes
selecting from the samples of 1 or 2 thousand cells that were isolated to initiate fitness competitions.
1062
To perform whole genome sequencing, cells from mixed pools were spread onto M3 agarose plates,
single colonies were selected and grown in YPD to saturation. DNA was extracted using the PureLink™
Genomic DNA Mini Kit (K182002). Sequencing libraries were made using Illumina DNA Prep kit by
diluting reactions by ⅕. Briefly, samples were prepared such that the starting concentration in 6 ul was
between 20 and 100 ng of DNA. 2µl of BLT and TB1were added to the starting material and incubated on
a thermocycler at 55°C (lid 100°C) for 15 min. 2 µl of TSB was added to each reaction and incubated at
37 C (lid 100°C) for 15 min. Beads were washed 2 times with 20 µl of TWB. Following the final wash, 4
µl of EPM, 4 µl of water and 2 µl of UD indexes were added to each sample. Depending on starting
concentration, PCR was performed based on Illumina guidelines as follows: lid 100°C, 68°C for 3 min,
98C for 3 min, [98°C for 45s, 62°C for 30s, 68°C for 2min] for 6 to 10 cycles, 68°C for 1 min, 10°C hold.
PCR products were cleaned with a double side sized selection as follows: 4 µl of each sample was pooled
together (32 µl total for 8 samples) and added to 28 µl of water plus 32 µl of SPB. After a 5 min
incubation 25 µl of supernatant was moved to a new tube containing 3 µl of SPB. Beads were washed
with fresh 80% ethanol and libraries were eluted in 12 µl RSB. Samples were multiplexed using
Illumina's unique dual (UD) index plates (A-D) and sequencing was performed with 2x150 paired end
sequencing on HiSeq X at Psomagen (Rockville, MD).
1079
In total 122 colonies were randomly picked and sequenced. As one might expect, barcodes that rose to
high frequency were more likely to be picked multiple times. In an attempt to find lineages with unique
attributes, some cultures were grown at 37°C or plated to high concentrations of drug prior to picking
isolated colonies for sequencing. Of the 122 genomes we sequenced, only 53 had unique barcodes that
pertained to the 774 lineages for which we obtained high enough barcode coverage to infer fitness. Only
two of these 53 had no sequenced mutations suggesting their fitness increase over ancestor is due to a
mutation we are unable to identify by sequencing, perhaps a change in ploidy. The other 51 all had at
least one single nucleotide mutation in a gene reported in **supplemental table 1**. Whole genome
sequences were deposited in GenBank under SRA reference PRJNA1023288.
1089
Variant calling was done using GATK as described here: https://github.com/gencorefacility/variant-
calling-pipeline-gatk4. Identified variants were annotated using SnpEff (Cingolani et al., 2012). Variant
call files from 132 (53 unique/in CS) sequenced lineages were analyzed in R and compared to reference
strain GCF_000146045.2 (Genome assembly 64: sacCer3). SNPs present in the ancestor (as well as all
evolved lineages) were ignored as these could not have caused the fitness differences we observed. We
also ignored SNPS that were present in a substantial number of evolved lineages, as these likely represent
background mutations that were present in a substantial portion of the cells representing the landing pad
strain (SHA185). These are reported in **supplemental table 1** and include: SRD1-Glu97Lys, RSC30-
Gly571Asp, OPT1-Val143Ile and LYS20-Thr29Met.
1099
Measuring growth curves of evolved lineages with unexpected fitness in multidrug conditions
Though fitness differences are not necessarily due to differences in maximum growth rate (Y. Li et al.,
2018), we measured growth curves for a few lineages. In one case, we did so to investigate a case where
an evolved lineage had unexpectedly low fitness in multidrug conditions (**Figure 7B**). Indeed, we found
that this mutant grew more slowly in those conditions (**Figure 7D; S11**). To perform this test, a lineage
with a mutation to GBP2, a lineage with a mutation to HDA1, and sometimes the ancestor strain were
streaked to YPD plates. We used the barcodeless ancestor strain, which is identical to the evolved
lineages in every way except for lacking a barcode, and is described above in the methods section
entitled, "**Initiating barcoded fitness competition experiments".** A single colony of each strain was
isolated from YPD plates and was used to inoculate an overnight YPD culture. After ~24 hours, a coulter
counter (BD) was used to determine the number of cells/ml present in each culture. Next, all cultures

were diluted such that the starting number of cells was 250,000 in 6 ml of M3 plus drug (either HR, LF,
LR, LRLF or HRLF, see **Table 1**). To measure growth curves, these samples were allowed to grow at 30
degrees C. OD was measured every 10 minutes as the cultures were grown to saturation using the
compact rocking incubator TVS062CA (Advantec Mfs). Raw growth curves for these conditions are
shown in **Figure S11B and C**. Maximum growth rate was calculated using a sliding window approach to
determine the region of each growth curve with the steepest log-linear slope. We used similar methods to
measure growth curves for 3 mutants from cluster 1 and 3 from cluster 3 in **Figure 5**.
1118
Determining ploidy
While our barcoded yeast strain is haploid, previous studies observed that some cells diploidize during the
course of evolution in M3 media and by doing so gain a fitness advantage (Levy et al., 2015; Venkataram
et al., 2016). To ensure that observed fitness effects in our experiments were not largely due to the effects
of diploids, we estimated the percent of diploid cells in each of our populations. We chose to make our
estimates from frozen samples taken at the same time points from which we sampled 1 or 2 thousand cells
to initiate fitness competitions (**Figure S1**). As such, our estimates also report on the percent of diploids
that were present at the start of the fitness competitions experiments (**Table S2**).
1127
To study ploidy, we used the nucleic acid stain SYTOX Green, which is capable of selectively staining
the nucleus of fixed cells and has been shown to be optimal for use in budding yeast (Haase, 2004). For
each of the 12 evolution experiments conditions, a small amount of freezer stock from the chosen
timepoints (**Figure S1**) was plated to YPD and grown for ~48 hours. Individual colonies were picked and
transferred to 96-well plates, one full plate per each condition, before being fixed with 95% ethanol for 1
hour. Plates were centrifuged at 4500 rpm and supernatant was discarded. 50 μL RNase A was added to
the samples at a concentration of 2mg/mL, and the plates were then incubated for 2 hours at 37°C. Cells
were pelleted by centrifuge and the supernatant was removed, which was followed by treatment with 20
μL of the protease pepsin at a concentration of 5mg/uL. Pepsin treated samples incubated at 37°C for 30
minutes before centrifugation and removal of supernatant. Finally, cells were resuspended in 50 μL
TrisCL (50 mM, pH 8) and stained with 100 μL of 1μM SYTOX Green. Known diploid and haploid
strains were used as controls alongside our samples to determine the expected fluorescence of stained
diploid vs. haploid cells. Analysis was performed using a ThermoFisher Attune NxT, housed in the Flow
Cytometry Core Facility at Arizona State University.
1142
Dimensional reduction
Our fitness inference procedure resulted in a data set consisting of nearly 10,000 fitness measurements
(774 lineages x 12 conditions = 9288 fitness measurements). Dimensional reduction was performed on
these data using UMAP (McInnes et al., 2018b). Clusters of similar mutants were identified and colored
using a gaussian mixed model (Fraley and Raftery, 2003); Bayesian Information Criteria (**Figure S6**) as
well as follow up genotyping and phenotyping studies (see **Figures 5 – 8**) were used to select the number
of clusters. These analyses were performed in R; code can be found
https://osf.io/pxyv9/?view_only=51241d8f00c24f7e83f1ece3ae31a53b.
1151
In order to prevent conditions with the most variation in fitness (e.g., high FLU) from dominating, we
normalized fitness measurements from each of the 12 environments to have the same overall mean and
variance (we transformed the data from every environment to have a mean of 0 and a standard deviation
of 1) before performing dimensional reduction. This normalization procedure did not have a dramatic
effect on the UMAP (**Figure S7A**). We also explored normalizing all data to account for magnitude
differences by setting the average fitness of each lineage across all 12 environments to 0. Doing so did not
significantly change the groupings present in the UMAP from those displayed in **Figure 4** (**Figure S10**)
other than in the ways we describe in **Figure 6**. Reducing our data set to 617 adaptive lineages with very
high sequencing coverage (**Figure S7B**) also did not significantly affect the way that mutants cluster into
groups, nor did using a different dimensional reduction algorithm altogether (**Figure S8** and see next

paragraph). In short, the clustering of mutants was robust to the different decisions we made when
choosing how to analyze these data.
1164
In order to assess whether clusters identified from the UMAP are robust to alternative clustering methods,
we also used hierarchical clustering to identify clusters of mutants with similar fitness profiles. First, we
computed the pairwise distance of all lineages across the fitness profiles. Then, we used Ward's method
from scikit-learn to iteratively cluster lineages such that the within-cluster variation is minimized
(Pedregosa et al., 2012; Ward, 1963). To test the consistency of lineage clustering, we chose a pairwise
cluster distance cutoff of 11, which results in the same number of clusters (7) as identified with the
UMAP clustering approach used in the main text. We then compared the identity of the lineages within
each of these clusters with the UMAP clusters. We found that, for most clusters, over 80% of lineages
from the UMAP cluster corresponded with a unique hierarchical cluster and labeled these hierarchical
clusters according to this correspondence (**Figure S8**). For UMAP cluster 1, lineages were more evenly
split between two clusters. 64% of these lineages clustered together in what is labeled as hierarchical
cluster 1 and 30% in hierarchical cluster 1/7 (**Figure S8**), which contains all of the control lineages that
comprise UMAP cluster 7. Despite these lineages clustering more closely with control lineages than the
remainder of the cluster 1, they do tend to cluster distinctly with the control lineages, suggesting they
have behavior that is distinguishable from the control lineages. If we consider these cluster 1 mutants that
end up in cluster 3/7 as "mis-clustered", we find that 85% of lineages from each UMAP cluster are
clustered together in the corresponding hierarchical cluster. If we consider these as "consistently
clustered", this metric increases to 90% of lineages correctly clustered. Similarly, clustering lineages
using principal component analysis also largely preserved the clusters reported in **Figure 4** (**Figure S9**).
Altogether, this analysis shows that the results we show are robust to alternative methods of clustering.
1185
Inferring where adaptive lineages originally evolved
All 774 adaptive lineages were isolated from one of the 12 evolution experiments at the timepoint
indicated in **figure S1** (see Methods section entitled, "**Isolating a large pool of adaptive mutants**"). The
sample we isolated from each evolution experiment was sequenced prior to pooling. This allows us to
computationally determine which barcoded lineages originated from which evolution experiment to
generate the pie charts in F**igures 3**, **5**, **6**, **7** and **8**.
1192
1193   If adaptive mutation arose independently during the course of each evolution experiment, it would be
1194   unlikely for any adaptive lineage we study to be present in more than one of the evolution conditions.
1195   This would make it very easy to assign each barcode to the evolution experiment from which it
1196   originated. However, this was not the case for many barcoded lineages.
1197
Previous work explained that the transformation procedure used to insert a barcode into the landing pad of
SHA185 is itself mutagenic, such that many mutations arise prior to the start of the evolution experiments
(Levy et al., 2015). Since all our evolution experiments were started from the same pool of barcoded
lineages, we thus expect that many adaptive lineages will be present in more than one condition.
However, it is not expected that these adaptive lineages will be present at the same frequency in every
condition; instead these frequencies change with the fitness of the mutation each lineage possesses.
Therefore, when an adaptive lineage appeared in multiple conditions, we weighted its origin to reflect its
frequency in each condition. In other words, adaptive lineages that were only present in the sample taken
from a single evolution condition were identified and assigned a single origin condition in the pie charts
in **Figures 3**, **5**, **6**, **7**, and **8**. But for adaptive lineages found in the samples taken from more than one
evolution condition, the proportions assigned to each origin condition in the pie charts was scaled to equal
the relative frequencies of that lineage in all evolution conditions where it was observed. Associated data
and code can be found here: https://osf.io/pxyv9/?view_only=51241d8f00c24f7e83f1ece3ae31a53b.
1211
1212

1214   **Supplemental Figures and Tables**

1215

1216
Figure S1: Twelve barcoded evolution experiments track ~300,000 lineages as they adapt to different
drug concentrations and combinations. In each plot, every line represents a unique barcoded lineage. The
vertical axis represents barcode frequency. When lineages reach a frequency of zero it means they were not
sampled at that time point of the experiment. The black boxes indicate the transfer number of each evolution
experiment from which evolved lineages are sampled; the number of cells sampled (colonies picked) is in the
upper right hand corner.

1223
1224

1225
Figure S2: Chosen drug concentrations do not dramatically reduce yeast's maximum growth rate. We
measured the growth rate of a single barcoded yeast lineage, prior to the evolution experiment, in different
concentrations and combinations of drugs using a plate reader to track changes in optical density over time.
Maximum growth rate is reported as a percentage of the maximum growth rate in conditions lacking FLU or
RAD. Maximum growth rate was calculated as the maximum log-linear slope of the change in optical density
over time.
1232

1234 Figure S3: Twenty four fitness competitions track evolved lineages as their barcodes change frequency.
In each plot, every line represents a barcoded lineage. Lineages with >5 reads per experiment are shown (4,815
lineages shown). Control lineages, known to possess no fitness differences relative to the ancestor, are
highlighted in black. Evolved lineages with higher fitness than controls are yellow, similar fitness to controls
(i.e., relative fitness ~0) are purple, and lower fitness than controls are orange. When a lineage declines in
frequency so much that its associated barcode is no longer observed, its line abruptly ends. Some lineages

appear to decline and then increase in frequency; this happens because low frequency lineages are subject to
sampling noise.
1242
1243
1244

1245
Figure S4: Fitness measurements are reproducible between replicates and closely related conditions.
A. Every point depicts one of the evolved lineages that was observed more than 500 times in both replicates of
the fitness competition experiment for a given condition. The average Pearson correlation across all pairs of
replicates in all conditions is 0.75. **B.** Every point depicts one of the 774 lineages that was observed more than
500 times in at least one replicate fitness competition. Here, instead of comparing the fitness of replicates, we
compare fitness across conditions for 3 closely related pairs of conditions. Comparisons across conditions may
be less noisy than those across replicates because fitness in a particular condition represents the average fitness
across replicates in that condition.
1254

1255
1256

Figure S5: **The two types of adaptive mutants depicted in figure 3 sort into different clusters on the UMAP.** The top 100 highest fitness lineages in high FLU (blue) and high RAD (red) largely cluster into different groups, with the 7 overlapping mutants (black) falling into the uppermost group.

1267
1268 **Figure S6: Bayesian information criteria (BIC) scores suggest the 774 mutants cluster into between**
1269 **6 and 13 groups.** We used a gaussian mixture model to distinguish clusters of mutants with unique
1270 fitness profiles. **A.** BIC scores for analysis performed on 774 mutant lineages that each were observed at
1271 minimum 500 times per environment. **B** BIC scores for analysis performed on 617 mutant lineages that
1272 each were observed at minimum 5000 times per environment.

1273
1274
1275

1276
Figure S7. UMAP structure is robust. A. In this UMAP plot, relative fitness of the 774 lineages is fed
directly into UMAP prior to normalizing the data to have equal variance across all environments as was done
in **figure 4D**. This has little impact on the appearance of the UMAP. **B.** UMAP made with 617 lineages that
were observed more than 5000 times in at least one replicate. While the appearance of the UMAP is inverted
relative to **figure 4D**, the clusters largely retain the same lineages. As lower coverage (noisier) data points are
removed from this dataset, the observation that the overall patterns are maintained suggests our original
clusters were not based on noisy fitness measurements.

1284

1285

1286
Figure S8. Clusters are robust to a hierarchical clustering method. A. In order to assess whether
clusters identified from the UMAP are robust to alternative clustering methods, we used hierarchical
clustering to identify clusters of mutants with similar fitness profiles. This figure depicts a dendrogram
where the branch length between each lineage (or group of lineages) represents the distance between the
clustered groups as measured across all measured conditions. The terminal branch leading to each lineage
is colored by that lineage's cluster from the UMAP clustering used in **figure 4** of the main text. Dark gray
lines show clusters identified using a between-cluster cutoff of 11, which creates 7 clusters, consistent
with the number of clusters we identified in **figure 4** via clustering on the UMAP space. Hierarchical
clusters are labeled and colored corresponding to whichever of the UMAP clusters they share the most
overlapping lineages with. Note that the y-axis is linearly scaled between 0 and 10 and log2 scaled above
10 for visualization purposes. **B.** A matrix quantifying how well the results of this hierarchical clustering
method correspond with those of the UMAP method presented in **figure 4**, depicting the percentage of a
given UMAP cluster that ends up with each corresponding hierarchical cluster. For the most part, mutants
that are clustered together in the UMAP in **figure 4** are also clustered together in **panel A**. Even though it
appears that mutants in cluster 1 (which are those most resistant to low FLU) move to a different cluster,
these mutants move together to the cluster pertaining to the control strains as they have the least severe
effects on fitness.
1304
1305

Figure S9. Clusters are robust to principal component analysis. In order to assess whether clusters identified in the UMAP are robust to alternative clustering methods, we used principal component analysis and plotted PC1 vs PC2 to visualize mutants in two dimensional space. Points representing each mutant are colored corresponding to their cluster identity in **figure 4**. The amount of variation captured by each component is stated in the axis label.

1312

Figure S10: UMAP on data that were normalized to account for magnitude differences (row means
set to 0). A. Every point in this plot represents one of the 774 barcoded lineages colored by cluster; clusters
were identified using a gaussian mixture model. After normalizing to account for magnitude differences,
the 774 adaptive lineages cluster into 6 (not 7) groups based on variation in their fitness profiles. The
number of clusters is reduced relative to the UMAP in **figure 4** because the control lineages no longer form
a separate group; they cluster into the middle cluster in light green. **B.** This plot shows the same UMAP as

above in **panel A**, but the color of each of the 774 points is changed to reflect its color in the UMAP in
figure 4. Despite coloring based on a different UMAP, differently colored points still largely fall into
separate clusters, except for those points colored blue or orange, which are discussed in **figure 6**. **C.** This
plot depicts all lineages in clusters 1 – 3 in either the **figure 4** UMAP (not normalized to lessen magnitude
differences) or the above UMAP (normalized to lessen magnitude differences). Lineages that do not switch
from one cluster to another are represented by diamonds while those that switch are noted with circles.
Across these three clusters, 67% of lineages remain in their original cluster.

1326

1327

1328

1329

Figure S11: Unexpected tradeoffs in evolved lineages in cluster 4 and 5 in response to combined
drugs. A. Cluster 5 (red) has an unexpected fitness disadvantage in the LRLF multidrug environment relative to
cluster 4 (green), given that cluster 5 lineages do not have a relative fitness disadvantage in the relevant single drug
environments. This plot is similar to the one in **figure 7B** except here the multidrug condition shown is LRLF rather
than HRLF. Boxplots summarize the distribution across all lineages within each cluster in each environment,
displaying the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5
$\times$ IQR (whiskers). **B.** Raw growth curve data from which relative maximum growth rates presented in **Figure 7D** were
calculated. Two mutants, one to GBP2 (cluster 4; green) and one to HDA1 (cluster 5; red) were each grown
individually in the conditions shown, as was the barcodeless ancestor strain (grey). All experiments were conducted
in duplicate. We initiated all cultures from the same number of starting cells. The mutant in cluster 5 (red) has an
obvious growth disadvantage in the combination condition (HRLF). **C.** Raw growth curve data for the same two
mutants in **panel B** in different conditions. These growth experiments were conducted similarly to those in **panel B**,
except they were tracked for a longer period of time, lacked the ancestor strain, and involved 3 replicates per mutant.
There appear to be obvious and reproducible differences in the growth curves of the two mutants surveyed.

1344
1345
1346

1347 **Table S1:** The sequenced mutations present in each of the adaptive lineages on which we performed whole
1348 genome sequencing:
1349 https://osf.io/pxyv9/

1350
1351

1352 **Table S2:** Estimated percentage of diploid cells in each evolution condition at the time point we sampled,
1353 determined using nuclear staining and flow cytometry as described in Materials and Methods. *For the
1354 High Radicicol + Low Fluconazole condition, 95 instead of 96 samples were measured due to undetectable
1355 growth in one well.

1356
1357

Evolution Conditions	Diploids detected (96 sampled)	Estimated % diploid
Low Rad	8	8.3%
Med Rad	0	0.0%
High Rad	0	0.0%
Low Flu	9	9.4%
Med Flu	0	0.0%
High Flu	1	1.0%

LR + LF	4	4.2%
HR +LF*	1	1.1%
LR + HF	1	1.0%
HR + HF	3	3.1%
DMSO	6	6.3%
No Drug	9	9.4%

1358
1359
1360
1361 **References**
1362

Abel zur Wiesch P, Kouyos R, Abel S, Viechtbauer W, Bonhoeffer S. 2014. Cycling empirical antibiotic
therapy in hospitals: meta-analysis and models. *PLoS Pathog* **10**:e1004225.
Aissa AF, Islam ABMMK, Ariss MM, Go CC, Rader AE, Conrardy RD, Gajda AM, Rubio-Perez C,
Valyi-Nagy K, Pasquinelli M, Feldman LE, Green SJ, Lopez-Bigas N, Frolov MV, Benevolenskaya
EV. 2021. Single-cell transcriptional changes associated with drug tolerance and response to
combination therapies in cancer. *Nat Commun* **12**:1628.
Algazi AP, Othus M, Daud AI, Lo RS, Mehnert JM, Truong T-G, Conry R, Kendra K, Doolittle GC,
Clark JI, Messino MJ, Moore DF Jr, Lao C, Faller BA, Govindarajan R, Harker-Murray A,
Dreisbach L, Moon J, Grossmann KF, Ribas A. 2020. Continuous versus intermittent BRAF and
MEK inhibition in patients with BRAF-mutated melanoma: a randomized phase 2 trial. *Nat Med*
26:1564–1568.
Allen RC, Angst DC, Hall AR. 2019. Resistance Gene Carriage Predicts Growth of Natural and Clinical
Escherichia coli Isolates in the Absence of Antibiotics. *Appl Environ Microbiol* **85**.
doi:10.1128/AEM.02111-18
Allen RC, Pfrunder-Cardozo KR, Hall AR. 2021. Collateral Sensitivity Interactions between Antibiotics
Depend on Local Abiotic Conditions. *mSystems* **6**:e0105521.
Amin AD, Rajan SS, Liang WS, Pongtornpipat P, Groysman MJ, Tapia EO, Peters TL, Cuyugan L,
Adkins J, Rimsza LM, Lussier YA, Puvvada SD, Schatz JH. 2015. Evidence Suggesting That
Discontinuous Dosing of ALK Kinase Inhibitors May Prolong Control of ALK+ Tumors. *Cancer
Res* **75**:2916–2927.
Andersson DI, Hughes D. 2010. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nat
Rev Microbiol* **8**:260–271.
Andersson DI, Nicoloff H, Hjort K. 2019a. Mechanisms and clinical relevance of bacterial
heteroresistance. *Nat Rev Microbiol* **17**:479–496.
Andersson DI, Nicoloff H, Hjort K. 2019b. Mechanisms and clinical relevance of bacterial
heteroresistance. *Nat Rev Microbiol* **17**:479–496.
Ardell SM, Kryazhimskiy S. 2021. The population genetics of collateral resistance and sensitivity. *Elife*
10. doi:10.7554/eLife.73250
Baker CM, Ferrari MJ, Shea K. 2018. Beyond dose: Pulsed antibiotic treatment schedules can maintain

individual benefit while reducing resistance. *Sci Rep* **8**:5866.

Bakerlee CW, Phillips AM, Nguyen Ba AN, Desai MM. 2021. Dynamics and variability in the pleiotropic effects of adaptation in laboratory budding yeast populations. *Elife* **10**. doi:10.7554/eLife.70918

Basra P, Alsaadi A, Bernal-Astrain G, O'Sullivan ML, Hazlett B, Clarke LM, Schoenrock A, Pitre S, Wong A. 2018. Fitness Tradeoffs of Antibiotic Resistance in Extraintestinal Pathogenic Escherichia coli. *Genome Biol Evol* **10**:667–679.

Baym M, Stone LK, Kishony R. 2016. Multidrug evolutionary strategies to reverse antibiotic resistance. *Science* **351**:aad3292.

Berkow E, Lockhart S. 2017. Fluconazole resistance in Candida species: a current perspective. *Infection and Drug Resistance*. doi:10.2147/idr.s118892

Berman J, Krysan DJ. 2020. Drug resistance and tolerance in fungi. *Nat Rev Microbiol* **18**:319–331.

Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* **3**. doi:10.3390/jof3040057

Bosch B, DeJesus MA, Poulton NC, Zhang W, Engelhart CA, Zaveri A, Lavalette S, Ruecker N, Trujillo C, Wallach JB, Li S, Ehrt S, Chait BT, Schnappinger D, Rock JM. 2021. Genome-wide gene expression tuning reveals diverse vulnerabilities of M. tuberculosis. *Cell* **184**:4579–4592.e24.

Bottery MJ, Pitchford JW, Friman V-P. 2021. Ecology and evolution of antimicrobial resistance in bacterial communities. *ISME J* **15**:939–948.

Boyer S, Hérissant L, Sherlock G. 2021. Adaptation is influenced by the complexity of environmental change during evolution in a dynamic environment. *PLoS Genet* **17**:e1009314.

Boyle EA, Li YI, Pritchard JK. 2017. An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169**:1177–1186.

Brandis G, Wrande M, Liljas L, Hughes D. 2012. Fitness-compensatory mutations in rifampicin-resistant RNA polymerase. *Mol Microbiol* **85**:142–151.

Brettner L, Eder R, Schmidlin K, Geiler-Samerotte K. 2024. An ultra high-throughput, massively multiplexable, single-cell RNA-seq platform in yeasts. *Yeast* **41**:242–255.

Brettner L, Ho WC, Schmidlin K, Apodaca S, Eder R, Geiler-Samerotte K. 2022. Challenges and potential solutions for studying the genetic and phenotypic architecture of adaptation in microbes. *Current Opinion in Genetics and Development* **75**.

Buechel ER, Pinkett HW. 2020. Transcription factors and ABC transporters: from pleiotropic drug resistance to cellular signaling in yeast. *FEBS Lett* **594**:3943–3964.

Cairns J, Borse F, Mononen T, Hiltunen T, Mustonen V. 2022. Strong selective environments determine evolutionary outcome in time-dependent fitness seascapes. *Evol Lett* **6**:266–279.

Cannataro VL, Gaffney SG, Stender C, Zhao Z-M, Philips M, Greenstein AE, Townsend JP. 2018. Heterogeneity and mutation in KRAS and associated oncogenes: evaluating the potential for the evolution of resistance to targeting of KRAS G12C. *Oncogene* **37**:2444–2455.

Centers for Disease Control and Prevention (U.S.). 2019. Antibiotic Resistance Threats in the United States, 2019. U.S. Department of Health and Human Services, Centres for Disease Control and Prevention.

Chen VK, Johnson MS, Hérissant L, Humphrey PT, Yuan DC, Li Y, Agarwala A, Hoelscher SB, Petrov DA, Desai MM, Sherlock G. 2023. Far From Home: Evolution of haploid and diploid populations reveals common, strong, and variable pleiotropic effects in non-home environments. *bioRxiv*. doi:10.1101/2023.02.28.530341

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly* **6**:80–92.

Cowen LE, Lindquist S. 2005. Hsp90 potentiates the rapid evolution of new traits: drug resistance in diverse fungi. *Science* **309**:2185–2189.

Cowen LE, Sanglard D, Howard SJ, Rogers PD, Perlin DS. 2014. Mechanisms of Antifungal Drug Resistance. *Cold Spring Harb Perspect Med* **5**:a019752.

Cowen LE, Singh SD, Köhler JR, Collins C, Zaas AK, Schell WA, Aziz H, Mylonakis E, Perfect JR,
Whitesell L, Lindquist S. 2009. Harnessing Hsp90 function as a powerful, broadly effective
therapeutic strategy for fungal infectious disease. *Proc Natl Acad Sci U S A* **106**:2818–2823.
Day T, Huijben S, Read AF. 2015. Is selection relevant in the evolutionary emergence of drug resistance?
Trends Microbiol **23**:126–133.
Eguchi Y, Bilolikar G, Geiler-Samerotte K. 2019. Why and how to study genetic changes with context-
dependent effects. *Curr Opin Genet Dev* **58-59**:95–102.
Fardeau V, Lelandais G, Oldfield A, Salin H, Lemoine S, Garcia M, Tanty V, Le Crom S, Jacq C,
Devaux F. 2007. The Central Role of PDR1 in the Foundation of Yeast Drug Resistance*. *J Biol
Chem* **282**:5063–5074.
Feder AF, Harper KN, Brumme CJ, Pennings PS. 2021. Understanding patterns of HIV multi-drug
resistance through models of temporal and spatial drug heterogeneity. *Elife* **10**.
doi:10.7554/eLife.69032
Flowers SA, Barker KS, Berkow EL, Toner G, Chadwick SG, Gygax SE, Morschhäuser J, Rogers PD.
2012. Gain-of-function mutations in UPC2 are a frequent cause of ERG11 upregulation in azole-
resistant clinical isolates of Candida albicans. *Eukaryot Cell* **11**:1289–1299.
Flynn JM, Joyce ME, Bolon DNA. 2024. Dominant negative mutations in yeast Hsp90 reveal triage
decision mechanism targeting client proteins for degradation. *bioRxiv*.
doi:10.1101/2024.01.02.573950
Flynn JM, Rossouw A, Cote-Hammarlof P, Fragata I, Mavor D, Hollins C 3rd, Bank C, Bolon DN. 2020.
Comprehensive fitness maps of Hsp90 show widespread environmental dependence. *Elife* **9**.
doi:10.7554/eLife.53810
Forsyth B, Torab P, Lee J-H, Malcom T, Wang T-H, Liao JC, Yang S, Kvam E, Puleo C, Wong PK.
2021. A Rapid Single-Cell Antimicrobial Susceptibility Testing Workflow for Bloodstream
Infections. *Biosensors* **11**. doi:10.3390/bios11080288
Fowler DM, Fields S. 2014. Deep mutational scanning: a new style of protein science. *Nat Methods*
11:801–807.
Fraley C, Raftery AE. 2003. Enhanced Model-Based Clustering, Density Estimation,and Discriminant
Analysis Software: MCLUST. *J Classification* **20**:263–286.
Geiler-Samerotte KA, Dion MF, Budnik BA, Wang SM, Hartl DL, Drummond DA. 2011. Misfolded
proteins impose a dosage-dependent fitness cost and trigger a cytosolic unfolded protein response in
yeast. *Proc Natl Acad Sci U S A* **108**:680–685.
Geiler-Samerotte KA, Li S, Lazaris C, Taylor A, Ziv N, Ramjeawan C, Paaby AB, Siegal ML. 2020.
Extent and context dependence of pleiotropy revealed by high-throughput single-cell phenotyping.
PLoS Biol **18**:e3000836.
Geiler-Samerotte KA, Zhu YO, Goulet BE, Hall DW, Siegal ML. 2016. Selection Transforms the
Landscape of Genetic Variation Interacting with Hsp90. *PLoS Biol* **14**:e2000465.
Gjini E, Wood KB. 2021. Price equation captures the role of drug interactions and collateral effects in the
evolution of multidrug resistance. *Elife* **10**. doi:10.7554/eLife.64851
Gomulkiewicz R, Kirkpatrick M. 1992. QUANTITATIVE GENETICS AND THE EVOLUTION OF
REACTION NORMS. *Evolution* **46**:390–411.
Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. 2017. The dynamics of molecular evolution
over 60,000 generations. *Nature* **551**:45–50.
Grier HE, Krailo MD, Tarbell NJ, Link MP, Fryer CJH, Pritchard DJ, Gebhardt MC, Dickman PS,
Perlman EJ, Meyers PA, Donaldson SS, Moore S, Rausen AR, Vietti TJ, Miser JS. 2003. Addition
of ifosfamide and etoposide to standard chemotherapy for Ewing's sarcoma and primitive
neuroectodermal tumor of bone. *N Engl J Med* **348**:694–701.
Haase SB. 2004. Cell cycle analysis of budding yeast using SYTOX Green. *Curr Protoc Cytom* **Chapter
7**:Unit 7.23.
Hall MD, Handley MD, Gottesman MM. 2009. Is resistance useless? Multidrug resistance and collateral
sensitivity. *Trends Pharmacol Sci* **30**:546–556.

Harris A, Wagner M, Du D, Raschka S, Nentwig L-M, Gohlke H, Smits SHJ, Luisi BF, Schmitt L. 2021.
Structure and efflux mechanism of the yeast pleiotropic drug resistance transporter Pdr5. *Nat*
Commun **12**:5254.
Herren CM, Baym M. 2022. Decreased thermal niche breadth as a trade-off of antibiotic resistance. *ISME*
J **16**:1843–1852.
Hill JA, O'Meara TR, Cowen LE. 2015. Fitness Trade-Offs Associated with the Evolution of Resistance
to Antifungal Drug Combinations. *Cell Rep* **10**:809–819.
Hinz A, Amado A, Kassen R, Bank C, Wong A. 2023. Unpredictability of the fitness effects of
antimicrobial resistance mutations across environments in Escherichia coli. *Mol Biol Evol* **41**.
doi:10.1101/2023.10.25.563906
Hsieh K, Mach KE, Zhang P, Liao JC, Wang T-H. 2022. Combating Antimicrobial Resistance via Single-
Cell Diagnostic Technologies Powered by Droplet Microfluidics. *Acc Chem Res* **55**:123–133.
Imamovic L, Ellabaan MMH, Dantas Machado AM, Citterio L, Wulff T, Molin S, Krogh Johansen H,
Sommer MOA. 2018. Drug-Driven Phenotypic Convergence Supports Rational Treatment Strategies
of Chronic Infections. *Cell* **172**:121–134.e14.
Iram S, Dolson E, Chiel J, Pelesko J, Krishnan N, Güngör Ö, Kuznets-Speck B, Deffner S, Ilker E, Scott
JG, Hinczewski M. 2020. Controlling the speed and trajectory of evolution with counterdiabatic
driving. *Nat Phys* **17**:135–142.
Iwasawa J, Maeda T, Shibai A, Kotani H, Kawada M, Furusawa C. 2022. Analysis of the evolution of
resistance to multiple antibiotics enables prediction of the Escherichia coli phenotype-based fitness
landscape. *PLoS Biol* **20**:e3001920.
Iyer KR, Robbins N, Cowen LE. 2022. The role of Candida albicans stress response pathways in
antifungal tolerance and resistance. *iScience* **25**:103953.
Jarosz DF, Lindquist S. 2010. Hsp90 and Environmental Stress Transform the Adaptive Value of Natural
Genetic Variation. *Science* **330**:1820–1824.
Jerison ER, Nguyen Ba AN, Desai MM, Kryazhimskiy S. 2020. Chance and necessity in the pleiotropic
consequences of adaptation for budding yeast. *Nat Ecol Evol* **4**:601–611.
Kaiser J. 2017. When less is more. *Science* **355**:1144–1146.
Kapitzky L, Beltrao P, Berens TJ, Gassner N, Zhou C, Wüster A, Wu J, Babu MM, Elledge SJ, Toczyski
D, Lokey RS, Krogan NJ. 2010. Cross-species chemogenomic profiling reveals evolutionarily
conserved drug mode of action. *Mol Syst Biol* **6**:451.
King ES, Maltas J, Weaver DT, Barker-Clarke R, Pelesko J, Dolson E, Scott JG. 2022. Fitness seascapes
are necessary for realistic modeling of the evolutionary response to drug therapy. *biorxiv*.
Kinsler G, Geiler-Samerotte K, Petrov DA. 2020. Fitness variation across subtle environmental
perturbations reveals local modularity and global pleiotropy of adaptation. *Elife* **9**.
doi:10.7554/eLife.61271
Kinsler G, Schmidlin K, Newell D, Eder R, Apodaca S, Lam G, Petrov D, Geiler-Samerotte K. 2023.
Extreme sensitivity of fitness to environmental conditions: Lessons from #1BigBatch. *J Mol Evol*
91:293–310.
Kolaczkowski M, van der Rest M, Cybularz-Kolaczkowska A, Soumillion JP, Konings WN, Goffeau A.
1996. Anticancer drugs, ionophoric peptides, and steroids as substrates of the yeast multidrug
transporter Pdr5p. *J Biol Chem* **271**:31543–31548.
Krishna A, Zere T, Mistry S, Ismaiel O, Stone H, Sacks LV, Weaver JL. 2022. Evaluation of a Sequential
Antibiotic Treatment Regimen of Ampicillin, Ciprofloxacin and Fosfomycin against Escherichia coli
CFT073 in the Hollow Fiber Infection Model Compared with Simultaneous Combination Treatment.
Antibiotics (Basel) **11**. doi:10.3390/antibiotics11121705
Ksiezopolska E, Schikora-Tamarit MÀ, Beyer R, Nunez-Rodriguez JC, Schüller C, Gabaldón T. 2021.
Narrow mutational signatures drive acquisition of multidrug resistance in the fungal pathogen
Candida glabrata. *Curr Biol* **31**:5314–5326.e10.
Kuchina A, Brettner LM, Paleologu L, Roco CM, Rosenberg AB, Carignano A, Kibler R, Hirano M,
DePaolo RW, Seelig G. 2021. Microbial single-cell RNA sequencing by split-pool barcoding.

Science **371**. doi:10.1126/science.aba5257

Larkins-Ford J, Degefu YN, Van N, Sokolov A, Aldridge BB. 2022. Design principles to assemble drug
combinations for effective tuberculosis therapy using interpretable pairwise drug response
measurements. *Cell Rep Med* **3**:100737.

Lässig M, Mustonen V, Walczak AM. 2017. Predicting evolution. *Nat Ecol Evol* **1**:77.

Lee JT, Coradini ALV, Shen A, Ehrenreich IM. 2019. Layers of Cryptic Genetic Variation Underlie a
Yeast Complex Trait. *Genetics* **211**:1469–1482.

Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. 2015. Quantitative evolutionary
dynamics using high-resolution lineage tracking. *Nature* **519**:181–186.

Li F, Salit ML, Levy SF. 2018. Unbiased Fitness Estimation of Pooled Barcode or Amplicon Sequencing
Studies. *Cell Syst* **7**:521–525.e4.

Li F, Tarkington J, Sherlock G. 2023. Fit-Seq2.0: An Improved Software for High-Throughput Fitness
Measurements Using Pooled Competition Assays. *J Mol Evol* **91**:334–344.

Li Y, Petrov DA, Sherlock G. 2019. Single nucleotide mapping of trait space reveals Pareto fronts that
constrain adaptation. *Nat Ecol Evol* **3**:1539–1551.

Li Y, Venkataram S, Agarwala A, Dunn B, Petrov DA, Sherlock G, Fisher DS. 2018. Hidden Complexity
of Yeast Adaptation under Simple Evolutionary Conditions. *Curr Biol* **28**:515–525.e6.

Logan A, Wolfe A, Williamson JC. 2022. Antifungal Resistance and the Role of New Therapeutic
Agents. *Curr Infect Dis Rep* **24**:105–116.

Lupetti A, Danesi R, Campa M, Del Tacca M, Kelly S. 2002. Molecular basis of resistance to azole
antifungals. *Trends Mol Med* **8**:76–81.

Maltas J, Wood KB. 2019. Pervasive and diverse collateral sensitivity profiles inform optimal strategies
to limit antibiotic resistance. *PLoS Biol* **17**:e3000515.

Martínez AA, Lang GI. 2023. Identifying Targets of Selection in Laboratory Evolution Experiments. *J
Mol Evol*. doi:10.1007/s00239-023-10096-2

McInnes L, Healy J, Melville J. 2018a. UMAP: Uniform Manifold Approximation and Projection for
Dimension Reduction. *arXiv [statML]*.

McInnes L, Healy J, Saul N, Großberger L. 2018b. UMAP: Uniform Manifold Approximation and
Projection. *J Open Source Softw* **3**:861.

Mehlhoff JD, Ostermeier M. 2023. Genes Vary Greatly in Their Propensity for Collateral Fitness Effects
of Mutations. *Mol Biol Evol* **40**. doi:10.1093/molbev/msad038

Mehlhoff JD, Stearns FW, Rohm D. 2020. Collateral fitness effects of mutations. *Proceedings of the*.

Melnikov SV, Stevens DL, Fu X, Kwok HS, Zhang J-T, Shen Y, Sabina J, Lee K, Lee H, Söll D. 2020.
Exploiting evolutionary trade-offs for posttreatment management of drug-resistant populations. *Proc
Natl Acad Sci U S A* **117**:17924–17931.

Mira PM, Crona K, Greene D, Meza JC, Sturmfels B, Barlow M. 2015. Rational design of antibiotic
treatment plans: a treatment strategy for managing evolution and reversing resistance. *PLoS One*
10:e0122283.

Mirkin B. 2011. Choosing the number of clusters. *Wiley Interdisc Rev Data Min Knowl Discov* **1**:252–
260.

Miyazaki H, Miyazaki Y, Geber A, Parkinson T, Hitchcock C, Falconer DJ, Ward DJ, Marsden K,
Bennett JE. 1998. Fluconazole resistance associated with drug efflux and increased transcription of a
drug transporter gene, PDH1, in Candida glabrata. *Antimicrob Agents Chemother* **42**:1695–1701.

Moye-Rowley WS. 2019. Multiple interfaces control activity of the Candida glabrata Pdr1 transcription
factor mediating azole drug resistance. *Curr Genet* **65**:103–108.

Nagasawa S, Kashima Y, Suzuki A, Suzuki Y. 2021. Single-cell and spatial analyses of cancer cells:
toward elucidating the molecular mechanisms of clonal evolution and drug resistance acquisition.
Inflamm Regen **41**:22.

Nichol D, Rutter J, Bryant C, Hujer AM, Lek S, Adams MD, Jeavons P, Anderson ARA, Bonomo RA,
Scott JG. 2019. Antibiotic collateral sensitivity is contingent on the repeatability of evolution. *Nat
Commun* **10**:334.

Nyhoegen C, Uecker H. 2023. Sequential antibiotic therapy in the laboratory and in the patient. *J R Soc Interface* **20**:20220793.

Ogbunugafor CB. 2022. The mutation effect reaction norm (mu-rn) highlights environmentally dependent mutation effects and epistatic interactions. *Evolution* **76**:37–48.

Osset-Trénor P, Pascual-Ahuir A, Proft M. 2023. Fungal Drug Response and Antimicrobial Resistance. *J Fungi (Basel)* **9**. doi:10.3390/jof9050565

Paaby AB, Rockman MV. 2013. The many faces of pleiotropy. *Trends Genet* **29**:66–73.

Paaby AB, White AG, Riccardi DD, Gunsalus KC, Piano F, Rockman MV. 2015. Wild worm embryogenesis harbors ubiquitous polygenic modifier variation. *Elife* **4**. doi:10.7554/eLife.09178

Padhy AA, Mavor D, Sahoo S, Bolon DNA, Mishra P. 2023. Systematic profiling of dominant ubiquitin variants reveals key functional nodes contributing to evolutionary selection. *Cell Rep* **42**:113064.

Pál C, Papp B, Lázár V. 2015. Collateral sensitivity of antibiotic-resistant microbes. *Trends Microbiol* **23**:401–407.

Palmer AC, Kishony R. 2014. Opposing effects of target overexpression reveal drug mechanisms. *Nat Commun* **5**:4296.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Müller A, Nothman J, Louppe G, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D, Brucher M, Perrot M, Duchesnay É. 2012. Scikit-learn: Machine Learning in Python. *arXiv [csLG]*.

Petti S, Reddy G, Desai MM. 2023. Inferring sparse structure in genotype-phenotype maps. *Genetics*. doi:10.1093/genetics/iyad127

Pinheiro F, Warsi O, Andersson DI, Lässig M. 2021. Metabolic fitness landscapes predict the evolution of antibiotic resistance. *Nat Ecol Evol* **5**:677–687.

Raymond B. 2019. Five rules for resistance management in the antibiotic apocalypse, a road map for integrated microbial management. *Evol Appl* **12**:1079–1091.

Read AF, Huijben S. 2009. Evolutionary biology and the avoidance of antimicrobial resistance. *Evol Appl* **2**:40–51.

Robbins N, Leach MD, Cowen LE. 2012. Lysine deacetylases Hda1 and Rpd3 regulate Hsp90 function thereby governing fungal drug resistance. *Cell Rep* **2**:878–888.

Rodrigues JV, Bershtein S, Li A, Lozovsky ER, Hartl DL, Shakhnovich EI. 2016. Biophysical principles predict fitness landscapes of drug resistance. *Proc Natl Acad Sci U S A* **113**:E1470–8.

Roemhild R, Linkevicius M, Andersson DI. 2020. Molecular mechanisms of collateral sensitivity to the antibiotic nitrofurantoin. *PLoS Biol* **18**:e3000612.

Roe SM, Prodromou C, O'Brien R, Ladbury JE, Piper PW, Pearl LH. 1999. Structural basis for inhibition of the Hsp90 molecular chaperone by the antitumor antibiotics radicicol and geldanamycin. *J Med Chem* **42**:260–266.

Rybak JM, Cuomo CA, Rogers PD. 2022. The molecular and genetic basis of antifungal resistance in the emerging fungal pathogen Candida auris. *Curr Opin Microbiol* **70**:102208.

Scarborough JA, McClure E, Anderson P, Dhawan A, Durmaz A, Lessnick SL, Hitomi M, Scott JG. 2020. Identifying States of Collateral Sensitivity during the Evolution of Therapeutic Resistance in Ewing's Sarcoma. *iScience* **23**:101293.

Schmidt F, Efferth T. 2016. Tumor Heterogeneity, Single-Cell Sequencing, and Drug Resistance. *Pharmaceuticals* **9**. doi:10.3390/ph9020033

Somarelli JA, Gardner H, Cannataro VL, Gunady EF, Boddy AM, Johnson NA, Fisk JN, Gaffney SG, Chuang JH, Li S, Ciccarelli FD, Panchenko AR, Megquier K, Kumar S, Dornburg A, DeGregori J, Townsend JP. 2020. Molecular Biology and Evolution of Cancer: From Discovery to Action. *Mol Biol Evol* **37**:320–326.

Sorgo AG, Heilmann CJ, Dekker HL, Bekker M, Brul S, de Koster CG, de Koning LJ, Klis FM. 2011. Effects of fluconazole on the secretome, the wall proteome, and wall integrity of the clinical fungus Candida albicans. *Eukaryot Cell* **10**:1071–1081.

Starr TN, Picton LK, Thornton JW. 2017. Alternative evolutionary histories in the sequence space of an ancient protein. *Nature* **549**:409–413.

Tanaka S, Tani M. 2018. Mannosylinositol phosphorylceramides and ergosterol coodinately maintain cell
wall integrity in the yeast Saccharomyces cerevisiae. *FEBS J* **285**:2405–2427.
Tan L, Chen L, Yang H, Jin B, Kim G, Im YJ. 2022. Structural basis for activation of fungal sterol
receptor Upc2 and azole resistance. *Nat Chem Biol* **18**:1253–1262.
Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, Gaut BS. 2012. The
molecular diversity of adaptive convergence. *Science* **335**:457–461.
Thomas DS, Cisneros LH, Anderson ARA, Maley CC. 2022. In Silico Investigations of Multi-Drug
Adaptive Therapy Protocols. *Cancers* **14**. doi:10.3390/cancers14112699
Uemura S, Moriguchi T. 2022. Pleiotropic roles of N-glycans for enzyme activities and stabilities of
MIPC synthases, Csh1 and Sur1/Csg1, in Saccharomyces cerevisiae. *Glycobiology* **32**:778–790.
Vasicek EM, Berkow EL, Flowers SA, Barker KS, Rogers PD. 2014. UPC2 is universally essential for
azole antifungal resistance in Candida albicans. *Eukaryot Cell* **13**:933–946.
Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, Ebel ER, Geiler-Samerotte K, Hérissant L, Blundell
JR, Levy SF, Fisher DS, Sherlock G, Petrov DA. 2016. Development of a Comprehensive Genotype-
to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell* **166**:1585–1596.e22.
Ventola CL. 2015. The antibiotic resistance crisis: part 1: causes and threats. *P T* **40**:277–283.
Vik Å, Rine J. 2001. Upc2p and Ecm22p, dual regulators of sterol biosynthesis in Saccharomyces
cerevisiae. *Mol Cell Biol* **21**:6395–6405.
Vu BG, Moye-Rowley WS. 2022. Nonidentical function of Upc2A binding sites in the Candida glabrata
CDR1 promoter. *Genetics* **222**. doi:10.1093/genetics/iyac135
Waller NJE, Cheung C-Y, Cook GM, McNeil MB. 2023. The evolution of antibiotic resistance is
associated with collateral drug phenotypes in Mycobacterium tuberculosis. *Nat Commun* **14**:1517.
Wang W-Y, Cai H-Q, Qu S-Y, Lin W-H, Liang C-C, Liu H, Xie Z-X, Yuan Y-J. 2022. Genomic
Variation-Mediating Fluconazole Resistance in Yeast. *Biomolecules* **12**. doi:10.3390/biom12060845
Wang X, Zhang H, Chen X. 2019. Drug resistance and combating drug resistance in cancer. *Cancer Drug*
Resist **2**:141–160.
Wang Y, Sang M, Feng L, Gragnoli C, Griffin C, Wu R. 2023. A pleiotropic–epistatic entangelement
model of drug response. *Drug Discov Today* 103790.
Ward JH. 1963. Hierarchical Grouping to Optimize an Objective Function. *J Am Stat Assoc* **58**:236–244.
Whitesell L, Santagata S, Mendillo ML, Lin NU, Proia DA, Lindquist S. 2014. HSP90 empowers
evolution of resistance to hormonal therapy in human breast cancer models. *Proc Natl Acad Sci U S*
A **111**:18297–18302.
Wilson BA, Garud NR, Feder AF, Assaf ZJ, Pennings PS. 2016. The population genetics of drug
resistance evolution in natural populations of viral, bacterial and eukaryotic pathogens. *Mol Ecol*
25:42–66.
Wortel MT, Agashe D, Bailey SF, Bank C, Bisschop K, Blankers T, Cairns J, Colizzi ES, Cusseddu D,
Desai MM, van Dijk B, Egas M, Ellers J, Groot AT, Heckel DG, Johnson ML, Kraaijeveld K, Krug
J, Laan L, Lässig M, Lind PA, Meijer J, Noble LM, Okasha S, Rainey PB, Rozen DE, Shitut S, Tans
SJ, Tenaillon O, Teotónio H, de Visser JAGM, Visser ME, Vroomans RMA, Werner GDA,
Wertheim B, Pennings PS. 2023. Towards evolutionary predictions: Current promises and
challenges. *Evol Appl* **16**:3–21.
Xie JL, Polvi EJ, Shekhar-Guturja T, Cowen LE. 2014. Elucidating drug resistance in human fungal
pathogens. *Future Microbiol* **9**:523–542.
Yadav A, Radhakrishnan A, Bhanot G, Sinha H. 2015. Differential regulation of antagonistic pleiotropy
in synthetic and natural populations suggests its role in adaptation. *G3* **5**:699–709.
Yang F, Scopel EFC, Li H, Sun L-L, Kawar N, Cao Y-B, Jiang Y-Y, Berman J. 2023. Antifungal
Tolerance and Resistance Emerge at Distinct Drug Concentrations and Rely upon Different
Aneuploid Chromosomes. *MBio* **14**:e0022723.
Yoon N, Krishnan N, Scott J. 2021. Theoretical modeling of collaterally sensitive drug cycles: shaping
heterogeneity to allow adaptive therapy. *J Math Biol* **83**:47.
Yoshida M, Reyes SG, Tsuda S, Horinouchi T, Furusawa C, Cronin L. 2017. Time-programmable drug

dosing allows the manipulation, suppression and reversal of antibiotic drug resistance in vitro. *Nat*
Commun **8**:15589.
Zhao L, Liu Z, Levy SF, Wu S. 2018. Bartender: a fast and accurate clustering algorithm to count barcode
reads. *Bioinformatics* **34**:739–747.
Zhao Q, Hautamaki V, Fränti P. 2008. Knee Point Detection in BIC for Detecting the Number of
ClustersAdvanced Concepts for Intelligent Vision Systems. Springer Berlin Heidelberg. pp. 664–
673.

1705

1706

1707