1 **Title:** Distinguishing mutants that resist drugs via different mechanisms by examining fitness tradeoffs
2 across hundreds of fluconazole-resistant yeast strains
3
4 **Keywords:** Fitness tradeoffs, drug resistance, DNA barcodes, fitness profiles, gene-by-environment
5 interactions, molecular mechanisms, genotype-phenotype map, azole resistance
6

7 **Authors:**
8 Schmidlin[1,2]*, Apodaca[1,2]*, Newell[1,2], Sastokas[1,2], Kinsler[3], Geiler-Samerotte[1,2]
9 * Co-first authors
10
11 **Affiliations:**
12 1 Biodesign Center for Mechanisms of Evolution, Arizona State University, Tempe, AZ
13 2 School of Life Sciences, Arizona State University, Tempe AZ
14 3 Department of Bioengineering, University of Pennsylvania, Philadelphia, PA
15
Corresponding Author:
kerry.samerotte@asu.edu

Abstract
There is growing interest in designing multidrug therapies that leverage tradeoffs to combat resistance.
Tradeoffs are common in evolution and occur when, for example, resistance to one drug results in sensitivity
to another. Major questions remain about the extent to which tradeoffs are reliable, specifically, whether
the mutants that provide resistance to a given drug all suffer similar tradeoffs. This question is difficult
because the drug-resistant mutants observed in the clinic, and even those evolved in controlled laboratory
settings, are often biased towards those that provide large fitness benefits. Thus, the mutations (and
mechanisms) that provide drug resistance may be more diverse than current data suggests. Here, we perform
evolution experiments utilizing lineage-tracking to capture a fuller spectrum of mutations that give yeast
cells a fitness advantage in fluconazole, a common antifungal drug. We then quantify fitness tradeoffs for
each of 774 evolved mutants across 12 environments, finding these mutants group into 6 classes with
characteristically different tradeoffs. Their unique tradeoffs may imply that each group of mutants affects
fitness through different underlying mechanisms. Some of the groupings we find are surprising. For
example, we find some mutants that resist single drugs do not resist their combination, and some mutants
to the same gene have different tradeoffs than others. These findings, on one hand, demonstrate the
difficulty in relying on consistent or intuitive tradeoffs when designing multidrug treatments. On the other
hand, by demonstrating that hundreds of adaptive mutations can be reduced to a few groups with
characteristic tradeoffs, our findings empower multidrug strategies that leverage tradeoffs to combat
resistance. Finally, by grouping mutants that likely affect fitness through similar underlying mechanisms,
our work guides efforts to map the phenotypic effects of mutation.

39
40
41
42

Introduction

How many different molecular mechanisms can a microbe exploit to adapt to a challenging environment? Answering this question is particularly urgent in the field of drug resistance because infectious populations are adapting to available drugs faster than new drugs are developed (Ventola 2015; Centers for Disease Control and Prevention (U.S.) 2019). Understanding the mechanistic basis of drug resistance can inform strategies for how to combine existing drugs in a way that prevents the evolution of resistance (Andersson and Hughes 2010; Melnikov et al. 2020; Pinheiro et al. 2021). For example, one strategy exposes an infectious population to one drug (Drug A) knowing that the mechanism of resistance to Drug A makes cells susceptible to Drug B (Hall et al. 2009; Pál et al. 2015; Baym et al. 2016; Roemhild et al. 2020). Problematically, these multi-drug strategies perform best when all mutants that resist Drug A have the same tradeoff in the second drug (**Figure 1A**). If there are multiple different mechanisms to resist Drug A, some of which lack this tradeoff, treatment strategies could fail (**Figure 1B**), and they sometimes do (Grier et al. 2003; Abel zur Wiesch et al. 2014; Wang et al. 2019; Scarborough et al. 2020).

Figure 1: A multidrug treatment strategy that relies on all mutants having the same tradeoffs. (A) All of the mutants that resist Drug A do so via a similar mechanism such that all are sensitive to Drug B. **(B)** There are multiple different types of mutants that resist Drug A, not all of which are sensitive to Drug B.

Laboratory experiments that have power to search for universal tradeoffs – where all the mutants that perform well in one environment perform poorly in another – often find there are mutants that violate trends or the absence of trends altogether (Hill et al. 2015; Nichol et al. 2019; Kinsler et al. 2020; Gjini and Wood 2021; Ardell and Kryazhimskiy 2021; Herren and Baym 2022). Another way to phrase this observation is to say that adaptive mutations often have collateral effects in environments other than the one in which they originally evolved (Pál et al. 2015). These effects, also referred to as pleiotropic effects, can be unpredictable and context dependent (Jerison et al. 2020; Geiler-Samerotte et al. 2020; Bakerlee et al. 2021; Chen et al. 2023). In simpler terms, some mutants that resist Drug A will suffer a tradeoff in Drug B, but others may suffer a tradeoff in Drug C. To sum, observations from many fields suggest that the mutations that provide a benefit in one environment do not always suffer similar tradeoffs. This begs questions about the extent of diversity among adaptive mutations: does each one suffer a unique set of tradeoffs or can many adaptive mutants be grouped by their common tradeoffs? If there are only a few

74   types of tradeoff present in a collection of adaptive mutations, multidrug treatments that target tradeoffs
75   may be more feasible.
76
The goal of the present study is to count how many different types of adaptive mutation, each type being
defined by its unique tradeoffs, exist in a population of drug-resistant yeast. This simple counting exercise
is surprisingly difficult. One reason why is that the mutations that provide the strongest fitness advantage
often dominate evolution. Thus, in the clinic, and in laboratory experiments, the same drug-resistant
mutations repeatedly emerge (Lupetti et al. 2002; Berkow and Lockhart 2017; Melnikov et al. 2020;
Ksiezopolska et al. 2021), potentially leading to the false impression that the mechanistic basis of
resistance to a particular drug, and the associated tradeoffs, are less varied than may be true. This problem
is amplified by the limitations of bulk DNA sequencing methods which often miss mutations that are
present in less than 10% of a population's cells (Good et al. 2017). A similar problem results from
strategies to disentangle adaptive from passenger mutations that rely on observing the adaptive ones
multiple times in multiple independent replicates (Martínez and Lang 2023). In order to design better
multidrug treatment strategies that thwart resistance, or to see if such strategies are feasible, we need
methods to survey a more complete set of mutations and mechanisms that can contribute to resistance.
90
Fortunately, single-cell and single-lineage DNA sequencing technologies are allowing us to more deeply
sample genetic diversity in evolving populations of microbes beyond the mutations that dominate
evolution (Schmidt and Efferth 2016). Here, we leverage a platform to perform massively-replicate
evolution experiments in yeast. This platform has been shown to reveal a fuller spectrum of mutations
underlying adaptation to a particular environment (Levy et al. 2015). One key to its success is that it uses
DNA barcodes to track all competing adaptive lineages, not just the ones that ultimately rise to
appreciable frequency. Another key feature is that it captures adaptive lineages before they accumulate
multiple mutations such that it is easy to pinpoint which mutation is adaptive. We apply this platform to
investigate mechanisms underlying resistance to a specific antifungal drug: fluconazole (FLU) (Wang et
al. 2022; Logan et al. 2022). Although serious fungal infections are most common in
immunocompromised individuals, their impact on global health is still striking, resulting in over 1.5
million deaths annually (Xie et al. 2014; Iyer et al. 2022). By focusing on mechanisms of FLU resistance,
we contribute to a growing literature about the tradeoffs that may be leveraged to design multidrug
antifungal treatment strategies (Cowen and Lindquist 2005; Hill et al. 2015; Ksiezopolska et al. 2021;
Iyer et al. 2022). However, our primary goal is more generic: we seek to explore the utility of a high-
throughput evolutionary approach to enumerate classes of drug resistant mutant and their associated
tradeoffs.
108
To enhance the diversity of drug resistant mutants in our experiment, we performed multiple laboratory
evolutions in a range of FLU concentrations and sometimes in combination with a second drug. We did
so because previous work has shown that different drug concentrations and combinations select for
different azole resistance mechanisms (Cowen and Lindquist 2005; Hill et al. 2015). Ultimately, we
obtained a large collection of 774 adaptive yeast strains. But how do we know whether we succeeded in
isolating diverse types of FLU-resistant mutants? Typical phenotyping methods, *e.g.,* quantifying
expression levels of drug export pumps (Miyazaki et al. 1998) or of the drug targets themselves (Palmer
and Kishony 2014), are low throughput and require some a priori knowledge of the phenotypes that may
be involved in drug resistance. Alternatively, many studies focus on identifying the genetic basis of

adaptation in order to glean insights about the underlying mechanisms of resistance (Tenaillon et al. 2012;
Cowen et al. 2014; Venkataram et al. 2016). However, genotyping lineages from barcoded pools is
technically challenging (Venkataram et al. 2016), and further, genotype does not necessarily predict
phenotype (Eguchi et al. 2019; Brettner et al. 2022b). For example, previous work using the same
barcoded evolution platform used here discovered that the mutations that provide an advantage under
glucose-limitation are in genes comprising a canonical glucose-sensing pathway (Venkataram et al.
2016). Yet despite this similarity at the genetic level, follow-up work showed that these mutants did not
experience the same tradeoffs when exposed to new environments (Li et al. 2018b; Kinsler et al. 2020).

Instead of trying to identify the phenotypic or even the genetic basis of adaptation, here we strive to
enumerate different classes of FLU-resistant mutants. We sort evolved FLU-resistant yeast strains into
classes based on whether they share similar tradeoffs across 12 different environments. Previous work
suggests that mutants with different fitness tradeoffs may affect fitness through different molecular
mechanisms (Rodrigues et al. 2016; Li et al. 2019; Pinheiro et al. 2021). Our work can thus be seen as
part of a growing push to flip the problem of mechanism on its head (Li et al. 2019; Kinsler et al. 2020;
Petti et al. 2023). Instead of using a mechanistic understanding to predict a microbe's fitness, here we use
how fitness varies across environments to enumerate classes of FLU-resistant mutant that likely affect
fitness via different mechanisms. Across our collection of 774 adaptive yeast lineages, we discovered at
least 6 distinct groups with characteristic tradeoffs. For example, we find some drug resistant mutants are
generally advantageous, while others are advantageous only in specific environments. And we find some
mutants that resist single drugs also resist combinations of those drugs, while others do not. By grouping
mutants with similar tradeoffs, we reduce the number of unique drug-resistant mutants from more than
can be easily phenotyped (774) to a manageable panel of 6 types for investigating the mechanistic basis of
resistance.

With regard to multidrug regimens that exploit tradeoffs (**Figure 1**), our finding of multiple mutant
classes with different tradeoffs suggests this may not be straightforward. The outlook is further
complicated by our finding that some classes of FLU-resistant mutant primarily emerge from evolution
experiments that did not contain FLU. This, as well as limits on our power to observe mutants with strong
tradeoffs, suggest there may be additional types of FLU-resistant mutant beyond those we sampled. These
observations suggest multidrug strategies that assume resistant mutants suffer consistent or common
tradeoffs will often fail.

On the other hand, nuanced strategies to forestall resistance that allow for multiple mutant types are
emerging (Maltas and Wood 2019; Gjini and Wood 2021; Wang et al. 2023). For example, one idea is to
apply a drug regimen that enriches for mutants that suffer a particular tradeoff before exploiting that
tradeoff (Iram et al. 2020). Another idea is to perform single-cell sequencing on infectious populations to
discover which classes of mutants are present (Nagasawa et al. 2021; Forsyth et al. 2021) and design
treatments specific to those (Maltas and Wood 2019; Aissa et al. 2021; Hsieh et al. 2022). Our findings
support that such ideas may be feasible by demonstrating that there are not as many unique fitness
tradeoffs as there are mutations.

More generally, our work – showing that 774 mutants fall into a much smaller number of groups –
contributes to growing literature suggesting that the phenotypic basis of adaptation is not as diverse as the

genetic basis (Kinsler et al. 2020; Iwasawa et al. 2022; Petti et al. 2023). This winnowing of diversity is
important: it may mean that evolutionary processes, for example, whether an infectious population will
adapt to resist a drug, are sometimes predictable (Rodrigues et al. 2016; Lässig et al. 2017; Kinsler et al.
2020; Yoon et al. 2021; King et al. 2022; Wortel et al. 2023).
166
Results
Barcoded evolution experiments uncover hundreds of yeast lineages with adaptive mutations
In order to create a sizable collection of drug-resistant mutants, we performed high-replicate evolution
experiments utilizing barcoded yeast (Levy et al. 2015; Li et al. 2018b; Boyer et al. 2021). This barcoding
system allows evolving hundreds of thousands of genetically identical yeast lineages together in a single
flask. Each lineage is tagged with a unique DNA barcode, which is a 26 base pair sequence of DNA
located within an artificial intron. Lineages with unique barcodes can be thought of as independent
replicates of an evolution experiment. This high-replicate system has the potential to generate many
different yeast lineages that differ by single adaptive mutations (Venkataram et al. 2016; Kinsler et al.
2020).
177
We performed a total of 12 barcoded evolution experiments, each of which started from the same pool of
approximately 300,000 barcoded yeast lineages (**Figure 2A – B; Figure S1**). These evolutions survey
how yeast cells adapt to different concentrations and combinations of two drugs: fluconazole (FLU) and
radicicol (RAD). FLU is a first line of defense against yeast infections, but over the past two decades
diverse resistant mutations have been identified (Bongomin et al. 2017; Rybak et al. 2022; Osset-Trénor
et al. 2023). Some earlier work suggested that FLU-resistant mutants are sensitive to the second drug we
study, radicicol (RAD) (Cowen and Lindquist 2005; Cowen et al. 2009), and more generally that RAD
can prevent the emergence of drug resistance in other systems (Whitesell et al. 2014). However, there are
some mutants that are cross resistant to both FLU and RAD (Hill et al. 2015), and the prominent
mechanism of resistance can differ with the intensity of selection and drug concentration (Cowen and
Lindquist 2005; Yang et al. 2023). We thus chose to evolve yeast to resist different concentrations and
combinations of FLU and RAD to generate a diverse pool of adaptive mutations comprising different
mechanisms of drug resistance.
191

Figure 2: An overview of the experimental design. (A) Yeast cells were barcoded to create 300,000 lineages. **(B)** These lineages were evolved in 12 different conditions (**Table 1**). **(C)** A small sample of evolved lineages was taken from each evolution experiment and sequenced. **(D)** These samples of evolved lineages were all pooled together with control strains representing the ancestral genotype. **(E)** The fitness of each lineage in this pool of ~21,000 was measured in each of the 12 evolution conditions via pooled fitness competitions. Fitness was measured by tracking changes in barcode frequencies over time relative to control strains. Two replicates per condition were performed. **(F)** The overall goal is to investigate the fitness tradeoffs for each evolved lineage. For example, the lineage depicted in dark blue has high fitness in some environments (HR, HF) but lower fitness in others (DMSO, ND). We were able to investigate fitness tradeoffs for 774 of the 21,000 lineages. We excluded lineages when we did not observe their associated barcode at least 500 times in all 12 environments. In other words, we only included lineages for which we obtained high quality fitness estimates in all 12 environments.

We evolved yeast to resist three different concentrations of either FLU and RAD for a total of six single-
drug conditions (**Table 1**). We also studied four conditions containing combinations of both drugs, as
well as two control conditions, for a total of 12 evolution experiments (**Table 1**). We chose to study
subclinical drug concentrations with the hope that no drug treatment would be strong enough to reduce
the population of yeast cells to only a handful of unique barcodes (**Figure S2**). We needed to maintain
barcode diversity in order to evolve a large number of unique lineages that each accumulate different
mutations.
223

Evolution Condition	Abbreviation	SYMBOL
4 µg/ml Fluconazole	Low Flu	Low Flu
6 µg/ml Fluconazole	Med Flu	Med Flu
8 µg/ml Fluconazole	High Flu	High Flu
5 µM Radicicol	Low Rad	Low Rad
15 µM Radicicol	Med Rad	Med Rad
20 µM Radicicol	High Rad	High Rad
5 µM Radicicol + 4 µg/ml Fluconazole	LRLF	LRLF
5 µM Radicicol + 8 µg/ml Fluconazole	LRHF	LRHF
15 µM Radicicol + 4 µg/ml Fluconazole	HRLF	HRLF
15 µM Radicicol +8 µg/ml Fluconazole	HRHF	HRHF
0.5% DMSO	DMSO	DMSO
No Drug	No Drug	No Drug

224
225 **Table 1:** A list of the environments included in this study and the symbol used to represent them in
226 subsequent figures.
227

With the goal of collecting adaptive lineages from each evolution experiment, we took samples from each
one after 3 - 6 growth/transfer cycles (**Figure S1**). This represents roughly 24 to 48 generations of growth
assuming 8 generations per growth/transfer cycle (Levy et al. 2015). We sampled early because previous
work using this barcoded evolution system demonstrated that the diversity of adaptive lineages peaks
after just a few dozen generations (Levy et al. 2015; Venkataram et al. 2016). We sampled about 2,000
cells from each evolution experiment except those 3 containing high concentrations of FLU from which
we sampled only 1,000 cells for a total of ~21,000 isolates (2,000 cells x 9 conditions + 1000 cells x 3
conditions) (**Figure 2C**).
236

Next, we measured the fitness of each isolate in each of the 12 evolution environments to quantify fitness
tradeoffs (e.g., whether mutants that perform well in one environment perform worse in another). This
process also indirectly screens isolates for adaptive mutations by comparing the fitness of each evolved
isolate to the ancestor of the evolution experiments (Venkataram et al. 2016). To do so, we pooled these
21,000 isolates and used this pool to initiate fitness competition experiments (**Figure 2D**). We competed

the pool against control strains, i.e., strains of the ancestral genotype that do not possess adaptive
mutations (Venkataram et al. 2016; Kinsler et al. 2020). We performed 24 such competitive fitness
experiments, 2 per each of the original 12 evolution conditions (**Figure 2E**). In each experiment, we
emulated the growth and transfer conditions of the original evolution experiments as precisely as possible,
tracking how barcode frequencies changed over 5 growth/transfer cycles (~40 generations). We used the
log-linear slope of this change, relative to the average slope for the control strains, to quantify relative
fitness.
249
We found that many barcoded lineages have higher fitness than the control strains in some conditions,
presumably because they possess adaptive mutations that improve their fitness in some conditions
(**Figure S3**). In fact, some of these adaptive lineages outcompeted the other lineages so quickly that it
posed a challenge. Barcodes pertaining to outcompeted lineages were often not present at high enough
coverage to track their fitness. We applied a conservative filter, preserving only 774 lineages with
barcodes that were observed >500 times in at least one replicate experiment per each of the 12 conditions.
The reason we required fitness measurements in all 12 conditions is that our goal is to examine each
lineage's fitness tradeoffs (**Figure 2F**) to see if different lineages have different tradeoffs. In order to
compare apples to apples, we need to measure each lineage's fitness in the same set of environments.
259
260   The 774 lineages we focus on are biased towards those that are reproducibly adaptive in multiple
261   environments we study. This is because lineages that have low fitness in a particular environment are
262   rarely observed >500 times in that environment (**Figure S4**). Despite this bias, we will go on to
263   demonstrate that there are different types of mutants with characteristically different fitness tradeoffs
264   present among these 774 lineages.
265
To provide additional evidence that these 774 barcoded yeast lineages indeed possess adaptive mutations,
we performed whole genome sequencing on a subset of 62 strains. Because we sampled these lineages
after only a few dozen generations of evolution, each lineage differs from the ancestor by just a few
mutations, making it easy to pinpoint the genetic basis of adaptation. Doing so revealed mutations that
have previously been shown to be adaptive in our evolution conditions (**Table S1**). For example, we
sequenced many FLU-resistant yeast lineages finding 35 with unique single nucleotide mutations in either
PDR1 or PDR3, and a few with mutations in SUR1 or UPC2, genes which have all been shown to
contribute to FLU resistance in previous work (Flowers et al. 2012; Vasicek et al. 2014; Tanaka and Tani
2018; Uemura and Moriguchi 2022; Vu and Moye-Rowley 2022). Similarly, lineages that have very high
fitness in RAD were found to possess single nucleotide mutations in genes associated with RAD
resistance, such as HDA1 (Robbins et al. 2012) and HSC82, which is the target of RAD (Roe et al. 1999).
We also observed several lineages with similar mutations to those observed in other studies using this
barcoded evolution regime, including mutations to IRA1, IRA2 and GPB2 (Venkataram et al. 2016;
Kinsler et al. 2020). Previous barcoded evolutions also observed that increases in ploidy were adaptive,
with 43% to 60% of cells becoming diploid during the course of evolution (Venkataram et al. 2016).
However, ploidy changes contributed less to adaptation in our experiment, with at most 9.4% of cells
becoming diploid by the time point when we sampled, but often less than 2% (**Table S2**).
283
284   In sum, we have created a diverse pool of 774 barcoded yeast lineages, most of which have a fitness
285   advantage in at least one of the conditions we study and are likely to possess a unique adaptive mutation.

286 The question we address for the rest of this study is to what extent these hundreds of mutant lineages
287 differ from one another in terms of their fitness tradeoffs and the mechanism/s underlying their fitness
288 advantages.
289
A unique mechanisms of FLU resistance emerges among mutants isolated in RAD evolutions
The majority of the 774 adaptive lineages that we study have higher fitness than the ancestral strains in
not one, but often in several drug conditions. This suggests that pleiotropy, and in particular cross-
resistance, is prevalent among the lineages we study. But not all lineages show the same patterns of cross
resistance (**Figure 3**). For example, the 100 most fit lineages in our highest concentration of fluconazole
are also beneficial in our highest concentration of radicicol (**Figure 3A**; leftmost two boxplots). As
expected, these 100 lineages also have high fitness in conditions where high concentrations of FLU and
RAD are combined (**Figure 3A**; third boxplot). And these 100 most-fit lineages in FLU lose their fitness
advantage in conditions where no drug is present (**Figure 3A**; rightmost boxplot).
299
Given their high fitness in conditions containing FLU, it seems likely that these 100 mutants originated
from evolution experiments containing FLU. We can trace every lineage back to the evolution
experiment/s it originated from because we sequenced the lineages we sampled from each evolution
experiment before pooling all 21,000 (**Figure 2C**). As we expected, these 100 best performing lineages in
high FLU largely originate from evolution experiments containing FLU (**Figure 3B**). Given that these
lineages have no fitness advantage in conditions containing no drug, it is also unsurprising that they are
underrepresented in evolution experiments lacking RAD and FLU (**Figure 3B**).
307
It might be tempting to generalize that most mutations that provide drug resistance are not beneficial in
environments without drugs. Afterall, we show this is true for 100 independent lineages (**Figure 3A**).
Further, many previous studies find a similar pattern, whereby drug resistant mutants often do not have
high fitness in the absence of drug (Andersson and Hughes 2010; Basra et al. 2018; Allen et al. 2019;
Melnikov et al. 2020), such that treatment strategies have emerged that cycle patients between drug and
no drug states, albeit with mixed success (Baker et al. 2018; Raymond 2019; Wang et al. 2019; Algazi et
al. 2020). However, this type of generalization is not supported by our data. We find that drug resistance
can sometimes come with an advantage, rather than a cost, in the absence of a drug (**Figure 3C**). The top
100 most fit mutants in our highest concentration of RAD provide a fitness advantage in high RAD, high
FLU, as well as in environments with no drug (**Figure 3C**). These observations suggest that there are at
least two different mechanisms by which to resist FLU that result in different tradeoffs in other
environments (**Figure 3A vs. 3C**).
320
321 Intriguingly, these FLU-resistant lineages that maintain their fitness advantage in the absence of drug
322 (**Figure 3C**) mainly originate from evolution experiments performed in conditions lacking FLU (**Figure
323 3D**). This highlights how the potential mechanisms by which a microbe can resist a drug may be more
324 varied than is often believed. Typically, one doesn't search for FLU-resistant mutants by evolving yeast
325 to resist RAD. Thus, typical studies might miss this unique class of FLU-resistant mutants.
326
In sum, there appear to be at least two different types of mutants present among our collection of 774
adaptive yeast lineages. One group has almost equally high fitness in RAD and FLU but has no fitness
advantage over the ancestral strain in conditions without either drug (**Figure 3A & 3E**). Another group is

defined by very high fitness in RAD, moderately high fitness in FLU and moderately high fitness in
conditions without either drug (**Figure 3C & 3E**). When comparing fitness in RAD vs. FLU across all
774 lineages, not only the top 100 best performing in each drug, we see some evidence that they largely
fall into the two main categories highlighted in **figures 3A and 3C** (**Figure 3E**). Thus, it might be
tempting to conclude that there are two different types of FLU-resistant mutant in our dataset. However,
sorting mutants into groups using a pairwise correlation plot (**Figure 3E**) excludes data from ten of our
twelve environments.
337

338
Figure 3: Two different classes of FLU-resistant mutants with unique tradeoffs. (A) This panel describes the 100
mutant lineages with the highest fitness relative to the control strains in the high FLU environment (8μg/ml FLU).
The vertical axis depicts the fitness advantages (log-linear slopes relative to control strains) for these 100 strains in
four selected environments, including the high FLU environment (boxed). Boxplots summarize the distribution across
all 100 lineages for each environment, displaying the median (center line), interquartile range (IQR) (upper and lower
hinges), and highest value within $1.5 \times$ IQR (whiskers). **(B)** The 100 lineages with highest fitness in high FLU were
most often sampled from evolution experiments in which FLU was present. In this pie-chart, colors correspond to the
evolution conditions listed in **table 1** and the blue outer ring highlights evolution conditions that contain FLU. The
size of each slice of pie represents the relative frequency with which these 100 lineages were found in each evolution
experiment. **(C)** Similar to panel A, this panel describes the 100 mutant lineages with the highest fitness relative to
the control strains in the high RAD environment (20uM Rad). **(D)** The 100 lineages with highest fitness in high RAD
were most often sampled from evolution experiments that did not contain FLU. **(E)** A pairwise correlation plot
showing that all 774 mutants, not just the two groups of 100 depicted in panels A and C, to some extent fall into two
groups defined by their fitness in high FLU and high RAD. The contours (black curves) were generated using kernel
density estimation with bins = 7. These contours describe the density of the underlying data, which is concentrated
into two clusters defined by the two smallest black circles. The 100 mutants with highest fitness in high FLU are blue,
highest fitness in high RAD are red, and the seven that overlap between the two aforementioned categories are black.
356
357

A strategy to differentiate classes of drug-resistant mutants with different tradeoffs

The observation of two distinct types of adaptive mutants (**Figure 3**) made us wonder whether there were additional unique types of FLU-resistant mutants with their own characteristic tradeoffs. This is difficult to tell by using pairwise correlation like that in **Figure 3E** because we are not studying pairs of conditions, as is somewhat common when looking for tradeoffs to leverage in multidrug therapies (Scarborough et al. 2020; Melnikov et al. 2020; Ardell and Kryazhimskiy 2021; Larkins-Ford et al. 2022). Instead, we have collected fitness data from 12 conditions to yield a more comprehensive set of gene-by-environment interactions for each mutant. This type of data, describing how a particular genotype responds to environmental change, is sometimes called a 'reaction norm' and can inform quantitative genetic models of how selection operates in fluctuating environments (Gomulkiewicz and Kirkpatrick 1992; Ogbunugafor 2022) and how much pleiotropy exists in nature (Yadav et al. 2015). More recent studies refer to the changing performance of a genotype across environments as a 'fitness profile' or in aggregate, a 'fitness seascape,' and suggest these type of dynamic measurements are the key to designing effective multi-drug treatments (King et al. 2022) and to predicting evolution (Lässig et al. 2017; Kinsler et al. 2020; Cairns et al. 2022; Iwasawa et al. 2022; Chen et al. 2023). And when the environments studied represent different drugs, these types of data are often referred to as "collateral sensitivity profiles" a term chosen to convey how resistance to one drug can have "collateral" effects on performance in other drugs (Pál et al. 2015; Maltas and Wood 2019; Gjini and Wood 2021). Despite the wide interest in this type of fitness data, it is technically challenging to generate, thus many previous studies of fitness profiles focus on a much smaller number of isolates (Imamovic et al. 2018; Nichol et al. 2019; Maltas and Wood 2019), sometimes with variation restricted to a single gene (Mira et al. 2015; King et al. 2022), or evolved in response to a single selection pressure (Li et al. 2018b; Kinsler et al. 2020). Here, we have generated fitness profiles for a large and diverse group of drug-resistant strains using the power of DNA barcodes. Now we seek to understand whether these mutants fall into distinct classes that each have characteristic fitness profiles (i.e., characteristic tradeoffs, characteristic reaction norms, or characteristic gene-by-environment interactions).

To address this question, we perform dimensional reduction, clustering mutants with fitness profiles that have a similar shape. It is in theory possible for all mutants to have similar profiles, perhaps implying they all affect fitness through similar underlying mechanisms (**Figure 4A**). However, the disparity reported in **Figure 3** suggests otherwise. It's also possible that every mutant will have a different profile. This could happen if each mutant affects different molecular-level phenotypes that underlie its drug resistance (**Figure 4B**). But previous work suggests that the phenotypic basis of adaptation is less diverse than the genotypic basis. This is important because it means that evolutionary outcomes are more predictable at the level of phenotype (Kinsler et al. 2020; Brettner et al. 2022b; Iwasawa et al. 2022). Even so, every mutant could have a slightly different fitness profile if each affects the same handful of molecular-level phenotypes but to relatively different degrees. This would allow every mutant to have a unique response to environmental change, without requiring that there be as many unique molecular mechanisms underlying drug resistance as there are mutants. A final possibility is that there exist discrete classes of drug-resistant mutants with characteristic tradeoffs (**Figure 4C**). This might imply that each class of mutants provides drug resistance via a different molecular mechanism, or a different set of mechanisms. In sum, our endeavor to enumerate mutants with different fitness profiles speaks to general questions about the extent of pleiotropy in the genotype-phenotype-fitness map (Boyle et al. 2017; Geiler-Samerotte et al. 2020; Bakerlee et al. 2021; Chen et al. 2023), the extent to which fitness tradeoffs are

universal (Andersson and Hughes 2010; Li et al. 2019; Herren and Baym 2022), and relatedly, the
predictability of evolution and potential for controlling evolutionary processes (Lässig et al. 2017; Iram et
al. 2020; Kinsler et al. 2020; King et al. 2022; Petti et al. 2023).

To see whether there are distinct classes of adaptive mutants among our drug-resistant yeast lineages, we
applied uniform manifold approximation and projection (UMAP) (McInnes et al. 2018a) to fitness
measurements for 774 yeast strains across all 12 environments. This method places mutants with similar
fitness profiles near each other in two-dimensional space. As might be expected, it largely places mutants
in each of the two categories described in **Figure 3** far apart, with drug-resistant mutants that lose their
benefit in the absence of drug in the top half of the graph, and those that maintain a benefit in the bottom
half (**Figure 4D and Figure S5**).

Beyond the obvious divide between the top and bottom clusters of mutants on the UMAP, we used a
gaussian mixture model (Fraley and Raftery 2003) to identify clusters. A common problem in this type of
analysis is the risk of dividing the data into clusters based on variation that represents measurement noise
rather than reproducible differences between mutants (Zhao et al. 2008; Mirkin 2011). One way we
avoided this was by ceasing to split out additional clusters before doing so no longer improved model
performance (**Figure S6**). The model we chose consists of seven clusters, including one pertaining to the
control strains, and six others pertaining to different classes of adaptive mutant (**Figure 4D**). We
investigated whether these clusters capture reproducible differences between mutants, rather than
measurement noise, by reducing the amount of noise in our data and asking if the same clusters are still
present. To do so, we reduced our collection of adaptive lineages from 774 to 617 by requiring 5,000
rather than 500 reads per lineage in order to infer fitness. This procedure reduced noise; the Pearson
correlation across replicate experiments improved from 0.756% to 0.813%. Despite this reduction in
variation, these 617 lineages cluster into the same six groups (plus a seventh pertaining to the control
strains) as do the original 774 (**Figure S7**). The groupings are also preserved when we perform alternate
methods for dimensionality reduction while adhering to a seven cluster model (**Figure S8**). Each of the
six clusters of adaptive mutants that we identify has a characteristic fitness profile (**Figure 4E**). In any
given environment, the fitness advantages of the mutants within each cluster are often very similar to one
another and often significantly different from other clusters (**Figure 4E**). Our follow-up investigations
provide additional evidence that these clusters of adaptive mutants each have characteristically different
tradeoffs, suggesting they affect fitness via different molecular mechanisms.

437
438
Figure 4: Clustering evolved lineages with similar fitness profiles. (A-C) Simulated data showing potential
fitness profiles when (**A**) all mutants have similar responses to environmental change and thus a similar fitness
profile, (**B**) every mutant has a different profile (five unique profiles are highlighted in color), or (**C**) every mutant
has one of a small number of unique profiles (two unique profiles are depicted). (**D**) Every point in this plot
represents one of the barcoded lineages colored by cluster; clusters were identified using a gaussian mixture model.
The 774 adaptive lineages cluster into 6 groups based on variation in their fitness profiles; the control lineages
cluster separately into the leftmost cluster in light green. (**E**) The fitness profiles of each cluster of adaptive lineages.
Boxplots summarize the distribution across all lineages within each cluster in each environment, displaying the
median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within $1.5 \times$ IQR
(whiskers).
449
A group of mutants with distinct genotypes are primarily resistant to low concentrations of FLU
The upper three clusters of mutants on the UMAP (**Figure 4D**) are all similar in that they have elevated
fitness in at least one FLU-containing environment but ancestor-like fitness in the absence of drug
(**Figure 4E**; upper three profiles). Despite these similarities, there are major differences between these
three groups of mutant lineages, both at the level of genotype and fitness profile (**Figure 5**). For example,

in cluster 1 (depicted in purple in **Figures 4 and 5**), the 3 sequenced lineages have single nucleotide
mutations to either SUR1 or UPC2 (**Figure 5A**). But in clusters 2 and 3 (depicted in blue and orange in
Figures 4 and 5), 35/36 sequenced lineages have unique single nucleotide mutations to one of two genes
associated with 'Pleiotropic Drug Resistance' (PDR1 or PDR3).
459
PDR1 and PDR3 are transcription factors that are well known to contribute to fluconazole resistance
through increased transcription of a drug pump (PDR5) that removes FLU from cells (Fardeau et al. 2007;
Osset-Trénor et al. 2023). However, SUR1 and UPC2 are less commonly mentioned in literature
pertaining to FLU resistance, and have different functions within the cell as compared to PDR1 and PDR3
(Kapitzky et al. 2010; Hill et al. 2015). SUR1 converts inositol phosphorylceramide to mannosylinositol
phosphorylceramide, which is a component of the plasma membrane (Uemura and Moriguchi 2022).
Similarly, UPC2 is a transcription factor with a key role in activating the ergosterol biosynthesis genes,
which contribute to membrane formation (Vik and Rine 2001; Tan et al. 2022). The presence of adaptive
mutations in genes involved in membrane synthesis is consistent with fluconazole's disruptive effect on
membranes (Sorgo et al. 2011).
470
Interestingly, the lineages with mutations to UPC2 and SUR1, and the unsequenced lineages in the same
cluster, do not consistently have cross resistance in RAD (**Figure 5B**; cluster 1). Oppositely, lineages
with mutations to PDR1 or PDR3, and the unsequenced lineages in the same clusters, are uniformly cross
resistant to RAD (**Figure 5B**; clusters 2 and 3). Perhaps this cross resistance is reflective of the fact that
the drug efflux pump that PDR1/3 regulates (PDR5) can transport a wide range of drugs and molecules
out of yeast cells (Kolaczkowski et al. 1996; Harris et al. 2021). Overall, the targets of adaptation in
cluster 1 have disparate functions within the cell as compared to the targets of adaptation in clusters 2 and
3. This may suggest that the mutants in cluster 1 confer FLU resistance via a different mechanism than
clusters 2 and 3.
480
The lineages in cluster 1 have additional important differences from clusters 2 and 3. The lineages in
cluster 1 perform best in the lowest concentration of FLU and have decreasing gains as the concentration
of FLU rises (**Figure 5C**). In fact, about 15% of these mutant lineages perform worse than their ancestor
in the highest concentration of FLU, suggesting the very mutations that provide resistance to low FLU are
costly in higher concentrations of the same drug. The mutants in clusters 2 and 3 show the opposite trend
from those in cluster 1: they perform best in the highest concentration of FLU and have reduced gains in
lower concentrations (**Figure 5C**). These findings provide additional evidence that a distinct mechanism
of FLU resistance distinguishes cluster 1 from clusters 2 and 3. The implication that different resistance
mechanisms will dominate evolution in slightly different concentrations of the same drug highlights the
complexity of adaptation and the need to more deeply understand the diversity of potential adaptive
mechanisms before designing treatment strategies (Berman and Krysan 2020; Yang et al. 2023).
492

493
494
Figure 5: Evolved lineages comprising cluster 1 have different genotypes and phenotypes from neighboring
clusters. (A) The three clusters on the top half of the UMAP differ in their targets of adaptation with cluster 1 being
unique in that it does not contain mutations to PDR1 or PDR3. **(B)** Evolved lineages comprising cluster 1 do not
have consistent fitness advantages in conditions containing RAD, while lineages comprising clusters 2 and 3 are
uniformly adaptive in RAD. Boxplots summarize the distribution across all lineages within each cluster in each
environment, displaying the median (center line), interquartile range (IQR) (upper and lower hinges), and highest
value within $1.5 \times$ IQR (whiskers). **(C)** Lineages comprising cluster 1 are most fit in low concentrations of FLU, and
this advantage dwindles as the FLU concentration increases. Lineages comprising clusters 2 and 3 show the opposite
trend.
504

Two groups of mutant lineages possessing similar adaptive mutations differ in sensitivity to RAD

While cluster 1 appears fairly different from its neighbors, it is not immediately obvious why the mutant lineages in clusters 2 and 3 are placed into separate groups. For one, the mutants in each cluster have fitness profiles with a very similar shape (**Figure 4E & 6A**). The sequenced lineages in each of these clusters also possess mutations to the same genes: PDR1 and PDR3 (**Figure 5A**). And finally, the lineages in each cluster originate from similar evolution experiments, largely those containing FLU (**Figure 6A**; pie charts). These observations made us wonder whether the difference between cluster 2 and 3 arose entirely because the mutants in cluster 3 have stronger effects than those in cluster 2 (**Figure 6A**; the solid blue line is above the solid orange line). In other words, we wondered whether the mutant lineages in clusters 2 and 3 affect fitness via the same mechanism, but to different degrees. To investigate this idea, we normalized all fitness profiles to have the same height on the vertical axis; this does not affect their shape (**Figure 6A**; dotted lines). Then we re-clustered and asked whether mutants pertaining to the original clusters 2 and 3 were now merged into a single cluster.

Normalizing in this way did not radically alter the UMAP, which still contains largely the same 6 clusters of mutants (**Figure S9**). Clusters 2 and 3, containing lineages with mutations to PDR1 or PDR 3, experienced the largest changes with 37% of mutants switching from one of these two groups to the other. The new clusters 2 and 3 now differ in the shape of their fitness profiles, whereby slight differences that existed between the original fitness profiles are exaggerated (**Figure 6B**). For example, mutants in cluster 3 perform better in high and medium concentrations of RAD (**Figure 6B**; line plot). This difference in fitness is reflected in the evolution experiments, with more mutant lineages in cluster 3 originating from the evolutions performed in RAD (**Figure 6B**; pie charts). Though cluster 3 mutants tend to have stronger RAD resistance, they tend to have reduced fitness in conditions containing neither FLU nor RAD as compared to cluster 2 lineages (**Figure 6B**; line plot). In sum, the differences between lineages in clusters 2 and 3 were not resolved upon normalizing fitness profiles to reduce magnitude differences, instead they were made more apparent (**Figure 6**). These differences do not appear to be random because they persist across experiments. For example, cluster 3 mutants are more fit in both medium and high RAD environments (**Figure 6B**; line plot) and were more often isolated from evolutions containing RAD (**Figure 6B**; pie charts). These observations beg a question: how can different mutations to the same gene affect fitness via different molecular mechanisms?

Asking this question forces us to consider what we mean by "mechanism." The mechanism by which mutations to PDR1 and PDR3 affect FLU resistance is well established: they increase transcription of an efflux pump that removes FLU from cells (Moye-Rowley 2019; Buechel and Pinkett 2020; Osset-Trénor et al. 2023). But if this is the only molecular-level effect of mutations to these genes, it is difficult to reconcile why PDR mutants fall into two distinct clusters with differently shaped fitness profiles. Others have also recently observed that mutants to PDR1 do not all behave the same way when exposed to novel drugs or changes in pH (Chen et al. 2023). This phenomenon is not reserved to PDR mutants, as adaptive missense mutations to another gene, IRA1, also do not share similarly shaped fitness profiles either (Kinsler et al. 2020). One explanation may be that, while all adaptive mutations within the same gene improve fitness via the same mechanism, not all mutants suffer the same costs. For example, perhaps the adaptive PDR mutations in cluster 2 cause misfolding of the PDR protein, resulting in lower fitness in RAD because this drug inhibits a chaperone that helps proteins to fold. In this case, it might be more correct to say that each of our six clusters affects fitness through a different, but potentially overlapping,

549 suite of mechanisms (Wang et al. 2023). Previous work demonstrating that mutations commonly affect
550 multiple traits supports this broader view of the mechanistic differences between clusters (Paaby and
551 Rockman 2013; Boyle et al. 2017; Geiler-Samerotte et al. 2020; Kinsler et al. 2020).
552
Alternatively, perhaps not all adaptive mutations to PDR improve fitness via the same mechanism. PDR1
and PDR3 regulate transcription of YOR1 and SNQ2 as well as PDR5, and maybe the different clusters
we observe represent mutants that upregulate one of these downstream targets more than the other (Osset-
Trénor et al. 2023). Or, the mutants in each cluster might harbor different aneuploidies or small, difficult
to sequence chromosomal insertions or deletions that affect fitness. We leave identification of the precise
mechanisms that differentiate these clusters for future work. Here, using the example of PDR mutants, we
showcase how genotype may not predict fitness tradeoffs, suggesting there is more to learn about the
mechanisms underlying FLU resistance.

566
Figure 6: Evolved lineages in clusters 2 and 3 have characteristic differences despite similarities at the genetic
level. (**A**) This panel shows the similarities between clusters 2 and 3. The upper right inset displays the same UMAP
from **Figure 4D** with only clusters 2 and 3 highlighted and with lineages possessing mutations to the PDR genes
depicted as blue diamonds. The line plot displays the same fitness profiles for clusters 2 and 3 as **Figure 4E**, plotting
the average fitness for each cluster in each environment and a 95% confidence interval. Dotted lines represent the
same data, normalized such that every lineage has an average fitness of 0 across all environments. These line plots
show that the fitness profiles for clusters 2 and 3 have a very similar shape. Pie charts display the relative frequency
with which lineages in clusters 2 and 3 were sampled from each of the 12 evolution conditions, colors match those
in the horizontal axis of the line plot and **table 1**. (**B**) This panel shows the differences between the new clusters 2
and 3 created after all fitness profiles were normalized to eliminate magnitude differences. The upper right inset
displays a new UMAP (also see **Figure S9**) that summarizes variation in fitness profiles after each profile was
normalized by setting its average fitness across all environments to 0. The line plot displays the fitness profiles for
the new clusters 2 and 3, which look different from those in panel A because 37% of mutants in the original clusters
2 and 3 switched identity from 2 to 3 or vice versa. The new clusters 2 and 3 are depicted in slightly different shades
of blue and orange to reflect that these are not the same groupings as those depicted in **Figure 4**. Pie charts display
the relative frequency with which lineages in new clusters 2 and 3 were sampled from each of the 12 evolution
conditions, colors match those in the horizontal axis of the line plot and **table 1**.

One group of RAD resistant mutants does not respond as expected to drug combinations

Though the three clusters of mutants on the bottom half of the UMAP are all advantageous in RAD and in conditions without any drug (**Figure 4E**; lower three plots), they differ in their fitness in conditions containing FLU. For example, the cluster of yeast lineages highlighted in green (cluster 4 in **Figures 4 and 7A**) is unique in that it has a slight advantage in the HRLF environment (**Figure 7B**). We found it especially strange that the neighboring cluster 5 does not also have a fitness advantage in this condition. Mutants in cluster 5 have a slight advantage in the LF condition, and a big advantage in the HR condition, thus we expect them to have at least some fitness advantage in the condition where these two drugs are combined (HRLF), but they do not (**Figure 7B**). The same is true for the combination of LRLF: cluster 5 mutants have an advantage in both single drug conditions which is lost when the drugs are combined (**Figure S10**). However, the mutants in cluster 4 (green) exhibit no such sensitivity to combined treatment. They have a slight advantage in all of the aforementioned single drug conditions, which is preserved in the relevant multidrug conditions (**Figure 7B & Figure S10**). To obtain an independent measure of the fitness of cluster 4 vs. cluster 5 lineages in these multidrug conditions, we asked from where the lineages in each cluster originate. About 10% of cluster 4 lineages originated from the HRLF evolution, while almost none of the lineages in cluster 5 came from this experiment, confirming that cluster 5 lineages are uniquely sensitive to this multidrug environment (**Figure 7C**).

The different fitness profiles of mutants in cluster 4 versus 5 (**Figure 7B & Figure S10**) might imply that they have different growth phenotypes. We performed a follow-up experiment that supports this observation. We asked whether there are differences in the growth phenotypes of cluster 4 versus 5 mutants by measuring a growth curve for the lineage we were able to isolate and sequence from cluster 5, comparing it to a growth curve from a cluster 4 lineage (**Figure 7D**). Indeed, the selected mutants in cluster 4 and 5 appear to have markedly different growth curves in some conditions (**Figure S10**). The growth differences echo those we see in the fitness data. For example, the cluster 5 mutant has a lower maximum growth rate in the HRLF multidrug condition, corresponding with its lower fitness in this condition (**Figure 7B & 7D**). The cluster 5 mutant also reaches a lower maximum cell density in the LRLF multidrug condition and has lower fitness in this condition (**Figure S10**). However, the growth curves and fitness of cluster 4 and 5 mutants are more similar in LF, LR and HR single drug conditions. The observation of reproducible growth differences between a cluster 4 and cluster 5 mutant provides evidence that our clustering approach is effective at separating mutants with different properties.

616
Figure 7: Evolved lineages in cluster 4 and 5 differ in response to combined drugs. (A) Adjacent clusters 4 and 5
each contain a small number of sequenced isolates depicted as diamonds; diamond colors correspond to the genes
containing adaptive mutations in each sequenced isolate. **(B)** Cluster 5 (red) has an unexpected fitness disadvantage
in the HRLF multidrug environment relative to cluster 4 (green), given that cluster 5 lineages do not have a fitness
disadvantage in the relevant single drug environments. Boxplots summarize the distribution across all lineages within
each cluster in each environment, displaying the median (center line), interquartile range (IQR) (upper and lower
hinges), and highest value within $1.5 \times$ IQR (whiskers). **(C)** Pie charts display the relative frequency with which
lineages in each cluster were sampled from each of the 12 evolution conditions, colors match those in table 1. **(D)** The
maximum exponential growth rate for a single lineage isolated from each of clusters 4 (green) and 5 (red), relative to
the ancestor. The growth rate of each lineage in each condition was measured twice by measuring changes in optical
density over time. Tested lineage from cluster 4 (in green) has a mutation to GBP2 (S317T) while the lineage from
cluster 5 (in red) has mutation to HDA1 (S600S).

One group of RAD resistant mutants is exceptionally adaptive in conditions without drug
One group of mutants in the lower half of the UMAP (cluster 6 in **Figure 8A**) appears distinct from the
other two in that it has the largest fitness advantage in conditions lacking any drug (**Figure 8B**). This
might imply that cluster 6 lineages rose to high frequency during our evolution experiments in
environments without either drug, specifically the "no drug" and "DMSO" control conditions. Indeed, this
is what we observe: over 50% of the lineages in cluster 6 were sampled from one of these two evolution
experiments (**Figure 8A**). On the contrary, the other clusters in the lower half of the UMAP consist
mainly of lineages sampled from one of the RAD evolutions (**Figure 7C**). Since our fitness experiments
were performed independently of the evolution experiments, this provides two independent pieces of
evidence suggesting that lineages in cluster 6 are defined by their superior performance in conditions
lacking any drug.

In line with the success of cluster 6 mutants in no drug conditions, the five sequenced mutants in this
cluster include three that have mutations to IRA1, which was the most common target of adaptation in
another evolution experiment in the condition we call "no drug" (**Figure 8A**) (Venkataram et al. 2016). In
that experiment, and in our no drug experiment, mutations to IRA1 result in a greater fitness advantage
than mutations to its paralog, IRA2, or mutations to other negative regulators of the RAS/PKA pathway
such as GPB2 (**Figure 8**). Previous work showed that sometimes IRA1 mutants have very strong
tradeoffs, for example, they become extremely maladaptive in environments containing salt or benomyl
(Kinsler et al. 2020). We do not observe this to be the case for either FLU or RAD. In fact, we observe
that cluster 6 mutants, including those in IRA1, maintain a fitness advantage in our highest concentration
of both drugs (**Figure 4**), being more fit in high FLU than mutants in either of the other clusters in the
lower half of the UMAP (**Figure 8B**). However, cluster 6 mutants are unique in that they lose their fitness
advantage in the lowest concentration of FLU (**Figure 8B**). Being singularly sensitive to a low
concentration of drug seems unusual, so much so that when this was observed previously for IRA1
mutants the authors added a note about the possibility of a technical error (Kinsler et al. 2020). Our results
suggest that there is indeed something uniquely treacherous about the low fluconazole environment, at
least for some genotypes.

660

Figure 8: Evolved lineages in cluster 6 have higher fitness than other lineages in the absence of FLU and
RAD. (A) Same UMAP as **Figure 4D** with clusters 4, 5, and 6 highlighted and sequenced isolates in these clusters
represented as diamonds. Diamond colors correspond to the targets of adaptation in the sequenced isolates. Pie
charts display the relative frequency with which lineages in cluster 6 were sampled from each of the 12 evolution
conditions; colors match those in **table 1**. **(B)** Of the three clusters on the bottom half of the UMAP, cluster 6
lineages perform best in conditions without any drug and with the highest concentration of FLU. Yet they perform
worst in conditions with the lowest concentration of FLU. Boxplots summarize the distribution across all lineages
within each cluster in each environment, displaying the median (center line), interquartile range (IQR) (upper and
lower hinges), and highest value within $1.5 \times$ IQR (whiskers).

Discussion:

Here, we present a barcoded collection of fluconazole (FLU) resistant yeast strains that is unique in its size, its diversity, and its tractability. One way we were able to isolate diverse types of FLU-resistance was by evolving yeast to resist diverse drug concentrations and combinations. But the more important tool used to increase both the number and type of mutants in our collection was DNA barcodes. These allowed us to sample beyond the drug resistant mutants that rise to appreciable frequency and to collect mutants that would eventually have been outcompeted by others. Our primary goal in collecting these mutants was to get a rough sense of how many different mechanisms of FLU resistance may exist. This question is relevant to evolutionary medicine (because more mechanisms of resistance make it harder to design strategies to avoid resistance), evolutionary theory (because more mechanisms of adaptation make it harder to predict how evolution will proceed), and genotype-phenotype mapping (because more mechanisms makes for a more complex map).

We distinguish mutants that likely act via different mechanisms by identifying those with different fitness tradeoffs across 12 environments, leveraging the mutants' barcodes to track their relative fitness following previous work (Kinsler et al. 2020). The 774 FLU-resistant mutants studied here clustered into a handful of groups (6) with characteristic tradeoffs. Some groupings are unintuitive in that they segregate mutations within the same gene (**Figure 6**) or are distinguished by unexpectedly low fitness in multidrug conditions (**Figure 7**). These findings are important because they challenge strategies in evolutionary medicine that rely on consistent tradeoffs or intuitive trends when designing sequential drug treatments. On the other hand, the observation that some mutants have very similar tradeoffs such that they cluster together is promising in that it suggests predicting the impact of some mutations by understanding the impacts of others is somewhat feasible.

Problematically, the clusters we present are incomplete and bound to change as additional data presents itself. For one, we have shown that additional FLU-resistant mutants emerge from evolution experiments in conditions lacking FLU (**Figure 3C & 3D**). This begs questions about what other FLU-resistant mutants might emerge in environments we have not studied here. Additionally, previous work has shown that some mutants that group together in our study (e.g., GPB2 and IRA2) have different fitness profiles in conditions that we did not include here (Kinsler et al. 2020). Also of note is that our evolution experiments were conducted for only a few generations and all started from the same genetic background. Additional types of FLU-resistant mutants with unique fitness profiles may emerge from other genetic backgrounds or arise after more mutations are allowed to accumulate (Brandis et al. 2012; Bosch et al. 2021; Allen et al. 2021). Finally, by requiring that all included mutants have sufficient sequencing coverage in all 12 environments, our study is underpowered to detect adaptive lineages that have low fitness in any of the 12 environments. This is bound to exclude large numbers of adaptive mutants. For example, previous work has shown some FLU resistant mutants have strong tradeoffs in RAD (Cowen and Lindquist 2005). Perhaps we are unable to detect these mutants because their barcodes are at too low a frequency in RAD environments, thus they are excluded from our collection of 774. All of the aforementioned observations combined suggest that there are more unique types of FLU-resistant mutations than those represented by these 6 clusters, and that the molecular mechanisms that can contribute to fitness in FLU are more diverse than we know.

On the up side for evolutionary medicine, not every infection harbors all possible types of mutants. This might explain why strategies that exploit one or two common tradeoffs have some, albeit mixed, success in delaying or preventing the emergence of resistance (Amin et al. 2015; Kaiser 2017; Imamovic et al. 2018; Wang et al. 2019; Krishna et al. 2022; Nyhoegen and Uecker 2023; Waller et al. 2023). Our results encourage more complex strategies to thwart drug resistance (Iram et al. 2020), such as those that focus on advance screening to determine the resistance mechanisms that are present (Andersson et al. 2019a), or on cycling a larger number of drugs to exploit a larger number of tradeoffs (Yoshida et al. 2017; Thomas et al. 2022). Problematically, these strategies often rely on knowledge about the diversity of mutants and tradeoffs that exist (or that can emerge) within an infectious population. This type of information about population heterogeneity, heteroresistance, and substructure is expensive and arduous to obtain (Andersson et al. 2019b; Bottery et al. 2021). Fortunately, new methods, in addition to the one presented in this study, are emerging (Kuchina et al. 2021; Aissa et al. 2021; Nagasawa et al. 2021; Forsyth et al. 2021; Hsieh et al. 2022; Brettner et al. 2022a). The richer data provided by these methods dovetails with emerging population genetic models that predict the likelihood of resistance to a given drug regimen (Read and Huijben 2009; Day et al. 2015; Wilson et al. 2016; Cannataro et al. 2018; Somarelli et al. 2020; Feder et al. 2021; King et al. 2022). In sum, our observation of numerous different types of drug-resistant mutations suggests that designing resistance-detering therapies is challenging, but perhaps not impossible.

Outside of predicting the evolution of resistance, our findings provide a tool to investigate the phenotypic impacts of mutation. This task has proven daunting in light of work demonstrating that mutations often have many phenotypic impacts (Paaby and Rockman 2013; Boyle et al. 2017) and that these impacts change with contexts including the environment (Paaby et al. 2015; Geiler-Samerotte et al. 2016, 2020; Lee et al. 2019; Eguchi et al. 2019). The approach presented in this study provides a way forward, not only by identifying mutations that cluster together such that the effects of some can be predicted from others, but also by identifying environments that differentiate one cluster of mutants from another. This suggests where to look to understand the phenotypes that differentiate each cluster of mutants. For example, we were able to show that the growth phenotypes of mutants from clusters 4 and 5 are different because we knew to look for these differences in multidrug environments (**Figure 7**). And our results suggest radicicol environments may be most helpful in teasing out any phenotypic differences that set apart some PDR mutations from others (**Figure 6**). Thus, our approach guides efforts to understand the phenotypic effects of mutation, while also guiding efforts to predict the effects of some mutations from others as well as efforts to predict the outcomes of adaptive evolution.

Acknowledgements:
We are grateful to Sasha Levy for providing plasmids and yeast strains, to Martin Mullis, Xianan Liu, Sasha Levy and Gavin Sherlock for advice about creating a high diversity library of barcoded yeast, Jamie Blundell, Sandeep Venkataram and Fangfei Li for discussions about how to infer adaptive lineages, and to the members of the Geiler-Samerotte lab as well as Michael Lynch for discussions about the manuscript. The authors acknowledge resources and support from the KE core facilities at Arizona State University.

Funding:
This work was supported by a National Institutes of Health grant R35GM133674 (to KGS), an Alfred P Sloan Research Fellowship in Computational and Molecular Evolutionary Biology grant FG-2021-15705 (to KGS), and a National Science Foundation Biological Integration Institution grant 2119963 (to KGS).

Methods:

Base yeast strains
All of the yeast lineages studied here originated from the same starting strain referred to as the "landing pad strain" (SHA185) in previous work (Levy et al. 2015). We transformed a barcode library (generously provided by Sasha Levy) into this strain as described below, creating a strain with the following genetic background:

MATα, ura3Δ0, ybr209w::Gal-Cre-KanMX-1/2URA3-loxP-Barcode-1/2URA3-HygMX-lox66/71.

Base media
All experiments were conducted in "M3" media defined in the same study as the landing pad strain (Levy et al. 2015), which is a glucose-limited media lacking uracil. In our study, we supplemented this media with fluconazole, radicicol, or DMSO when appropriate.

Selecting drug concentrations
Our goal was to choose concentrations of each drug that would not kill so many yeast cells as to dramatically decrease barcode diversity. We wanted to maintain a high number of unique barcodes so we could track a high number of yeast lineages as they independently evolved drug resistance. We measured the effect of each drug and drug combination on the growth rate of a single barcoded yeast strain using a plate reader to track changes in optical density (OD) over time. Ultimately we chose a "low" concentration of each drug that appeared to have no effect on growth rate, and a "high" concentration that appeared to reduce growth rate by about 15% (**Figure S2**). Though the lowest concentration of radicicol that we tested on a plate reader was 10 μM, we chose 5 μM as our low RAD concentration because previous work suggested this concentration had widespread effects on yeast physiology without affecting growth (Jarosz and Lindquist 2010; Geiler-Samerotte et al. 2016). To perform our plate reader experiment, a single colony was grown to saturation. From this culture, 5 μl was added to every well of a 96-well plate, where every well contained 195 μl of M3 media. Some wells also contained either fluconazole, radicicol, DMSO, or combinations of these drugs. The concentrations that were tested are listed on the horizontal axis of **figure S2**; each drug condition was replicated six times. The 96-well plate was incubated at 30°C for 48 hours on a plate reader and OD measurements were taken every 30 minutes. Raw OD values were exported and maximum exponential growth rates for all tested conditions were calculated from the log-linear changes in OD over time.

Inserting 300,000 unique DNA barcodes into otherwise genetically identical yeast cells
In order to track many yeast lineages as they independently develop drug resistance, we needed to insert unique DNA barcodes into many yeast cells. Plasmids harboring barcodes (pBar3) were the same as those used in a previous barcoded evolution experiment (Levy et al. 2015) and were generously provided to us by Sasha Levy. These barcodes are 25 base pairs in length. They are targeted to an artificial intron within the Ura3 gene, such that they must be retained in media lacking uracil but are not expressed and thus do not themselves affect fitness (Levy et al. 2015). We transformed this barcode library (pBar3) into the landing pad strain (SHA185) as was done previously, activating a Cre-lox recombination system by growing the cells in YP-galactose, which resulted in genomic integration of the barcode. However, our efforts to perform extremely high efficiency transformations from which we could isolate hundreds of

810 thousands of uniquely barcoded yeast were unsuccessful, despite manipulating the levels and timing of
811 the inducer (galactose). Ultimately we performed 24 separate transformations and pooled many of these
812 to obtain a large pool of barcoded yeast where every yeast cell was genetically identical except for its
813 DNA barcode.
814
Examining the frequency of each barcode in the starting pool of cells
We sequenced the barcode region of these 24 transformed yeast populations on the Hiseq X platform
using a dual index system (Kinsler et al. 2023) to discern barcode coverage, i.e., how many total unique
barcodes were successfully inserted into yeast cells and how evenly these barcodes were sampled. We
needed many uniquely barcoded yeast in order to observe many different adaptive lineages within each
evolution experiment. But barcodes with very high frequencies, referred to herein as monster lineages,
were present in 10 of the 24 transformations and present a problem. Monster lineages allow too many
cells to carry the same barcode, giving that barcode more chances to develop an adaptive mutation. This
could allow different cells harboring that same barcode to pick up different adaptive mutations,
destroying our ability to draw conclusions about adaptive mutations by using barcodes. Therefore, our
final library of barcoded lineages was created by pooling 14 individual transformations together, choosing
those 14 that lacked monster lineages, which we defined as lineages representing greater than 1% of all
transformants. Our sequencing results suggest that this library contains about 300,000 unique barcodes.
828
Initiating 12 barcoded evolution experiments
All evolution experiments started from the same pool of roughly 300,000 uniquely barcoded yeast
lineages. To start the evolution experiments, a pea sized amount of the frozen yeast barcode library was
grown up in 4 ml YPD for 4 hours at 30 °C in a shaking incubator at 220 rpm. Then, 300 µl of the grown
barcode library was added to each of 12 pre-prepared 500mL flasks representing the 12 evolution
experiments listed in **Table 1**. To prepare these flasks, first, 1.2 L of M3 media was warmed at 30°C.
Then, 100 ml was added to each of 12 flat bottom flasks. Next, 500 µl of the appropriate drug or drug
combination was added to each flask. Drugs were pre-diluted, aliquoted and frozen such that 500 µl of the
appropriate tube could be added to each flask to achieve the desired concentration as listed in **Table 1**. All
drugs were resuspended in DMSO such that the final concentration of DMSO in all experiments (except
the "no drug" control) was 0.5%.
840
Performing barcoded evolution experiments
Evolution experiments were performed following previous work (Levy et al. 2015). After initiation (see
above), the yeast in every flask were allowed to grow at 30°C with shaking at 200 RPM for 48 hours.
Then, the flasks were removed from the incubator and 400 – 1000 µl of each culture was transferred to a
new pre-prepared flask with identical conditions to the first. The reason we added more volume (1000 µl)
to some flasks than previous work was that the cell counts at the end of the 48 hours were lower for some
of our higher drug conditions. We adjusted the transfer volume to maintain a transfer population of 4×10^7
cells, which was the same as in previous work (Levy et al. 2015). We completed a total of 24
growth/transfer cycles, corresponding to 192 generations of growth assuming 8 generations per 48-hour
cycle (Levy et al. 2015). Following each transfer, the remaining culture from each flask were split into
two 50 ml conical vials, centrifuged for 3 minutes at 4000 rpm, and the supernatant was discarded. The
final pellet was resuspended in 30% glycerol up to a total volume of 6 ml before being split into three 2
ml cryovials and stored at -80°C. These frozen samples were later utilized for barcode sequencing and
isolating adaptive mutants.
855
Isolating a large pool of adaptive mutants
To generate a large pool of diverse adaptive mutants, our goal was to collect a sample from each
evolution experiment at a time point when there were many different adaptive lineages competing. If we
sampled too late, the adaptive lineage with the greatest fitness advantage would have already risen to high
frequency, thus reducing diversity. But if we sampled too early, adaptive lineages would not yet have

risen in frequency above other lineages. Therefore, we chose to sample cells from a time in each
evolution experiment when many barcoded lineages appeared to be rising in frequency (**Figure S1**). We
sampled either 1 or 2 thousand cells per each evolution experiment by spreading frozen stock from the
chosen time point onto agarose plates, scraping 1 or 2 thousand colonies into a 15mL conical tube
containing a final concentration of 30% glycerol, and freezing the pool pertaining to each of the 12
evolutions. We sampled 2,000 cells from most evolution experiments, but sampled only 1,000 from those
containing a high concentration of FLU as those evolutions appeared to have reduced barcode diversity
(**Figure S1**), presumably because high FLU represents a strong selective pressure. We sequenced the
barcodes from each of these 12 pools so that we could track which adaptive mutants originated from
which evolution experiment (see methods section below entitled, "**Inferring where adaptive lineages**
originally evolved").
872
Initiating barcoded fitness competition experiments
To assess the fitnesses of the 1 or 2 thousand barcoded lineages that we sampled from each evolution
experiment, we pooled all sampled lineages together into a larger pool of roughly 21,000 barcoded lineages.
We used this larger pool to initiate 24 fitness competition experiments, 2 replicates for each of the 12
conditions listed in **Table 1**. In this type of competition, we measure fitness by tracking changes in each
barcode's frequency over time. Barcodes that rise in frequency represent strains that have higher fitness
than others.
880
Our goal was to calculate the fitness effect of adaptive mutations. Therefore, we needed to calculate the
fitness of every evolved lineage relative to the unmutated ancestor of the evolution experiments. To do so,
we followed previous work by spiking in a large quantity of this unmutated ancestor strain into each fitness
competition, with this ancestor making up at least 90% of the final culture (Venkataram et al. 2016; Kinsler
et al. 2020). In environments containing a high concentration of FLU which resulted in the ancestral strain
having a more severe growth defect, we spiked in the ancestor such that it represented 95% of the final
pool.
888
889   To avoid wasting 90% or more of our sequencing reads on the ancestor strain's barcode, we created a
890   barcodeless ancestor strain. This strain was created by transforming SHA185 with a linear piece of DNA
891   such that the genetic background was identical to the strains of the barcoded library, but the homology to
892   the primers used to amplify the barcode was missing. Thus the DNA from these cells does not get amplified
893   or sequenced during subsequent steps.
894
895   In addition to this barcodeless ancestor, we also spiked in some barcoded ancestral strains at lower
896   frequency (1%) to use as "reference" or "control" strains, following previous work (Kinsler et al. 2020,
897   2023). These strains have been previously shown to possess no fitness differences from the ancestor. We
898   used these strains as a baseline when calculating relative fitness by setting the fitness of these strains to zero
899   during our fitness inference procedure (see methods section below entitled, "**Inferring fitness**").
900
All 24 fitness competitions were performed simultaneously in one big batch (Kinsler et al. 2023) and
initiated from the same pool of roughly 21,000 barcoded evolved yeast lineages, barcodeless ancestor, and
control strains. To initiate the competitions, $7x10^7$ cells from this pool were added to 24 pre-prepared
500mL flasks corresponding to the conditions listed in **Table 1**. These flasks were prepared exactly the
same way as was done for the evolution experiments (see above in "**Performing barcoded evolution**
experiments"). Each flask was allowed to grow for 48 hours at 30°C with shaking at 200 RPM.
907
Performing barcoded fitness competition experiments
Fitness competitions were performed following previous work (Kinsler et al. 2020). After the initial flasks
were allowed to grow for 48 hours, they were removed from the incubator and 400 µl from each culture
representing $4x10^7$ cells were transferred to a new flask with identical media. For each of 24 competitions,

we completed a total of 4 growth/transfer cycles, corresponding to 40 generations of growth assuming 8
generations per 48-hour cycle (Levy et al. 2015). Following each transfer, the remaining culture from each
flask was split into two 50 ml conical vials, centrifuged for 3 minutes at 4000 rpm, and the supernatant was
discarded. The final pellet was resuspended in 30% glycerol up to a total volume of 6 ml before being split
into three 2 ml cryovials and stored at -80°C. These frozen samples were later utilized for DNA extraction
and subsequent barcode sequencing.
918
Extracting genomic DNA
DNA was extracted from 500 μl of concentrated frozen stocks pertaining to the evolution experiments
and fitness competitions. Frozen cells were thawed and pelleted. Cells were treated with 250 μl of 0.1 M
Na2EDTA, 1M sorbitol and 5U/μl zymolyase for a minimum of 15 minutes at 37 °C to remove the cell
wall. Lysis was completed by adding 250 μl of 1% SDS, 0.2N NaOH and inverting to mix. Proteins and
cell debris were removed with 5M KOAc by spinning for 5 min at 15,000 rpm. Supernatant was moved to
a new tube and DNA was precipitated with 600 μl isopropanol by spinning for 5 min at 15,000 rpm. The
resulting pellet was washed 1 ml of 70% ethanol before being resuspended in 50 μl water plus 10μg/ml
RNAse. Extracted DNA was quantified using the NanoDrop spectrophotometer and all samples were
diluted to a concentration of 50 ng/μL for barcode amplification and sequencing library preparation.
929
Preparing barcodes for high-throughput multiplexed sequencing using PCR
Extracted DNA was prepared for sequencing using a two-step PCR that preserves information about the
relative frequency of each barcode in each sample (Venkataram et al. 2016; Kinsler et al. 2020, 2023).
Briefly, in the first step PCR, the barcode region is amplified from the genomic DNA, labeled with a
sample-specific combination of primers, and tagged with a UMI. This step utilizes a short 3 cycle PCR
with New England Biolabs OneTaq polymerase. Purification of the first step product to remove excess
reagents was performed using Thermo Scientific GeneJET PCR Purification Kit. The second step PCR
attached Illumina indices that were used to distinguish samples from different experiments and
timepoints. We utilized a dual indexing scheme to prevent index misassignment that is common when
sequencing amplicon libraries using patterned flow cell technology (Kinsler et al. 2023). Amplification of
this second step of PCR was done with a longer 23 cycle PCR using Q5 polymerase. Final libraries were
bead purified using 0.8X Quantabio sparQ Pure Mag beads. Quantification of the final PCR products was
done using the Invitrogen Qubit Fluorometer before all samples were pooled at equimolar ratios for
sequencing.
944
Sequencing and clustering barcodes
Next Generation Sequencing was performed at either Psomagen (Rockville, MD) or at the Translation
Genomics Research Institute (Phoenix, AZ) on patterned flow cells (either an Illumina HiSeqX or
NovaSeq) using 2 x 150 base pair paired end reads. Samples were dual indexed to allow multiplexing
while minimizing contamination from index misassignments (Kinsler et al. 2023). The 20 base pairs of
variable sequence referred to as a DNA barcode were identified and clustered to determine the number of
unique barcodes and the frequency of each barcode in each sample. For the evolution experiments, this
was done following our previous work (Venkataram et al. 2016; Kinsler et al. 2020). For the fitness
competition experiments, this was done using updated software (Zhao et al. 2018) with the following
command:
bartender_extractor_com -f ${SAMPLE}_R1_001.fastq -o ${SAMPLE}_extracted -q 0 -p
GTACC[5]AA[5]AA[5]TT[5]ATAAC -m 2 -d f -u 0,8
957
Inferring fitness
In fitness competition experiments, fitness is often inferred from the log-linear change in a strain's
frequency over time (Geiler-Samerotte et al. 2011; Bakerlee et al. 2021; Kinsler et al. 2023). Recently,
more advanced methods to infer fitness have emerged that take into account nonlinearities in frequency
changes over time, for example, nonlinearities that reflect changes in the mean fitness of the population

(Venkataram et al. 2016; Li et al. 2018a, 2023; Kinsler et al. 2020). We had trouble implementing these
newer methods on our fitness data, perhaps because many of our evolved lineages, and our control strains,
have low fitness in some drugs. This caused their barcodes to rapidly decline in frequency such that they
received low counts only at later time points. Their counts could become so low that these lineages would
seemingly disappear due to sampling error, and then reappear at a subsequent time point. This dramatic
(but false) late increase in frequency was sometimes interpreted as evidence of very high fitness,
especially when we inferred fitness using approaches that account for nonlinearities.
970
To contend with this issue, we applied strict coverage thresholds to every fitness measurement: we
required at least 500 counts across all timepoints in order to infer fitness for a given lineage in a given
environment. This is stricter than previous work that does not require a minimum number of reads per
each lineage and instead requires a minimum number of reads per time point (Kinsler et al. 2020). We
found that 774 lineages passed our threshold in at least one replicate experiment per all 12 environments.
Of these, 729 passed for both replicates and the final fitness value we report represents the average of
both replicates.
978
Even with our strict coverage threshold, fitness inference methods that account for nonlinearities still
interpreted minor stochastic fluctuations in fitness at later time points as evidence of a fitness advantage,
even if fitness dramatically declined in earlier time points. Therefore, we calculated fitness via the
traditional method, as the slope of the log-linear change in barcode frequency relative to the average slope
of the control strains. We found that this method is less sensitive to that type of error. Using this method,
we found that our fitness inferences were reproducible between replicates (**Figure S4A**), and between
experiments performed in similar conditions (e.g., medium vs. high concentrations of the same drug)
(**Figure S4B**). When we increased our coverage threshold to require an order of magnitude more reads
per lineage per measurement (from 500 to 5000), we lost 157 lineages (from 774 to 617), saw
reproducibility increase across replicates (from an average Pearson correlation of 0.756 to 0.813) and the
main conclusions of our study were unchanged in that the same 6 clusters were present on a UMAP
(**Figure S7**).
991
Identifying adaptive mutations using whole-genome sequencing
One downside of barcoded evolution experiments is that all lineages exist together in a pooled culture.
Fishing out adaptive lineages in order to perform whole genome sequencing is a major challenge
(Venkataram et al. 2016). Here, we randomly selected cells from these mixed pools for whole genome
sequencing, sometimes selecting from later time points in the evolution experiments and sometimes
selecting from the samples of 1 or 2 thousand cells that were isolated to initiate fitness competitions.
998
To perform whole genome sequencing, cells from mixed pools were spread onto M3 agarose plates,
single colonies were selected and grown in YPD to saturation. DNA was extracted using the PureLink™
Genomic DNA Mini Kit (K182002). Sequencing libraries were made using Illumina DNA Prep kit by
diluting reactions by ⅕. Briefly, samples were prepared such that the starting concentration in 6 ul was
between 20 and 100 ng of DNA. 2μl of BLT and TB1were added to the starting material and incubated on
a thermocycler at 55°C (lid 100°C) for 15 min. 2 μl of TSB was added to each reaction and incubated at
37 C (lid 100°C) for 15 min. Beads were washed 2 times with 20 μl of TWB. Following the final wash, 4
μl of EPM, 4 μl of water and 2 μl of UD indexes were added to each sample. Depending on starting
concentration, PCR was performed based on Illumina guidelines as follows: lid 100°C, 68°C for 3 min,
98C for 3 min, [98°C for 45s, 62°C for 30s, 68°C for 2min] for 6 to 10 cycles, 68°C for 1 min, 10°C hold.
PCR products were cleaned with a double side sized selection as follows: 4 μl of each sample was pooled
together (32 μl total for 8 samples) and added to 28 μl of water plus 32 μl of SPB. After a 5 min
incubation 25 μl of supernatant was moved to a new tube containing 3 μl of SPB. Beads were washed
with fresh 80% ethanol and libraries were eluted in 12 μl RSB. Samples were multiplexed using

Illumina's unique dual (UD) index plates (A-D) and sequencing was performed with 2x150 paired end
sequencing on HiSeq X at Psomagen (Rockville, MD).
1015
In total 122 colonies were randomly picked and sequenced. As one might expect, barcodes that rose to
high frequency were more likely to be picked multiple times. In an attempt to find lineages with unique
attributes, some cultures were grown at 37°C or plated to high concentrations of drug prior to picking
isolated colonies for sequencing. Of the 122 genomes we sequenced, only 53 had unique barcodes and
pertained to the 774 lineages for which we obtained high enough barcode coverage to infer fitness. Only
two of these 53 had no sequenced mutations suggesting their fitness increase over ancestor is due to a
mutation we are unable to identify by sequencing, perhaps a change in ploidy. The other 51 all had at
least one single nucleotide mutation in a gene reported in **supplemental table 1**.
1024
Variant calling was done using GATK as described here: https://github.com/gencorefacility/variant-
calling-pipeline-gatk4. Identified variants were annotated using SnpEff (Cingolani et al. 2012). Variant
call files from 132 (53 unique/in CS) sequenced lineages were analyzed in R and compared to reference
strain GCF_000146045.2 (Genome assembly 64: sacCer3). SNPs present in the ancestor (as well as all
evolved lineages) were ignored as these could not have caused the fitness differences we observed. We
also ignored SNPS that were present in a substantial number of evolved lineages, as these likely represent
background mutations that were present in a substantial portion of the cells representing the landing pad
strain (SHA185). These are reported in **supplemental table 1** and include: SRD1-Glu97Lys, RSC30-
Gly571Asp, OPT1-Val143Ile and LYS20-Thr29Met.
1034
Measuring growth curves of evolved lineages with unexpected fitness in multidrug conditions
Though fitness differences are not necessarily due to differences in maximum growth rate (Li et al.
2018b), we measured growth curves for a few lineages to investigate a case where an evolved lineage had
unexpectedly low fitness in multidrug conditions (**Figure 7B**). Indeed, we found that this mutant grew
more slowly in those conditions (**Figure 7D; S10**). To perform this test, a lineage with a mutation to
GBP2, a lineage with a mutation to HDA1, and sometimes the ancestor strain were streaked to YPD
plates. We used the barcodeless ancestor strain, which is identical to the evolved lineages in every way
except for lacking a barcode, and is described above in the methods section entitled, "**Initiating
barcoded fitness competition experiments".** A single colony of each strain was isolated from YPD
plates and was used to inoculate an overnight YPD culture. After ~24 hours, a coulter counter (BD) was
used to determine the number of cells/ml present in each culture. Next, all cultures were diluted such that
the starting number of cells was 250,000 in 6 ml of M3 plus drug (either HR, LF, LR, LRLF or HRLF,
see **Table 1**). To measure growth curves, these samples were allowed to grow at 30 degrees C. OD was
measured every 10 minutes as the cultures were grown to saturation using the compact rocking incubator
TVS062CA (Advantec Mfs). Raw growth curves for these conditions are shown in **Figure S10B and C**.
Maximum growth rate was calculated using a sliding window approach to determine the region of each
growth curve with the steepest log-linear slope.
1052
Determining ploidy
While our barcoded yeast strain is haploid, previous studies observed that some cells diploidize during the
course of evolution in M3 media and by doing so gain a fitness advantage (Levy et al. 2015; Venkataram
et al. 2016). To ensure that observed fitness effects in our experiments were not largely due to the effects
of diploids, we estimated the percent of diploid cells in each of our populations. We chose to make our
estimates from frozen samples taken at the same time points from which we sampled 1 or 2 thousand cells
to initiate fitness competitions (**Figure S1**). As such, our estimates also report on the percent of diploids
that were present at the start of the fitness competitions experiments (**Table S2**).
1061
To study ploidy, we used the nucleic acid stain SYTOX Green, which is capable of selectively staining
the nucleus of fixed cells and has been shown to be optimal for use in budding yeast (Haase 2004). For

each of the 12 evolution experiments conditions, a small amount of freezer stock from the chosen
timepoints (**Figure S1**) was plated to YPD and grown for ~48 hours. Individual colonies were picked and
transferred to 96-well plates, one full plate per each condition, before being fixed with 95% ethanol for 1
hour. Plates were centrifuged at 4500 rpm and supernatant was discarded. 50 µL RNase A was added to
the samples at a concentration of 2mg/mL, and the plates were then incubated for 2 hours at 37°C. Cells
were pelleted by centrifuge and the supernatant was removed, which was followed by treatment with 20
µL of the protease pepsin at a concentration of 5mg/uL. Pepsin treated samples incubated at 37°C for 30
minutes before centrifugation and removal of supernatant. Finally, cells were resuspended in 50 µL
TrisCL (50 mM, pH 8) and stained with 100 µL of 1µM SYTOX Green. Known diploid and haploid
strains were used as controls alongside our samples to determine the expected fluorescence of stained
diploid vs. haploid cells. Analysis was performed using a ThermoFisher Attune NxT, housed in the Flow
Cytometry Core Facility at Arizona State University.
1076
Dimensional reduction
Our fitness inference procedure resulted in a data set consisting of nearly 10,000 fitness measurements
(774 lineages x 12 conditions = 9288 fitness measurements). Dimensional reduction was performed on
these data using UMAP (McInnes et al. 2018b). Clusters of similar mutants were identified and colored
using a gaussian mixed model (Fraley and Raftery 2003); Bayesian Information Criteria were used to
select the number of clusters (**Figure S6**). These analyses were performed in R; code can be found
https://osf.io/pxyv9/?view_only=51241d8f00c24f7e83f1ece3ae31a53b.
1084
In order to prevent conditions with the most variation in fitness (e.g., high FLU) from dominating, we
normalized fitness measurements from each of the 12 environments to have the same overall mean and
variance (we transformed the data from every environment to have a mean of 0 and a standard deviation
of 1) before performing dimensional reduction. This normalization procedure did not have a dramatic
effect on the UMAP (**Figure S7A**). We also explored normalizing all data to account for magnitude
differences by setting the average fitness of each lineage across all 12 environments to 0. Doing so did not
significantly change the groupings present in the UMAP from those displayed in **Figure 4** (**Figure S9**)
other than in the ways we describe in **Figure 6**. Reducing our data set to 617 adaptive lineages with very
high sequencing coverage (**Figure S7B**) also did not significantly affect the way that mutants cluster into
groups, nor did using a different dimensional reduction algorithm altogether (**Figure S8** and see next
paragraph). In short, the clustering of mutants was robust to the different decisions we made when
choosing how to analyze these data.
1097
In order to assess whether clusters identified from the UMAP are robust to alternative clustering methods,
we also used hierarchical clustering to identify clusters of mutants with similar fitness profiles. First, we
computed the pairwise distance of all lineages across the fitness profiles. Then, we used Ward's method
from scikit-learn to iteratively cluster lineages such that the within-cluster variation is minimized (Ward
1963; Pedregosa et al. 2012). To test the consistency of lineage clustering, we chose a pairwise cluster
distance cutoff of 11, which results in the same number of clusters (7) as identified with the UMAP
clustering approach used in the main text. We then compared the identity of the lineages within each of
these clusters with the UMAP clusters. We found that, for most clusters, over 80% of lineages from the
UMAP cluster corresponded with a unique hierarchical cluster and labeled these hierarchical clusters
according to this correspondence (**Figure S8**). For UMAP cluster 1, lineages were more evenly split
between two clusters. 64% of these lineages clustered together in what is labeled as hierarchical cluster 1
and 30% in hierarchical cluster 1/7 (**Figure S8**), which contains all of the control lineages that comprise
UMAP cluster 7. Despite these lineages clustering more closely with control lineages than the remainder
of the cluster 1, they do tend to cluster distinctly with the control lineages, suggesting they have behavior
that is distinguishable from the control lineages. If we consider these cluster 1 mutants that end up in
cluster 3/7 as "mis-clustered", we find that 85% of lineages from each UMAP cluster are clustered
together in the corresponding hierarchical cluster. If we consider these as "consistently clustered", this

metric increases to 90% of lineages correctly clustered. Altogether, this analysis shows that the results we
show are robust to alternative methods of clustering.
1117
Inferring where adaptive lineages originally evolved
All 774 adaptive lineages were isolated from one of the 12 evolution experiments at the timepoint
indicated in **figure S1** (see Methods section entitled, "**Isolating a large pool of adaptive mutants**"). The
sample we isolated from each evolution experiment was sequenced prior to pooling. This allows us to
computationally determine which barcoded lineages originated from which evolution experiment to
generate the pie charts in F**igures 3**, **5**, **6**, **7** and **8**.
1124
1125 If adaptive mutation arose independently during the course of each evolution experiment, it would be
1126 unlikely for any adaptive lineage we study to be present in more than one of the evolution conditions.
1127 This would make it very easy to assign each barcode to the evolution experiment from which it
1128 originated. However, this was not the case for many barcoded lineages.
1129
Previous work explained that the transformation procedure used to insert a barcode into the landing pad of
SHA185 is itself mutagenic, such that many mutations arise prior to the start of the evolution experiments
(Levy et al. 2015). Since all our evolution experiments were started from the same pool of barcoded
lineages, we thus expect that many adaptive lineages will be present in more than one condition.
However, it is not expected that these adaptive lineages will be present at the same frequency in every
condition; instead these frequencies change with the fitness of the mutation each lineage possesses.
Therefore, when an adaptive lineage appeared in multiple conditions, we weighted its origin to reflect its
frequency in each condition. In other words, adaptive lineages that were only present in the sample taken
from a single evolution condition were identified and assigned a single origin condition in the pie charts
in **Figures 3**, **5**, **6**, **7**, and **8**. But for adaptive lineages found in the samples taken from more than one
evolution condition, the proportions assigned to each origin condition in the pie charts was scaled to equal
the relative frequencies of that lineage in all evolution conditions where it was observed. Associated data
and code can be found here: https://osf.io/pxyv9/?view_only=51241d8f00c24f7e83f1ece3ae31a53b.

1166 **Supplemental Figures and Tables**

1167

1168
Figure S1: Twelve barcoded evolution experiments track ~300,000 lineages as they adapt to different
drug concentrations and combinations. In each plot, every line represents a unique barcoded lineage. The
vertical axis represents barcode frequency. When lineages reach a frequency of zero it means they were not
sampled at that time point of the experiment. The black boxes indicate the transfer number of each evolution
experiment from which evolved lineages are sampled; the number of cells sampled (colonies picked) is in the
upper right hand corner.
1175

1177
Figure S2: Chosen drug concentrations do not dramatically reduce yeast's maximum growth rate. We
measured the growth rate of a single barcoded yeast lineage, prior to the evolution experiment, in different
concentrations and combinations of drugs using a plate reader to track changes in optical density over time.
Maximum growth rate is reported as a percentage of the maximum growth rate in conditions lacking FLU or
RAD. Maximum growth rate was calculated as the maximum log-linear slope of the change in optical density
over time.

1185
Figure S3: Twenty four fitness competitions track evolved lineages as their barcodes change frequency.
In each plot, every line represents a barcoded lineage. Lineages with >5 reads per experiment are shown (4,815
lineages shown). Control lineages, known to possess no fitness differences relative to the ancestor, are
highlighted in black. Evolved lineages with higher fitness than controls are yellow, similar fitness to controls
(i.e., relative fitness ~0) are purple, and lower fitness than controls are orange. When a lineage declines in
frequency so much that its associated barcode is no longer observed, its line abruptly ends. Some lineages
appear to decline and then increase in frequency; this happens because low frequency lineages are subject to
sampling noise.

Figure S4: Fitness measurements are reproducible between replicates and closely related conditions.
A. Every point depicts one of the evolved lineages that was observed more than 500 times in both replicates of
the fitness competition experiment for a given condition. The average Pearson correlation across all pairs of
replicates in all conditions is 0.75. **B.** Every point depicts one of the 774 lineages that was observed more than
500 times in at least one replicate fitness competition. Here, instead of comparing the fitness of replicates, we
compare fitness across conditions for 3 closely related pairs of conditions. Comparisons across conditions may
be less noisy than those across replicates because fitness in a particular condition represents the average fitness
across replicates in that condition.

1207
1208

Figure S5: **The two types of adaptive mutants depicted in figure 3 sort into different clusters on the UMAP.** The top 100 highest fitness lineages in high FLU (blue) and high RAD (red) largely cluster into different groups, with the 7 overlapping mutants (black) falling into the uppermost group.

Figure S6: Bayesian information criteria (BIC) scores suggest the 774 mutants cluster into between 6 and 13 groups. We used a gaussian mixture model to distinguish clusters of mutants with unique fitness profiles. **A.** BIC scores for analysis performed on 774 mutant lineages that each were observed at minimum 500 times per environment. **B** BIC scores for analysis performed on 617 mutant lineages that each were observed at minimum 5000 times per environment.

Figure S7. UMAP structure is robust. A. In this UMAP plot, relative fitness of the 774 lineages is fed directly into UMAP prior to normalizing the data to have equal variance across all environments as was done in **figure 4D**. This has little impact on the appearance of the UMAP. **B.** UMAP made with 617 lineages that were observed more than 5000 times in at least one replicate. While the appearance of the UMAP is inverted relative to **figure 4D**, the clusters largely retain the same lineages. As lower coverage (noisier) data points are removed from this dataset, the observation that the overall patterns are maintained suggests our original clusters were not based on noisy fitness measurements.

1237

1238
Figure S8. Clusters are robust to a hierarchical clustering method. A. In order to assess whether
clusters identified from the UMAP are robust to alternative clustering methods, we used hierarchical
clustering to identify clusters of mutants with similar fitness profiles. This figure depicts a dendrogram
where the branch length between each lineage (or group of lineages) represents the distance between the
clustered groups as measured across all measured conditions. The terminal branch leading to each lineage
is colored by that lineage's cluster from the UMAP clustering used in **figure 4** of the main text. Dark gray
lines show clusters identified using a between-cluster cutoff of 11, which creates 7 clusters, consistent
with the number of clusters we identified in **figure 4** via clustering on the UMAP space. Hierarchical
clusters are labeled and colored corresponding to whichever of the UMAP clusters they share the most
overlapping lineages with. Note that the y-axis is linearly scaled between 0 and 10 and log2 scaled above
10 for visualization purposes. **B.** A matrix quantifying how well the results of this hierarchical clustering
method correspond with those of the UMAP method presented in **figure 4**, depicting the percentage of a
given UMAP cluster that ends up with each corresponding hierarchical cluster. For the most part, mutants
that are clustered together in the UMAP in **figure 4** are also clustered together in **panel A**. Even though it
appears that mutants in cluster 1 (which are those most resistant to low FLU) move to a different cluster,
these mutants move together to the cluster pertaining to the control strains as they have the least severe
effects on fitness.
1256
1257
1258

1259
Figure S9: UMAP on data that were normalized to account for magnitude differences (row means
set to 0). A. Every point in this plot represents one of the 774 barcoded lineages colored by cluster; clusters
were identified using a gaussian mixture model. After normalizing to account for magnitude differences,
the 774 adaptive lineages cluster into 6 (not 7) groups based on variation in their fitness profiles. The
number of clusters is reduced relative to the UMAP in **figure 4** because the control lineages no longer form
a separate group; they cluster into the middle cluster in light green. **B.** This plot shows the same UMAP as
above in **panel A**, but the color of each of the 774 points is changed to reflect its color in the UMAP in
figure 4. Despite coloring based on a different UMAP, differently colored points still largely fall into

separate clusters, except for those points colored blue or orange, which are discussed in **figure 6**. **C.** This
plot depicts all lineages in clusters 1 – 3 in either the **figure 4** UMAP (not normalized to lessen magnitude
differences) or the above UMAP (normalized to lessen magnitude differences). Lineages that do not switch
from one cluster to another are represented by diamonds while those that switch are noted with circles.
Across these three clusters, 67% of lineages remain in their original cluster.
1273
1274
1275

Figure S10: Unexpected tradeoffs in evolved lineages in cluster 4 and 5 in response to combined
drugs. A. Cluster 5 (red) has an unexpected fitness disadvantage in the LRLF multidrug environment relative to
cluster 4 (green), given that cluster 5 lineages do not have a relative fitness disadvantage in the relevant single drug
environments. This plot is similar to the one in **figure 7B** except here the multidrug condition shown is LRLF rather
than HRLF. Boxplots summarize the distribution across all lineages within each cluster in each environment,
displaying the median (center line), interquartile range (IQR) (upper and lower hinges), and highest value within 1.5
$\times$ IQR (whiskers). **B.** Raw growth curve data from which relative maximum growth rates presented in **Figure 7D** were
calculated. Two mutants, one to GBP2 (cluster 4; green) and one to HDA1 (cluster 5; red) were each grown
individually in the conditions shown, as was the barcodeless ancestor strain (grey). All experiments were conducted
in duplicate. We initiated all cultures from the same number of starting cells. The mutant in cluster 5 (red) has an
obvious growth disadvantage in the combination condition (HRLF). **C.** Raw growth curve data for the same two
mutants in **panel B** in different conditions. These growth experiments were conducted similarly to those in **panel B**,
except they were tracked for a longer period of time, lacked the ancestor strain, and involved 3 replicates per mutant.
There appear to be obvious and reproducible differences in the growth curves of the two mutants surveyed.

1291
1292
1293

1294 **Table S1:** The sequenced mutations present in each of the adaptive lineages on which we performed whole
1295 genome sequencing
1296 https://osf.io/pxyv9/files/osfstorage/6516049b3d9bde04078d871f

1321 **Table S2:** Estimated percentage of diploid cells in each evolution condition at the time point we sampled,
1322 determined using nuclear staining and flow cytometry as described in Materials and Methods. *For the
1323 High Radicicol + Low Fluconazole condition, 95 instead of 96 samples were measured due to undetectable
1324 growth in one well.
1325
1326

Evolution Conditions	Diploids detected (96 sampled)	Estimated % diploid
Low Rad	8	8.3%
Med Rad	0	0.0%
High Rad	0	0.0%
Low Flu	9	9.4%
Med Flu	0	0.0%
High Flu	1	1.0%
LR + LF	4	4.2%
HR +LF*	1	1.1%
LR + HF	1	1.0%
HR + HF	3	3.1%
DMSO	6	6.3%
No Drug	9	9.4%

References

Abel zur Wiesch P, Kouyos R, Abel S, et al (2014) Cycling empirical antibiotic therapy in hospitals: meta-analysis and models. PLoS Pathog 10:e1004225

Aissa AF, Islam ABMMK, Ariss MM, et al (2021) Single-cell transcriptional changes associated with drug tolerance and response to combination therapies in cancer. Nat Commun 12:1628

Algazi AP, Othus M, Daud AI, et al (2020) Continuous versus intermittent BRAF and MEK inhibition in patients with BRAF-mutated melanoma: a randomized phase 2 trial. Nat Med 26:1564–1568

Allen RC, Angst DC, Hall AR (2019) Resistance Gene Carriage Predicts Growth of Natural and Clinical Escherichia coli Isolates in the Absence of Antibiotics. Appl Environ Microbiol 85.: https://doi.org/10.1128/AEM.02111-18

Allen RC, Pfrunder-Cardozo KR, Hall AR (2021) Collateral Sensitivity Interactions between Antibiotics Depend on Local Abiotic Conditions. mSystems 6:e0105521

Amin AD, Rajan SS, Liang WS, et al (2015) Evidence Suggesting That Discontinuous Dosing of ALK Kinase Inhibitors May Prolong Control of ALK+ Tumors. Cancer Res 75:2916–2927

Andersson DI, Hughes D (2010) Antibiotic resistance and its cost: is it possible to reverse resistance? Nat Rev Microbiol 8:260–271

Andersson DI, Nicoloff H, Hjort K (2019a) Mechanisms and clinical relevance of bacterial heteroresistance. Nat Rev Microbiol 17:479–496

Andersson DI, Nicoloff H, Hjort K (2019b) Mechanisms and clinical relevance of bacterial heteroresistance. Nat Rev Microbiol 17:479–496

Ardell SM, Kryazhimskiy S (2021) The population genetics of collateral resistance and sensitivity. Elife 10.: https://doi.org/10.7554/eLife.73250

Baker CM, Ferrari MJ, Shea K (2018) Beyond dose: Pulsed antibiotic treatment schedules can maintain individual benefit while reducing resistance. Sci Rep 8:5866

Bakerlee CW, Phillips AM, Nguyen Ba AN, Desai MM (2021) Dynamics and variability in the pleiotropic effects of adaptation in laboratory budding yeast populations. Elife 10.: https://doi.org/10.7554/eLife.70918

Basra P, Alsaadi A, Bernal-Astrain G, et al (2018) Fitness Tradeoffs of Antibiotic Resistance in Extraintestinal Pathogenic Escherichia coli. Genome Biol Evol 10:667–679

Baym M, Stone LK, Kishony R (2016) Multidrug evolutionary strategies to reverse antibiotic resistance. Science 351:aad3292

Berkow E, Lockhart S (2017) Fluconazole resistance in Candida species: a current perspective. Infection and Drug Resistance 10:237–245

Berman J, Krysan DJ (2020) Drug resistance and tolerance in fungi. Nat Rev Microbiol 18:319–331

Bongomin F, Gago S, Oladele RO, Denning DW (2017) Global and Multi-National Prevalence of Fungal

Diseases-Estimate Precision. J Fungi (Basel) 3.: https://doi.org/10.3390/jof3040057

Bosch B, DeJesus MA, Poulton NC, et al (2021) Genome-wide gene expression tuning reveals diverse vulnerabilities of M. tuberculosis. Cell 184:4579–4592.e24

Bottery MJ, Pitchford JW, Friman V-P (2021) Ecology and evolution of antimicrobial resistance in bacterial communities. ISME J 15:939–948

Boyer S, Hérissant L, Sherlock G (2021) Adaptation is influenced by the complexity of environmental change during evolution in a dynamic environment. PLoS Genet 17:e1009314

Boyle EA, Li YI, Pritchard JK (2017) An Expanded View of Complex Traits: From Polygenic to Omnigenic. Cell 169:1177–1186

Brandis G, Wrande M, Liljas L, Hughes D (2012) Fitness-compensatory mutations in rifampicin-resistant RNA polymerase. Mol Microbiol 85:142–151

Brettner L, Eder R, Schmidlin K, Geiler-Samerotte K (2022a) An ultra high-throughput, massively multiplexable, single-cell RNA-seq platform in yeasts. bioRxiv 2022.09.12.507686

Brettner L, Ho WC, Schmidlin K, Apodaca S (2022b) Challenges and potential solutions for studying the genetic and phenotypic architecture of adaptation in microbes. Current Opinion in

Buechel ER, Pinkett HW (2020) Transcription factors and ABC transporters: from pleiotropic drug resistance to cellular signaling in yeast. FEBS Lett 594:3943–3964

Cairns J, Borse F, Mononen T, et al (2022) Strong selective environments determine evolutionary outcome in time-dependent fitness seascapes. Evol Lett 6:266–279

Cannataro VL, Gaffney SG, Stender C, et al (2018) Heterogeneity and mutation in KRAS and associated oncogenes: evaluating the potential for the evolution of resistance to targeting of KRAS G12C. Oncogene 37:2444–2455

Centers for Disease Control and Prevention (U.S.) (2019) Antibiotic Resistance Threats in the United States, 2019. U.S. Department of Health and Human Services, Centres for Disease Control and Prevention

Chen VK, Johnson MS, Hérissant L, et al (2023) Far From Home: Evolution of haploid and diploid populations reveals common, strong, and variable pleiotropic effects in non-home environments. bioRxiv 2023.02.28.530341

Cingolani P, Platts A, Wang LL, et al (2012) A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6:80–92

Cowen LE, Lindquist S (2005) Hsp90 potentiates the rapid evolution of new traits: drug resistance in diverse fungi. Science 309:2185–2189

Cowen LE, Sanglard D, Howard SJ, et al (2014) Mechanisms of Antifungal Drug Resistance. Cold Spring Harb Perspect Med 5:a019752

Cowen LE, Singh SD, Köhler JR, et al (2009) Harnessing Hsp90 function as a powerful, broadly effective therapeutic strategy for fungal infectious disease. Proc Natl Acad Sci U S A 106:2818–2823

Day T, Huijben S, Read AF (2015) Is selection relevant in the evolutionary emergence of drug resistance? Trends Microbiol 23:126–133

Eguchi Y, Bilolikar G, Geiler-Samerotte K (2019) Why and how to study genetic changes with context-dependent effects. Curr Opin Genet Dev 58-59:95–102

Fardeau V, Lelandais G, Oldfield A, et al (2007) The Central Role of PDR1 in the Foundation of Yeast Drug Resistance*. J Biol Chem 282:5063–5074

Feder AF, Harper KN, Brumme CJ, Pennings PS (2021) Understanding patterns of HIV multi-drug resistance through models of temporal and spatial drug heterogeneity. Elife 10.: https://doi.org/10.7554/eLife.69032

Flowers SA, Barker KS, Berkow EL, et al (2012) Gain-of-function mutations in UPC2 are a frequent cause of ERG11 upregulation in azole-resistant clinical isolates of Candida albicans. Eukaryot Cell 11:1289–1299

Forsyth B, Torab P, Lee J-H, et al (2021) A Rapid Single-Cell Antimicrobial Susceptibility Testing Workflow for Bloodstream Infections. Biosensors 11.: https://doi.org/10.3390/bios11080288

Fraley C, Raftery AE (2003) Enhanced Model-Based Clustering, Density Estimation,and Discriminant Analysis Software: MCLUST. J Classification 20:263–286

Geiler-Samerotte KA, Dion MF, Budnik BA, et al (2011) Misfolded proteins impose a dosage-dependent fitness cost and trigger a cytosolic unfolded protein response in yeast. Proc Natl Acad Sci U S A 108:680–685

Geiler-Samerotte KA, Li S, Lazaris C, et al (2020) Extent and context dependence of pleiotropy revealed by high-throughput single-cell phenotyping. PLoS Biol 18:e3000836

Geiler-Samerotte KA, Zhu YO, Goulet BE, et al (2016) Selection Transforms the Landscape of Genetic Variation Interacting with Hsp90. PLoS Biol 14:e2000465

Gjini E, Wood KB (2021) Price equation captures the role of drug interactions and collateral effects in the evolution of multidrug resistance. Elife 10.: https://doi.org/10.7554/eLife.64851

Gomulkiewicz R, Kirkpatrick M (1992) QUANTITATIVE GENETICS AND THE EVOLUTION OF REACTION NORMS. Evolution 46:390–411

Good BH, McDonald MJ, Barrick JE, et al (2017) The dynamics of molecular evolution over 60,000 generations. Nature 551:45–50

Grier HE, Krailo MD, Tarbell NJ, et al (2003) Addition of ifosfamide and etoposide to standard chemotherapy for Ewing's sarcoma and primitive neuroectodermal tumor of bone. N Engl J Med 348:694–701

Haase SB (2004) Cell cycle analysis of budding yeast using SYTOX Green. Curr Protoc Cytom Chapter 7:Unit 7.23

Hall MD, Handley MD, Gottesman MM (2009) Is resistance useless? Multidrug resistance and collateral sensitivity. Trends Pharmacol Sci 30:546–556

Harris A, Wagner M, Du D, et al (2021) Structure and efflux mechanism of the yeast pleiotropic drug

resistance transporter Pdr5. Nat Commun 12:5254

Herren CM, Baym M (2022) Decreased thermal niche breadth as a trade-off of antibiotic resistance.
ISME J 16:1843–1852

Hill JA, O'Meara TR, Cowen LE (2015) Fitness Trade-Offs Associated with the Evolution of Resistance
to Antifungal Drug Combinations. Cell Rep 10:809–819

Hsieh K, Mach KE, Zhang P, et al (2022) Combating Antimicrobial Resistance via Single-Cell
Diagnostic Technologies Powered by Droplet Microfluidics. Acc Chem Res 55:123–133

Imamovic L, Ellabaan MMH, Dantas Machado AM, et al (2018) Drug-Driven Phenotypic Convergence
Supports Rational Treatment Strategies of Chronic Infections. Cell 172:121–134.e14

Iram S, Dolson E, Chiel J, et al (2020) Controlling the speed and trajectory of evolution with
counterdiabatic driving. Nat Phys 17:135–142

Iwasawa J, Maeda T, Shibai A, et al (2022) Analysis of the evolution of resistance to multiple antibiotics
enables prediction of the Escherichia coli phenotype-based fitness landscape. PLoS Biol
20:e3001920

Iyer KR, Robbins N, Cowen LE (2022) The role of Candida albicans stress response pathways in
antifungal tolerance and resistance. iScience 25:103953

Jarosz DF, Lindquist S (2010) Hsp90 and Environmental Stress Transform the Adaptive Value of Natural
Genetic Variation. Science 330:1820–1824

Jerison ER, Nguyen Ba AN, Desai MM, Kryazhimskiy S (2020) Chance and necessity in the pleiotropic
consequences of adaptation for budding yeast. Nat Ecol Evol 4:601–611

Kaiser J (2017) When less is more. Science 355:1144–1146

Kapitzky L, Beltrao P, Berens TJ, et al (2010) Cross-species chemogenomic profiling reveals
evolutionarily conserved drug mode of action. Mol Syst Biol 6:451

King ES, Maltas J, Weaver DT, et al (2022) Fitness seascapes are necessary for realistic modeling of the
evolutionary response to drug therapy. biorxiv

Kinsler G, Geiler-Samerotte K, Petrov DA (2020) Fitness variation across subtle environmental
perturbations reveals local modularity and global pleiotropy of adaptation. Elife 9.:
https://doi.org/10.7554/eLife.61271

Kinsler G, Schmidlin K, Newell D, et al (2023) Extreme sensitivity of fitness to environmental
conditions: Lessons from #1BigBatch. J Mol Evol 91:293–310

Kolaczkowski M, van der Rest M, Cybularz-Kolaczkowska A, et al (1996) Anticancer drugs, ionophoric
peptides, and steroids as substrates of the yeast multidrug transporter Pdr5p. J Biol Chem
271:31543–31548

Krishna A, Zere T, Mistry S, et al (2022) Evaluation of a Sequential Antibiotic Treatment Regimen of
Ampicillin, Ciprofloxacin and Fosfomycin against Escherichia coli CFT073 in the Hollow Fiber
Infection Model Compared with Simultaneous Combination Treatment. Antibiotics (Basel) 11.:
https://doi.org/10.3390/antibiotics11121705

Ksiezopolska E, Schikora-Tamarit MÀ, Beyer R, et al (2021) Narrow mutational signatures drive
acquisition of multidrug resistance in the fungal pathogen Candida glabrata. Curr Biol 31:5314–
5326.e10

Kuchina A, Brettner LM, Paleologu L, et al (2021) Microbial single-cell RNA sequencing by split-pool
barcoding. Science 371.: https://doi.org/10.1126/science.aba5257

Larkins-Ford J, Degefu YN, Van N, et al (2022) Design principles to assemble drug combinations for
effective tuberculosis therapy using interpretable pairwise drug response measurements. Cell Rep
Med 3:100737

Lässig M, Mustonen V, Walczak AM (2017) Predicting evolution. Nat Ecol Evol 1:77

Lee JT, Coradini ALV, Shen A, Ehrenreich IM (2019) Layers of Cryptic Genetic Variation Underlie a
Yeast Complex Trait. Genetics 211:1469–1482

Levy SF, Blundell JR, Venkataram S, et al (2015) Quantitative evolutionary dynamics using high-
resolution lineage tracking. Nature 519:181–186

Li F, Salit ML, Levy SF (2018a) Unbiased Fitness Estimation of Pooled Barcode or Amplicon
Sequencing Studies. Cell Syst 7:521–525.e4

Li F, Tarkington J, Sherlock G (2023) Fit-Seq2.0: An Improved Software for High-Throughput Fitness
Measurements Using Pooled Competition Assays. J Mol Evol 91:334–344

Li Y, Petrov DA, Sherlock G (2019) Single nucleotide mapping of trait space reveals Pareto fronts that
constrain adaptation. Nat Ecol Evol 3:1539–1551

Li Y, Venkataram S, Agarwala A, et al (2018b) Hidden Complexity of Yeast Adaptation under Simple
Evolutionary Conditions. Curr Biol 28:515–525.e6

Logan A, Wolfe A, Williamson JC (2022) Antifungal Resistance and the Role of New Therapeutic
Agents. Curr Infect Dis Rep 24:105–116

Lupetti A, Danesi R, Campa M, et al (2002) Molecular basis of resistance to azole antifungals. Trends
Mol Med 8:76–81

Maltas J, Wood KB (2019) Pervasive and diverse collateral sensitivity profiles inform optimal strategies
to limit antibiotic resistance. PLoS Biol 17:e3000515

Martínez AA, Lang GI (2023) Identifying Targets of Selection in Laboratory Evolution Experiments. J
Mol Evol. https://doi.org/10.1007/s00239-023-10096-2

McInnes L, Healy J, Melville J (2018a) UMAP: Uniform Manifold Approximation and Projection for
Dimension Reduction. arXiv [stat.ML]

McInnes L, Healy J, Saul N, Großberger L (2018b) UMAP: Uniform Manifold Approximation and
Projection. J Open Source Softw 3:861

Melnikov SV, Stevens DL, Fu X, et al (2020) Exploiting evolutionary trade-offs for posttreatment
management of drug-resistant populations. Proc Natl Acad Sci U S A 117:17924–17931

Mira PM, Crona K, Greene D, et al (2015) Rational design of antibiotic treatment plans: a treatment

strategy for managing evolution and reversing resistance. PLoS One 10:e0122283

Mirkin B (2011) Choosing the number of clusters. Wiley Interdiscip Rev Data Min Knowl Discov 1:252–
260

Miyazaki H, Miyazaki Y, Geber A, et al (1998) Fluconazole resistance associated with drug efflux and
increased transcription of a drug transporter gene, PDH1, in Candida glabrata. Antimicrob Agents
Chemother 42:1695–1701

Moye-Rowley WS (2019) Multiple interfaces control activity of the Candida glabrata Pdr1 transcription
factor mediating azole drug resistance. Curr Genet 65:103–108

Nagasawa S, Kashima Y, Suzuki A, Suzuki Y (2021) Single-cell and spatial analyses of cancer cells:
toward elucidating the molecular mechanisms of clonal evolution and drug resistance acquisition.
Inflamm Regen 41:22

Nichol D, Rutter J, Bryant C, et al (2019) Antibiotic collateral sensitivity is contingent on the
repeatability of evolution. Nat Commun 10:334

Nyhoegen C, Uecker H (2023) Sequential antibiotic therapy in the laboratory and in the patient. J R Soc
Interface 20:20220793

Ogbunugafor CB (2022) The mutation effect reaction norm (mu-rn) highlights environmentally
dependent mutation effects and epistatic interactions. Evolution 76:37–48

Osset-Trénor P, Pascual-Ahuir A, Proft M (2023) Fungal Drug Response and Antimicrobial Resistance. J
Fungi (Basel) 9.: https://doi.org/10.3390/jof9050565

Paaby AB, Rockman MV (2013) The many faces of pleiotropy. Trends Genet 29:66–73

Paaby AB, White AG, Riccardi DD, et al (2015) Wild worm embryogenesis harbors ubiquitous polygenic
modifier variation. Elife 4.: https://doi.org/10.7554/eLife.09178

Pál C, Papp B, Lázár V (2015) Collateral sensitivity of antibiotic-resistant microbes. Trends Microbiol
23:401–407

Palmer AC, Kishony R (2014) Opposing effects of target overexpression reveal drug mechanisms. Nat
Commun 5:4296

Pedregosa F, Varoquaux G, Gramfort A, et al (2012) Scikit-learn: Machine Learning in Python. arXiv
[cs.LG] 2825–2830

Petti S, Reddy G, Desai MM (2023) Inferring sparse structure in genotype-phenotype maps. Genetics.
https://doi.org/10.1093/genetics/iyad127

Pinheiro F, Warsi O, Andersson DI, Lässig M (2021) Metabolic fitness landscapes predict the evolution
of antibiotic resistance. Nat Ecol Evol 5:677–687

Raymond B (2019) Five rules for resistance management in the antibiotic apocalypse, a road map for
integrated microbial management. Evol Appl 12:1079–1091

Read AF, Huijben S (2009) Evolutionary biology and the avoidance of antimicrobial resistance. Evol
Appl 2:40–51

Robbins N, Leach MD, Cowen LE (2012) Lysine deacetylases Hda1 and Rpd3 regulate Hsp90 function
thereby governing fungal drug resistance. Cell Rep 2:878–888

Rodrigues JV, Bershtein S, Li A, et al (2016) Biophysical principles predict fitness landscapes of drug
resistance. Proc Natl Acad Sci U S A 113:E1470–8

Roemhild R, Linkevicius M, Andersson DI (2020) Molecular mechanisms of collateral sensitivity to the
antibiotic nitrofurantoin. PLoS Biol 18:e3000612

Roe SM, Prodromou C, O'Brien R, et al (1999) Structural basis for inhibition of the Hsp90 molecular
chaperone by the antitumor antibiotics radicicol and geldanamycin. J Med Chem 42:260–266

Rybak JM, Cuomo CA, Rogers PD (2022) The molecular and genetic basis of antifungal resistance in the
emerging fungal pathogen Candida auris. Curr Opin Microbiol 70:102208

Scarborough JA, McClure E, Anderson P, et al (2020) Identifying States of Collateral Sensitivity during
the Evolution of Therapeutic Resistance in Ewing's Sarcoma. iScience 23:101293

Schmidt F, Efferth T (2016) Tumor Heterogeneity, Single-Cell Sequencing, and Drug Resistance.
Pharmaceuticals 9.: https://doi.org/10.3390/ph9020033

Somarelli JA, Gardner H, Cannataro VL, et al (2020) Molecular Biology and Evolution of Cancer: From
Discovery to Action. Mol Biol Evol 37:320–326

Sorgo AG, Heilmann CJ, Dekker HL, et al (2011) Effects of fluconazole on the secretome, the wall
proteome, and wall integrity of the clinical fungus Candida albicans. Eukaryot Cell 10:1071–1081

Su M, Satola SW, Read TD (2019) Genome-Based Prediction of Bacterial Antibiotic Resistance. J Clin
Microbiol 57.: https://doi.org/10.1128/JCM.01405-18

Tanaka S, Tani M (2018) Mannosylinositol phosphorylceramides and ergosterol coodinately maintain cell
wall integrity in the yeast Saccharomyces cerevisiae. FEBS J 285:2405–2427

Tan L, Chen L, Yang H, et al (2022) Structural basis for activation of fungal sterol receptor Upc2 and
azole resistance. Nat Chem Biol 18:1253–1262

Tenaillon O, Rodríguez-Verdugo A, Gaut RL, et al (2012) The molecular diversity of adaptive
convergence. Science 335:457–461

Thomas DS, Cisneros LH, Anderson ARA, Maley CC (2022) In Silico Investigations of Multi-Drug
Adaptive Therapy Protocols. Cancers 14.: https://doi.org/10.3390/cancers14112699

Uemura S, Moriguchi T (2022) Pleiotropic roles of N-glycans for enzyme activities and stabilities of
MIPC synthases, Csh1 and Sur1/Csg1, in Saccharomyces cerevisiae. Glycobiology 32:778–790

Vasicek EM, Berkow EL, Flowers SA, et al (2014) UPC2 is universally essential for azole antifungal
resistance in Candida albicans. Eukaryot Cell 13:933–946

Venkataram S, Dunn B, Li Y, et al (2016) Development of a Comprehensive Genotype-to-Fitness Map of
Adaptation-Driving Mutations in Yeast. Cell 166:1585–1596.e22

Ventola CL (2015) The antibiotic resistance crisis: part 1: causes and threats. P T 40:277–283

Vik Å, Rine J (2001) Upc2p and Ecm22p, dual regulators of sterol biosynthesis in Saccharomyces
cerevisiae. Mol Cell Biol 21:6395–6405

Vu BG, Moye-Rowley WS (2022) Nonidentical function of Upc2A binding sites in the Candida glabrata
CDR1 promoter. Genetics 222.: https://doi.org/10.1093/genetics/iyac135

Waller NJE, Cheung C-Y, Cook GM, McNeil MB (2023) The evolution of antibiotic resistance is
associated with collateral drug phenotypes in Mycobacterium tuberculosis. Nat Commun 14:1517

Wang W-Y, Cai H-Q, Qu S-Y, et al (2022) Genomic Variation-Mediating Fluconazole Resistance in
Yeast. Biomolecules 12.: https://doi.org/10.3390/biom12060845

Wang X, Zhang H, Chen X (2019) Drug resistance and combating drug resistance in cancer. Cancer Drug
Resist 2:141–160

Wang Y, Sang M, Feng L, et al (2023) A pleiotropic–epistatic entangelement model of drug response.
Drug Discov Today 103790

Ward JH (1963) Hierarchical Grouping to Optimize an Objective Function. J Am Stat Assoc 58:236–244

Whitesell L, Santagata S, Mendillo ML, et al (2014) HSP90 empowers evolution of resistance to
hormonal therapy in human breast cancer models. Proc Natl Acad Sci U S A 111:18297–18302

Wilson BA, Garud NR, Feder AF, et al (2016) The population genetics of drug resistance evolution in
natural populations of viral, bacterial and eukaryotic pathogens. Mol Ecol 25:42–66

Wortel MT, Agashe D, Bailey SF, et al (2023) Towards evolutionary predictions: Current promises and
challenges. Evol Appl 16:3–21

Xie JL, Polvi EJ, Shekhar-Guturja T, Cowen LE (2014) Elucidating drug resistance in human fungal
pathogens. Future Microbiol 9:523–542

Yadav A, Radhakrishnan A, Bhanot G, Sinha H (2015) Differential regulation of antagonistic pleiotropy
in synthetic and natural populations suggests its role in adaptation. G3 5:699–709

Yang F, Scopel EFC, Li H, et al (2023) Antifungal Tolerance and Resistance Emerge at Distinct Drug
Concentrations and Rely upon Different Aneuploid Chromosomes. MBio 14:e0022723

Yoon N, Krishnan N, Scott J (2021) Theoretical modeling of collaterally sensitive drug cycles: shaping
heterogeneity to allow adaptive therapy. J Math Biol 83:47

Yoshida M, Reyes SG, Tsuda S, et al (2017) Time-programmable drug dosing allows the manipulation,
suppression and reversal of antibiotic drug resistance in vitro. Nat Commun 8:15589

Zhao L, Liu Z, Levy SF, Wu S (2018) Bartender: a fast and accurate clustering algorithm to count
barcode reads. Bioinformatics 34:739–747

Zhao Q, Hautamaki V, Fränti P (2008) Knee Point Detection in BIC for Detecting the Number of
Clusters. In: Advanced Concepts for Intelligent Vision Systems. Springer Berlin Heidelberg, pp
664–673

1624