Microtubule reorganization during mitotic cell division in the dinoflagellate

Ostreospis **cf.** *ovata*

David Velasquez-Carvajal[1,2], Flavie Garampon[1], Sebastian Schaub[1] and Stefania

Castagnetti[1*]

[1] Sorbonne Universités, CNRS, Laboratoire de Biologie du Développement de

Villefranche-sur-Mer (LBDV), 06230 Villefranche-sur-Mer, France.

[2] current address: Grupo de investigación en neurociencias y envejecimiento. Facultad de

Ciencias de la Salud. Corporación Universitaria Remington. calle 51 n.° 51-27, Medellín,

Colombia.

* corresponding author: stefania.castagnetti@imev-mer.fr

Running title: Microtubules in dinomitosis

Keywords: dinoflagellates, microtubules, mitosis, cytokinesis, centrin

Summary statement

Our study describes special features of mitosis and cytokinesis in dinoflagellates and uncovers a new alternative mechanism for cell division, highlighting the plasticity of cell biological process in eukaryotic cells.

Abstract

Dinoflagellates are marine organisms that undergo seasonal explosive proliferation events known as algal blooms. Vegetative cell proliferation is a main contributing factor in these events. However, mechanistical understanding of mitosis and cytokinesis in dinoflagellate remains rudimentary. Using an optimized immunofluorescence protocol, we analysed changes in microtubule organization occurring during the mitotic cycle of the toxic dinoflagellate *Ostreopsis* cf. *ovata.* This study revealed important features of dinoflagellate cell division. We find that the two flagella and the cortical microtubule array persist throughout the mitotic cycle. Two microtubule bundles are present in the cytoplasm originating from the ventral area, where the basal bodies are located: a cortical bundle and a cytoplasmic ventral bundle. The latter associates with the nucleus in the cell centre in preparation for mitosis and with the acentrosomal extranuclear spindle during mitosis. Analysis of tubulin post-translational modifications identifies two populations of spindle microtubules: polar acetylated microtubules whose length is stable throughout mitosis and central tyrosinated microtubules which elongate during chromosome segregation. During cell division a microtubule rich structure forms along the dorsal-ventral axis, associated with the site of cytokinesis, consistent with a cytokinetic mechanism independent of the actomyosin ring typical of animal and yeast cells.

Introduction

Dinoflagellates are unicellular eukaryotes that have adapted to most aquatic ecosystems, from tropical to polar waters. These protists are known to undergo large population expansions that can impact heavily on the marine ecosystem and on human health, a phenomenon known as Harmful Algae Bloom (HAB) (Driscoll et al., 2016). Despite the extensive investigations into the ecological and cellular basis of HABs, the mechanisms controlling dinoflagellate cell proliferation, a major process underlying HAB development, still remain elusive.

Dinoflagellate mitosis, also known as dinomitosis, is unique among eukaryotes. It is a form of closed mitosis, where chromosome segregation occurs within the nucleus, without nuclear envelope breakdown. Differently from yeasts, diatoms and euglenoids, which undergo closed mitosis with a nuclear spindle, in dinoflagellates spindle microtubules are cytoplasmic and traverse the nucleus through tunnels never entering the nucleoplasm (Gavelis et al., 2019). Segregating chromosomes are attached to the nuclear membrane surrounding the nuclear tunnels, through electron dense kinetochore-like structures, which in turn contact microtubules on the cytoplasmic side (Cachon and Cachon, 1977; Oakley and Dodge, 1974). Hence, the interaction between spindle microtubules and segregating chromosomes is indirect, across the nuclear membrane. This configuration poses the question of how the spindle is organized in these organisms and how chromosomes are segregated within the nuclear membrane. EM studies in *Syndinium* (Ris and Kubai, 1974) and in *Amphidinium carterae* (Oakley and Dodge, 1976) showed that the microtubules emanating from the kinetochore-like structures extend exclusively towards the nearby polar region of the spindle and the kinetochore to pole distance remains constant throughout mitosis. Thus, chromosome segregation was suggested to be accomplished by elongation of cytoplasmic microtubules running through the nuclear channels between the two spindle poles

Following nuclear division, the cell divides during cytokinesis. In animal and yeast cells, cytokinesis relies on the constriction of an actomyosin ring. The absence of identifiable myosin-II in many bikonts (Richards and Cavalier-Smith, 2005), including dinoflagellates,

however, suggests that an alternative mechanism for cytokinesis must be in place in these organisms. In plants a transient ring of cortical microtubules and actin filaments, the preprophase band, forms early in the cell cycle and marks the site of cytokinesis. During anaphase an equatorial array of microtubules, the phragmoplast, then directs Golgi-derived vesicles to the cortical division site to allow deposition of the new cell wall (Nishihama and Machida, 2001). Mechanisms of cytokinesis that do not involve an actomyosin ring have also been described for many single-celled protists. In *Trypanosoma brucei* cell division occurs by microtubule dependent formation of a unidirectional furrow from the anterior to the posterior of the cell (Sherwin et al., 1989). In *Chlamydomonas reinhardtii* and many related green algae, a microtubule structure, called the phycoplast, forms between the duplicated basal bodies at the anterior of the cell and then grows towards the posterior pole (Ehler and Dutcher, 1998). Actin also accumulates at this site, but treatment with actin depolymerizing drugs does not induce any defect in cell division (Onishi et al., 2020). For dinoflagellates, little is known on how cytokinesis takes place and the contribution of the cytoskeleton to this process. Early EM studies in *A. carterae* (Oakley and Dodge, 1976) and in *Ceratium tripos* (Wetherbee, 1975) showed that cell division occurs by pinching in of the membrane in the equatorial region where microtubules accumulate under the plasma-membrane. Treatment with actin depolymerizing drugs, however, was shown to interfere with completion of cell division in *Prorocentrum micans*, suggesting a contribution of the actin cytoskeleton to cytokinesis, at least in some species (Schnepf et al., 1990).

To gain further insights into the mechanism of dinomitosis and the organization of the mitotic spindle in dinoflagellates, we analysed the spatial and temporal changes in cytoskeletal organization during the mitotic cycle in the dinoflagellate *Ostreopsis* cf. *ovata* (*O.* cf. *ovata*), one of the most common HAB causing species in tropical and subtropical areas (Parsons et al., 2012; Rhodes, 2011) and the most abundant and widely distributed benthic dinoflagellate in the Mediterranean Sea (Ninčević Gladan et al., 2019). *O.* cf. *ovata* cells have a teardrop shape with a pointy ventral side, from which two flagella emanate and a rounded dorsal side, where the nucleus is located during interphase. In interphase cells, a cortical array of microtubules underlies the plasma membrane (cortical microtubule

array) and a microtubule bundle runs from the ventral sulcal area towards the inside of the

cell (Escalera et al., 2014). Here we show that the cortical microtubule array and the flagella

persist throughout the cell cycle. We also analyse changes in the organization of

microtubule-based structures during the mitotic cycle and describe novel ultrastructural

features of the microtubule cytoskeleton as cells undergo mitosis and cytokinesis. We find

that microtubules of the ventral bundle contact the nucleus while it localizes to the centre

of the cell in preparation for mitosis. An acentrosomal microtubule-based spindle then

forms in the cell centre, transverses the nucleus and separates chromosomes. Analysis of

tubulin posttranslational modifications identifies two regions in the anaphase spindle: a

stable polar spindle enriched in acetylated tubulin, linking chromosomes to the ventral

bundle and a dynamic central spindle, containing tyrosinated tubulin, between separating

chromosome masses. In dividing cells, a microtubule-rich structure, free of actin, forms

from the dorsal side along the longitudinal axis of the cell where cytokinesis takes place,

and divides the cell. Our observations reveal that both mitosis and cytokinesis in *O*. cf.

ovata are different from yeast and animal cells and support the idea that the mechanisms

of cell division are highly diverse outside of opisthokonts.

Results

Cortical microtubule array and flagella are maintained throughout the cell cycle

To analyse the changes in the organization of the microtubule cytoskeleton associated with

the mitotic cell cycle in *O*. cf. *ovata* we have optimized an immunofluorescence protocol

previously developed to stain cortical microtubules (Escalera et al., 2014). We applied it

to label microtubules in both cultured cells and wild samples collected in the Bay of

Villefranche-sur-mer during summer blooming and obtained similar results with both

sources of cells. Staining using the ß-tubulin D66 antibody labelled several structures in

the cell (Fig. 1A). Previous description of microtubules in *O*. cf. *ovata* had identified a

network of microtubules underlying the plasma membrane in interphase cells (Escalera et

al., 2014). We confirmed the presence of this cortical microtubule array (Fig. 1A and movie

1). On the posterior side of the cell, cortical microtubules ran vertically along the dorso-

ventral axis covering the entire surface of the cell. On the opposite anterior side, recognized
by the presence of the apical pore, instead, cortical microtubules run obliquely from the
apical pore to the side of the cell. The anterior and posterior halves are separated by the
cingulum, which circles the entire cell (Fig. 1A). The cingulum is delimited by two parallel
bands of microtubules, while in the groove of the cingulum tightly packed parallel
microtubules run transversally, perpendicular to the lateral bands (Fig. 1A, lateral view).
Two previously identified thick bundles of MTs originating from microtubular roots
present in the ventral sulcal area, can also be observed: one close to the cell surface on the
posterior side of the cell, which we named cortical bundle (CB), and a second deeper in the
cytoplasm, herein referred to as ventral bundle (VB, Fig. 1A and movie 1). Specificity of
the observed tubulin staining was confirmed by treatment of cells with the microtubule
depolymerizing drug colchicine for 1 hour prior to fixation. All microtubules were lost in
colchicine treated samples but not in control (dH2O) samples (Fig. 1B).

Using bloom samples collected between 12pm and 4am, which as previously showed are
enriched in mitotic and dividing cells (Pavaux et al., 2021), we then asked whether cortical
microtubules were maintained throughout the cell cycle. In *O*. cf. *ovata* different cell cycle
stages can be identified based on morphological features and on the characteristic position
of the nucleus within the cell (Bravo et al., 2012; Pavaux et al., 2021). The nucleus which
is located in the dorsal area during interphase (non-dividing cells) relocates to the cell
centre in preparation for mitosis (pre-mitotic cells), where the nucleus divides (mitotic
cells). Following mitosis binucleated cells divide (cytokinesis) by fission and, using
brightfield microscopy a furrow can be observed bisecting the cell to give rise to two post-
dividing cells recognized by the presence of the nucleus in lateral position.

We observed that the cortical array of microtubules is stable and persists throughout the
cell cycle: in interphase, pre-mitotic, mitotic and post-dividing cells (Fig. 1C). Similarly,
flagella, emanating from the ventral pole, were present at all stages of the cell cycle,
including mitosis and cytokinesis (Fig. S1). This is in contrast with what was previously
described in another dinoflagellate species *Crypthecodinium cohnii* (Perret et al., 1993) or

in other algal species like *C. reinhardtii*, where flagella are absent during mitotic cell
division (Cross and Umen, 2015; Rosenbaum et al., 1969).

Changes in microtubule organization during mitosis and cytokinesis

In most interphase cells (> 95%), the two microtubule bundles (VB and CB) extend for a
third to half of the cell length towards the cell centre (Fig. 2A, interphase). The cortical
bundle is a stable thick bundle of microtubules most likely associated with the flagellar
apparatus to anchor the basal bodies to the cell cortex, comparable to the rootlet
microtubule of *C. reinhardtii* (Rosenbaum et al., 1969). The ventral bundle instead is a
more dynamic structure which changes during the cell cycle. During interphase,
microtubules of the VB are compacted for most of the length of the bundle but thin
ramifications can be observed extending in the cell centre (Fig. 2A, interphase). In few
interphase cells (< 5%), the ventral bundle extends past the cell mid-line and reaches the
nucleus in dorsal position (Fig. 2A, interphase). In pre-mitotic cells, when the nucleus is
positioned in the cell centre, microtubules branching from the ventral bundle contact the
nucleus, forming a structure that resembles an inverted "Y" (Fig. 2A and Fig. S2, pre-
mitotic) and enveloping the ventral side of the nucleus.

In mitotic cells, microtubules then organize in 6 to 8 parallel bundles (n= 5 cells) that align
along the short axis of the cell to form the mitotic spindle (Fig. 2A and Fig. S2,
pro/metaphase), suggesting the presence of 6-8 nuclear tunnels. Spindle microtubules
converge at two opposite poles which are connected to the ventral area of the cell by thin
microtubule bundles (Fig. 2A), probably originating from splitting of the ventral bundle
present in interphase cells. No astral microtubules are visible in mitotic cells. As previously
reported for other dinoflagellate species (Bhaud et al., 2000), we observed that in early
mitosis, the DNA concentrates in the central region of the spindle onto a metaphase-like
plate (Fig. 2A, pro/metaphase), before separating into two discrete masses during anaphase
(Fig. 2A and Fig. S2, anaphase). Measurement of pole to pole distance showed that spindle
length is constant prior to chromosome segregation (pro/metaphase; average $\pm$ SD: 12,1 $\pm$
1,3 µm) and spindle elongation is observed only following chromosome segregation in
anaphase when spindle length extends up to 27,5 µm (Fig. 2B). However, we observed that

pole to DNA distance, measured as distance between the middle of each DNA mass and

the corresponding spindle pole, remains constant during mitosis from prophase to telophase

(Fig. 2C; average ± SD: pro/metaphase: 5,95 ± 0,6 µm; anaphase: 5,7± 0,6 µm; telophase:

6,1± 0,5 µm). Thus, chromosome segregation appears to be driven exclusively by extension

of the central spindle region.

As shown in Fig. 2A (anaphase, arrowhead), in many anaphase cells, microtubules also

accumulate in the dorsal area of the cell. By analysing cells at different mitotic stages, we

noticed that a dorsal enrichment in microtubules could be observed already at earlier

mitotic stages. This microtubule-rich structure corresponds to the site of cytokinesis (Fig.

2A and Fig. S2, telophase/cytokinesis and Fig. 3A). Three-D reconstruction of cells at

different stages of mitosis showed that this microtubule structure develops progressively.

In early mitotic cells, up to anaphase, microtubules first arrange in a ring, perpendicular to

the cingulum, under the cell surface (Fig. 3A, metaphase/anaphase). As cells progress

through anaphase microtubules fill the space between the dividing cells (Fig. 3A,

telophase) and by the end of mitosis, when chromosomes are fully segregated to opposite

sides of the cell and the central spindle is no longer visible, microtubules fill the entire

surface between the two daughter cells, forming a dense microtubule plate (Fig. 3A,

cytokinesis).

The presence of a microtubule rich plate at the site of cell division suggests that cytokinesis

is driven by microtubules in *O*. cf. *ovata*. This is consistent with observations in other

protists where microtubule-rich structures drive cell division and are essential for

cytokinesis (Ehler and Dutcher, 1998; Francia et al., 2015; Hammarton, 2019; Onishi et

al., 2020; Sherwin et al., 1989). Although microtubule enrichment at the cleavage furrow

was already observed in dinoflagellates, actin is thought to be required for completion of

cell division in some dinoflagellates species, like *P*. *micans* (Schnepf et al., 1990). We

therefore asked whether actin microfilaments are enriched at the cytokinetic plate. Similar

to what was previously observed in other dinoflagellate species (Schnepf, 1988; Villanueva

et al., 2014), staining with phalloidin showed a diffuse cytoplasmic grid-like pattern (Fig.

3B) which was lost following 5 minutes of treatment with the actin depolymerizing drug

latrunculin-A (10 μM, Fig. 3C). Interestingly latranculin A treated cells quickly lost their

shape and often become round within 5 minutes of treatment. In untreated cells, we

observed actin fibres connecting the cell cortex to the cell centre, where the phalloidin

staining is more concentrated. In all cells, irrespectively of cell cycle stage, actin also

accumulated around the nucleus (Fig. 3B), as previously described in *C. cohnii* (Perret et

al., 1993) and *L. polyedra* (Schnepf, 1988; Stires and Latz, 2018). No discrete actin-

containing structure was observed at the site of division, further supporting the existence

of a microtubule-based mechanism both for chromosome segregation and for cytokinesis

in *O.* cf. *ovata*.

Different microtubular sub-populations drive cell division

All microtubules are formed by α/ß-tubulin heterodimers whose properties are modulated

by post-translational modifications (PTMs) (Bulinski et al., 1988; Janke, 2013; Wolff et

al., 1992). Among all identified PTMs, acetylation, tyrosination and poly-glutamylation

(Fig. 4A) appear to be conserved across most eukaryotes. We reasoned that analysing

tubulin PTMs would allow to distinguish between different sub-populations of

microtubules and their changes during the cell cycle, providing insight into how the

cytoskeletal structures described above are generated and are related to one another. We

therefore analysed tubulin tyrosination, acetylation and poly-glutamylation during *O.* cf.

ovata cell division (dinomitosis and cytokinesis). We first tested whether these tubulin

PTMs could be detected in *O.* cf. *ovata* protein extracts by western blot using specific

antibodies. We used the YL1/2 monoclonal antibody which recognizes the C-terminal EEF

sequence of yeast α-tubulin (Kilmartin et al., 1982), conserved in *O.* cf. *ovata* α-tubulin; a

monoclonal antibody (T6793) which specifically recognizes acetylated lys40 of α-tubulin,

and the antibody GT335 against mouse poly-glutamylated tubulin (Wolff et al., 1992). As

shown in Fig. 4B, YL1/2 and T6793 antibodies recognized a single protein of the same

size as α-tubulin.

Tyrosinated α-tubulin corresponds usually to newly polymerized and dynamic

microtubules, whereas removal of the carboxy-terminal tyrosine residue (detyrosination)

marks microtubule aging and stabilization (Bulinski et al., 1988; Schulze et al., 1987).

Immunolabelling of cycling *O*. cf. *ovata* cells using the YL ½ antibody showed that

tyrosinated tubulin is present in the flagella, in the cortical array and in the ventral bundle

at all cell cycle stages (Fig. 4C, interphase and Fig. S3A). In mitotic cells both in early

(pro/metaphase), and late (anaphase) stages, the YL ½ antibody labelled the central region

of the spindle near the midplane, but no signal was detected in the polar spindle regions

until telophase (Fig. 4C, mitosis).

Acetylation of lysine 40 (lys40) on α-tubulin instead was shown to be often associated with

stable long-lived microtubules (Eshun-Wilson et al., 2019). Immunostaining with the

T6793 antibody showed that, at all cell cycle stages, both the cortical array and flagella

contain acetylated microtubules (Fig. S3B). Differently from tyrosinated tubulin, in early

mitotic cells, before chromosome segregation (pro/metaphase), tubulin acetylation extends

over the entire length of the spindle, including the polar region. However, in anaphase cells,

following chromosome segregation, acetylated α-tubulin is restricted to the polar region of

the spindle which is associated with the DNA (Fig. 4C), but is absent in the central spindle

(between the segregating chromosomes). Acetylated-tubulin was also detected at the site

of cytokinesis both in the early-cytokinetic ring and in the late-cytokinetic plate (Fig. 4C).

Co-staining of *O*. cf. *ovata* cells for acetylated and tyrosinated tubulin did not allow to

identify any difference in the cortical array (Fig. S3C), but confirmed that acetylated

tubulin is enriched in the polar region of the spindle but absent from the central spindle

when the spindle elongates in anaphase (Fig. 4D).

Finally, poly-glutamylation, which is the addition of glutamate side chains of variable

length to the carboxy-terminal tail of both α- and ß-tubulin (Bodakuntla et al., 2021; Eddé

et al., 1990; RadiBer et al., 1992), is thought to modulate the interaction between

microtubules and their partners, MAPs (Microtubule Associated Proteins) and molecular

motors, affecting cargo movement, microtubule stability and severing (Bodakuntla et al.,

2021). Using the antibody GT335 against glutamylated tubulin (Wolff et al., 1992) we

showed that in *O.* cf. *ovata* glutamylated-tubulin is undetectable in non-dividing cells (Fig.

S3E, interphase). We could observe tubulin glutamylation exclusively in mitotic cells,

starting from early mitosis. As for tyrosinated tubulin, glutamylated tubulin was enriched

in the central spindle (Fig. S3E). Differently from acetylated and tyrosinated tubulin,

glutamylated-tubulin is not present at the site of cytokinesis.

Centrin accumulates exclusively at basal bodies

In eukaryotic cells centrioles and basal bodies are essential for the organization of the

microtubule cytoskeleton during interphase and mitosis (Wu and Akhmanova, 2017). A

well conserved component of both these structures is centrin, a calmodulin-like Ca^{2+}-

binding protein present in protists, yeast and animal cells. In the green algae *C. reinhardtii*,

where it was first identified, centrin localizes to the region of the basal body associated

with the flagellum axoneme during interphase, and re-localizes to spindle poles during

mitosis to organize the mitotic spindle (Salisbury et al., 1988). As genomic data are not

available for *O.* cf. *ovata* we generated transcriptomic data from cultured *O.* cf. *ovata* cells

(strain MCCV54) during exponential growth and searched for centrin homologues.

Analysis of this transcriptomic dataset allowed the identification of a centrin transcript in

O. cf. *ovata* with 55% identity to human centrin and 53% to *C. reinhardtii* centrin (Fig.

5A). Using a polyclonal antibody against *C. reinhardtii* centrin (20H5), we then sought to

identify microtubule organizing centres (MTOCs) in *O.* cf. *ovata* cells. In western blot

assays the 20H5 antibody recognizes a single *O.* cf. *ovata* protein of the expected (20kDa)

molecular weight (Fig. 5B). In cells the 20H5 antibody intensely labelled a structure in the

ventral region of the cell, near the flagella insertion site where the basal bodies are located

(Fig. 5C). In line with our observation that flagella are maintained during the entire cell

cycle (Fig. S1), we observed that centrin, which is required for flagellar motility (Fenchel,

2001), was maintained in the ventral area throughout mitosis and cytokinesis. Consistently,

we did not observe accumulation of centrin at spindle poles at any stage of mitosis,

supporting earlier EM studies suggesting that dinoflagellate spindles are acentriolar. The

centrin antibody also stained fibrous structures which extend from the basal bodies towards

both the cell centre and the cell cortex (Fig. 5C).

Based on our observations we cannot determine when basal body duplication takes place.

However, as shown in Fig. 5C we observed that the two centrin-labelled basal bodies are

juxtaposed in interphase, separate in mitosis and then segregate to the two daughter cells

in cytokinesis. Quantification of the distance between the two peaks of fluorescent intensity

in centrin-labelled cells confirmed this observation (Fig. 5D), showing that the two basal

bodies separate when the spindle forms as the distance between the two peaks remains

constant in interphase (average $\pm$ SD: 1,4 $\pm$ 0,4µm) and pre-mitotic cells (1,4 $\pm$ 0,3µm) but

increases to 2,7 µm ($\pm$ 0,5 µm) in mitotic cells containing a fully formed spindle. In

telophase the cytokinetic plate then forms between the two fully segregated basal bodies

(5,6 $\pm$ 1,7 µm), generating two daughter cells with their own flagellar apparatus.

Discussion

Due to the unique characteristics of mitosis in dinoflagellates, the term dinomitosis has

been used to describe how chromosomes are segregated in those organisms. Here, we have

analysed cytoskeletal changes associated with dinomitosis and cytokinesis in the

dinoflagellate *O.* cf. *ovata*. which are summarized in Fig. 5 E. In interphase cells the

nucleus is located near the dorsal side of the cell and does not interact with microtubules

of the ventral bundle. Cortical microtubules run under the plasma membrane and as

previously shown for another dinoflagellate, *C. cohnii* (Perret et al., 1993), they persist

throughout the cell cycle, including mitosis and cytokinesis. This is similar to what was

observed in other unicellular organisms, such as *Trypanosoma brucei*, where the

subpellicular microtubule corset which is required for cell shape (Sinclair et al., 2021), is

maintained during cell division and is only remodelled locally at the site of furrow

formation during cytokinesis (Gull, 1999).

The first sign of preparation for mitosis in *O.* cf. *ovata* is the re-localization of the nucleus

to the centre of the cell. We observed that while in the cell centre, the nucleus is always

associated with microtubules of the ventral bundle and in few cells this association is also observed while the nucleus is in dorsal position. This configuration suggests that nuclear re-positioning is microtubule based and that extension and branching of the microtubule bundle towards the dorsal side of the cell might be a pre-requisite for nucleus-microtubule interaction and could be considered as an early indicator of preparation for mitosis. Centrin filaments are also associated with the ventral bundle and with the basal bodies where the ventral bundle originates from. It is therefore possible that nuclear movement relies not only on the mechanical action of microtubules and associated motors but also on the contractile activity of centrin, as already observed in nuclear repositioning during mitosis in *C. reinhardtii* (Salisbury et al., 1988). In this organism the two flagella are reabsorbed prior to division and during prophase two major microtubule fibres extend from the basal bodies toward the nucleus and reposition it from the middle of the cell to the apical side, where the basal bodies are located. Centrin which forms filamentous structures that link the basal bodies to the nucleus (Geimer and Melkonian, 2005; Wright et al., 1989) provides contractility to the microtubule bundles and contraction of centrin fibres at mitotic entry allows nuclear movement towards the apical cell surface (Salisbury et al., 1988). This movement is followed by separation of the basal bodies in prophase. Indeed, we observe a similar separation of centrin-labelled bodies in early mitotic stages in *O.* cf. *ovata*.

Upon localization of the nucleus to the cell centre, the mitotic spindle forms, with 6-8 spindle microtubule bundles traversing the nucleus along the short axis of the cell, in line with previous observations in other gonnyalucoid species, the dinoflagellate clade with the highest number of nuclear tunnels (Gavelis et al., 2019). Throughout mitosis, the spindle remains connected to the cell ventral cortex through two microtubule fibres that probably originate from splitting of the ventral bundle. This configuration is similar to the three-pronged fork described in *C. cohnii* where the spindle poles are connected to the basal bodies by a bundle of microtubules, the desmoses (Perret et al., 1993). In animal, yeast and several algal cells, spindle formation is usually driven by the centrosomes localized at spindle poles. Although more markers will need to be analysed, the geometry of the spindle with the association of spindle poles to the microtubules of the ventral bundle and the absence of centrin accumulation at spindle poles suggests that *O.* cf. *ovata* spindles are

acentrosomal. Indeed, EM studies in other dinoflagellate species have never identified
centrioles in the vicinity of the nuclear membrane, with the only exception of Syndiniales
(Bhaud et al., 2000; Drechsler and McAinsh, 2012; Moon et al., 2015). Acentrosomal
spindle formation has been observed in animal and plant cells. In those cells spindle
organization is driven by chromosomes, often through the formation of a local gradient of
RanGTP (Deng et al., 2007; Lee and Liu, 2019). However, this is difficult to envisage for
dinomitosis as spindle microtubules are cytoplasmic and indirectly contact the
chromosomes which are connected to the inner side of the nuclear membrane.

Our analysis of microtubule PTMs allowed the identification of two regions in anaphase
spindles with different characteristics: a polar region highly acetylated and constant in
length (constant pole-DNA distance), which is anchored to the ventral area by the ventral
bundle, and a central spindle region which is instead enriched in tyrosinated tubulin and
whose length increases during mitosis (Fig. 4D,E). Similar to kinetochore microtubules in
animal spindles (Wilson and Forer, 1989), the stable acetylated microtubules link
chromosomes to spindle poles and most likely correspond to the k-fibers observed by EM
both in other dinoflagellates (Oakley and Dodge, 1976; Ris and Kubai, 1974) and in the
flagellate hypermastigote *Barbulanympha* (Inoué and Ritter, 1978; Ritter et al., 1978).
Different from most animal spindles, instead, the interpolar microtubules running between
opposite spindle poles, correspond to the tyrosinated microtubules described here.
Interestingly the tyrosinated microtubules never reach the spindle poles. In anaphase
spindles, tyrosinated microtubules appear to overlap with acetylated microtubules only in
the vicinity of the segregating chromosomes, possibly at the level of the kinetochores.
Based on these studies we suggest that during pro/metaphase of dinomitosis mitotic
chromosomes become anchored to opposite spindle poles by stable acetylated polar
microtubules. During anaphase polymerization of the more dynamic tyrosinated interpolar
microtubules then drives chromosome segregation (anaphase B) without prior poleward
chromosome movement towards the spindle poles, thus suggesting a lack of anaphase A.
The exact organization of the central microtubule array as well as the nature of the
molecular motors generating the pushing force remains unclear. Further studies will be
required to determine whether chromosome segregation relies on microtubule

polymerization or sliding of interdigitating central spindle microtubules. Either way, this mechanism is reminiscent of the central-spindle driven chromosome segregation described during meiosis in *C. elegans* oocytes (Laband et al., 2017) and during mitosis in *C. elegans* embryos without centrosomes (Cowan and Hyman, 2004). In both cases, kinetochore microtubules act as a scaffold to anchor chromosomes to spindle poles and correctly localize them onto the metaphase plate during spindle polarization (Pitayu-Nugroho et al., 2023), while polymerization of microtubules of the central spindle generates an outward pushing force to allow chromosome segregation. The presence of this mechanism in distantly related nematodes and dinoflagellates suggests that it might have emerged early in evolution and could represent a primitive microtubule-based mechanism for chromosome segregation which predates the evolution of cortical interactions to generate pulling forces and ensure higher accuracy in chromosome segregation. Indeed, a similar mechanism can also be observed in bacteria where segregation of low copy number plasmids relies on the polymerization of antiparallel ParM filaments, whose elongation pushes plasmid clusters towards the poles of the cell, allowing their segregation before cell division (Gayathri et al., 2012).

Starting from anaphase, a microtubule-rich structure is formed from the dorsal side, opposite the basal bodies. This structure coincides with the cleavage plane, suggesting a direct implication for microtubules either in placement of the cleavage plane and/or cytokinesis. This observation is in line with the idea that actomyosin driven cytokinesis is a novel acquisition of unikont (animal, yeast and amoeba) cells. Indeed, the lack of Myosin II in most bikont species (Richards and Cavalier-Smith, 2005), including dinoflagellates, supports the use of alternative mechanisms for cytokinesis. In many unicellular protists and in plant cells microtubules have a central role in cytokinesis and form specialized structures associated with the site of cytokinesis. In plants phragmoplast microtubules form in the centre of the cell and then expand centrifugally towards the cell membrane. In *C. reinhardtii* and in other related green algae, instead, cytokinesis is associated with a special array of microtubules, the phycoplast, which forms at the anterior end of the cell where the microtubule rootlets are located and grows unidirectionally towards the opposite pole of the cell (Ehler and Dutcher, 1998). The cytokinetic plate we described here for *Ostreopsis*

bears similarities with the algal phycoplast as it forms at the dorsal side and then grows in a unidirectional fashion. However differently from *C. reinhardtii*, the cytokinetic plate described here originates opposite to the basal bodies and grows towards them, suggesting that, as already observed for the plant phragmoplast, the mechanism of microtubule nucleation in the dinoflagellate cytokinetic plate is most likely independent of centrosomes and basal bodies.

Materials and Methods

Cell culture and bloom sampling

Strains of O. cf. *ovata* (MCCV054) obtained from the Mediterranean Culture Collection of Villefranche-sur-Mer (MCCV), EMBRC-France, were cultured in autoclaved and filtered enriched seawater, at a salinity of 38 g/L, as described by Guillard and Hargraves (L1 enriched seawater medium) (Guillard and Hargraves, 1993). Cultures were incubated at 22 °C under a 14:10 light/dark cycle (250 μmol m-2 s-1). Cells were fixed for staining during the proliferative phase (day 6 to 10) in order to obtain the greatest number of cells in division.

Bloom samples were collected in Villefranche-sur-mer, on the Mediterranean French coast (43°41′34.83″ N and 7°18′31.66″ E), where yearly blooms occur. Samples used for immunofluorescence were collected in 2018, 2019 and 2020 between 12:00 pm and 4:00 am (Pavaux et al., 2021).

Actin staining and immunofluorescence

For immunofluorescence a previous published protocol was modified (Escalera et al., 2014). Cells were fixed in cold fixative solution (80% methanol, 10% DMSO (Sigma-Aldrich) and 0,5 mM EGTA pH 7.5 (from 0,5 M EGTA pH 7.5 stock solution in distilled water)) for at least 24 hours, at -20ºC to improve membrane permeabilization. Samples could be maintained in fixative solution for several months. When needed, samples were rehydrated progressively in methanol-PBS (60%, 40%, 20%) and finally in PBS, with a 10 minutes incubation at each step. Cells were then permeabilized by two consecutive

incubations in PBT (PBS with 0.1% Triton X-100) for 15 minutes at room temperature

(RT). Cells were recovered by centrifugation, 5 minutes at 0.2 RCF, and then blocked for

30 minutes at RT in PBT containing 5% BSA (PBT-BSA) and then incubated at 4°C in

PBT-BSA containing primary antibody at the appropriate dilution for 3-6 days to maximize

antibody penetration. With incubations shorter that 2 days structures deep into the

cytoplasm were not labelled. For microtubule staining, D66 anti β-tubulin antibody

(against tubulin from *Lytechinus pictus*, mouse, Sigma-Aldrich) was used at 1:400 dilution;

for acetylated tubulin, T6793 antibody (Clone 6- 11B-1, mouse, Sigma-Aldrich) was used

at 1:400 dilution; for tyrosinated tubulin, YL1/2 antibody (ab6160; rat, Abcam) was used

at 1:400 dilution; for polyglutamylated tubulin, monoclonal antibody anti

polyglutamylated (clone GT335, mouse, Adipogen); the anti-centrin antibody (clone

20H5; mouse, Merk) was used at 1:100 dilution.

After incubation with the primary antibody, cells were washed three times in PBT, and

then incubated over-night at 4°C in PBT-BSA with the appropriate secondary antibody at

a 1:400 dilution (fluorescently-conjugated anti-mouse and anti-rat antibodies, Jackson

ImmunoResearch). After over-night incubation, cells were washed twice with PBS, and

then incubated for 10 minutes with 1μg/ml Hoechst-33342 (Sigma-Aldrich) to label the

DNA, rinsed twice with PBS and finally resuspended in 100-200 μl of PBS. 50 μl of cells

were mixed with 50 μl of citiflour AF1 (Science Services) and mounted on slides for

imaging. Each experiment was repeated 3-5 times, 25-50 cells were counted for each

sample and 5-15 images were collected for further analysis.

Actin filaments were visualized using CytoPainter Phalloidin-iFluor 488 Reagent

(ab176753, Abcam). Cells were fixed in 4% paraformaldehyde in PBS for at least 24 hours

at 4°C. Cells were then washed three times in PBS and permeabilized as described for

immunofluorescence. After a 30-minute incubation in PBT-BSA, cells were then incubated

in PBT containing phalloidin at 1:400 dilution, for 1 hour at RT in the dark. Cells were

then washed twice in PBS, and mounted in citiflour AF1 (Science Services) for immediate

imaging.

Images were acquired with Leica SP8 and Stellaris 5 microscopes both equipped with a 63X/1.4 Oil and a 40X/1.1W objective, which are part of the Plateforme d'Imagerie Microscopique (PIM) of the Institute de la mer de Villefranche sur mer. On stellaris 5 images were deconvolved with Lightning. When present, Z-Drift was corrected using a custom-made tool (https://github.com/SebastienSchaub/SatAndDriftEstimator). Spindle measurements and image analysis were performed in Fiji (49) with home-made 3D distance tool (https://github.com/SebastienSchaub/3DtDist) and visualization on Imaris (Imaris, BitPlane, Zurich, Switzerland).

Protein extraction and western blot analysis

To test antibodies, proteins were extracted from cells of strain MCCV-054. 50 ml of cells were grown in L1 media and collected 14 days after inoculum by filtration through Durapore 5 μm PDVF membranes (Millipore) using a manifold system. Cells were recovered from the filters in 1 ml of PBS and then pelleted by centrifugation at 5000rpm for 5 minutes. Pellets were then fast frozen in liquid nitrogen, then resuspended in an equal volume of 2x Laemmli (100 mM Tris-HCl at pH 6.8, 4% SDS, 0.2% bromophenol blue, 20% glycerol, 200 mM dithiothreitol) and heated at 100ºC for 5 minutes. Samples were separated on a 10% SDS-polyacrylamide gel and then transferred to a nitrocellulose membrane. Membranes were blocked for 1h in Tris Buffer-Saline (TBS: 20 mM Tris Base, 150 mM NaCl) containing 2% milk, 0.1% tween-20 (TBS-Tween-milk) and then incubated in blocking solution containing the appropriate antibody overnight at 4°C. For centrin detection the 20H5 antibody (Merck) was diluted 1:2000, for detection of tubulin (DM1a antibody, Sigma-Aldrich, mouse) and of PTMs (tyrosinated tubulin: YL ½ antibody, ab6160; rat monoclonal, Abcam; acetylated tubulin: T6793 antibody, clone 6-11B-1, mouse, Sigma-Aldrich; polyglutamylated tubulin: GT335, mouse, Adipogen) all antibodies were diluted 1:3000. Following antibody incubation, membranes were washed 3 times with TBS-Tween (TBS with 0.1% tween-20) and incubated with an appropriate horseradish peroxidase-conjugated secondary antibody (Jackson ImmunoResearch) at 1:10000 dilution in TBS-Tween-milk for 1 hour at RT. After 3 washes with TBS-Tween, signal detection was carried out using the SuperSignal West Pico chemio-luminescent

substrate (Thermo Scientific) as described by the manufacturer and imaged in the Vilmen-
Loubet Fusion machine.

RNA extraction, transcriptome annotation and sequence analysis

For RNA sequencing total RNA was extracted from cells of strain MCCV-054. 150 ml of
cells were grown in L1 media and collected 6, 12 and 18 days after initial inoculum as for
protein preparation.

Filters with cells were placed in 1ml of TRI-reagent (Sigma-Aldrich) and incubated on ice
for five minutes. Cells were broken by beating in the presence of ~ 3 g of glass beads
(Sigma) in a 220 V bead beater at 4000 rpm for 40s. RNA was then extracted using a
RiboPureTM (Ambion) kit following the manufacturer's instructions. Residual DNA was
removed by treatment with DNase I (Q1 DNAse, Promega) for 1h at 37 °C (2 units per
sample) followed by purification with the RNeasy miniElute Cleanup kit (Qiagen).

RNA purity, quantity and integrity were assessed for each sample using a Nanodrop ND-
1000 and 2100 Bioanalyser. RNA samples were then sent to BGI-Tech, China for library
generation and sequencing with the DNBSEQ™ technology platforms to generate 100 base
pair (bp) paired end reads.

De novo assembly of reads was carried out with Trinity version 2.4.0 (Grabherr et al., 2011)
using default parameters. The full assembly was filtered to the longest transcript per gene
group making use of the '_i' designations of the Trinity transcript identifiers. Open reading
frames were predicted using TransDecoder v5.3. These protein sequences were searched
against the NCBI NR database and sequences whose best hit was to Dinophyceae were
retained. To annotate these sequences, the pantherScore2.1.pl script was used to search
them against the PANTHER hidden Markov model library (version 14.1). Multiple
sequence alignments were performed using Clustal Omega and viewed using Jalview Vs
2-a (Waterhouse et al., 2009).

Acknowledgments

We are grateful to L. Besnardeau for technical support, to R. Lemée, E. Christians, S. Marro

and M. Hagström for discussion and sharing of reagents, to J. Chenevert and R. Copley for

critical reading of the manuscript. We thank the imaging platform, PIM (member of the

MICA microscopy platform) where image acquisition was conducted. PIM is supported by

EMBRC-France, whose French state funds are managed by the ANR within the

investments for the future program under reference ANR-10-INSB-02. DV was supported

by Minciencias- Colombia - financial support 783, PhD program.

Competing interests

540 No competing interests declared

References

Bhaud, Y., Guillebault, D., Lennon, J., Defacque, H., Soyer-Gobillard, M.O., Moreau,
H., 2000. Morphology and behaviour of dinoflagellate chromosomes during the
cell cycle and mitosis. J Cell Sci 113, 1231–1239.
Bodakuntla, S., Janke, C., Magiera, M.M., 2021. Tubulin polyglutamylation, a regulator
of microtubule functions, can cause neurodegeneration. Neuroscience Letters 746,
135656. https://doi.org/10.1016/j.neulet.2021.135656
Bravo, I., Vila, M., Casabianca, S., Rodriguez, F., Rial, P., Riobó, P., Penna, A., 2012.
Life cycle stages of the benthic palytoxin-producing dinoflagellate *Ostreopsis* cf.
ovata (Dinophyceae). Harmful Algae 18, 24–34.
https://doi.org/10.1016/j.hal.2012.04.001
Bulinski, J.C., Richards, J.E., Piperno, G., 1988. Posttranslational modifications of alpha
tubulin: detyrosination and acetylation differentiate populations of interphase
microtubules in cultured cells. Journal of Cell Biology 106, 1213–1220.
https://doi.org/10.1083/jcb.106.4.1213
Cachon, J., Cachon, M., 1977. Observations on the mitosis and on the chromosome
evolution during the lifecycle of *Oodinium*, a parasitic dinoflagellate.
Chromosoma 60, 237–251. https://doi.org/10.1007/BF00329773
Cowan, C.R., Hyman, A.A., 2004. Centrosomes direct cell polarity independently of
microtubule assembly in *C. elegans* embryos. Nature 431, 92–96.
https://doi.org/10.1038/nature02825
Cross, F.R., Umen, J.G., 2015. The Chlamydomonas cell cycle. Plant J 82, 370–392.
https://doi.org/10.1111/tpj.12795

Deng, M., Suraneni, P., Schultz, R.M., Li, R., 2007. The Ran GTPase Mediates Chromatin Signaling to Control Cortical Polarity during Polar Body Extrusion in Mouse Oocytes. Developmental Cell 12, 301–308. https://doi.org/10.1016/j.devcel.2006.11.008

Drechsler, H., McAinsh, A.D., 2012. Exotic mitotic mechanisms. Open Biol 2. https://doi.org/10.1098/rsob.120140

Driscoll, W.W., Hackett, J.D., Ferrière, R., 2016. Eco-evolutionary feedbacks between private and public goods: evidence from toxic algal blooms. Ecol Lett 19, 81–97. https://doi.org/10.1111/ele.12533

Eddé, B., Rossier, J., Le Caer, J.-P., Desbruyères, E., Gros, F., Denoulet, P., 1990. Posttranslational Glutamylation of α-tubulin. Science 247, 83–85. https://doi.org/10.1126/science.1967194

Ehler, L.L., Dutcher, S.K., 1998. Pharmacological and genetic evidence for a role of rootlet and phycoplast microtubules in the positioning and assembly of cleavage furrows in *Chlamydomonas reinhardtii.* Cell Motil Cytoskeleton 40, 193–207. https://doi.org/10.1002/(SICI)1097-0169(1998)40:2<193::AID-CM8>3.0.CO;2-G

Escalera, L., Benvenuto, G., Scalco, E., Zingone, A., Montresor, M., 2014. Ultrastructural Features of the Benthic Dinoflagellate *Ostreopsis* cf. *ovata* (Dinophyceae). Protist 165, 260–274. https://doi.org/10.1016/j.protis.2014.03.001

Eshun-Wilson, L., Zhang, R., Portran, D., Nachury, M.V., Toso, D.B., Löhr, T., Vendruscolo, M., Bonomi, M., Fraser, J.S., Nogales, E., 2019. Effects of α-tubulin acetylation on microtubule structure and stability. Proc Natl Acad Sci USA 116, 10366–10371. https://doi.org/10.1073/pnas.1900441116

Fenchel, T., 2001. How Dinoflagellates Swim. Protist 152, 329–338. https://doi.org/10.1078/1434-4610-00071

Francia, M.E., Dubremetz, J.-F., Morrissette, N.S., 2015. Basal body structure and composition in the apicomplexans Toxoplasma and Plasmodium. Cilia 5, 3. https://doi.org/10.1186/s13630-016-0025-5

Gavelis, G.S., Herranz, M., Wakeman, K.C., Ripken, C., Mitarai, S., Gile, G.H., Keeling, P.J., Leander, B.S., 2019. Dinoflagellate nucleus contains an extensive endomembrane network, the nuclear net. Scientific Reports 9. https://doi.org/10.1038/s41598-018-37065-w

Gayathri, P., Fujii, T., Møller-Jensen, J., Van Den Ent, F., Namba, K., Löwe, J., 2012. A Bipolar Spindle of Antiparallel ParM Filaments Drives Bacterial Plasmid Segregation. Science 338, 1334–1337. https://doi.org/10.1126/science.1229091

Geimer, S., Melkonian, M., 2005. Centrin Scaffold in *Chlamydomonas reinhardtii* Revealed by Immunoelectron Microscopy. Eukaryot Cell 4, 1253–1263. https://doi.org/10.1128/EC.4.7.1253-1263.2005

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol 29, 644–652. https://doi.org/10.1038/nbt.1883

Guillard, R.R.L., Hargraves, P.E., 1993. S*tichochrysis immobilis* is a diatom, not a
chrysophyte. Phycologia 32, 234–236. https://doi.org/10.2216/i0031-8884-32-3-
234.1
Gull, K., 1999. The Cytoskeleton of Trypanosomatid Parasites. Annu. Rev. Microbiol.
53, 629–655. https://doi.org/10.1146/annurev.micro.53.1.629
Hammarton, T.C., 2019. Who Needs a Contractile Actomyosin Ring? The Plethora of
Alternative Ways to Divide a Protozoan Parasite. Front. Cell. Infect. Microbiol. 9,
397. https://doi.org/10.3389/fcimb.2019.00397
Inoué, S., Ritter, H., 1978. Mitosis in *Barbulanympha.* II. Dynamics of a two-stage
anaphase, nuclear morphogenesis, and cytokinesis. The Journal of cell biology 77,
655–684. https://doi.org/10.1083/jcb.77.3.655
Janke, C., 2013. Mysterious modification of tubulin. Nat Rev Mol Cell Biol 14, 692–692.
https://doi.org/10.1038/nrm3685
Kilmartin, J.V., Wright, B., Milstein, C., 1982. Rat monoclonal antitubulin antibodies
derived by using a new nonsecreting rat cell line. The Journal of cell biology 93,
576–582. https://doi.org/10.1083/jcb.93.3.576
Laband, K., Le Borgne, R., Edwards, F., Stefanutti, M., Canman, J.C., Verbavatz, J.-M.,
Dumont, J., 2017. Chromosome segregation occurs by microtubule pushing in
oocytes. Nat Commun 8, 1499. https://doi.org/10.1038/s41467-017-01539-8
Lee, Y.J., Liu, B., 2019. Microtubule nucleation for the assembly of acentrosomal
microtubule arrays in plant cells. New Phytol 222, 1705–1718.
https://doi.org/10.1111/nph.15705
Moon, E., Nam, S.W., Shin, W., Park, M.G., Coats, D.W., 2015. Do All Dinoflagellates
have an Extranuclear Spindle? Protist 166, 569–584.
https://doi.org/10.1016/j.protis.2015.08.005
Ninčević Gladan, Ž., Arapov, J., Casabianca, S., Penna, A., Honsell, G., Brovedani, V.,
Pelin, M., Tartaglione, L., Sosa, S., Dell'Aversano, C., Tubaro, A., Žuljević, A.,
Grbec, B., Čavar, M., Bužančić, M., Bakrač, A., Skejić, S., 2019. Massive
Occurrence of the Harmful Benthic Dinoflagellate *Ostreopsis* cf. *ovata* in the
Eastern Adriatic Sea. Toxins 11, 300. https://doi.org/10.3390/toxins11050300
Nishihama, R., Machida, Y., 2001. Expansion of the phragmoplast during plant
cytokinesis: a MAPK pathway may MAP it out. Current Opinion in Plant Biology
4, 507–512. https://doi.org/10.1016/S1369-5266(00)00208-9
Oakley, B.R., Dodge, J.D., 1976. Mitosis and cytokinesis in the dinoflagellate
Amphidinium carterae. Cytobios 17, 35–46.
Oakley, B.R., Dodge, J.D., 1974. Kinetochores associated with the nuclear envelope in
the mitosis of a dinoflagellate. J Cell Biol 63, 322–325.
Onishi, M., Umen, J.G., Cross, F.R., Pringle, J.R., 2020. Cleavage-furrow formation
without F-actin in *Chlamydomonas*. Proc Natl Acad Sci USA 117, 18511–18520.
https://doi.org/10.1073/pnas.1920337117
Parsons, M.L., Aligizaki, K., Bottein, M.-Y.D., Fraga, S., Morton, S.L., Penna, A.,
Rhodes, L., 2012. *Gambierdiscus* and *Ostreopsis*: Reassessment of the state of
knowledge of their taxonomy, geography, ecophysiology, and toxicology.
Harmful Algae 14, 107–129. https://doi.org/10.1016/j.hal.2011.10.017

Pavaux, A.-S., Velasquez-Carjaval, D., Drouet, K., Lebrun, A., Hiroux, A., Marro, S.,
Christians, E., Castagnetti, S., Lemée, R., 2021. Daily variations of *Ostreopsis* cf.
ovata abundances in NW Mediterranean Sea. Harmful Algae 110, 102144.
https://doi.org/10.1016/j.hal.2021.102144
Perret, E., Davoust, J., Albert, M., Besseau, L., Soyer-Gobillard, M.O., 1993.
Microtubule organization during the cell cycle of the primitive eukaryote
dinoflagellate *Crypthecodinium cohnii*. J Cell Sci 104 (Pt 3), 639–651.
Pitayu-Nugroho, L., Aubry, M., Laband, K., Geoffroy, H., Ganeswaran, T., Primadhanty,
A., Canman, J.C., Dumont, J., 2023. Kinetochore component function in *C.*
elegans oocytes revealed by 4D tracking of holocentric chromosomes. Nat
Commun 14, 4032. https://doi.org/10.1038/s41467-023-39702-z
RadiBer, M., Plessman, U., KlOppelh, K.-D., 1992. Class II tubulin, the major brain,6
tubulin isotype is polyglutamylated on glutamic acid residue 435. FEBS
LETTERS.
Rhodes, L., 2011. World-wide occurrence of the toxic dinoflagellate genus *Ostreopsis*
Schmidt. Toxicon 57, 400–407. https://doi.org/10.1016/j.toxicon.2010.05.010
Richards, T.A., Cavalier-Smith, T., 2005. Myosin domain evolution and the primary
divergence of eukaryotes. Nature 436, 1113–1118.
https://doi.org/10.1038/nature03949
Ris, H., Kubai, D.F., 1974. An unusual mitotic mechanism in the parasitic protozoan
Syndinium sp. Journal of Cell Biology 60, 702–720.
https://doi.org/10.1083/jcb.60.3.702
Ritter, H., Inoué, S., Kubai, D., 1978. Mitosis in *Barbulanympha*. I. Spindle structure,
formation, and kinetochore engagement. The Journal of cell biology 77, 638–654.
https://doi.org/10.1083/jcb.77.3.638
Rosenbaum, J.L., Moulder, J.E., Ringo, D.L., 1969. Flagellar elongation and shortening
in Chlamydomonas. Journal of Cell Biology 41, 600–619.
https://doi.org/10.1083/jcb.41.2.600
Salisbury, J.L., Baron, A.T., Sanders, M.A., 1988. The centrin-based cytoskeleton of
Chlamydomonas reinhardtii: distribution in interphase and mitotic cells. Journal
of Cell Biology 107, 635–641. https://doi.org/10.1083/jcb.107.2.635
Schnepf, E., 1988. Cytochalasin D inhibits completion of cytokinesis and affects theca
formation in dinoflagellates. Protoplasma 143, 22–28.
https://doi.org/10.1007/BF01282955
Schnepf, E., Winter, S., Storck, I., Quader, H., 1990. A complementary experimental
study of cell division in the dinoflagellate *Prorocentrum micans*. European
Journal of Protistology 25, 234–242. https://doi.org/10.1016/S0932-
4739(11)80175-6
Schulze, E., Asai, D.J., Bulinski, J.C., Kirschner, M., 1987. Posttranslational
modification and microtubule stability. Journal of Cell Biology 105, 2167–2177.
https://doi.org/10.1083/jcb.105.5.2167
Sherwin, T., Gull, K., Vickerman, K., 1989. The cell division cycle of T*rypanosoma*
brucei: timing of event markers and cytoskeletal modulations. Philosophical
Transactions of the Royal Society of London. B, Biological Sciences 323, 573–
588. https://doi.org/10.1098/rstb.1989.0037

Sinclair, A.N., Huynh, C.T., Sladewski, T.E., Zuromski, J.L., Ruiz, A.E., de Graffenried,
699   C.L., 2021. The *Trypanosoma brucei* subpellicular microtubule array is organized
700   into functionally discrete subdomains defined by microtubule associated proteins.
701   PLoS Pathog 17, e1009588. https://doi.org/10.1371/journal.ppat.1009588

702   Stires, J.C., Latz, M.I., 2018. Contribution of the cytoskeleton to mechanosensitivity
703   reported by dinoflagellate bioluminescence. Cytoskeleton 75, 12–21.
704   https://doi.org/10.1002/cm.21392

705   Villanueva, M.A., Arzápalo-Castañeda, G., Castillo-Medina, R.E., 2014. The actin
706   cytoskeleton organization and disorganization properties of the photosynthetic
707   dinoflagellate *Symbiodinium kawagutii* in culture. Can. J. Microbiol. 60, 767–775.
708   https://doi.org/10.1139/cjm-2014-0325

709   Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., Barton, G.J., 2009. Jalview
710   Version 2—a multiple sequence alignment editor and analysis workbench.
711   Bioinformatics 25, 1189–1191. https://doi.org/10.1093/bioinformatics/btp033

712   Wetherbee, R., 1975. The fine structure of *Ceratium tripos*, a marine armored
713   dinoflagellate. Journal of Ultrastructure Research 50, 77–88.
714   https://doi.org/10.1016/S0022-5320(75)90010-6

715   Wilson, P.J., Forer, A., 1989. Acetylated a-tubulin in spermatogenic cells of the crane fly
716   *Nephrotoma suturalis*: Kinetochore microtubules are selectively acetylated. Cell
717   Motil. Cytoskeleton 14, 237–250. https://doi.org/10.1002/cm.970140210

718   Wolff, A., de Néchaud, B., Chillet, D., Mazarguil, H., Desbruyères, E., Audebert, S.,
719   Eddé, B., Gros, F., Denoulet, P., 1992. Distribution of glutamylated alpha and
720   beta-tubulin in mouse tissues using a specific monoclonal antibody, GT335. Eur J
721   Cell Biol 59, 425–432.

722   Wright, R.L., Adler, S.A., Spanier, J.G., Jarvik, J.W., 1989. Nucleus-basal body
723   connector in Chlamydomonas: Evidence for a role in basal body segregation and
724   against essential roles in mitosis or in determining cell polarity. Cell Motil.
725   Cytoskeleton 14, 516–526. https://doi.org/10.1002/cm.970140409

726   Wu, J., Akhmanova, A., 2017. Microtubule-Organizing Centers. Annu. Rev. Cell Dev.
727   Biol. 33, 51–75. https://doi.org/10.1146/annurev-cellbio-100616-060615

Figure 1. The cortical microtubule array is maintained during mitosis and cytokinesis.

(A) Representative confocal images of *O*. cf. *ovata* cells stained for ß-tubulin (D66 anti-ß-tubulin antibody). F= flagellum, AP= apical pore; VB= ventral bundle; CB= cortical bundle; C= cingulum. **(B)** Representative confocal images of control (dH2O) and colchicine (2µM) treated *O*. cf. *ovata* cells stained for ß-tubulin (D66 antibody). **(C)** Representative confocal images of pre-mitotic, mitotic and post-dividing *O*. cf. *ovata* cells stained for ß-tubulin (top, D66 antibody) and with Hoechst (DNA, bottom). Hoechst also stains DNA in mitochondria and chloroplasts. All cells are oriented with the ventral side on top, with the only exception of dH2O treated cells in B. All images are projections of the confocal Z-stack covering the whole cell. Scale bars are 10µm.

Figure 2: Changes in microtubule organization during dinomitosis.

Figure 2. Changes in microtubule organization during dinomitosis.

(A) Representative confocal images of *O.* cf. *ovata* cells at different cell cycle stages

stained with Hoechst (DNA, magenta) and for tubulin (green) using the D66 anti-ß-tubulin

antibody. White squares indicate the cell region enlarged underneath each image. All cells

are oriented with the ventral side on top. All images are projections of the confocal Z-stack

covering the nucleus. Schematic representations of microtubules and nucleus at each

analysed cell cycle stage are reported at the bottom. Arrow points to DNA; arrowhead to

forming cytokinetic structure. Number of analysed cells (n) is given for each stage. **(B)**

DNA-to-spindle pole distance as a function of total spindle length (pole to pole) in

pro/metaphase (red) and anaphase (blue) cells. Each dot corresponds to half a spindle. **(C)**

Quantification of DNA-to-spindle pole distance in cells at different mitotic stages. Each

dot represents half a spindle. Boxes represent 25-75th percentiles, and the mean is shown.

T-test was performed using all data points and retrieved non-significant difference (ns).

Scale bars are 20 µm.

Figure 3. A microtubule-rich plate divides daughter cells in cytokinesis.
(A) Confocal images (top) and corresponding 3-D reconstructions (anterior view, middle; lateral view: rotation of anterior view by 90 degrees to the right, bottom) of mitotic cells stained for tubulin using the D66 anti-ß-tubulin antibody. Arrows point to telophase polar spindle. (B) Representative confocal images of *O.* cf. *ovata* cells at different cell cycle stages or (C) of cells treated with the actin depolymerizing drug latranculin A, stained with Hoechst (DNA, top) and with phalloidin (actin, bottom).
All cells are oriented with the ventral side on top. Scale bars are 10 μm.

Figure 4: Tubulin post-translational modifications identify different microtubule populations

Figure 4. Tubulin post-translational modifications identify different microtubule populations

(**A**) Schematic representation of a ⬜- (black) and ⬜- (grey) tubulin dimer with some posttranslational modifications. (**B**) Western blot analysis of ß-tubulin and tubulin post-translational modifications in whole protein extract from *O.* cf. *ovata* cells. (**C**) Representative confocal images of *O.* cf. *ovata* cells stained with Hoechst (DNA, magenta), and for tubulin with various PTMs (green): tyrosinated tubulin (YL1/2 antibody, top), or acetylated tubulin (T6793 antibody, bottom). The number of analysed cells is indicated on each image. (**D**) Confocal image of *O.* cf. *ovata* pro/metaphase (left) and anaphase (right) spindles co-stained for tyrosinated tubulin (YL1/2 antibody, green), or acetylated tubulin (T6793 antibody, magenta). (**E**) Schematic representation of mitotic nuclei with metaphase and anaphase spindles: chromosomes are in light grey, kinetochore like structures in dark grey, nuclear membrane in black, tyrosinated tubulin in green and acetylated tubulin in magenta. Only 4 chromosomes are represented for simplicity. Scale bars are 20 µm, unless otherwise stated.

Figure 5: Centrin localizes to basal bodies throughout the mitotic cycle.

Figure 5. Centrin localizes to basal bodies throughout the mitotic cycle.

(**A**) Alignment of centrin sequences from *O.* cf. *ovata, C. reinhardtii*, and *Homo sapiens*. Coloured residues (blue) are identical across species. (**B**) Western blot analysis of centrin in whole protein extract from *O.* cf. *ovata* cells. (**C**) Representative confocal images of *O.* cf. *ovata* cells stained with Hoechst (DNA, cyan), for tubulin using the YL1/2 rat anti-□-tubulin antibody (green) and for centrin using the mouse 20HA antibody against centrin (magenta). White squares indicate the cell region enlarged underneath each image. All cells are oriented with the ventral side on top. All images are projections of the confocal Z-stack covering the nucleus. (**D**) Quantification of distance between peaks of fluorescence corresponding to centrin staining at different cell cycle stages. T-test was performed using all data points and retrieved either non-significant difference (ns) or a p value $< 0,001$ (*). (**E**) Schematic representation of changes in nuclear and basal body position and microtubule organization during the mitotic cell division cycle of *O.* cf. *ovata*. Scale bars are 20 µm.