Bursts of coalescence within population pedigrees whenever big families occur

Dimitrios Diamantidis[1], Wai-Tong (Louis) Fan[1,2], Matthias Birkner[3], and John Wakeley[2,*]

[1]*Department of Mathematics, Indiana University, Bloomington*
[2]*Department of Organismic and Evolutionary Biology, Harvard University*
[3]*Institut für Mathematik, Johannes-Gutenberg-Universität Mainz*
[*]*Corresponding author: wakeley@fas.harvard.edu*

October 16, 2023

Abstract

We consider a simple diploid population-genetic model with potentially high variability of offspring numbers among individuals. Specifically, against a backdrop of Wright-Fisher reproduction and no selection there is an additional probability that a pair of individuals has a number of offspring on the order of the population size. We study how the pedigree of the population generated under this model affects the ancestral genetic process of a sample of size two at a single autosomal locus without recombination. Our population model is of the type for which multiple-mergers coalescent processes have been described. We prove that the conditional distribution of the pairwise coalescence time given the random pedigree converges to a limit law as the population size tends to infinity. This limit law may or may not be the usual exponential distribution of the Kingman coalescent, depending on the frequency of high-reproduction events. But because it includes the number and times of high-reproduction events it differs from the usual multiple-merger coalescent models. The usual multiple-merger coalescent models are seen as describing the ancestral process marginal to, or averaging over, the pedigree. In the limiting ancestral process conditional on the pedigree, the intervals between high-reproduction events can be modeled using the Kingman coalescent but the events themselves cause discrete jumps in the probability of coalescence. Analogous results should hold for larger samples and other population models. We illustrate these results with simulations and additional analysis, highlighting their implications for understanding multi-locus data.

Introduction

Population geneticists routinely make inferences about the past by applying statistical models to DNA sequences or other genetic data. Because past events have already occurred, these models describe what might have happened. They are necessary because patterns of variation in DNA provide only indirect evidence about the past. But the decisions made in building these statistical models have important consequences for inference. Which parts of the past, if any, should be considered fixed objects or quantities and which should be treated as random variables? For the random variables, should their distributions be pseudo priors or explicitly modeled random processes? Our particular concern here will be with the treatment of pedigrees, or the reproductive relationships among diploid individuals.

With limited exceptions the statistical models of population genetics have inherited the initial decisions which Fisher (1922, 1930) and Wright (1931) made in deriving allele frequency spectra and probability density functions of allele frequencies at stationarity. They modeled neutral alleles as well as those under selection in a large well-mixed population which in the simplest case was assumed to be of constant size over time. Accordingly it has been common in population genetics to think of population sizes as fixed, not random. Today's coalescent hidden Markov models, for example, infer a fixed trajectory of population sizes over time under the assumption of neutrality (Li and Durbin, 2011; Sheehan et al., 2013; Wang et al., 2020; Schweiger and Durbin, 2023).

Although coalescent models reflect later developments and were a significant shift in thinking for the field, fundamentally they depend on the same assumptions as the classical models of Fisher and Wright (Ewens, 1990; Möhle, 1999). This is clear even in the earliest treatments of ancestral genetic processes by Malécot (1941, 1946, 1948). What coalescent theory did was to broaden the scope of population genetics beyond forward-time models of changes in allele counts or frequencies to include gene genealogies constructed by series of common-ancestor events backward in time (Kingman, 1982; Hudson, 1983a,b; Tajima, 1983). Mathematically, the forward-time and backward-time models of population genetics are dual to each other (Möhle, 1999).

Most importantly for our purposes here, Fisher (1922, 1930) and Wright (1931) obtained their predictions about genetic variation by averaging over an assumed random process of reproduction. The particular random process they used is now called the Wright-Fisher model (Ewens, 2004). Because the outcome of the process of reproduction is a pedigree, their method is equivalent to averaging over the random pedigree of the population. That they did this without explanation in this context is somewhat curious given the attention to pedigrees in Fisher's infinitesimal model of quantitative genetics (Fisher, 1918; Barton et al., 2017) and in Wright's method of path coefficients whose very purpose was to make predictions conditional on pedigrees (Wright, 1921a,b,c,d,e, 1922).

The pedigree of the entire population is the set of reproductive relationships of all individuals for all time when reproduction is bi-parental. The corresponding graph is a genealogy in the usual sense. It has been referred to as an organismal pedigree (Ball et al., 1990) and the population pedigree (Wollenberg and Avise, 1998; Wakeley et al., 2012; Ralph, 2019). Here we simply call it the pedigree. Patterns of genetic variation depend on the pedigree because genetic inheritance happens within it. In particular, transmission of an autosomal genetic locus forward in time through the pedigree occurs by Mendel's law of independent segregation. Multi-locus transmission follows Mendel's law of independent assortment or is mediated by recombination if the loci are linked. These processes, which may also be view backward in time, are conditional on the pedigree.

Within any pedigree, many possible uni-parental paths can be traced backward in time from each individual. If there are two mating types, for example karyotypic females (F) and karyotypic males (M), then one such path might be depicted $F \to F \to M \to F \to M \to \cdots$ (Avise and Wollenberg, 1997). For the ancestry of a single allele at an autosomal locus in a single individual, applying Mendel's law of independent segregation backward in time generates these uni-parental paths with equal probabilities $1/2^g$ for any path extending g generations into the past. When two such paths meet in the same individual, then with equal probability, $1/2$, the alleles either coalesce in that individual or remain distinct. Thus coalescence is conditional on the pedigree, and many possible gene genealogies are embedded in any one pedigree. Some loci, such as the mitochondrial genome and the Y chromosome in humans, are strictly uni-parentally inherited. They follow only paths $F \to F \to F \to \cdots$ and $M \to M \to M \to \cdots$, respectively, and two such paths coalesce with probability one when they meet. For these loci, there is only one gene genealogy within the pedigree.

Under Wright-Fisher reproduction, parents are chosen at random uniformly from among all

possible parents. This determines the structure of the pedigree in that generation. Assume that there are N_f karyotypic females and N_m karyotypic males in every generation. For autosomal loci, the familiar effective population size $N_e = 4N_\text{f}N_\text{m}/(N_\text{f} + N_\text{m})$ from classical forward-time analysis (Wright, 1931) and its backward-time counterpart $1/(2N_e)$ for the pairwise coalescence probability (Möhle, 1998a,b) come from averaging over the possible outcomes of reproduction in a single generation. Sections 6.1 and 6.2 in Wakeley (2009) give a detailed illustration. For uni-parentally inherited loci, this averaging yields $1/N_\text{f}$ and $1/N_\text{m}$ for the pairwise coalescence probabilities. In the diploid monoecious Wright-Fisher model or by setting $N_\text{f} = N_\text{m} = N/2$, these average probabilities of coalescence become $2/N$ for uni-parentally inherited loci and $1/(2N)$ for autosomal loci. For simplicity in this work we will focus on the diploid monoecious Wright-Fisher model.

Averaging over pedigrees is what leads to the effective population size, N_e, being the primary determinant of forward-time and backward-time dynamics in neutral population genetic models. For very large populations, N_e becomes the only parameter of the Wright-Fisher diffusion (Ewens, 2004) and the standard neutral or Kingman coalescent process (Sjödin et al., 2005). In particular, N_e sets the timescale over which mutation acts to produce genetic variation. Such averaging removes the pedigree as a possible latent variable which could be important for structuring genetic variation. As a result, from the perspective of the standard neutral coalescent, information about the (marginal) gene genealogical process together with the mutation process is all we can hope to infer from genetic data (Sjödin et al., 2005).

The situation in which it makes the most sense to use this marginal process of coalescence is when the only data available come from a single non-recombining locus. In fact, the initial applications of ancestral inference to single-locus data, namely to restriction fragment length polymorphisms in human mitochondrial DNA (mtDNA) (Brown, 1980; Cann et al., 1987) then to sequences of the hyper-variable control region (Vigilant et al., 1989, 1991; Ward et al., 1991; Di Rienzo and Wilson, 1991), did not even use of the statistical machinery of population genetics. They instead took the gene genealogy and times to common ancestry to be fixed, and estimated them using traditional phylogenetic methods (Felsenstein, 2004). But this in turn spurred the development of likelihood-based methods of ancestral inference using coalescent prior distributions for gene genealogies (Lundstrom et al., 1992; Griffiths and Tavaré, 1994; Kuhner et al., 1995). We note that in the interim it has also become common to treat phylogenies as random variables using a wide variety of prior models (Ronquist et al., 2012; Suchard et al., 2018; Bouckaert et al., 2019).

The desirability of accounting for variation in gene genealogies became especially clear when the first sample DNA sequences of the human *ZFY* gene was obtained and was completely monomorphic (Dorit et al., 1995). The mutation rate is lower on the Y chromosome than in the hyper-variable region of mtDNA but it is not equal to zero (Brown et al., 1979; Wilson et al., 1985; Ingman et al., 2000; The 1000 Genomes Project Consortium, 2015). Using coalescent priors it was shown that the complete lack of variation in that first sample at *ZFY* was consistent with a wide range of times to common ancestry for the Y chromosome (Dorit et al., 1995; Donnelly et al., 1996; Fu and Li, 1996; Weiss and von Haeseler, 1996).

If instead data come from multiple loci, it is impossible to ignore variation in gene genealogies regardless of whether one thinks of the pedigree as fixed or random. Variation in gene genealogies across the genome is, for example, what coalescent hidden Markov models use to estimate trajectories of population sizes. The simplest illustrative case is when the loci are on different chromosomes or far enough apart on the same chromosome that they assort independently into gametes, and when within each locus there is no recombination. The gene genealogies of such loci will vary due to the particular outcomes of Mendelian segregation. They will also be independent due to

Mendelian assortment, but only given the pedigree. Mendel's law of independent assortment is a law of conditional independence. It applies once relationships have been specified.

However, throughout much of the history of population genetics, it was assumed that independently assorting loci would have completely independent evolutionary histories. In coalescent theory, this means independent gene genealogies. As Charlesworth (2022) recently noted, Fisher (1922, 1930) and Wright (1931) intended their results on allele frequency spectra and probability density functions of allele frequencies at stationarity to be descriptions of the behavior of large numbers of independently assorting loci in the same genome. This is evident in their application of these distributions to the multiple Mendelian factors of Fisher's infinitesimal model (Fisher, 1918) in their arguments about the Dominance Ratio (Fisher, 1922; Charlesworth, 2022).

An early application to multi-locus data was made by Cavalli-Sforza and Edwards (1967) and Felsenstein (1973) who developed likelihood-based methods to infer trees of populations within species from multi-locus allele-frequency data, specifically human blood group data, by modeling the forward-time process of random genetic drift independently at each locus conditional on the population tree. Felsenstein (1981) further developed and applied these methods to gel electrophoretic data. Today's methods of inferring admixture from single nucleotide polymorphism, or SNP, data using F-statistics are based on the same notion of independence (Patterson et al., 2012).

Like the population size itself, demographic features such as the splitting of populations have mostly been treated as fixed in population genetics. Cavalli-Sforza and Edwards (1967) and Felsenstein (1973) did discuss but did not implement prior models for trees of populations, specifically as outcomes of birth-death processes. More recently, Heled and Drummond (2009) did implement this in a coalescent framework for multi-locus sequence data, using the prior distribution of Gernhard (2008); see also Lambert and Stadler (2013). Yang (2002) and Rannala and Yang (2003) took a different approach, using gamma-distributed pseudo priors for times in trees.

Although the underlying assumption that unlinked loci have completely independent evolutionary histories is mistaken because it would require them having independent pedigrees, most theoretical work has followed the lead of Fisher (1922, 1930) and Wright (1931). Examples in which this is made explicit include Karlin and McGregor (1967), Kimura (1969), Ewens (1974), and Ewens and Maruyama (1975). Multiplying likelihoods across loci in applications to genetic data subsequently became common practice (Watterson, 1985; Padmadisastra, 1988; Sawyer and Hartl, 1992; Wakeley, 1999; Nielsen, 2000; Wooding and Rogers, 2002; Adams and Hudson, 2004). It is built into current inference packages, including $\partial a \partial i$ (Gutenkunst et al., 2009), $momi2$ (Kamm et al., 2020) and $fastsimcoal2$ (Excoffier et al., 2013, 2021).

As it happens, this conceptual mistake has almost no practical ramifications if the population is large and well mixed, and the variance of offspring numbers among individuals is not too large. Ball et al. (1990) were the first to address the question of gene genealogies within pedigrees. They used simulations to show that the distribution of gene genealogies and coalescence times among loci on a single pedigree do not differ substantially from their distributions among loci which have independent pedigrees. Their population model was similar to the Wright-Fisher model with population size $N = 100$: Poisson offspring numbers with strong density regulation to a carrying capacity of 100. Their results were based on simulations of 50 gene genealogies for each of 50 pedigrees and samples of size $n = 100$, in which a single gene copy was taken at random at each locus within each individual.

Wakeley et al. (2012) confirmed these results and related them to coalescent theory using simulations of 10^8 gene genealogies for $n = 2$ for each of 10^4 pedigrees and population sizes up to 10^5, together with more limited treatments of larger samples $n = 20$ and $n = 100$. Pedigrees

were constructed in three different ways: assuming Wright-Fisher reproduction, using empirically derived human family structures, and under a model in which the outcome of a single generation of Wright-Fisher reproduction was repeated over time, resulting in a so-called cyclical pedigree. These simulations showed that times to common ancestry conditional on the pedigree conform well to the probability law underlying coalescent theory, with a constant coalescence probability $1/(2N_e) = 1/(2N)$ each generation under the Wright-Fisher model with $N_{\mathrm{f}} = N_{\mathrm{m}} = N/2$, except for in the recent past where they differ greatly and depend on the pedigree.

Here "recent" means proportional to $\log_2(N)$ generations, which is the timescale for the first occurrence of a common ancestor of all present-day individuals (Chang, 1999) and for the complete overlap of all individuals' families in a well-mixed bi-parental population (Chang, 1999; Derrida et al., 1999, 2000a,b; Barton and Etheridge, 2011; Coron and Le Jan, 2022). This is much shorter than the N-generations timescale required for common ancestry of uni-parental genetic lineages (Chang, 1999; Donnelly et al., 1999). Further work on these properties of pedigrees include Rohde et al. (2004) and Lachance (2009) who showed that population structure and inbreeding do not strongly affect the time to the first occurrence of a common ancestor of all individuals. Blath et al. (2014) proved that the ancestries of the great majority of individuals overlap even in cyclical pedigrees as $N \to \infty$. Matsen and Evans (2008) and Gravel and Steel (2015) showed that ancestral genetic lineages pass through only a small minority of the shared pedigree ancestors. Sainudiin et al. (2016) constructed a model with recombination which interpolates between uni-parental common ancestry on the N-generations timescale and bi-parental common ancestry on the $\log_2(N)$-generations timescale.

Tyukin (2015) proved what was implied by the simulations of Ball et al. (1990) and Wakeley et al. (2012), that when the population is large and well mixed the pedigree-averaged coalescent process is a good substitute for the actual coalescent process conditional on the pedigree. Questions of this sort have a long history in mathematical physics and probability theory, where "quenched" and "annealed" are often used to refer to conditional as opposed to averaged processes. Molchanov (1994) and Bolthausen and Sznitman (2002b) provide background and developments in the classical context of random walks in random environments. What Tyukin (2015) proved is that the quenched coalescent process conditional on the pedigree converges to the pedigree-averaged standard neutral or Kingman coalescent process in the limit $N \to \infty$. Tyukin (2015) did this under a broader set of reproduction models with mating analogous to Wright-Fisher but with a general exchangeable distribution of offspring numbers (Cannings, 1974), assuming as Kingman (1982) did that the variance of this distribution is finite as $N \to \infty$.

Since time in the Kingman coalescent process is measured in units proportional to N generations, the result of Tyukin (2015) provides insight into the role of the pedigree in the recent ancestry of the sample ($\propto \log_2(N)$ generations) under the Cannings and Wright-Fisher models. Specifically, the chance of any events in the recent past which would dramatically alter the rate of coalescence must be negligible as $N \to \infty$. Intuitively we might surmise that (1) individuals randomly sampled from a large well mixed population are unlikely to be closely related, and barring coalescence for some small number of generations until their ancestries overlap does not affect the limit, and (2) by the time their ancestries do overlap in the pedigree, their numbers of ancestors are approaching the population size, making the chance of coalescence of order $1/N$.

Two cases have been identified using simulations where quenched and annealed results are noticeably different. The first is population subdivision, especially with limited migration. Wollenberg and Avise (1998) showed that as the migration distance decreases in a linear habitat, fewer independent loci are needed to accurately measure pairwise coefficients of coancestry on the pedigree.

Wilton et al. (2017) described increasingly strong pedigree effects as the migration rate decreased in a two-subpopulation model, specifically spikes in the distribution of pairwise coalescence times corresponding to the particular series of individual migration events that occurred in the ancestry. These results illustrate how even single gene genealogies may contain information about events in the ancestry of geographically structured populations, via the pedigree. Thus they are relevant for applications of ancestral inference to single-locus data, such as mtDNA, as well as to the broader field of intraspecific phylogeography (Avise et al., 1987; Avise, 1989, 2000). For recent empirical studies of spatiotemporally structured pedigrees and their effects on local patterns of genetic variation, see Aguillon et al. (2017) and Anderson-Trocmé et al. (2023).

The second situation in which pedigrees have a strong effect on coalescence times and gene genealogies is when there is a high variance of offspring numbers among individuals. This variance is comparatively low in the Wright-Fisher model, which has a multinomial distribution of offspring numbers (becoming Poisson as $N \to \infty$). In deriving the standard neutral coalescent process, Kingman (1982) started with the general exchangeable model of Cannings (1974) then assumed that the variance of offspring numbers was finite as $N \to \infty$. Without this assumption, the ancestral limit process is not the Kingman coalescent process but rather a coalescent process with multiple mergers (see below). In addition in this situation simulations have shown that the pedigree has a marked effect on genetic ancestries.

Wakeley et al. (2016) simulated pedigrees in which a single individual had a very large number of offspring in some past generation and otherwise there was Wright-Fisher reproduction. This large reproduction event greatly increased the probability of coalescence in the generation in which it occurred, causing a spike in the distribution of pairwise coalescence times and altering the allele frequency spectrum. A strong selective sweep at one locus gave similar effects at unlinked loci via the pedigree (Wakeley et al., 2016). Similar deviations from standard neutral coalescent predictions are produced by cultural transmission of reproductive success (Guez et al., 2023).

Here, we present a new quenched limit result for coalescent processes in fixed pedigrees under a modified Wright-Fisher model which allows for large reproduction events. Wright-Fisher reproduction on its own produces various kinds of large reproduction events but these are all extremely rare. Our model adds big families with two parents and numbers of offspring proportional to the population size. These are inserted into the pedigree either on the same N-generations timescale as coalescent events in the Wright-Fisher background model or much faster so that they completely dominate the ancestral process. In both cases, the limiting ancestral process conditional on the pedigree is different than the limiting ancestral process which averages over pedigrees. For simplicity, we focus on samples of size two. Consistent with the results of Tyukin (2015), our result reduces to the Kingman coalescent with $n = 2$ in the case where there are no big families.

Note that the corresponding averaged process is not the Kingman coalescent but rather a coalescent process with multiple mergers; see Tellier and Lemaire (2014) for an overview of these models in the context of population genetics. Multiple-mergers coalescent processes arise as $N \to \infty$ limits when the variance of offspring numbers is large, and so may be applicable to a broad range of species with the capacity for high fecundity (Eldon, 2020). They also arise from recurrent selective sweeps, when differences in offspring numbers are determined by individuals' genotypes (Durrett and Schweinsberg, 2004, 2005; Schweinsberg and Durrett, 2005). Whereas the Kingman coalescent includes only binary mergers of ancestral genetic lineages, these more general processes allow mergers of any size. At issue here is how these models should be interpreted and applied.

By averaging over the process of reproduction, two kinds of multiple-mergers coalescent processes have been described: Λ-coalescents which have asynchronous multiple-mergers (Donnelly and

Kurtz, 1999; Pitman, 1999; Sagitov, 1999) and Ξ-coalescents which have simultaneous multiple-mergers (Schweinsberg, 2000; Möhle and Sagitov, 2001; Sagitov, 2003). Multiple-mergers processes for diploid organisms are always Ξ-coalescents with even numbers of simultaneous mergers (Birkner et al., 2018). Our quenched limit result brings into question what seems like a natural extension from applications of the standard neutral coalescent model, namely to apply multiple-mergers models independently to independent loci as has been done both in theoretical explorations (Der and Plotkin, 2014; Eldon et al., 2015; Spence et al., 2016; Matuszewski et al., 2018) and in analyses of SNP data (Birkner et al., 2013a; Blath et al., 2016; Árnason et al., 2023; Freund et al., 2023).

To establish the quenched limit process, we adapt the method that Birkner et al. (2013c) used for a quenched limit of a random walk in a random environment. See also the earlier work of Bolthausen and Sznitman (2002a). In this approach, the problem of convergence in distribution is addressed by analyzing a pair of conditionally independent processes, here corresponding to the ancestries of samples at two independently assorting loci on the pedigree. As Koskela (2018) has pointed out, positive correlations of coalescence times for pairs of unlinked loci are a hallmark of (pedigree-averaged) multiple-mergers coalescent models. Our result frames this in terms of pedigrees, in which big families are the only elements that persist as $N \to \infty$. If a big family has occurred in a particular generation, the probability of coalescence is greatly increased in that generation for all loci. All other aspects of the pedigree, that is to say the outcomes of ordinary Wright-Fisher reproduction, "average out" such that the Kingman coalescent process describes the ancestral process during the times between big families.

Theory and results

In this section, we present the population model considered in this paper, the mathematical statement of our main result and its proof. This result (Theorem 1) is stated as a convergence of the *conditional* distribution of the coalescence time of a pair of gene copies, given that we know the pedigree and which individuals were sampled. We assume that the pedigree is the outcome of a random process of reproduction, the population model described in the following section, and that the two individuals are sampled without replacement from the current generation. To connect with known results and highlight the effect of conditioning, we first state and prove the corresponding result (Lemma 1) for the *un*conditional distribution of the coalescence time. This corresponds to fixing the sampled individuals and averaging over the pedigree. We close this section with simulations illustrating multi-locus genetic ancestry and further analysis showing how non-zero correlations of coalescence times at unlinked loci result from averaging over the pedigree.

The population model

We consider a diploid, monoecious, bi-parental, panmictic population of constant fixed size $N \in \mathbb{N}$ with discrete, non-overlapping generations. Implicitly there is no selection, but we do not in fact model mutation or genetic variation, only the generation of the pedigree and coalescence within it. There are two different types of reproduction. With high probability, reproduction follows the diploid bi-parental Wright Fisher model. With small probability α_N each generation, there is a highly reproductive pair whose offspring comprise a proportion $\psi \in [0, 1]$ of the population. Note that ψ is a fixed deterministic constant. More precisely, for each positive integer g, the reproductive dynamics between the parent generation $g + 1$ and the children generation g is given as follows:

1. (Wright Fisher reproduction) With probability $1 - \alpha_N$, each individual in the next generation
is formed by choosing two parents at random, uniformly with replacement from the adults of
the current generation. Note this allows the possibility of (random) selfing. Genetically, each
offspring is produced according to Mendel's laws which means each of the two gene copies
in a parent is equally likely to be the one transmitted to the offspring. In this case we call
g a "*Wright Fisher (WF) generation*". An example of this standard reproduction dynamics
between the parent generation $g + 1$ and the offspring generation g is depicted below for a
population of size $N = 7$.

2. (High reproduction) With probability α_N, a highly reproductive pair, chosen uniformly among
all possible pairs, appears whose offspring replace a fraction $\psi \in [0, 1]$ of the population. In
this case we refer to g as a "*high-reproduction (HR) generation*". The chosen pair has $[\psi N]$
children, and we call these "big families". The other $N - [\psi N]$ offspring are produced as
above according to the Wright-Fisher model. An example is depicted below for $N = 7$ and
$\psi = 0.72$ in which the highly reproductive pair $(I_1, I_2) = (4, 5)$ in generation $g + 1$ has
$[\psi N] = [0.72 \cdot 7] = 5$ children in generation g.

These two possibilities happen independently for all generations $g \in \mathbb{Z}_{\geq 0}$. The classical Wright-
Fisher model corresponds to the case when $\alpha_N = 0$. In this case every $g \in \mathbb{Z}_{\geq 0}$ is a WF generation.

The *parent assignment* between two generations $g+1$ (parents) and g (children) is the collection
of edges connecting the children with their parents. The diagram in (1) below shows a parent
assignment corresponding to the Wright-Fisher reproduction in the example above.

$$(1)$$

332 On the other hand, the diagram in (2) below shows a parent assignment between generations g and
333 $g + 1$ corresponding to the high-reproduction generation in the example above.

$$(2)$$

335 **Pedigree**   The collection of all the parent assignments among all pairs of consecutive generations is
336 called the *pedigree* and it is denoted as $\mathcal{A}^{(N)}$. The pedigree models the set of all family relationships
337 among the members of the population for all generations. The pedigree is shared among all loci.
338 It is the structure through which genetic lineages are transmitted. Patterns of ancestry, or gene
339 genealogies are outcomes of Mendelian inheritance in this single shared pedigree.

Frequency of HR events Recall that α_N denotes the probability of a big family to appear in a generation. We set

$$\alpha_N = \frac{\lambda}{N^\theta}, \tag{3}$$

where $\theta \in (0, 1]$ and $\lambda \in \mathbb{R}_{\geq 0}$ is a fixed parameter which determines the relative frequency of the high-reproduction events.

Timescale Suppose two individuals are sampled uniformly *without* replacement among the N individuals of the current generation $g = 0$ and we sample one gene copy from each. Let $\tau^{(N,2)}$ be the pairwise coalescence time, that is, the number of generations in the past until the two sampled gene copies coalesce. How long is the pairwise coalescence time $\tau^{(N,2)}$? This will depend on N and also on θ owing to our assumption (3). In considering the limiting ancestral process for the sample, we re-scale time so that it is measured in units of N^θ generations. We study the distribution of the re-scaled pairwise coalescence time, $\tau^{(N,2)}/N^\theta$, with different results depending on whether $\theta \in (0, 1)$ or $\theta = 1$. In the latter case, our timescale is N generations, which we note is $1/2$ the usual coalescent timescale for diploids. In the former case, where we may infer from (3) that high-reproduction events will dominate the ancestral process, the timescale is accordingly much shorter than the usual coalescent timescale. Coalescence times in both cases also depend on a combined parameter $\psi^2/4$ which is the limiting probability of coalescence when a big family occurs.

Limiting process by averaging over the pedigree

For reference and to illustrate our choice of timescale, we begin with a Kingman coalescent approximation for the pairwise coalescence time in the classical Wright-Fisher model, here the special case $\theta = 1$ and $\lambda = 0$ or $\alpha_N = 0$. Averaging over the process of reproduction in a single generation gives a coalescence probability of $1/(2N)$. With $\theta = 1$, we measure time in units of N generations. To parallel the derivation of our main result, we consider the probability that the coalescence time $\tau^{(N,2)}$ is more than $[tN]$ generations. The limiting ancestral process is obtained as

$$\mathbb{P}^{(N)}(\tau^{(N,2)} > [tN]) = \left(1 - \frac{1}{2N}\right)^{[tN]} \to e^{-t/2} \qquad \text{as } N \to \infty. \tag{4}$$

In words, the re-scaled coalescence time $\frac{\tau^{(N,2)}}{N}$ converges in distribution to an exponential random variable with rate parameter $1/2$.

Before stating our main result, we first prove Lemma 1 below, generalizing (4) to our population model, in the sense that for $\theta = 1$ with $\psi = 0$ or $\lambda = 0$ in Lemma 1 we recover (4).

Lemma 1. *Let $\lambda \in \mathbb{R}_{\geq 0}$, $\theta \in (0, 1]$, and set $\alpha_N = \frac{\lambda}{N^\theta}$. The re-scaled coalescence time $\frac{\tau^{(N,2)}}{N^\theta}$ converges in distribution to an exponential random variable with rate parameter*

$$\begin{cases} \lambda \frac{\psi^2}{4}, & when \ \theta \in (0, 1) \\ \\ \frac{1}{2} + \lambda \frac{\psi^2}{4}, & when \ \theta = 1. \end{cases} \tag{5}$$

We note that in Birkner et al. (2013b), the full ancestral recombination graph for samples of arbitrary size and genomes consisting of arbitrary numbers of linked loci is described for a population model nearly identical to ours here. The ancestral recombination graph (Hudson, 1983a; Griffiths and Marjoram, 1997), like the Kingman coalescent itself, averages over the pedigree. Lemma 1 describes the marginal ancestral process for a sample of size two at a single locus.

Proof of Lemma 1 The lineage dynamics of our model can be analyzed using a Markov chain. In any generation g in the past, the ancestral lineages of a pair of gene copies must be in one of the three states $\{\xi_0, \xi_1, \xi_2\}$, where

- $\xi_0 = (\bullet)(\bullet)$ represents two ancestral lineages in two distinct individuals,

- $\xi_1 = (\bullet\bullet)$ represents two ancestral lineages on different chromosomes in the same individual,

- $\xi_2 = (\bullet)$ represents that the ancestral lineages have coalesced.

The diploid ancestral process for a pair of gene copies can thus be represented as a Markov chain $(M_g)_{g \in \mathbb{Z}_{\geq 0}}$ with state space $\{\xi_0, \xi_1, \xi_2\}$, where M_g is the state of the two lineages g generations in the past. Its one-step transition matrix Π_N is given by

$$\Pi_N := (1 - \alpha_N)\Pi_N^{\mathrm{WF}} + \alpha_N \Pi_N^{\mathrm{HR}} \tag{6}$$

where

$$\Pi_N^{\mathrm{WF}} = \begin{array}{c} \\ \xi_0 \\ \xi_1 \\ \xi_2 \end{array} \begin{array}{ccc} \xi_0 & \xi_1 & \xi_2 \\ \left[\begin{array}{ccc} 1 - \frac{1}{N} & \frac{1}{2N} & \frac{1}{2N} \\ 1 - \frac{1}{N} & \frac{1}{2N} & \frac{1}{2N} \\ 0 & 0 & 1 \end{array}\right] \end{array} \tag{7}$$

and

$$\Pi_N^{\mathrm{HR}} = \begin{array}{c} \\ \xi_0 \\ \xi_1 \\ \xi_2 \end{array} \begin{array}{ccc} \xi_0 & \xi_1 & \xi_2 \\ \left[\begin{array}{ccc} 1 - \frac{\psi^2}{2} & \frac{\psi^2}{4} & \frac{\psi^2}{4} \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{array}\right] \end{array} + O\left(\frac{1}{N}\right). \tag{8}$$

The matrix Π_N^{WF} in (7) is the transition matrix for a Wright Fisher generation, whereas Π_N^{HR} in (8) is for a high-reproduction generation. The entries of Π_N^{HR} in (8) are derived by conditioning on the parent assignment(s) for the individual(s) containing the ancestral lineages, with respect to the highly reproductive pair. For instance, for ancestral lineages currently in two distinct individuals, the coalescence probability is $1/4$ if both individuals descend from the HR couple and $1/(2N)$ otherwise. Thus we have

$$\mathbb{P}(M_{g+1} = \xi_2 | M_g = \xi_0, \mathrm{HR}) = \frac{[\psi N]([\psi N] - 1)}{N(N-1)} \frac{1}{4} + \left(1 - \frac{[\psi N]([\psi N] - 1)}{N(N-1)}\right) \frac{1}{2N}$$

$$= \frac{\psi^2}{4} + O\left(\frac{1}{N}\right)$$

for the transition $\xi_0 \to \xi_2$ in Π_N^{HR}, and where we have also specified that this contribution to the overall probability in (6) is conditional on the HR couple.

The rest of the proof is a straightforward application of Möhle (1998a, Lemma 1). This is a separation-of-timescales result. To see how it works, using (3) we can rewrite (6) as

$$\Pi_N := A + \frac{1}{N^\theta} B_N + O\left(\frac{1}{N^{\theta+1}}\right) \tag{9}$$

where

$$A = \begin{array}{cc} & \begin{array}{ccc} \xi_0 & \xi_1 & \xi_2 \end{array} \\ \begin{array}{c} \xi_0 \\ \xi_1 \\ \xi_2 \end{array} & \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{array} \tag{10}$$

and

$$B_N = \begin{array}{cc} & \begin{array}{ccc} \xi_0 & \xi_1 & \xi_2 \end{array} \\ \begin{array}{c} \xi_0 \\ \xi_1 \\ \xi_2 \end{array} & \begin{bmatrix} -\frac{N^\theta}{N} - \frac{\psi^2}{2} & \frac{N^\theta}{N}\frac{1}{2} + \lambda\frac{\psi^2}{4} & \frac{N^\theta}{N}\frac{1}{2} + \lambda\frac{\psi^2}{4} \\ -\frac{N^\theta}{N} & \frac{N^\theta}{N}\frac{1}{2} & \frac{N^\theta}{N}\frac{1}{2} \\ 0 & 0 & 1 \end{bmatrix} \end{array}. \tag{11}$$

The matrix A contains the fastest parts of the process. The matrix B_N contains the next-fastest parts of the process, specifically those occurring on the timescale of N^θ generations.

Möhle's result depends on the existence of equilibrium stochastic matrix $P := \lim_{k\to\infty} A^k$ which in this case is equal to A. Möhle's result also requires the existence of the limiting infinitesimal generator $G := \lim_{N\to\infty} PB_N P$. Note that in our application $B := \lim_{N\to\infty} B_N$ itself converges. From (11) it is clear that the limiting result will differ depending on whether $\theta \in (0,1)$ or $\theta = 1$. If $\theta \in (0,1)$, the Wright-Fisher contribution to the coalescence rate shrinks to zero in the limit. If $\theta = 1$, the Wright-Fisher contribution, which is $1/2$ on our timescale, remains comparable to the HR contribution in the limit.

Applying Möhle (1998a, Lemma 1) to compute our probability of interest,

$$\mathbb{P}^{(N)}(\tau^{(N,2)} > [tN^\theta]) = (1,0,0)\,\Pi_N^{[tN^\theta]}\,(1,1,0)^T \tag{12}$$

$$= (1,0,0)\left(A + \frac{1}{N^\theta} B_N + O\left(\frac{1}{N^{\theta+1}}\right)\right)^{[tN^\theta]}(1,1,0)^T \tag{13}$$

$$\to (1,0,0)\,P e^{tG}\,(1,1,0)^T \qquad \text{as } N \to \infty. \tag{14}$$

The initial vector $(1,0,0)$ enforces our assumed starting state, ξ_0. The end vector $(1,1,0)^T$ enforces the requirement that the lineages remain distinct at generation $[tN^\theta]$, i.e. that the Markov chain $(M_g)_{g\in\mathbb{Z}_{\geq 0}}$ has not reached state ξ_2. Möhle's result $P e^{tG}$ is in the middle. Recall that P, which here is equal to A, instantaneously adjusts the sample so that the effective starting state is ξ_0 even if the sample state is ξ_1. The lineages then enter the continuous-time process with rate matrix G. Overall we have

$$P\,e^{tG}\,(1,1,0)^T = \begin{cases} \left(e^{-t\lambda\frac{\psi^2}{4}}, e^{-t\lambda\frac{\psi^2}{4}}, 0 \right)^T, & \text{if } \theta \in (0,1), \\[2ex] \left(e^{-t\left(\frac{1}{2}+\lambda\frac{\psi^2}{4}\right)}, e^{-t\left(\frac{1}{2}+\lambda\frac{\psi^2}{4}\right)}, 0 \right)^T, & \text{if } \theta = 1. \end{cases} \tag{15}$$

The right hand side of (14) is equal to (5), and the proof of Lemma 1 is complete.   □

Remark 1 (Robustness against perturbation of initial condition)**.** The form of P shows that the
limiting result in Lemma 1 holds regardless of whether the sample begins in state ξ_0, as we have
assumed, or in state ξ_1. So, other sampling schemes could be considered. In fact Lemma 1 still
holds if the initial distribution lies in the set $\mathcal{I} := \{(c, 1 - c, 0) \in [0, 1]^3 : c \in [0, 1]\}$. This can be
seen clearly in (15).

Limiting process by conditioning on the pedigree

437   Our main result is about the conditional distribution. We let

$$F_N(t, \mathcal{A}^{(N,2)}) := \mathbb{P}^{(N)} \left(\tau^{(N,2)} > [tN^\theta] \mid \mathcal{A}^{(N,2)} \right) \tag{16}$$

be the conditional probability of the event $\{\tau^{(N,2)} > [tN^\theta]\}$ given the (random) pedigree and the
sampled pair of individuals. Mathematically, $\mathcal{A}^{(N,2)}$ is the sigma-field (all information) generated
by the outcome of the random reproduction of the population and the knowledge which pair of
individuals was sampled.

Theorem 1. *Let $\lambda \in \mathbb{R}_{\geq 0}$, $\theta \in (0, 1]$, and set $\alpha_N = \frac{\lambda}{N^\theta}$. For all $t \in (0, \infty)$, we have the following*
convergence in distribution as $N \to \infty$

$$F_N(t, \mathcal{A}^{(N,2)}) \to \begin{cases} \left(1 - \frac{\psi^2}{4}\right)^{Y(t)}, & \text{when } \theta \in (0, 1), \\[2ex] e^{-t/2} \left(1 - \frac{\psi^2}{4}\right)^{Y(t)}, & \text{when } \theta = 1, \end{cases} \tag{17}$$

where $Y(t)$ is Poisson process with rate λ. In fact, the convergence in (17) holds jointly for all
$t > 0$, see the discussion in Remark 4 in the Appendix section A.4 for details.

Theorem 1 offers a description of the conditional distribution of the coalescence time $\tau^{(N,2)}$ for
a sample of two genes in a population of size N given the pedigree. It says that the law of $\frac{\tau^{(N,2)}}{N^\theta}$,
under the conditional probability $\mathbb{P}(\cdot \mid \mathcal{A}^{(N,2)})$, converges weakly as $N \to \infty$ to the law of a random
variable (call it T) under a probability measure $\mathbb{P}_Y$ that depends on the Poisson process Y with
rate λ. Furthermore, the survival function $\mathbb{P}_Y(T > t)$ is equal to the right hand side of (17). In
what follows, we will refer to $F_N(t, \mathcal{A}^{(N,2)})$ defined in (16) as the discrete survival function.

Theorem 1 has an intuitive interpretation. Taking the case $\theta = 1$, the $e^{-t/2}$ represents the prob-
ability that the two lineages have not coalesced by time t due to ordinary Wright-Fisher/Kingman
coalescence. Against this smooth backdrop there are $Y(t)$ points, representing essentially instanta-
neous HR events, in which the lineages have a large probability, $\psi^2/4$, of coalescing. Thus there is
an additional factor in the survival function representing the probability the pair does not coalesce
in any of these HR events. The case $\theta \in (0, 1)$ is analogous except the timescale is so short that
there is no chance of an ordinary Wright-Fisher/Kingman coalescent event.

Note that when $\lambda = 0$, there are no large reproduction events and $Y(t) \equiv 0$. Then for $\theta \in (0, 1)$,
the right hand side of (17) is 1, i.e. there is no coalescence with probability 1. For $\theta = 1$, the right
hand side of (17) is $e^{-t/2}$ which is expected from the cumulative distribution function (CDF) of
the Kingman coalescent for a sample of size 2, with our timescale. The degenerate case $\lambda > 0$ but
$\psi = 0$ effectively gives these same results for any $Y(t)$.

Proof of Theorem 1

Recall that each $g \in \mathbb{Z}_{\geq 0}$ is a WF generation (resp. HR generation) with probability $1 - \alpha_N$ (resp. α_N), independently for all $g \in \mathbb{Z}_{\geq 0}$. The number of HR generations in $\{0, 1, \ldots, G-1\}$, denoted by $H_N(G)$, therefore has the binomial distribution $\text{Bin}(G, \alpha_N)$.

We begin by addressing the technical point that we cannot actually know just by looking at the pedigree whether generation g is a HR generation. But we do know whether a "big family" occurs. Let us define this as the event that, spanning generations $g + 1$ and g, there exists a pair of parents with at least $[\psi N]$ children. A big family must happen if g is a an HR generation, but it can also happen with a small probability ϵ_N when g is a WF generation. Precisely,

$$\epsilon_N \leq \binom{N}{2} \binom{N}{[\psi N]} \left(\frac{1}{\binom{N}{2}} \right)^{[\psi N]} \leq \frac{2^{[\psi N]-1}}{N^{[\psi N]-2}}.$$

Let $Q_N(G)$ be the number of generations $g \in \{0, 1, \ldots, G-1\}$ such that a big family event occurs between generations $g + 1$ and g. Then $Q_N(G)$ is extremely close to the binomial variable $H_N(G) \sim \text{Bin}(G, \alpha_N)$ because

$$H_N(G) \leq Q_N(G) \quad \text{and} \quad Q_N(G) \leq \text{Bin}(G, \alpha_N + \epsilon_N), \tag{18}$$

where the first inequality holds almost surely and the second is a stochastic dominance. Since $\alpha_N = \frac{\lambda}{N^\theta}$, for each $t \in (0, \infty)$ and $\theta \in (0, 1]$ we have convergence in distribution

$$Q_N([tN^\theta]) \to Y(t) \qquad \text{as } N \to \infty \tag{19}$$

which is identical to the limiting result for $H_N([tN^\theta])$. In other words, ϵ_N is so small for any sizeable N, that we are safe in assuming that big families in the pedigree reliably signify HR generations.

Indeed, from the discussion above we have

$$\lim_{N \to \infty} \mathbb{E}^{(N)} \left[\left| \left(1 - \frac{\psi^2}{4} \right)^{H_N([tN])} - \left(1 - \frac{\psi^2}{4} \right)^{Q_N([tN])} \right|^2 \right] = 0 \tag{20}$$

so that we can (and will) in the following computations replace $H_N([tN])$ by $Q_N([tN])$ without changing any limit as $N \to \infty$.

Proof of (17) **when** $\theta = 1$ In this case it suffices to show that

$$\lim_{N \to \infty} \mathbb{E}^{(N)} \left[\left| F_N(t, \mathcal{A}^{(N,2)}) - e^{-t/2} \left(1 - \frac{\psi^2}{4} \right)^{Q_N([tN])} \right|^2 \right] = 0. \tag{21}$$

Expanding the square in (21) gives

$$\mathbb{E}^{(N)} \left[F_N^2(t, \mathcal{A}^{(N,2)}) - 2 e^{-t/2} F_N(t, \mathcal{A}^{(N,2)}) \left(1 - \frac{\psi^2}{4} \right)^{Q_N([tN])} + e^{-t} \left(1 - \frac{\psi^2}{4} \right)^{2 Q_N([tN])} \right], \tag{22}$$

which requires the computation of three expectations. The first is the expectation of the square of the discrete survival function,

$$\mathbb{E}^{(N)} \left[F_N^2(t, \mathcal{A}^{(N,2)}) \right]. \tag{23}$$

The second is the expectation of the discrete survival function times the probability that a single pair of lineages does not coalesce in any of the HR events in the pedigree up to time t,

$$\mathbb{E}^{(N)}\left[F_N(t, \mathcal{A}^{(N,2)})\left(1 - \frac{\psi^2}{4}\right)^{Q_N([tN])}\right], \tag{24}$$

The third is the expectation of the square of the same, latter probability that a single pair of lineages does not coalesce in any of the HR events in the pedigree up to time t,

$$\mathbb{E}^{(N)}\left[\left(1 - \frac{\psi^2}{4}\right)^{2\,Q_N([tN])}\right]. \tag{25}$$

First term in (22)   The expectation in (23) can be computed by considering two samples of size 2 whose lineage dynamics are conditionally independent given $\mathcal{A}^{(N,2)}$. Genetically, this corresponds to the ancestral processes of two unlinked loci given the pedigree and the two sampled individuals, and where one gene copy has been sampled at each locus from each of the individuals. Let τ and τ' be the coalescence times of these two pairs of sampled gene copies. Then it holds that

$$\mathbb{E}^{(N)}\left[F_N^2(t, \mathcal{A}^{(N,2)})\right] = \mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{\mathcal{A}^{(N,2)}}\left(\tau > [tN]\right)\,\mathbb{P}^{(N)}_{\mathcal{A}^{(N,2)}}\left(\tau' > [tN]\right)\right]$$

$$= \mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{\mathcal{A}^{(N,2)}}\left(\tau > [tN],\ \tau' > [tN]\right)\right]$$

$$= \mathbb{P}^{(N)}\left(\tau > [tN],\ \tau' > [tN]\right) \tag{26}$$

where the second equality follows from the assumed conditionally independence (and we use the short-hand notation $\mathbb{P}_{\mathcal{A}^{(N,2)}}(\cdot)$ for (16)). In order to compute the limit as $N \to \infty$ in (26), we introduce the ancestral process of two conditionally independent samples given the pedigree.

Joint diploid ancestral process   The stochastic dynamics of the two conditionally independent, given the pedigree, pairs of lineages are described by the *joint diploid ancestral process* $\widetilde{M} :=$ $(\widetilde{M}_g)_{g \in \mathbb{Z}_{\geq 0}}$. This is a Markov chain with state space $\mathcal{S} = \{\xi_{00}^{(4)}, \xi_{00}^{(3)}, \ldots, \xi_\Delta\}$ described below, where $\widetilde{M}_g$ is the state of the two pairs of lineages in a common pedigree g generations backwards in time. Denote by $\widetilde{\Pi}_N$ its transition matrix, the derivation of its entries is available at A.1.1 and its entries are available at A.1.2 for an HR generation and at A.1.3 for a WF generation.

Similarly to the proof of Lemma 1, denote by $\bullet$ an ancestral lineage of a gene copy in the first pair and by $\star$ the same for the second pair. Parentheses are used to denote individuals. More precisely, consider the following 10 states:

$$\xi_{00}^{(4)} = (\bullet)(\bullet)(\star)(\star)$$

$$\xi_{00}^{(3)} = (\bullet)(\bullet\star)(\star)$$

$$\xi_{00}^{(2)} = (\bullet\star)(\bullet\star)$$

$$\xi_{10}^{(2)} = (\bullet\bullet\star)(\star)$$

$$\xi_{01}^{(2)} = (\star\star\bullet)(\bullet)$$

$$\xi_{10}^{(3)} = (\bullet\bullet)(\star)(\star)$$

$$\xi_{01}^{(3)} = (\star\star)(\bullet)(\bullet)$$

$$\xi_{11}^{(2)} = (\bullet\bullet)(\star\star)$$

$$\xi_{11}^{(1)} = (\bullet\bullet\star\star)$$

$$\xi_\Delta = \text{coal.}$$

The superscript indicates the total number of individuals in which the 4 ancestral lineages reside. The two subscripts tell us the states of the two pairs respectively: 0 means a pair of lineages in state ξ_0 and 1 means a pair of lineages in state ξ_1, with these as defined in the proof of Lemma 1. For example, $\xi_{10}^{(3)}$ involves 3 individuals in which the first pair of lineages are in the the same individual and the second pair of lineages is in different individuals. Finally, the state ξ_Δ is an absorbing state which represents the event that at least one of the two pairs has coalesced.

By definition, the two pairs of gene copies are drawn from the same pair of individuals at the present generation $g = 0$, where for each pair one gene copy is picked from each of the individuals. Hence the initial state $\widetilde{M}_0$ must be $\xi_{00}^{(2)}$. In other words, the distribution $\vec{p}_0$ of $\widetilde{M}_0$ is given by

$$\vec{p}_0 = (0, 0, 1, \cdots, 0). \tag{27}$$

It follows from Lemma 2 in Appendix Section A.1.4 that

$$\lim_{N\to\infty} \mathbb{P}^{(N)}\left(\tau > [tN],\, \tau' > [tN]\right) = \lim_{N\to\infty} \vec{p}_0 \cdot \widetilde{\Pi}_N^{[tN]} (1, \cdots, 1, 0)^T \tag{28}$$

$$= (0, 0, 1, 0, \cdots, 0)\widetilde{P}e^{t\widetilde{G}}(1, \cdots, 1, 0)^T \tag{29}$$

$$= e^{-t}e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, \tag{30}$$

where (28) follows from the definition of $\widetilde{M}$ and $\vec{p}_0$, (29) from (Möhle, 1998a, Lemma 1) as explained in Section A.1.4 and (30) by Lemma 2. Note that the vector $(1, \cdots, 1, 0)^T$ in (28)-(29) amounts to the Markov chain $(\widetilde{M}_g)_{g\in\mathbb{Z}_{\geq 0}}$ not reaching state ξ_Δ, i.e. that neither pair has coalesced.

Importantly, Lemma 2 and Theorem 1 both hold for any initial state $\vec{p}_0$ whose last coordinate is zero. This is because the sample will undergo an instantaneous adjustment by $\widetilde{P}$ given in (A20), so that the effective starting state is always $\xi_{00}^{(4)}$. Also, owing to $\widetilde{G}$, the lineages will remain in state $\xi_{00}^{(4)}$ except for instants in which they have a chance to coalesce. This robustness against initial condition is analogous to (14). Probably the three states of most biological interest are $\xi_{00}^{(2)}$, $\xi_{11}^{(2)}$ and $\xi_{11}^{(1)}$, and Lemma 2 and Theorem 1 hold for these initial states as well.

Second term in (22) We now show that (24) converges to $e^{-t/2}e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}$ as $N \to \infty$. Through the use of the law of total expectation, (24) is equal to

$$\sum_{k=0}^{[tN]} \mathbb{E}^{(N)}\left[\mathbb{P}_{\mathcal{A}^{(N,2)}}^{(N)}\left(\tau > [tN]\right) \mid Q_N([tN]) = k\right] \left(1 - \frac{\psi^2}{4}\right)^k \mathbb{P}^{(N)}(Q_N([tN]) = k). \tag{31}$$

By the fact that $Q_N([tN])$ is known given the pedigree and an application of the tower property, the conditional expectation in (31) is equal to

$$\mathbb{P}^{(N)}\left(\tau > [tN] \mid Q_N([tN]) = k\right),$$

which is approximately equal to

$$\mathbb{P}^{(N)}\left(\tau > [tN] \mid H_N([tN]) = k\right),$$

by (18). That is to say

$$\mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{\mathcal{A}^{(N,2)}}(\tau > [tN])\left(1 - \frac{\psi^2}{4}\right)^{Q_N([tN])}\right] \approx \mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{H_N}(\tau > [tN])\left(1 - \frac{\psi^2}{4}\right)^{H_N([tN])}\right],$$

in the sense that (20) holds. By Lemma 3 in the Appendix, for $g = [tN]$, it follows that for each $N \geq 2$ and $t \in (0, \infty)$,

$$\mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{H_N}(\tau > [tN])\left(1 - \frac{\psi^2}{4}\right)^{H_N([tN])}\right] = (1,0,0)\left(\Pi^{\mathrm{mid}}_N\right)^{[tN]}(1,1,0)^T, \tag{32}$$

where Π^{mid}_N is defined as

$$\Pi^{\mathrm{mid}}_N := \alpha_N\left(1 - \frac{\psi^2}{4}\right)\Pi^{\mathrm{HR}}_N + (1 - \alpha_N)\Pi^{\mathrm{WF}}_N. \tag{33}$$

It now follows by Lemma 4 that the right hand side of (32) converges to $e^{-t/2}e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}$.

Third term in (22) Finally, (25) is computed by first noticing that the number of highly reproductive pairs up to generation T, $Q_N(T)$, is (almost) binomially distributed according to $\mathrm{Bin}([T], \alpha_N)$ for all $T \geq 1$, as observed by (18). Using the probability generating function of $Q_N(T)$ we get that the third term in (22) is equal to $e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}$.

Putting everything together As $N \to \infty$, (22) is equal to 0 since (25) multiplied by e^{-t} and (23) add up to $2e^{-t}e^{-\lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16})t}$ which cancel out with (24) multiplied by $-2e^{-t}$. This gives (21) which concludes the proof of Theorem 1 in the case of $\theta = 1$.

Convergence (17) **when** $\theta \in (0, 1)$ The proof is similar to the case of $\theta = 1$. In all of the above, substitute $[tN]$ by $[tN^\theta]$, and show instead of (22) that

$$\lim_{N \to \infty} \mathbb{E}^{(N)}\left[\left|\left|F_N(t, \mathcal{A}^{(N,2)}) - \left(1 - \frac{\psi^2}{4}\right)^{Q_N([tN^\theta])}\right|\right|^2\right] = 0. \tag{34}$$

Expanding (34) gives the same three terms as in (23)-(25). In this faster timescale, as $N \to \infty$, (23) is now equal to

$$\lim_{N \to \infty} \mathbb{E}^{(N)}\left[F_N^2(t, \mathcal{A}^{(N,2)})\right] = e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, \tag{35}$$

as available in Lemma 2. The limiting behavior of (24) is the same as before, that is

$$\lim_{N \to \infty} \mathbb{E}^{(N)}\left[F_N(t, \mathcal{A}^{(N,2)})\left(1 - \frac{\psi^2}{4}\right)^{Q_N([tN])}\right] = e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})} \tag{36}$$

and

$$\lim_{N \to \infty} \mathbb{E}^{(N)}\left[\left(1 - \frac{\psi^2}{4}\right)^{2Q_N([tN^\theta])}\right] = e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}. \tag{37}$$

Multiplying (36) by -2 and summing it up with (35) and (36) concludes the proof in the case of $\theta \in (0, 1)$.

The proof of Theorem 1 is complete. $\qquad\square$

Remark 2 (Only the high-reproduction events matter). Let $\vec{g}^{(N)} = (g_1, g_2, \cdots)$ be the (random) generations at which a big family event occurred (between generations $g_i + 1$ and g_i), where $0 \leq g_1 < g_2 < \cdots$. Then $\vec{g}^{(N)}$ is known if we know the pedigree. Similar to (16), we let

$$F_N(t, \vec{g}^{(N)}) := \mathbb{P}^{(N)}\left(\tau^{(N,2)} > [tN^\theta] \mid \vec{g}^{(N)}\right) \tag{38}$$

be the conditional probability of the event $\{\tau^{(N,2)} > [tN^\theta]\}$ given the (random) generations $\vec{g}^{(N)}$. Hence, here we condition on less information than on the left hand side of (17). We can show that Theorem 1 still holds (i.e. the weak convergence in (17) still holds) if we replace $F_N(t, \mathcal{A}^{(N,2)})$ by $F_N(t, \vec{g}^{(N)})$.

Remark 3 (Joint convergence of conditionally independent coalescence times). Recall from (26) that τ and τ' are the coalescence times of two samples of size 2 whose lineage dynamics are conditionally independent given $\mathcal{A}^{(N,2)}$. By Lemma 2, the pair $(\tau/N^\theta, \tau'/N^\theta)$ converges in distribution to a pair of non-negative random variables (T, T') that is conditionally independent given the Poisson process Y in Theorem 1.

More generally, suppose $\frac{1}{N^\theta}\vec{g}^{(N)}$ converges in probability to $\vec{t} = (t_1, t_2, \cdots)$ where $0 < t_1 < t_2 < \cdots$ is a strictly increasing positive sequence. We conjecture that then the law of the pair $(\tau/N^\theta, \tau'/N^\theta)$, under the conditional probability $\mathbb{P}(\cdot \mid \vec{g}^{(N)})$, converges as $N \to \infty$ to the law of a pair of random variables (T, T') under a probability $\mathbb{P}_{\vec{t}}$ such that

$$\mathbb{P}_{\vec{t}}(T > t, T' > s) = \mathbb{P}_{\vec{t}}(T > t)\,\mathbb{P}_{\vec{t}}(T > s) = \begin{cases} \left(1 - \frac{\psi^2}{4}\right)^{N(t)+N(s)}, & \text{when } \theta \in (0,1), \\[2ex] e^{\frac{-(t+s)}{2}}\left(1 - \frac{\psi^2}{4}\right)^{N(t)+N(s)}, & \text{when } \theta = 1, \end{cases} \tag{39}$$

where $N(t) = \max\{i \in \mathbb{Z}_{\geq 0} : t_i < t\}$, and $t_0 = 0$.

Coalescence times, gene genealogies and correlations

Here we briefly recap then provide three illustrations of our results. Our main result is Theorem 1 which describes two limiting ancestral processes conditional on the pedigree. In this case, the probability of coalescence in a generation depends on whether that particular generation includes a big family. For background and comparison, Lemma 1 presents the corresponding two limiting processes obtained by the usual method of averaging over pedigrees, i.e. over all possible outcomes of reproduction in a single generation, including the possibility of a big family. In this case, the probability of coalescence is the same in every generation.

Time is re-scaled in all of these limiting ancestral processes. It is measured in units of N^θ generations for some $\theta \in (0, 1]$. When $\theta \in (0, 1)$, the timescale for big families to occur is much shorter than the usual Wright-Fisher coalescent timescale of N generations. When $\theta = 1$, the timescales for big families and for ordinary Wright-Fisher coalescence are the same. Big families occur at rate λ in re-scaled time, and their offspring comprise a fraction $\psi \in [0, 1]$ of the population in that generation. Underpinning our results is the fact that as $N \to \infty$ ancestral genetic lineages spend the overwhelming majority of their time in separate individuals, i.e. in state ξ_0 for a pair of lineages at the same locus (cf. Lemma 1) or state $\xi_{00}^{(4)}$ for two pairs of lineages at two unlinked loci (cf. Theorem 1). Thus when a big family occurs, each lineage independently: (i) is among the offspring of the highly reproductive pair with probability ψ and (ii) if so, is equally likely to descend

from each of the four copies of the corresponding locus in the two parents. A pair of lineages at the same locus coalesces in the big family with probability $\psi^2/4$. Pairs of lineages at different, unlinked loci do this independently.

Our first illustration compares our limiting results to the cumulative distribution function (CDF, i.e. one minus the survival function) of pairwise coalescence times in the discrete model. Figure 1 displays CDFs for five simulated pedigrees for $N = 500$, assuming that the probability of a big family is equal to the expected pairwise coalescence probability, $1/(2N) = 0.001$, and the offspring make up the entire population in that generation. This corresponds to the limiting process in Theorem 1 with $\theta = 1$, $\lambda = 1/2$ and $\psi = 1$. This makes coalescence probability $(\psi^2/4)$ equal to $1/4$ in each HR generation. We computed coalescence probabilities on each pedigree in each generation starting from a pair of randomly sampled individuals using the method in Wakeley et al. (2012). The corresponding "expected" CDF of the pedigree-averaged process from Lemma 1, i.e. of an exponential random variable with rate parameter $5/8$, is shown for comparison.

The left panel of Figure 1 illustrates that the ancestral process conditional on the pedigree is quite close to limiting result in Theorem 1, even when $N = 500$. The CDFs make discrete jumps whenever big families occur. The magnitude of a jump is always $1/4$ of the remaining distance to 1. Between jumps the CDFs show a steady increase in the cumulative coalescence probability, in line with the limiting prediction with its rate of $1/2$. None of these CDFs are close to the prediction of the pedigree-averaged process in Lemma 1.

Figure 1: Cumulative distribution functions (CDFs) of pairwise coalescence times for $\theta = 1$ and $\lambda = 1/2$. Left: CDFs for five simulated pedigrees for populations of size $N = 500$ together with the corresponding expected CDF from Lemma 1. Right: The same five CDFs and the corresponding expectation from Lemma 1, only plotted over the most recent 40 generations.

The right panel of Figure 1 details the short-time behavior of the ancestral process conditional on the pedigree, displaying these same CDFs only over the most recent 40 generations. The scale on the vertical axis is such that the diagonal corresponds approximately to the prediction of the background Wright-Fisher model (not shown) and a line with slope 1.25 corresponds approximately to prediction of Lemma 1 which is shown. After a small number of generations, which from Chang (1999) should be of order $\log_2(N)$, the CDFs for the five pedigrees start to show the predicted Wright-Fisher slope of one. However, they start at different places depending on the particular ancestries of the sampled individuals, specifically whether there are very recent shared ancestors as in pedigree 5 or more likely there are no very recent shared ancestors as in pedigrees 1 through 4;

cf. also Wakeley et al. (2012). These differences are barely visible on the timescale of the left panel of Figure 1, and it is implicit in Theorem 1 that they become negligible as $N \to \infty$.

Next we illustrate the effects that big families have on the gene genealogies of larger samples, in particular the sharing of identical coalescence times at unlinked loci. Rather than simulating pedigrees for finite populations, we use the limiting model directly so that big families are the only possible cause of shared coalescence times. Here we use the HR-only model, $\theta \in (0,1)$, but still assuming $\psi = 1$. We set $\lambda = 1$ without loss of generality, as λ is arbitrary except when $\theta = 1$.

Based on Theorem 1, we model gene genealogies by generating a series of exponential waiting times between HR events and, since $\theta \in (0,1)$, disallowing coalescence between them. When the n ancestral lineages of the sample reach the first HR event, their distribution among the four parental gene copies will be multinomial with parameters n and $(1/4, 1/4, 1/4, 1/4)$. Anywhere from one to four simultaneous multiple mergers will occur. The number of ancestral lineages which emerge is also at most four. If more than one lineage emerges, the same process is repeated until a single lineage remains which is the most recent common ancestor of the entire sample. The only aspects of the pedigree which persist in the limit are the HR events (cf. Remark 2). Thus, independent runs of this multinomial coalescent process using the same series of exponential waiting times correspond to gene genealogies of unlinked loci conditional on the pedigree.

The top row of Figure 2 displays the gene genealogies of seven unlinked loci for a sample of size 16, assuming in this way that all loci share the same pedigree. The trees are oriented with the present-day samples at the bottom. Solid lines trace (unlabeled) ancestral lineages up into the past. Thin dotted lines show the times of the HR events. All seven gene genealogies have multiple-mergers at the first HR event in the past, and five have common ancestor events at the second HR event. In the more distant past when there are small numbers of ancestral lineages, there is less sharing of coalescence times among gene genealogies. This is expected; for example, the final two lineages only coalesce with probability $1/4$ at each HR event.

The bottom row of Figure 2 shows seven gene genealogies, again for samples of size 16, but now assuming that each locus has its own pedigree. These are equivalent to seven gene genealogies sampled from seven independent populations, each with its own series of exponential waiting times between HR events (not shown in the bottom row). These gene genealogies differ from the ones in the top row, most obviously in the different timings of their first common ancestor events. Clearly, the distribution of gene genealogies produced in this way will not be close to the distribution of gene genealogies of unlinked loci in the same genome which perforce come the same population.

Finally, we illustrate how averaging over pedigrees as in Lemma 1 results in positive correlations of coalescence times between unlinked loci. Explicitly modeling pedigrees as in Theorem 1 predicts these to be zero as might be expected for independently assorting loci. Based on the property of ancestral lineages spending the overwhelming majority of their time in separate individual, we consider the two-locus analogue of Lemma 1 with reduced state space

$$\xi_{00} = (\bullet)(\bullet)(\star)(\star)$$
$$\xi_{10} = (\bullet)(\star)(\star)$$
$$\xi_{01} = (\bullet)(\bullet)(\star)$$
$$\xi_{11} = (\bullet)(\star)$$

where now the first and second subscripts are indicators of whether locus 1 or locus 2 has coalesced. By extension from Lemma 1, the limiting ancestral process for two unlinked loci has transition rate

Figure 2: Simulated gene genealogies for seven independently assorting loci when all seven share the same pedigree (top row) versus when each locus has its own independently generated pedigree (bottom row). The sample size is $n = 16$ for every locus. Gene genealogies were generated as described in the text for the limiting model with $\theta \in (0,1)$ and $\lambda = \psi = 1$. Thin dotted lines in the top row show the particular series of times of big families in that population.

matrix

$$
Q = \begin{cases} \lambda Q^{\text{HR}}, & \text{if } \theta \in (0,1) \\[2ex] Q^{\text{WF}} + \lambda Q^{\text{HR}}, & \text{if } \theta = 1 \end{cases} \tag{40}
$$

where

$$
Q^{\text{HR}} = \begin{array}{c} \\ \xi_{00} \\ \xi_{10} \\ \xi_{01} \\ \xi_{11} \end{array} \begin{array}{cccc} \xi_{00} & \xi_{10} & \xi_{01} & \xi_{11} \\ \left[-\frac{\psi^2}{4}\left(2 - \frac{\psi^2}{4}\right) & \frac{\psi^2}{4}\left(1 - \frac{\psi^2}{4}\right) & \frac{\psi^2}{4}\left(1 - \frac{\psi^2}{4}\right) & \frac{\psi^4}{16} \right. \\ 0 & -\frac{\psi^2}{4} & 0 & \frac{\psi^2}{4} \\ 0 & 0 & -\frac{\psi^2}{4} & \frac{\psi^2}{4} \\ \left. 0 & 0 & 0 & 0 \right] \end{array} \tag{41}
$$

and

$$
Q^{\mathrm{WF}} = \begin{array}{c} \\ \xi_{00} \\ \xi_{10} \\ \xi_{01} \\ \xi_{11} \end{array} \begin{array}{c} \begin{array}{cccc} \xi_{00} & \xi_{10} & \xi_{01} & \xi_{11} \end{array} \\ \begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{array}.
\tag{42}
$$

Focusing on the case $\theta = 1$, the rate matrix Q is the sum of a Wright-Fisher (or Kingman coalescent) component and a high-reproduction component. We have factored the tuning parameter λ out of the latter to emphasize that, conditional on a HR event, samples at the two loci coalesce or do not coalesce independently of each other.

Let T_1 and T_2 be the coalescence times at the two loci. These correspond to the limiting random variables in Lemma 2, called T and T' in Remark 3. In the present context, these are the times to state ξ_{11} starting from states ξ_{01} and ξ_{10}, respectively. From the rate matrix Q in (40) or from Lemma 1 directly, T_1 and T_2 are identically distributed. In particular,

$$
T_1 \sim \begin{cases} \text{exponential}\left(\lambda \frac{\psi^2}{4}\right), & \text{if } \theta \in (0,1) \\[2mm] \text{exponential}\left(\frac{1}{2} + \lambda \frac{\psi^2}{4}\right), & \text{if } \theta = 1. \end{cases}
\tag{43}
$$

However, T_1 and T_2 are not necessarily independent. We use first-step analysis to compute the correlation coefficient, $\mathrm{Corr}[T_1, T_2]$. Let W be the waiting time to the first event in the ancestry of the two loci starting from state ξ_{00}, and T_1^* and T_2^* be the additional times to coalescence at each locus following the first event. In this formulation,

$$
T_i = W + T_i^*
\tag{44}
$$

for $i \in \{1, 2\}$. From (40), we have

$$
W \sim \begin{cases} \text{exponential}\left(\lambda \frac{\psi^2}{4}\left(2 - \frac{\psi^2}{4}\right)\right), & \text{if } \theta \in (0,1) \\[2mm] \text{exponential}\left(1 + \lambda \frac{\psi^2}{4}\left(2 - \frac{\psi^2}{4}\right)\right), & \text{if } \theta = 1. \end{cases}
\tag{45}
$$

and we point out that, since W is exponentially distributed,

$$
\mathbb{E}[W^2] = 2\mathbb{E}[W]^2.
\tag{46}
$$

Conditioning on the first step from state ξ_{00} and simplifying,

$$
\mathbb{E}[T_1 T_2] = \mathbb{E}[W^2] + \mathbb{E}[W]\mathbb{E}[T_1^*] + \mathbb{E}[W]\mathbb{E}[T_2^*] + \mathbb{E}[T_1^* T_2^*]
\tag{47}
$$

$$
= 2\mathbb{E}[W]\mathbb{E}[T_1].
\tag{48}
$$

Going from (47) to (48) uses (46), (44), $\mathbb{E}[T_1] = \mathbb{E}[T_2]$, and the fact that either T_1^* or T_2^* or both are equal to zero following the first event. Then for the correlation coefficient, we have simply

$$
\mathrm{Corr}[T_1, T_2] = \frac{2\mathbb{E}[W]\mathbb{E}[T_1] - \mathbb{E}[T_1]^2}{\mathrm{Var}[T_1]},
\tag{49}
$$

which, using (43) and (45), becomes

$$\text{Corr}[T_1, T_2] = \begin{cases} \dfrac{\psi^2}{8 - \psi^2}, & \text{if } \theta \in (0,1) \quad\quad\quad\text{(50a)} \\[3mm] \dfrac{\lambda\psi^4}{16 + \lambda\psi^2(8 - \psi^2)}, & \text{if } \theta = 1. \quad\quad\quad\text{(50b)} \end{cases}$$

Even though the loci assort independently, their ancestries in the pedigree-averaging model jointly depend on the random process of HR events occurring in the population. As a result, their coalescence times are positively correlated.

The correlation coefficient (50b), obtained here under the assumption that the loci are unlinked, corresponds to Equation 31 in Birkner et al. (2013b, p. 266), obtained there by modeling recombination explicitly and then taking the limit as the re-scaled recombination parameter tends to infinity. The timescales in these two works differ by a factor of two. Our (50b) becomes identical to Equation 31 in Birkner et al. (2013b) by putting $\lambda = c/2$.

For a given value of ψ, the correlation coefficient is smaller when $\theta = 1$ than when $\theta \in (0,1)$. In the case that both high-reproduction and Wright-Fisher/Kingman coalescence are possible ($\theta = 1$), the correlation tends to zero as λ tends to zero. As λ grows, (50b) grows until it approaches (50a). Thus, HR events may be said to be the source of positive correlations in coalescence times at unlinked loci. In a similar vein, $\text{Corr}[T_1, T_2]$ tends to zero as the fraction of the population replaced in each HR event, ψ, tends to zero. This is true even if $\theta \in (0,1)$, i.e. when there is no Wright-Fisher/Kingman component in the limit process. At the other extreme, as $\psi \to 1$, $\text{Corr}[T_1, T_2] \to 1/7$ which is considerably less than one. Even when all coalescence happens in HR events and the children of the big family replace the entire population, there are still two diploid parents and the loci will generally have different coalescence times.

The following alternate derivation of (50a) shows how these positive correlations arise. In short it is because T_1 and T_2 have a shared dependence on the times between big families in the pedigree. Implicitly, Lemma 1 averages over these times whereas Theorem 1 retains them.

When $\theta \in (0,1)$, coalescence can only happen when there is a high-reproduction event. Let K_1 and K_2 be the numbers of HR events it takes for locus 1 and locus 2 to coalesce, respectively. These do not depend on the times between HR events when $\theta \in (0,1)$. Further, K_1 and K_2 are independent because the loci are unlinked. They are geometric random variables with parameter $\psi^2/4$. Let X_i, $i \in \mathbb{Z}_{\geq 0}$, be the time from the $(i-1)$th to the ith HR event backward in time, with $X_0 \equiv 0$. In the context of Theorem 1, these times are independent and identically distributed exponential random variables with rate parameter λ. Under this formulation,

$$T_i = \sum_{j=1}^{K_i} X_j \quad\quad\quad (51)$$

for $i \in \{1,2\}$. There are two sources of variation in T_i: variation in K_i and variation in the lengths of the intervals, X_j, $j \in \{1, \ldots, K_i\}$. Starting with (51), it is straightforward to confirm that the distribution of T_i is exponential with rate parameter $\lambda\psi^2/4$ as in (43) or Lemma 1.

From (51) and the fact that X_i and X_j are independent for $i \neq j$, it is also clear that intervals in a common to T_1 and T_2 are a key source of their covariation. For given values of K_1 and K_2,

they are the only source. The first interval is always shared, as are all subsequent intervals until one or the other locus coalesces. Let K_{12} be the number of these shared intervals and

$$T_{12} = \sum_{i=1}^{K_{12}} X_i \tag{52}$$

be the corresponding total length of time. By definition $K_{12} = \min(K_1, K_2)$. The more ancient $K_1 - K_2$ or $K_2 - K_1$ intervals are only ancestral to one of the loci.

Applying the conditional covariance formula, or law of total covariance, we have

$$\mathrm{Cov}[T_1, T_2] = \mathbb{E}\big[\mathrm{Cov}[T_1, T_2 | K_1, K_2]\big] + \mathrm{Cov}\big[\mathbb{E}[T_1 | K_1, K_2], \mathbb{E}[T_2 | K_1, K_2]\big]$$

$$= \mathbb{E}\big[\mathrm{Var}[T_{12} | K_{12}]\big] + \mathrm{Cov}\big[\mathbb{E}[T_1 | K_1], \mathbb{E}[T_2 | K_2]\big]. \tag{53}$$

The outer expectation and covariance are with respect to the joint distribution of K_1 and K_2. Note that K_{12} is a marginal property of this distribution. The inner variance (or covariance) and expectations are with respect to the joint distributions of the X_i which are the only parts of T_1 and T_2 that vary conditional on K_1 and K_2.

At this point, in (53), we have not applied the fundamental property that K_1 and K_2 are independent since the loci are unlinked, nor have we assumed any particular distribution(s) for the X_i. We have only used the definitions of T_1 and T_2 as sums of random variables and the assumption that X_i and X_j are independent for $i \neq j$. So we may consider that the interval times are fixed numbers: $X_i \equiv x_i$, $i \in \mathbb{Z}_{\geq 0}$. They could be the outcomes of the exponential random times implicit in Theorem 1. Fixing the X_i means fixing the only aspects of the pedigree that persist in the limiting model. Conditioning on the pedigree, T_1 and T_2 are independent even in the limiting model; cf. Remark 3. The point we wish to emphasize here is that fixing the X_i removes one particular source of covariation of T_1 and T_2. It makes $\mathrm{Var}[T_{12} | K_{12}] = 0$.

Continuing from (53) and assuming that X_i, $i \in \mathbb{Z}_{\geq 0}$, are independent and identically distributed

$$\mathrm{Cov}[T_1, T_2] = \mathbb{E}[K_{12}]\mathrm{Var}[X_i] + \mathbb{E}[X_i]^2\mathrm{Cov}[K_1, K_2]$$

$$= \mathbb{E}[K_{12}]\mathrm{Var}[X_i], \tag{54}$$

the latter following from the independence of K_1 and K_2. Again, $X_i \sim \mathrm{exponential}(\lambda)$, and from the definition of K_{12} as the number of HR events it takes for one locus or the other to coalesce,

$$K_{12} \sim \mathrm{geometric}\left(1 - \left(1 - \frac{\psi^2}{4}\right)\right). \tag{55}$$

Putting the required quantities in (54) and simplifying gives

$$\mathrm{Cov}[T_1, T_2] = \frac{16}{\lambda^2 \psi^2 (8 - \psi^2)} \tag{56}$$

which is exactly the covariance needed to produce the correlation coefficient (50a). In sum, the model of Lemma 1 predicts a positive correlation of coalescence times at unlinked loci because it averages over the distributions of the intervals X_i. Starting instead with the model of Theorem 1 shows that the particular quantity controlling these positive correlations is $\mathrm{Var}[X_i]$.

Discussion

The use of random models to describe past events raises many questions in population genetics. Everything in the past has already occurred, including all instances and timings of reproduction and genetic transmission. For empirical work this may be a truism. But population genetics has always been concerned with evolutionary processes. How do mutation, recombination, selection, random genetic drift, non-random mating, limited dispersal, etc., conspire to produce observable patterns of variation? By emphasizing the fixed nature of the past, we highlight the subjectivity of theoretical work, specifically when the goal is to interpret data from natural populations. Ultimately, the choices one makes about modeling the past may be application-dependent.

Here, motivated by applications to multi-locus data, we singled out the pedigree as a key feature of the past and obtained a result (Theorem 1) concerning the application of neutral coalescent models in sexually reproducing species. We have the following sampling structure in mind. Processes of survival and reproduction result in a pedigree. Genetic transmission, including mutation and recombination across the entire genome, occurs within the pedigree. A number of individuals are sampled from the population and some or all of their genomes are sequenced. We modeled the single-locus coalescent process conditional on the pedigree. Our results specify the distribution of coalescence times given the pedigree and the sampled individuals. This distribution can be interpreted either as a prior for a single locus or as a prediction about the distribution of coalescence times among unlinked loci.

We contrasted our results conditional on the pedigree with results obtained by averaging over pedigrees, noting that the latter is the tradition of theoretical population genetics. We did not model mutation or recombination, but our fundamental conclusion—that some population processes cause the quenched and averaged processes to be very different—should be as important for genetic variation as it is for coalescence times. We also did not model selection, which is a special kind of population process in which survival and reproduction (and hence the pedigree) depends on the genotypes of individuals. But we note that for loci unlinked to the locus or loci under selection the pedigree contains all the information needed to predict patterns of coalescence and variation under neutral mutation and recombination.

We can compare our the framework with that of Ralph (2019). The two have a lot in common. Ralph (2019) takes the pedigree and the outcomes of genetic transmission, including recombination across the entire genome, to be fixed. The latter is referred to as the ancestral recombination graph (ARG), which we note differs slightly from the corresponding objects in Hudson (1983a) and Griffiths and Marjoram (1997) because it is embedded in the fixed pedigree. Without specifying a generative model for the pedigree, Ralph (2019) focuses on the ARG as the fixed but unknown object of interest in empirical population genetics. A sample is taken and some stretch of the genome is sequenced. Its ancestry is a collection of gene genealogies, a subset of the ARG. Implicitly, it is the outcome of the random process of genetic transmission within the fixed pedigree, but this too is not modeled.

The only randomness is in how the collection of gene genealogies of the sample is revealed by mutation. Ralph (2019) assumes the infinite-sites mutation process and uses this to show that predictions about summary statistics of DNA sequence variation, such as the average number of pairwise nucleotide differences or the F-statistics of Patterson et al. (2012), can be expressed in terms of the fixed branch lengths in the sampled subset of the ARG. This is the empirical version of what Slatkin (1991), Griffiths and Tavaré (1998), Nielsen (2000), McVean (2002) and Peter (2016) had done in the context of the standard neutral coalescent, where instead the moments of

summary statistics can be expressed in terms of corresponding moments of branch lengths. Ralph et al. (2020) describe a hybrid approach, with the ARG conceived as in Ralph (2019) and with times of events in the ARG for data from humans (The 1000 Genomes Project Consortium, 2015) estimated with the aid of the standard neutral coalescent (Speidel et al., 2019).

Whereas we model the production of the pedigree and the process of coalescence within it but do not model mutation, Ralph (2019) models only mutation on the fixed ARG. Consider a species in which recurrent selective sweeps across the genome have structured the ARG. An empirical estimate of the ARG would find regions of the genome with reduced variation due to reduced times to common ancestry. In order to relate these observations to an evolutionary process within the empirical framework of Ralph (2019), for example to describe them in terms of recurrent selective sweeps as in Durrett and Schweinsberg (2005), additional modeling would be needed. In contrast, in a theoretical approach such as ours here, recurrent sweeps would be included in the model at the outset, and this in turn would facilitate the interpretation of patterns in the data. Under our model, it is important to keep in mind that the ARG is in fact a fixed object and that the process of coalescence within the pedigree models the sampling of a locus in the ARG.

Today detailed estimates of the ARG for large samples of human genomes are available (Wohns et al., 2022; Zhang et al., 2023). These have been obtained, like other recent estimates (Kelleher et al., 2019; Speidel et al., 2019; Albers and McVean, 2020), using the standard neutral coalescent as a prior for gene genealogies and times to common ancestry. Our results and those of Tyukin (2015) help to justify using such a prior despite the fact that the pedigree is fixed, so long as the processes which laid down the pedigree are not too different from the Wright-Fisher or Cannings models with relatively low variation of offspring numbers. The empirically oriented interpretations in these works, for example in Wohns et al. (2022), connect features of the ARG with major events in human history, such as the out-of-Africa event which has been studied genetically since the first mtDNA discoveries (Cann et al., 1987; Vigilant et al., 1991) and the novel finding of an accumulation of ancestry in Papua New Guinea more than 100-thousand years ago. This is intraspecific phylogeography (Avise et al., 1987; Avise, 1989, 2000) at genome scale.

Our model for generating the pedigree includes the possibility of a high-reproduction (HR) event in each generation. We obtained different coalescent processes as $N \to \infty$, depending on the relative rate of HR events in the limit and whether the ancestral process is conditional on the pedigree (Theorem 1) or not (Lemma 1). This essentially negative result, that the averaged process cannot be used in place of the conditional process, includes the positive finding that the Kingman coalescent can be used between HR events in the case that both occur on the same timescale (Theorem 1 with $\theta = 1$). The numbers and timings of HR events are all that is left of the pedigree in the limit (cf. Remark 2). Needing to keep track of just these is much less daunting than the prospect of including entire pedigrees in all of our population-genetic models. There may be other circumstances in which aspects of the pedigree are important, but so far the only other instance identified is when sub-populations are connected by limited migration (Wilton et al., 2017).

Limiting coalescent processes for our model generally involve simultaneous multiple-mergers. Yet the familiar extensions of the Kingman coalescent to include multiple-mergers have been derived by averaging over the pedigree, not by conditioning on it. They begin with single-generation marginal probabilities of coalescence, whereas in truth the individuals in the sample either have or do not have common ancestors in any preceding generation and this is what determines probabilities of coalescence. Without HR events, our results and those of Tyukin (2015) provide belated justification for the early uses of the Kingman coalescent process as a prior model for the gene genealogy of a single locus (Lundstrom et al., 1992; Griffiths and Tavaré, 1994; Kuhner et al.,

1995). Our work also clarifies what is involved in using pedigree-averaged ancestral processes as single-locus priors in cases where big families can occur.

If the only data available were from a single locus without recombination, one could model the gene genealogy using the pedigree-averaged ancestral process. The logic would be that a single locus has one unknown random pedigree and one unknown random gene genealogy within that pedigree, and that the gene genealogies from multiple-mergers coalescent models are marginal predictions over both of these unknowns. For example, with $n = 2$, a single draw of a coalescence time from the appropriate exponential distribution in Lemma 1 accounts for both sources of variation. Implicit in this accounting is that repeated samples would each have their own pedigree and conditional gene genealogy. This two-fold sampling structure is precisely what Theorem 1 describes. It is straightforward to show that repeated sampling under Theorem 1 (each time drawing a new pedigree) gives the same exponential distributions as in Lemma 1. Yet even for a single locus, it may be preferable to record the additional information about HR events as in Theorem 1.

Applying this type of repeated sampling (i.e. including re-sampling the pedigree) to multiple loci is another matter. Population-genetic models should not allow the pedigree to vary among loci. Theorem 1 is a simple initial example of the kind of coalescent modeling required for multi-locus data generally but especially when multiple-mergers processes are implicated. In cases where HR events may occur with some frequency, it is crucial to retain the information about the pedigree which matters for the gene genealogies at all loci. All multiple-mergers coalescent models so far described, which implicitly average over the pedigree, are inadequate in this sense. The broader implication of Lemma 1 and Theorem 1 is that there exists a collection of quenched limits conditional on the pedigree which await description and are the appropriate models for multi-locus data.

The diploid exchangeable population models in Birkner et al. (2018) are a natural starting point for the description of general quenched-pedigree Ξ-coalescent models. Alternatively, parameterized models could be considered, controlling for example rates of monogamy and the distribution of offspring numbers as in the program SLiM 3 (Haller and Messer, 2019) or the Pólya urn scheme of Gasbarra et al. (2005). The latter was used for the prior in the Bayesian inference methods of Gasbarra et al. (2007a,b) and Ko and Nielsen (2019) for estimating the recent few generations of the pedigree from sequence data. Non-exchangeable models for generating pedigrees are possible, for example with recurrent selective sweeps (Durrett and Schweinsberg, 2005) or cultural transmission of reproductive success (Guez et al., 2023). It could also be of interest to describe these coalescent models directly in terms of the properties of pedigrees as directed graphs, and here we note the study of Blath et al. (2014) as a start in this direction.

The model underlying Lemma 1 and Theorem 1 is very simple. Only one type of big family is allowed, HR events occur as a Poisson process, and we only considered a sample of size two. We hypothesize that the basic principles of Theorem 1 will be robust to all of these. For example, other ways of generating HR events should be possible, such that $Y(t) \sim \text{Poisson}(\lambda t)$ would be replaced by some other distribution. Extensions to larger sample sizes and to variation in the numbers and types of big families seem straightforward in principle, though they will require a lot more bookkeeping. Finally we emphasize that it will be necessary to specify HR events in sufficient detail that Mendel's laws can be applied. For example, if four parents have $[\psi N]$ offspring, it will matter whether they form two monogamous pairs or comprise one big family with four parents, and in either case just how many offspring each pair has. In a more general model, each HR event will need to include these details.

The issues we raise here about pedigrees parallel those in recent work on population bottlenecks. Like the trajectories of population sizes through time in coalescent hidden Markov models,

bottlenecks have traditionally been considered fixed events of the past. But models of recurrent bottlenecks have recently been considered. A bottleneck is the event that a population ordinarily of size N_0 has size $N_B < N_0$ for a period of time. In Birkner et al. (2009, Section 6) it was shown that a Ξ-coalescent describes the limiting gene-genealogical process for a model with recurrent severe bottlenecks, specifically with the bottleneck duration going to zero and $N_B/N_0 \to 0$ as both N_B and N_0 go to infinity. González Casanova et al. (2022) used a similar framework but allowed that N_B could be finite. They described a new class of Ξ-coalescents they called the symmetric coalescent. Our model gives one of these when the offspring replace the entire population ($\psi = 1$), since this is identical to a short drastic bottleneck (González Casanova et al., 2022, Definition 3) with $N_B = 4$ and our θ and λ corresponding to their α and $k^{(N)}$. But as noted in Birkner et al. (2009), Ξ-coalescent models are only obtained for recurrent bottlenecks by averaging over the exponential process which generates them. When the times and severities of bottlenecks are fixed, the result will depend on these and will not be a time-homogeneous Markov process. Against this backdrop of similarities, a small but notable difference is that the bottleneck models in Birkner et al. (2009) and González Casanova et al. (2022) are haploid rather than diploid.

The proof of Theorem 1 uses the idea of two independent copies of the coalescent process on the same pedigree. Genetically, these are the gene-genealogies of two independently assorting loci conditional on their shared pedigree. Our results suggest a reinterpretation of population-genetic models which predict non-zero correlations or positive covariances of coalescence times at unlinked loci. In the context of high reproduction, Eldon and Wakeley (2008) found that the correlation could be positive in an effectively haploid model of recombination and HR events, and Birkner et al. (2013b) extended this to a diploid model very similar to the one we studied here. It appears that such correlations result from averaging over the pedigree. In the simple model we have considered, they arise from averaging over the times of HR events, cf. (53) and (54). On any fixed pedigree the correlation of coalescence times at unlinked loci must be zero.

The comparison with recurrent bottlenecks is apt here as well. Schaper et al. (2012) constructed a recurrent-bottleneck model for recombination and coalescence at two loci with recombination. They note that what is relevant for data is the covariance of coalescence times conditional on the series bottleneck events in the ancestry of the population, not the unconditional covariance which averages over these. They showed that the conditional covariance goes to zero as the recombination parameter goes to infinity. They also showed, in the Ξ-coalescent limit of Birkner et al. (2009) that the covariance could be positive even as the recombination parameter goes to infinity. We note an analogous finding for yet another model in Wakeley and Lessard (2003), in which non-zero correlations of coalescence times at unlinked resulted from taking the number of subpopulations to infinity in an island migration model, even though for any finite number of subpopulations the correlation goes to zero as the recombination parameter tends to infinity.

In sum, for a century it has been common practice in population genetics to compute probabilities of events by averaging over an assumed process of reproduction. What we have shown is that when big families occur with some frequency, or more generally when the descendants of a small number of individuals take over a sizable fraction of the population in a short period of time, this averaging is not justified and can produce spurious results. Instead, such extreme outcomes of reproduction need to be treated as fixed, and probabilities of coalescence and other events conditioned upon them. In light of this, existing multiple-mergers coalescent models must be reassessed and most likely replaced with conditional or quenched models. A comparison with how population size has been treated as fixed is of some interest because it too is an outcome of reproduction. In both cases, it is when population-genetic models are applied to explain variation among loci that the importance of conditioning on past events is most readily apparent.

Data availability

Mathematica notebooks containing some calculations, detailed in the Appendix, are available at:
`https://github.com/DimitriDiam/Bursts-of-coalescence.git`.

Acknowledgements

This work was support in part by National Science Foundation grants DMS-1855417 and DMS-2152103, and Office of Naval Research grant N00014-20-1-2411 to Wai-Tong (Louis) Fan. Dimitrios Diamantidis is supported by NSF DMS-2152103.

A   Appendix

A.1   Joint diploid ancestral process

Recall the joint ancestral process $\widetilde{M}$ describes the lineage dynamics of two conditionally independent samples of size 2. In section A.1.1 we establish its Markov property. This property should hold for any population model in which the outcomes of reproduction in different generations are independent. In section A.1.2 we provide its transition matrix for a high-reproduction generation and in section A.1.3 we study two extreme cases, $\psi = 0$ and $\psi = 1$. In the former case we recover the transition matrix of the joint diploid ancestral process for a Wright-Fisher generation and in the latter we consider the case when the highly reproductive couple replaces the entire population.

A.1.1   The joint diploid ancestral process with 10 states is Markovian

Recall the state space of the process $\widetilde{M} = (\widetilde{M}_g)_{g \in \mathbb{Z}_{\geq 0}}$ is $\mathcal{S} = \{\xi_{00}^{(4)}, \xi_{00}^{(3)}, \ldots, \xi_\Delta\}$ which has 10 states including the state ξ_Δ where one, the other or both pairs of lineages have coalesced.

Fix a pair of states $(\xi, \eta) \in \mathcal{S}^2$. It holds that

$$\mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \{\widetilde{M}_k\}_{k=0}^{g-1} \right)$$

$$= \sum_{\alpha^{(g)}} \mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \alpha^{(g)}, \ \{\widetilde{M}_k\}_{k=0}^{g-1} \right) \mathbb{P}^{(N)}(\alpha^{(g)} \mid \widetilde{M}_g = \xi, \{\widetilde{M}_k\}_{k=0}^{g-1}) \tag{A1}$$

$$= \sum_{\alpha^{(g)} \in A_{\xi,\eta}} \mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \alpha^{(g)} \right) \mathbb{P}^{(N)}(\alpha^{(g)}), \tag{A2}$$

where the last sum $\sum_{\alpha^{(g)} \in A_{\xi,\eta}}$ is over all possible parent assignments $\alpha^{(g)}$ that belong to the pattern $A_{\xi,\eta}$ described in Tables A1-A3. These are exactly the parent assignments for which transition to η is *possible* in the sense that $\mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \alpha^{(g)} \right) > 0$ if and only if $\alpha^{(g)} \in A_{\xi,\eta}$.

In the last equality above, we noted that a parent assignment $\alpha^{(h)}$ between generations g and $g+1$ is independent of $\{\widetilde{M}_k\}_{k=0}^{g}$ and so $\mathbb{P}^{(N)}(\alpha^{(g)} \mid \widetilde{M}_g = \xi, \{\widetilde{M}_k\}_{k=0}^{g-1}) = \mathbb{P}^{(N)}(\alpha^{(g)})$. Also, $\mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \alpha^{(g)}, \{\widetilde{M}_k\}_{k=0}^{g-1} \right) = \mathbb{P}^{(N)} \left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \ \alpha^{(g)} \right)$ because it is determined entirely by Mendelian randomness and not by the population model (whether it is Wright-Fisher or something else). This conditional probability is computed in Tables A4-A10, and we only

need to compute this for those $\alpha^{(g)}$ in patterns whose probability is of order $O(\frac{1}{N})$ or larger in view of Möhle (1998a, Lemma 1).

In particular, we have shown that the number $\widetilde{\pi}_{\xi,\eta}^{(N)} := \mathbb{P}^{(N)}\left(\widetilde{M}_{g+1} = \eta \mid \widetilde{M}_g = \xi, \{\widetilde{M}_k\}_{k=0}^{g-1}\right)$ does not depend on the "history" $\{\widetilde{M}_k\}_{k=0}^{g-1}$. Hence the process $\widetilde{M}$ is Markovian.

A.1.2 Transition matrix for a high-reproduction generation

For any $\xi \in \mathcal{S}$, let $\vec{r}_\xi = (\widetilde{\pi}_{\xi,\eta}^{(N)})_{\eta \in \mathcal{S}}$ be the ξth row of $\widetilde{\Pi_N}$. By (A2), each row of $\widetilde{\Pi_N}$ is computed as

$$\vec{r}_\xi = A_\xi \cdot \vec{p}_\xi,$$

where A_ξ is given by the appropriate matrix in Tables A4-A10 corresponding to ξ, and $\vec{p}_\xi$ contains the probabilities of the patterns from Tables A1-A3 used in all the transitions from ξ to $\eta \in \mathcal{S}$. When a highly reproductive pair appears, $\widetilde{\Pi}_N$ is given as

$$\widetilde{\Pi_N^{\mathrm{HR}}} = \begin{pmatrix} r_{\xi_{00}^{(4)}}^{\mathrm{HR}} \\ \vdots \\ r_{\xi_\Delta}^{\mathrm{HR}} \end{pmatrix} + O(N^{-2}), \tag{A3}$$

where r_ξ^{HR} in (A3) is the row of $\widetilde{\Pi}_N^{\mathrm{HR}}$ that corresponds to initial state ξ up to an error of order $O(N^{-2})$, given in (A4)-(A8). For example, $r_{\xi_{11}^{(1)}}^{\mathrm{HR}}$ is derived by multiplying the matrix $A_{\xi_{11}^{(1)}}$ given in Table A4 by the vector column $\vec{p} := \left(\frac{1-\psi}{N}, \psi + (1-\psi)(1-\frac{1}{N})\right)^T$ on the right. The calculations for each row are available at https://github.com/DimitriDiam/Bursts-of-coalescence.git in the **PHR_derivation+WF+x=1.pdf** file and can be reproduced in the corresponding Mathematica notebook **PHR_derivation+WF+x=1.nb**, where Mathematica version 13.1.0 (Wolfram Research, Inc., 2022) was used. Its rows are given as

$$r_{\xi_{11}^{(1)}}^{\mathrm{HR}} = \begin{array}{c} \xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta \end{array} \begin{pmatrix} 0 \\ 0 \\ \frac{\psi-1}{N}+1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{1-\psi}{4N} \\ \frac{3-3\psi}{4N} \end{pmatrix}^T, \quad r_{\xi_{11}^{(2)}}^{\mathrm{HR}} = \begin{array}{c} \xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_\Delta \end{array} \begin{pmatrix} \frac{4\psi^2+2\psi-6}{N}-\psi^2+1 \\ \frac{4-4\psi}{N} \\ \psi^2 \\ \frac{1-\psi^2}{2N} \\ 0 \\ \frac{1-\psi^2}{2N} \\ 0 \\ 0 \\ 0 \\ \frac{-3\psi^2+2\psi+1}{N} \end{pmatrix}^T, \tag{A4}$$

$$
r^{\mathrm{HR}}_{\xi^{(2)}_{10}} =
\begin{array}{c}
\xi^{(4)}_{00} \\
\xi^{(3)}_{00} \\
\xi^{(2)}_{00} \\
\xi^{(3)}_{01} \\
\xi^{(2)}_{01} \\
\xi^{(3)}_{10} \\
\xi^{(2)}_{10} \\
\xi^{(2)}_{11} \\
\xi^{(1)}_{11} \\
\xi_{\Delta}
\end{array}
\begin{pmatrix}
0 \\
\frac{3\psi^2}{N} - \frac{3}{N} - \psi^2 + 1 \\
-\frac{\psi}{N} + \frac{1}{N} + \frac{\psi^2}{2} \\
0 \\
-\frac{\psi}{N} + \frac{1}{N} + \frac{\psi^2}{4} \\
0 \\
\frac{1}{2N} - \frac{\psi^2}{2N} \\
0 \\
0 \\
-\frac{5\psi^2}{2N} + \frac{2\psi}{N} + \frac{1}{2N} + \frac{\psi^2}{4}
\end{pmatrix}^{T}
, \
r^{\mathrm{HR}}_{\xi^{(3)}_{10}} =
\begin{array}{c}
\xi^{(4)}_{00} \\
\xi^{(3)}_{00} \\
\xi^{(2)}_{00} \\
\xi^{(3)}_{01} \\
\xi^{(2)}_{01} \\
\xi^{(3)}_{10} \\
\xi^{(2)}_{10} \\
\xi^{(2)}_{11} \\
\xi^{(1)}_{11} \\
\xi_{\Delta}
\end{array}
\begin{pmatrix}
-\frac{45\psi^3}{2N} + \frac{83\psi^2}{2N} - \frac{13\psi}{N} - \frac{6}{N} + \frac{3\psi^3}{2} - \frac{5\psi^2}{2} + 1 \\
\frac{5\psi^3}{N} - \frac{\psi^2}{N} - \frac{8\psi}{N} + \frac{4}{N} - 2\psi^3 + 2\psi^2 \\
-\frac{\psi^3}{N} + \frac{\psi^2}{N} + \frac{\psi^3}{2} \\
\frac{\psi^3}{2N} - \frac{\psi}{N} + \frac{1}{2N} - \frac{\psi^3}{4} + \frac{\psi^2}{4} \\
-\frac{\psi^3}{4N} + \frac{\psi^2}{4N} + \frac{\psi^3}{4} \\
\frac{\psi^3}{4N} - \frac{\psi^2}{4N} - \frac{\psi}{2N} + \frac{1}{2N} \\
0 \\
\frac{\psi^2}{4N} - \frac{\psi^3}{4N} \\
0 \\
\frac{73\psi^3}{4N} - \frac{167\psi^2}{4N} + \frac{45\psi}{2N} + \frac{1}{N} + \frac{\psi^2}{4}
\end{pmatrix}^{T}
,
$$

$$(A5)$$

$$
r^{\mathrm{HR}}_{\xi^{(2)}_{01}} =
\begin{array}{c}
\xi^{(4)}_{00} \\
\xi^{(3)}_{00} \\
\xi^{(2)}_{00} \\
\xi^{(3)}_{01} \\
\xi^{(2)}_{01} \\
\xi^{(3)}_{10} \\
\xi^{(2)}_{10} \\
\xi^{(2)}_{11} \\
\xi^{(1)}_{11} \\
\xi_{\Delta}
\end{array}
\begin{pmatrix}
0 \\
\frac{3\psi^2}{N} - \frac{3}{N} - \psi^2 + 1 \\
-\frac{\psi}{N} + \frac{1}{N} + \frac{\psi^2}{2} \\
0 \\
\frac{1}{2N} - \frac{\psi^2}{2N} \\
0 \\
-\frac{\psi}{N} + \frac{1}{N} + \frac{\psi^2}{4} \\
0 \\
0 \\
-\frac{5\psi^2}{2N} + \frac{2\psi}{N} + \frac{1}{2N} + \frac{\psi^2}{4}
\end{pmatrix}^{T}
, \
r^{\mathrm{HR}}_{\xi^{(3)}_{01}} =
\begin{array}{c}
\xi^{(4)}_{00} \\
\xi^{(3)}_{00} \\
\xi^{(2)}_{00} \\
\xi^{(3)}_{01} \\
\xi^{(2)}_{01} \\
\xi^{(3)}_{10} \\
\xi^{(2)}_{10} \\
\xi^{(2)}_{11} \\
\xi^{(1)}_{11} \\
\xi_{\Delta}
\end{array}
\begin{pmatrix}
-\frac{45\psi^3}{2N} + \frac{83\psi^2}{2N} - \frac{13\psi}{N} - \frac{6}{N} + \frac{3\psi^3}{2} - \frac{5\psi^2}{2} + 1 \\
\frac{5\psi^3}{N} - \frac{\psi^2}{N} - \frac{8\psi}{N} + \frac{4}{N} - 2\psi^3 + 2\psi^2 \\
-\frac{\psi^3}{N} + \frac{\psi^2}{N} + \frac{\psi^3}{2} \\
\frac{\psi^3}{2N} - \frac{\psi}{N} + \frac{1}{2N} - \frac{\psi^3}{4} + \frac{\psi^2}{4} \\
0 \\
\frac{\psi^3}{4N} - \frac{\psi^2}{4N} - \frac{\psi}{2N} + \frac{1}{2N} \\
-\frac{\psi^3}{4N} + \frac{\psi^2}{4N} + \frac{\psi^3}{4} \\
\frac{\psi^2}{4N} - \frac{\psi^3}{4N} \\
0 \\
\frac{73\psi^3}{4N} - \frac{167\psi^2}{4N} + \frac{45\psi}{2N} + \frac{1}{N} + \frac{\psi^2}{4}
\end{pmatrix}^{T}
,
$$

$$(A6)$$

$$
r^{\mathrm{HR}}_{\xi^{(2)}_{00}} =
\begin{matrix}
\xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_{\Delta}
\end{matrix}
\begin{pmatrix}
\frac{\psi^2}{N} + \frac{\psi}{2N} - \frac{3}{2N} - \frac{\psi^2}{4} + \frac{1}{4} \\[4pt]
\frac{\psi^2}{N} - \frac{\psi}{2N} - \frac{1}{2N} - \frac{\psi^2}{2} + \frac{1}{2} \\[4pt]
-\frac{\psi}{4N} + \frac{1}{4N} + \frac{1}{4} \\[4pt]
\frac{1}{8N} - \frac{\psi}{8N} \\[4pt]
-\frac{\psi}{4N} + \frac{1}{4N} + \frac{\psi^2}{8} \\[4pt]
\frac{1}{8N} - \frac{\psi}{8N} \\[4pt]
-\frac{\psi}{4N} + \frac{1}{4N} + \frac{\psi^2}{8} \\[4pt]
\frac{\psi^2}{32} \\[4pt]
-\frac{\psi}{16N} + \frac{1}{16N} + \frac{\psi^2}{32} \\[4pt]
-\frac{2\psi^2}{N} + \frac{17\psi}{16N} + \frac{15}{16N} + \frac{7\psi^2}{16}
\end{pmatrix}^{T}
,\;
r^{\mathrm{HR}}_{\xi^{(3)}_{00}} =
\begin{matrix}
\xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_{\Delta}
\end{matrix}
\begin{pmatrix}
-\frac{25\psi^3}{2N} + \frac{22\psi^2}{N} - \frac{13\psi}{2N} - \frac{3}{N} + \psi^3 - \frac{3\psi^2}{2} + \frac{1}{2} \\[4pt]
-\frac{29\psi^3}{4N} + \frac{77\psi^2}{4N} - \frac{25\psi}{2N} + \frac{1}{2N} - \frac{\psi^3}{2} + \frac{1}{2} \\[4pt]
\frac{3\psi^3}{4N} - \frac{3\psi^2}{4N} - \frac{\psi^3}{4} + \frac{\psi^2}{2} \\[4pt]
\frac{5\psi^3}{16N} + \frac{3\psi^2}{16N} - \frac{\psi}{N} + \frac{1}{2N} - \frac{\psi^3}{8} + \frac{\psi^2}{8} \\[4pt]
\frac{\psi^2}{2N} - \frac{\psi}{N} + \frac{1}{2N} + \frac{\psi^2}{8} \\[4pt]
\frac{5\psi^3}{16N} + \frac{3\psi^2}{16N} - \frac{\psi}{N} + \frac{1}{2N} - \frac{\psi^3}{8} + \frac{\psi^2}{8} \\[4pt]
-\frac{\psi^3}{8N} + \frac{5\psi^2}{8N} - \frac{\psi}{N} + \frac{1}{2N} + \frac{\psi^2}{8} \\[4pt]
-\frac{\psi^3}{16N} + \frac{\psi^2}{16N} + \frac{\psi^3}{32} \\[4pt]
-\frac{3\psi^3}{32N} + \frac{3\psi^2}{32N} + \frac{\psi^3}{32} \\[4pt]
\frac{597\psi^3}{32N} - \frac{1349\psi^2}{32N} + \frac{23\psi}{N} + \frac{1}{2N} - \frac{\psi^3}{16} + \frac{\psi^2}{2}
\end{pmatrix}^{T}
$$

$$\tag{A7}$$

$$
\text{and}\quad r^{\mathrm{HR}}_{\xi^{(4)}_{00}} =
\begin{matrix}
\xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_{\Delta}
\end{matrix}
\begin{pmatrix}
\frac{\left(239\psi^2 - 350\psi - 24\right)(\psi-1)^2}{4N} + (2\psi+1)(\psi-1)^2 \\[4pt]
\frac{-59\psi^4 + 121\psi^3 - 18\psi^2 - 60\psi + 16}{4N} - 2(\psi-1)\psi^2 \\[4pt]
\frac{\psi^2 - \psi^4}{8N} + \frac{\psi^4}{4} \\[4pt]
\frac{-11\psi^4 + 23\psi^3 - \psi^2 - 15\psi + 4}{8N} - \frac{1}{4}(\psi-1)\psi^2 \\[4pt]
\frac{7(\psi-1)\psi^3}{8N} - \frac{1}{8}(\psi-2)\psi^3 \\[4pt]
\frac{-11\psi^4 + 23\psi^3 - \psi^2 - 15\psi + 4}{8N} - \frac{1}{4}(\psi-1)\psi^2 \\[4pt]
\frac{7(\psi-1)\psi^3}{8N} - \frac{1}{8}(\psi-2)\psi^3 \\[4pt]
\frac{\left(\psi^2 - 3\psi + 2\right)\psi^2}{16N} + \frac{\psi^4}{32} \\[4pt]
\frac{\psi^4}{32} - \frac{(\psi-1)\psi^3}{16N} \\[4pt]
\frac{-351\psi^4 + 1383\psi^3 - 1794\psi^2 + 754\psi + 8}{8N} - \frac{1}{16}\psi^2\left(\psi^2 - 8\right)
\end{pmatrix}^{T}
.
$$

$$\tag{A8}$$

A.1.3   Two extremes for the sizes of big families

Degenerate case of no offspring, $\psi = 0$. When a Wright Fisher generation is taking place, denote $\widetilde{\Pi}_N$ by $\widetilde{\Pi}^{\mathrm{WF}}_N$. The matrix $\widetilde{\Pi}^{\mathrm{WF}}_N$ can be recovered from $\widetilde{\Pi}^{\mathrm{HR}}_N$ by setting $\psi = 0$ up to a term of order $O(N^{-2})$. The additional $O(N^{-2})$ term represents cases in which both pairs of lineages coalesce and which become negligible in the limit. Note there is a single case of non-negligible double coalescence

 in the $O(N^{-1})$ transition $\xi_{00}^{(2)} \to \xi_\Delta$. The matrix $\widetilde{\Pi}_N^{WF}$ is given as

$$\widetilde{\Pi}_N^{WF} =$$

$\mathcal{S}$	$\xi_{00}^{(4)}$	$\xi_{00}^{(3)}$	$\xi_{00}^{(2)}$	$\xi_{10}^{(3)}$	$\xi_{10}^{(2)}$	$\xi_{01}^{(3)}$	$\xi_{01}^{(2)}$	$\xi_{11}^{(2)}$	$\xi_{11}^{(1)}$	ξ_Δ
$\xi_{00}^{(4)}$	$1-\frac{6}{N}$	$\frac{4}{N}$	0	$\frac{1}{2N}$	0	$\frac{1}{2N}$	0	0	0	$\frac{1}{N}$
$\xi_{00}^{(3)}$	$\frac{1}{2}-\frac{3}{N}$	$\frac{1}{2}+\frac{1}{2N}$	0	$\frac{1}{2N}$	$\frac{1}{2N}$	$\frac{1}{2N}$	$\frac{1}{2N}$	0	0	$\frac{1}{2N}$
$\xi_{00}^{(2)}$	$\frac{1}{4}-\frac{3}{2N}$	$\frac{1}{2}-\frac{1}{2N}$	$\frac{1}{4}+\frac{1}{4N}$	$\frac{1}{8N}$	$\frac{1}{4N}$	$\frac{1}{8N}$	$\frac{1}{4N}$	0	$\frac{1}{16N}$	$\frac{15}{16N}$
$\xi_{10}^{(3)}$	$1-\frac{6}{N}$	$\frac{4}{N}$	0	$\frac{1}{2N}$	0	$\frac{1}{2N}$	0	0	0	$\frac{1}{N}$
$\xi_{10}^{(2)}$	0	$1-\frac{3}{N}$	$\frac{1}{N}$	0	$\frac{1}{2N}$	0	$\frac{1}{N}$	0	0	$\frac{1}{2N}$
$\xi_{01}^{(3)}$	$1-\frac{6}{N}$	$\frac{4}{N}$	0	$\frac{1}{2N}$	0	$\frac{1}{2N}$	0	0	0	$\frac{1}{N}$
$\xi_{01}^{(2)}$	0	$1-\frac{3}{N}$	$\frac{1}{N}$	0	$\frac{1}{N}$	0	$\frac{1}{2N}$	0	0	$\frac{1}{2N}$
$\xi_{11}^{(2)}$	$1-\frac{6}{N}$	$\frac{4}{N}$	0	$\frac{1}{2N}$	0	$\frac{1}{2N}$	0	0	0	$\frac{1}{N}$
$\xi_{11}^{(1)}$	0	0	$1-\frac{1}{N}$	0	0	0	0	0	$\frac{1}{4N}$	$\frac{3}{4N}$
ξ_Δ	0	0	0	0	0	0	0	0	0	1

$$+\; O\left(\frac{1}{N^2}\right).$$

$$(A9)$$

Offspring make up the whole population, $\psi = 1$. When $\psi = 1$, the highly reproductive pair gives birth to the entire next population. This is the case of pure Mendelian randomness within a single family with two parents. The matrix $\widetilde{\Pi}_N^{HR}$ is then equal to

$\mathcal{S}$	$\xi_{00}^{(4)}$	$\xi_{00}^{(3)}$	$\xi_{00}^{(2)}$	$\xi_{10}^{(3)}$	$\xi_{10}^{(2)}$	$\xi_{01}^{(3)}$	$\xi_{01}^{(2)}$	$\xi_{11}^{(2)}$	$\xi_{11}^{(1)}$	ξ_Δ
$\xi_{00}^{(4)}$	0	0	$\frac{1}{4}$	0	$\frac{1}{8}$	0	$\frac{1}{8}$	$\frac{1}{32}$	$\frac{1}{32}$	$\frac{7}{16}$
$\xi_{00}^{(3)}$	0	0	$\frac{1}{4}$	0	$\frac{1}{8}$	0	$\frac{1}{8}$	$\frac{1}{32}$	$\frac{1}{32}$	$\frac{7}{16}$
$\xi_{00}^{(2)}$	0	0	$\frac{1}{4}$	0	$\frac{1}{8}$	0	$\frac{1}{8}$	$\frac{1}{32}$	$\frac{1}{32}$	$\frac{7}{16}$
$\xi_{01}^{(3)}$	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$	0	0	$\frac{1}{4}$
$\xi_{01}^{(2)}$	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$	0	0	$\frac{1}{4}$
$\xi_{10}^{(3)}$	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$	0	0	0	0	$\frac{1}{4}$
$\xi_{10}^{(2)}$	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$	0	0	0	0	$\frac{1}{4}$
$\xi_{11}^{(2)}$	0	0	1	0	0	0	0	0	0	0
$\xi_{11}^{(1)}$	0	0	1	0	0	0	0	0	0	0
ξ_Δ	0	0	0	0	0	0	0	0	0	1

$$.\qquad (A10)$$

The entries of the matrix above can be explained in terms of Mendel's laws of independent segregation and independent assortment, given that all individuals have the same two parents. For example, in the first row the initial state is $\xi_{00}^{(4)}$: all four lineages in distinct individuals. The transition to state $\xi_{11}^{(1)}$, where all four lineages are in the same individual but have not coalesced, takes place with probability $1/32$ and is computed as follows. Each lineage chooses a parent with probability $1/2$, then chooses a gene copy within that parent also with probability $1/2$. All four lineages choose the same parent (which can be either of the two) with probability $1/8$. Then each pair independently chooses distinct gene copies (i.e. does not coalesce) with probability $1/2$.

A.1.4 Limiting behavior of the joint diploid ancestral process

The joint diploid ancestral process has transition matrix

$$\widetilde{\Pi}_N := (1 - \alpha_N)\widetilde{\Pi}_N^{\mathrm{WF}} + \alpha_N \widetilde{\Pi}_N^{\mathrm{HR}}. \tag{A11}$$

One can write

$$\widetilde{\Pi}_N^{\mathrm{WF}} = \widetilde{A}^{\mathrm{WF}} + \frac{1}{N}\widetilde{B}^{\mathrm{WF}} + o\!\left(\frac{1}{N}\right), \tag{A12}$$

where

$$
\widetilde{A}^{\mathrm{WF}} :=
\begin{array}{c}
\begin{array}{ccccccccccc}
\mathcal{S} & \xi_{00}^{(4)} & \xi_{00}^{(3)} & \xi_{00}^{(2)} & \xi_{10}^{(3)} & \xi_{10}^{(2)} & \xi_{01}^{(3)} & \xi_{01}^{(2)} & \xi_{11}^{(2)} & \xi_{11}^{(1)} & \xi_\Delta
\end{array}\\
\begin{array}{c}
\xi_{00}^{(4)}\\ \xi_{00}^{(3)}\\ \xi_{00}^{(2)}\\ \xi_{10}^{(3)}\\ \xi_{10}^{(2)}\\ \xi_{01}^{(3)}\\ \xi_{01}^{(2)}\\ \xi_{11}^{(2)}\\ \xi_{11}^{(1)}\\ \xi_\Delta
\end{array}
\left[
\begin{array}{cccccccccc}
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
\frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
\frac{1}{4} & \frac{1}{2} & \frac{1}{4} & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0\\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{array}
\right]
\end{array}
\tag{A13}
$$

and

$$
\widetilde{B}^{\mathrm{WF}} :=
\begin{array}{c}
\begin{array}{ccccccccccc}
\mathcal{S} & \xi_{00}^{(4)} & \xi_{00}^{(3)} & \xi_{00}^{(2)} & \xi_{10}^{(3)} & \xi_{10}^{(2)} & \xi_{01}^{(3)} & \xi_{01}^{(2)} & \xi_{11}^{(2)} & \xi_{11}^{(1)} & \xi_\Delta
\end{array}\\
\begin{array}{c}
\xi_{00}^{(4)}\\ \xi_{00}^{(3)}\\ \xi_{00}^{(2)}\\ \xi_{10}^{(3)}\\ \xi_{10}^{(2)}\\ \xi_{01}^{(3)}\\ \xi_{01}^{(2)}\\ \xi_{11}^{(2)}\\ \xi_{11}^{(1)}\\ \xi_\Delta
\end{array}
\left[
\begin{array}{cccccccccc}
-6 & 4 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 & 1\\
-3 & \frac{1}{2} & 0 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 & 0 & \frac{1}{2}\\
-\frac{3}{2} & -\frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \frac{1}{4} & \frac{1}{8} & \frac{1}{4} & 0 & \frac{1}{16} & \frac{15}{16}\\
-6 & 4 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 & 1\\
0 & -3 & 1 & 0 & \frac{1}{2} & 0 & 1 & 0 & 0 & \frac{1}{2}\\
-6 & 4 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 & 1\\
0 & -3 & 1 & 0 & 1 & 0 & \frac{1}{2} & 0 & 0 & \frac{1}{2}\\
-6 & 4 & 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 & 1\\
0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & \frac{1}{4} & \frac{3}{4}\\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{array}
\right]
\end{array}.
\tag{A14}
$$

Similarly, we split

$$\widetilde{\Pi}_N^{\mathrm{HR}} = \widetilde{A}^{\mathrm{HR}} + \frac{1}{N}\widetilde{B}^{\mathrm{HR}} + o\!\left(\frac{1}{N}\right) \tag{A15}$$

with

$$\widetilde{A}^{\mathrm{HR}} = \left(r_{\xi_{00}^{(4)}}^{(1)}, \cdots, r_{\xi_\Delta}^{(1)}\right)^T, \quad \widetilde{B}^{\mathrm{HR}} = \frac{1}{N}\left(r_{\xi_{00}^{(4)}}^{(2)}, \cdots, r_{\xi_\Delta}^{(2)}\right)^T \tag{A16}$$

defined as $r_\xi^{(1)} := \lim_{N\to\infty} r_\xi^{\mathrm{HR}}$ and $r_\xi^{(2)} := \lim_{N\to\infty} N \cdot (r_\xi^{\mathrm{HR}} - r_\xi^{(1)})$ for all $\xi \in \mathcal{S}$. We provide only the rows of $\widehat{A}^{\mathrm{HR}}$ since those are the only ones we will need for the subsequent computations. In particular those are given as

$$
r_{\xi_{11}^{(1)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}^{T}
,\
r_{\xi_{11}^{(2)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} 1-\psi^2 \\ 0 \\ \psi^2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}^{T}
,\
r_{\xi_{10}^{(2)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} 0 \\ 1-\psi^2 \\ \frac{\psi^2}{2} \\ 0 \\ \frac{\psi^2}{4} \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{\psi^2}{4} \end{pmatrix}^{T}
,\
r_{\xi_{10}^{(3)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} \frac{3\psi^3}{2}-\frac{5\psi^2}{2}+1 \\ 2\psi^2(1-\psi) \\ \frac{\psi^3}{2} \\ \frac{\psi^2-\psi^3}{4} \\ \frac{\psi^3}{4} \\ 0 \\ 0 \\ 0 \\ 0 \\ \frac{\psi^2}{4} \end{pmatrix}^{T}
,
$$

(A17)

$$
r_{\xi_{01}^{(2)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} 0 \\ 1-\psi^2 \\ \frac{\psi^2}{2} \\ 0 \\ 0 \\ 0 \\ \frac{\psi^2}{4} \\ 0 \\ 0 \\ \frac{\psi^2}{4} \end{pmatrix}^{T}
,\
r_{\xi_{01}^{(3)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} \frac{3\psi^3}{2}-\frac{5\psi^2}{2}+1 \\ 2\psi^2(1-\psi) \\ \frac{\psi^3}{2} \\ \frac{\psi^2-\psi^3}{4} \\ 0 \\ 0 \\ \frac{\psi^3}{4} \\ 0 \\ 0 \\ \frac{\psi^2}{4} \end{pmatrix}^{T}
,\
r_{\xi_{00}^{(2)}}^{(1)} =
\begin{array}{c}
\xi_{00}^{(4)} \\ \xi_{00}^{(3)} \\ \xi_{00}^{(2)} \\ \xi_{01}^{(3)} \\ \xi_{01}^{(2)} \\ \xi_{10}^{(3)} \\ \xi_{10}^{(2)} \\ \xi_{11}^{(2)} \\ \xi_{11}^{(1)} \\ \xi_\Delta
\end{array}
\begin{pmatrix} \frac{1}{4}-\frac{\psi^2}{4} \\ \frac{1}{2}-\frac{\psi^2}{2} \\ \frac{1}{4} \\ 0 \\ \frac{\psi^2}{8} \\ 0 \\ \frac{\psi^2}{8} \\ \frac{\psi^2}{32} \\ \frac{\psi^2}{32} \\ \frac{7\psi^2}{16} \end{pmatrix}^{T}
,
$$

(A18)

$$
r^{(1)}_{\xi^{(3)}_{00}} =
\begin{array}{c}
\xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_\Delta
\end{array}
\begin{pmatrix}
\psi^3 - \frac{3\psi^2}{2} + \frac{1}{2} \\
\frac{1}{2} - \frac{\psi^3}{2} \\
\frac{\psi^2}{2} - \frac{\psi^3}{4} \\
\frac{\psi^2}{8} - \frac{\psi^3}{8} \\
\frac{\psi^2}{8} \\
\frac{\psi^2}{8} - \frac{\psi^3}{8} \\
\frac{\psi^2}{8} \\
\frac{\psi^3}{32} \\
\frac{\psi^3}{32} \\
\frac{\psi^2}{2} - \frac{\psi^3}{16}
\end{pmatrix}^T
\quad \text{and} \quad
r^{(1)}_{\xi^{(4)}_{00}} =
\begin{array}{c}
\xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_\Delta
\end{array}
\begin{pmatrix}
(2\psi + 1)(\psi - 1)^2 \\
-2(\psi - 1)\psi^2 \\
\frac{\psi^4}{4} \\
\frac{(1-\psi)\psi^2}{4} \\
\frac{(2-\psi)\psi^3}{8} \\
\frac{(1-\psi)\psi^2}{4} \\
\frac{(2-\psi)\psi^3}{8} \\
\frac{\psi^4}{32} \\
\frac{\psi^4}{32} \\
\frac{\psi^2(8-\psi^2)}{16}
\end{pmatrix}^T .
\tag{A19}
$$

Consider now the matrices $\widetilde{A} := \lim_{N\to\infty} \widetilde{\Pi}_N = \widetilde{A}^{\mathrm{WF}}$ and $\widetilde{B}^{(\theta)}_N := N^\theta(\widetilde{\Pi}_N - \widetilde{A})$. We see from (A13) that

$$
\widetilde{P} := \lim_{m\to\infty} \widetilde{A}^m =
\begin{array}{c}
\\ \xi^{(4)}_{00} \\ \xi^{(3)}_{00} \\ \xi^{(2)}_{00} \\ \xi^{(3)}_{10} \\ \xi^{(2)}_{10} \\ \xi^{(3)}_{01} \\ \xi^{(2)}_{01} \\ \xi^{(2)}_{11} \\ \xi^{(1)}_{11} \\ \xi_\Delta
\end{array}
\begin{array}{c}
\begin{matrix} \mathcal{S} & \xi^{(4)}_{00} & \xi^{(3)}_{00} & \xi^{(2)}_{00} & \xi^{(3)}_{10} & \xi^{(2)}_{10} & \xi^{(3)}_{01} & \xi^{(2)}_{01} & \xi^{(2)}_{11} & \xi^{(1)}_{11} & \xi_\Delta \end{matrix} \\
\begin{bmatrix}
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{bmatrix}
\end{array} .
\tag{A20}
$$

In fact, $\widetilde{A}$ is the transition matrix of a Markov chain on $\mathcal{S} = \{\xi^{(4)}_{00}, \xi^{(3)}_{00}, \ldots, \xi_\Delta\}$ with two absorbing states $\xi^{(4)}_{00}$ and ξ_Δ; furthermore, $\xi^{(4)}_{00}$ is reached from every $\xi \in \mathcal{S} \setminus \{\xi_\Delta\}$. Denoting $\widetilde{B}^{(\theta)}_N := N(\widetilde{\Pi}_N - \widetilde{A})$ we can write $\widetilde{\Pi}_N = \widetilde{A} + \frac{1}{N}\widetilde{B}^{(\theta)}_N$, for $\alpha_N = \lambda/N^\theta$ and $\theta \in (0,1]$. Also let

$$
\widetilde{B}^{(\theta)} := \lim_{N\to\infty} \widetilde{B}^{(\theta)}_N =
\begin{cases}
\lambda(\widetilde{A}^{\mathrm{HR}} - \widetilde{A}^{\mathrm{WF}}), & \text{if } \theta \in (0,1) \\
\lambda(\widetilde{A}^{\mathrm{HR}} - \widetilde{A}^{\mathrm{WF}}) + \widetilde{B}^{\mathrm{WF}}, & \text{if } \theta = 1
\end{cases},
$$

and observe that the first nine rows of

$$
\widetilde{G}^{(\theta)} := \lim_{N\to\infty} \widetilde{P}\widetilde{B}^{(\theta)}_N \widetilde{P} = \widetilde{P}\widetilde{B}^{(\theta)}\widetilde{P},
\tag{A21}
$$

are identical and equal to

$$
\left(-1 - \lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16}), \ 0, \ 0, \ldots, \ 0, \ 1 + \lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16})\right),
$$

1104 when $\theta = 1$, and equal to

$$\left(-\lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16}),\ 0,\ 0, \ldots,\ 0,\ \lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16})\right),$$

when $\theta \in (0,1)$. The last row of $G^{(\theta)}$ is equal to $(0, 0, \ldots, 0)$ for all $\theta \in (0,1]$. Note that for
computations involving $\widetilde{G}^{(\theta)}$, we can effectively collapse the state space $\mathcal{S}$ into two points: ξ_Δ
and $\mathcal{S} \setminus \{\xi_\Delta\}$. Now we have all the required terms to invoke Möhle (1998a, Lemma 1) for the
proof of Lemma 2. We note that $\sup_N ||\widetilde{B}_N^{(\theta)}|| < \infty$, where for any matrix $K = (k_{ij})_{i,j}$, the
norm $||K|| := \max_i \sum_j |k_{ij}|$ is used. The calculations for Lemma 2 are available at https:
//github.com/DimitriDiam/Bursts-of-coalescence.git in the **2pairs_mix_Möhle.pdf** file.
The computations can be reproduced using the **2pairs_mix_Möhle.nb** Mathematica notebook,
version 13.1.0 (Wolfram Research, Inc., 2022).

Lemma 2. Let τ and τ' be the coalescent times of the two pairs of genes that are conditionally
independent given $\mathcal{A}^{(N,2)}$. Suppose the initial distribution $\vec{p}_0$ of $\widetilde{M}_0$ puts zero mass on state ξ_Δ.
Then for all $0 < t \le t'$, as $N \to \infty$,

$$\mathbb{P}^{(N)}(\tau > [tN^\theta],\ \tau' > [t'N^\theta]) \to \begin{cases} \exp\left\{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16}) - \lambda\frac{\psi^2}{4}(t' - t)\right\}, & \text{if } \theta \in (0,1), \\ \exp\left\{-\left(1 + \lambda(\frac{\psi^2}{2} - \frac{\psi^4}{16})\right)t - \left(\frac{1}{2} + \lambda\frac{\psi^2}{4}\right)(t' - t)\right\}, & \text{if } \theta = 1. \end{cases}$$

1117
$$(A22)$$

1118 **Proof:** We first establish the result for the case $t = t'$, and then strengthen it to the general case
1119 $t \le t'$ by using the Markov property.

1120 Let $\vec{p}_0 = (0, 0, 1, 0, \cdots, 0)$ be the 1×13 row vector representing the distribution of $\widetilde{M}_0$, which
1121 corresponds to the initial distribution of the Markov chain. Then for each $N \in \mathbb{N}$ and each
1122 generation $g \in \mathbb{Z}_{\ge 0}$,

$$\mathbb{P}^{(N)}(\tau > g,\ \tau' > g) = \vec{p}_0\ \widetilde{\Pi}_N^g\ (1, 1, 1, 1, 1, 1, 1, 1, 1, 0)^T, \tag{A23}$$

1125 where $\widetilde{\Pi}_N$ was defined in (A11). Setting $g = [tN^\theta]$ in (A23), an application of Möhle (1998a,
1126 Lemma 1) gives

$$\lim_{N \to \infty} \vec{p}_0\ \widetilde{\Pi}_N^{[tN^\theta]}\ (1, 1, 1, 1, 1, 1, 1, 1, 1, 0)^T = \vec{p}_0\ \widetilde{P} e^{t\widetilde{G}}\ (1, 1, 1, 1, 1, 1, 1, 1, 1, 0)^T. \tag{A24}$$

1129 Substituting $\widetilde{P}$ from (A20) and $\widetilde{G}^{(\theta)}$ from (A21), the right hand side of (A24) is equal to

$$\vec{p}_0\ \widetilde{P} e^{t\widetilde{G}}\ (1, 1, 1, 1, 1, 1, 1, 1, 1, 0)^T \tag{A25}$$

$$= \begin{cases} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta \in (0,1), \\ \\ e^{-t} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta = 1, \end{cases} \tag{A26}$$

1133 for any $\vec{p}_0$ in the set $\mathcal{I}' := \{(x_1, \cdots, x_{13}) \in [0,1]^{13} : \sum_{i=1}^{12} x_i = 1,\ x_{13} = 0\}$. We have shown that

$$\lim_{N \to \infty} \mathbb{P}^{(N)}(\tau > [tN^\theta],\ \tau' > [tN^\theta]) = \begin{cases} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta \in (0,1), \\ \\ e^{-t} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta = 1. \end{cases} \tag{A27}$$

For each $N \in \mathbb{N}$ and generations $0 \le g \le g'$, by the Markov property,

$$\mathbb{P}^{(N)}(\tau > g, \, \tau' > g') = \mathbb{E}^{(N)}_{\widetilde{M}_0}[1_{\{\tau>g\}} 1_{\{\tau'>g'\}}] = \mathbb{E}^{(N)}_{\widetilde{M}_0}\left[1_{\{\tau>g\}} 1_{\{\tau'>g\}} \, \mathbb{P}^{(N)}_{\widetilde{M}_g}(\tau > g' - g)\right]. \quad (A28)$$

Set $g = [tN^\theta]$ and $g' = [t'N^\theta]$. Note that $\widetilde{M}_g \in \mathcal{I} := \{(c, 1-c, 0) \in [0,1]^3 : c \in [0,1]\}$ when $\tau > g$. Hence by Remark 1 and Lemma 1,

$$\lim_{N\to\infty} \mathbb{P}^{(N)}_{\widetilde{M}_g}(\tau > g' - g) = \begin{cases} e^{-\lambda \frac{\psi^2}{4}(t'-t)} & \text{when } \theta \in (0,1) \\[2ex] e^{-\left(\frac{1}{2}+\lambda\frac{\psi^2}{4}\right)(t'-t)} & \text{when } \theta = 1. \end{cases}$$

Putting this into (A28) and using (A27), we obtain the desired (A22).

$\square$

A.2   Supporting Lemmas

The second step in the proof of Theorem 1 was the computation of (24) for $\theta = 1$ and of (36) for $\theta \in (0,1)$. An important step in those calculations was to establish (32). Lemma 3 below establishes (32). Similarly to (16), for each $g \in \mathbb{Z}_{\ge 0}$ we let $F_N(g) := \mathbb{P}^{(N)}_{\mathcal{A}^{(N,2)}}(\tau > g)$ and $G_N(g) := \mathbb{P}^{(N)}_{H_N}(\tau > g)$.

Lemma 3. Recall the definition of Π_N^{mid} in (33). For all $g \in \mathbb{N}$, $\psi \in (0,1]$ and $N \ge 2$ and initial distribution $\vec{v}_0$ (represented as a 1×3 row vector),

$$\mathbb{E}^{(N)}\left[G_N(g)\left(1 - \frac{\psi^2}{4}\right)^{H_N(g)}\right] = \vec{v}_0 \left(\Pi_N^{\text{mid}}\right)^g (1,1,0)^T.$$

Proof: By the law of total expectation,

$$\mathbb{E}^{(N)}\left[G_N(g)\left(1 - \frac{\psi^2}{4}\right)^{H_N(g)}\right] = \mathbb{E}^{(N)}\left[\mathbb{P}^{(N)}_{H_N}(\tau > g)\left(1 - \frac{\psi^2}{4}\right)^{H_N(g)}\right]$$

$$= \sum_{k=0}^{g} \mathbb{P}^{(N)}(\tau > g \mid H_N(g) = k)\left(1 - \frac{\psi^2}{4}\right)^k \mathbb{P}^{(N)}(H_N(g) = k). \quad (A29)$$

Given g generations, k of which being HR generations, the total number of arrangements of those events in the pedigree is equal to C_k^g. Note that the arrangement is independent of the number of such events. Denote each such arrangement by $\vec{t}$ and by W_k the space of all possible arrangements of k HR generations within g generations. Let $\Pi_N^{\vec{t}}$ be the product of g terms, each term being either Π_N^{WF} or Π_N^{HR}. The order of the terms (from left to right) is determined by the vector $\vec{t}$ so that whenever a WF generation is taking place the corresponding factor in the product is Π_N^{WF} and Π_N^{HR} otherwise. Hence, the conditional probability in (A29) is equal to

$$\mathbb{P}^{(N)}(\tau > g \mid H_N(g) = k) = \frac{1}{\binom{g}{k}} \sum_{\vec{t} \in W_k} \mathbb{P}^{(N)}\left(\tau > g \mid \vec{t}\right) \quad (A30)$$

$$= \frac{1}{\binom{g}{k}} \sum_{\vec{t} \in W_k} \vec{v}_0 \, \Pi_N^{\vec{t}} \, (1,1,0)^T, \quad (A31)$$

where (A30) is true by the conditional law of total probability and (A31) by the definition of conditional probability. Plugging (A31) back into (A29), since $H_N(g) \sim \mathrm{Binom}(g, \alpha_N)$, it follows

$$\mathbb{E}^{(N)} \left[G_N(g) \left(1 - \frac{\psi^2}{4} \right)^{H_N(g)} \right]$$

$$= \vec{v}_0 \sum_{k=0}^{g} \sum_{\vec{t} \in W_k} \Pi_N^{\vec{t}} \left((1 - \frac{\psi^2}{4}) \alpha_N \right)^k (1 - \alpha_N)^{g-k} (1,1,0)^T$$

$$= \vec{v}_0 \left(\alpha_N (1 - \frac{\psi^2}{4}) \Pi_N^{\mathrm{HR}} + (1 - \alpha_N) \Pi_N^{\mathrm{WF}} \right)^g (1,1,0)^T$$

$$= \vec{v}_0 \left(\Pi_N^{\mathrm{mid}} \right)^g (1,1,0)^T, \tag{A32}$$

where the second last equality follows from distributing k terms of α_N and $g - k$ terms of $1 - \alpha_N$ into the product $\Pi_N^{\vec{t}}$.

$\square$

Finally, we compute the right hand of (32). The calculations for Lemma 4 are available at https://github.com/DimitriDiam/Bursts-of-coalescence.git in the **Mixed_Term_Moment.pdf** file and can be reproduced in the Mathematica notebook **Mixed_Term_Moment.nb**, where Mathematica version 13.1.0 (Wolfram Research, Inc., 2022) was used.

Lemma 4. Suppose the distribution $\vec{v}_0$ of the initial state M_0 puts zero mass at the coalescent state ξ_2 (recall 3-states Markov chain M in the proof of Lemma 1). For each $t \in (0, \infty)$, it holds that

$$\lim_{N \to \infty} \mathbb{E}^{(N)} \left[G_N([tN^\theta]) \left(1 - \frac{\psi^2}{4} \right)^{H_N([tN^\theta])} \right] = \begin{cases} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta \in (0,1), \\ \\ e^{-t/2} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta = 1. \end{cases}$$

Proof: Let Π_N^{mid} as in (33), by Lemma 3 for $g = [tN^\theta]$ it follows that

$$\mathbb{E}^{(N)} \left[G_N([tN^\theta]) \left(1 - \frac{\psi^2}{4} \right)^{H_N([tN^\theta])} \right] = \vec{v}_0 \left(\Pi_N^{\mathrm{mid}} \right)^{tN^\theta} (1,1,0)^T. \tag{A33}$$

Notice that Π_N^{mid} is not a stochastic matrix, consider instead

$$\widetilde{\Pi}_N^{\mathrm{mid}} := \left(1 - \frac{\lambda \psi^2}{4N^\theta} \right)^{-1} \Pi_N^{\mathrm{mid}},$$

so that $\widetilde{\Pi}_N^{\mathrm{mid}}$ is a stochastic matrix. For any $\theta \in (0,1]$, let

$$A := \lim_{N \to \infty} \widetilde{\Pi}_N^{\mathrm{mid}} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \text{ and}$$

$$B^{(\theta)} := \lim_{N \to \infty} N^\theta (\widetilde{\Pi}_N^{\mathrm{mid}} - A) \tag{A34}$$

$$= \begin{cases} \begin{pmatrix} \frac{\lambda\psi^4}{8} - \frac{\lambda\psi^2}{2} & \frac{\lambda\psi^2}{4} - \frac{\lambda\psi^4}{16} & \frac{\lambda\psi^2}{4} - \frac{\lambda\psi^4}{16} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \text{if } 0 < \theta < 1, \\[2em] \begin{pmatrix} \frac{\lambda\psi^4}{8} - \frac{\lambda\psi^2}{2} - 1 & -\frac{\lambda\psi^4}{16} + \frac{\lambda\psi^2}{4} + \frac{1}{2} & -\frac{\lambda\psi^4}{16} + \frac{\lambda\psi^2}{4} + \frac{1}{2} \\ -1 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 \end{pmatrix}, & \text{if } \theta = 1. \end{cases} \quad (A35)$$

Furthermore consider the matrix $P := \lim_{k \to \infty} A^k$, then $P = A$ and let $G^{(\theta)} := PB^{(\theta)}P$. Then for $\theta \in (0,1)$, the first two rows of $G^{(\theta)}$ are identical to

$$\left(\frac{1}{16}\lambda\psi^2 \left(\psi^2 - 4 \right) , \, 0 , \, -\frac{1}{16}\lambda\psi^2 \left(\psi^2 - 4 \right) \right),$$

and the last one is equal to $(0, 0, 0)$. For $\theta = 1$, the first two rows of $G^{(1)}$ are identical to

$$\left(\frac{1}{8}\lambda \left(\psi^2 - 4 \right) \psi^2 + \frac{1}{16} \left(8 - \lambda\psi^2 \left(\psi^2 - 4 \right) \right) - 1 , \, 0 , \, \frac{1}{16} \left(8 - \lambda\psi^2 \left(\psi^2 - 4 \right) \right) \right),$$

and the last one is equal to $(0, 0, 0)$. Writing $\widetilde{\Pi}_N^{\mathrm{mid}}$ as

$$\widetilde{\Pi}_N^{\mathrm{mid}} = A + \frac{B^{(\theta)}}{N^\theta} + o(\frac{1}{N^\theta}), \quad (A36)$$

we can apply Möhle's Lemma (Möhle, 1998a, Lemma 1) which gives

$$\lim_{N \to \infty} \vec{v}_0 \left(\widetilde{\Pi}_N^{\mathrm{mid}} \right)^{tN^\theta} (1,1,0)^T = \begin{cases} e^{-\lambda t(\frac{\psi^2}{4} - \frac{\psi^4}{16})}, & \text{for } \theta \in (0,1), \\[1em] e^{-t/2} e^{-\lambda t(\frac{\psi^2}{4} - \frac{\psi^4}{16})}, & \text{for } \theta = 1. \end{cases} \quad (A37)$$

Hence (A33), through the use of (A37), is equal to

$$\lim_{N \to \infty} \vec{v}_0 \left(\Pi_N^{\mathrm{mid}} \right)^{tN^\theta} (1,1,0)^T = \lim_{N \to \infty} \vec{v}_0 \left(\left(1 - \frac{\lambda\psi^2}{4N^\theta} \right) \widetilde{\Pi}_N^{\mathrm{mid}} \right)^{tN^\theta} (1,1,0)^T$$

$$= e^{-\lambda t \frac{\psi^2}{4}} \lim_{N \to \infty} \vec{v}_0 \left(\widetilde{\Pi}_N^{\mathrm{mid}} \right)^{tN^\theta} (1,1,0)^T$$

$$= \begin{cases} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta \in (0,1), \\[1em] e^{-t/2} e^{-\lambda t(\frac{\psi^2}{2} - \frac{\psi^4}{16})}, & \text{if } \theta = 1. \end{cases}$$

The last equality holds because $\vec{v}_0 = (x, 1 - x, 0)$ for some $x \in [0,1]$. $\qquad \square$

A.3   List of Patterns and Mendelian randomness tables

A.3.1   List of patterns

For the Tables A1-A3, let $x, y, z, w \in [N]$ represent distinct individuals. Circles represent the parent(s) of these individuals. For any individual, two arrows pointing to the same circle indicate

that the individual results from selfing. If two individuals each have one arrow pointing to the same circle, it means that the individuals have exactly one parent in common. The third column in each of the Tables A1-A3 contains the probabilities of the patterns in a Wright-Fisher generation, while the last column represents a generation with a highly reproductive pair. We only keep track of terms of order $O(1)$ and $O(N^{-1})$.

For example, the probabilities of the patterns A_{11} and A_{12} are explained as follows: In the offspring generation, an individual originates from the highly reproductive pair with probability ψ, signifying two distinct parents. Conversely, the same individual is *not* a result of the highly reproductive pair with a probability of $1 - \psi$. In this scenario, the individual comes from selfing with a probability of $1/N$, and from two distinct parents with probability $1 - 1/N$. Finally, the probability of pattern A_{22} is explained as follows: Each individual in the offspring generation originates from the highly reproductive pair with probability ψ. By independence, both individuals are offspring of the highly reproductive couple with probability ψ^2. Note that two individuals can share both their parents in various ways. However, in all other scenarios, the probability of such an event is of the order $O(N^{-2})$.

Pattern	Name	Wright-Fisher	High Reproduction
	A_{11}	$\frac{1}{N}$	$\frac{1-\psi}{N}$
	A_{12}	$1 - \frac{1}{N}$	$\psi + (1-\psi)\left(1 - \frac{1}{N}\right)$
	A_{22}	$O(N^{-2})$	$\psi^2 + O(N^{-2})$
	$A_{23,s}(x,y)$	$\frac{(N-1)(N-2)}{N^3}$	$\psi(1-\psi)\frac{1}{N} + (1-\psi)^2\frac{(N-1)(N-2)}{N^3}$
	A_{23}	$\frac{2(N-1)(N-2)}{N^3}$	$2\psi(1-\psi)\frac{2}{N}(1 - \frac{2}{N}) + (1-\psi)^2\frac{2(N-1)(N-2)}{N^3}$
	A_{24}	$\frac{(N-1)(N-2)(N-3)}{N^3}$	$2\psi(1-\psi)\frac{(N-2)(N-3)}{N^2} + (1-\psi)^2\frac{(N-1)(N-2)(N-3)}{N^3}$

Table A1: List of patterns when the sampled genes are distributed among one or two individuals.

Pattern	Name	Wright-Fisher	High Reproduction
	A_{32}	$O(N^{-2})$	$\psi^3 + O(N^{-2})$
	$A_{33,s}(z)$	$O(N^{-2})$	$\psi^2(1-\psi)(1-\frac{2}{N})\frac{1}{N}$
	$A_{33,2}(z)$	$O(N^{-2})$	$\psi^2(1-\psi)(1-\frac{2}{N})\frac{2}{N}$
	$A_{33}(z)$	$O(N^{-2})$	$\psi^2(1-\psi)(1-\frac{2}{N})(1-\frac{3}{N})$
	$A_{35,s}(x)$	$\frac{(N-1)\cdots(N-4)}{N^5}$	$\psi(1-\psi)^2\frac{(N-2)\cdots(N-4)}{N^4} + (1-\psi)^3\frac{(N-1)\cdots(N-4)}{N^5}$
	$A_{35}(x,y)$	$\frac{2(N-1)\cdots(N-4)}{N^5}$	$\psi(1-\psi)^2\frac{2(N-2)\cdots(N-4)}{N^4} + (1-\psi)^3\frac{2(N-1)\cdots(N-4)}{N^5}$
	A_{36}	$\frac{(N-1)\cdots(N-5)}{N^5}$	$3\psi(1-\psi)^2\frac{(N-2)\cdots(N-5)}{N^4} + (1-\psi)^3\frac{(N-1)\cdots(N-5)}{N^5}$

Table A2: List of patterns when the sampled genes are distributed among three individuals.

Pattern	Name	Wright-Fisher	High Reproduction
	A_{42}	$O(N^{-2})$	$\psi^4 + O(N^{-2})$
	$A_{43}(w)$	$O(N^{-2})$	$\psi^3(1-\psi)\frac{2}{N}(1-\frac{2}{N})$
	$A_{43,s}(w)$	$O(N^{-2})$	$\psi^3(1-\psi)(1-\frac{2}{N})\frac{1}{N}$
	$A_{44}(w)$	$O(N^{-2})$	$\psi^3(1-\psi)\frac{(N-2)(N-3)}{N^2}$
	$A_{45,s}(z,w)$	$O(N^{-2})$	$\psi^2(1-\psi)^2\frac{(N-2)\cdots(N-4)}{N^4}$
	$A_{45}(z,w)$	$O(N^{-2})$	$\psi^2(1-\psi)^2\frac{2(N-2)(N-3)(N-4)}{N^4}$
	$A_{46}(z,w)$	$O(N^{-2})$	$\psi^2(1-\psi)^2\frac{(N-2)(N-3)(N-4)(N-5)}{N^4}$
	$A_{47,s}(x)$	$\frac{(N-1)\cdots(N-6)}{N^7}$	$\psi(1-\psi)^3\frac{(N-2)\cdots(N-6)}{N^6} + (1-\psi)^4\frac{(N-1)\cdots(N-6)}{N^7}$
	$A_{47}(x,y)$	$\frac{2(N-1)\cdots(N-6)}{N^7}$	$\psi(1-\psi)^3\frac{2(N-2)\cdots(N-6)}{N^6} + (1-\psi)^4\frac{4(N-1)\cdots(N-6)}{N^7}$
	A_{48}	$\frac{(N-1)\cdots(N-7)}{N^7}$	$4\psi(1-\psi)^3\frac{(N-2)\cdots(N-7)}{N^6} + (1-\psi)^4\frac{(N-1)\cdots(N-7)}{N^7}$

Table A3: List of patterns when the sampled genes are distributed among four individuals.

A.3.2   Mendelian randomness tables

Tables A4-A10 below, contain the transition probabilities for every pair of states $(\xi, \eta) \in \mathcal{S}^2$, conditional on the patterns from Tables A1-A3. The derivation of the entries of Tables A4-A10 follow similarly to the entries of (A10) in subsection A.1.3. For example, in Table A10, the last row of the column labeled A_{47} is explained as follows. Lineages in distinct individuals sharing exactly one parent have descended from different parents with probability $3/4$. On the other hand, state $\xi_{00}^{(4)}$ consists of two distinct individuals, hence the pattern A_{47} is used in $\binom{4}{2} = 6$ ways. Finally we note that in Tables A4-A10, the fractional quantities stand for events determined purely by Mendelian randomness (conditional on each pattern) whereas the integer quantities refer to combinatorial terms corresponding to the appropriate use of the patterns.

$(\xi_{11}^{(1)}, \eta)$	A_{11}	A_{12}
$(\xi_{11}^{(1)}, \xi_{11}^{(1)})$	$\frac{1}{4}$	0
$(\xi_{11}^{(1)}, \xi_{11}^{(2)})$	0	0
$(\xi_{11}^{(1)}, \xi_{10}^{(2)})$	0	0
$(\xi_{11}^{(1)}, \xi_{10}^{(3)})$	0	0
$(\xi_{11}^{(1)}, \xi_{01}^{(2)})$	0	0
$(\xi_{11}^{(1)}, \xi_{01}^{(3)})$	0	0
$(\xi_{11}^{(1)}, \xi_{00}^{(2)})$	0	$\frac{1}{1}$
$(\xi_{11}^{(1)}, \xi_{00}^{(3)})$	0	0
$(\xi_{11}^{(1)}, \xi_{00}^{(4)})$	0	0

Table A4: Conditional probabilities given the pattern when the initial state is $\xi_{11}^{(1)}$.

$(\xi_{11}^{(2)}, \eta)$	A_{22}	$A_{23,s}$	A_{23}	A_{24}
$(\xi_{11}^{(2)}, \xi_{11}^{(1)})$	0	0	0	0
$(\xi_{11}^{(2)}, \xi_{11}^{(2)})$	0	0	0	0
$(\xi_{11}^{(2)}, \xi_{10}^{(2)})$	0	0	0	0
$(\xi_{11}^{(2)}, \xi_{10}^{(3)})$	0	$\frac{1}{2}$	0	0
$(\xi_{11}^{(2)}, \xi_{01}^{(2)})$	0	0	0	0
$(\xi_{11}^{(2)}, \xi_{01}^{(3)})$	0	$\frac{1}{2}$	0	0
$(\xi_{11}^{(2)}, \xi_{00}^{(2)})$	$\frac{1}{1}$	0	0	0
$(\xi_{11}^{(2)}, \xi_{00}^{(3)})$	0	0	$\frac{1}{1}$	0
$(\xi_{11}^{(2)}, \xi_{00}^{(4)})$	0	0	0	$\frac{1}{1}$

Table A5: Conditional probabilities given the pattern when the initial state is $\xi_{11}^{(2)}$.

$(\xi_{10}^{(2)}, \eta)$	A_{22}	$A_{23,s}$	A_{23}	A_{24}
$(\xi_{10}^{(2)}, \xi_{11}^{(1)})$	0	0	0	0
$(\xi_{10}^{(2)}, \xi_{11}^{(2)})$	0	0	0	0
$(\xi_{10}^{(2)}, \xi_{10}^{(2)})$	0	$\frac{1}{2}$	0	0
$(\xi_{10}^{(2)}, \xi_{10}^{(3)})$	0	0	0	0
$(\xi_{10}^{(2)}, \xi_{01}^{(2)})$	$\frac{1}{4}$	0	$\frac{1}{4}$	0
$(\xi_{10}^{(2)}, \xi_{01}^{(3)})$	0	0	0	0
$(\xi_{10}^{(2)}, \xi_{00}^{(2)})$	$\frac{1}{2}$	0	$\frac{1}{4}$	0
$(\xi_{10}^{(2)}, \xi_{00}^{(3)})$	0	$1 \cdot \frac{1}{1}$	$\frac{1}{2}$	$\frac{1}{1}$
$(\xi_{10}^{(2)}, \xi_{00}^{(4)})$	0	0	0	0

Table A6: Conditional probabilities given the pattern when the initial state is $\xi_{10}^{(2)}$.

$(\xi_{00}^{(2)}, \eta)$	A_{22}	$A_{23,s}$	A_{23}	A_{24}
$(\xi_{00}^{(2)}, \xi_{11}^{(1)})$	$\frac{1}{32}$	0	$\frac{1}{64}$	0
$(\xi_{00}^{(2)}, \xi_{11}^{(2)})$	$\frac{1}{32}$	0	0	0
$(\xi_{00}^{(2)}, \xi_{10}^{(2)})$	$\frac{1}{8}$	0	$\frac{1}{16}$	0
$(\xi_{00}^{(2)}, \xi_{10}^{(3)})$	0	0	$\frac{1}{32}$	0
$(\xi_{00}^{(2)}, \xi_{01}^{(2)})$	$\frac{1}{8}$	0	$\frac{1}{16}$	0
$(\xi_{00}^{(2)}, \xi_{01}^{(3)})$	0	0	$\frac{1}{32}$	0
$(\xi_{00}^{(2)}, \xi_{00}^{(2)})$	$\frac{1}{4}$	$2 \cdot \frac{1}{2}$	$1 \cdot \frac{3}{16}$	$\frac{1}{4}$
$(\xi_{00}^{(2)}, \xi_{00}^{(3)})$	0	$2 \cdot \frac{1}{2}$	$1 \cdot \frac{6}{16}$	$\frac{1}{2}$
$(\xi_{00}^{(2)}, \xi_{00}^{(4)})$	0	0	0	$\frac{1}{4}$

Table A7: Conditional probabilities given the pattern when the initial state is $\xi_{00}^{(2)}$.

$(\xi_{10}^{(3)}, \eta)$	A_{32}	$A_{33,s}$	$A_{33,2}$	A_{33}	$A_{35,s}$	A_{35}	A_{36}
$(\xi_{10}^{(3)}, \xi_{11}^{(1)})$	0	0	0	0	0	0	0
$(\xi_{10}^{(3)}, \xi_{11}^{(2)})$	0	$\frac14$	0	0	0	0	0
$(\xi_{10}^{(3)}, \xi_{10}^{(2)})$	0	0	0	0	0	0	0
$(\xi_{10}^{(3)}, \xi_{10}^{(3)})$	0	$\frac14$	0	0	$1\cdot\frac12$	0	0
$(\xi_{10}^{(3)}, \xi_{01}^{(2)})$	$\frac14$	0	$1\cdot\frac18$	0	0	0	0
$(\xi_{10}^{(3)}, \xi_{01}^{(3)})$	0	0	$3\cdot\frac18$	$1\cdot\frac14$	0	$1\cdot\frac18$	0
$(\xi_{10}^{(3)}, \xi_{00}^{(2)})$	$\frac12$	0	$2\cdot\frac14$	0	0	0	0
$(\xi_{10}^{(3)}, \xi_{00}^{(3)})$	0	$2\cdot1$	$3\cdot\frac12$	$2\cdot1$	0	$2\cdot\frac12$	0
$(\xi_{10}^{(3)}, \xi_{00}^{(4)})$	0	0	0	$1\cdot\frac12$	$2\cdot\frac11$	$1\cdot\frac34+2\cdot\frac12$	$\frac11$

Table A8: Conditional probabilities given the pattern when the initial state is $\xi_{10}^{(3)}$.

$(\xi_{00}^{(3)}, \eta)$	A_{32}	$A_{33,s}$	$A_{33,2}$	A_{33}	$A_{35,s}$	A_{35}	A_{36}
$(\xi_{00}^{(3)}, \xi_{11}^{(1)})$	$1\cdot\frac{1}{32}$	0	$3\cdot\frac{1}{64}$	0	0	0	0
$(\xi_{00}^{(3)}, \xi_{11}^{(2)})$	$1\cdot\frac{1}{32}$	0	$2\cdot\frac{1}{64}$	0	0	0	0
$(\xi_{00}^{(3)}, \xi_{10}^{(2)})$	$1\cdot\frac18$	$1\cdot\frac18$	$2\cdot\frac18+1\cdot\frac{1}{16}$	$1\cdot\frac18$	0	$1\cdot\frac18$	0
$(\xi_{00}^{(3)}, \xi_{10}^{(3)})$	0	$1\cdot\frac18$	$1\cdot\frac{1}{32}+1\cdot\frac{1}{16}$	$1\cdot\frac18$	0	$1\cdot\frac18$	0
$(\xi_{00}^{(3)}, \xi_{01}^{(2)})$	$1\cdot\frac18$	$1\cdot\frac18$	$2\cdot\frac{1}{16}+1\cdot\frac18$	$1\cdot\frac18$	0	$1\cdot\frac18$	0
$(\xi_{00}^{(3)}, \xi_{01}^{(3)})$	0	$1\cdot\frac18$	$1\cdot\frac{1}{32}+1\cdot\frac{1}{16}$	$1\cdot\frac18$	0	$1\cdot\frac18$	0
$(\xi_{00}^{(3)}, \xi_{00}^{(2)})$	$1\cdot\frac14$	$1\cdot\frac12$	$1\cdot\frac38+2\cdot\frac18$	$1\cdot\frac12$	0	0	0
$(\xi_{00}^{(3)}, \xi_{00}^{(3)})$	0	$3\cdot\frac12$	$1\cdot\frac38+2\cdot\frac14$	$3\cdot\frac12$	$1\cdot\frac11+2\cdot\frac12$	$1\cdot\frac12+2\cdot\frac12$	$1\cdot\frac12$
$(\xi_{00}^{(3)}, \xi_{00}^{(4)})$	0	0	0	0	$2\cdot\frac12$	$1\cdot\frac38+2\cdot\frac14$	$1\cdot\frac12$

Table A9: Conditional probabilities given the pattern when the initial state is $\xi_{00}^{(3)}$.

$(\xi_{00}^{(4)}, \eta)$	A_{42}	A_{43}	$A_{43,s}$	A_{44}	A_{45}	$A_{45,s}$	A_{46}	$A_{47,s}$	A_{47}	A_{48}
$(\xi_{00}^{(4)}, \xi_{11}^{(1)})$	$1\cdot\frac{1}{32}$	$2\cdot\frac{1}{64}$	0	0	0	0	0	0	0	0
$(\xi_{00}^{(4)}, \xi_{11}^{(2)})$	$1\cdot\frac{1}{32}$	$2\cdot\frac{1}{64}$	0	0	$2\cdot\frac{1}{16}$	0	0	0	0	0
$(\xi_{00}^{(4)}, \xi_{10}^{(2)})$	$1\cdot\frac{1}{8}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	0	0	0	0	0	0
$(\xi_{00}^{(4)}, \xi_{10}^{(3)})$	0	$2\cdot\frac{1}{32}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	$1\cdot\frac{1}{16}+1\cdot\frac{3}{16}$	$1\cdot\frac{1}{4}$	$1\cdot\frac{1}{4}$	0	$1\cdot\frac{1}{8}$	0
$(\xi_{00}^{(4)}, \xi_{01}^{(2)})$	$1\cdot\frac{1}{8}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	0	0	0	0	0	0
$(\xi_{00}^{(4)}, \xi_{01}^{(3)})$	0	$2\cdot\frac{1}{32}$	$2\cdot\frac{1}{16}$	$2\cdot\frac{1}{16}$	$1\cdot\frac{1}{16}+1\cdot\frac{3}{16}$	$1\cdot\frac{1}{4}$	$1\cdot\frac{1}{4}$	0	$1\cdot\frac{1}{8}$	0
$(\xi_{00}^{(4)}, \xi_{00}^{(2)})$	$1\cdot\frac{1}{4}$	$2\cdot\frac{1}{16}$	0	0	$1\cdot\frac{1}{8}$	0	0	0	0	0
$(\xi_{00}^{(4)}, \xi_{00}^{(3)})$	0	$2\cdot\frac{1}{16}$	$4\cdot\frac{4}{8}$	$4\cdot\frac{1}{2}$	$1\cdot\frac{8}{16}$	$2\cdot\frac{1}{2}$	$4\cdot\frac{1}{2}$	0	$4\cdot\frac{1}{4}$	0
$(\xi_{00}^{(4)}, \xi_{00}^{(4)})$	0	0	0	0	$6\cdot\frac{3}{8}$	$6\cdot\frac{1}{2}$	$6\cdot\frac{1}{2}$	$4\cdot\frac{1}{1}$	$6\cdot\frac{3}{4}$	$1\cdot\frac{1}{1}$

Table A10: Conditional probabilities given the pattern when the initial state is $\xi_{00}^{(4)}$.

A.4 Strengthening the convergence in Theorem 1

Remark 4. *The arguments used in the proof of Theorem 1 in fact allow to conclude the (stronger) statement that*

$$\left(F_N(t, \mathcal{A}^{(N,2)})\right)_{t \geq 0} \longrightarrow \begin{cases} \left(\left(1 - \frac{\psi^2}{4}\right)^{Y(t-)}\right)_{t \geq 0}, & when \ \theta \in (0,1), \\ \left(e^{-t/2}\left(1 - \frac{\psi^2}{4}\right)^{Y(t-)}\right)_{t \geq 0}, & when \ \theta = 1, \end{cases} \tag{A38}$$

in finite-dimensional distributions.

Proof. We will only consider the case $\theta = 1$ here, the case $\theta < 1$ is similar.

Fix $k \in \mathbb{N}$ and $0 < t_1 < t_2 < \cdots t_k$. Using (21) for $t_1, \ldots, t_k$, we have

$$\lim_{N \to \infty} \mathbb{E}^{(N)}\left[\sum_{j=1}^{k} \left|F_N(t_j, \mathcal{A}^{(N,2)}) - e^{-t_j/2}\left(1 - \frac{\psi^2}{4}\right)^{Q_N([t_j N])}\right|^2\right] = 0, \tag{A39}$$

which implies as in the proof of Theorem 1 that

$$\left(F_N(t_1, \mathcal{A}^{(N,2)}), \ldots, F_N(t_k, \mathcal{A}^{(N,2)})\right)$$

$$\longrightarrow \left(e^{-t_1/2}\left(1 - \frac{\psi^2}{4}\right)^{Y(t_1)}, \ldots, e^{-t_k/2}\left(1 - \frac{\psi^2}{4}\right)^{Y(t_k)}\right)$$

$$\stackrel{d}{=} \left(e^{-t_1/2}\left(1 - \frac{\psi^2}{4}\right)^{Y(t_1-)}, \ldots, e^{-t_k/2}\left(1 - \frac{\psi^2}{4}\right)^{Y(t_k-)}\right)$$

in distribution. This shows convergence of finite dimensional distributions. $\square$

Further we conjecture that we have weak convergence in the space of left-continuous functions with right limits, equipped with the Skorohod topology, but we do not pursue this here.

References

Adams A. M. and Hudson R. R. Maximum-likelihood estimation of demographic parameters using the frequency spectrum of unlinked single-nucleotide polymorphisms. *Genetics*, 168(3):1699–1712, 2004. doi: 10.1534/genetics.104.030171.

Aguillon S. M., Fitzpatrick J. W., Bowman R., Schoech S. J., Clark A. G., Coop G., and Chen N. Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLOS Genetics*, 13(8):1–27, 2017. doi: 10.1371/journal.pgen.1006911.

Albers P. K. and McVean G. Dating genomic variants and shared ancestry in population-scale sequencing data. *PLOS Biology*, 18(1):1–26, 2020. doi: 10.1371/journal.pbio.3000586.

Anderson-Trocmé L., Nelson D., Zabad S., Diaz-Papkovich A., Kryukov I., Baya N., Touvier M., Jeffery B., Dina C., Vézina H., Kelleher J., and Gravel S. On the genes, genealogies, and geographies of Quebec. *Science*, 380(6647):849–855, 2023. doi: 10.1126/science.add5300.

Árnason E., Koskela J., Halldórsdóttir K., and Eldon B. Sweepstakes reproductive success via pervasive and recurrent selective sweeps. *eLife*, 12:e80781, 2023. doi: 10.7554/eLife.80781.

Avise J. C. Gene trees and organismal histories: a phylogenetic approach to population biology. *Evolution*, 43(6):1192–1208, 1989. doi: 10.1111/j.1558-5646.1989.tb02568.x.

Avise J. C. *Phylogeography: The History and Formation of Species*. Harvard University Press, Cambridge, Massachusetts, 2000.

Avise J. C. and Wollenberg K. Phylogenetics and the origin of species. *Proceedings of the National Academy of Sciences USA*, 94(15):7748–7755, 1997. doi: 10.1073/pnas.94.15.7748.

Avise J. C., Arnold J., Ball R. M., Bermingham E., Lamb T., Neigel J. E., Reeb C. A., and Saunders N. C. Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. *Annual Review of Ecology and Systematics*, 18(1):489–522, 1987. doi: 10.1146/annurev.es.18.110187.002421.

Ball R. M., Neigel J. E., and Avise J. C. Gene genealogies within the organismal pedigrees of random-mating populations. *Evolution*, 44(2):360–370, 1990. doi: 10.1111/j.1558-5646.1990.tb05205.x.

Barton N. H., Etheridge A. M., and Véber. The infinitesimal model: Definition, derivation, and implications. *Theoretical Population Biology*, 118:50–73, 2017. doi: 10.1016/j.tpb.2017.06.001.

Barton N. H. and Etheridge A. M. The relation between reproductive value and genetic contribution. *Genetics*, 188(4):953–973, 2011. doi: 10.1534/genetics.111.127555.

Birkner M., Blath J., Möhle M., Steinrücken, and Tams J. A modified lookdown construction for the Xi-Fleming-Viot process with mutation and populations with recurrent bottlenecks. *ALEA Latin American Journal of Probability and Mathematical Statistics*, 6:35–61, 2009. URL https://alea.impa.br/articles/v6/06-02.pdf.

Birkner M., Blath J., and Eldon B. Statistical properties of the site-frequency spectrum associated with λ-coalescents. *Genetics*, 195(3):1037–1053, 2013a. doi: 10.1534/genetics.113.156612.

Birkner M., Blath J., and Eldon B. An ancestral recombination graph for diploid populations with skewed offspring distribution. *Genetics*, 193(1):255–290, 2013b. doi: 10.1534/genetics.112.144329.

Birkner M., Černý J., Depperschmidt A., and Gantert N. Directed random walk on the backbone of an oriented percolation cluster. *Electronic Journal of Probability*, 18:1–35, 2013c. doi: 10.1214/EJP.v18-2302.

Birkner M., Liu H., and Sturm A. Coalescent results for diploid exchangeable population models. *Electronic Journal of Probability*, 23:1–44, 2018. doi: 10.1214/18-EJP175.

Blath J., Kadow S., and Ortgiese M. The largest strongly connected component in the cyclical pedigree model of wakeley et al. *Theoretical Population Biology*, 98:28–37, 2014. doi: 10.1016/j.tpb.2014.10.001.

Blath J., Cronjäger M. C., Eldon B., and Hammer M. The site-frequency spectrum associated with xi-coalescents. *Theoretical Population Biology*, 110:36–50, 2016. doi: 10.1016/j.tpb.2016.04.002.

Bolthausen E. and Sznitman A. On the static and dynamic points of view for certain random walks in random environment. *Methods and Applications of Analysis*, 9(3):345–376, 2002a. URL https://projecteuclid.org/journals/methods-and-applications-of-analysis/volume-9/issue-3/On-the-Satic-and-Dynamic-Points-of-View-for-Certain/maa/1119027729.full.

Bolthausen E. and Sznitman A. Ten lectures on random media. In *DMV-Seminar, Vol 32*, Oberwolfach Seminars. Birkhäuser, Basel, 2002b. doi: 10.1007/978-3-0348-8159-3.

Bouckaert R., Vaughan T. G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A., Heled J., Jones G., Kühnert D., De Maio N., Matschiner M., Mendes F. K., Müller N. F., Ogilvie H. A., du Plessis L., Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchard M. A., Wu C., Xie D., Zhang C., Stadler T., and Drummond A. J. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLOS Computational Biology*, 15(4):1–28, 2019. doi: 10.1371/journal.pcbi.1006650.

Brown W. M. Polymorphism in mitochondrial DNA of humans as revealed by restriction endonuclease analysis. *Proceedings of the National Academy of Sciences USA*, 77(6):3605–3609, 1980. doi: 10.1073/pnas.77.6.360.

Brown W. M., George M., and Wilson A. C. Rapid evolution of animal mitochondrial DNA. *Proceedings of the National Academy of Sciences USA*, 76(4):1967–1971, 1979. doi: 10.1073/pnas.76.4.1967.

Cann R. L., Stoneking M., and Wilson A. C. Mitochondrial DNA and human evolution. *Nature*, 325(6099):31–36, 1987. doi: 10.1038/325031a0.

Cannings C. The latent roots of certain Markov chains arising in genetics: a new approach. I. Haploid models. *Advances in Applied Probability*, 6(2):260–290, 1974. doi: 10.2307/1426293.

Cavalli-Sforza L. L. and Edwards A. W. F. Phylogenetic analysis: Models and estimation procedures. *Evolution*, 21(3):550–570, 1967. doi: 10.1111/j.1558-5646.1967.tb03411.x. Also published as: Cavalli-Sforza LL, Edwards AWF (1967) Phylogenetic analysis. Models and estimation procedures. Am J Hum Genet 19(3 Pt 1):233–257.

Chang J. T. Recent common ancestors of all present-day individuals. *Advances in Applied Probability*, 31(4):1002–1026, 1999. doi: 10.1239/aap/1029955256.

Charlesworth B. Fisher's historic 1922 paper On the dominance ratio. *Genetics*, 220(3):iyac006, 2022. doi: 10.1093/genetics/iyac006.

Coron C. and Le Jan Y. Pedigree in the biparental Moran model. *Journal of Mathematical Biology*, 84(6):51, 2022. doi: 10.1007/s00285-022-01752-0.

Der R. and Plotkin J. B. The equilibrium allele frequency distribution for a population with reproductive skew. *Genetics*, 196(4):1199–1216, 2014. doi: 10.1534/genetics.114.161422.

Derrida B., Manrubia S. C., and Zanette D. H. Statistical properties of genealogical trees. *Physical Review Letters*, 82:1987–1990, 1999. doi: 10.1103/PhysRevLett.82.1987.

Derrida B., Manrubia S. C., and Zanette D. H. Distribution of repetitions of ancestors in genealogical trees. *Physica A: Statistical Mechanics and its Applications*, 281(1):1–16, 2000a. doi: 10.1016/S0378-4371(00)00031-5.

Derrida B., Manrubia S. C., and Zannette D. H. On the genealogy of a population of biparental individuals. *Journal of Theoretical Biology*, 203(3):303–315, 2000b. doi: 10.1006/jtbi.2000.1095.

Di Rienzo A. and Wilson A. C. Branching pattern in the evolutionary tree for human mitochondrial DNA. *Proceedings of the National Academy of Sciences USA*, 88(5):1597–1601, 1991. doi: 10.1073/pnas.88.5.1597.

Donnelly P. and Kurtz T. G. Particle representations for measure-valued population models. *The Annals of Probability*, 27(1):166–205, 1999. doi: 10.1214/aop/1022677258.

Donnelly P., Tavaré S., Balding D. J., and Griffiths R. C. Estimating the age of the common ancestor of men from the *ZFY* intron. *Science*, 272(5266):1357–1359, 1996. doi: 10.1126/science.272.5266.1357.

Donnelly P., Wiuf C., Hein J., Slatkin M., Ewens W. J., and Kingman J. F. C. Discussion: Recent common ancestors of all present-day individuals. *Advances in Applied Probability*, 31(4):1027–1035, 1999. doi: 10.1239/aap/1029955257.

Dorit R. L., Akashi H., and Gilbert W. Absence of polymorphism at the *ZFY* locus on the human Y chromosome. *Science*, 268(5214):1183–1185, 1995. doi: 10.1126/science.7761836.

Durrett R. and Schweinsberg J. Approximating selective sweeps. *Theoretical Population Biology*, 66(2):129–138, 2004. doi: 10.1016/j.tpb.2004.04.002.

Durrett R. and Schweinsberg J. A coalescent model for the effect of advantageous mutations on the genealogy of a population. *Stochastic Processes and their Applications*, 115(10):1628–1657, 2005. doi: 10.1016/j.spa.2005.04.009.

Eldon B. Evolutionary genomics of high fecundity. *Annual Review of Genetics*, 54(1):213–236, 2020. doi: 10.1146/annurev-genet-021920-095932.

Eldon B. and Wakeley J. Linkage disequilibrium under skewed offspring distribution among individuals in a population. *Genetics*, 178(3):1517–1532, 2008. doi: 10.1534/genetics.107.075200.

Eldon B., Birkner M., Blath J., and Freund F. Can the site-frequency spectrum distinguish exponential population growth from multiple-merger coalescents? *Genetics*, 199(3):841–856, 2015. doi: 10.1534/genetics.114.173807.

Ewens W. J. A note on the sampling theory for infinite alleles and infinite sites models. *Theoretical Population Biology*, 6(2):143–148, 1974. doi: 10.1016/0040-5809(74)90020-3.

Ewens W. J. and Maruyama T. A note on the variance of the number of loci having a given gene frequency. *Genetics*, 80(1):221–222, 1975. doi: 10.1093/genetics/80.1.221.

Ewens W. J. Population genetics theory – the past and the future. In Lessard S., editor, *Mathematical and Statistical Developments of Evolutionary Theory*, pages 177–227. Kluwer Academic Publishers, Amsterdam, 1990.

Ewens W. J. *Mathematical Population Genetics, Volume I: Theoretical Foundations*. Springer-Verlag, Berlin, 2004.

Excoffier L., Dupanloup I., Huerta-Sánchez E., Sousa V. C., and Foll M. Robust demographic inference from genomic and SNP data. *PLOS Genetics*, 9(10):1–17, 2013. doi: 10.1371/journal.pgen.1003905.

Excoffier L., Marchi N., Marques D. A., Matthey-Doret R., Gouy A., and Sousa V. C. fastsimcoal2: demographic inference under complex evolutionary scenarios. *Bioinformatics*, 37(24):4882–4885, 2021. doi: 10.1093/bioinformatics/btab468.

Felsenstein J. Maximum-likelihood estimation of evolutionary trees from continuous characters. *American Journal of Human Genetics*, 25(5):471–492, 1973. URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1762641/.

Felsenstein J. Evolutionary trees from gene frequencies and quantitative characters: Finding maximum likelihood estimates. *Evolution*, 35(6):1229–1242, 11 1981. doi: 10.1111/j.1558-5646.1981.tb04991.x.

Felsenstein J. *Inferring Phylogenies*. Sinauer Associates, Inc, Sunderland, MA, 2004.

Fisher R. A. The correlation between relatives on the supposition of Mendelian inheritance. *Transactions of the Royal Society of Edinburgh*, 52:399–433, 1918. URL https://hdl.handle.net/2440/15097.

Fisher R. A. On the dominance ratio. *Proceedings of the Royal Society of Edinburgh*, 42:321–341, 1922. URL https://hdl.handle.net/2440/15098.

Fisher R. A. The distribution of gene ratios for rare mutations. *Proceedings of the Royal Society of Edinburgh*, 50:205–220, 1930. URL https://hdl.handle.net/2440/15106.

Freund F., Kerdoncuff E., Matuszewski S., Lapierre M., Hildebrandt M., Jensen J. D., Ferretti L., Lambert A., Sackton T. B., and Achaz G. Interpreting the pervasive observation of U-shaped site frequency spectra. *PLOS Genetics*, 19(3):1–18, 2023. doi: 10.1371/journal.pgen.1010677.

Fu Y. and Li W. Estimating the age of the common ancestor of men from the *ZFY* intron. *Science*, 272(5266):1356–1357, 1996. doi: 10.1126/science.272.5266.1356.

Gasbarra D., J S. M., and Arjas E. Backward simulation of ancestors of sampled individuals. *Theoretical Population Biology*, 67(2):75–83, 2005. doi: 10.1016/j.tpb.2004.08.003.

Gasbarra D., Pirinen M., Sillanpää M. J., and Arjas E. Estimating genealogies from linked marker data: a Bayesian approach. *BMC Bioinformatics*, 8(1):411, 2007a. doi: 10.1186/1471-2105-8-411.

Gasbarra D., Pirinen M., Sillanpää M. J., Salmela E., and Arjas E. Estimating genealogies from unlinked marker data: a Bayesian approach. *Theoretical Population Biology*, 72(3):305–322, 2007b. doi: 10.1016/j.tpb.2007.06.004.

Gernhard T. The conditioned reconstructed process. *Journal of Theoretical Biology*, 253(4):769–778, 2008. doi: 10.1016/j.jtbi.2008.04.005.

González Casanova A., Miró Pina V., and Siri-Jégousse A. The symmetric coalescent and Wright-Fisher models with bottlenecks. *The Annals of Applied Probability*, 32(1):235 – 268, 2022. doi: 10.1214/21-AAP1676.

Gravel S. and Steel M. The existence and abundance of ghost ancestors in biparental populations. *Theoretical Population Biology*, 101:47–53, 2015. doi: 10.1016/j.tpb.2015.02.002.

Griffiths and Tavaré S. Ancestral inference in population genetics. *Statistical Science*, 9(3):307–319, 1994. doi: 10.1214/ss/1177010378.

Griffiths R. C. and Marjoram P. An ancestral recombination graph. In Donnelly P. and Tavaré S., editors, *Progress in Population Genetics and Human Evolution (IMA Volumes in Mathematics and its Applications, vol. 87)*, pages 257–270. Springer-Verlag, New York, 1997.

Griffiths R. C. and Tavaré S. The age of a mutation in a general coalescent tree. *Communications in Statistics. Stochastic Models*, 14(1-2):273–295, 1998. doi: 10.1080/15326349808807471.

Guez J., Achaz G., Bienvenu F., Cury J., Toupance B., Heyer É., Jay F., and Austerlitz F. Cultural transmission of reproductive success impacts genomic diversity, coalescent tree topologies, and demographic inferences. *Genetics*, 223(4), 2023. doi: 10.1093/genetics/iyad007. iyad007.

Gutenkunst R. N., Hernandez R. D., Williamson S. H., and Bustamante C. D. Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLOS Genetics*, 5(10):1–11, 2009. doi: 10.1371/journal.pgen.1000695.

Haller B. C. and Messer P. W. SLiM 3: Forward genetic simulations beyond the Wright-Fisher model. *Molecular Biology and Evolution*, 36(3):632–637, 2019. doi: 10.1093/molbev/msy228.

Heled J. and Drummond A. J. Bayesian inference of species trees from multilocus data. *Molecular Biology and Evolution*, 27(3):570–580, 2009. doi: 10.1093/molbev/msp274.

Hudson R. R. Properties of a neutral allele model with intragenic recombination. *Theoretical Population Biology*, 23(2):183–201, 1983a. doi: 10.1016/0040-5809(83)90013-8.

Hudson R. R. Testing the constant-rate neutral allele model with protein sequence data. *Evolution*, 37(1):203–217, 1983b. doi: 10.1111/j.1558-5646.1983.tb05528.x.

Ingman M., Kaessmann H., Pääbo S., and Gyllensten U. Mitochondrial genome variation and the origin of modern humans. *Nature*, 408(6813):708–713, 2000. doi: 10.1038/35047064.

Kamm J., Terhorst J., Durbin R., and Song Y. S. Efficiently inferring the demographic history of many populations with allele count data. *Journal of the American Statistical Association*, 115 (531):1472–1487, 2020. doi: 10.1080/01621459.2019.1635482.

Karlin S. and McGregor J. The number of mutant forms maintained in a population. In Le Cam L. M. and Neyman J., editors, *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability: held at the Statistical Laboratory, University of California, June 21-July 18, 1965 and December 27, 1965-January 7, 1966*, pages 415–538. University of California Press, 1967.

Kelleher J., Wong Y., Wohns A. W., Fadil C., Albers P. K., and McVean G. Inferring whole-genome histories in large population datasets. *Nature Genetics*, 51(9):1330–1338, 2019. doi: 10.1038/s41588-019-0483-y.

Kimura M. The number of heterozygous nucleotide sites maintained in a finite population due to the steady flux of mutations. *Genetics*, 61(4):893–903, 1969. doi: 10.1093/genetics/61.4.893.

Kingman J. F. C. On the genealogy of large populations. *Journal of Applied Probability*, 19(A): 27–43, 1982. doi: 10.2307/3213548.

Ko A. and Nielsen R. Joint estimation of pedigrees and effective population size using Markov chain Monte Carlo. *Genetics*, 212(3):855–868, 2019. doi: 10.1534/genetics.119.302280.

Koskela J. Multi-locus data distinguishes between population growth and multiple merger coalescents. *Statistical Applications in Genetics and Molecular Biology*, 17(3):20170011, 2018. doi: doi:10.1515/sagmb-2017-0011.

Kuhner M. K., Yamato J., and Felsenstein J. Estimating effective population size and mutation rate from sequence data using Metropolis-Hastings sampling. *Genetics*, 140(4):1421–1430, 1995. doi: 10.1093/genetics/140.4.1421.

Lachance J. Inbreeding, pedigree size, and the most recent common ancestor of humanity. *Journal of Theoretical Biology*, 261(2):238–247, 2009. doi: 10.1016/j.jtbi.2009.08.006.

Lambert A. and Stadler T. Birth-death models and coalescent point processes: The shape and probability of reconstructed phylogenies. *Theoretical Population Biology*, 90:113–128, 2013. doi: 10.1016/j.tpb.2013.10.002.

Li H. and Durbin R. Inference of population history from individual whole-genome sequences. *Nature*, 475(7357):493–496, 2011. doi: 10.1038/nature10231.

Lundstrom R., Tavaré S., and Ward R. H. Estimating substitution rates from molecular data using the coalescent. *Proceedings of the National Academy of Sciences USA*, 89(13):5961–5965, 1992. doi: 10.1073/pnas.89.13.5961.

Malécot G. Etude mathématique des populations Mendélienne. *Annales de l'Université de Lyon, Sciences A*, 4:45–60, 1941.

Malécot G. La consanguinité dans une population limitée. *Comptes Rendus de l'Académie des Sciences, Paris*, 222:841–843, 1946.

Malécot G. *Les Mathématiques de l'Hérédité*. Masson, Paris, 1948. URL https://wellcomecollection.org/works/msfaxgkw. Extended translation: *The Mathematics of Heredity*. W.H. Freeman, San Francisco (1969).

Matsen F. A. and Evans S. N. To what extent does genealogical ancestry imply genetic ancestry? *Theoretical Population Biology*, 74(2):182–190, 2008. doi: 10.1016/j.tpb.2008.06.003.

Matuszewski S., Hildebrandt M. E., Achaz G., and Jensen J. D. Coalescent processes with skewed offspring distributions and nonequilibrium demography. *Genetics*, 208(1):323–338, 2018. doi: 10.1534/genetics.117.300499.

McVean G. A. T. A genealogical interpretation of linkage disequilibrium. *Genetics*, 162(2):987–991, 2002. doi: 10.1093/genetics/162.2.987.

Möhle M. A convergence theorem for Markov chains arising in population genetics and the coalescent with selfing. *Advances in Applied Probability*, 30(2):493–512, 1998a. doi: 10.1239/aap/1035228080.

Möhle M. Coalescent results for two-sex population models. *Advances in Applied Probability*, 30(2):513–520, 1998b. doi: 10.1239/aap/1035228081.

Möhle M. The concept of duality and applications to Markov processes arising in neutral population genetics models. *Bernoulli*, 5:761–777, 1999.

Möhle M. and Sagitov S. A classification of coalescent processes for haploid exchangeable population models. *The Annals of Probability*, 29(4):1547–1562, 2001. doi: 10.1214/aop/1015345761.

Molchanov S. A. Lectures on random media. In Bernard P., editor, *École d'Été de Probabilités de Saint-Flour XXII – 1992*, volume 1581 of *Lectures Notes in Mathematics*, pages 242–411. Springer-Verlag, Berlin, 1994. doi: 10.1007/BFb0073871.

Nielsen R. Estimation of population parameters and recombination rates from single nucleotide polymorphisms. *Genetics*, 154(2):931–942, 2000. doi: 10.1093/genetics/154.2.931.

Padmadisastra S. Estimating divergence times. *Theoretical Population Biology*, 34(3):297–319, 1988. doi: 10.1016/0040-5809(88)90026-3.

Patterson N., Moorjani P., Luo Y., Mallick S., Rohland N., Zhan Y., Genschoreck T., Webster T., and Reich D. Ancient admixture in human history. *Genetics*, 192(3):1065–1093, 11 2012. doi: 10.1534/genetics.112.145037.

Peter B. M. Admixture, population structure, and f-statistics. *Genetics*, 202(4):1485–1501, 2016. doi: 10.1534/genetics.115.183913.

Pitman J. Coalescents with multiple collisions. *The Annals of Probability*, 27(4):1870–1902, 1999. doi: 10.1214/aop/1022874819.

Ralph P., Thornton K., and Kelleher J. Efficiently summarizing relationships in large samples: A general duality between statistics of genealogies and genomes. *Genetics*, 215(3):779–797, 2020. doi: 10.1534/genetics.120.303253.

Ralph P. L. An empirical approach to demographic inference with genomic data. *Theoretical Population Biology*, 127:91–101, 2019. doi: 10.1016/j.tpb.2019.03.005.

Rannala B. and Yang Z. Bayes estimation of species divergence times and ancestral population sizes using DNA sequences from multiple loci. *Genetics*, 164(4):1645–1656, 2003. doi: 10.1093/genetics/164.4.1645.

Rohde D. L. T., Olson S., and Chang J. T. Modelling the recent common ancestry of all living humans. *Nature*, 431(7008):562–566, 2004. doi: 10.1038/nature02842.

Ronquist F., Teslenko M., van der Mark P., Ayres D. L., Darling A., Höhna S., Larget B., Liu L., Suchard M. A., and Huelsenbeck J. P. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, 61(3):539–542, 2012. doi: 10.1093/sysbio/sys029.

Sagitov S. The general coalescent with asynchronous mergers of ancestral lines. *Journal of Applied Probability*, 36(4):1116–1125, 1999. doi: 10.1239/jap/1032374759.

Sagitov S. Convergence to the coalescent with simultaneous multiple mergers. *Journal of Applied Probability*, 40(4):839–854, 2003. doi: 10.1239/jap/1067436085.

Sainudiin R., Thatte B., and Véber A. Ancestries of a recombining diploid population. *Journal of Mathematical Biology*, 72(1):363–408, 2016. doi: 10.1007/s00285-015-0886-z.

Sawyer S. A. and Hartl D. L. Population genetics of polymorphism and divergence. *Genetics*, 132 (4):1161–1176, 1992. doi: 10.1093/genetics/132.4.1161.

Schaper E., Eriksson A., Rafajlovic M., Sagitov S., and Mehlig B. Linkage disequilibrium under recurrent bottlenecks. *Genetics*, 190(1):217–229, 2012. doi: 10.1534/genetics.111.134437.

Schweiger R. and Durbin R. Ultrafast genome-wide inference of pairwise coalescence times. *Genome Research*, 33:1023–1031, 2023. doi: 10.1101/gr.277665.123.

Schweinsberg J. Coalescents with simultaneous multiple collisions. *Electronic Journal of Probability*, 5:1–50, 2000. doi: 10.1214/EJP.v5-68.

Schweinsberg J. and Durrett R. Random partitions approximating the coalescence of lineages during a selective sweep. *Annals of Applied Probability*, 15(3):1591–1651, 2005. doi: 10.1214/105051605000000430.

Sheehan S., Harris K., and Song Y. S. Estimating variable effective population sizes from multiple genomes: A sequentially Markov conditional sampling distribution approach. *Genetics*, 194(3):647–662, 2013. doi: 10.1534/genetics.112.149096.

Sjödin P., Kaj I., Krone S., Lascoux M., and Nordborg M. On the meaning and existence of an effective population size. *Genetics*, 169(2):1061–1070, 2005. doi: 10.1534/genetics.104.026799.

Slatkin M. Inbreeding coefficients and coalescence times. *Genetics Research*, 58(2):167–175, 1991. doi: 10.1017/S0016672300029827.

Speidel L., Forest M., Shi S., and Myers S. R. A method for genome-wide genealogy estimation for thousands of samples. *Nature Genetics*, 51(9):1321–1329, 2019. doi: 10.1038/s41588-019-0484-x.

Spence J. P., Kamm J. A., and Song Y. S. The site frequency spectrum for general coalescents. *Genetics*, 202(4):1549–1561, 2016. doi: 10.1534/genetics.115.184101.

Suchard M. A., Lemey P., Baele G., Ayres D. L., Drummond A. J., and Rambaut A. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evolution*, 4(1):vey016, 2018. doi: 10.1093/ve/vey016.

Tajima F. Evolutionary relationship of DNA sequences in finite populations. *Genetics*, 105(2):437–460, 1983. URL https://www.genetics.org/content/105/2/437.

Tellier A. and Lemaire C. Coalescence 2.0: a multiple branching of recent theoretical developments and their applications. *Molecular Ecology*, 23(11):2637–2652, 2014. doi: 10.1111/mec.12755.

The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*, 526(7571):68–74, 2015. doi: 10.1038/nature15393.

The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*, 526:68–74, 2015. doi: 10.1038/nature15393.

Tyukin A. Quenched limits of coalescents in fixed pedigrees. Master's thesis, Johannes-Gutenberg-Universität Mainz, Germany, 2015. URL https://www.glk.uni-mainz.de/files/2018/08/andrey_tyukin_msc.pdf.

Vigilant L., Pennington R., Harpending H., Kocher T. D., and Wilson A. C. Mitochondrial DNA sequences in single hairs from a southern african population. *Proceedings of the National Academy of Sciences USA*, 86(23):9350–9354, 1989. doi: 10.1073/pnas.86.23.9350.

Vigilant L., Stoneking M., Harpending H., Hawkes K., and Wilson A. C. African populations and the evolution of human mitochondrial DNA. *Science*, 253(5027):1503–1507, 1991. doi: 10.1126/science.1840702.

Wakeley J. Nonequilibrium migration in human history. *Genetics*, 153(4):1863–1871, 1999. doi: 10.1093/genetics/153.4.1863.

Wakeley J. *Coalescent Theory: An Introduction*. Roberts & Company Publishers, Greenwood Village, Colorado, 2009. Current publisher: Macmillan Learning, New York, NY.

Wakeley J. and Lessard S. Theory of the effects of population structure and sampling on patterns of linkage disequilibrium applied to genomic data from humans. *Genetics*, 164(3):1043–1053, 2003. doi: 10.1093/genetics/164.3.1043.

Wakeley J., King L., Low B. S., and Ramachandran S. Gene genealogies within a fixed pedigree, and the robustness of Kingman's coalescent. *Genetics*, 190(4):1433–1445, 2012. doi: 10.1534/genetics.111.135574.

Wakeley J., King L., and Wilton P. R. Effects of the population pedigree on genetic signatures of historical demographic events. *Proceedings of the National Academy of Sciences USA*, 113(29): 7994–8001, 2016. doi: 10.1073/pnas.160108011.

Wang K., Mathieson I., O'Connell J., and Schiffels S. Tracking human population structure through time from whole genome sequences. *PLOS Genetics*, 16(3):1–24, 03 2020. doi: 10.1371/journal. pgen.1008552.

Ward R. H., Frazier B. L., Dew-Jager K., and Pääbo S. Extensive mitochondrial diversity within a single amerindian tribe. *Proceedings of the National Academy of Sciences USA*, 88(19):8720–8724, 1991. doi: 10.1073/pnas.88.19.8720.

Watterson G. A. Estimating species divergence times using multi-locus data. In Ohta T. and Aoki K., editors, *Population Genetics and Molecular Evolution: Papers Marking the Sixtieth Birthday of Motoo Kimura*, pages 163–183. Japan Scientific Societies Press; Springer-Verlag, Tokyo; Berlin, New York, 1985.

Weiss G. and von Haeseler A. Estimating the age of the common ancestor of men from the *ZFY* intron. *Science*, 272(5266):1359–1360, 1996. doi: 10.1126/science.272.5266.1359.

Wilson A. C., Cann R. L., Carr S. M., George M., Gyllensten U. B., Helm-Bychowski K. M., Higuchi R. G., Palumbi S. R., Prager E. M., Sage R. D., and Stoneking M. Mitochondrial DNA and two perspectives on evolutionary genetics. *Biological Journal of the Linnean Society*, 26(4): 375–400, 1985. doi: 10.1111/j.1095-8312.1985.tb02048.x.

Wilton P. R., Baduel P., Landon M. M., and Wakeley J. Population structure and coalescence in pedigrees: Comparisons to the structured coalescent and a framework for inference. *Theoretical Population Biology*, 115:1–12, 2017. doi: 10.1016/j.tpb.2017.01.004.

Wohns A. W., Wong Y., Ben J., Akbari A., Mallick S., Pinhasi R., Patterson N., Reich D., Kelleher J., and McVean G. A unified genealogy of modern and ancient genomes. *Science*, 375(6583): eabi8264, 2022. doi: 10.1126/science.abi8264.

Wolfram Research, Inc. Mathematica, version 13.1, 2022. Version 13.1.

Wollenberg K. and Avise J. C. Sampling properties of genealogical pathways underlying population pedigrees. *Evolution*, 52(4):957–966, 1998. doi: 10.1111/j.1558-5646.1998.tb01825.x.

Wooding S. and Rogers A. The matrix coalescent and an application to human single-nucleotide polymorphisms. *Genetics*, 161(4):1641–1650, 2002. doi: 10.1093/genetics/161.4.1641.

Wright S. Systems of mating. I. The biometric relations between parent and offspring. *Genetics*, 6 (2):111–123, 1921a. doi: 10.1093/genetics/6.2.111.

Wright S. Systems of mating. II. The effects of inbreeding on the genetic composition of a population. *Genetics*, 6(2):124–143, 1921b. doi: 10.1093/genetics/6.2.124.

Wright S. Systems of mating. III. Assortative mating based on somatic resemblance. *Genetics*, 6 (2):144–161, 1921c. doi: 10.1093/genetics/6.2.144.

Wright S. Systems of mating. IV. The effects of selection. *Genetics*, 6(2):162–66, 1921d. doi: 10.1093/genetics/6.2.162.

Wright S. Systems of mating. V. General considerations. *Genetics*, 6(2):167–178, 1921e. doi: 10.1093/genetics/6.2.167.

Wright S. Coefficients of inbreeding and relationship. *The American Naturalist*, 56(645):330–338, 1922. doi: 10.1086/279872.

Wright S. Evolution in Mendelian populations. *Genetics*, 16(2):97–159, 1931. URL `https://www.genetics.org/content/16/2/97`.

Yang Z. Likelihood and Bayes estimation of ancestral population sizes in hominoids using data from multiple loci. *Genetics*, 162(4):1811–1823, 12 2002. doi: 10.1093/genetics/162.4.1811.

Zhang B. C., Biddanda A., Gunnarsson Á. F., Cooper F., and Palamara P. F. Biobank-scale inference of ancestral recombination graphs enables genealogical analysis of complex traits. *Nature Genetics*, 55(5):768–776, 2023. doi: 10.1038/s41588-023-01379-x.