1 An RNAi screen for conserved kinases that enhance microRNA activity after dauer in *C.*
2 *elegans*
3

4 Himal Roka Pun[1,2], Xantha Karp[1,2]
5
6 1) Department of Biology
7 2) Biochemistry, Cell and Molecular Biology Program
8   Central Michigan University, Mount Pleasant, MI 48859
9
10 Correspondence: karp1x@cmich.edu

11
12 Running title: Post-dauer kinase screen
13
14
15

Keywords: Diapause, dauer, *C. elegans,* microRNA, Argonaute, kinase, chromosome
segregation, molting, checkpoint-kinase

Abstract:

Gene regulation in changing environments is critical for maintaining homeostasis.

Some animals undergo a stress-resistant diapause stage to withstand harsh

environmental conditions encountered during development. MicroRNAs (miRNAs) are

one mechanism for regulating gene expression during and after diapause. MicroRNAs

downregulate target genes post-transcriptionally through the activity of the miRNA-

Induced Silencing Complex (miRISC). Argonaute is the core miRISC protein that binds

to both the miRNA and to other miRISC proteins. The two major miRNA Argonautes in

the *C. elegans* soma are ALG-1 and ALG-2, which function partially redundantly. Loss

of *alg-1 (alg-1(0))* causes penetrant developmental phenotypes including vulval defects

and the reiteration of larval cell programs in hypodermal cells. However, these

phenotypes are essentially absent if *alg-1(0)* animals undergo a diapause stage called

dauer. Levels of the relevant miRNAs are not higher during or after dauer, suggesting

that activity of the miRISC may be enhanced in this context. To identify genes that are

required for *alg-1(0)* mutants to develop without vulval defects after dauer, we

performed an RNAi screen of genes encoding conserved kinases. We focused on

kinases because of their known role in modulating miRISC activity. We found RNAi

knockdown of four kinase-encoding genes, *air-2*, *bub-1*, *chk-1,* and *nekl-3*, caused

vulval defects and reiterative phenotypes in *alg-1(0)* mutants after dauer, and that these

defects were more penetrant in an *alg-1(0)* background than in wild type. Our results

implicate these kinases as potential regulators of miRISC activity during post-dauer

development in *C. elegans*.

Introduction:

The regulation of gene expression in response to changing environments is critical to maintaining homeostasis. One mechanism for modulating gene expression in different contexts is microRNA-mediated gene silencing. MicroRNAs (miRNAs) are ~22 nucleotide non-coding RNAs that regulate mRNA transcripts post-transcriptionally via the miRNA Induced Silencing Complex (miRISC). MiRNAs bind to the 3'UTR of the target mRNA via imperfect complementarity, thereby recruiting miRISC. Argonaute is the core miRNA-interacting protein which in turn binds with other protein cofactors (Bartel 2018). The protein factors in miRISC inhibit translation and cause destabilization of the transcript, thus silencing the expression of the target (Jonas and Izaurralde 2015; Bartel 2018).

Changes in miRNA activity can influence gene expression in a variety of physiological or environmental contexts (Leung and Sharp 2010; Galagali and Kim 2020). When confronted with adverse environments, some animals can enter a stress-resistant and developmentally arrested diapause stage (Hand *et al.* 2016). MicroRNAs have been implicated in the regulation of diapause in insects and nematodes (Alvarez-Saavedra and Horvitz 2010; Karp *et al.* 2011; Meuti *et al.* 2018). Although changes in miRNA levels have been documented before, during, and after diapause, little is known about changes in miRISC activity and how they may impact development in the diapause life history.

C. elegans larval development is a useful model system to explore the modulation of miRISC activity during diapause. *C. elegans* larvae develop through four larval stages (L1-L4) before becoming adults (Byerly *et al.* 1976). Larval development

occurs continuously when the environment is favorable for reproduction. In unfavorable

environments, larvae can pause their development after the second larval molt in the

stress-resistant dauer diapause stage (Cassada and Russell 1975). If dauer larvae

encounter favorable environmental conditions, they can recover and resume

development through post-dauer larval stages that are developmentally equivalent to

those occurring in continuously developing larvae (Liu and Ambros 1991; Euling and

Ambros 1996).

One developmental system regulated by miRNAs is the hypodermal stem-cell-

like seam cells. Seam cells divide in a particular pattern and sequence at each larval

stage. At adulthood, seam cells cease dividing and differentiate (Sulston and Horvitz

1977; Ambros and Horvitz 1984). Larval vs. adult seam programs are regulated by

heterochronic genes that encode miRNAs and their targets (Ambros and Horvitz 1984;

Rougvie and Moss 2013; Galagali and Kim 2020). The loss of heterochronic miRNAs or

impaired function of miRISC proteins causes a reiterative phenotype whereby the

pattern and sequence of seam cell divisions appropriate to earlier developmental stages

is reiterated in later stages (Ambros and Horvitz 1984; Reinhart *et al.* 2000; Grishok *et*

al. 2001; Abbott *et al.* 2005).

ALG-1 and ALG-2 are the two major somatic miRNA-specific Argonaute proteins

in *C. elegans* and act partially redundantly (Grishok *et al.* 2001; Tops *et al.* 2006;

Vasquez-Rifo *et al.* 2012). While loss of both *alg-1* and *alg-2* results in embryonic

lethality, loss of *alg-1* alone *(alg-1(0))* causes a reiterative phenotype, suggesting *alg-2*

is not sufficient to mediate the activity of the heterochronic miRNAs (Grishok *et al.* 2001;

Zinovyeva *et al.* 2014). Strikingly, this phenotype is strongly suppressed in post-dauer

alg-1(0) mutant animals (Karp and Ambros 2012). This finding suggests that miRNA

activity is enhanced after dauer, such that *alg-2* activity becomes sufficient. Prior work

has demonstrated that levels of heterochronic miRNAs are the same or reduced in post-

dauer larvae compared to continuously developing larvae, suggesting that the

enhanced miRNA activity may arise from modulation of miRISC (Karp *et al.* 2011; Karp

and Ambros 2012).

Kinases are good candidates to mediate miRISC enhancement after dauer

because they are known modulators of miRISC activity. In *C. elegans* and other

animals, kinases phosphorylate Argonaute and other miRISC components, thereby

altering miRISC activity (Wilczynska and Bushell 2015; Frédérick and Simard 2022). To

identify conserved kinases that may enhance miRNA activity after dauer, we carried out

an RNAi screen for conserved kinase-encoding genes that are necessary for the post-

dauer suppression of *alg-1(0)* reiterative phenotypes. We found that *air-2, bub-1, chk-1,*

and *nekl-3* are required to prevent vulval defects and reiterative phenotypes in *alg-1(0)*

mutants after dauer. Furthermore, RNAi of these genes causes more penetrant

phenotypes in an *alg-1* background than in wild-type, suggesting that the kinases they

encode may enhance miRNA function after dauer.

Methods:

***C. elegans* strain and maintenance.** All strains were maintained on Nematode Growth

Medium (NGM) plates seeded with *E. coli* strain OP50 at 15˚C or 20˚C (Brenner 1974).

Strains used in the study were: XV88 *daf-7(e1372); maIs105[col-19p::gfp]; alg-*

1(gk214), VT1777 *daf-7(e1372); maIs105,* XV85 *alg-2(ok302); daf-7(e1372); maIs105,*

and VT1274 *alg-2(ok302); maIs105.*

RNAi. A list of genes encoding kinases conserved between *C. elegans* and humans

was obtained from (Deng *et al.* 2019). Their list was generated using the Ortholist tool

(Shaye and Greenwald 2011; Kim *et al.* 2018). A complete list of RNAi clones used, and

their sources is provided in Table S1. RNAi bacteria were grown in LB media containing

50µL/mL carbenicillin shaking overnight at 37˚C. RNAi bacteria were then seeded on

NGM plates containing 200µg/mL IPTG and 50µg/mL carbenicillin ("RNAi plates"). For

the primary screen, 12-well plates were used, and for all other experiments, 60mm

plates were used. The primary screen was conducted in duplicate. RNAi clones that

caused reiterative alae defects in post-dauer adults were verified by Sanger sequencing

using the M13F primer (Eurofins Genomics).

Embryo Isolation. Gravid adult hermaphrodites were treated with a bleach solution

(0.4% sodium hypochlorite dissolved in 1M NaOH) for two 2-minute incubations at room

temperature. The embryos were washed with sterile water and then added to RNAi

plates.

Dauer Induction. The *daf-7(e1372)* allele was used to induce dauer formation (Vowels

and Thomas 1992; Karp 2018). Embryos were incubated on RNAi plates at 24°C for 48

– 50 hours, corresponding to the time just after the molt into dauer (Karp 2018).

Dauer recovery. For the primary screen, dauer recovery was induced by washing

dauer larvae off the RNAi plates with sterile water, adding the dauer larvae to fresh

RNAi plates, and shifting the worms to 20°C. For all other experiments, dauer larvae

were selected by incubating worms with 1% (w/v) SDS (sodium dodecyl sulfate) for 30

minutes at room temperature (Cassada and Russell 1975; Karp 2018). The dauer

larvae were then washed with sterile water, added to fresh RNAi plates, and shifted to

20°C to induce dauer recovery. Post-dauer larvae were incubated at 20°C for 48 hours

to obtain young post-dauer adults.

New RNAi clones. RNAi clones for *air-2, chk-1,* and *bub-1* were created based on the

protocol described in (Kamath and Ahringer 2003). Briefly, primers were designed to

amplify regions of these genes that were distinct from those contained in the Ahringer

RNAi clones. The amplified sequences were cloned into the L4440 vector by TA

cloning. The resulting plasmids were sequenced, then transformed into HT115 bacteria.

Primers used for cloning are listed below.

air-2 forward: 5'-TACTCCACAGAAGGGAGGGT-3'

air-2 reverse: 5'-ATGTTGGCCACTAAGCTGAAATC-3'

chk-1 forward: 5'-GGCGGAGAGACAGAATGCTT-3'

chk-1 reverse: 5'-CCGAGTGCTCCACATTGACT-3'

bub-1 forward: 5' CCGTCGACATGTGGTCTTGA-3'

bub-1 reverse: 5'-GAGGTTTGCGTCACTGGAGA-3'

Microscopy. For the primary screen, bursting and/or protrusion of the vulva (Rup or Pvl

phenotypes) of the post-dauer adults were assessed using a dissecting microscope

(Zeiss Stereo V12 fitted with M2 Bio for fluorescence). In the secondary screen, the *alg-1(0)* mutant phenotypes were assessed using a Zeiss AxioImager D2 compound

microscope. Post-dauer animals were immobilized using 0.1M levamisole on 2%

agarose pads. DIC and fluorescence images were taken using an AxioCam MRm Rev 3

camera and ZEN 3.2 software. GFP was visualized with a high-efficiency GFP shift-free

filter at 63x with an exposure time of 9ms.

Data availability. All *C. elegans* strains and newly made RNAi clones are available

upon request. The authors state that all data necessary for confirming the conclusions

of the article are present within the article, figures, and tables, including supplementary

information.

Results and discussion.

RNAi screen of the conserved kinome to identify miRISC regulators acting after dauer.

If *C. elegans* larvae lacking *alg-1* develop continuously from embryo to adult, the

adults display reiterative phenotypes. In contrast, these phenotypes are suppressed if

alg-1(0) mutant larvae develop through the dauer diapause stage (Karp and Ambros

2012). We hypothesized that modulation of miRISC activity in post-dauer animals could

account for this difference. Since kinases are known to modulate miRISC in various

physiological and environmental contexts (Wu *et al.* 2011; Alessi *et al.* 2015; Olejniczak

et al. 2018), we designed an RNAi screen of all of the conserved kinases in *C. elegans*

(Kim *et al.* 2018; Deng *et al.* 2019). If post-dauer *alg-1(0)* mutants display reiterative

phenotypes after depletion of a particular kinase, that kinase is a candidate for

modulating miRISC function in post-dauer animals.

To perform our RNAi screen, we used a strain that contained the temperature-

sensitive *daf-7(e1372)* allele to induce dauer formation at 24˚C and allow recovery at

lower temperatures (Vowels and Thomas 1992; Karp 2018). Embryos were added to

RNAi plates and incubated at 24˚C for 2 days, or until just after dauer formation. Dauer

larvae were washed off with water, transferred to fresh RNAi plates, and shifted to 20˚C

to promote recovery. Post-dauer *alg-1(0)* adults were then screened for the vulval

defects associated with compromised miRISC function (Figure 1A). We used *lacZ* RNAi

as a negative control and *kin-3* RNAi as a positive control because we previously

showed that RNAi of *kin-3* caused reiterative phenotypes in post-dauer *alg-1(0)* mutants

(Alessi *et al.* 2015). *kin-3* encodes the catalytic subunit of Casein Kinase II (CK2), and

CK2 phosphorylates the miRISC proteins ALG-1 and CGH-1 to promote miRISC

function (Alessi *et al.* 2015; Shah *et al.* 2023). We used a previously published list of the

247 *C. elegans* kinases with human orthologs (Kim *et al.* 2018; Deng *et al.* 2019). Some

kinases are represented by more than one RNAi clone, giving us 286 RNAi clones in

total (Table S1).

In our primary screen, performed in duplicate, we used a dissecting microscope

to look for vulval defects in *alg-1(0)* post-dauer adults. Vulval defects included both

bursting (Rup) and protruding vulva (Pvl) phenotypes. We focused on vulval defects

because these phenotypes are easily visible in the dissecting microscope. As expected,

we saw almost no vulval defects after treatment with *lacZ* RNAi (Figure 1B). By

contrast, *kin-3* RNAi caused a high penetrance of vulval defects (Figure 1B). We found

12 additional conserved kinases that caused similar phenotypes. We observed a broad

range of penetrance of vulval defects which could be due to different degrees of

requirement for vulval development and/or variable effectiveness of RNAi.

Fig. 1. Kinases identified in a primary screen for enhancement of an *alg-1(0)* phenotype after dauer. A. Strategy for RNAi screening. The *daf-7(e1372)* allele was used to induce dauer formation at 24°C and allow recovery at lower temperatures. See Methods for details. **B.** RNAi clones that produced vulval defects (Rup and/or Pvl phenotypes) in the primary screen. The screen was performed in duplicate and clones that produced vulval defects in both plates are shown here. Table S1 shows data for all clones that produced vulval defects in at least one well. The remaining clones produced no vulval defects in either well (≥20 adults scored per well). Numbers in B and C indicated the number of post-dauer (PD) adults displaying vulval defects over the total PD adults scored. **C.** The 12 hits from the primary screen were retested for vulval defects. Seven genes caused bursting and/or protrusion of the vulva in post-dauer *alg-1(0)* adults. Among these 7 genes, 6 were novel *alg-1* interactors.

To ensure our results were reproducible, we re-tested the 12 hits from the

primary screen. In addition, in this validation experiment, we selected dauer larvae by

treatment with 1% SDS (Cassada and Russell 1975). Treatment with SDS ensured that

the adults we screened were indeed post-dauer and not displaying vulval defects

because the larvae had escaped dauer formation. Upon retesting, seven genes

displayed vulval defects (Figure 1C). One of these genes, *kin-10*, encodes the

regulatory subunit of the of CK2 protein and was therefore expected (Alessi *et al.* 2015;

Shah *et al.* 2023). We, therefore, focused on the remaining six kinase-encoding genes.

Five kinase-encoding genes contribute to adult alae formation in *alg-1(0)* post-

dauer adults.

In addition to vulval defects, *alg-1(0)* mutant adults display gapped or missing

alae after continuous development (Grishok *et al.* 2001; Vasquez-Rifo *et al.* 2012). This

reiterative phenotype occurs in heterochronic mutants because the underlying seam

cells repeat larval cell programs (Ambros and Horvitz 1984). In addition to gapped or

missing alae, *alg-1(0)* animals that develop continuously can display other defects in

alae formation, including small breaks and growth that deviates from the straight line

observed in wild-type adults (Alessi *et al.* 2015; Shah *et al.* 2023). Furthermore, the loss

of *alg-1* can reduce the expression of the adult cell fate marker *col-19p::gfp,* albeit at

low penetrance (Zinovyeva *et al.* 2015). If the kinases we identified were important for

miRNA function we would expect depletion of these genes to cause alae defects and

possibly reduced *col-19p::gfp* in the hypodermis of post-dauer *alg-1(0)* adults.

To test the hypothesis that these kinases are required to prevent reiterative

phenotypes in post-dauer *alg-1(0)* adults, we performed RNAi on the six hits and looked

at adult alae formation and *col-19p::gfp* expression with DIC and fluorescence

microscopy. We found that knocking down five of the six genes with RNAi resulted in

penetrant alae defects (Figure 2). The most penetrant alae phenotypes were caused by

bub-1 RNAi. RNAi of *cdk-9* produced no reiterative phenotypes and occasional other

alae defects. While *lacZ* RNAi produced no alae defects at all, *cdk-9* RNAi was not

significantly different from the *lacZ* control. In contrast to the alae defects produced by

RNAi, *col-19p::gfp* expression was not reduced by RNAi of any of the six candidate

genes and only moderately downregulated in our *kin-3* positive control (Figure S1).

Uncoupling of alae defects and *col-19p::gfp* expression has been previously observed in

some mutant backgrounds (Hada *et al.* 2010; Hansen *et al.* 2022).

RNAi of some kinases produces a stronger phenotype in a miRISC-sensitized

Fig. 2 RNAi of five kinase-encoding genes produces defects in adult alae formation in post-dauer *alg-1(0)* mutants. **A**. Comparison of alae phenotypes in post-dauer *alg-1(0)* adults. A compound microscope with DIC optics was used to visualize the presence of alae in post-dauer adult animals. White lines represent complete alae, purple lines represent gapped alae, and black lines show other alae defects including small breaks or curved or swirly alae. **B**. Quantification of alae defects. **** = P <0.0002; ns = P > 0.05, Fisher Exact Test.

background and during post-dauer development.

Vulval defects can be caused by the depletion of genes that regulate processes

other than miRISC activity. For example, loss of genes involved in regulation of uterine

development can produce similar phenotypes (Seydoux and Greenwald 1989). If the

kinases identified in our screen enhance miRISC activity after dauer, we would expect

that RNAi of these kinases would produce a stronger phenotype in *alg-1(0)* adults than

when *alg-1* was wild type. To test this hypothesis, we performed RNAi on *alg-1(0)* and

alg-1(+) animals in parallel. RNAi of *bub-1, air-2, nekl-3,* and *chk-1* each displayed a

more penetrant phenotype in the miRISC-compromised background (Figure 3A). RNAi

of *trr-1* or *cdk-9* caused a low-penetrance phenotype in both backgrounds, and there

was no significant difference between the backgrounds (Figure 3A). While these data do

not rule out the involvement of any of these genes in regulating miRISC activity, they

provide higher confidence in *bub-1, air-2, nekl-3,* and *chk-1* than *trr-1* or *cdk-9.*

Since *alg-1(0)* mutants display vulval defects during continuous development but

not after dauer, we next wondered whether the kinases we identified play a greater role

in preventing vulval defects in post-dauer hermaphrodites than in those that develop

continuously, particularly when miRISC is compromised. We were unable to test this

question using *alg-1(0)* mutants because they display penetrant phenotypes during

continuous development (Figure 3B). However, animals lacking *alg-2* appear

superficially wild-type in both life histories (Grishok *et al.* 2001; Karp and Ambros 2012).

We therefore compared phenotypes in each life history following RNAi of *bub-1, nekl-3,*

air-2, and *chk-1* in an *alg-2(0)* mutant background. We found that RNAi of *bub-1* and

nekl-3 both produced statistically more penetrant phenotypes after dauer (Figure 3C),

consistent with these kinases playing a larger role in post-dauer development.

Fig. 3. RNAi of some kinases caused more penetrant phenotypes after dauer and when miRISC was compromised. A. Vulval defects in post-dauer adults with wild-type miRISC were compared to animals with compromised miRISC function *(alg-1(0)* mutants). **B.** During continuous development, the penetrance of vulval defects in *alg-1(0)* adults is high such that no further enhancement is seen when *kin-3* is knocked down. **C.** Vulval defects in *alg-2(0)* young adults after dauer (PD = post-dauer) or after continuous development (CD). Dauer was induced by *daf-7(e1372),* whereas continuously developing animals were wild-type for *daf-7*. *p*-value >0.05 (ns), <0.0332 (*), <0.0021 (**), <0.0002 (****), Fisher exact test. **D)** Model for role of miRISC activity in different developmental contexts to control larval vs. adult seam cell developmental programs.

By contrast, RNAi of *air-2* and *chk-1* produced low-penetrance phenotypes in both life

histories in the *alg-2(0)* background (Figure 3C). While in this latter case there was no

statistical difference between post-dauer and continuously developing animals,

observing a decrease in penetrance in adults that had developed continuously would be

difficult given the low penetrance phenotype in post-dauer adults. Thus, we are unable

to draw strong conclusions about the relative role of *air-2* and *chk-1* in post-dauer vs.

continuous development.

Finally, we performed RNAi of *bub-1, air-2, nekl-3,* and *chk-1* using independent

clones that target different regions of each gene than the region targeted in the original

clones. We observed vulval defects using each set of clones, albeit at reduced

penetrance (Figure S2). The percent of vulval defects observed for each clone was

statistically different from the empty vector control, except the newly created *chk-1* clone

(P = 0.088). Both original and newly created *chk-1* clones produced low-penetrance

phenotypes, whereas the negative control produced no vulval defects out of 226 adults

examined.

Kinases identified in this screen

We have identified four kinases that, when depleted, reproducibly enhanced two

different *alg-1(0)* phenotypes in post-dauer animals and whose depletion caused more

severe defects in a miRISC-compromised background than in wild type (Table 1). The

known roles for these kinases are described below.

Among the four genes identified as the highest confidence candidates in our

screen, *nekl-3* is the only one to have been previously implicated in the regulation of

Table 1: Conservation and function of genes isolated in the screen

C. elegans gene	Human ortholog	Connection to microRNA function	Characterized function	Reference
air-2	AURKB (Aurora kinase B)	None	Chromosome segregation and cytokinesis in mitosis and meiosis	(Schumacher *et al.* 1998; Rogers *et al.* 2002)
bub-1	BUB1 (BUB1 mitotic checkpoint serine/threonine kinase)	None	Chromosome alignment and segregation, spindle assembly, and promotion of anaphase in mitosis and meiosis	(Sharp-Baker and Chen 2001; Kim *et al.* 2015; Edwards *et al.* 2018)
chk-1	CHEK1 (checkpoint kinase 1)	None	DNA damage checkpoint and embryonic S-M checkpoint kinase	(Kalogeropoulos *et al.* 2004)
nekl-3	NEK6/7 (NIMA-related kinase 6/7)	Identified as a weak enhancer of the *let-7(mg279)* Rup phenotype during continuous development.	Clathrin-mediated endocytosis of cargo necessary for molting	(Parry *et al.* 2007; Joseph *et al.* 2020)

miRNA activity. *nekl-3* was isolated in a large-scale RNAi screen for genes that

enhance the Rup phenotype of mutants with reduced activity of the *let-7* miRNA (Parry

et al. 2007). Although *nekl-3* was a relatively weak enhancer and the screen was

performed during continuous (non-dauer) development, this observation is consistent

with the hypothesis that *nekl-3* can promote miRISC function. This role for *nekl-3*

remains unexplored. In contrast, a role for *nekl-3* in the regulation of molting has been

well-characterized. *nekl-3* is required for the internalization via clathrin-mediated

endocytosis of cargo important for molting (Yochem *et al.* 2015; Joseph *et al.* 2020).

Specifically*, nekl*-3 is required for the L1-L2 and L2-L3 molts but not later molts during

continuous development*.* In the dauer life history, *nekl*-3 has a reduced requirement for

the L2d-dauer molt and is dispensable for post-dauer molting (Binti *et al.* 2022).

In mammals, the *nekl-3* orthologs, NEK family proteins, are best characterized

for their role in mitosis and chromosome segregation (Fry *et al.* 2012). Notably, *air-2*

and *bub-1* both encode conserved proteins that are critical for chromosome segregation

and mitosis and meiosis across species (Schumacher *et al.* 1998; Sharp-Baker and

Chen 2001; Rogers *et al.* 2002; Kim *et al.* 2015). Neither of these genes have been

previously implicated in miRNA activity. Previous large-scale screens conducted during

continuous development found that RNAi knockdown of *bub-1* produced Pvl and

bursting phenotypes (Fraser *et al.* 2000). Loss-of-function *bub-1* mutants display a

vulvaless (Vul) phenotype and a low-penetrance Pvl phenotype (Wang *et al.* 2009).

However, whether these phenotypes are related to the characterized role in

chromosome segregation or whether they may indicate a role for *bub-1* in regulating

miRNA activity is unknown. Finally, *chk-1* encodes a serine-threonine kinase that plays

a role in the checkpoint response to DNA damage and in the S-M checkpoint in

embryos (Kalogeropoulos *et al.* 2004).

Concluding Remarks

The goal of this study was to identify kinase-encoding genes that may enhance

miRISC function in post-dauer development. This study initially found six candidate

genes required to prevent developmental phenotypes in post-dauer *alg-1(0)* animals.

Four of these genes, *bub-1, air-2, nekl-3,* and *chk-1*, showed higher penetrance

phenotypes in a miRISC-compromised background, making these four genes the most

likely to affect miRISC activity after dauer. *bub-1* and *nekl-3* produced more penetrant

phenotypes after dauer than during continuous development in an *alg-2(0)* background,

whereas *air-2* and *chk-1* produced low-penetrance phenotypes in both life histories in

an *alg-2(0)* background. Prior to this work, the only kinase-encoding gene known to

enhance *alg-1(0)* phenotypes after dauer was *kin-3,* used as a positive control in this

study (Alessi *et al.* 2015). During continuous development, three kinases have been

shown to phosphorylate miRISC components to modulate different aspects of miRNA

activity: KIN-1/PKA, casein kinase 1 (CK1) and CK2 (Alessi *et al.* 2015; Huberdeau *et*

al. 2022; Shah *et al.* 2023). In our screen, only the CK2-encoding genes *kin-3* and *kin-*

10 were found to be necessary to prevent vulval defects in *alg-1(0)* mutants after dauer,

whereas RNAi of *kin-1* and *kin-19,* encoding PKA and CK1 respectively, did not cause

vulval defects in our primary screen (*kin-1,* 0/40; *kin-19,* 0/60). While these findings may

be explained by incomplete knockdown of *kin-1* and *kin-19,* it is also possible that

distinct kinases modulate miRISC in the dauer life history. This study lays the

foundation for future studies that can dissect the molecular relationship between these

genes and miRISC in *C. elegans* and other species.

Acknowledgments

We thank Karp lab members, Jennifer Schisa, and Steven Gorsich (Central Michigan

University) for their helpful advice throughout this project. We are grateful to BIO 546

students (Central Michigan University) who helped with pilot experiments. We thank

WormBase. Some strains were provided by the CGC, which is funded by NIH Office of

Research Infrastructure Programs (P40 OD010440). This project was funded by

CAREER 1652283 to XK from the National Science Foundation.

347
348
349
350

References:

Abbott, A. L., E. Alvarez-Saavedra, E. A. Miska, N. C. Lau, D. P. Bartel *et al.*, 2005 The
let-7 microRNA family members *mir-48, mir-84*, and *mir-241* function together to
regulate developmental timing in *Caenorhabditis elegans*. Developmental Cell 9:
403–414.

Alessi, A. F., V. Khivansara, T. Han, M. A. Freeberg, J. J. Moresco *et al.*, 2015 *Casein*
kinase II promotes target silencing by miRISC through direct phosphorylation of the
DEAD-box RNA helicase CGH-1. Proceedings of the National Academy of Sciences
201509499.

Alvarez-Saavedra, E., and H. R. Horvitz, 2010 Many families of *C. elegans* microRNAs
are not essential for development or viability. Current Biology 1–7.

Ambros, V., and H. R. Horvitz, 1984 Heterochronic mutants of the nematode
Caenorhabditis elegans. Science 226: 409–416.

Bartel, D. P., 2018 Metazoan MicroRNAs. Cell 173: 20–51.

Binti, S., R. V. Melinda, B. B. Joseph, P. T. Edeen, S. D. Miller *et al.*, 2022 A life cycle
alteration can correct molting defects in *Caenorhabditis elegans*. Dev Biol.

Brenner, S., 1974 The genetics of *Caenorhabditis elegans*. Genetics 77: 71–94.

Byerly, L., R. C. Cassada, and R. L. Russell, 1976 The life cycle of the nematode
Caenorhabditis elegans. I. Wild-type growth and reproduction. Developmental
Biology 51: 23–33.

Cassada, R. C., and R. L. Russell, 1975 The dauerlarva, a post-embryonic
developmental variant of the nematode *Caenorhabditis elegans*. Developmental
Biology 46: 326–342.

Deng, Y., K. L. Luo, D. D. Shaye, and I. Greenwald, 2019 A screen of the conserved
kinome for negative regulators of LIN-12 negative regulatory region ("NRR")-
missense activity in *Caenorhabditis elegans*. G3 Genes Genomes Genetics 9: 3567–
3574.

Edwards, F., G. Maton, N. Gareil, J. C. Canman, and J. Dumont, 2018 BUB-1 promotes
amphitelic chromosome biorientation via multiple activities at the kinetochore. Elife 7:
e40690.

Euling, S., and V. Ambros, 1996 Reversal of cell fate determination in *Caenorhabditis*
elegans vulval development. Development 122: 2507–2515.

Frédérick, P., and M. J. Simard, 2022 Regulation and different functions of the animal
microRNA-induced silencing complex. Wiley Interdiscip Rev RNA 13: e1701.

Fraser, A. G., R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann *et al.*,
2000 Functional genomic analysis of *C. elegans* chromosome I by systematic RNA
interference. Nature 408: 325–330.

Fry, A. M., L. O'Regan, S. R. Sabir, and R. Bayliss, 2012 Cell cycle regulation by the
NEK family of protein kinases. J. Cell Sci. 125: 4423–4433.

Galagali, H., and J. K. Kim, 2020 The multifaceted roles of microRNAs in differentiation.
Curr. Opin. Cell Biol. 67: 118–140.

Grishok, A., A. E. Pasquinelli, D. Conte, N. Li, S. Parrish *et al.*, 2001 Genes and
mechanisms related to RNA interference regulate expression of the small temporal
RNAs that control *C. elegans* developmental timing. Cell 106: 23–34.

Hada, K., M. Asahina, H. Hasegawa, Y. Kanaho, F. J. Slack *et al.*, 2010 The nuclear
receptor gene nhr-25 plays multiple roles in the *Caenorhabditis elegans*
heterochronic gene network to control the larva-to-adult transition. Developmental
Biology 344: 1100–1109.

Hand, S. C., D. L. Denlinger, J. E. Podrabsky, and R. Roy, 2016 Mechanisms of animal
diapause: recent developments from nematodes, crustaceans, insects, and fish.
American journal of physiology. Regulatory, integrative and comparative physiology
ajpregu.00250.2015.

Hansen, M. A., A. Dahal, T. A. Bernstein, C. Kohtz, S. Ali *et al.*, 2022 ztf-16 is a novel
heterochronic modulator that opposes adult cell fate in dauer and continuous life
histories in *Caenorhabditis elegans*. bioRxiv 2022.06.20.496913.

Huberdeau, M. Q., V. N. Shah, S. Nahar, J. Neumeier, F. Houle *et al.*, 2022 A specific
type of Argonaute phosphorylation regulates binding to microRNAs during *C. elegans*
development. Cell Rep. 41: 111822.

Jonas, S., and E. Izaurralde, 2015 Towards a molecular understanding of microRNA-
mediated gene silencing. Nat Rev Genet 16: 421–433.

Joseph, B. B., Y. Wang, P. Edeen, V. Lažetić, B. D. Grant *et al.*, 2020 Control of
clathrin-mediated endocytosis by NIMA family kinases. Plos Genet 16: e1008633.

Kalogeropoulos, N., C. Christoforou, A. J. Green, S. Gill, and N. R. Ashcroft, 2004 *chk-1*
is an essential gene and is required for an S-M checkpoint during early
embryogenesis. Cell Cycle 3: 1194–1198.

Kamath, R. S., and J. Ahringer, 2003 Genome-wide RNAi screening in *Caenorhabditis*
elegans. Methods 30: 313–321.

Karp, X., 2018 Working with dauer larvae. WormBook 2018: 1–19.

Karp, X., and V. Ambros, 2012 Dauer larva quiescence alters the circuitry of microRNA
pathways regulating cell fate progression in *C. elegans*. Development 139: 2177–
2186.

Karp, X., M. Hammell, M. C. Ow, and V. Ambros, 2011 Effect of life history on
microRNA expression during *C. elegans* development. RNA 17: 639–651.

Kim, T., M. W. Moyle, P. Lara-Gonzalez, C. D. Groot, K. Oegema *et al.*, 2015
Kinetochore-localized BUB-1/BUB-3 complex promotes anaphase onset in *C.*
elegans. J. Cell Biol. 209: 507–517.

Kim, W., R. S. Underwood, I. Greenwald, and D. D. Shaye, 2018 OrthoList 2: a new
comparative genomic analysis of human and *Caenorhabditis elegans* Genes.
Genetics 210: 445–461.

Leung, A. K. L., and P. A. Sharp, 2010 MicroRNA functions in stress responses.
Molecular Cell 40: 205–215.

Liu, Z., and V. Ambros, 1991 Alternative temporal control systems for hypodermal cell
differentiation in *Caenorhabditis elegans*. Nature 350: 162–165.

Meuti, M. E., R. Bautista-Jimenez, and J. A. Reynolds, 2018 Evidence that microRNAs
are part of the molecular toolkit regulating adult reproductive diapause in the
mosquito, Culex pipiens. PLoS ONE 13: e0203015.

Olejniczak, M., A. Kotowska-Zimmer, and W. Krzyzosiak, 2018 Stress-induced changes
in miRNA biogenesis and functioning. Cell Mol Life Sci 75: 177–191.

Parry, D. H., J. Xu, and G. Ruvkun, 2007 A Whole-Genome RNAi Screen for *C. elegans*
miRNA Pathway Genes. Curr Biol 17: 2013–2022.

Reinhart, B. J., F. J. Slack, M. Basson, A. E. Pasquinelli, J. C. Bettinger *et al.*, 2000 The
21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*.
Nature 403: 901–906.

Rogers, E., J. D. Bishop, J. A. Waddle, J. M. Schumacher, and R. Lin, 2002 The aurora
kinase AIR-2 functions in the release of chromosome cohesion in *Caenorhabditis*
elegans meiosis. J. Cell Biol. 157: 219–229.

Rougvie, A. E., and E. G. Moss, 2013 Developmental transitions in *C. elegans* larval
stages. Current topics in developmental biology 105: 153–180.

Schumacher, J. M., A. Golden, and P. J. Donovan, 1998 AIR-2: An Aurora/Ipl1-related protein kinase associated with chromosomes and midbody microtubules is required for polar body extrusion and cytokinesis in *Caenorhabditis elegans* embryos. J. Cell Biol. 143: 1635–1646.

Required for Polar Body Extrusion and Cytokinesis in *Caenorhabditis elegans* Embryos. J Cell Biology 143: 1635–1646.

Rual, J.-F., J. Ceron, J. Koreth, T. Hao, A.-S. Nicot *et al.*, 2004 Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. Genome Res. 14: 2162–2168.

Seydoux, G., and I. Greenwald, 1989 Cell autonomy of lin-12 function in a cell fate decision in *C. elegans*. Cell 57: 1237–1245.

Shah, V. N., J. Neumeier, M. Q. Huberdeau, D. M. Zeitler, A. Bruckmann *et al.*, 2023 *Casein kinase 1* and *2* phosphorylate Argonaute proteins to regulate miRNA-mediated gene silencing. EMBO Rep.

Sharp-Baker, H., and R.-H. Chen, 2001 Spindle checkpoint protein Bub1 Is required for kinetochore localization of Mad1, Mad2, Bub3, and Cenp-E, independently of its kinase activity. J. Cell Biol. 153: 1239–1250.

Shaye, D. D., and I. Greenwald, 2011 OrthoList: a compendium of *C. elegans* genes with human orthologs. PLoS ONE 6: e20085.

Sulston, J. E., and H. R. Horvitz, 1977 Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. Developmental Biology 56: 110–156.

Tops, B. B. J., R. H. A. Plasterk, and R. F. Ketting, 2006 The *Caenorhabditis elegans* Argonautes ALG-1 and ALG-2: almost identical yet different. Cold Spring Harb. Symp. Quant. Biol. 71: 189–194.

Vasquez-Rifo, A., G. Jannot, J. Armisen, M. Labouesse, S. I. A. Bukhari *et al.*, 2012 Developmental characterization of the microRNA-specific *C. elegans* Argonautes *alg-1* and *alg-2*. PLoS ONE 7: e33750.

Vowels, J. J., and J. H. Thomas, 1992 Genetic analysis of chemosensory control of dauer formation in *Caenorhabditis elegans*. Genetics 130: 105–123.

Wang, X., M. Liu, W. Li, C. D. Suh, Z. Zhu *et al.*, 2009 The function of a spindle checkpoint gene *bub-1* in *C. elegans* development. PLoS ONE 4: e5912.

Wilczynska, A., and M. Bushell, 2015 The complexity of miRNA-mediated repression. Cell death and differentiation 22: 22–33.

Wu, C., J. So, B. N. Davis-Dusenbery, H. H. Qi, D. B. Bloch *et al.*, 2011 Hypoxia
Potentiates MicroRNA-Mediated Gene Silencing through Posttranslational
Modification of Argonaute2. Mol Cell Biol 31: 4760–4774.

Yochem, J., V. Lažetić, L. Bell, L. Chen, and D. Fay, 2015 C. elegans NIMA-related
kinases NEKL-2 and NEKL-3 are required for the completion of molting. Dev Biol
398: 255–266.

Zinovyeva, A. Y., S. Bouasker, M. J. Simard, C. M. Hammell, and V. Ambros, 2014
Mutations in conserved residues of the *C. elegans* microRNA Argonaute ALG-1
identify separable functions in ALG-1 miRISC loading and target repression. PLoS
Genetics 10: e1004286.

Zinovyeva, A. Y., I. Veksler-Lublinsky, A. A. Vashisht, J. A. Wohlschlegel, and V. R.
Ambros, 2015 *Caenorhabditis elegans* ALG-1 antimorphic mutations uncover
functions for Argonaute in microRNA guide strand selection and passenger strand
disposal. Proceedings of the National Academy of Sciences 112: E5271-80.