A Histomorphometric and Computational Investigation of the Stabilizing Role of Pecti-
nate Ligaments in the Aqueous Outflow Pathway

Babak N. Safa[1#], Nina Sara Fraticelli Guzmán[2,3#], Guorong Li[4],

W. Daniel Stamer[4], Andrew J. Feola[1,3,5], and C. Ross Ethier[1,2,3*]

[1] Wallace H. Coulter Department of Biomedical Engineering,
Georgia Institute of Technology/Emory University, Atlanta, GA, USA
7
8 [2] George W. Woodruff School of Mechanical Engineering,
9 Georgia Institute of Technology, Atlanta, GA, USA
10
11 [3] Department of Ophthalmology, Emory University, Atlanta, GA, USA
12
13 [4] Department of Ophthalmology, Duke University, Durham, NC, USA
14
[5] Atlanta VA Center for Visual and Neurocognitive Rehabilitation,
Atlanta Veterans Affair Healthcare System, Decatur, GA, USA

#Shared co-first authors

***Corresponding author:**
Petit Biotechnology Building (IBB)
315 Ferst Drive, Room 2306
Atlanta, GA 30332-0363
ross.ethier@bme.gatech.edu
23

24 *Submitted to Journal of Biomechanical Engineering as an Original Research Article.*

Abstract

Murine models are commonly used to study glaucoma, the leading cause of irreversible blindness in humans. Blindness in glaucoma is associated with intraocular pressure (IOP), which is generated and regulated by the tissues of the aqueous outflow pathway. One of the main components of the aqueous humor pathway are the pectinate ligaments (PLs) that connect iris and trabecular meshwork at the anterior chamber angle. However, their role in helping maintain the biomechanical stability of murine aqueous outflow pathway is unknown, thus motivating this study. In this work, we conducted histomorphometric analysis and optical coherence tomographic-imaging-based finite element (FE) modeling on three age cohorts of C57BL/6 mice: 'young' (2-6 months), 'middle-aged' (11-16 months), and 'elderly' (29-32 months). We also evaluated the age-specific morphometric characteristics of the outflow pathway tissues, specifically the trabecular meshwork (TM) and Schlemm's canal (SC). Further, because of the known pressure-dependent narrowing of the SC lumen in humans and mice, we assessed the sensitivity of the SC lumen area (SCLA) to varying IOPs in age-specific FE models in a physiological range of TM and PL stiffness values. We found age-dependent changes in morphological characteristics of outflow tissues; notably, the PLs were more developed in older compared to younger mice. In addition, our FE analysis demonstrated that the patency of SC in the mouse is highly dependent on the presence of PLs, and that increased IOP caused SC collapse only when TM/PL stiffness was sufficiently low. In particular, young and middle-aged mice showed a lower susceptibility to SC collapse under elevated IOP compared to elderly mice. In conclusion, our study has elucidated the previously unexplored role of PLs in the aqueous outflow pathway, indicating their function in biomechanically supporting TM and SC under elevated IOP.

Keywords: Glaucoma, Biomechanics, IOP, Pectinate Ligament, Finite Element Method

Introduction

Glaucoma is a group of eye diseases that are collectively responsible for the majority of irreversible blindness worldwide (World Health Organization, 2019). Glaucoma is often associated with increased intraocular pressure (IOP), i.e., ocular hypertension (Safa et al., 2022c; Weinreb et al., 2014), and lowering of IOP helps preserve vision (Gordon et al., 2002; Heijl et al., 2002; Kass et al., 2002; Leske et al., 2007). Therefore, understanding the mechanisms by which IOP is regulated has great importance in developing new glaucoma treatments.

A critical determinant of IOP is the hydrodynamic resistance of the conventional outflow pathway tissues, including the inner wall of Schlemm's canal (SC) and trabecular meshwork (TM; Figure 1). Further, the biomechanical properties of these tissues are thought to play a key role in IOP homeostasis (Pedrigi et al., 2011; Stamer et al., 2015). For example, several studies have observed TM stiffening in glaucoma and mechanobiological factors are known to affect aqueous humor drainage across the inner wall of SC (Last et al., 2011; Overby et al., 2014; Vahabikashi et al., 2019; Wang et al., 2017). Another integral part of the outflow pathway in many species is the pectinate ligaments, yet, little is known about their mechanical properties and their role in ocular hypertension and the regulation of IOP (Bergmanson, 1985; De Geest et al., 1990; Flügel et al., 1991; McMenamin and Steptoe, 1991; Samuelson and Gelatt, 1984; Smith et al., 2001).

The pectinate ligaments (PLs), as their name implies, form a comb-like fibrous structure located at the angle of the anterior chamber, bridging between the iris and the TM. PL structure varies substantially between species (Simones et al., 1996), suggesting potential evolutionary significance for this tissue. For example, piscine PLs (or annular ligaments) tend to be more extensive than in mammals (Chen et al., 2008; Soules and Link, 2005). Notably, in primates, including humans, the PLs are less extensive, with a porous presentation that is visible by gonioscopy.

We hypothesize that the PLs play a role in mechanically stabilizing the TM and SC by

imposing mechanical tension that resists the normal pressure differential from the anterior

chamber to SC lumen. Our objective in this study was to test this hypothesis via a combination

of histology-based morphometry (histomorphometry), *in vivo* optical coherence tomography

(OCT) imaging, and computational modeling of the conventional outflow pathway tissues. All

work was carried out on mice (Li et al., 2023), commonly used to study aqueous humor dy-

namics.

Methods

We first histomorphometrically characterized key features of the primary outflow path-

way in mice of different ages. We then augmented this histomorphometric data with OCT im-

aging to create age-specific computational (finite element [FE]) models of the murine outflow

pathway from mice of different ages. Finally, we conducted forward FE analyses to investigate

the effect of PL structure and mechanics on the biomechanical response of TM/SC to IOP as a

function of age. The experimental procedures are described in detail elsewhere with a specific

focus on morphological and biological effect of aging on aqueous outflow parameters (Li et

al., 2023). Here, we pursue an in-depth analysis of morphology and biomechanics of the PLs.

Animals

All experimental procedures were approved by the Institutional Animal Care and Use

Committees (IACUC) at Duke University and observed the tenets of the Association for Re-

search in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic

and Vision Research. Three groups of wild-type C57BL/6 mice were used in this study, as

reported elsewhere (Li et al., 2023).

• Group 1 was used for histomorphometry and consisted of seven young (2-3 months), six

middle-aged (12-13 months), and seven elderly (29 months) mice. All mice for histomor-

phometry were male except for two females in the young cohort.

• Group 2 was used for high spatial resolution OCT imaging of the TM and SC ("TM-OCT")

and consisted of five young (4-6 months), seven middle-aged (11-16 months), and five

elderly (25-31 months) mice. These mice were all male except for two females in the young

cohort and two mice of unknown sex in the middle-aged cohort.

• Group 3 was used to evaluate iridial deformation via OCT imaging of the anterior segment

("Iris-OCT"). The young (3-6 months), middle-aged (9-11 months), and elderly (29-32

months) cohorts included six, eight, and six mice, respectively. In the middle-aged and

elderly cohorts, all animals were male except for two females in the middle-aged cohort,

while in the young cohort, all animals were female except for one male.

Histomorphometric analysis

Not all relevant anatomic features were visible on OCT images. Thus, we used sagittal

histological sections through the outflow tract of mice in Group 1 to obtain some of the ana-

tomical information needed to create the FE models. Anesthetized mice were euthanized by

decapitation, and the eyes were immediately harvested. Next, the eyes were immersion-fixed

in 4% paraformaldehyde + 1% glutaraldehyde at 4 °C overnight. The eyes were then bisected,

and the posterior segments and lenses were discarded. The anterior segments were cut into four

quadrants and each quadrant was embedded in Epon, after which 0.5 µm thick sagittal sections

were cut using a microtome, stained with methylene blue, and imaged (Axioplan2, Carl Zeiss

MicroImaging, Thornwood, NY, USA) with an image resolution of ca. 0.26 µm/pixel capturing

the outflow pathway tissues (Figure 1). For further details of these experimental procedures,

please see Li *et al.* (Li et al., 2023, 2021, 2019).

We manually identified and segmented the following features of interest from the re-

sulting images (Figure 1):

• TM cross-sectional area ("TM area"), defined as the area of TM tissue bounded by the

posterior terminus of Descemet's membrane and by the most posterior point of SC (Figure 1A).

- TM length, measured as the linear distance from the two (anterior-posterior) extremes of the TM (Figure 1B).

- PL thickness, obtained by identifying the most anterior PL and measuring its thickness perpendicular to the local axis of the PL (Figure 1C).

- Active TM cross-sectional area ("Active TM area"), defined as the TM area lying directly adjacent to the lumen of SC, excluding any parts of the TM adjacent to the sclera (Figure 1D).

- SC length, measured from the most posterior to the most anterior point of the SC lumen using a straight line (Figure 1E).

- PL attachment site, measured as the straight-line distance parallel to the line defining the SC length from the posterior end of SC to the identified PL (Figure 1F).

Given the subjectivity in identifying some of these features, the segmentations were independently carried out by four separate annotators who were masked to mouse age and sample identity. All segmentations were performed using ImageJ (Fiji distribution; various 2023 versions (Schindelin et al., 2012)). The results were then compiled, and the annotators met to discuss the segmentation results and resolve any disagreements. The annotators then performed a second round of segmentation under the same masking conditions. After the second round of segmentation and discussion, three images were removed from the data set (two from the young cohort and one from the elderly cohort) because the tissues were damaged during the histologic processing and could not be reliably segmented. As a result, a total of 6 histological images were analyzed per age cohort, corresponding to 24 possible measurements (4 annotators × 6 images) per feature of interest. We then calculated the average "active TM

thickness" in each image by dividing, for each annotator, the active TM area by the corresponding SC length, generating an additional 24 values per age cohort. Of note, due to the complex structure of the outflow tissues and inter-annotator variability, there were several unsuccessful annotations in some images. For instance, in the young cohort, a PL was not discernible in several samples (13 unmeasurable PL widths out of a total of 24), suggesting under-development of PLs in young animals (see below).

As mentioned above, it was necessary to combine the morphometric data with both OCT and histologic images to generate the FE models. We reasoned that such a combination would be most robust if we scaled all dimensions by a common length apparent in both OCT and histologic images; therefore, we selected SC anterior-posterior length ("SC length") as the common feature for this purpose. Hence, we calculated two ratios from histologic images: the "TM ratio," defined as (TM length)/(SC length), and the "PL ratio," defined as (PL attachment distance)/(SC length). We accounted for age variation by using age-specific TM ratios, i.e., we computed the ratio of average TM length divided by the average SC length for each age cohort, and we repeated the same process for the PL ratio. These ratios were later used in creating the FE models (see *Model geometry creation and FE mesh generation* section, below).

OCT Imaging

For OCT imaging, mice were anesthetized using an intraperitoneal injection of ketamine (100 mg/kg) and xylazine (10 mg/kg) plus a drop of 0.5% proparacaine applied to the mouse eye (Li et al., 2023). Mice were secured on a custom-made platform, and a single pulled glass microneedle filled with PBS was inserted into the anterior chamber of one eye. The microneedle was connected to both a manometric column to adjust IOP and a pressure transducer to continuously monitor IOP levels using PowerLab software.

Two types of OCT imaging were performed using a high-resolution spectral domain OCT system (SD-OCT Envisu R2200, Bioptigen Inc., Research Triangle Park, NC, USA). In

the first type of OCT image (TM-OCT, Group 2), we acquired sagittal scans focused on the

iridocorneal angle, including the TM and SC lumen, at IOPs of 10, 15, and 20 mmHg. We

identified the iridocorneal angle structures following our previously established techniques (Li

et al., 2019). In the second type of OCT image (Iris-OCT, Group 3), the OCT scan head was

placed near the corneal apex to acquire images of half of the anterior chamber at IOPs of 10,

15, and 20 mmHg. Additional IOP steps were also used but were not needed in this study (Lee

et al., 2021; Li et al., 2023).

Model geometry creation and FE mesh generation

We created age-specific models of the conventional outflow pathway tissues by com-

bining the morphometric measurements and the TM-OCT images. The TM-OCT images from

10 mmHg served as the reference state for creating all models (Figure 2A, D, and G). For each

age-specific model, we delineated the corneal and iridial margins in one representative TM-

OCT image (Figure 2B, E, and H) using the 2D spline function in SolidWorks (version 2022

Dassault Systèmes, Waltham, MA, USA).

The details of the iridocorneal angle tissues were not resolved in these OCT images,

and thus, as mentioned earlier, we relied on histomorphometry to reconstruct details of the

outflow pathway tissues as follows. Using the segmented SC lumen, we created the TM by

assuming that the TM's inner margin was offset from the inner wall of SC by the age-specific

average active TM thickness (Figure 2F, feature '*ii*'). This offset distance was measured normal

to the line defining SC length (Figure 2F, feature '*i*'). Then, to complete the TM profile, we

connected the ends of this inner TM margin to the cornea and root of the iris using linear ex-

trapolations (Figure 2F, features '*iv*' and '*v*', respectively). We used a constant TM thickness

(averaged active TM thickness), rather than a spatially-variable thickness, as a practical ap-

proximation to keep the model tractable.

We modeled the PLs as a diffuse (distributed) material rather than attempting to model

the complex, interconnected, and networked structure of the mouse PLs (Chalupa and Wil-

liams, 2008). We recognize that this homogenized approach to modeling the PLs is an approx-

imation, and we therefore created two additional models to assess its validity. In the first, we

did not include the PLs at all, while in the second we only included a single PL "beam" (the

anterior-most PL), as previously described (Li et al., 2019). For consistency with previous

work, we used the young model for these additional simulations (see supplementary Figure S2

for details). In both of these models, SC collapsed at low IOPs (IOP < 11 mmHg), which is

contradictory to observations during OCT imaging (Li et al., 2023, 2019). This simple result

suggested that treating the PL network as a diffuse (effective) material, coupling the iris and

TM, was the most physiologically-relevant option for our study, as discussed below.

The "distributed" PL material filled the iridocorneal angle, extending from the apex of

the angle to the anterior-most PL attachment site (Figure 2F, feature '*iii'*), as determined for

each age model using the "PL ratio" based on the histomorphometric analysis (Figure 1C and

2). The PL region was bounded by the posterior surface of the TM and the anterior surface of

the iris.

The final modeling step was to create a 3D CAD model by extruding the cornea, iris,

TM, and PL as separate bodies, each 10 µm in height. This approach was preferred over creat-

ing an axisymmetric model due to the plane strain assumption (see the *Boundary conditions*

section for further details).

To generate the FE mesh, we exported the CAD model as a STEP file (ASCII version

2) which was then imported into the Gmsh meshing software (version 4.11.1; (Geuzaine and

Remacle, 2009)). We tessellated the volume using TET10 quadratic elements with overall

mesh density set at 7.5 µm, confirmed to provide adequate numerical accuracy in a preliminary

mesh refinement study. To enhance numerical accuracy, we used higher mesh densities in se-

lect locations. Specifically, we decreased the elemental characteristic edge length in the TM by

a factor of 4 and in the PL region by a factor of 8. The meshes were generated using the Frontal

algorithm, and ten steps of smoothing and optimization were conducted to enhance mesh qual-

ity (Safa et al., 2022b).

Boundary conditions

We used a plane strain modeling approach, justified by the conventional outflow re-

gion's small size relative to its distance from the anterior chamber's axis of symmetry. Further,

we constrained the posterior (scleral) end of the model and the anterior corneoscleral surface

(Figure 3), which was justified based on the significantly higher stiffness of the sclera and

cornea relative to the key tissues of our model (see *Tissue constitutive models* section below).

In addition, the right (central) edge of the cornea and iris were constrained to move only in the

'y' direction (Figure 3), which allowed for anterior chamber deepening during IOP elevation.

Increasing IOP caused posterior displacement of the iris, as observed from the OCT

images of the anterior chamber (see supplementary Figure S1). We used the lower-resolution

OCT images (Iris-OCT) to extract iris deformations, which we then applied as age-specific

boundary conditions in the FE model. These OCT scans were from a different set of animals

(Group 3) than the high magnification OCT images (TM-OCT; Group 2) due to technical dif-

ficulties in maintaining a consistent OCT field of view at all IOP levels and in aligning the field

of view with the anterior chamber's central axis in the high-magnification TM-OCT scans.

Therefore, we measured the iris displacement along the 'y' axis of Iris-OCT images (Figure

S1; same definition as in Figure 2A for TM-OCT images) at IOP levels of 15 and 20 mmHg

relative to the configuration at 10 mmHg and prescribed this displacement in the FE model as

an input (see supplementary Figure S1 for more details).

The pressure in the anterior chamber was specified based on the IOP levels imposed

experimentally. For numerical modeling, IOP values were specified relative to the baseline

pressure of 10 mmHg, i.e., we imposed a relative IOP of 5 mmHg (corresponding to 15 mmHg

absolute) and of 10 mmHg (corresponding to 20 mmHg absolute). We set the posterior cham-

ber pressure (PCP) to zero (10 mmHg absolute) based on the results of a prior study evaluating

PCP as a data-fitting variable, which showed that PCP remained at 10 mmHg when IOP ex-

ceeded 10 mmHg (Lee et al., 2021). Finally, SC lumen pressure (SCLP) was calculated based

on IOP and the total resistance of the primary outflow pathway (Li et al., 2019), which indicated

SCLPs of 2 mmHg (12 mmHg absolute) and 4 mmHg (14 mmHg absolute) at IOP levels of 15

mmHg and 20 mmHg, respectively.

Finally, we imposed a contact boundary condition between the inner and outer walls of

SC, which occurred when SC collapsed, using a node-on-facet algorithm and a penalty method

to enforce contact (Maas et al., 2023). Specifically, we used an automatic penalty approach

with the scaling factor set at 100 times TM stiffness.

Tissue constitutive models

We used an incompressible neo-Hookean ("INH") constitutive relation to model the

cornea and iris.

$$\Psi_{\text{INH}}(\boldsymbol{C}) = \frac{E}{6}(\tilde{I}_1 - 3) + \frac{K}{2}\ln{(J)}^2 \tag{1}$$

where Ψ_{INH} is the Helmholtz free energy, $\boldsymbol{C}$ is the right Cauchy-Green strain tensor, $\tilde{I}_1$ is the

first invariant of the deviatoric right Cauchy-Green strain tensor, and $J = \det(\boldsymbol{F})$ is the Jaco-

bian of the deformation gradient tensor. The stiffness E was set to 2700 kPa (Li et al., 2019)

for the cornea/sclera and 96.1 kPa for the iris (Lee et al., 2021). The bulk modulus K was set

to 100 times the shear modulus to enforce incompressibility (Weiss et al., 1996).

Due to the fibrous structure of the PLs, we did not model them as incompressible; in-

stead, they were modeled as a homogenized compressible neo-Hookean material ("NH")

$$\Psi_{\text{NH}}(\boldsymbol{C}) = \frac{E}{4(1+v)}(I_1 - 3) - \frac{E}{2(1+v)}\ln(J)$$

$$+ \frac{vE}{2(1+v)(1-2v)}(\ln J)^2 \tag{2}$$

268   Here, I_1 is the first invariant of the right Cauchy-Green strain tensor, and v is Poisson's ratio,

269   which we set to zero due to the approximately parallel structure of the PLs noted above, as-

270   suming the PL beams behave as 1D elements lacking lateral contraction under axial tension.

271   Therefore,

$$\Psi_{\text{NH}}(\boldsymbol{C}) = \frac{E}{4}(I_1 - 3) - \frac{E}{2}\ln(J) \tag{3}$$

We assumed that the intrinsic stiffnesses of TM and PL were the same, based on their similar

collagenous composition, but adjusted the stiffness value for the PLs to account for the high

porosity of the PL network. In brief, we calculated the effective stiffness of the PL relative to

the TM using a linear relationship $E_{PL} = \beta E_{TM}$, where β is the solid fraction of the PL network.

We computed β from an analysis of scanning electron micrographs of rabbit PLs (Ramos et

al., 2017; Simones et al., 1996) since suitable high-resolution scanning electron micrographs

from mouse eyes do not exist to the best of our knowledge, obtaining $\beta = 2.3\%$ (supplemen-

tary Figure S3). We then scaled this value down by a factor of 10 for mouse, since frontal

sections through the mouse eye showed a more rarified PL network in mouse vs. rabbit (Cha-

lupa and Williams, 2008). The same value $\beta = 0.23\%$ was used for all ages.

Forward finite element simulations and data analysis

We solved the FE models using FEBio (version 4.3.0, (Maas et al., 2012)), and devel-

oped a custom Python class to automate the simulations and post-processing. We specifically

analyzed the deformations of aqueous outflow pathway tissues as a function of IOP in models

with different TM and PL stiffnesses, for which a key metric was SC lumen cross-sectional

area (SCLA). We varied E_{TM} in a physiological range (5-150 kPa) (Wang et al., 2017), with

E_{PL} calculated from E_{TM} as described above. We simulated 15 cases with varying E_{TM} and

E_{PL}.

To calculate SCLA, we used the *concave_hull* function (Shapely version 2.0.1, (Gillies,

2023)) to fit a concave hull shape to the coordinates of the nodes on the inner and outer walls

of SC (Figure 1). For consistency between models, we then computed a normalized SCLA,

defined as

$$nSCLA = \frac{SCLA}{SCLA_{@10 \text{ mmHg}}} \tag{4}$$

where $SCLA_{@10 \text{ mmHg}}$ is $SCLA$ at the reference IOP (10 mmHg).

Finally, for each age-specific model and set of TM/PL stiffness values, we assessed SC

lumen 'patency' using a quantity we term "Ultimate nSCLA," defined as the value of nSCLA

at IOP = 20 mmHg. An Ultimate SCLA value greater than one indicated a SC luminal area

expansion at 20 mmHg relative to SCLA at 10 mmHg; conversely, Ultimate SCLA values less

than one indicated an area reduction. In addition, as another measure of SC patency, we deter-

mined the IOP at which SC 'collapse' occurred, which was defined as the first state when

$nSCLA < 33\%$. These definitions were based on informed estimations of the state of SC as it

pertains to its capability to function within normal aqueous outflow, where collapsed regions

of SC would not be able to function normally.

Statistical analysis

We used a linear mixed effects (LME) model to study the effect of age on the morpho-

metric characteristics of the mouse outflow pathway (JMP, 2023). Specifically, we used 'age'

as the fixed effect and 'annotator' as a random effect. The significance threshold (α) was set at

5%. We also conducted a *post hoc* multiple comparison test on 'age' using Tukey's HSD test

with Bonferroni correction (critical $\alpha = 5\%/3$). In addition, we conducted one-way ANOVA

to test for the effect of age on TM- and PL-ratios ($\alpha = 5\%$), followed with *post hoc* Tukey's

HSD test (critical $\alpha = 5\%/3$). Statistical analysis was performed using JMP (Pro version 16,

SAS Institute Inc., Cary, NC, USA)

Results

Morphometrical characterization of TM and PL in C57BL/6 mice

Histomorphometric analysis indicated that all measured dimensions changed with age, except PL thickness (Figure 4 and Table 1; $p < 0.05$ for the effect of age), with no effect of annotator ($p > 0.05$, Table 1). *Post hoc* multiple comparison tests indicated that, compared to the young cohort, the elderly cohort had a larger TM area ($p = 0.014$), Active TM area ($p = 0.001$), and SC length ($p = 0.001$) (Figure 4, Table 1, and Table S1). In addition, the middle-aged cohort had a smaller SC length compared to the elderly cohort ($p < 0.0001$) (Figure 3, Table 1, and Table S1). Overall, there was a positive correlation between age and the size of the features.

Furthermore, the TM- and PL-ratios were significantly affected by age ($p < 0.05$; Table 2). More specifically, the elderly mice had smaller TM- and PL-ratios relative to the young cohort (Table S3), indicating a moderately posterior localization of TM and PLs relative to SC lumen in elderly animals.

Dependence of SC lumen area on TM/PL stiffness

All models (for all three age groups) demonstrated IOP-induced SC lumen collapse, as was also observed experimentally (Li et al., 2023, 2019, 2016, 2014). However, large TM/PL stiffness values caused unphysiological SC expansion at elevated IOP (Figure 5), which was not observed experimentally. More specifically, SC lumen area showed a highly nonlinear dependence on IOP and TM/PL stiffness in all age-specific models (Figure 5D). Interestingly, elevation of IOP did not guarantee reduction of SC luminal area. Instead, collapse only occurred at lower TM stiffnesses (Figure 5B, C, and D). At higher stiffnesses, SC lumen−counterintuitively−expanded with increasing IOP (Figure 5A and D). Further, the dependence of normalized SC cross-sectional lumen area, *nSCLA*, on IOP was not always even monotonic. For instance, in the middle-aged model, there was an initial phase of nSCLA reduction with increasing IOP followed by a sudden expansion of the SC at *ca.* IOP = 11 mmHg (Figure 5D

middle). But, as IOP further increased beyond 15 mmHg, SC luminal area gradually shrank. In

a similar fashion, albeit more gradually, such behavior was observed in the elderly model, and

to a lesser extent, in the young model (Figure 5D).

Patency of SC lumen

To investigate the patency of SC, we evaluated: (i) the Ultimate nSCLA for each age-

specific FE model vs. TM/PL stiffness (Figure 6A) and (ii) the IOP at which SC lumen first

collapsed for each age-specific model vs. TM/PL stiffness (Figure 6B). Recall that in our ap-

proach, increasing TM stiffness also proportionally increased PL stiffness; thus, we simply

denote stiffness as "TM/PL stiffness". We observed that the Ultimate nSCLA strongly de-

pended on TM/PL stiffness, with a lower stiffness giving a smaller Ultimate nSCLA (Figure

6A). In the young and middle-aged models, the Ultimate nSCLA was greater than 1 (expansion

of SC lumen) for cases with $E_{TM} > 46$ kPa ($E_{PL} > 0.11$ kPa) in the young model, and $E_{TM} >$

67 kPa ($E_{PL} > 0.15$ kPa) in the middle-aged model (Figure 6A). For the cases where the Ulti-

mate nSCLA was smaller than 1, which are physiologically relevant, the elderly model showed

a smaller Ultimate nSCLA for any given E_{TM} compared to both young and middle-aged mod-

els.

Further, when investigating the collapse IOP for different TM/PL stiffnesses (Figure

6B), the highest E_{TM} at which SC collapsed in the elderly model was $E_{TM} = 88$ kPa ($E_{PL} =$

0.20 kPa), which was higher than that of the young and middle-aged models $E_{TM} = 26$ kPa

($E_{PL} = 0.06$ kPa; Figure 6B). As expected, all models collapsed at a low IOP of 10~11 mmHg

at the lowest allowable E_{TM} stiffness of 5 kPa ($E_{PL} = 0.01$ kPa; Figure 6B). Yet, surprisingly,

at the same TM/PL stiffness levels, the elderly model collapsed at the lowest IOP compared to

both other models (Figure 6B), except for the limiting case of $E_{TM} = 5$ kPa ($E_{PL} = 0.01$ kPa),

where the models had a similar collapse IOPs.

Discussion

In this study, we investigated the stabilizing role of pectinate ligaments (PLs) on the

conventional outflow pathway in the mouse eye, a very common model for studying aqueous

humor dynamics. We showed that the stiffness and morphology of the PLs play a significant

role in stabilizing the TM, and hence, in the patency of SC lumen and how it deforms as IOP

changes. These observations reveal a previously unexplored role for the PLs in murine aqueous

humor dynamics. Indeed, in our computational models in which the PLs were not present, SC

was predicted to collapse at very low IOPs, in opposition to what is observed experimentally.

Considering the anatomy of the mouse TM (few beams and proportionately much thinner in

the interior-exterior direction than in larger eyes), it is perhaps not surprising that mechanical

stabilization of the TM in the mouse is needed to avoid complete SC collapse.

One interesting observation was the non-monotonic dependence of Schlemm's canal

luminal area on IOP (Figure 5D), which illustrates the complex biomechanics of the outflow

pathway. This behavior was particularly evident in the middle-aged model (Figure 5D, middle)

which illustrates two opposing factors impacting the configuration of outflow tissues: (i) IOP

elevation tends to collapse SC lumen by outwards displacement of the TM, whereas (ii) poste-

rior iridial movement with increasing IOP tends to pull the TM inward and open SC. The com-

plex shapes of the specific curves depend on TM stiffness, PL stiffness, and geometric factors

such as TM thickness and PL anterior-posterior extent (Figure 5 A-C). The complexity of these

effects is reinforced by observing the slope discontinuities in the nSCLA vs. IOP curves, which

were due to progressively larger contact between the inner and outer walls of SC. In future

work, we should also investigate the role of septae in the SC lumen (Bentley et al., 2016),

known to be present in human eyes but not yet reported in the murine eye.

Although PL biomechanics has not been the focus of many investigations, PL morphol-

ogy has been studied in a variety of domestic mammals, including dogs, horses, and rabbits,

via scanning electron microscopy (SEM) and other types of histology (Chen et al., 2008; Simones et al., 1996). PL morphology varies between species via differences in PL width, length, spacing, and insertion angle into the TM. Furthermore, PL morphology also varies locally within the same eye. One factor suggested to influence PL structure is age (Simones et al., 1996). While the effect of aging on PLs was not a primary focus of this work (see (Li et al., 2023)), our models were based on images from different aged animals. Therefore, it is worth exploring our results from that perspective.

Our histomorphometric analysis demonstrated a dependence of structural features on age (Figure 2 and Table 1) with certain features becoming more prominent with age (Figure 2), as expected. For instance, in the young cohort, the PL was relatively underdeveloped. In fact, we were unable to observe PLs in 13 out of 24 annotations from young eyes. Additionally, age also affected the TM and PL ratios (Table 2), with older eyes having larger SCs, and thus, smaller TM and PL ratios (Figure 2). We hypothesize these differences might be related to development of more prominent PLs with age, thereby providing more support to the TM, preventing SC collapse in aged eyes.

In addition, our simulations indicated a connection between iris deformation and TM/PL stiffness (Figure 5A-C), which was demonstrated by differences in the slope of the iris midline at the pupillary (right) edge of our models (Figure 5A-C). This observation suggests a connection between iris deformations and aging, which can influence the association between anterior chamber deepening (AC deepening) and IOP. We know AC deepening significantly affects outflow facility and also can artificially elevate facility in experiments (Boussommier-Calleja et al., 2015; Brubaker, 1975; Ellingsen and Grant, 1971). Thus, it would be worth exploring the connection between TM/PL stiffness and iris deformations more closely under, for example, various physiologically occurring iris contractions (e.g., at different light levels (Safa et al., 2022a)), and at different ages to study connections between iris deformation, age, and

outflow pathway function.

As a further age-dependent observation, our simulations indicated a strong effect of age

on the biomechanical properties of outflow tissues. For example, by comparing modeled SCLA

vs. experimentally observed SCLA changes with IOP, a significant stiffening of TM was ob-

served with age (Li et al., 2023). Of note, our predicted TM stiffness for young mice (~29 kPa

(Li et al., 2023)) was consistent with previous results on young mice (Li et al., 2019). We also

predicted that elderly eyes had the smallest Ultimate nSCLA for a given TM/PL stiffness (Fig-

ure 6A). Overall, these observations indicate an increased resistance to IOP-induced defor-

mations of outflow tissues with age, due to a combination of changes in biomechanical prop-

erties and tissue geometry.

It is important to note that we here used quadratic finite elements (TET10) that have

superior performance in modeling incompressible materials (Bonet and Wood, 1997) than the

linear elements that we previously used (TET4; (Li et al., 2019)). This element type was essen-

tial for making comparisons between different FE models in this study and should be used as

a suitable element type for future modeling work in studying outflow tissue biomechanics.

It is worth noting this study had some limitations. First, due to a lack of information

about PL biomechanical properties, we were forced to estimate the PL stiffness based on in-

formation about PL porosity in a rabbit eye which required several assumptions. Fortunately,

we kept the PL to TM stiffness ratio consistent between models, thus eliminating variability in

this ratio as a source of difference between models. Nonetheless, more direct estimates/meas-

urements of PL biomechanics properties, while challenging, should be undertaken in future.

Second, because of the diminutive size of the mouse eye, the manual positioning of the OCT

scan head, and IOP-induced changes in ocular geometry, there was inevitable shifting of the

relative position of the scan probe and the eye between scans of the same eye at different IOPs.

This added variability to our analysis, which was partially addressed by using lower magnifi-

cation Iris-OCT images for certain parts of the modeling process. In the future, hardware to

accurately position the OCT scan head might improve the repeatability of high-resolution OCT

images. Finally, although sex differences are likely important in glaucoma (Douglass et al.,

2023; Lee et al., 2021), due to the scarcity of female samples in our data set, we did not consider

sex as a biological variable in our analysis. Therefore, caution is advised interpreting these

results for both sexes.

In conclusion, our simulations show that PL biomechanics plays an essential role in

maintaining stability of the outflow pathway tissues in the mouse eye. In the future, it will be

important to directly characterize PLs' biomechanical properties, preferably by direct biome-

chanical testing of PL beams, e.g., using atomic force microscopy. We note that PLs are less

prominent in healthy human eyes and may thus be less clinically important than the situation

in pre-clinical studies in mouse eyes. However, structures like pectinate ligaments can persist

(see e.g. Figure 11 in (Anderson, 1981)) in human eyes with developmental pathologies leading

to congenital (infantile) glaucoma, and it would be of interest to evaluate their role in such

cases.

Conflict of interest

454   The authors declare no conflicts of interest.

Acknowledgments

We gratefully acknowledge support from The BrightFocus Foundation (postdoctoral

fellowship G2021005F, BNS), NIH (K99EY035360 (BNS), T32EY007092 (NSFG),

R01EY030871 (AJF), P30EY006360, P30EY005722, R01EY031710 (CRE and WDS),

5R01EY030124 (WDS) the Alfred P. Sloan Foundation G-2019-11435 (NSFG), the Georgia

Research Alliance (CRE), and Department of Veterans Affairs Rehab R&D Service Career

Development Awards (AJF; CDA-2; RX002342).

References

Anderson, D.R., 1981. The development of the trabecular meshwork and its abnormality in primary infantile glaucoma. Trans. Am. Ophthalmol. Soc. 79, 458–485.

Bentley, M.D., Hann, C.R., Fautsch, M.P., 2016. Anatomical Variation of Human Collector Channel Orifices. Invest. Ophthalmol. Vis. Sci. 57, 1153–1159. https://doi.org/10.1167/iovs.15-17753

Bergmanson, J.P., 1985. The anatomy of the rabbit aqueous outflow pathway. Acta Ophthalmol. (Copenh.) 63, 493–501. https://doi.org/10.1111/j.1755-3768.1985.tb05234.x

Bonet, J., Wood, R.D., 1997. Nonlinear Continuum Mechanics for Finite Element Analysis.

Boussommier-Calleja, A., Li, G., Wilson, A., Ziskind, T., Scinteie, O.E., Ashpole, N.E., Sherwood, J.M., Farsiu, S., Challa, P., Gonzalez, P., Downs, J.C., Ethier, C.R., Stamer, W.D., Overby, D.R., 2015. Physical Factors Affecting Outflow Facility Measurements in Mice. Invest. Ophthalmol. Vis. Sci. 56, 8331–8339. https://doi.org/10.1167/iovs.15-17106

Brubaker, R.F., 1975. The effect of intraocular pressure on conventional outflow resistance in the enucleated human eye. Invest. Ophthalmol. Vis. Sci. 14, 286–292.

Chalupa, L.M., Williams, R.W., 2008. Eye, retina, and visual system of the mouse, Eye, retina, and visual system of the mouse. Boston Review, Cambridge, MA, US.

Chen, C.-C., Yeh, L.-K., Liu, C.-Y., Kao, W.W.-Y., Samples, J.R., Lin, S.-J., Hu, F.-R., Wang, I.-J., 2008. Morphological Differences between the Trabecular Meshworks of Zebrafish and Mammals. Curr. Eye Res. 33, 59–72. https://doi.org/10.1080/02713680701795026

De Geest, J.P., Lauwers, H., Simoens, P., De Schaepdrijver, L., 1990. The morphology of the equine iridocorneal angle: a light and scanning electron microscopic study. Equine Vet. J. Suppl. 30–35. https://doi.org/10.1111/j.2042-3306.1990.tb04708.x

Douglass, A., Dattilo, M., Feola, A.J., 2023. Evidence for Menopause as a Sex-Specific Risk Factor for Glaucoma. Cell. Mol. Neurobiol. 43, 79–97. https://doi.org/10.1007/s10571-021-01179-z

Ellingsen, B., Grant, M., 1971. The Relationship of Pressure and Aqueous Outflow in Enucleated Human Eyes. Invest. Ophthalmol. Vis. Sci. 10, 430–437.

Flügel, C., Tamm, E., Lütjen-Drecoll, E., 1991. Different cell populations in bovine trabecular meshwork: an ultrastructural and immunocytochemical study. Exp. Eye Res. 52, 681–690. https://doi.org/10.1016/0014-4835(91)90020-f

Geuzaine, C., Remacle, J.-F., 2009. Gmsh: A 3-D finite element mesh generator with built-in pre- and post-processing facilities. Int. J. Numer. Methods Eng. 79, 1309–1331. https://doi.org/10.1002/nme.2579

Gillies, S., 2023. The Shapely User Manual — Shapely 2.0.1 documentation [WWW Document]. URL https://shapely.readthedocs.io/en/stable/manual.html (accessed 10.9.23).

JMP, 2023. Mixed Model Report and Options [WWW Document]. URL https://www.jmp.com/support/help/en/17.2/index.shtml#page/jmp/mixed-model-report-and-options.shtml (accessed 10.16.23).

Last, J.A., Pan, T., Ding, Y., Reilly, C.M., Keller, K., Acott, T.S., Fautsch, M.P., Murphy, C.J., Russell, P., 2011. Elastic modulus determination of normal and glaucomatous human trabecular meshwork. Invest. Ophthalmol. Vis. Sci. 52, 2147–2152. https://doi.org/10.1167/iovs.10-6342

Lee, C., Li, G., Stamer, W.D., Ethier, C.R., 2021. In vivo estimation of murine iris stiffness using finite element modeling. Exp. Eye Res. 202, 108374. https://doi.org/10.1016/j.exer.2020.108374

Li, G., Farsiu, S., Chiu, S.J., Gonzalez, P., Lütjen-Drecoll, E., Overby, D.R., Stamer, W.D.,

2014. Pilocarpine-Induced Dilation of Schlemm's Canal and Prevention of Lumen Collapse at Elevated Intraocular Pressures in Living Mice Visualized by OCT. Invest. Ophthalmol. Vis. Sci. 55, 3737–3746. https://doi.org/10.1167/iovs.13-13700

Li, G., Lee, C., Agrahari, V., Wang, K., Navarro, I., Sherwood, J.M., Crews, K., Farsiu, S., Gonzalez, P., Lin, C.-W., Mitra, A.K., Ethier, C.R., Stamer, W.D., 2019. In vivo measurement of trabecular meshwork stiffness in a corticosteroid-induced ocular hypertensive mouse model. Proc. Natl. Acad. Sci. 116, 1714–1722. https://doi.org/10.1073/pnas.1814889116

Li, G., Lee, C., Read, A.T., Wang, K., Ha, J., Kuhn, M., Navarro, I., Cui, J., Young, K., Gorijavolu, R., Sulchek, T., Kopczynski, C., Farsiu, S., Samples, J., Challa, P., Ethier, C.R., Stamer, W.D., 2021. Anti-fibrotic activity of a rho-kinase inhibitor restores outflow function and intraocular pressure homeostasis. eLife 10, e60831. https://doi.org/10.7554/eLife.60831

Li, G., Mukherjee, D., Navarro, I., Ashpole, N.E., Sherwood, J.M., Chang, J., Overby, D.R., Yuan, F., Gonzalez, P., Kopczynski, C.C., Farsiu, S., Stamer, W.D., 2016. Visualization of conventional outflow tissue responses to netarsudil in living mouse eyes. Eur. J. Pharmacol. 787, 20–31. https://doi.org/10.1016/j.ejphar.2016.04.002

Li, G., van Batenburg-Sherwood, J., Safa, B.N., Fraticelli Guzmán, N.S., Wilson, A., Bahrani Fard, M.R., Choy, K., De Leso, M.L., J. Serena, C., Feola, A.J., Weisz, T., Kuhn, M., Bowes Rickman, C., Farsiu, S., Ross Ethier, C., Stamer, W.D., 2023. Aging and intraocular pressure homeostasis in mice. Submitt. BioRxiv.

Maas, S., Weiss, J.A., Ateshian, G.A., 2023. FEBio User's Manual 4.1.

Maas, S.A., Ellis, B.J., Ateshian, G.A., Weiss, J.A., 2012. FEBio: Finite Elements for Biomechanics. J. Biomech. Eng. 134, 011005–011005. https://doi.org/10.1115/1.4005694

McMenamin, P.G., Steptoe, R.J., 1991. Normal anatomy of the aqueous humour outflow system in the domestic pig eye. J. Anat. 178, 65–77.

Overby, D.R., Zhou, E.H., Vargas-Pinto, R., Pedrigi, R.M., Fuchshofer, R., Braakman, S.T., Gupta, R., Perkumas, K.M., Sherwood, J.M., Vahabikashi, A., Dang, Q., Kim, J.H., Ethier, C.R., Stamer, W.D., Fredberg, J.J., Johnson, M., 2014. Altered mechanobiology of Schlemm's canal endothelial cells in glaucoma. Proc. Natl. Acad. Sci. 111, 13876–13881. https://doi.org/10.1073/pnas.1410602111

Pedrigi, R.M., Simon, D., Reed, A., Stamer, W.D., Overby, D.R., 2011. A model of giant vacuole dynamics in human Schlemm's canal endothelial cells. Exp. Eye Res. 92, 57–66. https://doi.org/10.1016/j.exer.2010.11.003

Ramos, M.F., Teixeira, L., Brandt, C.R., Auyeung-Kim, D., 2017. Ocular Immunopathology, in: Parker, G.A. (Ed.), Immunopathology in Toxicology and Drug Development: Volume 2, Organ Systems, Molecular and Integrative Toxicology. Springer International Publishing, Cham, pp. 695–762. https://doi.org/10.1007/978-3-319-47385-7_14

Safa, B.N., Bahrani Fard, M.R., Ethier, C.R., 2022a. In vivo biomechanical assessment of iridial deformations and muscle contractions in human eyes. J. R. Soc. Interface 19, 20220108. https://doi.org/10.1098/rsif.2022.0108

Safa, B.N., Bleeker, A., Berdahl, J.P., Ethier, C.R., 2022b. The Effects of Negative Periocular Pressure on Biomechanics of the Optic Nerve Head and Cornea: A Computational Modeling Study. https://doi.org/10.1101/2022.05.17.492338

Safa, B.N., Wong, C.A., Ha, J., Ethier, C.R., 2022c. Glaucoma and biomechanics. Curr. Opin. Ophthalmol. https://doi.org/10.1097/ICU.0000000000000829

Samuelson, D.A., Gelatt, K.N., 1984. Aqueous outflow in the beagle. I. Postnatal morphologic development of the iridocorneal angle: pectinate ligament and uveal trabecular meshwork. Curr. Eye Res. 3, 783–794. https://doi.org/10.3109/02713688409000790

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

560   S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V.,
561   Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for bio-
562   logical-image analysis. Nat. Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

563 Simones, P., De Geest, J.P., Lauwers, H., 1996. Comparative morphology of the pectinate lig-
564   aments of domestic mammals, as observed under the dissecting microscope and the
565   scanning electron microscope. J. Vet. Med. Sci. 58, 977–982.
566   https://doi.org/10.1292/jvms.58.10_977

567 Smith, R.S., Zabaleta, A., Savinova, O.V., John, S.W., 2001. The mouse anterior chamber an-
568   gle and trabecular meshwork develop without cell death. BMC Dev. Biol. 1, 3.
569   https://doi.org/10.1186/1471-213x-1-3

570 Soules, K., Link, B., 2005. Morphogenesis of the anterior segment in the zebrafish eye. BMC
571   Dev. Biol. 5, 12. https://doi.org/10.1186/1471-213X-5-12

572 Stamer, W.D., Braakman, S.T., Zhou, E.H., Ethier, C.R., Fredberg, J.J., Overby, D.R., Johnson,
573   M., 2015. Biomechanics of Schlemm's canal endothelium and intraocular pressure re-
574   duction. Prog. Retin. Eye Res. 44, 86–98. https://doi.org/10.1016/j.pret-
575   eyeres.2014.08.002

576 Vahabikashi, A., Gelman, A., Dong, B., Gong, L., Cha, E.D.K., Schimmel, M., Tamm, E.R.,
577   Perkumas, K., Stamer, W.D., Sun, C., Zhang, H.F., Gong, H., Johnson, M., 2019. In-
578   creased stiffness and flow resistance of the inner wall of Schlemm's canal in glauco-
579   matous human eyes. Proc. Natl. Acad. Sci. U. S. A. 116, 26555–26563.
580   https://doi.org/10.1073/pnas.1911837116

581 Wang, K., Read, A.T., Sulchek, T., Ethier, C.R., 2017. Trabecular meshwork stiffness in glau-
582   coma. Exp. Eye Res. 158, 3–12. https://doi.org/10.1016/j.exer.2016.07.011

583 Weinreb, R.N., Aung, T., Medeiros, F.A., 2014. The Pathophysiology and Treatment of Glau-
584   coma. JAMA 311, 1901–1911. https://doi.org/10.1001/jama.2014.3192

585 Weiss, J.A., Makerc, B.N., Govindjeed, S., 1996. Finite element implementation of incom-
586   pressible , isotropic hyperelasticity transversely 7825.

587 World Health Organization, 2019. World report on vision, World health Organisation.
588
589

Table 1: Results of the linear mixed effects model (LMM) testing on the histomorphometric characterization of the C57BL/6 mouse outflow pathway. The columns provide F-statistics for the fixed effect ('age'; **Bold**: $p < 0.05$), and Wald p-value for the random factor ('annotators'), which tests the null hypothesis that the random effect was significantly different from zero (*Wald p-value < 0.05*) (JMP, 2023).

Feature of interest	Age		Annotators
	F-statistics	*p-value*	*Wald p-value*
TM Area (μm²)	$f(2, 66) = 4.34$	**0.017**	0.419
TM length (μm)	$f(2, 66) = 3.77$	**0.028**	0.523
PL thickness (μm)	$f(2, 41.3) = 0.97$	0.389	0.410
Active TM area (μm²)	$f(2, 66) = 6.95$	**0.002**	0.479
SC length (μm)	$f(2, 65.1) = 12.40$	**< 0.0001**	0.911
PL attachment site (μm)	$f(2, 45.2) = 3.36$	**0.044**	0.082

Table 2: Ratios of the features measured from histomorphometric analysis used for creating the FE models. (*$p < 0.05$)

	Young	Middle-aged	Elderly	F-statistics	p-value
TM-ratio (TM length/SC length)	1.40 ± 0.26	1.35 ± 0.43	1.16 ± 0.19	$f(2,68)$ = 3.86	**0.026***
PL-ratio (PL attachment site location/SC length)	1.36 ± 0.46	1.11 ± 0.36	1.00 ± 0.25	$f(2,45)$ = 3.83	**0.029***

593 **Figure 1:** A representative sagittal section (from a young mouse) including the conventional outflow
594 pathway tissues, showing features that were annotated (red lines) for morphometric characterization.
595 TM = trabecular meshwork, PL = pectinate ligament, SC = Schlemm's canal, CB = ciliary body. Scale
596 bar = 50 µm.

599
600
Figure 2: TM-OCT images, outlined features, and FE meshes for the (A-C) 'young', (B-F) 'middle-
aged', and (G-I) 'elderly' models. Panels A, D and G show OCT images acquired at an IOP of 10 mHg
with the lumen of SC overlain in blue. Panels B, E and H show key structures segmented from OCT
images or deduced from histomorphometric analysis (red overlaid lines), while panels C, F, and I show
the resulting FE meshes. (J) Demonstration of a set of features used to create models, with emphasis on
features based on histomorphometric analysis, including: (*i*) the line connecting the posterior and ante-
rior ends of SC, (*ii*) an offset to create the inner margin of the TM based on the contour of the inner
wall of SC, (*iii*) the age-specific calculated distance of the posterior end of SC to the anterior-most PL
attachment site, (*iv*) a linear interpolation of the TM inner margin from the PL attachment site to cornea,
(*v*) a linear interpolation of the inner TM margin connecting the posterior end of the TM to the apex of
the irideocorneal angle, and (*vi*) the age-specific calculated TM length. Colors in C, F, and I: orange =
cornea and sclera; yellow = iris and other uveal tissue; blue = pectinate ligament region; purple = TM.
Scale bar = 100 µm.

Figure 3: Boundary conditions for the FE model. Here, the young model is used as an example for
illustrative purposes. Note the colored lines indicating the location where pressure loads were applied:
SCLP = SC lumen pressure, IOP = intraocular pressure, PCP = posterior chamber pressure, and Δy =
prescribe displacement of iris. Pinned support means no displacement along the 'x' and 'y' directions;
rolling support indicates no displacement along the 'x' direction. Scale bar = 100 µm.

Figure 4: Results of the histomorphometric analysis of mouse outflow pathway tissue dimensions. For each feature, the data is shown using violin plots for each age cohort, with the mean indicated by a short black horizontal line. The results from different annotators are shown with colored dots, where each color corresponds to a unique annotator. Statistically significant differences (after Bonferroni correction) are shown using solid horizontal lines ($p < \alpha/3 = 0.017$), and trending differences are also shown using dashed horizontal lines ($p < 0.05$).

631
632 **Figure 5**: The results of forward FE simulations for each age-specific model, showing representative simulation results for the (A) young, (B) middle-aged, and
633 (C) elderly models. In addition, (D) normalized SC lumen area (nSCLA; SC area normalized by the SCLA at 10 mmHg) vs. IOP is plotted vs. E_{TM}, in the range
634 of 5 kPa to 150 kPa. Interestingly, SCLA was not monotonically reduced by increasing IOP in the cases with higher E_{TM}, explainable by tension on the TM
635 delivered by the PL from the posterior deformations of the iris. Additionally, in the middle-aged simulations, there was an interesting rapid initial increase in
636 SCLA followed by a gradual decrease, which was not evident in the simulations in the other age-specific models.

Figure 6: The patency of the SC lumen was characterized in each age-specific model by evaluating (A) the dependence of Ultimate nSCLA (nSCLA at 20 mmHg) on TM stiffness, and (B) the IOP at which SC lumen collapsed vs. TM stiffness. Note that in the young and middle-aged models SC collapse (B) did not occur in cases with stiffness higher that 26 kPa, which is the reason for the continuation of the associated curves beyond that point.

Table S1: Results of the multiple comparison test on the 'age' factor on the histomorphometric characterization of the C57BL/6 mouse outflow pathway. (*p<0.05, **p<0.017*)

Feature of interest	Age comparison		p-value
TM Area (µm²)	Young	Middle-aged	0.575
	Young	Elderly	**0.014****
	Middle-aged	Elderly	0.149
TM length (µm)	Young	Middle-aged	0.243
	Young	Elderly	0.513
	Middle-aged	Elderly	**0.022***
PL thickness (µm)	Young	Middle-aged	0.445
	Young	Elderly	0.417
	Middle-aged	Elderly	1.000
Active TM area (µm²)	Young	Middle-aged	0.386
	Young	Elderly	**0.001****
	Middle-aged	Elderly	0.055
SC length (µm)	Young	Middle-aged	0.809
	Young	Elderly	**0.001****
	Middle-aged	Elderly	**< 0.001****
PL attachment site (µm)	Young	Middle-aged	0.229
	Young	Elderly	0.861
	Middle-aged	Elderly	**0.040***

Table S2: Summary statistics for the histomorphometric characterization of C57BL/6 mouse outflow pathway features by age cohort. Data is shown as mean and 95% confidence interval (95% CI).

Feature of interest	Age	Mean	95% CI
TM Area (μm^2)	Young	1445.49	[1311.43, 1579.56]
	Middle-aged	1619.64	[1311.58, 1927.71]
	Elderly	1946.82	[1638.26, 2255.38]
TM length (μm)	Young	194.29	[182.89, 205.70]
	Middle-aged	177.86	[163.05, 192.68]
	Elderly	205.50	[187.07, 223.94]
PL thickness (μm)	Young	4.46	[3.72, 5.21]
	Middle-aged	5.59	[3.77, 7.41]
	Elderly	5.77	[4.28, 7.26]
Active TM area (μm^2)	Young	1159.28	[1012.10, 1306.45]
	Middle-aged	1364.23	[1073.60, 1654.86]
	Elderly	1728.41	[1490.55, 1966.28]
SC length (μm)	Young	142.86	[131.86, 153.86]
	Middle-aged	137.39	[122.86, 151.92]
	Elderly	177.58	[164.61, 190.55]
PL attachment site (μm)	Young	162.95	[140.77, 185.13]
	Middle-aged	139.49	[127.86, 151.13]
	Elderly	171.07	[150.02, 192.12]

Table S3: Results of multiple comparison statistical testing on TM- and PL-ratios. (*p<0.05, **p<0.017)

Feature of interest	Age		p-value
TM-ratio	Young	Elderly	**0.030***
	Young	Middle-aged	0.094
	Middle-aged	Elderly	0.887
PL-ratio	Young	Elderly	**0.023***
	Young	Middle-aged	0.144
	Middle-aged	Elderly	0.620

647
648

649
Figure S1: Iris-OCT images, used to evaluate the IOP-induced deformations of the iris. We drew a
line segment (cyan) along the midline of the cornea (yellow lines), starting at the apex of the iridocor-
neal angle and extending to the edge of the cornea typically visible in TM-OCT images. We then con-
structed a perpendicular line segment connecting the end of the cyan line to the midline of the iris
(purple). We determined the length of this purple line segment along the 'y' axis (see IOP 10 mmHg,
the first row, for the definition of 'y' axis) for different IOPs, and based on the change in the length of
the line we evaluated iris displacement in the FE model.

Figure S2: Computed tissue deformations vs. IOP in three cases with different PL structures. **(A)** No PL, **(B)** a single PL at the anterior end of the PL region, and **(C)** the homogenized (diffuse) PL model described in the main text. The top row shows that, in the absence of a PL, SC collapse is predicted even at low values of IOP (< 15 mmHg); such collapse is not observed experimentally. The second row shows that a single PL is also not sufficient to prevent collapse at low IOP, especially when the PL is located at the anterior aspect of the TM, which is often is. Only the diffuse (distributed) model of the PL ligaments avoids SC collapse at low IOPs. Colors separate tissues (e.g., PL is blue)

Figure S3: Description of the technique used to estimate PL solid fraction, β. **(A)** A Scanning electron
micrograph of rabbit PL (Simones et al., 1996) with the PLs outlined by a dashed rectangle (red). White
scale bar = 200 µm. **(B)** The radii of the PLs were determined by manually drawing line segments (red)
in ImageJ, and the mean PL beam radius (r_{avg}) was computed.

The solid fraction within the relevant region of interest (ROI) was then determined by:

(i) Projecting the red rectangle in panel (A) into the plane of the page by a distance L, deter-
mined from a histological image of the New Zealand white rabbit angle (Ramos et al.,
2017), showing the distance from the iridocorneal angle to the most anterior PL as $L =$
$455 \, \mu m$.

(ii) Computing $\beta = \sum V_{\mathrm{PL}} / V_{\mathrm{ROI}}$, where N is the number of PLs annotated in panel B ($N =$
15); $\sum V_{\mathrm{PL}} = N \times \pi r_{avg}{}^2 H$ is the total volume of the PLs within the ROI, assuming cylin-
drically-shaped PLs; and $V_{\mathrm{ROI}} = H \times W \times L$ is the volume of the ROI.

This procedure yielded $\beta = 2.3\%$, which is clearly only an approximate estimate and should be
refined in future work.