1 **A Histomorphometric and Computational Investigation of the Stabilizing Role of**
2 **Pectinate Ligaments in the Aqueous Outflow Pathway**

3 Babak N. Safa[1#], Nina Sara Fraticelli Guzmán[2,3#], Guorong Li[4],

4 W. Daniel Stamer[4], Andrew J. Feola[1,3,5], and C. Ross Ethier[1,2,3*]

[1] Wallace H. Coulter Department of Biomedical Engineering,
Georgia Institute of Technology/Emory University, Atlanta, GA, USA

[2] George W. Woodruff School of Mechanical Engineering,
Georgia Institute of Technology, Atlanta, GA, USA

[3] Department of Ophthalmology, Emory University, Atlanta, GA, USA

[4] Department of Ophthalmology, Duke University, Durham, NC, USA

[5] Atlanta VA Center for Visual and Neurocognitive Rehabilitation,
Atlanta Veterans Affair Healthcare System, Decatur, GA, USA

#Shared co-first authors
Underlined text indicates surname.

*Corresponding author:
Petit Biotechnology Building (IBB)
315 Ferst Drive, Room 2306
Atlanta, GA 30332-0363
ross.ethier@bme.gatech.edu
24

Abstract

Murine models are commonly used to study glaucoma, the leading cause of irreversible blindness. Glaucoma is associated with elevated intraocular pressure (IOP), which is regulated by the tissues of the aqueous outflow pathway. In particular, pectinate ligaments (PLs) connect the iris and trabecular meshwork (TM) at the anterior chamber angle, with an unknown role in maintenance of the biomechanical stability of the aqueous outflow pathway, thus motivating this study. We conducted histomorphometric analysis and optical coherence tomography-based finite element (FE) modeling on three cohorts of C57BL/6 mice: 'young' (2-6 months), 'middle-aged' (11-16 months), and 'elderly' (25-32 months). We evaluated the age-specific morphology of the outflow pathway tissues. Further, because of the known pressure-dependent Schlemm's canal (SC) narrowing, we assessed the dependence of the SC lumen area to varying IOPs in age-specific FE models over a physiological range of TM/PL stiffness values. We found age-dependent changes in morphology of outflow tissues; notably, the PLs were more developed in older mice compared to younger ones. In addition, FE modeling demonstrated that murine SC patency is highly dependent on the presence of PLs, and that increased IOP caused SC collapse only with sufficiently low TM/PL stiffness values. Moreover, the elderly model showed more susceptibility to SC collapse compared to the younger models. In conclusion, our study elucidated the previously unexplored role of PLs in the aqueous outflow pathway, indicating their function in supporting TM and SC under elevated IOP.

Keywords: Glaucoma, Biomechanics, IOP, Pectinate Ligament, Finite Element Method

Introduction

Glaucoma is a group of eye diseases that are collectively responsible for the majority of irreversible blindness worldwide (World Health Organization, 2019). Glaucoma is often associated with increased intraocular pressure (IOP), i.e., ocular hypertension (Safa et al., 2022c; Weinreb et al., 2014), and lowering of IOP helps preserve vision (Gordon et al., 2002; Heijl et al., 2002; Kass et al., 2002; Leske et al., 2007). Therefore, understanding the mechanisms by which IOP is regulated has great importance in developing new glaucoma treatments.

A critical determinant of IOP is the hydrodynamic resistance of the conventional outflow pathway tissues, including the inner wall of Schlemm's canal (SC) and trabecular meshwork (TM; Figure 1). Further, the biomechanical properties of these tissues are thought to play a key role in IOP homeostasis (Pedrigi et al., 2011; Stamer et al., 2015). For example, several studies have observed TM stiffening in glaucoma and mechanobiological factors are known to affect aqueous humor drainage across the inner wall of SC (Last et al., 2011; Overby et al., 2014; Vahabikashi et al., 2019; Wang et al., 2017). Another integral part of the outflow pathway in many species is the pectinate ligaments (also known as iris processes (Smith et al., 2001)), yet little is known about their biomechanical properties and their role in ocular hypertension and the regulation of IOP (Bergmanson, 1985; De Geest et al., 1990; Flügel et al., 1991; McMenamin and Steptoe, 1991; Samuelson and Gelatt, 1984; Smith et al., 2001).

The pectinate ligaments (PLs), as their name implies, form a comb-like fibrous structure located at the angle of the anterior chamber, bridging between the iris and the TM. PL structure varies substantially between species (Simones et al., 1996), suggesting potential evolutionary significance for this tissue. For example, piscine PLs (or annular ligaments) tend to be more extensive than in mammals (Chen et al., 2008; Soules and Link, 2005). Notably,

in primates, including humans, the PLs are less extensive, with a porous presentation that is

visible by gonioscopy.

Due to the less complex structure of the TM in mice, we hypothesize that the PLs play

a role in mechanically stabilizing the TM and SC by imposing mechanical tension that resists

the normal pressure differential from the anterior chamber to SC lumen. Our objective in this

study was to test this hypothesis via a combination of histology-based morphometry

(histomorphometry), *in vivo* optical coherence tomography (OCT) imaging, and computa-

tional modeling of the conventional outflow pathway tissues. All work was carried out on

mice (Li et al., 2023), a common model to study aqueous humor dynamics.

Methods

We first histomorphometrically characterized key features of the primary outflow

pathway in mice of different ages. We then augmented this histomorphometric data with

OCT imaging to create age-specific computational (finite element [FE]) models of the murine

outflow pathway from mice of different ages. Finally, we conducted forward FE analyses to

investigate the effect of PL structure and mechanics on the biomechanical response of

TM/SC to IOP as a function of age. The experimental procedures are described in detail

elsewhere with a specific focus on morphological and biological effect of aging on aqueous

outflow parameters (Li et al., 2023). Here, we pursue an in-depth analysis of morphology and

biomechanics of the PLs.

Animals

All the experimental procedures were approved by the Institutional Animal Care and

Use Committees (IACUC) at Duke University and followed the tenets of the Association for

Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Oph-

thalmic and Vision Research. Three groups of wild-type C57BL/6 mice were used in this

study, as reported elsewhere (Li et al., 2023).

• Group 1 was used for histomorphometry and consisted of eight young (2-3 months), six

middle-aged (12-13 months), and seven elderly (29 months) mice. All mice for

histomorphometry were male except for two females in the young cohort.

• Group 2 was used for high spatial resolution OCT imaging of the TM and SC ("TM-

OCT") and consisted of five young (4-6 months), seven middle-aged (11-16 months), and

five elderly (25-31 months) mice. These mice were all male except for three females in

the young cohort and two mice of unknown sex in the middle-aged cohort.

• Group 3 was used to evaluate IOP-induced iridial deformation via OCT imaging of the

anterior segment ("Iris-OCT"). The young (3-6 months), middle-aged (9-13 months), and

elderly (29-32 months) cohorts included five, seven, and five mice, respectively. In the

middle-aged and elderly cohorts, all animals were male except for two females in the

middle-aged cohort, while in the young cohort, all animals were female except for one

male.

Please note that the age grouping used in this study essentially corresponded to those reported

in our related paper (Li et al., 2023).

Histomorphometric analysis

Not all relevant anatomic features were visible on OCT images. Thus, we used sagit-

tal histological sections through the outflow tract of mice in Group 1 to obtain some of the

anatomical information needed to create the FE models. Anesthetized mice were euthanized

by decapitation, and the eyes were immediately enucleated and immersion-fixed in 4% para-

formaldehyde + 1% glutaraldehyde at 4 °C overnight. The eyes were then bisected, and the

posterior segments and lenses were discarded. The anterior segments were cut into four quad-

rants and each quadrant was embedded in Epon, after which 0.5 µm thick sagittal sections

were cut using a microtome, stained with methylene blue, and imaged (Axioplan2, Carl Zeiss

MicroImaging, Thornwood, NY, USA; image resolution of ca. 0.26 µm/pixel) to visualize the

outflow pathway tissues (Figure 1). For further details of these experimental procedures,

please see Li *et al.* (Li et al., 2023, 2021, 2019).

We manually identified and segmented the following features of interest from the re-

sulting images (Figure 1):

- TM cross-sectional area ("TM area"), defined as the area of TM tissue bounded by the posterior terminus of Descemet's membrane and by the most posterior point of SC (Figure 1A).

- TM length, measured as the distance between the two (anterior-posterior) extremes of the TM, measured using a straight line (Figure 1B).

- PL thickness, obtained by identifying the most anterior PL and measuring its thickness perpendicular to the local axis of the PL (Figure 1C).

- Active TM cross-sectional area ("Active TM area"), defined as the area of TM tissue lying directly adjacent to the lumen of SC (Figure 1D). Note that "Active TM area" is $\leq$ "TM area".

- SC length, measured from the most posterior to the most anterior point of the SC lumen using a straight line (Figure 1E).

- PL attachment site(s), measured as the straight-line distance parallel to the line defining the SC length from the posterior end of SC to the identified PL (Figure 1F).

Given the subjectivity in identifying some of these features, the segmentations were

independently carried out by four separate annotators who were masked to mouse age and

sample identity. All segmentations were performed using ImageJ (Fiji distribution; various

2023 versions (Schindelin et al., 2012)). The results were then compiled, and the annotators

met to discuss the segmentation results and resolve any disagreements. The annotators then

performed a second round of segmentation under the same masking conditions. After the se-

cond round of segmentation and discussion, three images were removed from the data set

(two from the young cohort and one from the elderly cohort) because the tissues could not be

reliably segmented. As a result, a total of 6 histological images were analyzed per age cohort,

corresponding to 24 possible measurements (4 annotators $\times$ 6 images) per feature of interest.

We then calculated the average "active TM thickness" in each image by dividing, for each

annotator, the active TM area by the corresponding SC length, generating an additional 24

values per age cohort. Of note, due to the complex structure of the outflow tissues and inter-

annotator variability, there were several unsuccessful annotations in some images. For in-

stance, in the young cohort, a PL was not discernible in several samples (13 unmeasurable PL

widths out of a total of 24), suggesting under-development of PLs in young animals (see the

Discussion section).

As mentioned above, it was necessary to combine the morphometric data with both

OCT and histologic images to generate the FE models. We reasoned that such a combination

would be most robust if we scaled all dimensions by a common length apparent in both OCT

and histologic images. We selected SC anterior-posterior length ("SC length") as the common

feature for this purpose. Hence, we calculated two ratios from histologic images: the "TM

ratio," defined as (TM length)/(SC length), and the "PL ratio," defined as (PL attachment dis-

tance)/(SC length). We accounted for age variation by using age-specific TM ratios, i.e., we

computed the ratio of average TM length divided by the average SC length for each age co-

hort, and we repeated the same process for the PL ratio. These ratios were later used in creat-

ing the FE models (see *Model geometry creation and FE mesh generation* section, below).

OCT Imaging

Mice were anesthetized using an intraperitoneal injection of ketamine (100 mg/kg)

and xylazine (10 mg/kg). A drop of 0.5% proparacaine was applied to the eye (Li et al.,

2023). Animals were secured on a custom-made platform, and a single pulled glass

microneedle filled with phosphate-buffered saline was inserted into the anterior chamber of

one eye. The microneedle was connected to both a manometric column to adjust IOP and a

pressure transducer to continuously monitor IOP levels using LabChart software

(ADInstruments, Colorado Springs CO).

Two types of OCT imaging were performed using a high-resolution spectral domain

OCT system (SD-OCT Envisu R2200, Bioptigen Inc., Research Triangle Park, NC, USA). In

the first type of OCT image (TM-OCT, Group 2), we acquired sagittal scans focused on the

iridocorneal angle, including the TM and SC lumen, at IOPs of 10, 15, and 20 mmHg. We

identified the iridocorneal angle structures following our previously established techniques

(Li et al., 2019). In the second type of OCT image (Iris-OCT, Group 3), the OCT scan head

was placed near the corneal apex to acquire images of half of the anterior chamber at IOPs of

10, 15, and 20 mmHg. Additional IOP steps were also used but were not needed in this study

(Lee et al., 2021; Li et al., 2023).

Model geometry creation and FE mesh generation

We created age-specific models of the conventional outflow pathway tissues by com-

bining the morphometric measurements and the TM-OCT images. Due to physiological rele-

vance, the TM-OCT images from 10 mmHg served as the reference state for creating all

models (Figure 2A, D, and G). For each age-specific model, we delineated the corneal and

iridial margins in one representative TM-OCT image (Figure 2B, E, and H) using the 2D

spline function in SolidWorks (version 2022 Dassault Systèmes, Waltham, MA, USA).

The details of the iridocorneal angle tissues were not resolved in these OCT images,

and thus, as mentioned earlier, we relied on histomorphometry to reconstruct details of the

outflow pathway tissues as follows. Using the segmented SC lumen, we created the TM by

assuming that the TM's inner margin was offset from the inner wall of SC by the age-specific

average active TM thickness (Figure 2J, feature '*ii*'). This offset distance was measured nor-

mal to the line defining SC length (Figure 2J, feature '*i*'). Then, to complete the TM profile,

we connected the ends of this inner TM margin to the cornea and root of the iris using linear

extrapolations (Figure 2J, features '*iv*' and '*v*', respectively). In the young model the calculat-

ed PL and TM ratios were almost identical, and thus we extended the calculated TM length

by c. 5% to facilitate meshing, which we note is well within the uncertainty in TM dimen-

sions (Table 2). Finally, we used a constant TM thickness (averaged active TM thickness),

rather than a spatially-variable thickness, as a practical approximation to keep the model trac-

table.

We modeled the PLs as a diffuse (distributed) material rather than attempting to mod-

el the complex, interconnected, and networked structure of the mouse PLs (Chalupa and Wil-

liams, 2008; Smith et al., 2001). We recognize that this homogenized approach to modeling

the PLs is an approximation, and we therefore created two additional models to assess its va-

lidity. In the first, we did not include the PLs at all, while in the second we only included a

single PL "beam" (the anterior-most PL), as previously described (Li et al., 2019). For con-

sistency with previous work, we used the young model for these additional simulations (see

supplementary Figure S2 for details). In both of these models, SC collapsed at low IOPs

(IOP < 11 mmHg), which is contradictory to observations during OCT imaging (Li et al.,

2023, 2019). This simple result suggested that treating the PL network as a diffuse (distribut-

ed) material, coupling the iris and TM, was the most physiologically-relevant option for our

study, as discussed below.

The "distributed" PL material filled the iridocorneal angle, extending from the apex of

the angle to the anterior-most PL attachment site (Figure 2F, feature '*iii*'), as determined for

each age model using the "PL ratio" based on the histomorphometric analysis (Figure 1C and

2). The PL region was bounded by the interior surface of the TM and the anterior surface of

the iris.

The final modeling step was to create a 3D CAD model by extruding the cornea, iris,

TM, and PL as separate bodies, each 10 µm in height. This approach was preferred over cre-

ating an axisymmetric model due to the plane strain assumption (see the *Boundary conditions*

section for further details).

To generate the FE mesh, we exported the CAD model as a STEP file (ASCII version

2) which was then imported into the Gmsh meshing software (version 4.11.1; (Geuzaine and

Remacle, 2009)). We tessellated the volume using TET10 quadratic elements with overall

mesh density set at 7.5 µm, confirmed to provide adequate numerical accuracy in a prelimi-

nary mesh refinement study. To enhance numerical accuracy, we used higher mesh densities

in select locations. Specifically, we decreased the elemental characteristic edge length in the

TM by a factor of 4 and in the PL region by a factor of 8. The meshes were generated using

the Frontal algorithm, and ten steps of smoothing and optimization were conducted to en-

hance mesh quality (Safa et al., 2022b).

Boundary conditions

We used a plane strain modeling approach, justified by the conventional outflow re-

gion's small size relative to its distance from the anterior chamber's axis of symmetry. Fur-

ther, we constrained the posterior (scleral) end of the model and the anterior corneoscleral

surface (Figure 3), which was justified based on the significantly higher stiffness of the sclera

and cornea relative to the key tissues of our model (see *Tissue constitutive models* section be-

low). In addition, the right (central) edge of the cornea and iris were constrained to move only

in the 'y' direction (Figure 3), which allowed for anterior chamber deepening during IOP ele-

vation.

Increasing IOP caused posterior displacement of the iris, as observed from the OCT

images of the anterior chamber (see supplementary Figure S1). We used the lower-resolution

OCT images (Iris-OCT) to extract iris deformations, which we then applied as age-specific

boundary conditions in the FE model. These OCT scans were from a different set of animals

(Group 3) than the high magnification OCT images (TM-OCT; Group 2) due to technical dif-

ficulties in maintaining a consistent OCT field of view at all IOP levels and in aligning the

field of view with the anterior chamber's central axis in the high-magnification TM-OCT

scans. Therefore, we measured the iris displacement along the 'y' axis of Iris-OCT images

(Figure S1; same definition as in Figure 2A for TM-OCT images) at IOP levels of 15 and 20

mmHg relative to the configuration at 10 mmHg and prescribed this displacement in the FE

model as an input (see supplementary Figure S1 for more details).

The pressure in the anterior chamber was specified based on the IOP levels imposed

experimentally. For numerical modeling, IOP values were specified relative to the baseline

pressure of 10 mmHg, i.e., we imposed a relative IOP of 5 mmHg (corresponding to 15

mmHg absolute) and of 10 mmHg (corresponding to 20 mmHg absolute). We set the posteri-

or chamber pressure (PCP) to zero (10 mmHg absolute) based on the results of a prior study

evaluating PCP as a data-fitting variable, which showed that PCP remained at 10 mmHg

when IOP exceeded 10 mmHg (Lee et al., 2021). Finally, SC lumen pressure (SCLP) was

calculated based on IOP and the total resistance of the primary outflow pathway (Li et al.,

2019), which indicated SCLPs of 2 mmHg (12 mmHg absolute) and 4 mmHg (14 mmHg ab-

solute) at IOP levels of 15 mmHg and 20 mmHg, respectively.

Finally, we imposed a contact boundary condition between the inner and outer walls

of SC, which occurred when SC collapsed, using a node-on-facet algorithm and a penalty

method to enforce contact. Specifically, we used an automatic penalty approach with the scal-

ing factor set at 100 times TM stiffness.

Tissue constitutive models

We used an incompressible neo-Hookean ("INH") constitutive relation to model the

cornea and iris.

$$\Psi_{\text{INH}}(\boldsymbol{C}) = \frac{E}{6}(\tilde{I}_1 - 3) + \frac{K}{2}\ln{(J)}^2 \tag{1}$$

where Ψ_{INH} is the Helmholtz free energy, $\boldsymbol{C}$ is the right Cauchy-Green strain tensor, $\tilde{I}_1$ is the

first invariant of the deviatoric right Cauchy-Green strain tensor, and $J = \det(\boldsymbol{F})$ is the

Jacobian of the deformation gradient tensor. The stiffness E was set to 2700 kPa (Li et al.,

2019) for the cornea/sclera and 96.1 kPa for the iris (Lee et al., 2021). The bulk modulus K

was set to 100 times the shear modulus to enforce incompressibility (Weiss et al., 1996).

Due to the fibrous structure of the PLs, we did not model them as incompressible; in-

stead, they were modeled as a homogenized compressible neo-Hookean material ("NH")

$$\Psi_{\text{NH}}(\boldsymbol{C}) = \frac{E}{4(1+v)}(I_1 - 3) - \frac{E}{2(1+v)}\ln(J)$$
$$+ \frac{vE}{2(1+v)(1-2v)}(\ln J)^2 \tag{2}$$

278   Here, I_1 is the first invariant of the right Cauchy-Green strain tensor, and v is Poisson's ratio,

279   which we set to zero due to the approximately parallel structure of the PLs noted above, as-

280   suming the PL beams behave as 1D elements lacking lateral contraction under axial tension.

281   Therefore,

$$\Psi_{\text{NH}}(\boldsymbol{C}) = \frac{E}{4}(I_1 - 3) - \frac{E}{2}\ln(J) \tag{3}$$

We assumed that the intrinsic stiffnesses of TM and PL were the same, based on their similar

collagenous composition, but adjusted the stiffness value for the PLs to account for the high

porosity of the PL network. In brief, we specified the effective stiffness of the PL relative to

the TM using a linear relationship $E_{PL} = \beta E_{TM}$, where β was the solid fraction of the PL

network. We computed β from an analysis of scanning electron micrographs of rabbit PLs

(Ramos et al., 2017; Simones et al., 1996) since suitable high-resolution scanning electron

micrographs from mouse eyes do not exist to the best of our knowledge, obtaining $\beta =$

2.3% (supplementary Figure S3). We then scaled this value down by a factor of 10 for

mouse, since frontal sections through the mouse eye showed a more rarified PL network in

mouse vs. rabbit (Chalupa and Williams, 2008). The same value $\beta = 0.23\%$ was used for all

ages.

Forward finite element simulations and data analysis

We solved the FE models using FEBio (version 4.3.0, (Maas et al., 2012)), and devel-

oped a custom Python class to automate the simulations and post-processing. We specifically

analyzed the deformations of aqueous outflow pathway tissues as a function of IOP in models

with different TM and PL stiffnesses, for which a key metric was SC lumen cross-sectional

area (SCLA). We varied E_{TM} in a physiological range (5-150 kPa) (Wang et al., 2017), with

E_{PL} calculated from E_{TM} as described above. We simulated 15 cases with varying E_{TM} and

E_{PL}.

To calculate SCLA, we used the *concave_hull* function (Shapely version 2.0.1, (Gil-

lies, 2023)) to fit a concave hull shape to the coordinates of the nodes on the inner and outer

walls of SC (Figure 1). For consistency between models, we then computed a normalized

SCLA, defined as

$$nSCLA = \frac{SCLA}{SCLA_{@10\text{ mmHg}}}$$

    4)

where $SCLA_{@10\text{ mmHg}}$ is $SCLA$ at the reference IOP (10 mmHg).

Finally, for each age-specific model and set of TM/PL stiffness values, we assessed

SC lumen 'patency' using a quantity we term "Ultimate nSCLA," defined as the value of

nSCLA at IOP = 20 mmHg. An Ultimate SCLA value greater than one indicated a SC lu-

minal area expansion at 20 mmHg relative to SCLA at 10 mmHg; conversely, Ultimate

SCLA values less than one indicated an area reduction. In addition, as another measure of SC

patency, we determined the lowest IOP at which SC 'collapse' occurred, which was defined

as $nSCLA < 33\%$. These definitions were based on informed estimations of the state of SC

as it pertains to its capability to function within normal aqueous outflow, where collapsed re-

gions of SC would not be able to function normally.

Statistical analysis

We used a linear mixed effects (LME) model to study the effect of age on the mor-

phometric characteristics of the mouse outflow pathway. Specifically, we used 'age' as the

fixed effect and 'annotator' as a random effect. The significance threshold (α) was set at 5%.

We also conducted a *post hoc* multiple comparison test on 'age' using Tukey's HSD test with

Bonferroni correction (critical $\alpha = 5\%/3$). In addition, we conducted a one-way ANOVA to

test for the effect of age on TM- and PL-ratios ($\alpha = 5\%$), followed with *post hoc* Tukey's

HSD test (critical $\alpha = 5\%/3$). Statistical analysis was performed using JMP (Pro version 16,

SAS Institute Inc., Cary, NC, USA)

Results

Morphometrical characterization of TM and PL in C57BL/6 mice

Histomorphometric analysis indicated that all measured dimensions changed with age,

except PL thickness (Figure 4 and Table 1; $p < 0.05$ for the effect of age), with no effect of

annotator ($p > 0.05$, Table 1). *Post hoc* multiple comparison tests indicated that, compared to

the young cohort, the elderly cohort had a larger TM area ($p = 0.014$), Active TM area

($p = 0.001$), and SC length ($p = 0.001$) (Figure 4, Table 1, and Table S1). In addition, the

middle-aged cohort had a smaller SC length compared to the elderly cohort ($p < 0.0001$)

(Figure 3, Table 1, and Table S1). Overall, there was a positive correlation between age and

the size of the features.

Furthermore, the TM- and PL-ratios were significantly affected by age ($p < 0.05$;

Table 2). More specifically, the elderly mice had smaller TM- and PL-ratios relative to the

young cohort (Table S3), indicating a moderately posterior localization of TM and PLs rela-

tive to SC lumen in elderly animals.

Dependence of SC lumen area on TM/PL stiffness

All models (for all three age groups) demonstrated IOP-induced SC lumen collapse,

as was also observed experimentally (Li et al., 2023, 2019, 2016, 2014). However, large

TM/PL stiffness values caused unphysiological SC expansion at elevated IOP (Figure 5),

which was not observed experimentally. More specifically, SC lumen area showed a highly

nonlinear dependence on IOP and TM/PL stiffness in all age-specific models (Figure 5D).

Interestingly, elevation of IOP did not guarantee reduction of SC luminal area. Instead, col-

lapse only occurred at lower TM stiffnesses (Figure 5B, C, and D). At higher stiffnesses, SC

lumen—counter-intuitively—expanded with increasing IOP (Figure 5A and D). Further, the

dependence of normalized SC cross-sectional lumen area, nSCLA, on IOP was not always

even monotonic. For instance, in the middle-aged model, there was an initial phase of nSCLA

reduction with increasing IOP followed by a sudden expansion of the SC at *ca.* IOP = 11

mmHg (Figure 5D middle). But, as IOP further increased beyond 15 mmHg, SC luminal area

gradually shrank. In a similar fashion, albeit more gradually, such behavior was observed in

the elderly model, and to a lesser extent, in the young model (Figure 5D).

Patency of SC lumen

To investigate the patency of SC, we evaluated: (i) the Ultimate nSCLA for each age-

specific FE model vs. TM/PL stiffness (Figure 6A) and (ii) the IOP at which SC lumen first

collapsed for each age-specific model vs. TM/PL stiffness (Figure 6B). Recall that in our ap-

proach, increasing TM stiffness also proportionally increased PL stiffness; thus, we simply

denote stiffness as "TM/PL stiffness". We observed that the Ultimate nSCLA strongly de-

pended on TM/PL stiffness, with a lower stiffness giving a smaller Ultimate nSCLA (Figure

6A). In the young and middle-aged models, the Ultimate nSCLA was greater than 1 (expan-

sion of SC lumen) for cases with $E_{TM} > 46.4$ kPa ($E_{PL} > 0.11$ kPa) in the young model, and

$E_{TM} > 67.1$ kPa ($E_{PL} > 0.15$ kPa) in the middle-aged model (Figure 6A). For the cases where

the Ultimate nSCLA was smaller than 1, which are physiologically relevant, the elderly mod-

el showed a smaller Ultimate nSCLA for any given E_{TM} compared to both young and middle-

aged models.

Further, when investigating the collapse IOP for different TM/PL stiffnesses (Figure

6B), the highest E_{TM} at which SC collapsed in the elderly model was $E_{TM} = 87.9$ kPa

($E_{PL} = 0.20$ kPa), which was higher than that of the young and middle-aged models $E_{TM} =$

25.7 kPa ($E_{PL} = 0.06$ kPa; Figure 6B). As expected, all models collapsed at a low IOP of

10~11 mmHg at the lowest simulated E_{TM} stiffness of 5 kPa ($E_{PL} = 0.01$ kPa; Figure 6B).

Yet, surprisingly, for the same TM/PL stiffness levels, the elderly model collapsed at the

lowest IOP compared to both other models (Figure 6B), except for the limiting case of

$E_{TM} = 5$ kPa ($E_{PL} = 0.01$ kPa), where the models had a similar collapse IOPs.

Discussion

In this study, we investigated the stabilizing role of the pectinate ligaments (PLs) on

the conventional outflow pathway in the mouse eye, a ubiquitous model for studying aqueous

humor dynamics. We showed that the stiffness and morphology of the PLs play a significant

role in stabilizing the TM, and hence, in the patency of SC lumen and how it deforms as IOP

changes. These observations reveal a previously unexplored role for the PLs in murine aque-

ous humor dynamics. Indeed, in our computational models in which the PLs were not pre-

sent, SC was predicted to collapse at very low IOPs, in opposition to what is observed exper-

imentally. Considering the anatomy of the mouse TM (few beams and proportionately much

thinner in the interior-exterior direction than in larger eyes), it is perhaps not surprising that

mechanical stabilization of the TM in the mouse is needed to avoid complete SC collapse,

which would cause partial loss of functionality of the primary aqueous humor outflow path-

way and significant elevation of IOP.

It is interesting that in one of the few studies of PL mechanics (Morrison and Van

Buskirk, 1982) reported that severing the PLs in a dog led to an expansion of the TM and a

corresponding increase in outflow facility, i.e. opposite to our predictions. This contradiction

is likely due to a species effect: dogs have a more robust TM and iris that may be more re-

sistant to collapse and bowing, respectively, so that the canine PLs likely play less of a TM

stabilizing role than in the mouse, but rather serve to maintain the angle between the TM and

iris. Consistent with this interpretation, canine PLs appear to have a subtle, but important dif-

ference in morphology vs. the mouse: in the dog, the PLs attach to the iris immediately prox-

imal to the root and then run anteriorly so that their anterior-most attachment point lies in the

corneal limbal region rather than in the TM (see Figure 2 from Morrison and Van Buskirk

(Morrison and Van Buskirk, 1982)). The net effect is that the course of the PLs in the canine

eye is more parallel to the beams of the TM than in the murine eye, where there seems to be a

more interior-exterior alignment of the PLs leading to iris bowing directly pulling the TM

interiorly. We conclude that inter-species differences in PL morphology and function may be

important and merit future attention.

One interesting observation was the non-monotonic dependence of Schlemm's canal

luminal area on IOP (Figure 5D), which illustrates the complex biomechanics of the outflow

pathway. This behavior was particularly evident in the middle-aged model (Figure 5D, mid-

dle) which illustrates two opposing factors impacting the configuration of outflow tissues: (i)

IOP elevation tends to collapse SC lumen by outwards displacement of the TM, whereas (ii)

posterior iridial movement with increasing IOP tends to pull the TM inward and open SC.

The complex shapes of the specific curves in Figure 5D depend on TM stiffness, PL stiffness,

and geometric factors such as TM thickness and PL anterior-posterior extent (Figure 5 A-C).

The complexity of these effects is reinforced by observing the slope discontinuities in the

nSCLA vs. IOP curves, which were due to progressively larger contact between the inner and

outer walls of SC. In future work, we should also investigate the role of septae in the SC lu-

men (Bentley et al., 2016), known to be present in human eyes but largely uninvestigated in

the murine eye.

Although PL biomechanics has not been the focus of many investigations, PL mor-

phology has been studied in a variety of domestic mammals, including dogs, horses, and rab-

bits, via scanning electron microscopy (SEM) and other types of histology (Chen et al., 2008;

Simones et al., 1996). PL morphology varies between species via differences in PL width,

length, spacing, and insertion angle into the TM. Furthermore, PL morphology also varies

locally within the same eye. One factor suggested to influence PL structure is age (Simones et

al., 1996). While the effect of aging on PLs was not a primary focus of this work (see (Li et

al., 2023)), our models were based on images from different aged animals. Therefore, it is

worth exploring our results from that perspective.

Our histomorphometric analysis demonstrated a dependence of structural features on

age (Figure 2 and Table 1) with certain features becoming more prominent with age (Figure

2), as expected. For instance, in the young cohort, the PL was relatively underdeveloped. In

fact, we were unable to observe PLs in 13 out of 24 annotations from young eyes. Additional-

ly, age also affected the TM and PL ratios (Table 2), with older eyes having larger SCs, and

thus, smaller TM and PL ratios (Figure 2). We hypothesize these differences might be related

to development and growth of more prominent PLs with age, thereby providing more support

to the TM, preventing SC collapse in middle-aged and older eyes.

In addition, our simulations indicated a connection between iris deformation and

TM/PL stiffness (Figure 5A-C), which was demonstrated by differences in the slope of the

iris midline at the pupillary (right) edge of our models (IOP = 15 mmHg and 20 mmHg col-

umns; Figure 5A-C). This observation suggests a connection between iris deformations and

aging, which can influence the association between anterior chamber deepening (AC deepen-

ing) and IOP. We know AC deepening significantly affects outflow facility and can also arti-

ficially elevate facility in experiments (Boussommier-Calleja et al., 2015; Brubaker, 1975;

Ellingsen and Grant, 1971). Therefore, in perfusion preparations where the iris is intact, the

association between AC deepening and changes in outflow facility may be explained by PL

tension, where with more AC deepening the PLs open TM and prevent SC collapse (as ob-

served in some cases in our study; Figure 5A), therefore increasing facility. Thus, it would be

worth more closely exploring whether there are connection between TM/PL stiffness and iris

deformations under, for example, various physiologically occurring iris contractions (e.g., at

different light levels (Safa et al., 2022a)), and at different ages to study connections between

iris deformation, age, and outflow pathway function.

As a further age-dependent observation, our simulations indicated a strong effect of

age on the biomechanical properties of outflow tissues. For example, by comparing modeled

SCLA vs. experimentally observed SCLA changes with IOP, a significant stiffening of TM

was observed with age (Li et al., 2023). Of note, our predicted TM stiffness for young mice

(~29 kPa (Li et al., 2023)) was consistent with previous results on young mice (Li et al.,

2019). We also predicted that elderly eyes had the smallest Ultimate nSCLA for a given

TM/PL stiffness (Figure 6A). Overall, these observations indicate an increased resistance to

IOP-induced deformations of outflow tissues with age, due to a combination of changes in

biomechanical properties and tissue geometry.

It is important to note that we here used quadratic finite elements (TET10) that have

superior performance in modeling incompressible materials (Bonet and Wood, 1997) than the

linear elements that we previously used (TET4; (Li et al., 2019)). This element type was es-

sential for making comparisons between different FE models in this study and should be used

as for future modeling work in studying outflow tissue biomechanics. This difference in ele-

ment type likely explains differences between our current results and those seen in earlier

studies from our lab, e.g. (Li et al., 2019), where we previously predicted that a single PL

could stabilize the TM.

It is worth noting this study had some limitations. First, due to a lack of information

about PL biomechanical properties, we were forced to estimate PL stiffness based on infor-

mation about PL porosity in a rabbit eye which required several assumptions. Fortunately, we

kept the PL to TM stiffness ratio consistent between models, thus eliminating variability in

this ratio as a source of difference between models. Nonetheless, more direct esti-

mates/measurements of PL biomechanical properties, while challenging, should be undertak-

en in future. Second, OCT imaging of mouse eye is subject to some technical limitations: *i)*

because of the diminutive size of the mouse eye, the manual positioning of the OCT scan

head, and IOP-induced changes in ocular geometry, there was inevitable shifting of the rela-

tive position of the scan probe and the eye between scans of the same eye at different IOPs,

and *ii)* OCT imaging is not able to capture all details of the angle structures, leaving us no

choice but to use histomorphometry to "inject" those structures into FEM models based on

OCT images. This process, together with the anatomic variability between mice, inevitably

led to some mismatches between histologically-determined dimensions and the individual

OCT images (e.g. Figure 2H). These technical limitations added variability to our analysis,

which was addressed by using lower magnification Iris-OCT images for certain parts of the

modeling process, and by using age-specific average histomorphometric feature dimensions.

In the future, hardware to accurately position the OCT scan head, and improved resolution

(penetration depth) to better image angle structures might improve the repeatability of high-

resolution OCT images. Third, we based our analysis on one strain of mice (i.e., C57BL/6;

however, PL in other strains, such as FVB/N, may demonstrate somewhat different behavior.

For example, there is a different correlation between AC deepening/outflow facility vs. pres-

sure in C57BL/6 as compared to FVB/N mice (Reina-Torres et al., 2022). Hence, future in-

vestigations should consider other mouse strains (and species) to better elucidate the role of

PL in aqueous outflow. Finally, although sex differences are likely important in glaucoma

(Douglass et al., 2023; Lee et al., 2021), due to the scarcity of female samples in our data set,

we did not consider sex as a biological variable in our analysis. Therefore, caution is advised

interpreting these results for both sexes.

In conclusion, our simulations showed that PL biomechanics plays an essential role in

maintaining stability of the outflow pathway tissues in the mouse eye. In the future, it will be

important to directly characterize PLs' biomechanical properties, preferably by direct biome-

chanical testing of PL beams, e.g., using atomic force microscopy. We note that PLs are less

prominent in healthy human eyes and may thus be less clinically important relative to the pre-

clinical studies in mouse eyes. However, structures like pectinate ligaments can persist (see

e.g. Figure 11 in (Anderson, 1981)) in human eyes with developmental pathologies leading to

congenital (infantile) glaucoma, and it would be of interest to evaluate their role in such cas-

es.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgments

We gratefully acknowledge Prof. Darryl Overby of Imperial College London for his

insightful comments regarding interpreting of our results. Further, we note the generous fi-

nancial support from The BrightFocus Foundation (postdoctoral fellowship G2021005F,

BNS), NIH (K99EY035360 [BNS], T32EY007092 [NSFG], R01EY030871 [AJF],

P30EY006360, P30EY005722, R01EY031710 [CRE and WDS], and R01EY030124

[WDS]), the Alfred P. Sloan Foundation G-2019-11435 (NSFG), the Georgia Research Alli-

ance (CRE), and Department of Veterans Affairs Rehab R&D Service Career Development

Awards (AJF; CDA-2; RX002342).

References

Anderson, D.R., 1981. The development of the trabecular meshwork and its abnormality in primary infantile glaucoma. Trans. Am. Ophthalmol. Soc. 79, 458–485.

Bentley, M.D., Hann, C.R., Fautsch, M.P., 2016. Anatomical Variation of Human Collector Channel Orifices. Invest. Ophthalmol. Vis. Sci. 57, 1153–1159. https://doi.org/10.1167/iovs.15-17753

Bergmanson, J.P., 1985. The anatomy of the rabbit aqueous outflow pathway. Acta Ophthalmol. (Copenh.) 63, 493–501. https://doi.org/10.1111/j.1755-3768.1985.tb05234.x

Bonet, J., Wood, R.D., 1997. Nonlinear Continuum Mechanics for Finite Element Analysis.

Boussommier-Calleja, A., Li, G., Wilson, A., Ziskind, T., Scinteie, O.E., Ashpole, N.E., Sherwood, J.M., Farsiu, S., Challa, P., Gonzalez, P., Downs, J.C., Ethier, C.R., Stamer, W.D., Overby, D.R., 2015. Physical Factors Affecting Outflow Facility Measurements in Mice. Invest. Ophthalmol. Vis. Sci. 56, 8331–8339. https://doi.org/10.1167/iovs.15-17106

Brubaker, R.F., 1975. The effect of intraocular pressure on conventional outflow resistance in the enucleated human eye. Invest. Ophthalmol. Vis. Sci. 14, 286–292.

Chalupa, L.M., Williams, R.W., 2008. Eye, retina, and visual system of the mouse, Eye, retina, and visual system of the mouse. Boston Review, Cambridge, MA, US.

Chen, C.-C., Yeh, L.-K., Liu, C.-Y., Kao, W.W.-Y., Samples, J.R., Lin, S.-J., Hu, F.-R., Wang, I.-J., 2008. Morphological Differences between the Trabecular Meshworks of Zebrafish and Mammals. Curr. Eye Res. 33, 59–72. https://doi.org/10.1080/02713680701795026

De Geest, J.P., Lauwers, H., Simoens, P., De Schaepdrijver, L., 1990. The morphology of the equine iridocorneal angle: a light and scanning electron microscopic study. Equine Vet. J. Suppl. 30–35. https://doi.org/10.1111/j.2042-3306.1990.tb04708.x

Douglass, A., Dattilo, M., Feola, A.J., 2023. Evidence for Menopause as a Sex-Specific Risk Factor for Glaucoma. Cell. Mol. Neurobiol. 43, 79–97. https://doi.org/10.1007/s10571-021-01179-z

Ellingsen, B., Grant, M., 1971. The Relationship of Pressure and Aqueous Outflow in Enucleated Human Eyes. Invest. Ophthalmol. Vis. Sci. 10, 430–437.

Flügel, C., Tamm, E., Lütjen-Drecoll, E., 1991. Different cell populations in bovine trabecular meshwork: an ultrastructural and immunocytochemical study. Exp. Eye Res. 52, 681–690. https://doi.org/10.1016/0014-4835(91)90020-f

Geuzaine, C., Remacle, J.-F., 2009. Gmsh: A 3-D finite element mesh generator with built-in pre- and post-processing facilities. Int. J. Numer. Methods Eng. 79, 1309–1331. https://doi.org/10.1002/nme.2579

Gillies, S., 2023. The Shapely User Manual — Shapely 2.0.1 documentation [WWW Document]. URL https://shapely.readthedocs.io/en/stable/manual.html (accessed 10.9.23).

Gordon, M.O., Beiser, J.A., Brandt, J.D., Heuer, D.K., Higginbotham, E.J., Johnson, C.A., Keltner, J.L., Miller, J.P., Parrish, R.K., II, Wilson, M.R., Kass, M.A., for the Ocular Hypertension Treatment Study Group, 2002. The Ocular Hypertension Treatment Study: Baseline Factors That Predict the Onset of Primary Open-Angle Glaucoma. Arch. Ophthalmol. 120, 714–720. https://doi.org/10.1001/archopht.120.6.714

Heijl, A., Leske, M.C., Bengtsson, Bo, Hyman, L., Bengtsson, Boel, Hussein, M., Early Manifest Glaucoma Trial Group, 2002. Reduction of intraocular pressure and glaucoma progression: results from the Early Manifest Glaucoma Trial. Arch. Ophthalmol. Chic. Ill 1960 120, 1268–1279. https://doi.org/10.1001/archopht.120.10.1268

JMP, 2023. Mixed Model Report and Options [WWW Document]. URL

https://www.jmp.com/support/help/en/17.2/index.shtml#page/jmp/mixed-model-
report-and-options.shtml (accessed 10.16.23).

Kass, M.A., Heuer, D.K., Higginbotham, E.J., Johnson, C.A., Keltner, J.L., Miller, J.P., Par-
rish, R.K., Wilson, M.R., Gordon, M.O., 2002. The Ocular Hypertension Treatment
Study: a randomized trial determines that topical ocular hypotensive medication de-
lays or prevents the onset of primary open-angle glaucoma. Arch. Ophthalmol. Chic.
Ill 1960 120, 701–713; discussion 829-830.
https://doi.org/10.1001/archopht.120.6.701

Last, J.A., Pan, T., Ding, Y., Reilly, C.M., Keller, K., Acott, T.S., Fautsch, M.P., Murphy,
C.J., Russell, P., 2011. Elastic modulus determination of normal and glaucomatous
human trabecular meshwork. Invest. Ophthalmol. Vis. Sci. 52, 2147–2152.
https://doi.org/10.1167/iovs.10-6342

Lee, C., Li, G., Stamer, W.D., Ethier, C.R., 2021. In vivo estimation of murine iris stiffness
using finite element modeling. Exp. Eye Res. 202, 108374.
https://doi.org/10.1016/j.exer.2020.108374

Leske, M.C., Heijl, A., Hyman, L., Bengtsson, B., Dong, L., Yang, Z., EMGT Group, 2007.
Predictors of long-term progression in the early manifest glaucoma trial. Ophthalmol-
ogy 114, 1965–1972. https://doi.org/10.1016/j.ophtha.2007.03.016

Li, G., Farsiu, S., Chiu, S.J., Gonzalez, P., Lütjen-Drecoll, E., Overby, D.R., Stamer, W.D.,
2014. Pilocarpine-Induced Dilation of Schlemm's Canal and Prevention of Lumen
Collapse at Elevated Intraocular Pressures in Living Mice Visualized by OCT. Invest.
Ophthalmol. Vis. Sci. 55, 3737–3746. https://doi.org/10.1167/iovs.13-13700

Li, G., Lee, C., Agrahari, V., Wang, K., Navarro, I., Sherwood, J.M., Crews, K., Farsiu, S.,
Gonzalez, P., Lin, C.-W., Mitra, A.K., Ethier, C.R., Stamer, W.D., 2019. In vivo
measurement of trabecular meshwork stiffness in a corticosteroid-induced ocular hy-
pertensive mouse model. Proc. Natl. Acad. Sci. 116, 1714–1722.
https://doi.org/10.1073/pnas.1814889116

Li, G., Lee, C., Read, A.T., Wang, K., Ha, J., Kuhn, M., Navarro, I., Cui, J., Young, K.,
Gorijavolu, R., Sulchek, T., Kopczynski, C., Farsiu, S., Samples, J., Challa, P., Ethier,
C.R., Stamer, W.D., 2021. Anti-fibrotic activity of a rho-kinase inhibitor restores out-
flow function and intraocular pressure homeostasis. eLife 10, e60831.
https://doi.org/10.7554/eLife.60831

Li, G., Mukherjee, D., Navarro, I., Ashpole, N.E., Sherwood, J.M., Chang, J., Overby, D.R.,
Yuan, F., Gonzalez, P., Kopczynski, C.C., Farsiu, S., Stamer, W.D., 2016. Visualiza-
tion of conventional outflow tissue responses to netarsudil in living mouse eyes. Eur.
J. Pharmacol. 787, 20–31. https://doi.org/10.1016/j.ejphar.2016.04.002

Li, G., van Batenburg-Sherwood, J., Safa, B.N., Fraticelli Guzman, N.S., Wilson, A., Fard,
M.R.B., Choy, K., Ieso, M.L. de, Cui, S., Feola, A.J., Weisz, T., Kuhn, M., Rickman,
C.B., Farsiu, S., Ethier, C.R., Stamer, W.D., 2023. Aging and intraocular pressure
homeostasis in mice. bioRxiv 2023.10.17.562768.
https://doi.org/10.1101/2023.10.17.562768

Maas, S.A., Ellis, B.J., Ateshian, G.A., Weiss, J.A., 2012. FEBio: Finite Elements for Bio-
mechanics. J. Biomech. Eng. 134, 011005–011005. https://doi.org/10.1115/1.4005694

McMenamin, P.G., Steptoe, R.J., 1991. Normal anatomy of the aqueous humour outflow sys-
tem in the domestic pig eye. J. Anat. 178, 65–77.

Morrison, J.C., Van Buskirk, E.M., 1982. The canine eye: pectinate ligaments and aqueous
outflow resistance. Invest. Ophthalmol. Vis. Sci. 23, 726–732.

Overby, D.R., Zhou, E.H., Vargas-Pinto, R., Pedrigi, R.M., Fuchshofer, R., Braakman, S.T.,
Gupta, R., Perkumas, K.M., Sherwood, J.M., Vahabikashi, A., Dang, Q., Kim, J.H.,
Ethier, C.R., Stamer, W.D., Fredberg, J.J., Johnson, M., 2014. Altered

mechanobiology of Schlemm's canal endothelial cells in glaucoma. Proc. Natl. Acad.
Sci. 111, 13876–13881. https://doi.org/10.1073/pnas.1410602111

Pedrigi, R.M., Simon, D., Reed, A., Stamer, W.D., Overby, D.R., 2011. A model of giant
vacuole dynamics in human Schlemm's canal endothelial cells. Exp. Eye Res. 92, 57–
66. https://doi.org/10.1016/j.exer.2010.11.003

Ramos, M.F., Teixeira, L., Brandt, C.R., Auyeung-Kim, D., 2017. Ocular Immunopathology,
in: Parker, G.A. (Ed.), Immunopathology in Toxicology and Drug Development: Vol-
ume 2, Organ Systems, Molecular and Integrative Toxicology. Springer International
Publishing, Cham, pp. 695–762. https://doi.org/10.1007/978-3-319-47385-7_14

Reina-Torres, E., van Batenburg-Sherwood, J., Overby, D.R., 2022. The relationship between
pressure-dependent changes in outflow facility and dimensional changes in the anteri-
or segment during perfusion of enucleated mouse eyes. Invest. Ophthalmol. Vis. Sci.
63, 3301-A0401.

Safa, B.N., Bahrani Fard, M.R., Ethier, C.R., 2022a. In vivo biomechanical assessment of
iridial deformations and muscle contractions in human eyes. J. R. Soc. Interface 19,
20220108. https://doi.org/10.1098/rsif.2022.0108

Safa, B.N., Bleeker, A., Berdahl, J.P., Ethier, C.R., 2022b. The Effects of Negative Periocular
Pressure on Biomechanics of the Optic Nerve Head and Cornea: A Computational
Modeling Study. https://doi.org/10.1101/2022.05.17.492338

Safa, B.N., Wong, C.A., Ha, J., Ethier, C.R., 2022c. Glaucoma and biomechanics. Curr.
Opin. Ophthalmol. https://doi.org/10.1097/ICU.0000000000000829

Samuelson, D.A., Gelatt, K.N., 1984. Aqueous outflow in the beagle. I. Postnatal morpholog-
ic development of the iridocorneal angle: pectinate ligament and uveal trabecular
meshwork. Curr. Eye Res. 3, 783–794. https://doi.org/10.3109/02713688409000790

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source
platform for biological-image analysis. Nat. Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019

Simones, P., De Geest, J.P., Lauwers, H., 1996. Comparative morphology of the pectinate
ligaments of domestic mammals, as observed under the dissecting microscope and the
scanning electron microscope. J. Vet. Med. Sci. 58, 977–982.
https://doi.org/10.1292/jvms.58.10_977

Smith, R.S., Zabaleta, A., Savinova, O.V., John, S.W., 2001. The mouse anterior chamber
angle and trabecular meshwork develop without cell death. BMC Dev. Biol. 1, 3.
https://doi.org/10.1186/1471-213x-1-3

Soules, K., Link, B., 2005. Morphogenesis of the anterior segment in the zebrafish eye. BMC
Dev. Biol. 5, 12. https://doi.org/10.1186/1471-213X-5-12

Stamer, W.D., Braakman, S.T., Zhou, E.H., Ethier, C.R., Fredberg, J.J., Overby, D.R., John-
son, M., 2015. Biomechanics of Schlemm's canal endothelium and intraocular pres-
sure reduction. Prog. Retin. Eye Res. 44, 86–98.
https://doi.org/10.1016/j.preteyeres.2014.08.002

Vahabikashi, A., Gelman, A., Dong, B., Gong, L., Cha, E.D.K., Schimmel, M., Tamm, E.R.,
Perkumas, K., Stamer, W.D., Sun, C., Zhang, H.F., Gong, H., Johnson, M., 2019. In-
creased stiffness and flow resistance of the inner wall of Schlemm's canal in glauco-
matous human eyes. Proc. Natl. Acad. Sci. U. S. A. 116, 26555–26563.
https://doi.org/10.1073/pnas.1911837116

Wang, K., Read, A.T., Sulchek, T., Ethier, C.R., 2017. Trabecular meshwork stiffness in
glaucoma. Exp. Eye Res. 158, 3–12. https://doi.org/10.1016/j.exer.2016.07.011

Weinreb, R.N., Aung, T., Medeiros, F.A., 2014. The Pathophysiology and Treatment of

Glaucoma. JAMA 311, 1901–1911. https://doi.org/10.1001/jama.2014.3192

Weiss, J.A., Makerc, B.N., Govindjeed, S., 1996. Finite element implementation of incom-
pressible , isotropic hyperelasticity transversely 7825.

World Health Organization, 2019. World report on vision, World health Organisation.

Table 1: Results of the linear mixed effects (LME) model testing on the histomorphometric characterization of the C57BL/6 mouse outflow pathway. The columns provide F-statistics for the fixed effect ('age'; **Bold**: $p < 0.05$), and Wald p-value for the random factor ('annotator'), which tests the null hypothesis that the random effect was significantly different from zero (*Wald p-value < 0.05*) (JMP, 2023).

Feature of interest	Age		Annotator
	F-statistics	*p-value*	*Wald p-value*
TM area (μm^2)	$f(2, 66) = 4.34$	**0.017**	0.419
TM length (μm)	$f(2, 66) = 3.77$	**0.028**	0.523
PL thickness (μm)	$f(2, 41.3) = 0.97$	0.389	0.410
Active TM area (μm^2)	$f(2, 66) = 6.95$	**0.002**	0.479
SC length (μm)	$f(2, 65.1) = 12.40$	< 0.0001	0.911
PL attachment site (μm)	$f(2, 45.2) = 3.36$	**0.044**	0.082

Table 2: Ratios of the features measured from histomorphometric analysis used for creating the FE models. (**Bold**: *$p < 0.05$)

	Young	*Middle-aged*	*Elderly*	*F-statistics*	*p-value*
TM-ratio (TM length/SC length)	1.40 ± 0.26	1.35 ± 0.43	1.16 ± 0.19	$f(2,68) = 3.86$	**0.026***
PL-ratio (PL attachment site /SC length)	1.36 ± 0.46	1.11 ± 0.36	1.00 ± 0.25	$f(2,45) = 3.83$	**0.029***

670 **Figure 1:** A representative sagittal section (from a young mouse) including the conventional outflow
671 pathway tissues, showing definitions of the annotated features (red lines/boundaries) for morphomet-
672 ric characterization. TM = trabecular meshwork, PL = pectinate ligament, SC = Schlemm's canal, CB
673 = ciliary body. Scale bar = 50 µm.

676
677
Figure 2: TM-OCT images, outlined features, and FE meshes for the (A-C) 'young', (B-F) 'middle-
aged', and (G-I) 'elderly' models. Panels A, D, and G show OCT images acquired at an IOP of 10
mmHg with the lumen of SC overlain in blue. Panels B, E, and H show key structures segmented
from OCT images or deduced from histomorphometric analysis (red overlaid curves), while panels C,
F, and I show the resulting FE meshes. (J) Demonstration of a set of features used to create models,
with emphasis on features based on histomorphometric analysis, including: (*i*) the line connecting the
posterior and anterior ends of SC, (*ii*) an offset to create the inner margin of the TM based on the con-
tour of the inner wall of SC, (*iii*) the age-specific calculated distance of the posterior end of SC to the
anterior-most PL attachment site, (*iv*) a linear interpolation of the TM inner margin from the PL at-
tachment site to cornea, (*v*) a linear interpolation of the inner TM margin connecting the posterior end
of the TM to the apex of the irideocorneal angle, and (*vi*) the age-specific calculated TM length. Col-
ors in C, F, I and J: orange = cornea and sclera; yellow = iris and other uveal tissue; blue = pectinate
ligament region; purple = TM. Scale bar = 100 μm.

Figure 3: Boundary conditions for the FE model. Here, the young model is used as an example for illustrative purposes. Note the colored lines indicating the location where pressure loads were applied: SCLP = SC lumen pressure, IOP = intraocular pressure, PCP = posterior chamber pressure, and ⫽ = prescribed displacement of the iris. Pinned support means no displacement in the 'x' and 'y' directions; rolling support indicates no displacement in the 'x' direction. Colors of individual sub-regions are the same as in Figure 2. Scale bar = 100 µm.

Figure 4: Results of the histomorphometric analysis of mouse outflow pathway tissue dimensions. For each feature, the data is shown using violin plots for each age cohort, with the mean indicated by a short black horizontal line. The results from different annotators are shown with colored dots, where each color corresponds to a unique annotator. Statistically significant differences by *post hoc* testing (after Bonferroni correction) are shown using solid horizontal lines (   ), and trending differences are also shown using dashed horizontal lines (.

Figure 5: The results of forward FE simulations for each age-specific model, showing representative simulation results for the (A) young, (B) middle-aged, and (C) elderly models. In addition, (D) normalized SC lumen area (nSCLA; SC area normalized by the SCLA at 10 mmHg) vs. IOP is plotted for a range of values (color bar; 5 kPa ≤ E_{TM} ≤ 150 kPa). Interestingly, SCLA was not monotonically reduced by increasing IOP in the cases with higher E_{TM}, explainable by tension on the TM delivered by the PL due to the posterior deformations of the iris. Additionally, in the middle-aged simulations, there was an interesting rapid initial increase in SCLA followed by a gradual decrease, which was not evident in the simulations in the other age-specific models.

716 **Figure 6:** The patency of the SC lumen was characterized in each age-specific model by evaluating
717 (A) the dependence of Ultimate nSCLA (nSCLA at 20 mmHg) on TM stiffness, and (B) the IOP at
718 which SC lumen collapsed vs. TM stiffness. Note that in the young and middle-aged models SC col-
719 lapse (B) did not occur in cases with stiffness higher that 26 kPa, which is the reason for the termina-
720 tion of the associated curves at that point.

Table S1: Results of the multiple comparison Tukey's HSD test on the 'age' factor on histomorphometric characterization of the C57BL/6 mouse outflow pathway. (*p<0.05, **p<0.017*)

Feature of interest	Age comparison		p-value
TM area (µm^2)	Young	Middle-aged	0.575
	Young	Elderly	**0.014****
	Middle-aged	Elderly	0.149
TM length (µm)	Young	Middle-aged	0.243
	Young	Elderly	0.513
	Middle-aged	Elderly	**0.022***
PL thickness (µm)	Young	Middle-aged	0.445
	Young	Elderly	0.417
	Middle-aged	Elderly	1.000
Active TM area (µm^2)	Young	Middle-aged	0.386
	Young	Elderly	**0.001****
	Middle-aged	Elderly	0.055
SC length (µm)	Young	Middle-aged	0.809
	Young	Elderly	**0.001****
	Middle-aged	Elderly	< 0.001**
PL attachment site (µm)	Young	Middle-aged	0.229
	Young	Elderly	0.861
	Middle-aged	Elderly	**0.040***

Table S2: Summary statistics for the histomorphometric characterization of C57BL/6 mouse outflow pathway features by age cohort. Data is shown as mean and 95% confidence interval (95% CI).

Feature of interest	Age	Mean	95% CI
TM area (μm^2)	Young	1445.49	[1311.43, 1579.56]
	Middle-aged	1619.64	[1311.58, 1927.71]
	Elderly	1946.82	[1638.26, 2255.38]
TM length (μm)	Young	194.29	[182.89, 205.70]
	Middle-aged	177.86	[163.05, 192.68]
	Elderly	205.50	[187.07, 223.94]
PL thickness (μm)	Young	4.46	[3.72, 5.21]
	Middle-aged	5.59	[3.77, 7.41]
	Elderly	5.77	[4.28, 7.26]
Active TM area (μm^2)	Young	1159.28	[1012.10, 1306.45]
	Middle-aged	1364.23	[1073.60, 1654.86]
	Elderly	1728.41	[1490.55, 1966.28]
SC length (μm)	Young	142.86	[131.86, 153.86]
	Middle-aged	137.39	[122.86, 151.92]
	Elderly	177.58	[164.61, 190.55]
PL attachment site (μm)	Young	162.95	[140.77, 185.13]
	Middle-aged	139.49	[127.86, 151.13]
	Elderly	171.07	[150.02, 192.12]

Table S3: Results of multiple comparison Tukey's HSD test on TM- and PL-ratios. (*p<0.05, **p<0.017*)

Feature of interest	Age		p-value
TM-ratio	Young	Elderly	**0.030***
	Young	Middle-aged	0.094
	Middle-aged	Elderly	0.887
PL-ratio	Young	Elderly	**0.023***
	Young	Middle-aged	0.144
	Middle-aged	Elderly	0.620

724
725

726
Figure S1: Iris-OCT images, used to evaluate the IOP-induced deformations of the iris. We drew a
line segment (cyan) along the midline of the cornea (yellow lines), starting at the apex of the
iridocorneal angle and extending to the edge of the cornea typically visible in TM-OCT images. We
then constructed a perpendicular line segment connecting the end of the cyan line to the midline of the
iris (purple). We determined the length of this purple line segment along the 'y' axis for different
IOPs, and we evaluated iris displacement in the FE model based on the length change of this line
segment. See top-row panels for the definition of the 'y' axis.

Figure S2: Computed tissue deformations vs. IOP in three cases with different PL structures. **(A)** No
PL, **(B)** a single PL at the anterior end of the PL region, and **(C)** the homogenized (diffuse) PL model
described in the main text. The top row shows that, in the absence of a PL, SC collapse is predicted
even at low values of IOP (< 15 mmHg); such collapse is not observed experimentally. The second
row shows that a single PL is also not sufficient to prevent collapse at low IOP, especially when the
PL is located at the anterior aspect of the TM, which it often is. Only the diffuse (distributed) model
of the PL ligaments avoids SC collapse at low IOPs. Colors represent separate tissues (e.g., PL is
blue)

743
Figure S3: Description of the technique used to estimate PL solid fraction, . **(A)** A Scanning elec-
tron micrograph of rabbit PL (Simones et al., 1996) with the PLs outlined by a dashed rectangle (red).
White scale bar = 200 µm. **(B)** The radii of the PLs were determined by manually drawing line seg-
ments (red) in ImageJ, and the mean PL beam radius ( ) was computed.

The solid fraction within the relevant region of interest (ROI) was then determined by:

(i)   Projecting the red rectangle in panel (A) into the plane of the page by a distance , deter-
mined from a histological image of the New Zealand white rabbit angle (Ramos et al.,
2017), showing the distance from the iridocorneal angle to the most anterior PL as
.

(ii)   Computing   , where   is the number of PLs annotated in panel B
(;   is the total volume of the PLs within the ROI, assuming
cylindrically-shaped PLs; and   is the volume of the ROI.

This procedure yielded   , which is clearly only an approximate estimate and should be
refined in future work.