**Title:  Modulating the Unfolded Protein Response with ISRIB Mitigates Cisplatin Ototoxicity**

**Authors:**

Jiang Li[1]*, Stephanie L. Rouse[2,3]*, Ian R. Matthews[2], Elliott H. Sherr[1,4]**, and Dylan K. Chan[2]**

**Affiliations:**

1.  Department of Neurology, UCSF

2.  Department of Otolaryngology-Head and Neck Surgery, University of California, San Francisco (UCSF)

3.  Department of Otolaryngology & Neurology, Boston Children's Hospital and Harvard Medical School

4.  Department of Pediatrics, Institute of Human Genetics, Weill Institute for Neurosciences, UCSF

*These two authors share first position

**These two authors share last position

To whom correspondence should be addressed:

Dylan K. Chan, MD, PhD

Department of Otolaryngology-Head and Neck Surgery, UCSF

513 Parnassus Ave, Rm 719

San Francisco, CA 94143

dylan.chan@ucsf.edu

**Conflict of interest statement:**  EHS and DKC are co-owners of Jacaranda Biosciences, Inc.

EHS is co-inventor on a US provisional patent application: "Novel Methods of Treating Hearing

Loss."

**Abstract**

Cisplatin is a common chemotherapy drug with a nearly universal side effect of ototoxicity. The cellular mechanisms underlying cisplatin ototoxicity are poorly understood. Efforts in drug development to prevent or reverse cisplatin ototoxicity have largely focused on pathways of oxidative stress and apoptosis. An effective treatment for cisplatin ototoxicity, sodium thiosulfate, is associated with reduced survival in disseminated hepatoblastoma, highlighting the need for more specific drugs. The unfolded protein response (UPR) and endoplasmic reticulum (ER) stress pathways have been shown to be involved in the pathogenesis of noise-induced hearing loss and cochlear synaptopathy *in vivo*, and these pathways have been implicated broadly in cisplatin cytotoxicity. This study sought to determine whether the UPR can be targeted to prevent cisplatin ototoxicity. Neonatal cochlear cultures and HEK cells were exposed to cisplatin and UPR-modulating drugs, and UPR marker gene expression and cell death measured. Treatment with ISRIB, a drug that activates eif2B and downregulates the pro-apoptotic PERK/CHOP pathway of the UPR, was tested in an *in vivo* mouse model of cisplatin ototoxicity and well as a head and neck squamous cell carcinoma (HNSCC) cell-based assay of cisplatin cytotoxicity. Cisplatin exhibited a biphasic, non-linear dose-response of cell death and apoptosis that correlated with different patterns of UPR marker gene expression in HEK cells and cochlear cultures. ISRIB treatment protected against cisplatin-induced hearing loss and hair-cell death, but did not impact cisplatin's cytotoxic effects on HNSCC cell viability. These findings demonstrate that targeting the pro-apoptotic PERK/CHOP pathway with ISRIB can mitigate cisplatin ototoxicity without reducing anti-cancer cell effects, suggesting that this may be a viable strategy for drug development.

**Keywords:** cisplatin, ototoxicity, unfolded protein response, endoplasmic reticulum stress

**Introduction**

Cisplatin is an effective chemotherapeutic agent used to treat many malignant solid tumors, yet treatment is limited by ototoxicity. Irreversible sensorineural hearing loss affects 75-100% of patients treated with cisplatin, with young children more susceptible than adults (1-4). Despite some efficacy for sodium thiosulfate in the treatment of cisplatin ototoxicity in non-metastatic hepatoblastoma in children (5,6), significant concerns have been raised that STS reduces overall survival, possibly due to interfering with cisplatin's anti-cancer effects (7). There remains, therefore, a significant need for more specific pharmacological intervention to prevent ototoxicity while preserving cancer-cell cytotoxicity.

The cellular mechanism underlying cisplatin ototoxicity is poorly understood. Upon exposure, cisplatin is taken up into hair cells principally through the copper transporter Ctr1 (8) and binds DNA, leading to generation of excessive ROS, inflammation, transcriptional and cell cycle arrest, ultimately causing apoptosis and necrosis of sensory hair cells as well as cells in the stria vascularis and spiral ganglion (9-13). Efforts in drug development to prevent or reverse cisplatin ototoxicity have largely focused on pathways of oxidative stress and apoptosis (14). Exogenous administration of antioxidants mitigates hearing loss, but antioxidants are not a viable solution as antioxidant stress is also harmful to the cochlea (14).

There is extensive crosstalk between oxidative stress and endoplasmic reticulum (ER) stress (15), as up to 25% of ROS are generated in the ER (16), suggesting that ER dysfunction may be crucial to understanding cisplatin-induced cochlear cell apoptosis. ER stress occurs when improperly folded proteins accumulate in the ER lumen. In response, the unfolded protein response (UPR) is activated to either restore ER proteostasis, in the case of more transient stress, or induce apoptosis in severe stress (17).

Cisplatin induces ER stress (18); it directly interacts with proteins (19) and promotes their unfolding (20), and damage done to other organelles by cisplatin may act as an indirect inducer of ER stress (21). Cisplatin ototoxicity is correlated with cytosolic protein synthesis inhibition (22). In a rat model of cisplatin ototoxicity, BiP expression was broadly upregulated, indicating induction of ER stress. Enhancement of ER chaperone function and proteostasis with tauroursodexoycholic acid protected against cisplatin ototoxicity *in vivo*, indicating the UPR is a viable target for therapy against cisplatin ototoxicity (23). However, while ER stress has been implicated, the precise role of the UPR in cisplatin ototoxicity and cytotoxicity is not fully understood.

Our group implicated the UPR and ER stress in rapidly progressive genetic hearing loss due to absence of the novel deafness gene, TMTC4, as well as in noise induced hearing loss (NIHL) and cochlear synaptopathy (24,25). We demonstrated that the UPR is activated within the cochlea of adult mice within two hours of noise exposure, and that treating mice with ISRIB (Integrated Stress Response inhIBitor), a small molecule activator of eIF2B that results in downregulation of CHOP and the pro-apoptotic arm of the UPR (26), protects against NIHL and synapse loss.

Based on our findings implicating the UPR in genetic and noise-induced hearing loss, and the work of others implicating ER stress and the UPR in cisplatin ototoxicity (27,28), we sought to further explore the role of the UPR in the pathogenesis and treatment of cisplatin-induced ototoxicity. In this study, we extend our finding that the UPR is involved in NIHL, and that ISRIB can prevent NIHL, by showing that the UPR also mediates cisplatin ototoxicity. Most significantly, we demonstrate that ISRIB reduces cisplatin ototoxicity *in vivo* without affecting cisplatin's cytotoxic effects on head-and-neck squamous cell carcinoma (HNSCC) cells *in vitro*. Our findings support the idea that the UPR is a common early pathway in NIHL and cisplatin ototoxicity and may be an appealing target for therapeutic intervention.

**Results**

**Non-linear cisplatin-induced cell death is correlated with UPR gene expression**

To explore the mechanistic relationship between cisplatin cytotoxicity and UPR gene expression, we first measured UPR marker gene expression in response to a broad range of cisplatin doses in HEK cells. UPR apoptosis/cell fate is regulated predominately through the actions of the IRE1 and PERK arms sending pro-homeostatic and pro-apoptotic signals, respectively (**Figure 1A**). Death receptor 5 (DR5) integrates the opposing UPR signals of these two arms to control ER-stress-induced apoptosis: PERK/CHOP activity induces *DR5* transcription leading to formation of caspase 8 activating complex and ultimately apoptosis, whereas IRE1α/S-XBP1 promotes *DR5* mRNA decay, giving the cell time to restore homeostasis (29). Thus, DR5 levels are a measure of the persistence of ER stress and likelihood of apoptosis and are controlled by the opposing activities of the PERK/CHOP and IRE1α/S-XBP1 pathways.

Surprisingly, HEK cells exhibited a biphasic, bell-shaped response of UPR markers Chop, S-XBP1, and DR5 with increasing cisplatin dose, with a peak in expression of pro-apoptotic Chop and trough in expression of pro-homeostatic S-XBP1 at 50-100 $\mu$M cisplatin (**Figure 1B-C**). The Chop/S-XBP1 ratio and DR5 expression were similarly elevated in this range, consistent with integration of the Chop and S-XBP1 pathways towards apoptosis (**Figure 1D-E**). In accordance with this, we observed a peak in apoptosis, measured by caspase 3/7 activity, at 50 $\mu$M cisplatin (**Figure 1F**).

Observation of this correlation of UPR gene expression and apoptosis across this non-linear, bi-phasic dose response led us to hypothesize that the UPR is causally involved in, and may be targeted to reduce, cisplatin-induced apoptosis. To test this, we treated cells with 0.2 $\mu$M ISRIB, an eIF2B activator that inhibits the PERK/CHOP arm of the UPR (26) (**Figure 1A**)

and measured the effect on apoptosis across the same range of cisplatin doses. ISRIB reduced cisplatin-induced apoptosis at low cisplatin doses but had no effect at high cisplatin doses (**Figure 1G**), suggesting that it may be an effective drug to reduce cisplatin ototoxicity.

**Cisplatin treatment induces the Unfolded Protein Response in the cochlea *in vitro***

ER stress and the UPR has been broadly implicated in cytotoxicity induced by cisplatin treatment (18). We treated neonatal cochlear explant cultures with 100 $\mu$M cisplatin for 20h and performed RNAScope to detect Chop. Cisplatin-induced Chop expression was detected principally in cells of the stria vascularis and inner and outer sulcus supporting cells, but not in the hair-cell region (**Figure 2**).

We then sought to quantify the effect of cisplatin on UPR markers in the cochlea. Explant cultures of the P3-5 neonatal cochleae from wild-type C57BL/6J mice were treated with escalating concentrations of cisplatin for 20h and then subjected to whole-mount immnunohistochemistry against Myo7a, a hair-cell marker. Loss of inner and outer hair cells occurred with increasing cisplatin dose up to 100 $\mu$M; with higher doses, hair cells were paradoxically preserved up to 1000 $\mu$M cisplatin (**Figure 3A**).

Multiple groups have observed this bell-shaped dose-response of hair-cell death upon exposure of neonatal cochlear cultures to cisplatin: low cisplatin concentrations induce hair-cell death that increases with increasing cisplatin dose, whereas high cisplatin concentrations cause minimal hair-cell loss (22, 27-28). Based on our findings in HEK cells (**Figure 1**), we hypothesized that the UPR may play a role in this response.

We performed qPCR on murine cochlear explant cultures exposed to the same escalating doses of cisplatin for 6h, and measured Chop, S-XBP1, and DR5 expression. Expression of these three markers exhibited a similar bell-shaped dose-response curve with inflection at cisplatin doses of 25-100 $\mu$M cisplatin dose (**Figure 3B-E**). The highest expression

level of the pro-apoptotic marker Chop was seen at a cisplatin dose of 100 $\mu$M, whereas the lowest expression level of the homeostatic marker S-XBP1 was seen at cisplatin doses of 25 and 100 $\mu$M. The ratio of Chop to S-XBP1, which reflects the balance between pro-apoptotic and pro-homeostatic arms of the UPR, demonstrated the same bell-shaped dose-response, and DR5, which integrates the two pathways and therefore reflects this Chop/S-XBP1 ratio, was maximally expressed at 25 $\mu$M cisplatin.  The similarity between the unusual bell-shaped dose-response relationships of cisplatin with hair-cell death and pro-apoptotic UPR marker expression suggests that UPR activity may be involved in mediating cisplatin-induced hair-cell death.

**ISRIB protects against cisplatin-induced hearing loss and hair cell loss *in vivo***

These *in vitro* experiments suggest that cisplatin may induce hair-cell death via ER stress and induction of the UPR, and that ISRIB, an eIF2B activator, is effective at reducing cisplatin-induced apoptosis.  Based on previous work suggesting that targeting ER stress can reduce cisplatin ototoxicity (23) and our finding that ISRIB protects against ER stress and NIHL (24), we hypothesized that ISRIB would also be effective at mitigating cisplatin ototoxicity. To test this, we used a model of cisplatin ototoxicity in Balb/cJ mice, a strain that has been shown to have low mortality with cisplatin treatment (30).  A single intraperitoneal (IP) dose of 5.5 mg/kg cisplatin administered to 8-week-old female wild-type Balb/cJ mice caused reproducible, isolated 32-kHz ABR threshold elevation at post-injection-day (PID) 21, with minimal morbidity and no mortality (**Figure 4**).  Cisplatin-treated mice experienced a transient loss of 10% body weight at PID7, with full recovery to baseline by PID21, with no mortality observed (N=23), suggesting no overall impact on Balb/cJ morbidity.

10 mg/kg IP ISRIB was administered concurrent with cisplatin and serial ABR performed at PID7, 14, and 21.  Cisplatin-induced 32 kHz ABR threshold elevation was reduced in ISRIB-

treated mice compared with vehicle-treated controls at all 3 timepoints (**Figure 5A**, p < 0.01).

21 days after cisplatin injection (PID21), ISRIB-treated animals had a mean 32-kHz ABR

threshold of 58.3 ± 21.8 (mean ± SD, N=6), compared with 89.2 ± 17.4 dB SPL for vehicle-

treated controls, a statistically significant difference (p = 0.02, unpaired two-tailed t test).  In fact,

32-kHz ABR ISRIB-treated animals exhibited no significant elevation in 32-kHz ABR thresholds

(44.2 ± 10.7 dB SPL at baseline compared with 58.3 ± 21.8 dB SPL at PID21, p = 0.18).

Overall, ISRIB treatment was associated with a 69% reduction in cisplatin-induced threshold

shift compared to vehicle.

To evaluate further for ototoxicity, whole-mount immunohistochemistry was performed

after 21 days of cisplatin exposure to quantify outer-hair-cell (OHC) loss.  Whereas 57% ± 10%

(mean ± s.e.m., N=4) of basal-turn OHCs were lost in mice treated with cisplatin and vehicle,

mice co-treated with ISRIB and cisplatin exhibited little to no OHC loss (7.9% ± 5.9%, N=4, p

<0.01 compared to vehicle; **Figure 5B,C**).  These results demonstrate that ISRIB co-treatment

protects against cisplatin-induced hearing loss and OHC death in this *in vivo* model of cisplatin

ototoxicity.

**ISRIB does not affect the cytotoxicity of cisplatin against cancer cells *in vitro***

A principal concern for any drug treatment for ototoxicity is potential interference with the anti-

tumor effects of cisplatin.  To test this, we used the FaDu head and neck squamous-cell

carcinoma (HNSCC) cell line (gift of D.J.) (31).  Cisplatin treatment reduces FaDu cell viability,

with an $IC_{50}$ of 38 $\mu$M.  Co-treatment with 10 $\mu$M z-VAD-FMK, a caspase inhibitor known to

inhibit cisplatin-induced hair-cell apoptosis (32) as well as cisplatin chemotoxicity in cancer cells

(33), significantly increased the $IC_{50}$ of cisplatin (52 $\mu$M; $IC_{50}$ shift re: control = 1.205 (95% CI:

1.001-1.476); p < 0.05, **Figure 6**).  In contrast, co-treatment with 0.2 $\mu$M ISRIB, which prevented

cisplatin-induced apoptosis in HEK cells (**Figure 1A**), did not affect the $IC_{50}$ of cisplatin (37 $\mu$M; $IC_{50}$ shift re: control = 0.985 (95% CI: 0.905-1.073) in FaDu cells.  These findings demonstrate that ISRIB does not reduce the chemo-efficacy of cisplatin in this HNSCC cell line *in vitro*.

**Discussion**

Cisplatin is a common chemotherapeutic with the dose-limiting side effect of sensorineural hearing loss. However, the cellular mechanism underlying cisplatin ototoxicity is poorly understood.  Here, we present evidence suggesting that the UPR may play an important role in mediating cisplatin-induced ototoxicity.  Using a novel mouse model of cisplatin ototoxicity that produces consistent high-frequency hearing threshold elevation with minimal morbidity, we additionally show that modulation of the UPR with ISRIB protects against cisplatin-induced hearing loss and hair-cell death.  Unlike the pan-caspase inhibitor z-VAD-fmk, ISRIB does not impair the cytotoxicity of cisplatin in a HNSCC cell line.  These findings suggest that targeting eIF2b using ISRIB may be a promising strategy to treat cisplatin ototoxicity without interfering with cisplatin chemo-efficacy.

*Non-linear dose-response of cisplatin and hair-cell death*

We observed a bell-shaped dose response of hair-cell death upon exposure of neonatal cochlear cultures to cisplatin: low cisplatin concentrations induce hair-cell death that increases with increasing cisplatin dose, whereas high cisplatin concentrations give rise to little or no hair-cell loss. This phenomenon has been observed previously (27-28), but the mechanism is unknown. One possibility is that cisplatin uptake may be altered at high cisplatin concentrations (27).  Other studies have suggested that this phenomenon is an artifact of retention of hair cell

"corpses" that retain hair-cell markers (34-35). However, a more recent study showed this is not the case: 500 µm cisplatin treatment caused arrest of protein synthesis, but subsequent washout permitted recovery of protein synthesis and hair-cell survival (22).

We observed in HEK cells and cochlear explant cultures that expression of Chop, S-XBP1, and DR5, markers of UPR activation specifically in the apoptotic, homeostatic, and integrated pathways, respectively, all displayed similar, unusual bell-shaped dose-response curves. This correlation suggests that the UPR plays a mechanistic role in cisplatin ototoxicity. The increase in S-XBP1 expression at high cisplatin concentrations, in particular, suggests that this protective response may underlie the reduction in hair-cell loss at these high concentrations. ISRIB, an eIF2B activator that suppresses the pro-apoptotic activity of the UPR, protected against cisplatin-induced apoptosis in the lower, physiologically relevant range of cisplatin doses, further supporting a mechanistic role of the UPR in cisplatin ototoxicity.

Identification of different control pathways for cisplatin-induced cytotoxicity in different cell types could be of significant value in development of drugs to protect against cisplatin ototoxicity while retaining its cytotoxic properties for cancer chemotherapy. Studies in multiple cancer cell types have suggested that the homeostatic arm of the UPR plays a central role in chemoresistance (36-38). *In vitro*, cisplatin affects cancer cells and hair cells differently: whereby cochlear hair cells exhibit the non-linear, bell-shaped dose response, cancer cell lines such as head and neck squamous cell carcinoma exhibit a typical monotonic dose-response (39). It is possible that leveraging different dose-response characteristics relating to the UPR could lead to the development of UPR-targeting drugs that protect the cochlear while retaining anti-tumor activity. Indeed, we found that ISRIB, which specifically targets eIF2B, did not alter cisplatin's cytotoxic effect on a HNSCC cell line, providing strong initial evidence supporting this strategy. Multiple additional small molecule drugs have been developed to target different points in the homeostatic and apoptotic arms of the UPR. These can be used *in vitro* and *in vivo* to

further test the hypothesis that the UPR is a control mechanism and target for treatment of cisplatin ototoxicity.

*ISRIB protects against cisplatin ototoxicity*

The UPR is a potential target pathway for treatment of cisplatin ototoxicity.  For this study, we refined a single-dose mouse model of acute cisplatin ototoxicity.  Murine models of cisplatin ototoxicity are often complicated by high rates of mortality and a narrow ototoxic window compared with guinea pig and rat models (40). Prior studies have suggested that Balb/cJ mice have increased susceptibility to cisplatin ototoxicity and reduced mortality compared with CBA/CaJ and C57BL/6J strains (30,41).  Owing to its selective high-frequency hearing loss, simplicity, and minimal morbidity, the Balb/cJ strain may be useful for more rapid evaluation and screening of candidate drugs, which can subsequently be tested in a more clinically relevant chronic administration paradigm.

We found that a single IP dose of ISRIB, administered concurrently with cisplatin, significantly reduced cisplatin-induced high-frequency hearing loss by 69%, with near-complete protection of cochlear hair cells.  ISRIB, a potent activator of eiF2B and thereby UPR-mediated apoptosis, has been found to effectively modulate the UPR in many disease models, including neurodegeneration (42), memory loss (43), and pulmonary fibrosis (44).  We found that ISRIB effectively protects against NIHL (24) and cochlear synaptopathy (25).  The current findings suggest that the UPR may be a common mechanism in ototoxicity relating to noise and cisplatin. The efficacy of ISRIB corroborates prior findings: tauroursodeoxycholic acid, a non-specific modulator of the UPR, was effective at mitigating cisplatin ototoxicity *in vivo* (45-46), and salubrinal, which targets eIF2a and, like ISRIB, modulates the PERK/CHOP pathway,

protects against cisplatin toxicity *in vitro* (47).  Taken together, these findings suggest that targeting the UPR may be a valuable broad strategy to treat acquired hearing loss.

*Limitations and Future Directions*

This study demonstrates that the UPR is involved in cisplatin ototoxicity; however, several limitations exist. We are unable to resolve which cochlear cell types are involved in this response. RNAscope labelling suggests Chop is upregulated in strial and supporting cells of the inner sulcus and outer sulcus in response to cisplatin treatment, but more work is required to assess cell-type specificity. However, the UPR is a highly conserved mechanism, and we found that HEK cells exhibited similar behavior, including a similar inflection point of the biphasic response.  A more significant limitation is that the doses in which the paradoxical response was seen (> 100 µM) are well above those thought to be clinically relevant.  However, even if these high cisplatin doses are not clinically relevant, the mechanisms uncovered by these supratherapeutic conditions that lead to hair-cell protection may be relevant when considering drug development for cisplatin ototoxicity.

Despite these *in vitro* limitations, the importance of the UPR in cisplatin ototoxicity is most strongly illustrated by our finding that ISRIB, which inhibits the pro-apoptotic arm of the UPR, can robustly protect against cisplatin-induced high-frequency hearing loss and hair cell death *in vivo*. We used a single-dose acute cisplatin model, which is in contrast with the chronic model of cisplatin administration and ototoxicity that occurs clinically in humans and that has been described in mice (48).  Corroboration of this finding in a more clinically relevant chronic administration model of cisplatin ototoxicity would be valuable.

*Conclusion*

Cisplatin ototoxicity is a significant challenge with incompletely understood pathophysiology. This study provides evidence for the involvement of the UPR in cisplatin ototoxicity and demonstrates that targeting the pro-apoptotic arm of the UPR with ISRIB mitigates cisplatin-induced hearing loss and hair-cell death in mice without affecting cisplatin chemo-efficacy in a HSNCC cell line. Further investigation of the role of the UPR in cisplatin ototoxicity, and the potential for targeting it for treatment, is warranted.

**Methods**

*In vivo mouse model of cisplatin-induced ototoxicity*

As a single-dose model for acute cisplatin ototoxicity, 8-week-old female wild-type Balb/cJ mice (Jackson Laboratories, Strain #000651) were administered 5.5 mg/kg cisplatin (Sigma-Aldrich, 1134357 in normal saline) via intraperitoneal (IP) injection. To assess the effect of UPR modulation on cisplatin ototoxicity, we treated animals with ISRIB concurrent with cisplatin administration. Mice were injected intraperitoneally with 10 mg/kg ISRIB (Sigma, SML0843) or vehicle (50% DMSO, 50% PEG-400) immediately before cisplatin administration. Animals were administered 0.5 mL IP normal saline (NS) bolus for hydration concurrent with cisplatin and drug delivery, and subsequently on post injection days 1, 7, 14, and 21.

*Auditory testing*

Hearing was tested in mice by measuring auditory brainstem response (ABR) thresholds in response to broadband tone pips at 8, 16, and 32 kHz in the sound field using a standard

commercial system (RZ6, Tucker-Davis Technologies) in a soundproof chamber as described

(24). All ABR measurements were performed by an investigator blinded to drug treatment.

*Organotypic culture of the neonatal mouse cochlea*

Organotypic cultures from wild-type P3-5 C57BL/6J mice (Harlan, 057) were established as

described (24).  Briefly, the cochlear duct was isolated from P3 mice, opened, and plated on

glass coverslips with Cell-Tak (Corning, 354240) with the apical surface of the epithelium facing

up.  Cultures were used for experiments after 24 hours in culture. Cisplatin was applied to

cultures in media in increasing doses of cisplatin from 0-1000 $\mu$M for 20 hours.

*UPR gene expression analysis in cell culture*

Human embryonic kidney (HEK) cells were grown in DMEM supplemented with 10%

FBS, at 37 °C in 5% $CO_2$. To measure UPR marker gene expression in response to a

broad range of cisplatin doses, HEK cells were treated with cisplatin (0-400 $\mu$M) for 40

h, and then total RNA was isolated using TRIzol Reagent (Invitrogen, Carlsbad, CA). 1

µg total RNA was used for first-strand cDNA synthesis using SuperScript IV VILO

master mix (Invitrogen, Carlsbad, CA). Real-time PCR was performed by 7900HT FAST

Real-time PCR system (Applied Biosystems, Foster City, CA) with a SYBR green I mix.

The mRNA primers for UPR markers were as follows: *Chop*: CCACCACACCTGAAAGCAGAA

(forward), AGGTGAAAGGCAGGGACTCA (reverse); *S-XBP1*:

CTGAGTCCGAATCAGGTGCAG, GTCCATGGGAAGATGTTCTGG; *BiP*:

TTCAGCCAATTATCAGCAAACTCT, TTTTCTGATGTATCCTCTTCACCAGT; *DR5*:

TGCTGCTTGCTGTGCTACAGGCTGT, TTCTGACAGGTACTGGCCTGCTAG. Relative

quantification by the $2^{-\mathrm{DDCT}}$ method was used for analysis as described previously (24). To assess activation of Caspase 3 and 7, the Caspase-Glo® 3/7 assay was performed according to manufacturer's instructions (Promega) on HEK cells after 40 h treatment with cisplatin. To test if UPR directly causes cisplatin-induced apoptosis, HEK cells treated for 40 h with 0-400 $\mu$M cisplatin were additionally treated with 3 doses (at 0, 16 and 24 h) of 0.2 $\mu$M ISRIB and the Caspase-Glo® 3/7 assay performed.

*Immunohistochemistry*

Evaluation and quantification of hair-cell loss was performed with whole-mount cochlear immunohistochemistry as previously described (24). Briefly, cochlear explant cultures were treated as indicated and fixed in 4% paraformaldehyde. 8-week-old cisplatin-treated mice were euthanized 21 days after cisplatin treatment and temporal bones dissected, isolating the apical, middle, and basal cochlear turns. Specimens were incubated with rabbit anti-myosin7a antibody (a hair-cell-specific marker; 1:200 dilution in PBS; 25-6790, Proteus Biosciences) overnight at 4°C. Cochleae were rinsed three times for 15 min with PBS and then incubated for 2 hours with a goat anti-rabbit secondary antibody conjugated to Alexa Fluor 488 (1:1,000 dilution in PBS; Life Technologies). Whole mounts were rinsed in PBS three times for 15 min and mounted on glass slides in VectaShield antifade mounting medium (Vector Laboratories). The hair-cell region of the organ of Corti was imaged on a Nikon A1R upright line-scanning confocal microscope as previously described. For 8-week-old cochleae, the imaging locations for the apical, middle, and basal turns corresponded approximately to the 8, 16, and 32-kHz regions of the cochlea (25), and missing outer hair cells were counted from anti-Myosin7

immunostain in 200 μm cochlear segments and expressed as a percentage of missing outer hair cells.

*In situ hybridization using RNAScope*

To visualize UPR upregulation in the cochlea, we utilized RNAscope to detect CHOP RNA, as a marker of the UPR, and Myosin7a as a hair cell marker in cochlear explants treated with cisplatin. C57BL/6J wild-type mice (Harlan, 057) were dissected as described above and cultured overnight. 100 μM cisplatin was bath applied to cultures for 20 h. Thapsigargin (Sigma), an irreversible inhibitor of SERCA2b which leads to the induction of the UPR, was used as a positive control for induction of the UPR. 1 μM thapsigargin was bath applied to cultures for 2 h, followed by a 1 h fixation in 4% paraformaldehyde.

*In situ* RNA expression analysis was performed using RNAscope Fluorescent Multiplex Reagent Kit (Advanced Cell Diagnostics, Cat. No. 320850) and probes specific to Myosin7a and Chop (RNAscope Probe-Mm-Ddit3-C1, 317661; RNAscope Probe-Mm-Myo7a-C2, 462771-C2) using a protocol adapted from manufacturer's instructions and previously described protocol for cochlear tissue (49). Cultures were imaged by line-scanning confocal microscopy (Nikon A1R).

Cell viability in FaDu cells

FaDu cells, a head and neck squamous cell carcinoma line, were cultured in 96-well plates at a density of 10,000 cells per well. Cells were allowed to adhere and grow for 24 h before treatment. Serial dilutions of cisplatin were prepared in cell culture medium at a range of concentrations from 0-200 $\mu$M, with either 0.2 $\mu$M ISRIB or 10 $\mu$M z-VAD-FMK. Plates were incubated for 48 h and cell viability assay performed (Promega CellTiter-Glo 2.0).

Luminescence was recorded using a SynergyH4 microplate reader (Agilent Technologies, Hayward, CA) and IC$_{50}$ curves analyzed (Prism 9, GraphPad).

*Statistics*

For comparison between treatment groups for UPR gene expression, we used one-way ANOVA followed by Tukey multiple comparison tests. For pairwise comparison of ABR thresholds between groups of mice, we used unpaired two-tailed Student's t-test. Statistical analyses were performed with GraphPad Prism v9.

*Study approval*

All experimental protocols were approved by the UCSF Institutional Animal Care and Use Committee.

## Author Contributions

JL designed and conducted experiments, acquired and analyzed data, and co-wrote the manuscript.

SLR designed and conducted experiments, acquired and analyzed data, and co-wrote the manuscript.

IRM designed and conducted experiments, acquired and analyzed data, and reviewed and revised the manuscript.

EHS designed experiments, analyzed data, reviewed and revised the manuscript, and provided funding.

DKC designed and conducted experiments, acquired and analyzed data, co-wrote the manuscript, and provided funding.

## Acknowledgements

This study was funded by NIDCD R01 DC018583-01 (to DKC and EHS).

**References**

1. McKeage MJ. Comparative adverse effect profiles of platinum drugs. *Drug Saf.* 1995; 13:228–244.

2. Li Y, Womer RB, Silber JH. Predicting cisplatin ototoxicity in children: influence of age and the cumulative dose. *Eur. J. Cancer.* 2004; 40:2445–2451.

3. Rybak LP, Mukherjea D, Jajoo S, Ramkumar V. Cisplatin ototoxicity and protection: clinical and experimental studies. *Tohoku J Exp Med.* 2009; 219(3):177–186.

4. Langer T, am Zehnhoff-Dinnesen A, Radtke S, Meitert J, Zolk O. Understanding platinum-induced ototoxicity. Trends Pharmacol Sci. 2013 Aug;34(8):458-69.

5. Brock PR, Maibach R, Childs M, Rajput K, Roebuck D, Sullivan MJ, Laithier V, Ronghe M, Dall'Igna P, Hiyama E, Brichard B, Skeen J, Mateos ME, Capra M, Rangaswami AA, Ansari M, Rechnitzer C, Veal GJ, Covezzoli A, Brugières L, Perilongo G, Czauderna P, Morland B, Neuwelt EA. Sodium Thiosulfate for Protection from Cisplatin-Induced Hearing Loss. N Engl J Med. 2018 Jun 21;378(25):2376-2385.

6. Freyer DR, Brock PR, Chang KW, Dupuis LL, Epelman S, Knight K, Mills D, Phillips R, Potter E, Risby D, Simpkin P, Sullivan M, Cabral S, Robinson PD, Sung L. Prevention of cisplatin-induced ototoxicity in children and adolescents with cancer: a clinical practice guideline. Lancet Child Adolesc Health. 2020 Feb;4(2):141-150.

7. Orgel E, Villaluna D, Krailo MD, Esbenshade A, Sung L, Freyer DR. Sodium thiosulfate for prevention of cisplatin-induced hearing loss: updated survival from ACCL0431. Lancet Oncol. 2022 May;23(5):570-572. doi: 10.1016/S1470-2045(22)00155-3. PMID: 35489339; PMCID: PMC9635495.

8. Howell SB, Safaei R, Larson CA, Sailor MJ. Copper transporters and the cellular pharmacology of the platinum-containing cancer drugs. Mol Pharmacol. 2010; 77:887–894.

9. Zheng JL, Stewart RR, Gao WQ. Neurotrophin-4/5, brain-derived neurotrophic factor, and neurotrophin-3 promote survival of cultured vestibular ganglion neurons and protect them against neurotoxicity of ototoxins. J Neurobiol. 1995 Nov;28(3):330-40.

10. Kopke RD, Liu W, Gabaizadeh R, Jacono A, Feghali J, Spray D, Garcia P, Steinman H, Malgrange B, Ruben RJ, Rybak L, Van de Water TR. Use of organotypic cultures of Corti's organ to study the protective effects of antioxidant molecules on cisplatin-induced damage of auditory hair cells. Am J Otol. 1997 Sep;18(5):559-71.

11. Devarajan P, Savoca M, Castaneda MP, Park MS, Esteban-Cruciani N, Kalinec G, Kalinec F. Cisplatin-induced apoptosis in auditory cells: role of death receptor and mitochondrial pathways. Hear Res. 2002 Dec;174(1-2):45-54.

12. Rybak LP, Whitworth CA, Mukherjea D, Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and prevention. Hear Res. 2007 Apr;226(1-2):157-67.

13. St Germain C, Niknejad N, Ma L, Garbuio K, Hai T, Dimitroulakos J. Cisplatin induces cytotoxicity through the mitogen-activated protein kinase pathways and activating transcription factor 3. Neoplasia. 2010 Jul;12(7):527-38.

14. Sheth S, Mukherjea D, Rybak LP, Ramkumar V. Mechanisms of Cisplatin-Induced Ototoxicity and Otoprotection. *Front Cell Neurosci.* 2017;11:338.

15. Chaudhari N, Talwar P, Parimisetty A, Lefebvre d'Hellencourt C, Ravanan P. A molecular web: endoplasmic reticulum stress, inflammation, and oxidative stress. *Front Cell Neurosci.* 2014; 8:213.

16. Tu BP, Weissman JS. Oxidative protein folding in eukaryotes: mechanisms and consequences. J Cell Biol. 2004 Feb 2;164(3):341-6.

17. Walter P and Ron D. The unfolded protein response: from stress pathway to homeostatic regulation. *Science.* 2011; 334(6059):1081-6.

18. Foufelle F, Fromenty B. Role of endoplasmic reticulum stress in drug-induced toxicity. *Pharmacol Res Perspect*. 2016;4(1):e00211.

19. Karasawa T, Sibrian-Vazquez M, Strongin RM, Steyger PS. Identification of Cisplatin-Binding Proteins Using Agarose Conjugates of Platinum Compounds. 2013. *PLoS* One. 8(6): e66220

20. Mezencev, R. Interactions of cisplatin with non-DNA targets and their influence on anticancer activity and drug toxicity:the complex world of the platinum complex. *Curr Cancer Drug Targets.* 2015; 14(9):794-816.

21. Sancho-Martínez SM, Prieto-García L, Prieto M, López-Novoa JM, López-Hernández FJ. Subcellular targets of cisplatin cytotoxicity: an integrated view. Pharmacol Ther. 2012 Oct;136(1):35-55.

22. Nicholas BD, Francis S, Wagner EL, Zhang S, Shin JB. Protein Synthesis Inhibition and Activation of the c-Jun N-Terminal Kinase Are Potential Contributors to Cisplatin Ototoxicity. Front Cell Neurosci. 2017 Sep 27;11:303.

23. Zong S, Liu T, Wan F, Chen P, Luo P, Xiao H: Endoplasmic Reticulum Stress Is Involved in Cochlear Cell Apoptosis in a Cisplatin-Induced Ototoxicity Rat Model. *Audiol Neurotol*. 2017; 22:160-168.

24. Li J, Akil O, Rouse SL, McLaughlin CW, Matthews IR, Chan DK, Sherr EH. Deletion of Tmtc4 activates the unfolded protein response and causes postnatal hearing loss. *J Clin Invest*. 2018; 128(11):5150-5162.

25. Rouse SL, Matthews IR, Li J, Sherr EH, Chan DK. Integrated stress response inhibition provides sex-dependent protection against noise-induced cochlear synaptopathy. Sci Rep. 2020 Oct 22;10(1):18063.

26. Sidrauski C, McGeachy A, Ingolia N, Walter P. The small molecule ISRIB reverses the effect of eIF2a phosphorylation on translation and stress granule assembly. *eLife*. 2015; 4:e05033.

27. Ding D, He J, Allman BL, Yu D, Jiang H, Seigel GM, Salvi RJ. Cisplatin ototoxicity in rat cochlear organotypic cultures. Hear Res. 2011; 282:196–203.

*28.* Zeng S, Sun X, Chen Z, Yu D, Chen B, Yin S. Low, but Not High, Doses of Cisplatin Damage Cochlear Hair Cells in C57 Mouse Organotypic Cultures. *ORL J Otorhinolaryngol Relat Spec.* 2016; 78:177-186.

29. Lu M, Lawrence DA, Marsters S *et al.* Opposing unfolded-protein-response signals converge on death receptor 5 to control apoptosis. *Science.* 2014; 345:98–101.

30. DeBacker JP, Harrison RY, Bielefeld EC (2020). Cisplatin-induced threshold shift in the CBA/CaJ, C57BL/6J, BALB/cJ mouse models of hearing loss. Hear Res 387:107878.

31. Lin CJ, Grandis JR, Carey TE, Gollin SM et al. Head and neck squamous cell carcinoma cell lines: established models and rationale for selection. *Head Neck.* 2007; 29(2):163-88.

32. Li Y, Li A, Wu J, He Y, Yu H, Chai R, Li H. MiR-182-5p protects inner ear hair cells from cisplatin-induced apoptosis by inhibiting FOXO3a. Cell Death Dis. 2016 Sep 8;7(9):e2362. doi: 10.1038/cddis.2016.246. PMID: 27607577; PMCID: PMC5059852.

33. Kim JS, Lee JH, Jeong WW, Choi DH, Cha HJ, Kim DH, Kwon JK, Park SE, Park JH, Cho HR, Lee SH, Park SK, Lee BJ, Min YJ, Park JW. Reactive oxygen species-dependent EndoG release mediates cisplatin-induced caspase-independent apoptosis in human head and neck squamous carcinoma cells. Int J Cancer. 2008 Feb 1;122(3):672-80. doi: 10.1002/ijc.23158. PMID: 17955488.

34. Monzack EL, May LA, Roy S, Gale JE, Cunningham LL. Live imaging the phagocytic activity of inner ear supporting cells in response to hair cell death. Cell Death Differ. 2015 Dec;22(12):1995-2005.

35. Anttonen T, Belevich I, Kirjavainen A, Laos M, Brakebusch C, Jokitalo E, Pirvola U. How to bury the dead: elimination of apoptotic hair cells from the hearing organ of the mouse. J Assoc Res Otolaryngol. 2014 Dec;15(6):975-92.

36. Cubillos-Ruiz JR, Bettigole SE, Glimcher LH.  Tumorigenic and immunosuppressive effects of endoplasmic reticulum stress in cancer.  *Cell.* 2017; 168(4):692-706.

37. Aran G, Sanjurjo L, Barcena C *et al.* CD5L is upregulated in hepatocellular carcinoma and promotes liver cancer cell proliferation and antiapoptotic responses by binding to HSPA5 (GRP78). *FASEB J.* 2018; 32:3878-3891.

38. Xu D, Liang SQ, Yang H *et al.* Increased sensitivity to apoptosis upon endoplasmic reticulum stres-induced activation of the unfolded protein response in chemotherapy-resistant malignant pleural mesothelioma. *Br J Cancer.* 2018; 119:65-75.

39. Schuler PJ, Trellakis S, Greve J *et al.* In vitro chemosensitivity of head and neck cancer cell lines.  *Eur J Med Res.* 2010; 15:337-344.

40. Poirrier AL, Van den Ackerveken P, Kim TS, Vandenbosch R, Nguyen L, Lefebvre PP, Malgrange B. Ototoxic drugs: difference in sensitivity between mice and guinea pigs. Toxicol Lett. 2010 Mar 1;193(1):41-9.

41. Chang L, Wang A (2013).  Calpain mediated cisplatin-induced ototoxicity in mice.  Neural Regen Res 8_21):1995-2002.

42. Bugallo R, Marlin E, Baltanás A, Toledo E, Ferrero R, Vinueza-Gavilanes R, Larrea L, Arrasate M, Aragón T. Fine tuning of the unfolded protein response by ISRIB improves neuronal survival in a model of amyotrophic lateral sclerosis. Cell Death Dis. 2020 May 26;11(5):397.

43. Krukowski K, Nolan A, Frias ES, Boone M, Ureta G, Grue K, Paladini MS, Elizarraras E, Delgado L, Bernales S, Walter P, Rosi S. Small molecule cognitive enhancer reverses age-related memory decline in mice. Elife. 2020 Dec 1;9:e62048.

44. Watanabe S, Markov NS, Lu Z, Piseaux Aillon R, Soberanes S, Runyan CE, Ren Z, Grant RA, Maciel M, Abdala-Valencia H, Politanska Y, Nam K, Sichizya L, Kihshen HG, Joshi N, McQuattie-Pimentel AC, Gruner KA, Jain M, Sznajder JI, Morimoto RI, Reyfman PA,

Gottardi CJ, Budinger GRS, Misharin AV. Resetting proteostasis with ISRIB promotes epithelial differentiation to attenuate pulmonary fibrosis. Proc Natl Acad Sci U S A. 2021 May 18;118(20):e2101100118.

45. Lee, C.H.; Park, S.S.; Lee, D.H.; Lee, S.M.; Kim, M.Y.; Choi, B.Y.; Kim, S.Y. Tauroursodeoxycholic acid attenuates cisplatin-induced hearing loss in rats. *Neurosci. Lett.* 2020, *722*, 134838.

46. Wen Y, Zong S, Liu T, Du P, Li H, Xiao H. Tauroursodeoxycholic acid attenuates cisplatin-induced ototoxicity by inhibiting the accumulation and aggregation of unfolded or misfolded proteins in the endoplasmic reticulum. Toxicology. 2021 Apr 15;453:152736.

47. Lu W, Ni K, Li Z, Xiao L, Li Y, Jiang Y, Zhang J, Shi H. Salubrinal Protects Against Cisplatin-Induced Cochlear Hair Cell Endoplasmic Reticulum Stress by Regulating Eukaryotic Translation Initiation Factor 2α Signalling. Front Mol Neurosci. 2022 May 30;15:916458.

48. Fernandez K, Wafa T, Fitzgerald TS, Cunningham LL. An optimized, clinically relevant mouse model of cisplatin-induced ototoxicity. Hear Res. 2019 Apr;375:66-74.

49. Kersigo J, Pan N, Lederman JD (2018).  A RNAscope whole mount approach that can be combined with immunofluorescence to quantify differential distribution of mRNA.  Cell Tissue Res 374(2):251-262.

**Figure Legends**

**Figure 1. Non-linear cisplatin dose-response of UPR gene expression and apoptosis in HEK cells.** **A.** In the unfolded protein response (UPR), two major pathways, labeled by their initiating effectors (IRE1a, and PERK) respond to accumulation of misfolded or unfolded proteins by facilitating homeostasis or, if the load is too great, apoptosis. The downstream effectors S-XBP1 and Chop are commonly used markers of the homeostatic (S-XBP1) and apoptotic (Chop) pathways. Depending on persistence and severity of stress, indicated by DR5, which integrates S-XBP1 and Chop signals, the UPR halts translation and allows for an adaptive response and return to proteostasis, or terminal response ending in apoptosis. ISRIB inhibits the PERK pathway by activating eiF2B and inhibiting ATF4. **B-E. UPR gene expression**.  Wild-type HEK cells were treated with multiple doses of cisplatin, and mRNA levels of Chop, S-XBP1, the ratio of Chop/S-XBP1, and DR5 were measured by qPCR and normalized to GAPDH by the $2^{-\Delta\Delta CT}$ method. **F. Apoptosis.** Caspase 3/7 activity was quantified using Caspase-Glo, demonstrating a non-linear cisplatin dose-dependence of HEK cells that matches DR5 expression. **B-F.** Data are means ± SEM, with individual data shown as grey dots. * $p < 0.05$; ** $p < 0.001$ by one-way ANOVA followed by pairwise t-test relative to control (0 $\mu$M cisplatin). Each graph is from a single experiment with 3 (**B-E**) or 5 (**F**) independent samples for each cisplatin dose. The experiment was repeated three times with identical results. **G. ISRIB treatment in cisplatin-treated cells reduces apoptosis.** HEK cells treated for 40 h with cisplatin were additionally treated with 3 doses (at 0, 16, and 24 h) of 0.2 $\mu$M ISRIB (red), or vehicle (black), and Caspase 3/7 activity measured. Increasing cisplatin concentration induced an initial increase, then decrease, in apoptosis in the absence of ISRIB, consistent with the non-linear pattern of hair-cell death seen in cisplatin-treated cochlear cultures. At low cisplatin

doses, ISRIB attenuated cisplatin-induced apoptosis and eliminated cisplatin dose-dependence; at high cisplatin doses, ISRIB had no effect. Data are means ± SEM, with individual values as pink or grey dots. *p<0.05; **p<0.01 compared to no drug. N=3 for each condition.

**Figure 2. Chop expression detected in cochlear cultures in response to TG and cisplatin.** RNAscope probing for Myo7a (green) and Chop (red) in P2 C57 cochlear cultures after 2 h treatment with thapsigargin (TG, a known inducer of the UPR, middle) shows increased expression of Chop in TG-treated cultures, particularly in the hair-cell region (HC). Cultures treated with 100 $\mu$M cisplatin (right) for 20h demonstrated upregulation of Chop in stria vascularis (SV) and supporting cells of the inner and outer sulcus (IS/OS), but not hair cells. Scale bars: 50 $\mu$m.

**Figure 3. UPR modulation in cisplatin-treated cochleae. A.** Non-linear hair-cell death in response to cisplatin. Bell-shaped dose response in hair-cell death is observed in P3 WT C57BL/6J organotypic cochlear cultures treated with cisplatin at increasing doses (0-1000 $\mu$M) for 24 h then fixed and stained with anti-Myo7a to detect hair cells. At low concentrations of cisplatin, hair-cell death increases with increasing cisplatin concentration, but above 100 $\mu$M, cisplatin-treated cochleae had preserved hair cells. **B-E.** Non-linear UPR gene expression in cochlear cultures quantified by qPCR after 6 h exposure to cisplatin. Chop (**C**), a marker of the pro-apoptotic arm of the UPR, and S-XBP1 (**C**), a pro-homeostatic UPR marker, displayed inverse non-linear dose-dependences. **D.** The ratio of Chop/S-XBP1 demonstrated elevated balance towards pro-apoptotic Chop expression consistent with the pattern of hair-cell death seen in **A.** The dose-dependency of this ratio was consistent with the expression pattern of DR5, a downstream marker that integrates the Chop and S-XBP1 signals and signals towards

apoptosis. **B-E**. Data are $2^{-\Delta\Delta CT}$ measurements compared internally to GAPDH and normalized

to control, non-treated cultures from the same experimental run, and are presented as means ±

SEM, with individual data shown as grey dots. *p < 0.05; **p < 0.001 by one-way ANOVA

relative to control (0 μM cisplatin).  Scale bars: 20 μm.

**Figure 4.  Balb/cJ model of murine cisplatin ototoxicity.**  A single intraperitoneal dose of 5.5

mg/kg cisplatin (CP) or saline was administered to 8-week-old female Balb/cJ mice.  **A.**

Compared to control mice (N=17), weight decreased slightly in the CP group (N=23) but

returned to baseline by post-injection-day (PID)21. No mortality occurred. **B-D.** ABR thresholds

to 8, 16, and 32 kHz pure-tone pips were recorded at PID3, 7, 14, and 21.  Data shown are

means ± SEM, with individual data shown in gray dots.

**Figure 5. ISRIB protects against cisplatin-induced hearing and hair-cell loss**.  8-week-old

Balb/cJ mice were treated with a single intraperitoneal dose of 5.5 mg/kg cisplatin, and either 10

mg/kg ISRIB or vehicle. ABR thresholds to pure-tone pips were measured 7, 14, and 21 days

after cisplatin treatment. After 21 days, mice were euthanized and temporal bones harvested. **A.**

32-kHz ABR threshold measurements demonstrated significant protection against cisplatin-

induced hearing loss by in ISRIB-treated animals (gray) compared to controls (white).  **B.**

Cochlear whole mounts from the apical, middle, and basal turns were stained with Myo7a and

visualized by confocal microscopy, demonstrating significant hair-cell loss in cisplatin + vehicle

treated animals (left), worse in the middle and basal turns. Cochleae from animals treated with

ISRIB had better hair-cell preservation after cisplatin exposure.  **C.** Hair cells were quantified as

the percentage of missing hair cells in a fixed-length segment from the cochlear basal, middle,

or apical turns from mice 21 days after treatment with cisplatin and either ISRIB (gray, N=4) or

vehicle (white, N=4).  Data shown are means ± SEM, with individual data shown in grey dots. **

*p* < 0.01; NS; not significant.  Scale bars: 20 $\mu$m.

**Figure 6. ISRIB does not interfere with cisplatin cytotoxicity against cancer cells.**  Viability

of FaDu cells, a head and neck squamous-cell carcinoma (HNSCC) cell line, was measured in

presence of escalating doses of cisplatin alone (black) or in combination with 10 $\mu$M z-VAD-

FMK (red), an apoptosis inhibitor, or 0.2 $\mu$M ISRIB (blue).  Data shown are means ± SEM (N = 3

for each condition).

**Figure 1. Non-linear cisplatin dose-response of UPR gene expression and apoptosis in HEK cells. A.** In the unfolded protein response (UPR), two major pathways, labeled by their initiating effectors (IRE1a, and PERK) respond to accumulation of misfolded or unfolded proteins by facilitating homeostasis or, if the load is too great, apoptosis. The downstream effectors S-XBP1 and Chop are commonly used markers of the homeostatic (S-XBP1) and apoptotic (Chop) pathways. Depending on persistence and severity of stress, indicated by DR5, which integrates S-XBP1 and Chop signals, the UPR halts translation and allows for an adaptive response and return to proteostasis, or terminal response ending in apoptosis. ISRIB inhibits the PERK pathway by activating eiF2B and inhibiting ATF4. **B-E. UPR gene expression**. Wild-type HEK cells were treated with multiple doses of cisplatin, and mRNA levels of Chop, S-XBP1, the ratio of Chop/S-XBP1, and DR5 were measured by qPCR and normalized to GAPDH by the $2^{-\Delta\Delta CT}$ method. **F. Apoptosis.** Caspase 3/7 activity was quantified using Caspase-Glo, demonstrating a non-linear cisplatin dose-dependence of HEK cells that matches DR5 expression. **B-F.** Data are means ± SEM, with individual data shown as grey dots. * $p < 0.05$; ** $p < 0.001$ by one-way ANOVA followed by pairwise t-test relative to control (0 µM cisplatin). Each graph is from a single experiment with 3 (**B-E**) or 5 (**F**) independent samples for each cisplatin dose. The experiment was repeated three times with identical results. **G. ISRIB treatment in cisplatin-treated cells reduces apoptosis.** HEK cells treated for 40 h with cisplatin were additionally treated with 3 doses (at 0, 16, and 24 h) of 0.2 µM ISRIB (red), or vehicle (black), and Caspase 3/7 activity measured. Increasing cisplatin concentration induced an initial increase, then decrease, in apoptosis in the absence of ISRIB, consistent with the non-linear pattern of hair-cell death seen in cisplatin-treated cochlear cultures. At low cisplatin doses, ISRIB attenuated cisplatin-induced apoptosis and eliminated cisplatin dose-dependence; at high cisplatin doses, ISRIB had no effect. Data are means ± SEM, with individual values as pink or grey dots. *$p<0.05$; **$p<0.01$ compared to no drug. N=3 for each condition.

**Figure 2. Chop expression detected in cochlear cultures in response to TG and cisplatin.** RNAscope probing for Myo7a (green) and Chop (red) in P2 C57 cochlear cultures after 2 h treatment with thapsigargin (TG, a known inducer of the UPR, middle) shows increased expression of Chop in TG-treated cultures, particularly in the hair-cell region (HC). Cultures treated with 100 $\mu$M cisplatin (right) for 20h demonstrated upregulation of Chop in stria vascularis (SV) and supporting cells of the inner and outer sulcus (IS/OS), but not hair cells. Scale bars: 50 $\mu$m.

**Figure 3. UPR modulation in cisplatin-treated cochleae. A.** Non-linear hair-cell death in response to cisplatin. Bell-shaped dose response in hair-cell death is observed in P3 WT C57BL/6J organotypic cochlear cultures treated with cisplatin at increasing doses (0-1000 µM) for 24 h then fixed and stained with anti-Myo7a to detect hair cells. At low concentrations of cisplatin, hair-cell death increases with increasing cisplatin concentration, but above 100 µM, cisplatin-treated cochleae had preserved hair cells. **B-E.** Non-linear UPR gene expression in cochlear cultures quantified by qPCR after 6 h exposure to cisplatin. Chop (**C**), a marker of the pro-apoptotic arm of the UPR, and S-XBP1 (**C**), a pro-homeostatic UPR marker, displayed inverse non-linear dose-dependences. **D.** The ratio of Chop/S-XBP1 demonstrated elevated balance towards pro-apoptotic Chop expression consistent with the pattern of hair-cell death seen in **A.** The dose-dependency of this ratio was consistent with the expression pattern of DR5, a downstream marker that integrates the Chop and S-XBP1 signals and signals towards apoptosis. **B-E**. Data are $2^{-\Delta\Delta CT}$ measurements compared internally to GAPDH and normalized to control, non-treated cultures from the same experimental run, and are presented as means ± SEM, with individual data shown as grey dots. *p < 0.05; **p < 0.001 by one-way ANOVA relative to control (0 µM cisplatin). Scale bars: 20 µm.

**Figure 4. Balb/cJ model of murine cisplatin ototoxicity.** A single intraperitoneal dose of 5.5 mg/kg cisplatin (CP) or saline was administered to 8-week-old female Balb/cJ mice. **A.** Compared to control mice (N=17), weight decreased slightly in the CP group (N=23) but returned to baseline by post-injection-day (PID)21. No mortality occurred. **B-D.** ABR thresholds to 8, 16, and 32 kHz pure-tone pips were recorded at PID3, 7, 14, and 21. Data shown are means ± SEM, with individual data shown in gray dots.

**Figure 5. ISRIB protects against cisplatin-induced hearing and hair-cell loss**. 8-week-old Balb/cJ mice were treated with a single intraperitoneal dose of 5.5 mg/kg cisplatin, and either 10 mg/kg ISRIB or vehicle. ABR thresholds to pure-tone pips were measured 7, 14, and 21 days after cisplatin treatment. After 21 days, mice were euthanized and temporal bones harvested. **A.** 32-kHz ABR threshold measurements demonstrated significant protection against cisplatin-induced hearing loss by in ISRIB-treated animals (gray) compared to controls (white). **B.** Cochlear whole mounts from the apical, middle, and basal turns were stained with Myo7a and visualized by confocal microscopy, demonstrating significant hair-cell loss in cisplatin + vehicle treated animals (left), worse in the middle and basal turns. Cochleae from animals treated with ISRIB had better hair-cell preservation after cisplatin exposure. **C.** Hair cells were quantified as the percentage of missing hair cells in a fixed-length segment from the cochlear basal, middle, or apical turns from mice 21 days after treatment with cisplatin and either ISRIB (gray, N=4) or vehicle (white, N=4). Data shown are means +/- SEM, with individual data shown in grey dots. ** $p < 0.01$; NS; not significant. Scale bars: 20 μm.

Cisplatin    IC$_{50}$=38$\mu$M   R$^2$=0.95
Cisplatin+ISRIB    IC$_{50}$=37$\mu$M   R$^2$=0.96
Cisplatin+Z-VAD-FMK  IC$_{50}$=52$\mu$M   R$^2$=0.92

**Figure 6. ISRIB does not interfere with cisplatin cytotoxicity against cancer cells.** Viability of FaDu cells, a head and neck squamous-cell carcinoma (HNSCC) cell line, was measured in presence of escalating doses of cisplatin alone (black) or in combination with 10 $\mu$M z-VAD-FMK (red), an apoptosis inhibitor, or 0.2 $\mu$M ISRIB (blue). Data shown are means $\pm$ SEM (N = 3 for each condition).