Locus coeruleus neuromelanin accumulation and dissipation across the lifespan

Authors: Elizabeth Riley[1], Nicholas Cicero[1], Khena Swallow[1], Eve De Rosa[1], Adam Anderson[1]
1: Department of Psychology, Cornell University

Corresponding author: Elizabeth Riley, er482@cornell.edu

Keywords: neuromelanin, locus coeruleus, aging, cognition

Abstract:
The pigment neuromelanin, produced in the locus coeruleus (LC) as a byproduct of catecholamine synthesis, gives the "blue spot" its name, and both identifies LC neurons and is thought to play an important yet complex role in normal and pathological aging. Using neuromelanin-sensitive T1-weighted turbo spin echo MRI scans we characterized volume and neuromelanin signal intensity in the LC of 96 participants between the ages of 19 and 86. Although LC volume did not change significantly throughout the lifespan, LC neuromelanin signal intensity increased from early adulthood, peaked around age 60 and precipitously declined thereafter. Neuromelanin intensity was greater in the caudal relative to rostral extent and in women relative to men. With regard to function, rostral LC neuromelanin intensity was associated with fluid cognition in older adults (60+) only in those above the 50th percentile of cognitive ability for age. The gradual accumulation of LC neuromelanin across the lifespan, its sudden dissipation in later life, and relation to preserved cognitive function, is consistent with its complex role in normal and pathological aging.

Introduction:

The locus coeruleus, located in the pons, is part of the reticular activating system and the major source of modulatory neurochemical norepinephrine for the brain (Aston-Jones & Cohen, 2005; Berridge & Waterhouse, 2003; Sara & Bouret, 2012). Research has linked the status of the locus coeruleus norepinephrine system to cognitive health and the underpinnings of multiple neurodegenerative diseases (Feinstein et al., 2016), including Alzheimer's Disease (Beardmore et al., 2021; Cassidy et al., 2022; Chan-Palay & Asan, 1989) and Parkinson's Disease (Gesi et al., 2000; Zarow et al., 2003). The LC is vulnerable to pathological change extremely early in the disease process (Braak et al., 2011; Cassidy et al., 2022), especially in its accumulation of hyperphosphorylated tau (Chalermpalanupap et al., 2017), which can spread throughout its extensive dendritic arbor to the cortex (Ghosh et al., 2019). In addition to this role in spreading pathology, the loss of LC neurons results in dysfunction due to the crucial role that the norepinephrine system plays in sleep (Van Egroo et al., 2022), arousal (Ross & Van Bockstaele, 2021), attention (Plini et al., 2021), cognitive reserve (Clewett et al., 2015), and cognitive control (Aston-Jones et al., 2000). Pathological changes in the LC take place years or even decades before symptomatic illness and may be detectable in those at risk for disease (Granholm et al., 2017; Kremen et al., 2019; Plini et al., 2021). For all these reasons, there is great interest in assessing the LC in vivo so that disease states may be predicted or prevented.

LC measurement in vivo rests primarily on neuromelanin-sensitive MRI sequences (Clewett et al., 2015; Sulzer et al., 2018; Trujillo et al., 2023). Neuromelanin is a pigment that is

produced as a byproduct of catecholamine synthesis, in a reaction that requires iron and results in the storage of iron-containing neuromelanin polymers in permanent granules within neurons (Nelson et al., 2009; Shibata et al., 2006; van der Pluijm et al., 2021). Neuromelanin cannot be efficiently cleared by the cell, and thus builds up throughout the lifespan, at least until neurons are lost (Haining & Achat-Mendes, 2017). LC can be reliably visualized as hyperintense on MRI with neuromelanin-specific sequences (Langley et al., 2016).

The relationship between neuromelanin and neuron health is complex. Since neuromelanin serves as a de facto record of former norepinephrine production, it could be said to cumulatively index prior mental engagement and arousal levels (Clewett et al., 2015). Altered LC activity and norepinephrine release are also associated with psychiatric conditions such as depression (Guinea-Izquierdo et al., 2021; Klimek et al., 1997) and anxiety or PTSD (Celebi, 2022; Ross & Van Bockstaele, 2021) In the context of cognitive aging, sustained mental engagement is associated with better cognitive outcomes (Ryff et al., 2016). Likely for this reason, some research has uncovered a positive relationship between neuromelanin intensity and cognitive performance (Clewett et al., 2015), including future cognitive performance (Dahl et al., 2023), in older adults. However, neuromelanin's complex role within LC neurons complicates this picture.

Granules of neuromelanin can absorb oxidative radicals in the brain, thus protecting the LC from various toxins (Haining & Achat-Mendes, 2017; Moreno-García et al., 2021; Zucca et al., 2017). The LC has inevitable toxin exposure due to its role in making dopamine and norepinephrine, both of which progress to highly toxic metabolites (Goldstein et al., 2013; Kang et al., 2020), despite their necessity and benefits to cognition. Additionally, the LC accumulates environmental toxins such as heavy metals more than surrounding structures, potentially due to increased exposure through its extensive dendritic arbor in addition to its ability to sequester such toxins (Pamphlett et al., 2018, 2020). These toxins remain sequestered in the neuromelanin granules indefinitely (Haining & Achat-Mendes, 2017; Iannitelli et al., 2023). If the granules are damaged, toxins then spill out resulting in inflammation and cell death (Zecca et al., 2003; Zucca et al., 2017). The conditions that cause damage to LC neuromelanin granules appear to include hyperphosphorylated tau accumulation (Jacobs et al., 2019), and at least in animal models, and an overaccumulation of neuromelanin itself, beyond some unknown saturation point (Carballo-Carbajal et al., 2019; Iannitelli et al., 2023). Thus, the presence of neuromelanin granules is neither unequivocally good or bad, and their value or risk to LC neurons is difficult to determine especially in older adults.

Within an individual, there may come a tipping point, before which neuromelanin accumulation is the happy result of cumulative mental engagement which provides detoxification of a vital brain nucleus, and after which neurons are being lost due to overload of either toxins or neuromelanin itself. This may explain why some previous post mortem and in vivo studies have detected a peak in LC neuromelanin around age 50-60 (Liu et al., 2019; Manaye et al., 1995; Mann et al., 1985; Shibata et al., 2006). A similar inverted-U life course has been found for the substantia nigra (Xing et al., 2018), the other neuromelanin-containing structure in the brain. For older adults, larger amounts of remaining neuromelanin may not necessarily be beneficial, but may indicate that such individuals have not yet reached their neuromelanin tipping point, leaving their LC vulnerable.

In attempting to understand the diagnostic or prognostic value of neuromelanin imaging, it is crucial to understand the typical life course of this enigmatic molecule, and to attempt to refine the relationship between its presence and cognitive function. Recent efforts by others have strongly suggested an inverted-U shaped curve of neuromelanin intensity across the lifespan (Liu et al., 2019). This result is important to replicate particularly from middle to older age. Given demographic disparities in Alzheimer's related dementias it is also imperative to examine groups from diverse racial backgrounds (Babulal et al., 2019). Most complex is understanding the relationship between neuromelanin intensity and cognition, which is unclear, given neuromelanin can represent both a protective factor and a vulnerability depending on the life stage. These important factors deserve exploration. To that end, we present an analysis of the LC that maximizes neuromelanin contrast to characterize its LC volume and neuromelanin signal intensity in 96 individuals aged 19-86, with comparisons by sex and race, and differences by cognitive functioning, using a method of neuromelanin intensity measurement (Turker et al., 2021) with high sensitivity and reliability across the lifespan.

Results:

Localization of the LC: We identified the locus coeruleus (LC) in 96 participants between the ages of 19-86 using neuromelanin-sensitive turbo spin echo T1-weighted MRI scans. The underlying mechanism for contrast on such scans remains unclear (Oshima et al., 2021; Priovoulos et al., 2020) but has previously thought to be related to the T1-shortening effects of neuromelanin, thanks to its iron binding ability (Sulzer et al., 2018). All LC intensity and volume measurements were made using individually hand-drawn ROIs made by two independent raters for each participant according to our previously published methods (Turker et al., 2021). Using the individual LC ROIs, we created a probabilistic map of the location of the locus coeruleus (LC) in MNI space. The probabilistic map spanned from coronal coordinates y = -39 to -32, axial coordinates z = -32 to -21, and sagittal coordinates x = -7 to -2 and x = 3 to 8 (MNI coordinates, **Figure 1**).

LC volume: The average total LC volume (both left and right) across all participants was 37.3 mm3, with a SD of 22.2 mm3 and a range from 6.1 to 133.1 mm3 (**Figure. 2A and 2B**). A linear model (using lme4 in R, REF) tested the effect of age group on total volume. LC volume did not differ significantly between age groups ($F(4) = 0.21$, $p = 0.92$).

LC neuromelanin intensity: Neuromelanin intensity was expressed in terms of multiples of the maximum intensity of non-LC voxels in the same brainstem slices where LC was identified. The average LC neuromelanin intensity across all participants was 2.16, or 2.16 times higher than the calculated maximum of non-LC voxels in the image, with a SD of 0.16 and a range from 1.81 to 2.46 **(Fig. 2C and 2D)**. Locus coeruleus neuromelanin intensity increased from young adulthood, was highest around age 60, and declined thereafter **(Fig. 2C)**. A linear model was constructed to test the effect of age group on average neuromelanin intensity; intensity varied significantly with age group ($F(4) = 4.9$, $p = 0.0011$). Age groups 30-50, 50-60, and 60-70 all differed significantly from the reference age group 18-30 (all $t > 2.3$, all $p < 0.03$), while age group 70+ did not ($t = 0.55$, $p = 0.57$). Since neuromelanin intensity across the

lifespan appeared to follow an inverted-U shaped curve, we also fitted a generalized additive model (Sørensen et al., 2021) to test the relationship between age (by year, rather than by group) and neuromelanin intensity, and found a significant relationship (for the age term, edf = 2.49, F = 4.46, p = 0.0053, R squared = 0.13).

To ensure that there was no systematic bias in the measurement of LC neuromelanin intensity across age groups, we tested whether minimum neuromelanin intensity (the lowest-intensity voxel included within each individual's hand-drawn LC ROI) varied with age. Overall, it did not (F(4) = 1.92, p = 0.12), although the 60-70 age group did have marginally greater minimum voxel values compared with the youngest age group (t = 1.99, p = 0.05). The 70+ age group, in which we saw a marked decrease in average neuromelanin intensity, did not have significantly decreased minimum neuromelanin intensity. The distribution of minimum voxel values was similar in all age groups and did not match the distribution of maximum voxel values **(Fig. 2E)**. Furthermore, our Dice similarity coefficients for the creation if individual LC ROIs did not differ between younger (< 60) and older (>60) participants (see Methods).These reassuring analyses suggested that our measurements of volume were not affected by systematically including less-intense voxels in some groups vs. others.

Differences between left and right LC: In all age groups, the average volume of the left LC was observed to be larger than the volume of the right LC. The average volume of the left LC was 19.8 mm3, while the average volume of the right LC was 16.6 mm3. A linear model was constructed to test whether side affected volume in any age groups; the effect of side on volume was marginally significant (F(1) = 3.15, p = 0.07, Fig. 3A and 3B), but there was no interaction between age group and side on volume (F(4) = 0.26, p = 0.89). A similar model was constructed to test whether the left and right LC intensity differed significantly in any age group; they did not (F(1) = 0.14, p = 0.69, Figs. 3C and 3D).

Rostral vs. caudal LC: Because the intensity of the rostral LC has often been associated with cognition more than the caudal LC (Chen et al., 2023; Dahl et al., 2019), we repeated the analysis of neuromelanin intensity across age groups using rostral LC intensity only, which also varied significantly with age (F(4) = 4.55, p = 0.0021, with age groups 30-50, 50-60, and 60-70 differing significantly from age group 18-30, all p < 0.04, as in the previous model). In all age groups but one, the average intensity of the caudal LC was observed to be greater than the intensity of the rostral LC. A linear model was constructed to determine whether intensity depended on LC region (rostral vs caudal) in any age group; caudal intensity was found to be significantly higher (F(1) = 4.92, p = 0.027). Average caudal intensity was 2.16 with SD = 0.18, while average rostral intensity was 2.12 with SD = 0.16 (Figs. 3E and 3F).

LC neuromelanin intensity and cognitive performance: We measured fluid cognition using the NIH Toolbox (Fluid Cognition score) and crystallized cognition using the National Adult Reading Test (NART). Across our sample, raw fluid cognition scores declined sharply across the lifespan. A linear model with raw (non-age normalized) fluid cognition standard score as the dependent variable and age as the independent variable demonstrated that cognitive ability depended on age (F(1) = 56.3, p < 0.000001, Pearson correlation coefficient = -0.67). A corresponding model tested whether crystallized cognition (raw NART score) depended on age;

it did not (F(1) = 2.43, p = 0.12, Pearson correlation coefficient = 0.16). Thus, as expected (Moran, 2013), fluid cognition declined with age but crystallized cognition did not. Given that we previously found that neuromelanin intensity increased with age (until the clear downturn in our oldest age group, aged 70+), we expected to find a significant negative correlation between rostral neuromelanin intensity and raw fluid cognition scores in those under 70, and this was indeed the case (r = -0.35, one sample t-test t(74) = -3.2, p = 0.0019).

In order to control for the very strong effect of age on fluid cognition, we next asked whether age-normalized fluid cognition scores (national percentile for age) were associated with neuromelanin intensity. Previous work has found an association between neuromelanin intensity and measures of fluid (REF) and crystallized cognition (REF) only in older adults. Since our data suggested a peak in LC intensity around age 60, we considered that age the lowest for inclusion in this analysis. First we ensured that our normalized cognitive measure did not depend on age by constructing a linear model with age-normalized fluid cognition score as the dependent variable and age as the independent variable, and we found no association (F(1) = 0.27, p = 0.6). We then constructed a linear model to test whether age-normalized fluid cognition ability depended on rostral LC intensity. We found no significant effect (Pearson correlation coefficient = 0.05 ,F(1) = 0.13, p = 0.73, Fig. 4A). However, upon closer examination , we noticed a number of very low-performing individuals, and that low- and high-performing older adults seemed to have a different relationship between neuromelanin intensity and cognition.

We repeated the analysis for adults over 60 within each of two cognitive groups (below 50th percentile for age (n=11), and above 50th percentile for age (n=32)). Among adults performing above average for their age, we found a significant positive association between rostral LC intensity and global cognitive performance (Pearson correlation coefficient = 0.39, F(1) = 5.48, p = 0.025, Fig. 4B). However, among adults performing below average for their age, there was no significant relationship (Pearson correlation coefficient = -0.04, F(1) = 0.3, p = 0.57, Fig. 4C). This is despite the fact that high-performing and low-performing groups did not differ in average rostral LC intensity (2.16 ± 0.16 vs 2.16 ± 0.21, respectively).

We used a similar linear model to test whether crystallized cognition (NART score; this score was not corrected for age because it did not depend significantly on age) depended on rostral LC intensity in adults over 60. We found no significant effect in high- or low-performing participants. There was no significant relationship between rostral LC intensity and cognitive performance in adults under 60, nor was there any relationship between caudal LC intensity and cognitive performance in any group. We also found no significant relationship between any cognitive subtests and cognitive performance.

Neuromelanin intensity differences by race and sex: In all age groups (reduced to 3 because of uneven gender balance within decade age groups), women had higher LC intensity than men. We constructed a linear model to determine whether these differences were significant and whether there was an interaction with age group. There was a significant effect of sex (F(1) = 16.04, p = 0.00012, Fig. 5A) and no significant interaction between sex and age group (F(2) = 0.43, p = 0.65).

We constructed a linear model to test whether there were differences in rostral LC intensity by race, and whether there was any interaction between race and age group. We did not find any significant differences in rostral (or average) LC intensity between participants of

black (n= 16) and white (n = 55) race (F(1) = 0.4, p = 0.52, Fig. 5B) in any age group, and there was no significant interaction between race and age group (F(2) = 1.4, p = 0.24). We were unable to test the effect of Asian or Latinx heritage by age group as there were not enough participants in each age category.

Discussion:

Employing neuromelanin-sensitive MRI, we found that neuromelanin intensity, defined relative to the surrounding pons, increased from early adulthood through approximately age 60, and declined thereafter. The caudal portion of the LC had greater neuromelanin intensity than the rostral portion. Demographically, women had greater neuromelanin intensity in the LC than men but we did not find any reliable differences by race. While the caudal portion of the LC had greater intensity, it was the rostral portion of the LC that was associated with cognitive performance. This relation applied only to older adults with above-average cognition for their age and was selective to fluid rather than crystalized measures of cognition.

Previous analyses of neuromelanin across the lifespan have also detected a peak in neuromelanin intensity around age 50-60, including early post mortem studies and two in-vivo MRI studies (Liu et al., 2019; Manaye et al., 1995; Mann et al., 1985; Shibata et al., 2006). However, other studies have found no change in neuromelanin concentration across the lifespan (Hämmerer et al., 2018; Ohm et al., 1997; Zucca et al., 2006), or simply an increase between younger and older adults (Clewett 2016 in vivo, Betts 2017 in vivo). Thus, even recent examinations of LC neuromelanin intensity have not yet arrived at a completely consistent result. In studies on the substantia nigra, the other neuromelanin-containing brain structure, the inverted-U with a peak around X has been found using neuromelanin-sensitive MRI (Xing et al., 2018). Differences by sex are even less clear, with some analyses (Clewett et al., 2015) showing markedly lower intensity in women, and some showing no difference (Betts et al., 2017; Shibata et al., 2006). By contrast, our analysis shows *increased* signal intensity in women. Though our sample is relatively small (N = 96), it is larger than the sample size in all studies cited in this paragraph except for (Liu et al., 2019), with which it most closely agrees. That our life course of neuromelanin intensity was internally replicated between left and right LC, between rostral and caudal regions, within racial groups and for both males and females, gives some further confidence in its accuracy. As far as we are aware, no other published studies have examined neuromelanin intensity across different races.

We found a positive association between rostral LC intensity and fluid, but not crystallized, cognition specific to older adults with above-average cognition. Related results have previously been reported by others (Bell et al., 2023; Clewett et al., 2015; Dahl et al., 2019, 2023), though not with perfect alignment. Clewett et al., 2015 found a significant association between crystallized intelligence and overall LC intensity, Bell et al., 2023 found an significant association between subjective cognition and rostral LC intensity, and Dahl et al., 2019 found a significant association between memory performance and rostral LC intensity, and then in 2023 found that LC intensity significantly predicted future episodic memory performance. This growing literature will benefit from continued study with large and diverse sample sizes. Our study had uneven group sizes, due to the relative difficulty of recruiting middle-aged participants, and while we had significant diversity within our sample, we were unable to ensure that every age, sex, and racial group was matched in all ways that might theoretically impact

neuromelanin intensity (e.g. education). Furthermore, even though we detected a significant positive relationship between neuromelanin and fluid cognition, it was not strong or specific enough such that neuromelanin measurement would tell a clear story about an individual.

The cause of loss of neuromelanin in older adults is not clear. It could be either death of catecholaminergic cells, or neuromelanin loss without cell death. It is difficult to distinguish between these two possibilities. There is some evidence for reduced neuromelanin content in surviving cells in older adults (Faucheux et al., 2003; Mann et al., 1985), although neuromelanin loss without cell death is considered less likely because previous work has linked neuromelanin directly to surviving LC neurons (Keren et al., 2009, 2015; Kitao et al., 2013). (This work comes from neuromelanin-sensitive MRI in patients with Parkinson's disease, a literature which has produced more than a dozen replications of reduced neuromelanin signal, see (Sulzer et al., 2018) Although we measured LC volume, diffuse neuronal loss across the LC could result in an unchanged volume even after substantial cell death. Furthermore, although neuromelanin clearly accumulates in granules within cells, it also exists in the extracellular space, for an undetermined amount of time prior to clearance, likely by microglia (Zhang et al., 2011; Zucca et al., 2006). Recent quantitative susceptibility mapping has made clear that iron resulting from neuromelanin granule loss is taken up by surrounding cells, where it is subsequently visible on MRI using iron-sensitive sequences that are *not* sensitive to neuromelanin (He et al., 2023; Takahashi et al., 2018).

The question of how neuromelanin impacts neuron health, and what to make of its decrease in old age, is being actively investigated and has been labeled a paradox (Moreno-García et al., 2021), an enigma (Youdim et al., 1994) and a riddle (Iannitelli & Weinshenker, 2023). Since neuromelanin unquestionably results from the oxidation of intermediate products in the norepinephrine production pathway, the amount produced might be considered a reporter of total NE and DA production in the locus coeruleus (Clewett et al., 2015), although very recent work suggests the possibility of active neuromelanin clearance (Bachman et al., 2023). Because NE exerts numerous neuroprotective effects on the brain and is essential for cognition (Feinstein et al., 2016; Robertson, 2013), and since neuromelanin is a highly effective quencher of toxins that might otherwise damage the LC (Moreno-García et al., 2021), greater production or preservation of this neurochemical could account for the positive relationship between neuromelanin intensity and cognition, which we also observed. On the other hand, it is very clear from recent animal studies that overexpression of human tyrosinase in both the substantia nigra and LC in animal models results in abundant neuromelanin production and very quickly induces neurodegeneration and associated phenotypes as a direct consequence of neuromelanin accumulation (Carballo-Carbajal et al., 2019; Iannitelli et al., 2023).

The implication of these studies is that once neuromelanin accumulates past a tipping point, its beneficial activities cease and it results in a neuroinflammatory response that damages or kills cells. From this point of view, the association between cognition and neuromelanin intensity could simply reflect the fact that the tipping point has not yet been reached. The LC norepinephrine system is essential for maintaining optimal cognitive function with age. However, in our data, we found that older adults with below-average cognition did not have reduced neuromelanin, but lacked any relationship between neuromelanin intensity and cognition. This highlights that lower cognitive function has many sources beyond the LC, but maintaining higher cognitive function, especially related to fluid cognition, may depend on integrity of the LC

norepinephrine system (Hayes & Petrov, 2016; Tsukahara et al., 2016; Tsukahara & Engle, 2021), though longitudinal data are needed to address this. This is an active topic of research, especially since the LC is the first structure to show tau pathology in the earliest stages of Alzheimer's disease (Braak stage 0, (Braak et al., 2011), and could serve as either an early warning sign, or perhaps as a literal spreader of disease. However, the implications for LC damage go beyond AD, since substantial LC damage is also found in Parkinson's disease (Zarow et al., 2003), and also other diseases such as multiple sclerosis (Polak et al., 2011), and pathological anxiety (Morris et al., 2020). What role neuromelanin, especially in its capacity to sequester environmental toxins, plays in all of these diseases is yet to be discovered.

Very recent work in the field is addressing crucial questions: whether individual neurons with the most neuromelanin are at the highest risk for imminent death (Nagatsu et al., 2023), whether neuromelanin content of an individual's LC relates to its catecholamine production capacity and/or connectivity (Chen et al., 2023), and the true source of the neuromelanin MRI signal and how it can be processed more effectively to get a more precise look in vivo at the state of the LC norepinephrine system in humans (Trujillo et al., 2023). These studies should help to further clarify the meaning of neuromelanin as a functional biomarker and its ability to affect neural activity and health in the LC. Furthermore, other types of neuromelanin-sensitive MRI sequences, including magnetization transfer and quantitative susceptibility mapping sequences, are being tested and have the potential to bring converging (or diverging) evidence about the importance of neuromelanin. All of these advances should bring us closer to being able to understand when neuromelanin is beneficial vs. harmful and how the associated benefits or risks might be most productively managed.

Methods:

Participants: We examined 96 participants between the ages of 19 and 86. For examination of neuromelanin across the lifespan, participants were broken up into 6 age groups: 18-29 (n = 32), 30-49 (n = 10), 50-59 (n = 9), 60-69 (n = 30), and 70+ (n = 15). Due to smaller sample size in middle-aged groups, for analyses by race and sex, participants were broken into 3 standard US Census groups instead: younger (19-44 years of age, n = 37, 27 female, 13 white, 5 Black), middle aged (45-64 years of age, n = 20, 12 female, 11 White, 6 Black), and older (65+ years of age, n = 37, 21 female, 31 White, 6 Black). Participants were screened for diagnosed cognitive impairment, neurological disease, head injury, ocular disease, and had vision and hearing that were normal or correctible to normal. All were fluent speakers of English. The sample included 55 white participants (57%), 16 Black participants (17%), 12 Asian participants (13%), 8 Latinx participants (8%), and 4 participants of other or multiple races or ethnicities (4%). The sample included 35 men (36%) and 61 women (64%). This research protocol was approved by the Cornell University Institutional Review Board for Human Participants with protocol number 1910009087.

Cognitive assessment: Participants were assessed with the NIH Toolbox Cognition Battery (Hodes et al., 2013). Subsets used include Picture Sequence Memory Test, Flanker Inhibitory Control and Attention Test, Dimension Change Card Sort Test, List Sorting Working

Memory Test, and Pattern Comparison Processing Speed Test. Uncorrected Standard Score was used to compare raw performance across age groups, and National Percentile (age-adjusted) was used to measure cognition relative to expectation. Participants were also assessed with the National Adult Reading Test (Venegas & Clark, 2011).

MRI: Volunteers were scanned at the Cornell Magnetic Resonance Imaging Facility on a General Electric 3.0 Tesla MR750 scanner (Waukesha, WI, software version DV24 and DV29.1) with a 32 channel receive-only head coil. Magnitude images were saved to disk. Anatomical data were acquired with a T1-weighted MPRAGE sequence (TR = 7.7 ms; TE = 3.42 ms; 7° flip angle; 1.0 mm isotropic voxels, 176 slices). For the localization of the LC we used a protocol that provided the best LC contrast in a reasonably short scan time. It was inspired by previous publications with LC MRI protocols (Jacobs et al., 2021; Keren et al., 2009; Liu et al., 2020; Watanabe et al., 2019) and was optimized to fit our specific 3.0 Tesla MRI scanner model and its available pulse sequences. A 2D T1-weighted fast spin-echo sequence with ETL = 3 was used, with 18 slices of 3 mm thickness, field-of-view = 22 × 22 cm, acquisition and output matrix size = 512 × 512 (resolution = 0.43 × 0.43 mm), 60% FOV, TR = 701 ms, TE = 11.44 ms, refocusing flip angle = 120 degrees, 6 averages, bandwidth = 15.63 kHz, and extended dynamic range. Slices for this volume were oriented perpendicular to the brain stem to provide high resolution data in the axial plane, where the dimensions of the LC are smallest, and positioned to cover the most rostral portion of the pons. Participants laid supine on the scanner bed for all scans with their head supported and immobilized. Ear plugs, headphones, and a microphone were used to reduce scanner noise and to allow the participant to communicate with the experimenters.

Analysis of images: In order to ensure that any voxels contained within the 4th ventricle were not used in neuromelanin analysis, MPRAGE images were processed using FreeSurfer *recon-all* (Fischl, 2012) in order to produce an automated ROI corresponding to the 4th ventricle. The MPRAGE image, and corresponding parcellation, was then aligned with the TSE image with AFNI (Cox, 1996) using the *3dAllineate* function. After definition of the LC ROI, any voxels overlapping with the 4th ventricle were removed. Measurement of neuromelanin intensity for each participant was then conducted as described below using the TSE image.

Probabilistic map: A probabilistic map illustrating the location and distribution of the LC ROIs in all participants was created by registering each person's LC ROI to MNI152 space. To do so, alignment parameters were calculated for each individual by combining the inverse of the transformation that aligned the native MPRAGE and TSE images and the nonlinear transformation from the individual's native MPRAGE to MNI152 standard calculated by AFNI functions *3dQwarp* and *3dNwarpApply*. Successful alignment of the brainstem to the individual MPRAGE and the MNI152 template was visually confirmed. LC ROIs in MNI152 space were thresholded to remove scattered non-zero voxels introduced by the nonlinear warping process. A heatmap was then created using AFNI function *3dMerge*.

Measurement of NM intensity: NM intensity was measured by exactly following the procedure described in our previous published work (Turker et al., 2021). Briefly, raw data was processed for correction of artifacts and optimized for contrast. Then, LC ROIs were

hand-drawn using *FreeView* software according to clear guidelines (Turker et al., 2021) by two independent raters. Only the overlap of both ROIs was used, less any voxels belonging to the 4th ventricle mask, after ensuring that the Dice similarity coefficient between raters was greater than 0.6. If the coefficient was less than 0.6, both raters reviewed the ROI drawing procedure and independently redid the ROI. This was necessary for only 4 out of 96 participants. The average Dice coefficient was 0.84 for all participants, 0.839 for participants under age 60 and 0.841 for participants over 60, demonstrating good agreement of the ROI drawing procedure across age groups.

After creation of the ROIs, neuromelanin images were then rescaled to a range of 6 standard deviations. This transformation was chosen to be as similar as possible to min-max scaling (min 0, max 1), without undue influence of extreme outliers caused by scanner noise. The scaling range was calculated using all voxels in the brainstem (as opposed to a specific reference region of the pontine tegmentum) in slices where the LC was defined, except for those within the LC ROI. The transformation was then applied to all voxels, including those within the LC, such that the neuromelanin intensity could then be interpreted relative to the range of non-LC values within all slices in which LC was observed. For example, a value of 2 indicates that the LC was twice as intense as the maximum (actually 3 SD above mean) of the non-LC values. Neuromelanin intensity was defined as the sum of all voxel values within the LC mask divided by the number of voxels. Rostral vs. caudal regions were defined by a 50/50 split along the rostral-caudal axis of the overall length of the individually defined LC ROI for each person. There was one participant with an outlier value, an average neuromelanin intensity 5.8 SD above the total sample average. This participant was removed from analysis, but inclusion or exclusion of this individual did not change the result of any statistical tests.

Statistics: Linear models were run in R using the *lme4* package (Bates et al., 2015). Summary statistics such as p values and F statistics, were calculated using the *anova* function, while p values and t statistics for levels within a predictor (e.g., to determine which age groups differ from the reference group in a model in which age group was a significant predictor) were calculated using the *summary* function. One general additive model was produced using the *mgcv* package (Sørensen et al., 2021).

Funding: This work was supported by the National Institutes of Health (National Institute on Aging grants F32 AG058479 to ER and R01AG066430 to EDR and AKA). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgements: The authors wish to thank Henning Voss of the Cornell MRI Facility for consultation, optimization and troubleshooting in the use of neuromelanin-sensitive MRI sequences.

Data Availability Statement: For privacy reasons, neither full anatomical MRI scans nor full neuromelanin-sensitive MRI scans can be made available. However, neuromelanin-sensitive MRI scans limited to the brainstem, in addition to all cognitive and demographic data necessary to replicate the analyses described here, are available at the Open Science Framework at https://osf.io/tf9br

Works Cited:

Aston-Jones, G., & Cohen, J. D. (2005). AN INTEGRATIVE THEORY OF LOCUS

   COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal

   Performance. *Annual Review of Neuroscience*, *28*(1), 403–450.

   https://doi.org/10.1146/annurev.neuro.28.061604.135709

Aston-Jones, G., Rajkowski, J., & Cohen, J. (2000). Locus coeruleus and regulation of

   behavioral flexibility and attention. *Progress in Brain Research*, *126*, 165–182.

   https://doi.org/10.1016/S0079-6123(00)26013-5

Babulal, G. M., Quiroz, Y. T., Albensi, B. C., Arenaza-Urquijo, E., Astell, A. J., Babiloni, C.,

   Bahar-Fuchs, A., Bell, J., Bowman, G. L., Brickman, A. M., Chételat, G., Ciro, C., Cohen,

   A. D., Dilworth-Anderson, P., Dodge, H. H., Dreux, S., Edland, S., Esbensen, A., Evered,

   L., … O'Bryant, S. E. (2019). Perspectives on ethnic and racial disparities in Alzheimer's

   disease and related dementias: Update and areas of immediate need. *Alzheimer's &*

   Dementia, *15*(2), 292–312. https://doi.org/10.1016/j.jalz.2018.09.009

Bachman, S. L., Cole, S., Yoo, H. J., Nashiro, K., Min, J., Mercer, N., Nasseri, P., Thayer, J. F.,

   Lehrer, P., & Mather, M. (2023). Daily heart rate variability biofeedback training

   decreases locus coeruleus MRI contrast in younger adults in a randomized clinical trial.

   International Journal of Psychophysiology, *193*, 112241.

   https://doi.org/10.1016/j.ijpsycho.2023.08.014

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models

   using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

   https://doi.org/10.18637/jss.v067.i01

Beardmore, R., Hou, R., Darekar, A., Holmes, C., & Boche, D. (2021). The Locus Coeruleus in

Aging and Alzheimer's Disease: A Postmortem and Brain Imaging Review. *Journal of*

Alzheimer's Disease, *83*(1), 5–22. https://doi.org/10.3233/JAD-210191

Bell, T. R., Elman, J. A., Beck, A., Fennema-Notestine, C., Gustavson, D. E., Hagler, D. J.,

Jack, A. J., Lyons, M. J., Puckett, O. K., Toomey, R., Franz, C. E., & Kremen, W. S.

(2023). Rostral-middle locus coeruleus integrity and subjective cognitive decline in early

old age. *Journal of the International Neuropsychological Society*, *29*(8), 763–774.

https://doi.org/10.1017/S1355617722000881

Berridge, C. W., & Waterhouse, B. D. (2003). The locus coeruleus-noradrenergic system:

Modulation of behavioral state and state-dependent cognitive processes. *Brain*

Research Reviews, *42*(1), 33–84. https://doi.org/10.1016/S0165-0173(03)00143-7

Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Jessen, F., & Düzel, E. (2017). In vivo MRI

assessment of the human locus coeruleus along its rostrocaudal extent in young and

older adults. *NeuroImage*, *163*, 150–159.

https://doi.org/10.1016/j.neuroimage.2017.09.042

Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the pathologic

process in alzheimer disease: Age categories from 1 to 100 years. *Journal of*

Neuropathology and Experimental Neurology, *70*(11), 960–969.

https://doi.org/10.1097/NEN.0b013e318232a379

Carballo-Carbajal, I., Laguna, A., Romero-Giménez, J., Cuadros, T., Bové, J., Martinez-Vicente,

M., Parent, A., Gonzalez-Sepulveda, M., Peñuelas, N., Torra, A., Rodríguez-Galván, B.,

Ballabio, A., Hasegawa, T., Bortolozzi, A., Gelpi, E., & Vila, M. (2019). Brain tyrosinase

overexpression implicates age-dependent neuromelanin production in Parkinson's

disease pathogenesis. *Nature Communications*, *10*(1), 973–973.

https://doi.org/10.1038/s41467-019-08858-y

Cassidy, C. M., Therriault, J., Pascoal, T. A., Cheung, V., Savard, M., Tuominen, L., Chamoun, M., McCall, A., Celebi, S., Lussier, F., Massarweh, G., Soucy, J. P., Weinshenker, D., Tardif, C., Ismail, Z., Gauthier, S., & Rosa-Neto, P. (2022). Association of locus coeruleus integrity with Braak stage and neuropsychiatric symptom severity in Alzheimer's disease. *Neuropsychopharmacology*, *47*(5), 1128–1136. https://doi.org/10.1038/s41386-022-01293-6

Celebi, S. N. (2022). *REM Sleep is Associated with Neuromelanin-Sensitive MRI Signal in the Locus Coeruleus in Veterans with a History of Post-Traumatic Stress Disorder* [Thesis, Université d'Ottawa / University of Ottawa]. https://doi.org/10.20381/ruor-28527

Chalermpalanupap, T., Weinshenker, D., & Rorabaugh, J. M. (2017). Down but Not Out: The Consequences of Pretangle Tau in the Locus Coeruleus. *Neural Plasticity*, *2017*. https://doi.org/10.1155/2017/7829507

Chan-Palay, V., & Asan, E. (1989). Alterations in catecholamine neurons of the locus coeruleus in senile dementia of the Alzheimer type and in Parkinson's disease with and without dementia and depression. *Journal of Comparative Neurology*, *287*(3), 373–392. https://doi.org/10.1002/cne.902870308

Chen, H.-Y., Parent, J. H., Ciampa, C. J., Dahl, M. J., Hämmerer, D., Maass, A., Winer, J. R., Yakupov, R., Inglis, B., Betts, M. J., & Berry, A. S. (2023). Interactive effects of locus coeruleus structure and catecholamine synthesis capacity on cognitive function. *Frontiers in Aging Neuroscience*, *15*, 1236335. https://doi.org/10.3389/fnagi.2023.1236335

Clewett, D. V., Lee, T. H., Greening, S., Ponzio, A., Margalit, E., & Mather, M. (2015). Neuromelanin marks the spot: Identifying a locus coeruleus biomarker of cognitive reserve in healthy aging. *Neurobiology of Aging*, *37*, 117–126. https://doi.org/10.1016/j.neurobiolaging.2015.09.019

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic
    resonance neuroimages. *Computers and Biomedical Research, an International Journal*,
    29(3), 162–173. https://doi.org/10.1006/cbmr.1996.0014

Dahl, M. J., Bachman, S. L., Dutt, S., Düzel, S., Bodammer, N. C., Lindenberger, U., Kühn, S.,
    Werkle-Bergner, M., & Mather, M. (2023). The integrity of dopaminergic and
    noradrenergic brain regions is associated with different aspects of late-life memory
    performance. *Nature Aging*, *3*(9), Article 9. https://doi.org/10.1038/s43587-023-00469-z

Dahl, M. J., Mather, M., Düzel, S., Bodammer, N. C., Lindenberger, U., Kühn, S., & Werkle-
    Bergner, M. (2019). Rostral locus coeruleus integrity is associated with better memory
    performance in older adults. *Nature Human Behaviour*, 1–12.
    https://doi.org/10.1038/s41562-019-0715-2

Faucheux, B. A., Martin, M.-E., Beaumont, C., Hauw, J.-J., Agid, Y., & Hirsch, E. C. (2003).
    Neuromelanin associated redox-active iron is increased in the substantia nigra of
    patients with Parkinson's disease. *Journal of Neurochemistry*, *86*(5), 1142–1148.
    https://doi.org/10.1046/j.1471-4159.2003.01923.x

Feinstein, D. L., Kalinin, S., & Braun, D. (2016). Causes, consequences, and cures for
    neuroinflammation mediated via the locus coeruleus: Noradrenergic signaling system.
    Journal of Neurochemistry, *139*, 154–178. https://doi.org/10.1111/jnc.13447

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.
    https://doi.org/10.1016/j.neuroimage.2012.01.021

Gesi, M., Soldani, P., Giorgi, F. S., Santinami, A., Bonaccorsi, I., & Fornai, F. (2000). The role of
    the locus coeruleus in the development of Parkinson's disease. *Neuroscience and
    Biobehavioral Reviews*, *24*(6), 655–668. https://doi.org/10.1016/S0149-7634(00)00028-2

Ghosh, A., Torraville, S. E., Mukherjee, B., Walling, S. G., Martin, G. M., Harley, C. W., & Yuan,
    Q. (2019). An experimental model of Braak's pretangle proposal for the origin of

Alzheimer's disease: The role of locus coeruleus in early symptom development. *Alzheimer's Research & Therapy*, *11*(1), 59. https://doi.org/10.1186/s13195-019-0511-2

Goldstein, D. S., Sullivan, P., Holmes, C., Miller, G. W., Alter, S., Strong, R., Mash, D. C., Kopin, I. J., & Sharabi, Y. (2013). Determinants of buildup of the toxic dopamine metabolite DOPAL in Parkinson's disease. *Journal of Neurochemistry*, *126*(5), 591–603. https://doi.org/10.1111/jnc.12345

Granholm, E. L., Panizzon, M. S., Elman, J. A., Jak, A. J., Hauger, R. L., Bondi, M. W., Lyons, M. J., Franz, C. E., & Kremen, W. S. (2017). Pupillary Responses as a Biomarker of Early Risk for Alzheimer's Disease. *Journal of Alzheimer's Disease*, *56*(4), 1419–1428. https://doi.org/10.3233/JAD-161078

Guinea-Izquierdo, A., Giménez, M., Martínez-Zalacaín, I., Del Cerro, I., Canal-Noguer, P., Blasco, G., Gascón, J., Reñé, R., Rico, I., Camins, A., Aguilera, C., Urretavizcaya, M., Ferrer, I., Menchón, J. M., Soria, V., & Soriano-Mas, C. (2021). Lower Locus Coeruleus MRI intensity in patients with late-life major depression. *PeerJ*, *9*, e10828. https://doi.org/10.7717/peerj.10828

Haining, R. L., & Achat-Mendes, C. (2017). Neuromelanin, one of the most overlooked molecules in modern medicine, is not a spectator. *Neural Regeneration Research*, *12*(3), 372–375. https://doi.org/10.4103/1673-5374.202928

Hämmerer, D., Callaghan, M. F., Hopkins, A., Kosciessa, J., Betts, M., Cardenas-Blanco, A., Kanowski, M., Weiskopf, N., Dayan, P., Dolan, R. J., & Düzel, E. (2018). Locus coeruleus integrity in old age is selectively related to memories linked with salient negative events. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(9), 2228–2233. https://doi.org/10.1073/pnas.1712268115

Hayes, T. R., & Petrov, A. A. (2016). Pupil Diameter Tracks the Exploration–Exploitation Trade-off during Analogical Reasoning and Explains Individual Differences in Fluid Intelligence.

Journal of Cognitive Neuroscience, *28*(2), 308–318.

https://doi.org/10.1162/jocn_a_00895

He, N., Chen, Y., LeWitt, P. A., Yan, F., & Haacke, E. M. (2023). Application of Neuromelanin

MR Imaging in Parkinson Disease. *Journal of Magnetic Resonance Imaging*, *57*(2), 337–

352. https://doi.org/10.1002/jmri.28414

Hodes, R. J., Insel, T. R., & Landis, S. C. (2013). The NIH Toolbox. *Neurology*, *80*(11 Suppl 3),

S1. https://doi.org/10.1212/WNL.0b013e3182872e90

Iannitelli, A. F., Segal, A., Pare, J.-F., Mulvey, B., Liles, L. C., Sloan, S. A., McCann, K. E.,

Dougherty, J. D., Smith, Y., & Weinshenker, D. (2023). *Tyrosinase-induced*

neuromelanin accumulation triggers rapid dysregulation and degeneration of the mouse

locus coeruleus (p. 2023.03.07.530845). bioRxiv.

https://doi.org/10.1101/2023.03.07.530845

Iannitelli, A. F., & Weinshenker, D. (2023). Riddles in the dark: Decoding the relationship

between neuromelanin and neurodegeneration in locus coeruleus neurons.

Neuroscience & Biobehavioral Reviews, *152*, 105287.

https://doi.org/10.1016/j.neubiorev.2023.105287

Jacobs, H. I. L., Becker, A., Sperling, R. A., Guzman-Velez, E., Baena, A., Uquillas, F. d'Oleire,

Artola, A., Pardilla-Delgado, E., Gatchel, J. R., Reiman, E. M., Johnson, K. A., Lopera,

F., & Quiroz, Y. T. (2019). LOCUS COERULEUS INTENSITY IS ASSOCIATED WITH

EARLY AMYLOID AND TAU PATHOLOGY IN PRECLINICAL AUTOSOMAL

DOMINANT ALZHEIMER'S DISEASE. *Alzheimer's & Dementia*, *15*(7), P774–P775.

https://doi.org/10.1016/j.jalz.2019.06.2832

Jacobs, H. I. L., Becker, J. A., Kwong, K., Engels-Domínguez, N., Prokopiou, P. C., Papp, K. V.,

Properzi, M., Hampton, O. L., d'Oleire Uquillas, F., Sanchez, J. S., Rentz, D. M., El

Fakhri, G., Normandin, M. D., Price, J. C., Bennett, D. A., Sperling, R. A., & Johnson, K.

A. (2021). In vivo and neuropathology data support locus coeruleus integrity as indicator

of Alzheimer's disease pathology and cognitive decline. *Science Translational Medicine*, *13*(612), eabj2511. https://doi.org/10.1126/scitranslmed.abj2511

Kang, S. S., Liu, X., Ahn, E. H., Xiang, J., Manfredsson, F. P., Yang, X., Luo, H. R., Liles, L. C., Weinshenker, D., & Ye, K. (2020). Norepinephrine metabolite DOPEGAL activates AEP and pathological Tau aggregation in locus coeruleus. *The Journal of Clinical Investigation*, *130*(1), 422–437. https://doi.org/10.1172/JCI130513

Keren, N. I., Lozar, C. T., Harris, K. C., Morgan, P. S., & Eckert, M. A. (2009). In vivo mapping of the human locus coeruleus. *NeuroImage*, *47*(4), 1261–1267. https://doi.org/10.1016/J.NEUROIMAGE.2009.06.012

Keren, N. I., Taheri, S., Vazey, E. M., Morgan, P. S., Granholm, A. C. E., Aston-Jones, G. S., & Eckert, M. A. (2015). Histologic validation of locus coeruleus MRI contrast in post-mortem tissue. *NeuroImage*, *113*, 235–245. https://doi.org/10.1016/J.NEUROIMAGE.2015.03.020

Kitao, S., Matsusue, E., Fujii, S., Miyoshi, F., Kaminou, T., Kato, S., Ito, H., & Ogawa, T. (2013). Correlation between pathology and neuromelanin MR imaging in Parkinson's disease and dementia with Lewy bodies. *Neuroradiology*, *55*(8), 947–953. https://doi.org/10.1007/s00234-013-1199-9

Klimek, V., Stockmeier, C., Overholser, J., Meltzer, H. Y., Kalka, S., Dilley, G., & Ordway, G. A. (1997). Reduced Levels of Norepinephrine Transporters in the Locus Coeruleus in Major Depression. *Journal of Neuroscience*, *17*(21), 8451–8458. https://doi.org/10.1523/JNEUROSCI.17-21-08451.1997

Kremen, W. S., Panizzon, M. S., Elman., J. A., Granholm, E. L., Andreassen, O. A., Dale, A. M., Gillespie, N. A., Gustavson, D. E., Logue, M. W., Lyons, M. J., Neale, M. C., Reynolds, C. A., Whitsel, N., & Franz, C. E. (2019). Pupillary dilation responses as a midlife indicator of risk for Alzheimer's Disease: Association with Alzheimer's disease polygenic risk. *Neurobiology of Aging*. https://doi.org/10.1016/j.neurobiolaging.2019.09.001

Langley, J., Huddleston, D. E., Liu, C. J., & Hu, X. (2016). Reproducibility of locus coeruleus

    and substantia nigra imaging with neuromelanin sensitive MRI. *Magnetic Resonance*

    Materials in Physics, Biology and Medicine, 1–5. https://doi.org/10.1007/s10334-016-

    0590-z

Liu, K. Y., Acosta-Cabronero, J., Cardenas-Blanco, A., Loane, C., Berry, A. J., Betts, M. J.,

    Kievit, R. A., Henson, R. N., Düzel, E., Howard, R., & Hämmerer, D. (2019). In vivo

    visualization of age-related differences in the locus coeruleus. *Neurobiology of Aging*,

    74, 101–111. https://doi.org/10.1016/J.NEUROBIOLAGING.2018.10.014

Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Tyler, L. K.,

    Brayne, C., Bullmore, E. T., Calder, A. C., Cusack, R., Dalgleish, T., Duncan, J.,

    Henson, R. N., Matthews, F. E., Marslen-Wilson, W. D., Shafto, M. A., Campbell, K.,

    Cheung, T., … Hämmerer, D. (2020). Noradrenergic-dependent functions are associated

    with age-related locus coeruleus signal intensity differences. *Nature Communications*,

    11(1), 1–9. https://doi.org/10.1038/s41467-020-15410-w

Manaye, K. F., McIntire, D. D., Mann, D. M., & German, D. C. (1995). Locus coeruleus cell loss

    in the aging human brain: A non-random process. *The Journal of Comparative*

    Neurology, *358*(1), 79–87. https://doi.org/10.1002/cne.903580105

Mann, D. M. A., Yates, P. O., & Marcyniuk, B. (1985). Correlation between senile plaque and

    neurofibrillary tangle counts in cerebral cortex and neuronal counts in cortex and

    subcortical structures in Alzheimer's disease. *Neuroscience Letters*, *56*(1), 51–55.

    https://doi.org/10.1016/0304-3940(85)90439-2

Moran, J. M. (2013). Lifespan development: The effects of typical aging on theory of mind.

    Behavioural Brain Research, *237*, 32–40. https://doi.org/10.1016/j.bbr.2012.09.020

Moreno-García, A., Kun, A., Calero, M., & Calero, O. (2021). The Neuromelanin Paradox and Its

    Dual Role in Oxidative Stress and Neurodegeneration. *Antioxidants*, *10*(1), Article 1.

    https://doi.org/10.3390/antiox10010124

Morris, L. S., McCall, J. G., Charney, D. S., & Murrough, J. W. (2020). The role of the locus coeruleus in the generation of pathological anxiety. *Brain and Neuroscience Advances*, *4*, 2398212820930321. https://doi.org/10.1177/2398212820930321

Nagatsu, T., Nakashima, A., Watanabe, H., Ito, S., Wakamatsu, K., Zucca, F. A., Zecca, L., Youdim, M., Wulf, M., Riederer, P., & Dijkstra, J. M. (2023). The role of tyrosine hydroxylase as a key player in neuromelanin synthesis and the association of neuromelanin with Parkinson's disease. *Journal of Neural Transmission*, *130*(5), 611–625. https://doi.org/10.1007/s00702-023-02617-6

Nelson, L. A., Yoash-Gantz, R. E., Pickett, T. C., & Campbell, T. A. (2009). Relationship Between Processing Speed and Executive Functioning Performance Among OEF/OIF Veterans. *Journal of Head Trauma Rehabilitation*, *24*(1), 32–40. https://doi.org/10.1097/HTR.0b013e3181957016

Ohm, T. G., Busch, C., & Bohl, J. (1997). Unbiased Estimation of Neuronal Numbers in the Human Nucleus Coeruleus during Aging. *Neurobiology of Aging*, *18*(4), 393–399. https://doi.org/10.1016/S0197-4580(97)00034-1

Oshima, S., Fushimi, Y., Okada, T., Nakajima, S., Yokota, Y., Shima, A., Grinstead, J., Ahn, S., Sawamoto, N., Takahashi, R., & Nakamoto, Y. (2021). Neuromelanin-Sensitive Magnetic Resonance Imaging Using DANTE Pulse. *Movement Disorders*, *36*(4), 874–882. https://doi.org/10.1002/mds.28417

Pamphlett, R., Bishop, D. P., Jew, S. K., & Doble, P. A. (2018). Age-related accumulation of toxic metals in the human locus ceruleus. *PLOS ONE*, *13*(9), e0203627. https://doi.org/10.1371/journal.pone.0203627

Pamphlett, R., Mak, R., Lee, J., Buckland, M. E., Harding, A. J., Jew, S. K., Paterson, D. J., Jones, M. W. M., & Lay, P. A. (2020). Concentrations of toxic metals and essential trace elements vary among individual neurons in the human locus ceruleus. *PLOS ONE*, *15*(5), e0233300. https://doi.org/10.1371/journal.pone.0233300

Plini, E. R. G., O'Hanlon, E., Boyle, R., Sibilia, F., Rikhye, G., Kenney, J., Whelan, R., Melnychuk, M. C., Robertson, I. H., & Dockree, P. M. (2021). Examining the Role of the Noradrenergic Locus Coeruleus for Predicting Attention and Brain Maintenance in Healthy Old Age and Disease: An MRI Structural Study for the Alzheimer's Disease Neuroimaging Initiative. *Cells*, *10*(7), Article 7. https://doi.org/10.3390/cells10071829

Polak, P. E., Kalinin, S., & Feinstein, D. L. (2011). Locus coeruleus damage and noradrenaline reductions in multiple sclerosis and experimental autoimmune encephalomyelitis. *Brain*, *134*(3), 665–677. https://doi.org/10.1093/brain/awq362

Priovoulos, N., van Boxel, S. C. J., Jacobs, H. I. L., Poser, B. A., Uludag, K., Verhey, F. R. J., & Ivanov, D. (2020). Unraveling the contributions to the neuromelanin-MRI contrast. *Brain Structure and Function*, *225*(9), 2757–2774. https://doi.org/10.1007/S00429-020-02153-Z/FIGURES/6

Robertson, I. H. (2013). A noradrenergic theory of cognitive reserve: Implications for Alzheimer's disease. *Neurobiology of Aging*, *34*(1), 298–308. https://doi.org/10.1016/j.neurobiolaging.2012.05.019

Ross, J. A., & Van Bockstaele, E. J. (2021). The Locus Coeruleus- Norepinephrine System in Stress and Arousal: Unraveling Historical, Current, and Future Perspectives. *Frontiers in Psychiatry*, *11*. https://www.frontiersin.org/articles/10.3389/fpsyt.2020.601519

Ryff, C. D., Heller, A. S., Schaefer, S. M., van Reekum, C., & Davidson, R. J. (2016). Purposeful Engagement, Healthy Aging, and the Brain. *Current Behavioral Neuroscience Reports*, *3*(4), 318–327. https://doi.org/10.1007/s40473-016-0096-z

Sara, S. J., & Bouret, S. (2012). Orienting and Reorienting: The Locus Coeruleus Mediates Cognition through Arousal. *Neuron*, *76*(1), 130–141. https://doi.org/10.1016/j.neuron.2012.09.011

Shibata, E., Sasaki, M., Tohyama, K., Kanbara, Y., Otsuka, K., Ehara, S., & Sakai, A. (2006). Age-related Changes in Locus Ceruleus on Neuromelanin Magnetic Resonance Imaging

at 3 Tesla. *Magnetic Resonance in Medical Sciences*, *5*(4), 197–200.
https://doi.org/10.2463/mrms.5.197

Sørensen, Ø., Walhovd, K. B., & Fjell, A. M. (2021). A recipe for accurate estimation of lifespan
brain trajectories, distinguishing longitudinal and cohort effects. *NeuroImage*, *226*,
117596. https://doi.org/10.1016/j.neuroimage.2020.117596

Sulzer, D., Cassidy, C., Horga, G., Kang, U. J., Fahn, S., Casella, L., Pezzoli, G., Langley, J.,
Hu, X. P., Zucca, F. A., Isaias, I. U., & Zecca, L. (2018). Neuromelanin detection by
magnetic resonance imaging (MRI) and its promise as a biomarker for Parkinson's
disease. *Npj Parkinson's Disease*, *4*(1), Article 1. https://doi.org/10.1038/s41531-018-
0047-3

Takahashi, H., Watanabe, Y., Tanaka, H., Mihara, M., Mochizuki, H., Liu, T., Wang, Y., &
Tomiyama, N. (2018). Quantifying changes in nigrosomes using quantitative
susceptibility mapping and neuromelanin imaging for the diagnosis of early-stage
Parkinson's disease. *The British Journal of Radiology*, *91*(1086), 20180037.
https://doi.org/10.1259/bjr.20180037

Trujillo, P., Aumann, M. A., & Claassen, D. O. (2023). Neuromelanin-sensitive MRI as a
promising biomarker of catecholamine function. *Brain*, awad300.
https://doi.org/10.1093/brain/awad300

Tsukahara, J. S., & Engle, R. W. (2021). Fluid intelligence and the locus coeruleus–
norepinephrine system. *Proceedings of the National Academy of Sciences*, *118*(46),
e2110630118. https://doi.org/10.1073/pnas.2110630118

Tsukahara, J. S., Harrison, T. L., & Engle, R. W. (2016). The relationship between baseline
pupil size and intelligence. *Cognitive Psychology*, *91*, 109–123.
https://doi.org/10.1016/j.cogpsych.2016.10.001

Turker, H. B., Riley, E., Luh, W. M., Colcombe, S. J., & Swallow, K. M. (2021). Estimates of
locus coeruleus function with functional magnetic resonance imaging are influenced by

localization approaches and the use of multi-echo data. *NeuroImage*, *236*. https://doi.org/10.1016/j.neuroimage.2021.118047

van der Pluijm, M., Cassidy, C., Zandstra, M., Wallert, E., de Bruin, K., Booij, J., de Haan, L., Horga, G., & van de Giessen, E. (2021). Reliability and Reproducibility of Neuromelanin-Sensitive Imaging of the Substantia Nigra: A Comparison of Three Different Sequences. *Journal of Magnetic Resonance Imaging*, *53*(3), 712–721. https://doi.org/10.1002/JMRI.27384

Van Egroo, M., Koshmanova, E., Vandewalle, G., & Jacobs, H. I. L. (2022). Importance of the locus coeruleus-norepinephrine system in sleep-wake regulation: Implications for aging and Alzheimer's disease. *Sleep Medicine Reviews*, *62*, 101592–101592. https://doi.org/10.1016/J.SMRV.2022.101592

Venegas, J., & Clark, E. (2011). National Adult Reading Test. In J. S. Kreutzer, J. DeLuca, & B. Caplan (Eds.), *Encyclopedia of Clinical Neuropsychology* (pp. 1705–1705). Springer. https://doi.org/10.1007/978-0-387-79948-3_1467

Watanabe, T., Tan, Z., Wang, X., Martinez-Hernandez, A., & Frahm, J. (2019). Magnetic resonance imaging of noradrenergic neurons. *Brain Structure and Function*, *224*(4), 1609–1625. https://doi.org/10.1007/s00429-019-01858-0

Xing, Y., Sapuan, A., Dineen, R. A., & Auer, D. P. (2018). Life span pigmentation changes of the substantia nigra detected by neuromelanin-sensitive MRI. *Movement Disorders*, *33*(11), 1792–1799. https://doi.org/10.1002/mds.27502

Youdim, M. B., Ben-Shachar, D., & Riederer, P. (1994). The enigma of neuromelanin in Parkinson's disease substantia nigra. *Journal of Neural Transmission Supplementum*, *43*, 113–122.

Zarow, C., Lyness, S. A., Mortimer, J. A., & Chui, H. C. (2003). Neuronal loss is greater in the locus coeruleus than nucleus basalis and substantia nigra in Alzheimer and Parkinson

diseases. *Archives of Neurology*, *60*(3), 337–341.

https://doi.org/10.1001/archneur.60.3.337

Zecca, L., Zucca, F. A., Wilms, H., & Sulzer, D. (2003). Neuromelanin of the substantia nigra: A neuronal black hole with protective and toxic characteristics. *Trends in Neurosciences*, *26*(11), 578–580. https://doi.org/10.1016/j.tins.2003.08.009

Zhang, W., Phillips, K., Wielgus, A. R., Liu, J., Albertini, A., Zucca, F. A., Faust, R., Qian, S. Y., Miller, D. S., Chignell, C. F., Wilson, B., Jackson-Lewis, V., Przedborski, S., Joset, D., Loike, J., Hong, J.-S., Sulzer, D., & Zecca, L. (2011). Neuromelanin Activates Microglia and Induces Degeneration of Dopaminergic Neurons: Implications for Progression of Parkinson's Disease. *Neurotoxicity Research*, *19*(1), 63–72.

https://doi.org/10.1007/s12640-009-9140-z

Zucca, F. A., Bellei, C., Giannelli, S., Terreni, M. R., Gallorini, M., Rizzio, E., Pezzoli, G., Albertini, A., & Zecca, L. (2006). Neuromelanin and iron in human locus coeruleus and substantia nigra during aging: Consequences for neuronal vulnerability. *Journal of Neural Transmission*, *113*(6), 757–767. https://doi.org/10.1007/s00702-006-0453-2

Zucca, F. A., Segura-Aguilar, J., Ferrari, E., Muñoz, P., Paris, I., Sulzer, D., Sarna, T., Casella, L., & Zecca, L. (2017). Interactions of iron, dopamine and neuromelanin pathways in brain aging and Parkinson's disease. *Progress in Neurobiology*, *155*, 96–119.

https://doi.org/10.1016/j.pneurobio.2015.09.012

| Contrast-enhanced Turbo Spin Echo image | Sagittal x = 6 (whole brain) | Axial z = -30 (brainstem) | Coronal y = -36 (brainstem) |

Fig. 1. On the left, an example of contrast-enhanced turbo spin echo image in (axial view, limited to brainstem) for localization of the locus coeruleus (bright spots indicated by purple arrows). On the right, three views of a probabilistic map of LC location using data from 96 individuals. MNI coordinates.

Fig. 2. (A) Boxplots showing distribution of total LC volume (left and right combined) across age groups. (B) Histogram of total LC volume for all participants. (C) Boxplots showing distribution of neuromelanin intensity (average per person, including left and right) across age groups. (D) Histogram of average neuromelanin intensity for all participants. (E) Density plot showing distribution of minimum neuromelanin (NM) intensity values by age group. Minimum NM intensity is the least intense voxel that was included in each individual's hand-drawn LC ROI. In box plots, the middle line shows median, the top and bottom of the box show 75th and 25th percentiles, the top and bottom whiskers show 95th and 5th percentiles, and dots are outliers. Neuromelanin intensity is measured in multiples of the maximum non-LC, non-outlier voxel value in the brainstem.

Fig. 3. At left, comparison of volume of left and right LC, overall (A) and by age group (B). In the middle, comparison of neuromelanin intensity of left and right LC, overall (C) and by age group (D). On the right, comparison of neuromelanin intensity between rostral and caudal LC, overall (E) and by age group (F). All plots show mean and standard error. Neuromelanin intensity is measured in multiples of the maximum non-LC, non-outlier voxel value in the brainstem.

Fig. 4. (A) Scatterplot comparing cognitive performance (national percentile for age in fluid cognition composite from NIH toolbox) and rostral LC neuromelanin intensity. (B) Scatterplot comparing cognitive performance amongst above-average (greater than 50th national percentile for age) performers only to rostral LC neuromelanin intensity. (C) Scatterplot comparing cognitive performance amongst below-average (less than 50th national percentile for age) performers only to rostral LC neuromelanin intensity. Lines show linear fit and 95% confidence interval. r is Pearson's correlation coefficient. Neuromelanin intensity is measured in multiples of the maximum non-LC, non-outlier voxel value in the brainstem.

Fig. 5. (A) Boxplot showing distribution of rostral LC neuromelanin intensity across condensed age groups and between female and male participants. (B) Boxplot showing distribution of rostral LC neuromelanin intensity across condensed age groups and between Black and White participants. In box plots, the middle line shows median, the top and bottom of the box show 75th and 25th percentiles, the top and bottom whiskers show 95th and 5th percentiles, and dots are outliers. Neuromelanin intensity is measured in multiples of the maximum non-LC, non-outlier voxel value in the brainstem.