The IRE1-bZIP60 branch of Unfolded Protein Response is required for *Arabidopsis* immune response to *Botrytis cinerea*

Blanchard Cécile[1], Aimé Sébastien[1], Ducloy Amélie[2], Hichami Siham[1], Azzopardi Marianne[2], Cacas Jean-Luc[2] and Lamotte Olivier[1*]

[1]Agroécologie, INRAE, Institut Agro Dijon, CNRS, Université de Bourgogne, Université de Bourgogne Franche-Comté, DIJON, France

[2]Institut Jean-Pierre Bourgin, Université Paris-Saclay, INRAE, AgroParisTech, VERSAILLES, France.

*Corresponding author: olivier.lamotte@inrae.fr

Abstract

The Unfolded Protein Response (UPR) is a retrograde signalling pathway which is activated when endoplasmic reticulum (ER) proteostasis is disturbed. Here, we have investigated by reverse genetics the contribution of such pathway in *Arabidopsis thaliana* response to two necrotrophic fungi of agricultural importance, *Botrytis cinerea* which is responsible for the development of grey mold disease, and *Alternaria brassicicola* which triggers black spot disease. We found that the branch of UPR dependent on the INOSITOL-REQUIRING ENZYME 1 (IRE1) and the transcription factor (TF) bZIP60 is required to restrict foliar necrotic symptoms induced by both fungi. Accordingly, focussing on *B. cinerea*, we provided evidence for the production of the active bZIP60 form during infection. This activation was accompanied by an increased expression of UPR-responsive genes coding for ER-localized chaperones and co-chaperones that belong to the ER-Quality Control (ER-QC) system. Furthermore, mutants deficient for two ER-QC components were also more susceptible to infection. By contrast, investigating the involvement of CELL DIVISION CYCLE 48 (CDC48) AAA+-ATPAses that assist ER-Associated Degradation (ERAD) pathway for disposal of luminal unfolded proteins, we showed that a series of mutants and transgenics are more resistant to grey mold disease. Seeking for molecular insights into how the ER could shape Arabidopsis immune response to *B. cinerea*, we quantified the expression of defence gene and cell death markers in single *bzip60* and double *ire1* mutants. However, none of those genes were mis-regulated in mutant genetic backgrounds, indicating that IRE1-bZIP60 branch of UPR modulates the Arabidopsis response to *B. cinerea* by a yet-to-be-identified mechanism. Interestingly, we identified the NAC053/NTL4 TF as a potential actor of this unknown mechanism, linking the UPR and proteasome stress regulon.

Author summary

Necrotrophic fungi are one of the most economically significant plant pathogens worldwide, inflicting massive pre- and post-harvest losses on a wide range of fruit and vegetable crops. They adopt a necrotrophic lifestyle, deriving their nutrients predominantly from dead plant tissues to complete their life cycle. *Botrytis cinerea* is the causal agent of grey mold and no plant shows complete resistance towards this pathogen. The use of genetic models such as the plant *Arabidopsis thaliana* has partially enabled the understanding of the immunity mechanisms involved in the plant's response to *B. cinerea*. Our work provides new insights into the cellular mechanisms of how plants cope with this pathogen. In this context, by means of a reverse genetic approach, we explored the role of the Unfolded Protein Response (UPR), a cell signalling pathway regulating protein homeostasis within the endoplasmic reticulum (ER) and thus protecting cells from a harmful over-accumulation of aberrant or misfolded proteins.

Introduction

The maintenance of protein homeostasis is one of the cornerstones of cellular functions. It involves the precise regulation of translation, the folding of newly synthesized proteins, their post-translational modifications, their sorting and trafficking within the cell, and their degradation (1). The endoplasmic reticulum (ER) plays crucial roles in these processes, supporting the synthesis of one third to a quarter of total proteins in eukaryotic cells (2). Environmental factors, such as pathogen infection, intensify the workload on the protein folding machinery. When the endoplasmic reticulum (ER) is unable to meet the cell's demand for protein folding, unfolded or misfolded polypeptides accumulate within the ER lumen, leading to a proteotoxic stress known as ER stress. This situation actuates the Unfolded Protein Response (UPR), a retrograde, ER-to-nucleus, signalling pathway, that is conserved across kingdoms. In Arabidopsis, the canonical UPR invokes three bZIP transcription factors (TF) that are anchored in ER membranes (bZIP17, bZIP28 and bZIP60) and made soluble, therefore active, through two distinct mechanisms when ER stress occurs. One mechanism is the non-conventional splicing of *bZIP60* mRNAs that results in a shift in the open reading frame, eventually causing the translated proteins to be devoid of their transmembrane domains. Such splicing process relies on the ribonuclease activity of two ER-resident transmembrane proteins, the INOSITOL-REQUIRING ENZYME 1A (IRE1A) and B (IRE1B) (3,4). When ER stress is persistent, IRE1 ribonuclease activity can also be employed to dispose of ER-associated mRNAs , a phenomenon known as Regulated IRE1-Dependant Decay (RIDD; (5,6); the latter of which reduces the amount of nascent polypeptides crossing ER membranes, thus preventing the accumulation of unfolded proteins within the ER lumen. The other activation mechanism of UPR involves the translocation of bZIP17 and/or bZIP28 TF to the Golgi apparatus through vesicular trafficking, their intramembrane proteolytic cleavage and subsequent release of their transcription-activating domains that can then reach the nucleus (7,8). Target genes of UPR TF encompasses those coding for chaperones, co-chaperones and additional factors involved in the ER-Quality Control (ER-QC) system, increasing ER folding capacity. This includes calreticulin (CRT) and calnexin (CNX), as well as BINDING-IMMUNOGLOBULIN PROTEINS (BiP) and its interacting proteins STROMAL CELL-DERIVED FACTOR 2

(SDF2) and ENDOPLASMIC RETICULUM DNAJ 3 (ERdj3) (9). In support of the transcriptional
reprogramming mediated by the UPR TF, mis-folded and aberrant proteins can also be retro-
translocated from ER lumen to cytoplasm where they are ubiquitinated for proteasomal degradation.
This process, so-called ER-associated degradation (ERAD), involves the HRD1 (3-Hydroxy-3-
methylglutaryl Reductase Degradation 1) complex and the AAA+-ATPase family of proteins CDC48
(Cell Division Control protein 48, (9). It aims at restoring ER proteostasis. When adaptive UPR
pathway fails to resolve ER stress, a genetic program that ends with the death of malfunctioning cells
is engaged. In this context, it has been demonstrated that ER stress-induced cell death (ERSID) is
under the control of NAC089 TF (10) and requires several cathepsin B proteases to be executed (11).

A plethora of environmental situations can activate the UPR in plants, including heat, salt and
drought stress, as well as pathogen infection (12,13). However, only a few data report on UPR
involvement in the host plant response to necrotrophic pathogens. For instance, it was
demonstrated that inoculating *Nicotiana attenuata* with *Alternaria alternata* resulted in the
activation of UPR, and silencing *NaIRE1* or *NabZIP60* genes rendered plants more susceptible to the
fungus (14). It was also showed that impairment of the IRE1/bZIP60 branch enhanced Arabidopsis
susceptibility to *Drechslera gigantea*, another necrotrophic fungus (15). Although diseases caused by
the necrotrophic fungi *Botrytis cinerea* and *Alternaria brassicicola* rank among the most devastating
plant diseases worldwide, there are no data documenting the role of UPR in response to these two
necrotrophic pathogens. *Botrytis cinerea* causes grey mold disease in a wide range of crops, and
Alternaria brassicicola is responsible for black spot disease in numerous Brassica species. Both
pathogens have the ability to infect the plant model *Arabidopsis thaliana*, making it a valuable tool
for unravelling the intricate signalling pathways involved in plant immune response to these
pathogens. Innate immunity in plants is orchestrated through signalling cascades involving key
hormones like salicylic acid (SA), jasmonic acid (JA), and ethylene (ET), inducing significant
modifications in gene expression that underlie enhanced resistance (16). The JA signalling pathway is
commonly linked to the establishment of plant defence against necrotrophic fungus (17). Arabidopsis
mutants with impaired JA perception, production or signalling are notably vulnerable to *B. cinerea* or
A. brassisicola (18,19). SA is recognized as a pivotal hormone that initiates immune responses against
biotrophic pathogens (16), but SA-mediated signalling has also been shown to be involved in
resistance against *B. cinerea* in Arabidopsis (19) or against *Alternaria solani* in potato (20). For a
comprehensive probing of hormonal signalling cascades, studying the expression profile of marker
genes is a widely established method. Activation of JA-dependent signalling pathway is frequently
assessed by investigating the expression of genes such as *PLANT DEFENSIN 1.2* (*PDF1.2*) or
OCTADECANOID-RESPONSIVE ARABIDOPSIS 59 (*ORA59*), whereas *PATHOGENESIS-RELATED-1 (PR1)*
serves as a marker gene for SA (21). Another typical response of plants facing necrotrophic
pathogens is the production in Arabidopsis of the phytoalexin camalexin through SA or JA signalling
pathways, depending on the invading pathogen (22). The last step of camalexin synthesis is catalysed
by the cytochrome P450 CYP71B15 (PHYTOALEXIN DEFICIENT 3, PAD3) which mRNA accumulation
correlates with camalexin production (23). Camalexin has been shown to be toxic for *B. cinerea* and
A. brassicicola (24,25), and *pad3* mutant shows enhanced susceptibility to both pathogens (19,26).
During infection process, necrotrophic pathogens secrete compounds that enable rapid host cell
death and disease spreading. *B. cinerea* produces cell-death inducing proteins, but also manipulates
the plant regulated cell death to promote host cell death (27,28).

In this study, we explored how the ER could shape *Arabidopsis thaliana* immune response to
necrotrophic pathogens, with a specific emphasis on *Botrytis cinerea*. The role of UPR, ER-QC and
ERAD in this context was investigated using a reverse genetic approach in order to gain insights into
molecular events that govern plant susceptibility and defence

136
137 **Results**
138
139 **The IRE1-bZIP60 branch of Arabidopsis UPR restricts lesions induced by necrotrophic pathogens.**
140
To investigate a potential role of the UPR pathway in Arabidopsis response to necrotrophic
fungi, we inoculated mutants defective in canonical UPR actors (*bzip17-1, bzip28-2, bzip60-3, ire1a-2*
ire1b-4) with *B. cinerea* or *A. brassicicola*. Mutant susceptibility to *B. cinerea* was examined 3 days
upon drop-inoculation of leaves with a solution of conidia using the lesion diameter as proxy. The
necrotic lesion diameter was averaging 5 mm in the WT genetic background, as previously reported
(29), so was it in the *bzip28-2* mutant. By contrast, a significant decrease of 18 % in lesion size was
recorded in *bzip17-1* mutant when compared to WT. Interestingly, both the single *bzip60-3* and
double *ire1a-2 ire1b-4* mutants were more susceptible to the infection, displaying a 17 % increase in
lesion diameter with respect to the WT genotype (Figure 1A, Supplemental Figure S1A). *A.*
brassicicola susceptibility was also determined by inoculating leaves with a spore solution and lesion
diameter were measured at 5 days post-inoculation. In WT plants, *A. brassicicola* induced necrotic
lesions of 2.5 mm diameter in average. Both *bzip17-1* and *bzip28-2* mutations had no impact on
symptoms. Genetic inactivation of *bZIP60* or *IRE1A* and *IRE1B* led to a 37 % or 27 % increase in lesion
diameter with regard to WT, respectively (Figure 1B, Supplemental Figure S1B). Altogether, our data
suggest that the IRE1-bZIP60 branch of UPR could be activated and participate either in defence
mounting or PCD inhibition triggered by the two necrotrophic fungal pathogens tested.
157
158 **The IRE1-bZIP60 branch of UPR is activated in response to *Botrytis cinerea*.**
159
To get into molecular mechanisms involving UPR, we focussed on the interaction between *A.*
thaliana and *B. cinerea*. The expression of *IRE1A, IRE1B, bZIP17, bZIP28* and *bZIP60* was followed by
RT-qPCR over time upon infection. For that purpose, a solution of *B. cinerea* spores prepared in PDB
medium or PDB alone (mock) was sprayed on leaves and samples were collected at 16, 24, 48 and 72
h post-infection (hpi) for further processing. Those experimental conditions were validated by
checking the expression of defence genes previously reported to be markedly induced in response to
B. cinerea, i.e. PAD3, PR1, PDF1.2a and *PATATIN-LIKE PROTEIN 2 (PLP2)* (19,24,30). As expected, all
four genes were up-regulated during infection (Supplemental Figure S2). As for UPR actors-encoded
genes, *IRE*1A and *IRE1B* were not differentially expressed between mock and challenged conditions.
A slight, but significant accumulation of *bZIP17* and *bZIP28* transcripts was only observed at 48 hpi
(Figure 2). However, in accordance to mutant over-susceptibility to *B. cinerea* (Figure 1A), the
strongest transcriptional response was observed for *bZIP60* gene with the unspliced mRNAs
accumulating at early time points (16 and 24 hpi) whereas the spliced mRNA levels kept increasing
from 16 hpi to 72 hpi. In addition to confirming the activation of the IRE1-bZIP60 branch, these
results suggest that *B. cinerea* infection induces an ER stress.
175
176 **Default in ER-QC machinery leads to enhanced necrotic lesions caused by *Botrytis cinerea***
177
We next determined by RT-qPCR the expression kinetic of ER stress gene hallmarks (*BIP1, BiP2, BiP3,*
ERDJ3A, ERDJ3B and *SDF2*) in mock and inoculated WT plants (Figure 3A). Consistent with a proteo-
toxic ER stress induced by *B. cinerea*, all genes coding for ER-QC components were up-regulated upon
infection, yet with distinct expression profiles. While *BIP1* and *BIP2* mRNAs were over-accumulating
as early as 16 hpi, the expression of *BIP3* gene tends to increase at 48 hpi. The expression level of the
three BiP-encoding genes were back to that of mock conditions at 72hpi. When compared to mock

plants, the infection by *B. cinerea* was also stimulating the expression of *ERDJ3A*, *ERDJ3B* and *SDF2*
genes at 24 and/or 48 hpi depending on the gene tested. These results prompted us to investigate
whether SDF2 and ERDJ3B could be involved in Arabidopsis response to *B. cinerea*. For that purpose,
we used *erdj3b-1* and *sdf2-2* mutants (31) that we challenged with a spore solution and lesion
diameters were measured at 3 days post-inoculation (Figure 3B, Supplemental Figure S3). Significant
increases in lesion diameter were recorded for both mutants (24 % for *erdj3b-1* and 18 % for *sdf2-2*)
in comparison to WT plants, indicating that they are more susceptible to *B. cinerea*. These data
indicate that both SDF2 and ERDJ3B could be acting in the same response pathway as that of IRE1-
bZIP60.
193
194
195   **Mutation in ERAD-related *CDC48* genes decreases susceptibility to *Botrytis cinerea***
196
The ERAD utilizes proteasome to dispose of misfolded proteins which have been retro-translocated
from ER lumen to cytoplasm. With ER-QC system which is reinforced to increase ER folding capacity
under ER stress, ERAD is part of the pro-adaptive mode of UPR that aims at restoring ER homeostasis.
The CDC48 family of AAA+-ATPase proteins have been described as ERAD actors in yeast (32),
mammals (33) and plants (34). To investigate a potential involvement of ERAD in response to *B.*
cinerea, we first functionally characterized Arabidopsis CDC48 homologs. In Arabidopsis, CD48
proteins are, indeed, likely to be encoded by five genes (35). CDC48A, B and C are the closest
sequence orthologs of human and yeast proteins, with which they share approximately 90 %
similarity. By contrast, CDC48D and E are more divergent, exhibiting no or little conservation within
their N and C-terminal domains (35,36). In accordance with its role in ERAD, CDC48A was shown to
control the turnover of two immune receptors (36,37), as well as that of a mutated carboxypeptidase
Y protein which is an ERAD substrate (34). It was also capable of complementing yeast *cdc48*
conditional mutant (37). The four Arabidopsis homologs of *CDC48A* were thus tested by yeast
functional complementation using the *Saccharomyces cerevisiae cdc48* mutant strains KFY189 and
DBY2030. KFY189 is a temperature-sensitive mutant which does not grow at 37°C whereas DBY2030
is cold-sensitive and fail to grow at 16°C (38,39). Both strains were transformed with the yeast
expression vector pDRF1-GW containing either the cDNA coding for CDC48A, B, C, D or E or with the
empty vector alone. All transformed strains were able to grow under permissive conditions at 30°C
(Figure 4A, C). As previously described (38), CDC48A complemented both strains under non-
permissive conditions while the empty vector did not, validating our experimental setup (Figure 4B,
D). Whereas *CDC48B* or *CDC48C* were able to restore the growth of DBY2030 strain at 16°C and that
of KFY189 strain at 37°C (Figure 4B, D), *CDC48D* and *CDC48E* could not rescue the temperature-
dependent phenotypes of both strains (Figure 4B, D). These data demonstrate that only CDC48A, B
and C can functionally substitute to the yeast protein, and suggest a role for CDC48B and C in ERAD.
Focusing on *CDC48A*, *B* and *C*, we next found that only *CDC48A* and *CDC48B* showed an
increased expression in response to *B. cinerea*, at 48 and 72 hpi (Figure 5A). We then evaluated the
susceptibility of different *cdc48* mutants to *B. cinerea*. We used *cdc48a-4/muse8* which is a partial
loss-of-function EMS mutant previously described (36), the knock-out *cdc48b3* (GABI_104F08) and
knock-down *cdc48b4* (GABI_485G04) mutants that show no or reduced *CDC48B* expression,
respectively (Supplemental Figure S4A), and the two knock-out *cdc48c2* (SALK_102955C) and *cdc48c3*
(SALK_123409) that do not express *CDC48C* (Supplemental Figure S4B). When compared to
challenged WT plants, *cdc48a-4/muse8* and *cdc48b-4* mutants were slightly, but significantly more
resistant to *B. cinerea*, as judged by the lesion size (Figure 5B, Supplemental Figure S5). *cdc48b-3* and
cdc48c-2 mutants were even more resistant, and *cdc48c-3* exhibited the strongest resistance with a
37% reduction in lesion diameter. We also produced Arabidopsis lines overexpressing either a WT

AtCDC48B protein or a mutated one (AtCDC48B^{E308QE581Q} further abbreviated AtCDC48B-QQ) under
the control of a CaMV35S promoter, the latter being a negative dominant allele (40). Two
overexpressing lines were selected for each construct (Supplemental Figure S4C). Upon challenged
with *B. cinerea*, plants overexpressing AtCDC48B-QQ were more resistant to the infection whereas
plants overexpressing AtCDC48B-WT were as susceptible as WT plants (Figure 5B, Supplemental
Figure S5). These data are consistent with results obtained using T-DNA insertion lines (Figure 5B),
and further indicate that CDC48 activity, likely though ERAD pathway, is required for full Arabidopsis
susceptibility to *B. cinerea.*
240
241 **Inactivation of IRE1/bZIP60 branch impacts neither the expression of *B. cinerea*-induced defence**
242 **genes, nor the expression of ER-stress-induced cell death genes.**
243
Following *B. cinerea* perception, plant cells activate a complex array of defence mechanisms
that relies, at least in part, on a transcriptional reprogramming. Among genuine defence gene
markers induced upon B. cinerea infection, one may cite *PR1, PDF1.2a, ORA59, FLG22-INDUCED*
RECEPTOR-LIKE KINASE 1 (*FRK1*), *PAD3* and *GLUTHATIONE S-TRANSFERASE 6* (*GSTF6*) (Supplemental
Figure S2; (24,41,42)). We reasoned that *bzip60-3 and ire1a-2 ire1b-4* mutants could have a reduced
defence response if bZIP60 transcription factor positively regulates defence gene expression. To test
this hypothesis, changes in the aforementioned gene expression was measured by qRT-PCR in the
single and double mutants and compared to WT. In a WT genetic background, infection triggered an
increase in expression for all genes at 24 and 48 hpi, except for *PR1* whose enhanced transcript level
was only detected at 48 hpi (Supplemental Figure S6). This expression pattern is consistent with
previous work (41). However, neither *ire1a-2 ire1b-4*, nor *bzip60-3* mutations impacted the
expression profiles of the tested defence marker genes in response to *B. cinerea* (Supplemental
Figure S6).
Evidence have been accumulating that necrotrophic pathogens, like *B. cinerea*, actively
induce host PCD for feeding on cell debris and subsequent colonization of dead tissues (27). With this
regard, we hypothesized that ERSID could be over-induced in the *bzip60-3 and ire1a-2 ire1b-4*
mutants, potentially explaining their enhanced susceptibility to the fungus (Figure 1A). ERSID was
found to be positively controlled by the NAC089 transcription factor (10) and the protease family of
cathepsins (11) under chemically-induced ER stress. Whereas *NAC089* and *CATHEPSIN B1, B2, B3*
gene expression was previously shown to be significantly increased during ERSID triggered by
tunicamycin (10,11), none of those genes were differentially expressed between mock and infection
(24 and 48 hpi) conditions in a WT genetic background (Supplemental Figure S7). In addition, no or
minor expression changes for those genes could be recorded in *bzip60-3 and* double *ire1a-2 ire1b-4*
mutants compared to WT, whether plants were inoculated or not (Supplemental Figure S7).
268
269 **The expression of NAC053, a negative regulator of Arabidopsis defence against *B. cinerea*, is**
270 **dependent on IRE1 proteins.**
271
To explain the susceptibility phenotype of *ire1a-2 ire1b-4* and *bzip60* mutants and the
resistant phenotype of *cdc48* mutants, we tested the hypothesis that the IRE1/bZIP60 branch of the
UPR negatively control *CDC48* gene expression and its regulators in response to *B. cinerea*. None of
the *CDC48A, B, C* genes were mis-regulated at 24hpi and 48hpi in *bzip60-3* and *ire1a-2-ire1b-4*
mutants (Supplemental Figure S8). The expression of the two NAC TF (NAC053/NTL4 and
NAC078/NTL11) that have been shown to regulate proteasome stress regulon-encoding genes,
including *CDC48A* (Gladman *et al.*, 2016), was next checked in the same genetic backgrounds (Figure
6). The expression level of *NAC078* was not altered by the infection in any of the genotypes for the

two time points tested (Figure 6A-B). By contrast, in response to *B. cinerea*, *NAC053* expression was
induced earlier in the *bzip60-3* mutant (24hpi, Figure 6A) when compared to WT (48hpi, Figure 6B),
and was totally abolished in the double *ire1a-2-ire1b-4* mutant at both time points (Figure 6A-B).
Infected *nac053-1* and *nac078-1* mutants showed smaller lesions at 3 dpi with respect to WT plants
(Figure 6C, Supplemental Figure S9), indicating that both mutants are more resistant to *B. cinerea*.
These results indicate that *NAC053* expression is dependent on IRE1 proteins, and that both NAC053
and NAC078 act as negative regulator of Arabidopsis defence against *B. cinerea*.

287
288
289

Discussion

The UPR pathway plays a crucial role in plant-pathogen interactions, irrespective of the
pathogen nature, *i.e.* bacteria, fungi and viruses (13,44). However, our understanding of its specific
role during plant infection by necrotrophic fungi remains elusive. Here we present a study that
identifies UPR pathway as a critical element in *Arabidopsis thaliana* response to two necrotrophic
fungi, *Botrytis cinerea* and *Alternaria brassicicola*. The RT-qPCR results demonstrated that *B. cinerea*
infection activated UPR pathways in WT plants. During infection, we observed an accumulation of
mRNAs coding the three BIP isoforms, along with its co-chaperones SDF2.2, ERDJ3A and ERDJ3B.
Additionally, we detected an increased level of spliced bZIP60 mRNA variants coding the active TF.
The importance of UPR has been confirmed by challenging plants mutated in the genes coding the
canonical UPR regulators IRE1A/IRE1B, bZIP60, as well as those of RIP-UPR bZIP17 and bZIP28, with a
spore solutions of *B. cinerea*. Mutation in the genes coding *IRE1A* and *IRE1B* or *bZIP60* results in a
greater plant susceptibility to *B. cinerea*. Similar observations were made when these mutants were
challenged with *Alternaria brassicicola* indicating that IRE1-bZIP60 pathway plays important role in
limiting the growth of both pathogens. However, it is worth noting that mutation in *bZIP28* did not
have a significant impact on the plant susceptibility to these fungi. Overall, our findings support the
hypothesis that the IRE1-bZIP60 branch is an important element in the establishment of plant
immunity to necrotrophic pathogens. This is in accordance with previous work on *Nicotiana*
benthamiana, where silencing of *NbbZIP60*, *NbIRE1a*, and *NbIRE1b* expression through a VIGS
approach led to increased plant susceptibility to the necrotrophic fungus *Alternaria alternata* (14).
Furthermore, infection of the double mutant *bzip28-2-bzip60-1* with *Drechslera gigantea*, a
necrotrophic fungus causing eyespot disease in crop plants, leads to increase disease symptoms
compared to WT plants in Arabidopsis (15). However, it is noteworthy to mention that neither the
knock down *bzip60.1* nor the double mutant *bzip28-2-bzip60-1* showed increased susceptibility when
infected with *B. cinerea* (Supplemental Figure S10). The significance of the IRE1-bZIP60 branch of the
UPR in controlling plant immunity is further demonstrated by the work of Tateda *et al.* (45), who
showed that silencing of *IRE1a/1b* or *bZIP60* resulted in disease symptoms in *Nicotiana tabacum*
caused by the non-host bacterial pathogen *Pseudomonas cichorri*. Additionally, concerning the
virulent bacteria *Pseudomonas syringae pv maculicola* ES4326, bacterial growth was substantially
more pronounced in *ire1a ire1b* or *bzip60* infected mutants compared to the infected WT plants (46).
It is must be pointed out that our study revealed distinct responses to *B. cinerea* and *A. Brassicicola*
depending on the genotype tested. Notably, the *bzip17* mutant demonstrated higher resistance to *B.*
cinerea compared to WT plants, while it exhibited a similar level of infection as WT plants when
exposed to *A. brassicicola*. Studies demonstrating a role for bZIP17 during biotic stresses are scarce.
One may cite the study of Li (47) where the silencing of both *NbbZIP17* and *NbbZIP28* delayed Rice
Stripe Virus infection and decreased the accumulation of viral RNA and proteins in *N. benthamiana*,

indicating that RIP-UPR promotes virus development. However, in Arabidopsis infected with the
Plantago asiatica mosaic virus, bZIP60 and bZIP17 acts synergistically to restrict viral infection (48).
Understanding the role of bZIP17 during *B. cinerea* infection will need further investigations.

Activation of UPR leads to an increase of the ER folding capacity, preventing an excessive
build-up of misfolded or aberrant proteins that may occur during stressful situations like pathogen
attacks. Unfolded and misfolded proteins can also be degraded through ERAD. During this process,
misfolded proteins are tagged with polyubiquitin chains and extracted from the ER membrane with
the assistance of the CDC48 AAA+ATPase complex (9). The number of gene coding CDC48 proteins in
Arabidopsis genome is unclear (35). Through yeast *cdc48* mutant complementation experiments, our
findings unequivocally establish that only three proteins - AtCDC48A (At3g09840), AtCDC48B
(At3g53230), and AtCDC48C (At5g03340) - are functional orthologs of ScCDC48 (as demonstrated in
our study and by Feiler *et al.* for CDC48A (38)). Thus, the two closely related AAA+-ATPase proteins
(At2g03670 and At3g01610), previously referred to as AtCDC48D and E (35,36), are not ScCDC48
functional orthologues and should not be categorized as such.

An increase in the accumulation of *AtCDC48A* and *AtCDC48B* mRNA have been detected
during *B. cinerea* infection. Interestingly, plants mutated in either *AtCDC48A*, *AtCD48B* or *AtCDC48C*
genes are more resistance to *B. cinerea* infection. Furthermore, the overexpression of an inactive
form of AtCDC48B (*eg* AtCDC48B^{QQ}) also results in an increased resistance phenotype toward *B.*
cinerea. The resistant phenotype of *atcdc48a-4* mutant to *B. cinerea* could be tentatively explained
by a constitutive immune response characterized by the stronger accumulation of *PR2* mRNA in this
mutant (36). Ao et *al.* (49) have proposed a model to explain this autoimmune phenotype related to
the role of AtCDC48A during plant pathogen interaction. AtCDC48A is recognized by the SFC^{SNIPER7}
complex and targeted for proteasomal degradation. It results in the accumulation of the NLR SNC1
(and likely other NLR) which may serve to enhance defence to the virulent oomycete pathogen *H.*
arabidopsidis NOCO2. The existence of such a mechanism in the context of *B. cinerea* infection
remains to be demonstrated. Regarding AtCDC48A paralogs, AtCDC48B and AtCDC48C, it is unlikely
that they have redundant functions with AtCDC48A. This is supported by the observations that
mutation in each of the *AtCDC48* gene results in resistant phenotypes against *B. cinerea*.
Nonetheless, mutations in *AtCDC48B* or *AtCDC48C* do not affect the resistance of Arabidopsis to *H.*
Arabidopsis NOCO2 (36). However, the importance of the IRE1-bZIP60 pathway in basal resistance to
B. cinerea infection is unlikely linked to CDC48 because *AtCDC48* mRNAs do not accumulate
differentially in *atire1a/ire1b* or *atbzip60* genetic backgrounds. Understanding the involvement of
each of the *AtCDC48* genes in plant immunity will necessitate additional in-depth investigations.

To identify potential mechanisms that play a role in the defence response against *B. cinerea*
via the IRE1-bZIP60 pathway, we have undertaken a RT-qPCR approach. None of the tested genes
had their expression altered in the *ire1a-2 ire1b-4* and *bzip60-3*. It concerns the SA (*PR1*) and JA
(*PDF1.2, ORA59*) responsive genes, indicating that the IRE1-bZIP60 pathway does not influence the
expression of SA and JA-dependant gene expression in response to *B. cinerea* infection. However, we
did not investigate the hypothesis whether SA or JA signalling pathways were necessary to regulate
the activation of UPR pathways as observed in other studies. In tomato, JA treatment induced the
expression of *BIP1* and tunicamycin-induced *BIP1* expression was strongly reduced in the JA signalling
mutant *jai1* indicating that JA pathway plays a role in ER stress signalling (50). Similar observations
were made in *N. attenuata* in which MeJA treatment induced the accumulation of mRNA coding
chaperone proteins such as BIP, PDI, CNX and CRT as well as the spliced form of *bZIP60* mRNA (14).
Using JA-deficient and JA-insensitive plants, these authors have also shown that JA controls
chaperone gene expression in response to *A. alternata*.

In addition, our data indicate that *PAD3* gene, which encode a cytochrome P450 enzyme that
catalyses the last step of camalexin biosynthesis is not regulated by UPR. To the contrary, the *F6'H1*

gene was found less induced in *N. attenuata* plants impaired in the IRE1-bZIP60 pathway during *A.*
alternata infection (14). It encodes the key enzyme which catalyzes the synthesis of scopoletin and
scopolin, two phytoalexin previously shown to be crucial for *N. attenuata* resistance to *A. alternata*.
These results indicate that the role of UPR in regulating plant resistance to one particular class of
pathogen is different from one plant-pathogen interaction model to another.

Necrotrophic pathogens induce plant cell death to their own benefits either by secreting
plant cell death inducers or by manipulating host regulated cell death (27,28). Genes known to
orchestrate ERSID, such as *NAC089* and *CATHEPSIN B*, are not induced after *B. cinerea* infection
whereas these genes are up-regulated following plant treatment with the ER stress-inducing agent
tunicamycin (10,11). We can formulate two hypotheses. Firstly, increased susceptibility of *ire1a-2*
atire1b-4 and *atbzip60-3* to *B. cinerea* is unlikely the consequence of an increase ERSID phenomenon
because *NAC089* or *Cathepsin B* genes are not upregulated in these mutant genotypes. It suggests
that a NAC089- and CATHEPSIN B-dependant ERSID is unlikely to be involved during Arabidopsis-*B.*
cinerea interaction. Secondly, we can hypothesized that ER stress is not involved in *B. cinerea*-
induced cell death. In any case, further experiments are needed to clarify the role of ER stress in *B.*
cinerea-induced cell death.

Our data indicate that, in *bzip60* mutant infected with *B. cinerea*, the expression of
NAC053/NTL4 is upregulated. Infected-*bzip60* mutant accumulates much more *NAC053* mRNA than
WT plant and this accumulation occurs earlier in *bzip60* (at 24 hpi) than in WT plant (at 48 hpi).
However, the expression of the NAC053 closest relative NAC078/NTL11 is not modulated by *B.*
cinerea in any of the genotypes tested. Together, both transcription factors are required to control
the expression of the proteasome stress regulon including *CDC48A* (43). Nevertheless, the expression
of genes coding CDC48 A, B or C proteins are not modulated in Botrytis-infected *bzip60* mutant. Very
surprisingly, results are different when using *ire1a-ire1b* double mutant. *NAC053* mRNA does not
accumulate neither at 24 hpi nor at 48 hpi in this mutant infected with *B. cinerea* in comparison to
WT plants in which *NAC053* accumulate at 48 hpi. It indicates that *NAC053* expression depends on
IRE1 proteins during *B. cinerea* infection. NAC053 is anchored to the plasma membrane (51) and
activated following stress perception in a ROS-dependant mechanism (52). Once activated, NAC053
control ROS production through the activation of *RBOH* genes (53). In our model, it is reasonable to
suggest that a disturbance in ROS homeostasis is likely present in the *bzip60* mutant, whereas such a
disturbance might not exist in the *ire1a/ire1b* mutant. Further research is necessary to substantiate
this hypothesis.

Summarizing, the infection of wild type Col-0 plant by *Botrytis cinerea* induces ER stress and
activates the UPR pathways (Figure 7). The activation of the IRE1-bZIP60 branch and ERQC is crucial
for the plant's response to the pathogen, as mutations in genes encoding key players in these
pathways result in increased plant susceptibility. Further work is however necessary to understand
how the IRE1/bZIP60 branch contributes to defence against this necrotrophic fungus, particularly its
potential involvement in the secretion of defence proteins. On the other hand, our results indicate
that mutations in the bZIP17 branch of the UPR, as well as mutations in genes encoding the ERAD-
involved proteins CDC48, lead to greater resistance of the plant. This suggests either that these
proteins act as negative regulators of immunity against this pathogen, or that these pathways might
be manipulated by the pathogen during infection. Similarly, further work is needed for a detailed
understanding of these mechanisms.

418
419
420
421

422
423
424
425

Experimental procedures

Plant materials

A. thaliana seeds were grown on jiffy-7 pellets in a growth chamber under controlled conditions (10h light/14h dark, 22°C/18°C, 60–70% humidity, light intensity of 100 and 120 μE.m^{-2}.sec^{-1}). All the mutants used are listed in Supplemental table S1. To screen T-DNA homozygous plant mutants by PCR, genomic DNA was isolated using Phire Plant Direct PCR kit (Thermoscientific) and PCR was performed using primers pairs as indicated in Supplemental table S2.

Pathogen infection

Botrytis cinerea strain BMM (29) was grown 10 days on sterile V8 agar plate (50% Campbells original V8 juice (v/v), 0.5 % KH2PO4 (m/v), 1.5 % agar (m/v), pH 6) at 20°C in the dark. Spores were harvested by scraping the plate with sterile milliQ water and filtered through sterile gaze. Infection tests are carried out by placing 6 µL of a solution of botrytis spores (5.10^4 spores.mL^{-1} diluted in quarter-strength Difco potato dextrose broth PDB) on 4 leaves of four weeks old plants. Lesion diameters were measured after 3 days. For qPCR analysis, plants were either sprayed with a spore solution (2.5.10^5 spores.mL^{-1}) or with quarter-strength PDB as described previously (41). For both experiments, the inoculated plants are placed in a tray closed with a transparent lid to maintain a high-humidity environment.

Alternaria brassicicola strain MIAE01824 was provided by Dr. Christian Steinberg (INRAe, Dijon, France) and grown 15-20 days on PDA medium (19 g.L^{-1}) supplemented with sucrose (20 g.L^{-1}) and CaCO$_3$ (30 g.L^{-1}) at 20°C in the dark. Spores were harvested by scraping the plate with sterile infection medium (GamborB5 medium (Duchefa), sucrose 10 mM, KH$_2$PO$_4$ 10 mM) and filtered through sterile gaze. Four leaves of four weeks old plants were drop inoculated (6 µL) with a solution of 1.10^6 spores.mL^{-1} in infection medium. Inoculated plants are placed in a tray closed with a transparent lid to maintain a high-humidity environment and lesion diameters were scored after five days.

RNA extraction and RT-qPCR analysis

Inoculated and mock-treated leaves were harvested at different time points, flash frozen and finely grinded in liquid nitrogen. Total RNA extraction and DNAse treatment were performed using SV Total RNA Isolation System (Promega) as described by the manufacturer. RNA integrity was analysed on a 1% agarose gel and concentration was measured using a nanodrop system (Thermoscientific). RNA (500 ng) were retrotranscribed into cDNA (High-Capacity cDNA Reverse Transcription Kit, Thermoscientific) using random hexamer and 17-mer oligodT primers. mRNA accumulation was assessed by real-time qPCR (GoTaq® qPCR Master Mix, Promega) and expression values were normalized to the expression of plant genes *At4g26410* and *At3g01150* as previously described as a stable reference genes (54). Levels of transcripts were calculated using efficiency-weighted $\Delta\Delta Cq^\omega$ method (55). All primers are listed in Supplemental table S3.

Statistical methods

For infection tests, significant differences from WT plant were determined by a Kruskal–Wallis one-way anova on ranks followed by a comparison with the Dunnett's method. For qPCR analysis, ΔCq^ω value from one infection time-point is compared to the data of the control condition at the same

470 time point using by a one-way ANOVA followed by a Tukey HSD Test (55). Then, data are represented
471 on the figures as fold induction to the control condition ($\Delta\Delta Cq^{(\omega)}$).
472

AtCDC48B cloning procedures and transgenic lines generation
Arabidopsis lines overexpressing either AtCDC48B or AtCDC48B mutated in its two ATPase sites
(AtCDC48B^{E308QE581Q}) under the control of a CaMV35S promoter were produced as follows.
PDONR/Zeo containing either AtCDC48B^{WT} or AtCDC48B^{E308QE581Q} cDNA were kindly provided by Drs.
Annette Niehl and Manfred Heinlein (IBMP, Strasbourg, France). pDNOR/Zeo were BP-recombined
(Thermoscientific) in pB2GW7 vector (56) and resulting vectors were introduced in *Agrobacterium*
tumefaciens strain GV3101. WT *A. thaliana* Columbia plants (N60,000, Eurasian Arabidopsis Stock
Center) were transformed using the floral dipping method (57). T1 and T2 transgenic plants were
grown on jiffy-7 pellets and selected using BASTA herbicide after 10 days of growth. To identify
homozygous plants overexpressing *AtCDC48B^{WT}* or *AtCDC48B^{E308QE581Q}*, seeds from T2 plants were
surface-sterilized and sown on MS Agar plates (4.4 g.L^{-1} of Duchefa MS powder M0222.0050, 1%
sucrose, 10 mM MES, 1% agar pH 5.7) containing 50 µg/mL of glufosinate ammonium (Sigma-
Aldrich). Expression of mRNA coding AtCDC48B^{WT} or AtCDC48B^{E308QE581Q} in transgenics plants were
analysed by RT-qPCR using primers described in supplemental table S3.
487

CDC48 expression in *cdc48* yeast mutant
CDC48 cDNA were cloned from an Arabidopsis thaliana cDNA pools made from total RNA extracted
from Arabidopsis Col-0 plants. AtCDC48A (At3g09840; forward primer
ATGTCTACCCCAGCTGAATCTTC; reverse primer CTAATTGTAGAGATCATCATCGTCCC), AtCDC48C
(At5g03340; forward primer ATGTCAAACGAACCGGAATC; reverse primer
CTAACTGTAGAGATCGTCGTCATC), AtCDC48D (At2g03670; forward primer ATGTTGGAAACCGAAAGC;
reverse primer TCATGTAGCAGAAGCTACTAGTAA), AtCDC48E (At3g01610; forward primer
ATGGGGAGGAGAGGTCGC; reverse primer TTACTCGAGGGTAAAAGATGGCC) were amplified by PCR
using Phusion DNA polymerase (Thermoscientific) and cloned into PCR8 vector (Thermoscientific).
PDONR/Zeo containing CDC48B (At3g53230) was kindly provided by Drs. Annette Niehl and Manfred
Heinlein (Institut de biologie moléculaire des plantes, Strasbourg, France; (40)). PCR8 and
pDNOR/Zeo vectors were then recombined in pDRF1-GW (58) and resulting vectors or empty vector
were introduced in *cdc48* deficient *Saccharomyces cerevisiae*. Two *cdc48 S. cerevisiae* mutants (39)
were kindly provided by K.-U. Fröhlich (University of Graz, Austria): the cold-sensitive strain DBY2030
(MATa ade2-101 lys2-801 ura3-52 cdc48-1;) or the temperature-sensitive strain KFY189 (MATa lys2
leu2 ura3 cdc48-8). Transformed *S. cerevisiae* cells were plated on minimal selective medium
containing 2% glucose, 0,77g/L SD-URA drop out mix (Clontech), 0.67 g/L Yeast Nitrogen Base (Sigma-
Aldrich), 15 g/L agar (Sigma-Aldrich) and grown at 30°C. Transformed colonies were checked by
colony PCR and then grown overnight at 30°C under shaking in YPD medium (10 g/L yeast extract, 20
g/L peptone, 2% glucose). Yeast cells were then streaked out on YDP agar plate (YPD medium
containing 15g/L agar) and grown at 30°C and 16 °C for DBY2030 transformed strain or 30°C and 37°C
for KFY189 transformed strain.
510

Acknowledgments

513 We would like to thank all the people who provided us with invaluable assistance in carrying out this
514 work and particularly Pr. Nathalie Leborgne-Castel, Dr. Mathieu Gayral, Dr. Stéphane Bourque and Pr.
515 David Wendehenne for helpful discussions. Pr. Steven H Howell, Pr. Cyril Zipfel and Pr. Xin Li are
516 kindly thanked for sharing Arabidopsis mutants with us, Dr Kai-Uwe Fröhlich for providing Yeast
517 strains, and Dr. Christian Steinberg for giving us *Alternaria brassicicola* strain MIAE01824.

518
519
520

References

1. Dissmeyer N, Coux O, Rodriguez MS, Barrio R. PROTEOSTASIS: A European Network to Break Barriers and Integrate Science on Protein Homeostasis. Trends Biochem Sci. 2019 May 1;44(5):383–7.

2. Kumari D, Brodsky JL. The Targeting of Native Proteins to the Endoplasmic Reticulum-Associated Degradation (ERAD) Pathway: An Expanding Repertoire of Regulated Substrates. Biomolecules. 2021;11(8).

3. Deng Y, Humbert S, Liu JX, Srivastava R, Rothstein SJ, Howell SH. Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in Arabidopsis. Proc Natl Acad Sci. 2011 Apr 26;108(17):7247–52.

4. Nagashima Y, Mishiba K ichiro, Suzuki E, Shimada Y, Iwata Y, Koizumi N. Arabidopsis IRE1 catalyses unconventional splicing of bZIP60 mRNA to produce the active transcription factor. Sci Rep. 2011 Jul 1;1(1):29.

5. Mishiba K ichiro, Nagashima Y, Suzuki E, Hayashi N, Ogata Y, Shimada Y, et al. Defects in IRE1 enhance cell death and fail to degrade mRNAs encoding secretory pathway proteins in the Arabidopsis unfolded protein response. Proc Natl Acad Sci. 2013 Apr 2;110(14):5713–8.

6. Pastor-Cantizano N, Ko DK, Angelos E, Pu Y, Brandizzi F. Functional Diversification of ER Stress Responses in Arabidopsis. Trends Biochem Sci. 2020 Feb;45(2):123–36.

7. Che P, Bussell JD, Zhou W, Estavillo GM, Pogson BJ, Smith SM. Signaling from the Endoplasmic Reticulum Activates Brassinosteroid Signaling and Promotes Acclimation to Stress in Arabidopsis. Sci Signal. 2010 Sep 28;3(141):ra69–ra69.

8. Liu JX, Srivastava R, Che P, Howell SH. An Endoplasmic Reticulum Stress Response in Arabidopsis Is Mediated by Proteolytic Processing and Nuclear Relocation of a Membrane-Associated Transcription Factor, bZIP28. Plant Cell. 2007 Dec 1;19(12):4111–9.

9. Strasser R. Protein Quality Control in the Endoplasmic Reticulum of Plants. Annu Rev Plant Biol. 2018 Apr 29;69(1):147–72.

10. Yang ZT, Wang MJ, Sun L, Lu SJ, Bi DL, Sun L, et al. The Membrane-Associated Transcription Factor NAC089 Controls ER-Stress-Induced Programmed Cell Death in Plants. Vol. 10, PLOS GENETICS. 1160 BATTERY STREET, STE 100, SAN FRANCISCO, CA 94111 USA: PUBLIC LIBRARY SCIENCE; 2014.

11. Cai YM, Yu J, Ge Y, Mironov A, Gallois P. Two proteases with caspase-3-like activity, cathepsin B and proteasome, antagonistically control ER-stress-induced programmed cell death in Arabidopsis. New Phytol. 2018 May 1;218(3):1143–55.

12. Reyes-Impellizzeri S, Moreno AA. The Endoplasmic Reticulum Role in the Plant Response to Abiotic Stress. Front Plant Sci [Internet]. 2021;12. Available from: https://www.frontiersin.org/articles/10.3389/fpls.2021.755447

13. Verchot J, Pajerowska-Mukhtar KM. UPR signaling at the nexus of plant viral, bacterial, and
fungal defenses. Curr Opin Virol. 2021 Apr 1;47:9–17.

14. Xu Z, Song N, Ma L, Wu J. IRE1-bZIP60 Pathway Is Required for Nicotiana attenuata Resistance to
Fungal Pathogen Alternaria alternata. Front Plant Sci [Internet]. 2019;10. Available from:
https://www.frontiersin.org/articles/10.3389/fpls.2019.00263

15. Samperna S, Boari A, Vurro M, Salzano AM, Reveglia P, Evidente A, et al. Arabidopsis Defense
against the Pathogenic Fungus Drechslera gigantea Is Dependent on the Integrity of the Unfolded
Protein Response. Biomolecules [Internet]. 2021;11(2). Available from:
https://www.mdpi.com/2218-273X/11/2/240

16. Bürger M, Chory J. Stressed Out About Hormones: How Plants Orchestrate Immunity. Cell Host
Microbe. 2019 Aug 14;26(2):163–72.

17. Macioszek VK, Jęcz T, Ciereszko I, Kononowicz AK. Jasmonic Acid as a Mediator in Plant Response
to Necrotrophic Fungi. Cells. 2023;12(7).

18. Thomma BPHJ, Eggermont K, Penninckx IAMA, Mauch-Mani B, Vogelsang R, Cammue BPA, et al.
Separate jasmonate-dependent and salicylate-dependent defense-response pathways in
Arabidopsis are essential for resistance to distinct microbial pathogens. Proc Natl Acad Sci. 1998
Dec 8;95(25):15107–11.

19. Ferrari S, Plotnikova JM, De Lorenzo G, Ausubel FM. Arabidopsis local resistance to Botrytis
cinerea involves salicylic acid and camalexin and requires EDS4 and PAD2, but not SID2, EDS5 or
PAD4. Plant J. 2003 Jul 1;35(2):193–205.

20. Brouwer SM, Odilbekov F, Burra DD, Lenman M, Hedley PE, Grenville-Briggs L, et al. Intact
salicylic acid signalling is required for potato defence against the necrotrophic fungus Alternaria
solani. Plant Mol Biol. 2020 Sep 1;104(1):1–19.

21. Zander M, La Camera S, Lamotte O, Métraux JP, Gatz C. Arabidopsis thaliana class-II TGA
transcription factors are essential activators of jasmonic acid/ethylene-induced defense
responses. Plant J. 2010 Jan 1;61(2):200–10.

22. Nguyen NH, Trotel-Aziz P, Villaume S, Rabenoelina F, Clément C, Baillieul F, et al. Priming of
camalexin accumulation in induced systemic resistance by beneficial bacteria against Botrytis
cinerea and Pseudomonas syringae pv. tomato DC3000. J Exp Bot. 2022 Jun 2;73(11):3743–57.

23. Glawischnig E. Camalexin. Phytochemistry. 2007 Feb 1;68(4):401–6.

24. Chassot C, Buchala A, Schoonbeek H jan, Métraux JP, Lamotte O. Wounding of Arabidopsis leaves
causes a powerful but transient protection against Botrytis infection. Plant J. 2008 Aug
1;55(4):555–67.

25. Sellam A, Iacomi-Vasilescu B, Hudhomme P, Simoneau P. In vitro antifungal activity of brassinin,
camalexin and two isothiocyanates against the crucifer pathogens Alternaria brassicicola and
Alternaria brassicae. Plant Pathol. 2007 Apr 1;56(2):296–301.

26. Thomma BPHJ, Nelissen I, Eggermont K, Broekaert WF. Deficiency in phytoalexin production
causes enhanced susceptibility of Arabidopsis thaliana to the fungus Alternaria brassicicola. Plant
J. 1999 Jul 1;19(2):163–71.

27. Bi K, Liang Y, Mengiste T, Sharon A. Killing softly: a roadmap of Botrytis cinerea pathogenicity.
Trends Plant Sci. 2023 Feb;28(2):211–22.

28. Govrin EM, Levine A. The hypersensitive response facilitates plant infection by the necrotrophic
pathogen Botrytis cinerea. Curr Biol. 2000 Jun 1;10(13):751–7.

29. Zimmerli L, Métraux JP, Mauch-Mani B. beta-Aminobutyric acid-induced protection of
Arabidopsis against the necrotrophic fungus Botrytis cinerea. Plant Physiol. 2001 Jun;126(2):517–
23.

30. La Camera S, Geoffroy P, Samaha H, Ndiaye A, Rahim G, Legrand M, et al. A pathogen-inducible
patatin-like lipid acyl hydrolase facilitates fungal and bacterial host colonization in Arabidopsis.
Plant J. 2005 Dec 1;44(5):810–25.

31. Nekrasov V, Li J, Batoux M, Roux M, Chu ZH, Lacombe S, et al. Control of the pattern-recognition
receptor EFR by an ER protein complex in plant immunity. EMBO J. 2009 Nov 4;28(21):3428–38.

32. Braun S, Matuschewski K, Rape M, Thoms S, Jentsch S. Role of the ubiquitin-selective
CDC48UFD1/NPL4 chaperone (segregase) in ERAD of OLE1 and other substrates. EMBO J. 2002
Feb 15;21(4):615–21.

33. Jarosch E, Taxis C, Volkwein C, Bordallo J, Finley D, Wolf DH, et al. Protein dislocation from the ER
requires polyubiquitination and the AAA-ATPase Cdc48. Nat Cell Biol. 2002 Feb 1;4(2):134–9.

34. Yamamoto M, Kawanabe M, Hayashi Y, Endo T, Nishikawa S ichi. A vacuolar carboxypeptidase
mutant of Arabidopsis thaliana is degraded by the ERAD pathway independently of its N-glycan.
Biochem Biophys Res Commun. 2010 Mar 12;393(3):384–9.

35. Bègue H, Jeandroz S, Blanchard C, Wendehenne D, Rosnoblet C. Structure and functions of the
chaperone-like p97/CDC48 in plants. Biochim Biophys Acta BBA - Gen Subj. 2017 Jan 1;1861(1,
Part A):3053–60.

36. Copeland C, Woloshen V, Huang Y, Li X. AtCDC48A is involved in the turnover of an NLR immune
receptor. Plant J. 2016 Oct 1;88(2):294–305.

37. Müller J, Piffanelli P, Devoto A, Miklis M, Elliott C, Ortmann B, et al. Conserved ERAD-Like Quality
Control of a Plant Polytopic Membrane Protein. Plant Cell. 2005 Jan 1;17(1):149–63.

38. Feiler HS, Desprez T, Santoni V, Kronenberger J, Caboche M, Traas J. The higher plant Arabidopsis
thaliana encodes a functional CDC48 homologue which is highly expressed in dividing and
expanding cells. EMBO J. 1995 Nov 1;14(22):5626–37.

39. Fröhlich KU, Fries HW, Rüdiger M, Erdmann R, Botstein D, Mecke D. Yeast cell cycle protein
CDC48p shows full-length homology to the mammalian protein VCP and is a member of a protein
family involved in secretion, peroxisome formation, and gene expression. J Cell Biol. 1991 Aug
1;114(3):443–53.

40. Niehl A, Amari K, Gereige D, Brandner K, Mély Y, Heinlein M. Control of Tobacco mosaic virus
Movement Protein Fate by CELL-DIVISION-CYCLE Protein48. Plant Physiol. 2012 Dec
1;160(4):2093–108.

41. La Camera S, L'Haridon F, Astier J, Zander M, Abou-Mansour E, Page G, et al. The glutaredoxin
ATGRXS13 is required to facilitate Botrytis cinerea infection of Arabidopsis thaliana plants. Plant
J. 2011 Nov 1;68(3):507–19.

42. Pré M, Atallah M, Champion A, De Vos M, Pieterse CMJ, Memelink J. The AP2/ERF Domain
Transcription Factor ORA59 Integrates Jasmonic Acid and Ethylene Signals in Plant Defense. Plant
Physiol. 2008 Jul 1;147(3):1347–57.

43. Gladman NP, Marshall RS, Lee KH, Vierstra RD. The Proteasome Stress Regulon Is Controlled by a
Pair of NAC Transcription Factors in Arabidopsis. Plant Cell. 2016 Jun 1;28(6):1279–96.

44. Jing M, Wang Y. Plant Pathogens Utilize Effectors to Hijack the Host Endoplasmic Reticulum as
Part of Their Infection Strategy. Engineering. 2020 May 1;6(5):500–4.

45. Tateda C, Ozaki R, Onodera Y, Takahashi Y, Yamaguchi K, Berberich T, et al. NtbZIP60, an
endoplasmic reticulum-localized transcription factor, plays a role in the defense response against
bacterial pathogens in Nicotiana tabacum. J Plant Res. 2008 Nov 1;121(6):603–11.

46. Moreno AA, Mukhtar MS, Blanco F, Boatwright JL, Moreno I, Jordan MR, et al. IRE1/bZIP60-
Mediated Unfolded Protein Response Plays Distinct Roles in Plant Immunity and Abiotic Stress
Responses. PLOS ONE. 2012 Feb 16;7(2):e31944.

47. Li C, Zhang T, Liu Y, Li Z, Wang Y, Fu S, et al. Rice stripe virus activates the bZIP17/28 branch of
the unfolded protein response signalling pathway to promote viral infection. Mol Plant Pathol.
2022 Mar 1;23(3):447–58.

48. Gayral M, Arias Gaguancela O, Vasquez E, Herath V, Flores FJ, Dickman MB, et al. Multiple ER-to-
nucleus stress signaling pathways are activated during Plantago asiatica mosaic virus and Turnip
mosaic virus infection in Arabidopsis thaliana. Plant J. 2020 Aug 1;103(3):1233–45.

49. Ao K, Tong M, Li L, Lüdke D, Lipka V, Chen S, et al. SCFSNIPER7 controls protein turnover of
unfoldase CDC48A to promote plant immunity. New Phytol. 2021 Mar 1;229(5):2795–811.

50. Czékus Z, Csíkos O, Ördög A, Tari I, Poór P. Effects of Jasmonic Acid in ER Stress and Unfolded
Protein Response in Tomato Plants. Biomolecules. 2020;10(7).

51. Kim SG, Lee S, Ryu J, Park CM. Probing protein structural requirements for activation of
membrane-bound NAC transcription factors in Arabidopsis and rice. Plant Sci. 2010 Mar
1;178(3):239–44.

52. Lee S, Lee HJ, Huh SU, Paek KH, Ha JH, Park CM. The Arabidopsis NAC transcription factor NTL4
participates in a positive feedback loop that induces programmed cell death under heat stress
conditions. Plant Sci. 2014 Oct 1;227:76–83.

53. Lee S, Seo PJ, Lee HJ, Park CM. A NAC transcription factor NTL4 promotes reactive oxygen species
production during drought-induced leaf senescence in Arabidopsis. Plant J. 2012 Jun
1;70(5):831–44.

54. Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR. Genome-Wide Identification and
Testing of Superior Reference Genes for Transcript Normalization in Arabidopsis. Plant Physiol.
2005 Sep 1;139(1):5–17.

55. Ganger MT, Dietz GD, Ewing SJ. A common base method for analysis of qPCR data and the
application of simple blocking in qPCR experiments. BMC Bioinformatics. 2017 Dec 1;18(1):534.

56. Karimi M, Inzé D, Depicker A. GATEWAY™ vectors for Agrobacterium-mediated plant
transformation. Trends Plant Sci. 2002 May 1;7(5):193–5.

57. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium -mediated transformation
of Arabidopsis thaliana. Plant J. 1998 Dec 1;16(6):735–43.

58. Loqué D, Lalonde S, Looger LL, von Wirén N, Frommer WB. A cytosolic trans-activation domain
essential for ammonium uptake. Nature. 2007 Mar 1;446(7132):195–8.

680
681
682
683   **Figure legends**
684
685
686
Figure 1: Mutants deficient for IRE1-bZIP60 branch of UPR show enhanced necrotic symptoms
induced by *B. cinerea* and *A. brassicicola*
(A) Lesion diameters were measured 3 days post-inoculation with *B. cinerea* and (B) 5 days post-
inoculation with *A. brassicicola*. Experiments were repeated seven to eight times for *B. cinerea*
infection and four times for *A. brassicicola* infection depending on the genotype tested. Seven plants
per genotype were infected in each experiment. Violin plots display the distribution of lesion
diameters for all experiments and diamonds indicate the median. Stars indicate significant
differences compared to the wild type and were assessed using Kruskal-Wallis' method followed by
Dunnett's post-hoc test (*** $p < 0.005$; ns : no significant difference).
696
Figure 2: Expression kinetic of UPR genes in response to *B. cinerea*.
Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B.*
cinerea spores (black bars). For each experiment, four leaves of three plants were collected at 16, 24,
48 and 72 hours post-inoculation. Level of transcripts coding IRE1A, IRE1B, bZIP17, bZIP28 and bZIP60
(unspliced or spliced forms) were quantified by RT-qPCR and normalized to those of two reference
genes *AT4G26410* and *AT3G01150* (54). Experiments were repeated five times and mean expression
for all experiments are presented. Significant differences between non-infected and infected plants
were determined by a one-way ANOVA followed by a Tukey HSD Test (*** $p<0.005$; ** $p<0.01$; *
$p<0.05$; • $p<0.1$).
706
707
Figure 3: Defect in ER-QC machinery enhances *B. cinerea*-induced symptoms.
(A) Expression of genes coding ER-QC machinery over *B. cinerea* infection. Plants were sprayed with
¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars).
For each experiment, four leaves of three plants were collected at 16, 24, 48 and 72 hours post-
inoculation and used for total RNAs extraction. Transcript levels were quantified by RT-qPCR and
normalized to those of two reference genes, *AT4G26410* and *AT3G01150* (54). The represented fold
change is the mean of six independent experiments. Significant differences between non-infected
and infected plants were determined by one-way ANOVA followed by a Tukey HSD post-hoc test (***
$p<0.005$; ** $p<0.01$; * $p<0.05$; • $p<0.1$).

(B) Disease phenotype of *erdj3b-1* and *sdf2-2* mutants infected by *B. cinerea*. Lesion diameters
observed in wild-type (Col-0) and mutants were measured 3 days after *B. cinerea* infection.
Experiments were repeated four times. Seven plants per genotype were infected in each experiment.
Violin plots display the distribution of lesion diameters for all experiments and diamonds indicate the
median. Stars indicate significant differences compared to the wild type and were using Kruskal-
Wallis' method followed by Dunnett's post-hoc test (*** p < 0.005).

Figure 4: Functional complementation of yeast *cdc48* mutants.
The yeast *cdc48* mutant strains which are cold-sensitive (strain DBY2030) or heat-sensitive (strain
KFY189) were transformed with pDRF1-GW vector containing or not the cDNA coding Arabidopsis
CDC48A-E proteins. Yeast strains were then streaked out on YPD agar plates and grown at 30°C (A, C)
or 16°C for DBY2030 (B) or 37°C for KFY189 (D). Experiments were repeated three times with similar
results.

Figure 5: CDC48 proteins are involved in disease susceptibility caused by *B. cinerea*
(A) Expression of *CDC48A, CDC48B and CDC48C genes* in response to *B. cinerea* infection. Plants were
sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores
(black bars). For each experiment, four leaves of three plants were collected at 16, 24, 48 and 72
hours post-inoculation and used for total RNAs extraction. Transcript levels were quantified by RT-
qPCR and normalized to those of two reference genes, *AT4G26410* and *AT3G01150* (54). The
represented fold change is the mean of 6 independent experiments. Significant differences between
non-infected and infected plants were assessed by one-way ANOVA followed by a Tukey HSD post-
hoc test (*** p<0.005; * p<0.05; • p<0.1).
(B) Disease phenotypes of *cdc48* mutants, *CDC48B-WT* overexpressors and *CDC48B-QQ* dominant
negative overexpressors. Lesion diameters observed in wild-type (Col-0) and transgenics were
measured 3 days after *B. cinerea* infection. Experiments were repeated seven to nine times. Seven
plants per genotype were infected in each experiment. The stars indicate significant differences
compared to the wild type and were assessed using Kruskal-Wallis' method followed by Dunnett's
post-hoc test (*** p<0.005; * p<0.05; • p<0.1).

Figure 6: Defect in NAC053 and NAC078 transcription factors reduces *B. cinerea*-induced
symptoms
(A-B) Expression of *NAC053* and *NAC078* genes in response to *B. cinerea* infection in WT, *ire1a/ire1b*
and *bzip60* mutants. Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB
solution containing *B. cinerea* spores (black bars). For each experiment, four leaves of three plants
were collected at 24 h (A) or 48 h (B) post infection and used for total RNA extraction. Transcript
levels were quantified by RT-qPCR and normalized to those of two reference genes, *AT4G26410* and
AT3G01150 (54). The represented fold changes are the mean of six independent experiments.
Different letters represent groups which were significantly different from one another as determined
by a one-way ANOVA followed by a multiple comparison with a Fisher's Least Significant Difference
(LSD) test.
(C) Disease phenotype of *nac053* and *nac078* mutants infected with *B. cinerea*. Lesion diameters
observed in wild-type (Col) and mutants were measured 3 days after *B. cinerea* infection.
Experiments were repeated four times. Seven plants per genotype were infected in each experiment.

The stars indicate significant differences compared to the wild type. Significant differences between
a mutant genotype and WT plants were assessed using Kruskal-Wallis' method followed by Dunnett's
post-hoc test (*** p < 0.005; ns : non-significant difference).
768
769
770
Figure 7: schematic representation of the role of UPR, ERQC and ERAD during *Botrytis cinerea*
infection of *Arabidopsis thaliana* plants.
Botrytis cinerea infection induces an ER stress which activates the canonical IRE1-bZIP60 branch of
the UPR and the expression of the genes coding the ERQC machinery (BIP, ERDJ3, SDF2). We
hypothesized the activation of these signalling pathways might control the secretion of defence
proteins which are necessary to establish defence against *B. cinerea*. Our data also indicate that
active bZIP60 suppresses *NAC053* expression, a negative regulator of defence against *B. cinerea*.
However, *NAC053* expression depends on IRE1. Regarding the bZIP17 arm of the UPR or the ERAD
proteins CDC48, we hypothesized that they act as negative regulators of immunity against B. cinerea
or might be the target of *B. cinerea* effectors (purple diamonds) thus facilitating infection process.
781
782
783
784

Figure 1 : Mutants deficient for IRE1-bZIP60 branch of UPR show enhanced necrotic symptoms induced by *B. cinerea* and *A. brassicicola*

(A) Lesion diameters were measured 3 days post-inoculation with *B. cinerea* and (B) 5 days post-inoculation with *A. brassicicola*. Experiments were repeated seven to eight times for *B. cinerea* infection and four times for *A. brassicicola* infection depending on the genotype tested . Seven plants per genotype were infected in each experiment. Violin plots display the distribution of lesion diameters for all experiments and diamonds indicate the median. Stars indicate significant differences compared to the wild type and were assessed using Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** p < 0.005 ; ns : no significant difference).

Figure 2 : Expression kinetic of UPR genes in response to *B. cinerea*.
Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 16, 24, 48 and 72 hours post-inoculation. Level of transcripts coding IRE1A, IRE1B, bZIP17, bZIP28 and bZIP60 (unspliced or spliced forms) were quantified by RT-qPCR and normalized to those of two reference genes *AT4G26410* and *AT3G01150* (Czechowski et al., 2005). Experiments were repeated five times and mean expression for all experiments are presented. Significant differences between non-infected and infected plants were determined by a one-way ANOVA followed by a Tukey HSD Test (*** p<0.005; ** p<0.01; * p<0.05; • p<0.1).

Figure 3 : Defect in ER-QC machinery enhances *B. cinerea*-induced symptoms.
(A) Expression of genes coding ER-QC machinery over *B. cinerea* infection. Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 16, 24, 48 and 72 hours post-inoculation and used for total RNA extraction. Transcript levels were quantified by RT-qPCR and normalized to those of two reference genes, *AT4G26410* and *AT3G01150* (Czechowski et al., 2005). The represented fold change is the mean of six independent experiments. Significant differences between non-infected and infected plants were determined by one-way ANOVA followed by a Tukey HSD post-hoc test (*** p<0.005; ** p<0.01; * p<0.05; • p<0.1).
(B) Disease phenotype of *erdj3b-1* and *sdf2-2* mutants infected by *B. cinerea*. Lesion diameters observed in wild-type (Col-0) and mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated four times. Seven plants per genotype were infected in each experiment. Violin plots display the distribution of lesion diameters for all experiments and diamonds indicate the median. Stars indicate significant differences compared to the wild type and were assessed using Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** p < 0.005).

Figure 4: Functional complementation of yeast *cdc48* mutants.
The yeast *cdc48* mutant strains which are cold-sensitive (strain DBY2030) or heat-sensitive (strain KFY189) were transformed with pDRF1-GW vector containing or not the cDNA coding Arabidopsis CDC48A-E proteins. Yeast strains were then streaked out on YPD agar plates and grown at 30°C (A, C) or 16°C for DBY2030 (B) or 37°C for KFY189 (D). Experiments were repeated three times with similar results.

Figure 5 : CDC48 proteins are involved in disease susceptibility caused by *B. cinerea*

(A) Expression of *CDC48A*, *CDC48B and CDC48C genes* in response to *B. cinerea* infection. Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 16, 24, 48 and 72 hours post-inoculation and used for total RNAs extraction. Transcript levels were quantified by RT-qPCR and normalized to those of two reference genes, *AT4G26410* and *AT3G01150* (Czechowski et al., 2005). The represented fold change is the mean of six independent experiments. Significant differences between non-infected and infected plants were assessed by one-way ANOVA followed by a Tukey HSD post-hoc test (*** $p<0.005$; * $p<0.05$; • $p<0.1$).

(B) Disease phenotypes of *cdc48* mutants, *CDC48B-WT* overexpressors and *CDC48B-QQ* dominant negative overexpressors. Lesion diameters observed in wild-type (Col-0) and transgenics were measured 3 days after *B. cinerea* infection. Experiments were repeated seven to nine times. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type and were assessed using Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** $p<0.005$; * $p<0.05$; • $p<0.1$).

Figure 6 : Defect in NAC053 and NAC078 transcription factors reduces *B. cinerea*-induced symptoms

(A-B) Expression of *NAC053* and *NAC078* genes in response to *B. cinerea* infection in WT, *ire1a/ire1b* and *bzip60* mutants. Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 24 h (A) or 48 h (B) post infection and used for total RNA extraction. Transcript levels were quantified by RT-qPCR and normalized to those of two reference genes, *AT4G26410* and *AT3G01150* (Czechowski et al., 2005). The represented fold changes are the mean of six independent experiments. Different letters represent groups which were significantly different from one another as determined by a one-way ANOVA followed by a multiple comparison with a Fisher's Least Significant Difference (LSD) test.

(C) Disease phenotype of *nac053* and *nac078* mutants infected with *B. cinerea*. Lesion diameters observed in wild-type (Col) and mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated four times. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type. Significant differences between a mutant genotype and WT plants were assesed using Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** $p < 0.005$; ns : non significant difference).

Figure 7 : schematic representation of the role of UPR, ERQC and ERAD during *Botrytis cinerea* infection of *Arabidopsis thaliana* plants.

Botrytis cinerea infection induces an ER stress which activates the canonical IRE1-bZIP60 branch of the UPR and the expression of the genes coding the ERQC machinery (BIP, ERDJ3, SDF2). We hypothesized the activation of these signalling pathways might control the secretion of defence proteins which are necessary to establish defence against *B. cinerea*. Our data also indicate that active bZIP60 suppresses *NAC053* expression, a negative regulator of defence against *B. cinerea*. However, *NAC053* expression depends on IRE1. Regarding the bZIP17 arm of the UPR or the ERAD proteins CDC48, we hypothesized that they act as negative regulators of immunity against B. cinerea or might be the target of *B. cinerea* effectors (purple diamonds) thus facilitating infection process.

Supplemental Figure S1 : Mean lesion diameters observed on UPR-deficient mutants infected by *B. cinerea* or *A. brassicicola*.

Lesion diameters were measured 3 days after *B. cinerea* infection (A) and 5 days after *A. brassicicola* infection (B). Experiments were repeated seven to eight times for *B. cinerea* infection depending on the genotype tested and four times for *A. brassicicola* infection. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type which were identified using a Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** p < 0.005 ; ns : non significant difference).

Supplemental Figure S2: Expression kinetic of the classical Botrytis-induced defense genes *PLP2*, *PR1*, *PAD3* and *PDF1.2* in our experimental conditions.

Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 16, 24, 48 and 72 hours post-inoculation. Transcript levels were quantified by RT-qPCR and normalized to the plant reference genes *AT4G26410* and *AT3G01150* transcript levels (Czechowski et al., 2005). The represented fold change is the mean of five independent experiments. Significant differences between non-infected and infected plants were determined by a one-way ANOVA followed by a Tukey HSD post-hoc test (*** $p<0.005$; ** $p<0.01$; * $p<0.05$; • $p<0.1$).

Supplemental Figure S3: Mean lesion diameters observed on ERQC mutants infected by *B. cinerea*.
Lesion diameters observed on wild-type Col plants and ERQC mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated three to four times. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type and were identified using Kruskal-Wallis' method followed by Dunnett's post-hoc test (*** p < 0.005).

Supplemental Figure S4: Expression of *CDC48* in *cdc48* mutants

A, B, C. Expression of CDC48 mRNAs in *cdc48b* and *cdc48c* mutants and in CDC48B overexpressing lines. Four leaves of three plants were collected and *CDC48B* or *CDC48C* transcript levels were quantified by RTqPCR in *cdc48b3* and *cdc48b4* (A) or in *cdc48c2* and *cdc48c3* (B) mutants, respectively. *CDC48B* transcript level was quantified in transgenic lines overexpressing CDC48B-WT or CDC48B^{E308QE581Q} (CDC48-QQ). All transcript level were normalized to the plant reference genes *AT4G26410* and *AT3G01150* transcript levels (Czechowski et al., 2005). The represented fold change is the mean of five independent experiment. Significant differences between non-infected and infected plants were determined by a one-way ANOVA followed by a Tukey HSD post-hoc test (*** $p < 0.005$; * $p < 0.05$).

Supplemental Figure S5: Mean lesion diameters observed on CDC48 mutants and plants expressing a WT (CDC48B-WT) or an inactive isoforms of CDC48B (CDC48B-QQ) infected by *B. cinerea*.
Lesion diameters observed on wild-type Col plants and mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated seven to nine times. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type which were identified using Kruskal-Wallis' method followed by Dunnett's post-hoc test(*** p<0.005; * p<0.05; • p<0.1).

Supplemental Figure S6 : Accumulation of mRNA coding classical defense genes in response to *B. cinerea* infection in WT, *ire1a/ire1b* and *bzip60* mutants.

Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 24 h (A) or 48 h (B) post-treatment. Transcript levels were quantified by RT-qPCR and normalized to the plant reference genes *AT4G26410* and *AT3G01150* transcript levels (Czechowski et al., 2005). The represented fold change is the mean of six independent experiments. Different letters represent groups which were significantly different from one another as determined by a one-way ANOVA followed by a multiple comparison with a Fisher's Least Significant Difference (LSD) post-hoc test.

Supplemental Figure S7 : Accumulation of mRNA coding cathepsin and NAC089 transcription factor in response to *B. cinerea* infection in WT, *ire1a/ire1b* and *bzip60* mutants.

Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 24 h (A) or 48 h (B) post-treatment. Transcript levels were quantified by RT-qPCR and normalized to the plant reference genes *AT4G26410* and *AT3G01150* transcript levels (Czechowski et al., 2005). The represented fold change is the mean of six independent experiments. Different letters represent groups which were significantly different from one another as determined by a one-way ANOVA followed by a multiple comparison with a Fisher's Least Significant Difference (LSD) post-hoc test.

Supplemental Figure S8: Accumulation of mRNA coding CDC48 in response to *B. cinerea* infection in WT, *ire1a/ire1b* and *bzip60* mutants.

Plants were sprayed with ¼ PDB solution (Mock, white bars) or with ¼ PDB solution containing *B. cinerea* spores (black bars). For each experiments, four leaves of three plants were collected at 24 h (A) or 48 h (B) post-treatment. Transcript levels were quantified by RT-qPCR and normalized to the plant reference genes *AT4G26410* and *AT3G01150* transcript levels (Czechowski et al., 2005). The represented fold change is the mean of six independent experiments. Different letters represent groups which were significantly different from one another as determined by a one-way ANOVA followed by a multiple comparison with a Fisher's Least Significant Difference (LSD) post-hoc test.

Supplemental Figure S9: Mean lesion diameters observed on *nac053* and *nac078* mutants infected by *B. cinerea*.

Lesion diameters observed on wild-type Col plants and mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated four times. Seven plants per genotype were infected in each experiment. The stars indicate significant differences compared to the wild type which were identified using Kruskal-Wallis' method followed by Dunnett's post-hoc test(*** $p<0.005$).

Supplemental Figure S10: Mean lesion diameters observed on *bzip60.1-bzip28.2 double mutant* infected by *B. cinerea*.

Lesion diameters observed on wild-type Col plants and mutants were measured 3 days after *B. cinerea* infection. Experiments were repeated six times. Seven plants per genotype were infected in each experiment. Significant differences compared to the wild type were identified using Kruskal-Wallis' method followed by Dunnett's post-hoc test (ns: not significant).

Supplemental table S1 : list of mutants used in this study

mutant name	gene locus	mutant reference	references	origin	
Atbzip17-1	At2g40950	SALK_104326	Liu et al., 2007a	Pr. Steven H Howell	https://doi.org/10.1111/j.1365-313X.2007.03195.x
Atbzip28-2	At3g10800	SALK_132285C	Liu et al., 2007b	Pr. Steven H Howell	https://doi.org/10.1105/tpc.106.050021
Atbzip60-3	At1g42990	GABI_326A12	Bao et al., 2012	Pr. Steven H Howell	https://doi.org/10.1080/15548627.2018.1462426
Atire1a-2	At2g17520	SALK_018112	Deng et al., 2011	Pr. Steven H Howell	https://doi.org/10.1073/pnas.1102117108
Atire1b-4	At5g24360	SAIL_238_F07	Deng et al., 2011	Pr. Steven H Howell	https://doi.org/10.1073/pnas.1102117108
Atire1a-2/Atire1b-4		SALK_018112 x SAIL_238_F07	Moreno et al., 2012	Pr. Steven H Howell	https://doi.org/10.1371/journal.pone.0031944
Aterdj3b-1	At3g62600	SALK_113364	Nekrasov et al, 2009	Pr. Cyril Zipfel	https://doi.org/10.1038/emboj.2009.262
Atsdf2.2	At2g25110	SALK_141321	Nekrasov et al, 2009	Pr. Cyril Zipfel	https://doi.org/10.1038/emboj.2009.262
Atcdc48a-4/muse8	At3g09840	EMS mutant	Copeland et al., 2016	Pr. Xin Li	https://doi.org/10.1111/tpj.13251
Atcdc48b-3	At3g53230	Gabi_104F08	our study	NASC	
Atcdc48b-4	At3g53230	Gabi_485G04	our study	NASC	
Atcdc48c-2	At5g03340	SALK_102955C	our study	NASC	
Atcdc48c-3	At5g03340	SALK_123409	our study	NASC	
Atnac53-1	At3g10500	SALK_009578C	Gladman et al., 2016	NASC	https://doi.org/10.1105/tpc.15.01022
Atnac78-1	At5g04410	SALK_025098	Gladman et al., 2016	NASC	https://doi.org/10.1105/tpc.15.01022

NASC : Nottingham Arabidopsis Stock Centre (https://arabidopsis.info/)

Supplemental table S2 : Primers used for mutant genotyping

mutant name	line	LB primer name	LB primer sequence	gene specific primer 1	gene specific primer 2
Atcdc48b1	GABI_104F08	GABI-LB-o8474	ATAATAACGCTGCGGACATCTACATTTT	TTCTCATCCCTCCTCTAACTAGGA	TAAATATGAACACCGAGCTCCTCA
Atcdc48b2	GABI_485G04	GABI-LB-o8474	ATAATAACGCTGCGGACATCTACATTTT	TTCTCATCCCTCCTCTAACTAGGA	TAAATATGAACACCGAGCTCCTCA
Atcdc48c1	SALK_123409	Lbe	GGAACAACACTCAACCCTATCTCG	GGTTGCTCTCACTCTCACCAG	TATGTCAAACGAACCGGAATC
Atcdc48c2	SALK_102955C	Lbe	GGAACAACACTCAACCCTATCTCG	GGTTGCTCTCACTCTCACCAG	TATGTCAAACGAACCGGAATC
Atnac053-1	SALK_009578C	Lbe	GGAACAACACTCAACCCTATCTCG	TCAAGAAGCAAACATGTGGTG	GATGACGACGCTTCTTTTCAG
Atnac078-1	SALK_025098	Lbe	GGAACAACACTCAACCCTATCTCG	TCTTTCGCATTTGCGATATTC	TTCAAGTTCTGGTTTTCACCG
Atbzip17-1	SALK_104326	Lbe	GGAACAACACTCAACCCTATCTCG	AAGGAGCAGCCTCCTCCAC	GCCTTTCTCGAAGCCACGG

Supplemental table S3 : RT-qPCR primers used in this work

name	locus		RT-qPCR primers	ref	
AtCDC48A	At3g09840	forward	AAACTCGAATTCTCTACTTTTtCDC48 A	this work	
		reverse	TTTTTCCATTTGTAATGAACGACC		
AtCDC48B	At3g53230	forward	GGTTGTGGATGAAGCCATTA/CDC48 B	this work	
		reverse	CGGTTCCCATACTTGACATCAG		
AtCDC48C	At5g03340	forward	GGCGATACAATTCTCATCAAC CDC48 C	this work	
		reverse	GACGGGGCATTCTTCTCAGC		
reference gene	*At4g26410*	forward	GAGCTGAAGTGGCTTCCATGAC	Czechowski *et al.* , 2005	https://doi.org/10.1104/pp.105.063743
		reverse	GGTCCGACATACCCATGATCC		
reference gene	*At3g01150*	forward	GATCTGAATGTTAAGGCTTTTAGCG	Czechowski *et al.* , 2005	https://doi.org/10.1104/pp.105.063743
		reverse	GGCTTAGATCAGGAAGTGTATAGTCTCTG		
AtIRE1A	AT2G17520	forward	GCTTCAGACCTCATATCCCG	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
		reverse	AGCATCACGAAGGAAAGACAG		
AtIRE1B	AT5G24360	forward	GGTGGGATGAGAAACTGGATA	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
		reverse	AGTTTGTTCCGTATGACCCG		
AtBzip60	AT1G42990	forward	GGAGACGATGATGCTGTGGCT	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
AtBzip60 unspliced Rev		reverse	CAGGGATTCCAACAAGAGCACAG		
AtBzip60 spliced Rev		reverse	CAGGGAACCCAACAGCAGACT		
AtbZIP28	AT3G10800	forward	GCCAGTGATCCTCTCTTTGC	Qiang *et al.* , 2012	https://doi.org/10.1105/tpc.111.093260
		reverse	CAGAAGACAGTGCACCAGGA		
AtbZIP17	AT2G40950	forward	ACAGGAGATCGGGAGGAGGAT	Qiang *et al.* , 2012	https://doi.org/10.1105/tpc.111.093260
		reverse	GCTCCTCGACGTAATGCTTC		
AtBIP1	AT5G28540	forward	GCTCGTTTGGAGCTAACAGTACC	this work	
		reverse	GAGGACAACGCAAATAAACATCCG		
AtBIP2	AT5G42020	forward	TTGCAGAGGAAGACAAGAAGGTG	this work	
		reverse	CACGTATGTCTCCAGGGCATTC		
AtBIP3	AT1G09080	forward	CACGGTTCCAGCGTATTTCAAT	Liu *et al.* , 2007b	https://doi.org/10.1105/tpc.106.050021
		reverse	ATAAGCTATGGCAGCACCCGTT		
AtSDF2	AT2G25110	forward	TCAAGAGTGGAGCAACCATTAG	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
		reverse	CCAAAGCAGCTAACCTCTAAGT		
AtERDJ3A	AT3G08970	forward	GGTAGCTCATCGAATGCTGAA	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
		reverse	GGTCCACAACGTCCTTCTTATAG		
AtERDJ3B	AT3G62600	forward	CAAATACGAACGGGAGGGATAC	Afrin *et al.* , 2020	https://doi.org/10.1038/s41598-020-76114-1
		reverse	GGTTCGCCGTCTTCATAGAAA		
AtPLP2	AT2G26560	forward	GTAGCTGGTTGGGGACTATTGA	La Camera *et al.* , 2005	https://doi.org/10.1111/j.1365-313X.2005.02578.x
		reverse	CGGTAGCGATATCAACAGAAGC		
AtPR1	AT2G14610	forward	AAGGGTTCACAACCAGGCAC	Anderson *et al.* , 2004	https://doi.org/10.1105/tpc.104.025833
		reverse	CACTGCATGGGACCTACGC		
AtPDF1.2a	AT5G44420	forward	TTTGCTGCTTTCGACGCAC	Anderson *et al.* , 2004	https://doi.org/10.1105/tpc.104.025833
		reverse	CGCAAACCCCTGACCATG		
AtNAC053/AtNTL4	AT3G10500	forward	TGTTGGGTGCTATTCCTGCT	Gladman *et al.* , 2016	https://doi.org/10.1105/tpc.15.01022
		reverse	CTGATGATTGTGCTGCGTGT		
AtNAC078/AtNTL11	AT5G04410	forward	TTGAAGAAAGCTGGTGTGCC	Gladman *et al.* , 2016	https://doi.org/10.1105/tpc.15.01022