Title: Pathogenic CD8 T cell responses are driven by neutrophil-mediated

hypoxia in cutaneous leishmaniasis

Authors: Erin A. Fowler[1], Camila Farias Amorim[2], Klauss Mostacada[1], Allison Yan[1], Laís

Amorim Sacramento[2], Rae A. Stanco[1], Emily D. S. Hales[1], Aditi Varkey[1], Wenjing Zong[3]†,

Gary D. Wu[3], Camila I. de Oliveira[4], Patrick L. Collins[1], Fernanda O. Novais[1]*

Affiliations:

8
[1]Department of Microbial Infection and Immunity, College of Medicine, The Ohio State

University; Columbus, USA

[2]Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania;

Philadelphia, USA

[3]Division of Gastroenterology and Hepatology, Perelman School of Medicine, University of

Pennsylvania; Philadelphia, USA.

[4]Centro de Pesquisas Gonçalo Moniz, Fundação Oswaldo Cruz, Salvador, Brazil

*Corresponding author. Email: Fernanda.Novais@osumc.edu

†Current address: Division of Gastroenterology/Hepatology/Nutrition, Children's Hospital of

Philadelphia; Philadelphia, USA.

One Sentence Summary: CD8 T cell-mediated pathogenesis in cutaneous leishmaniasis is

driven by the recruitment of neutrophils that act as an oxygen sink, thereby eliciting cytotoxic

effector function in response to hypoxia within the inflamed tissue.

Abstract: Cutaneous leishmaniasis caused by *Leishmania* parasites exhibits a wide range of

clinical manifestations. Although parasites influence disease severity, cytolytic CD8 T cell

responses mediate disease. While these responses originate in the lymph node, we find that

expression of the cytolytic effector molecule granzyme B is restricted to lesional CD8 T cells in

Leishmania-infected mice, suggesting that local cues within inflamed skin induce cytolytic

function. Expression of Blimp-1 (*Prdm1*), a transcription factor necessary for cytolytic CD8 T

cell differentiation, is driven by hypoxia within the inflamed skin. Hypoxia is further enhanced

by the recruitment of neutrophils that consume oxygen to produce reactive oxygen species,

ultimately increasing granzyme B expression in CD8 T cells. Importantly, lesions from

cutaneous leishmaniasis patients exhibit hypoxia transcription signatures that correlate with the

presence of neutrophils. Thus, targeting hypoxia-driven signals that support local differentiation

of cytolytic CD8 T cells may improve the prognosis for patients with cutaneous leishmaniasis, as

well as other inflammatory skin diseases where cytolytic CD8 T cells contribute to pathogenesis.

INTRODUCTION

Cutaneous leishmaniasis is caused by *Leishmania* parasites and exhibits a wide range of clinical manifestations, from self-healing lesions to chronic debilitating infections *(1)*. No vaccines exist for leishmaniasis, and the anti-parasitic drugs are often ineffective *(2)*. Although the parasites influence disease severity, immune responses are often the major driver of disease. For example, CD8 T cells in lesions are recruited to target parasite-infected cells but also induce host cell death through granule-mediated cytotoxicity. Rather than killing *Leishmania*, cells release parasites that metastasize to distant cutaneous sites *(3)*. In addition, cell death induced by CD8 T cells activates the NLRP3 inflammasome, IL-1β release, chronic neutrophil recruitment, and enhanced inflammation *(3–6)*. The importance of this pathway is supported by our previous transcriptional analysis of lesions from patients, in which granzyme B (*GZMB*), perforin (*PRF1*), and *IL1B* expression were associated with treatment failure *(7)*. While both murine and human studies underscore the pathogenic consequences of cytotoxicity, the microenvironmental cues and downstream mechanisms that drive cytolytic programs in lesional CD8 T cells remain unknown.

In response to *Leishmania* infection, CD8 T cells are activated and expand in draining lymph nodes (dLNs). The activated cells then travel to inflammatory sites where they can control pathogens, often by killing infected cells. While infections also induce differentiation of effector CD4 T cells in secondary lymphoid organs, CD8 T cells with cytotoxic effector function are frequently absent in LNs *(8–11)*. A prior study implicated PD-1 ligation on dLN CD8 T cells in suppressing their expression of granzyme B, a process that is likely advantageous since it would limit the killing of antigen-presenting cells *(11)*. While such findings suggest that CD8 T cell cytolytic effector function is only triggered once CD8 T cells enter inflammatory sites *(12, 13)*, the local cues that drive this response remain unclear. As the CD8 T cell cytolytic program can

have both positive and negative consequences for the host *(14–33)*, understanding the

mechanisms that drive cytotoxic CD8 T cell programs remains an important goal for unraveling

pathogenesis of numerous diseases such as alopecia, vitiligo, bullous pemphigoid, Steven-

Johnson syndrome and toxic epidermal necrolysis, where cytolytic CD8 T cells contribute to

pathogenesis *(15, 32, 34–37)*.

One potential mechanism relevant to inflamed tissues is hypoxia, which occurs when

oxygen consumption by infiltrating cells exceeds the supply *(38)* and differentially impacts

immune cell types. In *Leishmania* infection, hypoxia exclusively has been studied in

macrophages and dendritic cells *(39, 40)*, where it can either promote or block effector

responses, depending on the parasite species and the cell type *(41–49)*. Here, we find that

hypoxia within *Leishmania*-infected lesions directly promotes cytolytic effector function in

newly recruited CD8 T cells. Mechanistically, we identify a feed-forward loop, initiating with

early recruitment of neutrophils that consume oxygen to produce reactive oxygen species (ROS),

which promotes a persistently hypoxic environment, elevates the number of cytolytic CD8 T

cells, perpetuates chronic neutrophil recruitment, and exacerbates tissue damage. Thus, our

findings indicate that targeting cytolytic CD8 T cells driven by hypoxic microenvironments will

improve the prognosis for patients with cutaneous leishmaniasis and other inflammatory skin

diseases.

RESULTS

CD8 T cells express granzyme B in leishmanial lesions but not in the draining lymph nodes.

Cytotoxic CD8 T cells are pathogenic in cutaneous leishmaniasis *(3, 4, 6, 7, 50–54)*, but their anatomical distribution and mechanisms that regulate their induction remain unclear. In this regard, parasites are found in both dLNs and skin lesions. To determine if CD8 T cell cytotoxic effector function is present in both tissues, we assessed the expression of the cytotoxic marker granzyme B. C57BL/6 mice were infected with *Leishmania* in the ear, and two weeks post-infection, the frequency of granzyme B^{pos} antigen-experienced (CD44^{high}) CD8 T cells was assessed by flow cytometry. While granzyme B-expressing CD8 T cells were abundant in the infected skin, granzyme B expression was nearly absent in dLNs (Fig 1a). To determine if granzyme B^{pos} CD8 T cells were preferentially recruited to the skin, we treated mice with FTY720, a sphingosine 1-phosphate receptor agonist that blocks the egress of T cells from the dLNs. If granzyme B^{pos} CD8 T cells are preferentially recruited to cutaneous lesions, granzyme B^{pos} CD8 T cells should accumulate in the dLNs of FTY720-treated mice. However, there was no difference in the frequency of granzyme B^{pos} CD8 T cells in the dLNs of mice treated with FTY720 or vehicle control (Fig. 1b), suggesting that rather than preferential recruitment, exposure to the lesion microenvironment induces a cytotoxic profile in CD8 T cells. To directly test this, we infected C57BL/6 CD45.1 or CD45.2 congenic mice with *Leishmania*, and purified CD8 T cells from dLNs of CD45.2 donor mice (granzyme B^{neg}) three weeks post-infection. Donor dLN CD8 T cells were then transferred directly into the lesions of recipient CD45.1 infected animals (Fig. 1c - schematic). Notably, we found that granzyme B^{neg} CD8 T cells became granzyme B-expressing CD8 T cells in the lesions (Fig. 1c). In contrast, CD8 T cells that migrated back to dLNs remained granzyme B^{neg} (Fig. 1c). These results demonstrated that the lesion microenvironment triggers cytotoxic programs in CD8 T cells.

To determine the mechanisms by which CD8 T cell cytotoxic programs are induced in

lesions, we performed RNA-seq analysis on purified antigen-experienced CD8 T cells from

dLNs and lesions of mice infected with *Leishmania* for five weeks. Principal component analysis

(PCA) showed that almost half (Principal component 1, PC1 47%) of the differences on the

whole transcriptional profiles were associated with the organ from which the CD8 T cells were

derived (Fig. 1d). Differentially expressed gene (DEG) analysis with thresholds of false

discovery rate (FDR) < 0.05 and fold change > 1.5 between tissues revealed 118 genes

overexpressed in CD8 T cells from lesions compared to dLNs and 26 DEGs overexpressed in

dLNs compared to lesions (Data File S1 and Fig. S1). *Gzmb* was the top DEG in CD8 T cells

from lesions compared to dLNs, FDR=0.002 (Fig. 1e). Several transcription factors play

important roles in effector CD8 T cell biology, including BATF, ID-2, IRF4, Stat4, T-bet, Zeb2,

and Blimp-1 (encoded by *Prdm1*) *(55–57)*, and we found that *Id2*, *Irf4*, and *Prdm1* expressions

were significantly higher in lesions than in dLNs, and *Prdm1* had the most significant fold

change between tissues (>100 higher in lesions) (Fig. 1f). To further investigate the signals

received by CD8 T cells within lesions, we performed Gene Set Enrichment Analysis (GSEA),

which exhibited hypoxia-related signatures, as assessed by several pathway databases (Hallmark,

Biocarta, and Reactome), as well as the Harris et al. hypoxia-specific pathway described in *(58)*,

(Fig. 1g and Data File 2). These data indicate that, after their exit from dLNs, CD8 T cells

recruited to cutaneous leishmaniasis lesions are exposed to hypoxic conditions, inducing the

expression of key cytotoxicity effector mediators, including *Prdm1*.

Cutaneous leishmaniasis lesions are hypoxic and alter CD8 T cell function.

Normal skin is naturally low in oxygen *(59, 60)*, and inflammatory environments are

frequently hypoxic. To evaluate hypoxia in *Leishmania*-infected lesions, we employed

pimonidazole, a 2-nitroimidazole reporter molecule that is reductively activated and forms

covalent bonds with macromolecules, specifically in hypoxic cells. Mice were injected with

pimonidazole two weeks after *Leishmania* infection and one hour before euthanasia.

Pimonidazole staining was assessed in infected and contralateral ears by confocal microscopy.

As expected, we found pimonidazole staining within the epidermis, dermis, and hair follicles of

naïve mice (Fig. 2a and 2b – top images). Within lesions, we observed enlargement of the

epidermis and dermis associated with robust pimonidazole staining (Fig. 2a and 2b – bottom

images). Surprisingly, pimonidazole staining was absent from ulcerated areas. Representative

images in lower (Fig. 2a and Fig. S2a) and higher (Fig. 2b and Fig. S2b) magnification depict the

differences in pimonidazole staining distribution between the naïve and infected tissue.

To compare O_2 levels in lesions and dLNs, we infected mice with *Leishmania,* and two

weeks post-infection, mice received Oxyphor G4, a phosphorescent probe quenched by oxygen

(61). Infected skin exhibited a significant decrease in oxygen compared to dLNs (Fig. 2c). These

data directly demonstrated skin lesions present a hypoxic microenvironment to CD8 T cells upon

their recruitment from dLNs.

We tested whether exposure to hypoxia was sufficient to promote granzyme B expression

in dLN CD8 T cells. We isolated dLN cells from infected mice and stimulated them with anti-

CD3 and anti-CD28 in the presence of Dimethyloxalylglycine (DMOG), a compound that

mimics hypoxia at normal oxygen tension. As shown in Fig. 2d, DMOG induced the expression

of granzyme B in CD8 T cells, further demonstrating that hypoxia promotes upregulation of their

cytolytic program. Notably, DMOG treatment also increased *Prdm1* mRNA expression

compared to vehicle-treated CD8 T cells (Fig. 2e). Collectively, these data demonstrate that the

hypoxic state of lesions stimulates differentiation of recruited CD8 T cells into cytotoxic

effectors.

Blimp-1 expression is restricted to granzyme B expressing CD8 T cells.

Blimp-1 deficient CD8 T cells produce less granzyme B *(62)*, so we next sought to test

cause-effect relationships between hypoxia, Blimp-1, and cytotoxicity. To address this question,

we infected Blimp-1 Yellow Fluorescent Protein (YFP) reporter mice with *Leishmania* and

assessed Blimp-1 expression in CD8 T cells. We found that Blimp-1 was highly expressed in

lesional CD8 T cells but not in CD8 T cells from the dLNs (Fig. 3a). We also found that Blimp-1

expression was significantly higher in CD8 T cells that express granzyme B (Fig. 3b), directly

linking Blimp-1 and cytotoxicity. Finally, to assess if granzyme B expression in lesions resulted

from hypoxic induction of Blimp-1, we used mice expressing a fusion protein bearing an

oxygen-dependent degradation (ODD) domain from human HIF1A (hypoxia-inducible factor 1,

alpha subunit) fused to a tamoxifen-inducible Cre recombinase gene (cre/ERT2) *(63)*. When

expressed under normoxic conditions, the O2CreER fusion protein is degraded rapidly but is

stabilized by hypoxia when combined with tamoxifen injection. These mice were crossed with

Blimp-1$^{flox/flox}$ mice (here called ODD^{cre}Blimp-1$^{flox/flox}$ mice) to generate mice in which Blimp-1

expression is deleted specifically in hypoxic cells upon tamoxifen injection (Fig. 3c). Since the

skin has significantly less oxygen than dLNs, these mice provide lesion-specific deletion of

Blimp-1, thus ensuring the preservation of Blimp-1-dependent development of effector T cells in

dLNs (normoxic) and their recruitment to the inflamed tissue. *Leishmania*-infected

ODD[cre]Blimp-1[flox/flox] mice were treated with tamoxifen or left untreated for one week, and

granzyme B expression was determined. We found that specific deletion of Blimp-1 in hypoxic

cells significantly decreased granzyme B expression in lesional CD8 T cells compared to

untreated mice (Fig. 3d). Taken together, these results directly demonstrate that upregulation of

Blimp-1 by the hypoxic environment leads to increased granzyme B expression and CD8 T cell

cytotoxicity in leishmanial lesions.

Blimp-1 expression is necessary for CD8 T cells to mediate disease.

Based on our data, we predicted that Blimp-1 expression was required for cytotoxic CD8

T cell-mediated disease in leishmaniasis. Since Blimp-1 plays a vital role in CD4 T cell subsets

(64) and regulates IL-10 production *(65)*, total T cell deletion of Blimp-1 would complicate

interpretations. Therefore, we used our well-characterized mouse model of chronic leishmaniasis

in which severe disease develops following *Leishmania* infection of RAG-deficient mice

reconstituted with CD8 T cells *(3, 4, 66, 67)*. RAG-deficient mice infected with *Leishmania* were

reconstituted with WT or CD8 T cells lacking Blimp-1 expression (here called Blimp-1[cKO]) or

control animals receiving no T cells. As described previously *(3)*, RAG-deficient mice

reconstituted with WT CD8 T cells developed severe pathology characterized by granzyme B-

expressing CD8 T cells in lesions, while RAG-deficient mice that received no cells showed no

signs of pathology (Fig. 4a). Importantly, RAG-deficient mice reconstituted with Blimp-1[cKO]

CD8 T cells had minimal disease (Fig. 4a). As expected, there were similar numbers of parasites

(Fig. 4b) and a similar frequency of CD8 T cells (Fig. 4c) in lesions of RAG-deficient mice

reconstituted with WT and Blimp-1[cKO] CD8 T cells. Notably, granzyme B expression was

significantly diminished in cells lacking Blimp-1 (Fig. 4d). Collectively, these data indicate that

Blimp-1 expression is driven by the hypoxic microenvironment of the lesion, triggering

granzyme B expression and the pathogenic state of CD8 T cells.

Neutrophils generate the hypoxic microenvironment of cutaneous leishmaniasis lesions.

Hypoxia occurs in tissues when oxygen supply does not meet demand, including

scenarios characterized by defective tissue vascularization or increased demand by cells present

within a tissue *(38)*. *Leishmania*-infected lesions are highly vascularized *(68)*, so we

hypothesized that oxygen consumption by inflammatory cells recruited to cutaneous

leishmaniasis lesions promoted hypoxia. Neutrophils are the first cells recruited to skin upon

Leishmania infection *(69)*, but their recruitment does not control *L. major* parasites. Indeed, their

chronic presence is associated with worsening disease *(3, 4, 70–74)*. Therefore, we tested

whether neutrophil recruitment contributed to the hypoxic state of inflamed skin. *Leishmania*-

infected mice were injected with pimonidazole 1 hour before euthanasia at multiple time points

after infection: 1) soon after parasite challenge (2 and 5 hours), 2) when lesions started to

develop, and CD8 T cells were present and expressed granzyme B (2 weeks), and 3) when

lesions began to heal (9 weeks). As expected *(69)*, neutrophils were quickly recruited to

Leishmania-infected lesions and decreased in frequency when lesions resolved (Fig. 5a).

Surprisingly, neutrophil (CD11b^{pos}Ly6G^{pos}) staining by pimonidazole was significantly lower

compared to other myeloid cells (CD11b^{pos}Ly6G^{neg}) in the skin at all time points analyzed (Fig.

5b). To confirm these observations, lesions from pimonidazole-injected mice were stained for

Ly6G (clone 1A8), revealing intense neutrophil recruitment in ulcered regions in the epidermis

and clusters of neutrophils within the skin dermis (Fig. 5c and Fig. S3a). Notably, regions

without neutrophils had significantly more pimonidazole staining than those containing

neutrophil clusters. We also observed that pimonidazole staining in regions surrounding

neutrophil clusters was more intense. Quantitative assessment of pixel intensities (Fig. S3b)

revealed a negative correlation between pimonidazole and Ly6G expression (Fig. 5d). Higher

magnification of regions with intense neutrophil recruitment at two different time points showed

a lack of pimonidazole staining in neutrophils (Fig. 5e and Fig. S3c).

These results suggest that areas surrounding neutrophils are relatively hypoxic.

Accordingly, we hypothesized that neutrophils were responsible for promoting the hypoxic state

of the skin by competing for oxygen, thereby controlling CD8 T cell-effector function. To test

this, we depleted mice of neutrophils for the first two weeks of infection using anti-Ly6G (clone

1A8) (Fig. 5f) at the early stage of disease development and assessed granzyme B expression in

CD8 T cells. We found that neutrophil depletion decreased the frequency and the mean

fluorescence intensity (MFI) of granzyme B expressing CD8 T cells compared to isotype

control-treated mice (Fig. 5g). These data suggest that neutrophils consume oxygen in lesions,

altering the phenotype of CD8 T cells.

In this regard, activated neutrophils assemble NADPH oxidase to produce reactive

oxygen species (ROS), which requires oxygen *(75)*. Thus, we tested whether oxygen

consumption to generate ROS resulted in a hypoxic state that impacted CD8 T cell function. For

this purpose, we compared the ability of mice sufficient (WT) or deficient in the $gp91^{phox}$ subunit

of the NADPH oxidase ($Cybb^{-/-}$) to express granzyme B in lesional CD8 T cells upon infection

with *Leishmania*. We found significantly less granzyme B expressing CD8 T cells in Cybb[-/-]

mice than in WT (Fig. 5h), forging a link between NADPH oxidase-dependent ROS production

and the development of pathogenic CD8 T cells in cutaneous leishmaniasis lesions.

The magnitude of hypoxia correlates with the presence of neutrophils in human cutaneous

leishmaniasis lesions.

To determine whether hypoxia is a feature of human disease, we compared

transcriptional signatures in blood and skin from healthy subjects (HS) and cutaneous

leishmaniasis (CL) patients using previously published datasets *(51, 76)*. To estimate levels of

hypoxia-gene expression in clinical samples, we performed single-sample GSEA using the

Harris et al. hypoxia gene signature *(58)*, and referred to this variable as the "hypoxia score"

(Data File S3). While there was no difference in the hypoxia scores between blood from

cutaneous leishmaniasis patients and healthy subjects (Fig. 6a and Data File S4), the cutaneous

leishmaniasis lesions are significantly more hypoxic than healthy skin (Fig. 6b, Data File S3).

Importantly, we observed that the degree of the hypoxia-related gene expression in patients'

lesions was variable. To understand the biological implication of this variability, we performed

unsupervised hierarchical clustering (HC) to classify lesions and healthy skin samples according

to their hypoxic gene expression accessed by Harris et al., Hallmark, Biocarta and Reactome

sources (Fig. 6c). HC analysis revealed two groups of lesion samples: (1) 'baseline' lesions (n =

22) with a hypoxia phenotype comparable to control skin samples, and (2) 'hypoxic' lesions (n =

12) with a high hypoxic phenotype relative to the entire cohort (Fig. 6b and Fig 6c). The

remaining lesions had an intermediate phenotype. PCA revealed that hypoxic lesions are robustly

segregated from baseline counterparts (Permanova, Pr(>F)=0.001) (Fig. 6d). Gene Ontology

performed for genes overexpressed in hypoxic compared to baseline lesions (FDR>0.01 and

FC=1.5 thresholds) revealed statistically significant enrichment for neutrophil chemotaxis and

signaling (Data File S5 and Data File S6). Indeed, estimating cell abundances from this

unstructured bulk RNA-seq data revealed an increase in neutrophil counts in hypoxic compared

to baseline lesions (Fig 6e). The top DEGs between hypoxic compared to baseline lesions

included genes associated with neutrophil chemotaxis and survival *CXCL5*, *CXCL8*, *CXCR1*,

CSF3, and genes encoding for pro-inflammatory cytokines such as *IL1B*, *IL6,* and *OSM* (Fig. 6f

and Data File S5). Therefore, our analysis of patients' samples corroborates our mouse studies,

implicating the presence of neutrophils to the degree of hypoxia in lesions.

DISCUSSION

Cytolytic CD8 T cells cause severe inflammation in various settings, such as alopecia,

malaria, Steven-Johnson syndrome, toxic epidermal necrolysis, cutaneous leishmaniasis, and

many others *(14, 15, 21, 24, 31–34, 77)*. However, the factors that promote cytolysis by CD8 T

cells in tissue microenvironments are still poorly understood. Here, we show that neutrophils

recruited to the skin upon *Leishmania* infection consume oxygen to produce ROS, generating

chronic inflammatory hypoxia. The resultant competition for oxygen changes the aerobic

microenvironment and promotes the expression of Blimp-1 in recruited CD8 T cells, which

results in the expression of the cytolytic molecule granzyme B. Subsequent cell death within the

infected area results in tissue damage and chronic neutrophil recruitment (Fig. 6g). Because there

are numerous skin diseases associated with cytotoxicity, our findings provide evidence that local

inhibition of this pathway should be considered as a strategy to ameliorate these diseases.

In this regard, preferential expression of cytolytic genes by CD8 T cells in peripheral

tissues but not in dLNs is observed in numerous pathogenic conditions. For example, in models

of graft-versus-host disease and graft-versus-leukemia, granzyme B is absent in dLNs but present

in the skin *(9, 11)*. Additionally, memory CD8 T cells acquire cytolytic activity upon entry into

non-lymphoid tissues in vesicular stomatitis virus infections *(10)*. The site-specific function of

CD8 T cells is not limited to granzyme B expression. For example, in mice chronically infected

with *Trypanosoma cruzi*, CD8 T cells in the spleen produced IFN-γ, while CD8 T cells within

the muscle did not *(78)*. Similarly, we found previously that in cutaneous leishmaniasis, CD8 T

cells in dLNs produced IFN-γ while CD8 T cells in the skin lacked IFN-γ production *(67)*. Since

hypoxic dendritic cells produce less IL-12 *(46, 48)*, this could explain the decrease in CD8 T cell

IFN-γ production in lesions *(67)* and suggests that hypoxia may not only promote pathogenic

responses but also prevent protective ones. Our confocal microscopy data suggest that hypoxia

impacts many other cells beyond CD8 T cells; therefore, it is likely that hypoxia has a much

greater role in tissue immunity to *Leishmania*.

There is substantial clinical and experimental evidence that hypoxia contributes to tumor

progression, metastasis, and high mortality *(79)*. While low oxygen levels support myeloid

suppressor cell accumulation *(80)* and tumor cell metastasis *(81)*, hypoxia-inducible factors

(HIF) provide CD8 T cells with greater anti-tumor potential *(82–84)*. HIF increases CD8 T cell

effector function during chronic viral infections and reduces viral titers but leads to higher

mortality *(85)*. These findings agree with the observed increases in perforin expression and

killing ability of CD8 T cells cultured in hypoxia *(84, 86, 87)*. Similar results were observed in

lupus skin disease, where HIF drove granzyme B expression but, in this case, was independent of

hypoxia *(88)*. Hypoxia also alters CD8 T cell metabolism *(86)* and induces tissue residency *(82,*

89) and exhaustion *(90–92)* programs. We found a strong correlation between the hypoxic state

of CD8 T cells and Blimp-1 expression. Consistent with our findings, hypoxia also triggered

Blimp-1 expression in pancreatic tumor cells *(81)*. Since Blimp-1 can instruct tissue residency

and exhaustion *(62, 93–95)*, we speculate that hypoxia-driven Blimp-1 expression may also

initiate these CD8 T cell programs within cutaneous leishmaniasis lesions, which requires further

investigation.

Another immune cell type implicated in the hypoxia-cytotoxicity axis revealed in our

study is neutrophils, which rapidly migrate to infection sites and, in many infections, eliminate

invading microbes. However, it is well-recognized that the impact of neutrophils extends beyond

pathogen killing, as they also shape the function of other cells within tissues by producing

cytokines and chemokines, interacting with dendritic cells and T cells, and consequently

affecting disease outcomes. For example, neutrophils producing IL-10 *(96)* or expressing PD-L1

(97) suppress CD4 T cells. Additionally, neutrophils suppress effector responses by tumor-

specific T cells in lymph nodes *(98)* and within tumors *(99)*. In chronic viral infections,

suppressive neutrophil subsets reduce the antiviral capabilities of CD8 T cells *(100)*. In contrast,

neutrophils facilitate dendritic cell migration to dLNs in contact hypersensitivity and prime

allergen-specific CD8 T cells *(101)*. In cutaneous leishmaniasis, neutrophils recruited to lesions

can have paradoxical functions depending on the parasite species and mouse model. For

example, neutrophils kill *Leishmania* directly by extracellular trap formation or cooperate with

macrophages to activate microbicidal activities *(102, 103)*. In contrast, neutrophils reduce

costimulatory molecule expression in dendritic cells and silently deliver parasites to

macrophages *(69, 104, 105)*. Despite these disparities, it is clear that chronic recruitment of

neutrophils is associated with tissue destruction in cutaneous leishmaniasis *(3, 72, 74, 106)*.

Here, we uncovered a new role for neutrophils in cutaneous leishmaniasis lesions by showing

that their consumption of oxygen promotes a hypoxic environment, resulting in a phenotypic

switch in CD8 T cells.

Cytotoxic CD8 T cells promote increased inflammation by inducing excessive cell death

in lesions and enhancing chronic neutrophil recruitment to the skin *(3, 4)*. In what may be related

findings, neutrophil depletion decreases the hypoxic state of herpes stromal keratitis lesions and

ameliorates disease *(107)*. In a mouse model of colitis, the respiratory burst by Gr1[pos] cells -

encompassing neutrophils and inflammatory monocytes - induced hypoxia in intestinal epithelial

cells, which was critical for disease resolution *(108)*. One limitation of our work is that it does

not address the impact of other innate cells and their ability to deplete lesions of oxygen, which

could also impact CD8 T cell function. However, our work strongly supports oxygen

consumption and neutrophil recruitment as modulators of CD8 T cell granzyme B expression,

which fosters the chronic presence of neutrophils in lesions. While the negative impact of this

cascade of events is clear in cutaneous leishmaniasis, it could be beneficial in other contexts.

To be successful, host-directed therapies for infectious diseases should not augment

immune responses that damage tissues and must not interfere with responses that control the

pathogen *(2)*. Therefore, it is critical to define immune pathways that are pathogenic rather than

protective. In cutaneous leishmaniasis, cytotoxicity is an ideal therapeutic target since it

associates with treatment failure in patients who receive anti-parasitic drugs *(7)* and does not

control the parasite *(3)*. Patients affected by cutaneous leishmaniasis are frequently treated with

anti-parasitic drugs that require intravenous or intramuscular injections every day over the course

of 21 days. Because failure rates are high, many patients must go through two and sometimes

three rounds of treatment. Our data provide evidence that local inhibition of the cytolytic

program can circumvent the use of systemic drugs, which generate considerable side effects, and

topical targets should be considered as a new therapeutic strategy for cutaneous leishmaniasis.

MATERIALS AND METHODS

Ethics statement

371  This study was carried out per the recommendations in the Guide for the Care and Use of

372 Laboratory Animals of the National Institutes of Health. The Institutional Animal Care and Use

373 Committee, The Ohio State University, and The University of Pennsylvania approved the

374 protocol.

376 **Mice**

377  C57BL/6 CD45.2 and CD45.1 mice (6 weeks old) were purchased from Charles River,

and *Prdm1* YFP (here called Blimp-1 reporter), ODD^{cre}/ERT2 *(63)*, Gzmb^{cre}, Blimp-1$^{flox/flox}$, and

RAG$^{-/-}$ (B6.12957-RAG1^{tm1Mom}) were purchased from The Jackson Laboratory and crossed in

our mouse facility. Cybb$^{-/-}$ mice *(109)* were obtained from Dr. Juhi Bagaitkar at Nationwide

Children's Hospital, Columbus, USA. Experiments assessing Blimp-1 expression with Blimp-1

YFP reporter were reproduced using Blimp-1 GFP reporter mice (a kind gift from Stephen Nutt,

WEHI, Melbourne, Australia). Gzmb[cre] mice have the human *GZMB* promoter directing Cre

recombinase expression to activated T cells; therefore, it does not directly impact mouse

granzyme B expression. Gzmb[cre] and Blimp-1[flox/flox] were crossed to generate the deletion of

Blimp-1 in activated T cells, and here we called purified CD8 T cells Blimp-1[cKO]. ODD[cre]/ERT2

and Blimp-1[flox/flox] were crossed to generate mice in which Blimp-1 deletion was restricted to

hypoxic cells in tamoxifen-treated mice. All mice were maintained in a specific pathogen-free

environment at The Ohio State University or The University of Pennsylvania Animal Care

Facilities.

Parasites

L. major (strain WHO/MHOM/IL/80/Friedlin) and *L. braziliensis*

(MHOM/BR/01/BA788) were grown in Schneider's insect medium (GIBCO) supplemented with

20% heat-inactivated FBS (Sigma, St. Louis, MO, USA) and 2 mM glutamine (Thermo-fisher,

Waltham, MA, USA). Metacyclic-enriched promastigotes were used for infection *(110)*. Mice

were infected with either 10^6 *L. major* or 10^5 *L. braziliensis* intradermally in the ear, and the

lesion progression was monitored weekly by measuring the diameter of the ear with a digital

caliper.

Cell purification and adoptive transfer

Intralesional cell transfer mouse model: CD45.2 and CD45.1 mice were infected with *L.*

major. CD45.2 mice were euthanized three weeks post-infection, and CD8 T cells from dLNs

were purified using a magnetic bead separation kit (Miltenyi Biotec, Germany). 10^6 CD8 T cells

were injected into the lesions of CD45.1 mice. Two days post-injection, recipient mice were

euthanized, and granzyme B expression in CD8 T cells in the ear and dLN were analyzed by

flow cytometry directly ex vivo. RAG$^{-/-}$ mouse model: splenocytes from WT or Blimp-1^{cKO} mice

were collected, red blood cells lysed with ACK lysing buffer (LONZA, Switzerland) and CD8 T

cells were purified using a magnetic bead separation kit (Miltenyi Biotec, Germany). Three

million CD8 T cells were transferred intravenously into RAG$^{-/-}$ mice subsequently infected with

L. braziliensis. Mice reconstituted with CD8 T cells received four injections of 250 μg of anti-

CD4 (clone GK1.5, BioXCell, Lebanon, NH, USA) within the first two weeks.

In vivo treatment

FTY720 (Cayman Chemical, Ann Arbor, Michigan, USA) was administered

intraperitoneally (3 mg/kg) daily 12 days post-infection and for ten days before euthanasia. To

induce Cre expression in the ODD^{cre}/ERT2 mice, tamoxifen (Sigma, St. Louis, MO, USA) was

dissolved in corn oil and administered intraperitoneally (67mg/kg) for seven days following

infection. Pimonidazole (Hypoxyprobe Inc, Burlington, Massachusetts, USA) was administered

intraperitoneally (9 mg/mL, 200 μL) one hour before euthanasia. For neutrophil depletion, mice

received injections every three days of 250 μg of anti-Ly6G (clone 1A8, BioXCell, Lebanon,

NH, USA) within the first two weeks.

Tissue single-cell suspension preparation

Infected ears were harvested, the dorsal and ventral layers of the ear separated, and the

ears were incubated in RPMI 1640 (Gibco, Canada) with 250 µg/mL of Liberase TL (Roche

Diagnostics, Chicago, IL, USA) for 60-90 mins in a shaker at 37°C and 5% CO_2. Ears were

dissociated using a cell strainer (40 µm, BD Pharmingen, Franklin Lakes, NJ, USA), and an

aliquot of the cell suspension was used for parasite titration. dLNs were homogenized using a

cell strainer (40 µm, BD Pharmingen, Franklin Lakes, NJ, USA) to obtain single-cell

suspensions.

Parasite titration

434   The parasite burden in the ears was quantified as described previously *(111)*. Briefly, the

435   homogenate was serially diluted and incubated at 26°C. The number of viable parasites was

436   calculated from the highest dilution at which parasites were observed after seven days.

In vitro stimulation

dLN single-cell suspensions from C57BL/6 infected mice were cultured in 96 well plates

"U" bottom with 5µg/mL of plate-bound anti-CD3 (clone 145-2C11, Invitrogen, Waltham, MA,

USA), 0.5 µg/mL soluble anti-CD28 (clone 37.51, Invitrogen, Waltham, MA, USA), 37°C and

5% CO_2 for 48 hours in RPMI 1640 containing 100 Units of penicillin and 0.1 mg/mL of

Streptomycin (Sigma, St. Louis, MO, USA), 2mM L-Glutamine (Thermo-fisher, Waltham, MA,

USA) and 10% FBS (Sigma, St. Louis, MO, USA). Cells were cultured in 1 mM of DMOG

(Sigma, St. Louis, MO, USA) dissolved in DMSO or an equivalent volume of DMSO (vehicle)

for 48 hours. For mRNA isolation, dLN single-cell suspensions from C57BL/6 infected mice

were obtained, and CD8 T cells were purified using a magnetic bead separation kit (Miltenyi

Biotec, Germany). CD8 T cells were cultured in 96 well plates "U" bottom with 5μg/mL of

plate-bound anti-CD3 (clone 145-2C11, Invitrogen, Waltham, MA, USA), 0.5 μg/mL soluble

anti-CD28 (clone 37.51, Invitrogen, Waltham, MA, USA), 20 U/mL recombinant human IL-2 at

37ºC and 5% CO_2 for 24 hours in RPMI 1640 (Gibco, Canada) containing 100 Units of penicillin

and 0.1 mg/mL of Streptomycin (Sigma, St. Louis, MO, USA), 2mM L-Glutamine (Thermo-

fisher, Waltham, MA, USA) and 10% FBS (Sigma, St. Louis, MO, USA). Cells were cultured in

1 mM of DMOG (Sigma, St. Louis, MO, USA) dissolved in DMSO or an equivalent volume of

DMSO (vehicle).

Flow cytometric analysis

Before surface and intracellular staining, cell suspensions were stained with a

LIVE/DEAD fixable aqua dead cell stain kit (Thermo-fisher, Waltham, MA, USA), according to

manufacturer instructions. Analysis was performed using the FlowJo Software (Tree Star,

Ashland, OR, USA), and gates were drawn based on fluorescence minus one control. List of

antibodies: CD45 (clone 30-F11), GzmB (clone GB11), CD11b (clone M1/70) (all Invitrogen,

Waltham, MA, USA). CD44 (clone IM7, BD Biosciences, Franklin Lakes, NJ, USA), CD45.2

(clone 104), CD45.1 (clone A20), CD90 (clone 53-2), CD3 (clone 17A2), CD8β (clone

YTS156.7.7), and Ly6G (clone 1A8) (all from BioLegend, San Diego, CA, USA). The stained

cells were run on BD FACSymphony A3 or FACSCanto II (BD Biosciences, San Jose, CA,

USA).

Quantification of *Prdm1* mRNA by qPCR

RNA extraction of in vitro stimulated CD8 T cells was performed using nulceospin RNA, Mini kit (Macherery-Magel, Germany) and used to prepare complementary DNA (cDNA) using iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad, Hercules, CA, USA) on ABS Proflex Thermal cycler (Thermo-fisher, Waltham, MA, USA). qPCR was carried out on a C1000 RT-PCR using iTaq Universal SYBR Green Supermix (all from Bio-Rad, Hercules, CA, USA) and primers targeting *Prdm1* (forward 5'-TTC TCT TGG AAA AAC GTG TGG G-3' and reverse 5'-GGA GCC GGA GCT AGACTT G-3'). The qPCR results were normalized to β-Actin (forward 5'-CGC TGT ATT CCC CTC CAT CG-3' and reverse 5'-CCA GTT GGT AAC AAT GCC ATG T-3'). All reactions were carried out in duplicates, and data is represented as fold-change (FC) over the average expression of vehicle-treated cells.

Histology

Two weeks after infection, mice were euthanized, and infected skin (lesions) and contralateral (naïve skin) ears were collected, immediately immersed in 2% paraformaldehyde (PFA) (Thermo-fisher, Waltham, MA, USA), and incubated at room temperature for one hour. Ears were divided in half and re-immersed in PFA for another hour at room temperature. Samples were rinsed and immersed in phosphate-buffered saline (PBS, pH 7.4) (Gibco, Canada) for two hours, followed by cryoprotection in sucrose 30% for 48 hours. Cryoprotected samples were preserved in optimal cutting temperature (OCT) compound (Thermo-fisher, Waltham, MA, USA), and serial sections of 15 µm were obtained on a Tissue-Tek Cryo3 Flex cryostat (Sakura Finetek, Torrance, CA). Sections were collected onto Superfrost Plus microscope slides (Thermo-fisher, Waltham, MA, USA), air-dried, and stored at -80°C.

Immunofluorescence, Image acquisition, and Analysis

Slides were dried at room temperature for one hour, rinsed in PBS, and incubated for one

hour in blocking solution (4% bovine serum albumin and 0.1% Triton X-100 (both from

Thermo-fisher, Waltham, MA, USA) dissolved in PBS). Sections were incubated overnight at

4°C with rabbit anti-Pimonidazole antisera (Hypoxyprobe Inc, Burlington, Massachusetts, USA).

The slides were washed with PBS and incubated for 4 hours at room temperature with rat anti-

mouse Ly6G (cone 1A8, BioLegend, San Diego, CA, USA). Slides were washed with PBS and

incubated separately with Alexa Fluor 568 goat anti-rabbit (Invitrogen, Waltham, MA, USA) and

Alexa Fluor 488 goat anti-rat (Invitrogen, Waltham, MA, USA). Sections were covered in each

secondary antibody for two hours at room temperature. Finally, the sections were counter-stained

with the nuclear probe 4',6-diamino-2-phenylindole (DAPI, Biotechne-Tocris, UK). Slides were

covered with Prolong™ Gold Antifade Mountant (Invitrogen, Waltham, MA, USA) and kept

protected from the light at room temperature for a minimum of 24 hours to allow the mounting

media to dry before imaging. Representative images were acquired with 10x and 20x objectives

in an Olympus FV3000 confocal microscopy (Olympus Corporation, Japan). The color of the

different structures was digitally inverted to better match the presented data in this manuscript.

Using ImageJ software (version 1.53, National Institute of Health), 27 regions of interest (ROIs)

measuring 600x40 µm were created. From the low signal edge to the other, the pixel intensity

values of each ROI were obtained.

Transcriptional profiling of purified mouse CD8 T cells

C57BL/6 mice were infected with *L. major,* and five weeks post-infection, mice were

euthanized; cells from the infected ear and dLN cells were purified and stained for flow

cytometric analysis. Antigen-experienced CD8 T cells were purified using FACSAria cell sorter

(gating strategy: live, singlets, CD90$^+$, CD8$^+$CD44^{high}), and RNA was extracted using the

RNeasy Plus Mini Kit (QIAGEN, Germany) according to the manufacturer's instructions and

used to prepare Poly(A)+-enriched cDNA libraries Illumina TruSeq Stranded mRNA library

prep workflow. Ribo-Zero Gold rRNA depletion (Illumina, San Diego, CA, USA) removed

ribosomal content. Quality assessment and quantification of RNA preparations and libraries were

performed using an Agilent 4200 TapeStation and Qubit 3, respectively. Samples were

sequenced on an Illumina NextSeq 500 to produce 75–base pair single-end reads. The raw reads

were mapped to the mouse reference transcriptome (Ensembl; Mus musculus version 108) using

Kallisto version 0.46.0, and MultiQC v1.9 was used to check the quality of the alignment.

All subsequent analyses were conducted using the statistical computing environment R

version 4.1.0, RStudio version 1.4.1717, and Bioconductor version 3.13. Briefly, transcript

quantification data were summarized to genes using the BiomaRt and tximport package and

normalized using the trimmed mean of M values (TMM) method in edgeR. Genes with <1 CPM

in at least three samples were filtered out. Normalized filtered data were variance-stabilized

using the voom function in limma, and Differential Gene Expression analysis was performed

with linear modeling using limma after correcting for multiple testing using Benjamini-Hochberg

FDR correction. The list of 7 genes encoding for transcription factors associated with effector

like T cell functions was gathered from *(55–57).* Hypoxic enrichment per RNA-seq sample was

calculate with single-sample GSEA using the GSVA R package. The 5 Hypoxia-related

signatures used to estimate those levels were downloaded from: Harris et al. 2022 *(58)*, Hallmark

systematic name (sn) #M5891, Biocarta sn #M13324, Biocarta sn #M14863 and Reactome sn

#M641.

Hypoxia-related transcriptional analyses on lesion biopsies from cutaneous leishmaniasis

patients

Data in Fig. 6 are derived from published transcriptional profiling *(51, 76)*. The analysis

carried out for this current study was performed from the filtered, normalized gene expression

matrix available for download as a text file on NCBI GEO accession #GSE162760 and

#GSE214397. All subsequent analyses were conducted similarly to the purified mouse CD8 T

cells RNA-seq dataset. Hypoxia scores for the human RNA-seq dataset were calculated using the

Harris et al. gene signature *(58)*. Unsupervised hierarchical clustering classifying lesion samples

according to their "hypoxic scores" was performed with maximum distances and Ward's D2

minimum variance. Permanova statistical testing was performed with the vegan R package. Gene

Set Enrichment Analysis (GSEA) was performed using the Broad Institute software version

4.3.2. Gene Ontology (GO) analyses were carried out using DAVID Bioinformatics Resources

(2021 Update) from NIAID/NIH and biological process terms, clustering genesets by annotation

similarity. MCP-counter and immundeconv R packages were combined to estimate neutrophil

abundances from the unstructured RNA-seq dataset. Data visualization was mostly performed in

GraphPad Prism version 10 and in R programming language using tidyverse and gplots tools.

List of Supplementary Materials

Fig. S1. Genes overexpressed in CD8 T cells purified from dLNs compared to lesions.

Fig. S2. Pimonidazole expression in Leishmania-infected skin.

Fig. S3. Neutrophil and pimonidazole expression in Leishmania-infected skin.

Data File S1. Differentially expressed genes between CD8 T cells isolated from the lesions

compared to the dLN.

Data File S2. Gene Set Enrichment Analysis (GSEA) between CD8 T cells isolated from the

lesions compared to the dLN accessing the C2 CP collection of pathways.

Data File S3. Clinical metadata from lesions of patients with cutaneous leishmaniasis, including

the associated Harris et al. Hypoxia scores and Neutrophil abundances.

Data File S4. Clinical metadata from blood of patients with cutaneous leishmaniasis, including

the associated Harris et al. Hypoxia scores.

Data File S5. Differentially expressed genes between hypoxic and baseline lesions from patients

with cutaneous leishmaniasis.

Data File S6. Gene Ontology (GO) performed on DEGs between hypoxic and baseline lesions

from patients with cutaneous leishmaniasis accessing the Biological Process database.

References

1. P. Scott, F. O. Novais, Cutaneous leishmaniasis: immune responses in protection and pathogenesis., *Nat. Rev. Immunol.* **16**, 581–592 (2016).

2. F. O. Novais, C. F. Amorim, P. Scott, Host-Directed Therapies for Cutaneous Leishmaniasis., *Front. Immunol.* **12**, 660183 (2021).

3. F. O. Novais, L. P. Carvalho, J. W. Graff, D. P. Beiting, G. Ruthel, D. S. Roos, M. R. Betts, M. H. Goldschmidt, M. E. Wilson, C. I. de Oliveira, P. Scott, Cytotoxic T cells mediate pathology and metastasis in cutaneous leishmaniasis., *PLoS Pathog.* **9**, e1003504 (2013).

4. F. O. Novais, A. M. Carvalho, M. L. Clark, L. P. Carvalho, D. P. Beiting, I. E. Brodsky, E. M. Carvalho, P. Scott, CD8+ T cell cytotoxicity mediates pathology in the skin by inflammasome activation and IL-1β production., *PLoS Pathog.* **13**, e1006196 (2017).

5. F. O. Novais, P. Scott, CD8+ T cells in cutaneous leishmaniasis: the good, the bad, and the ugly., *Semin Immunopathol* **37**, 251–259 (2015).

6. E. J. Crosby, M. Clark, F. O. Novais, E. J. Wherry, P. Scott, Lymphocytic Choriomeningitis Virus Expands a Population of NKG2D+CD8+ T Cells That Exacerbates Disease in Mice Coinfected with Leishmania major., *J. Immunol.* **195**, 3301–3310 (2015).

7. C. F. Amorim, F. O. Novais, B. T. Nguyen, A. M. Misic, L. P. Carvalho, E. M. Carvalho, D. P. Beiting, P. Scott, Variable gene expression and parasite load predict treatment outcome in cutaneous leishmaniasis., *Sci. Transl. Med.* **11** (2019), doi:10.1126/scitranslmed.aax4204.

8. J. C. Silverio, I. R. Pereira, M. da C. Cipitelli, N. F. Vinagre, M. M. Rodrigues, R. T. Gazzinelli, J. Lannes-Vieira, CD8+ T-cells expressing interferon gamma or perforin play antagonistic roles in heart injury in experimental Trypanosoma cruzi-elicited cardiomyopathy., *PLoS Pathog.* **8**, e1002645 (2012).

9. P. Santos E Sousa, S. Ciré, T. Conlan, L. Jardine, C. Tkacz, I. R. Ferrer, C. Lomas, S. Ward, H. West, S. Dertschnig, S. Blobner, T. K. Means, S. Henderson, D. H. Kaplan, M. Collin, V. Plagnol, C. L. Bennett, R. Chakraverty, Peripheral tissues reprogram CD8+ T cells for pathogenicity during graft-versus-host disease., *JCI Insight* **3** (2018), doi:10.1172/jci.insight.97011.

10. A. L. Marzo, H. Yagita, L. Lefrançois, Cutting edge: migration to nonlymphoid tissues results in functional conversion of central to effector memory CD8 T cells., *J. Immunol.* **179**, 36–40 (2007).

11. D. Michonneau, P. Sagoo, B. Breart, Z. Garcia, S. Celli, P. Bousso, The PD-1 Axis Enforces an Anatomical Segregation of CTL Activity that Creates Tumor Niches after Allogeneic Hematopoietic Stem Cell Transplantation., *Immunity* **44**, 143–154 (2016).

12. S. P. Weisberg, B. B. Ural, D. L. Farber, Tissue-specific immunity for a changing world., *Cell* **184**, 1517–1529 (2021).

13. J. M. Schenkel, K. E. Pauken, Localization, tissue biology and T cell state - implications for cancer immunotherapy., *Nat. Rev. Immunol.* (2023), doi:10.1038/s41577-023-00884-8.

14. M. Ammar, M. Mokni, S. Boubaker, A. El Gaied, A. Ben Osman, H. Louzir, Involvement of granzyme B and granulysin in the cytotoxic response in lichen planus., *J. Cutan. Pathol.* **35**, 630–634 (2008).

15. S. Hiroyasu, M. R. Zeglinski, H. Zhao, M. A. Pawluk, C. T. Turner, A. Kasprick, C. Tateishi, W. Nishie, A. Burleigh, P. A. Lennox, N. Van Laeken, N. J. Carr, F. Petersen, R. I. Crawford, H. Shimizu, D. Tsuruta, R. J. Ludwig, D. J. Granville, Granzyme B inhibition reduces disease severity in autoimmune blistering diseases., *Nat. Commun.* **12**, 302 (2021).

16. L.-C. Kaminski, M. Riehn, A. Abel, C. Steeg, D. D. Yar, O. Addai-Mensah, F. Aminkiah, E. Owusu Dabo, T. Jacobs, M. S. Mackroth, Cytotoxic T Cell-Derived Granzyme B Is Increased in Severe Plasmodium Falciparum Malaria., *Front. Immunol.* **10**, 2917 (2019).

17. I. Voskoboinik, J. C. Whisstock, J. A. Trapani, Perforin and granzymes: function, dysfunction and human pathology., *Nat. Rev. Immunol.* **15**, 388–400 (2015).

18. M. Walch, F. Dotiwala, S. Mulik, J. Thiery, T. Kirchhausen, C. Clayberger, A. M. Krensky, D. Martinvalet, J. Lieberman, Cytotoxic cells kill intracellular bacteria through granulysin-mediated delivery of granzymes., *Cell* **157**, 1309–1323 (2014).

19. P. Eeka, P. B. Phanithi, Cytotoxic T Lymphocyte Granzyme-b mediates neuronal cell death during Plasmodium berghei ANKA induced experimental cerebral malaria., *Neurosci. Lett.* **664**, 58–65 (2018).

20. N.-N. Ji, L. Wu, B.-M. Shao, Q.-X. Meng, J.-N. Xu, H.-W. Zhu, Y.-M. Zhang, CTL-Derived Granzyme B Participates in Hippocampal Neuronal Apoptosis Induced by Cardiac Arrest and Resuscitation in Rats., *Front.*

Neurol. **10**, 1306 (2019).

21. A. Gleave, D. J. Granville, Granzyme B in autoimmune skin disease., *Biomolecules* **13** (2023),
doi:10.3390/biom13020388.

22. F. Dotiwala, S. Mulik, R. B. Polidoro, J. A. Ansara, B. A. Burleigh, M. Walch, R. T. Gazzinelli, J. Lieberman,
Killer lymphocytes use granulysin, perforin and granzymes to kill intracellular parasites., *Nat. Med.* **22**, 210–216
(2016).

23. S. S. Metkar, C. Menaa, J. Pardo, B. Wang, R. Wallich, M. Freudenberg, S. Kim, S. M. Raja, L. Shi, M. M.
Simon, C. J. Froelich, Human and mouse granzyme A induce a proinflammatory cytokine response., *Immunity* **29**,
720–733 (2008).

24. Y. Shen, M. R. Zeglinski, C. T. Turner, S. A. Raithatha, Z. Wu, V. Russo, C. Oram, S. Hiroyasu, L. Nabai, H.
Zhao, T. Bozin, K. Westendorf, I. Kopko, R. Huang, S. Arns, J. Tan, H. Zeng, A. Boey, R. Liggins, J. Jaquith, D. R.
Cameron, A. Papp, D. J. Granville, Topical small molecule granzyme B inhibitor improves remodeling in a murine
model of impaired burn wound healing., *Exp Mol Med* **50**, 1–11 (2018).

25. S. Park, B. Griesenauer, H. Jiang, D. Adom, P. Mehrpouya-Bahrami, S. Chakravorty, M. Kazemian, T. Imam, R.
Srivastava, T. A. Hayes, J. Pardo, S. C. Janga, S. Paczesny, M. H. Kaplan, M. R. Olson, Granzyme A–producing T
helper cells are critical for acute graft-versus-host disease, *JCI Insight* (2020).

26. V. R. Parasa, H. Forsslund, T. Enger, D. Lorenz, S. Kullberg, A. Eklund, M. Sköld, J. Wahlström, J. Grunewald,
S. Brighenti, Enhanced CD8+ cytolytic T cell responses in the peripheral circulation of patients with sarcoidosis and
non-Löfgren's disease., *Respir. Med.* **138S**, S38–S44 (2018).

27. C. Junqueira, C. R. R. Barbosa, P. A. C. Costa, A. Teixeira-Carvalho, G. Castro, S. Sen Santara, R. P. Barbosa,
F. Dotiwala, D. B. Pereira, L. R. Antonelli, J. Lieberman, R. T. Gazzinelli, Cytotoxic CD8+ T cells recognize and
kill Plasmodium vivax-infected reticulocytes., *Nat. Med.* **24**, 1330–1336 (2018).

28. S. J. Balin, M. Pellegrini, E. Klechevsky, S. T. Won, D. I. Weiss, A. W. Choi, J. Hakimian, J. Lu, M. T. Ochoa,
B. R. Bloom, L. L. Lanier, S. Stenger, R. L. Modlin, Human antimicrobial cytotoxic T lymphocytes, defined by NK
receptors and antimicrobial proteins, kill intracellular bacteria., *Sci. Immunol.* **3** (2018),
doi:10.1126/sciimmunol.aat7668.

29. P. Trivedi, K. L. Graham, B. Krishnamurthy, S. Fynch, R. M. Slattery, T. W. H. Kay, H. E. Thomas, Perforin
facilitates beta cell killing and regulates autoreactive CD8+ T-cell responses to antigen in mouse models of type 1
diabetes., *Immunol. Cell Biol.* **94**, 334–341 (2016).

30. M. Vičić, S. Peternel, E. Simonić, V. Sotošek-Tokmadžić, D. Massari, I. Brajac, M. Kaštelan, L. Prpić-Massari,
Cytotoxic T lymphocytes as a potential brake of keratinocyte proliferation in psoriasis., *Med. Hypotheses* **87**, 66–68
(2016).

31. V. Patra, J. Strobl, D. Atzmüller, B. Reininger, L. Kleissl, A. Gruber-Wackernagel, J.-F. Nicolas, G. Stary, M.
Vocanson, P. Wolf, Accumulation of Cytotoxic Skin Resident Memory T Cells and Increased Expression of IL-15
in Lesional Skin of Polymorphic Light Eruption., *Front Med (Lausanne)* **9**, 908047 (2022).

32. W.-H. Chung, S.-I. Hung, J.-Y. Yang, S.-C. Su, S.-P. Huang, C.-Y. Wei, S.-W. Chin, C.-C. Chiou, S.-C. Chu,
H.-C. Ho, C.-H. Yang, C.-F. Lu, J.-Y. Wu, Y.-D. Liao, Y.-T. Chen, Granulysin is a key mediator for disseminated
keratinocyte death in Stevens-Johnson syndrome and toxic epidermal necrolysis., *Nat. Med.* **14**, 1343–1350 (2008).

33. A. Haque, S. E. Best, K. Unosson, F. H. Amante, F. de Labastida, N. M. Anstey, G. Karupiah, M. J. Smyth, W.
R. Heath, C. R. Engwerda, Granzyme B expression by CD8+ T cells is required for the development of
experimental cerebral malaria., *J. Immunol.* **186**, 6148–6156 (2011).

34. L. Xing, Z. Dai, A. Jabbari, J. E. Cerise, C. A. Higgins, W. Gong, A. de Jong, S. Harel, G. M. DeStefano, L.
Rothman, P. Singh, L. Petukhova, J. Mackay-Wiggan, A. M. Christiano, R. Clynes, Alopecia areata is driven by
cytotoxic T lymphocytes and is reversed by JAK inhibition., *Nat. Med.* **20**, 1043–1049 (2014).

35. A. Zloza, G. E. Lyons, L. K. Chlewicki, F. J. Kohlhapp, J. A. O'Sullivan, A. T. Lacek, T. V. Moore, M. C.
Jagoda, V. Kumar, J. A. Guevara-Patiño, Engagement of NK receptor NKG2D, but not 2B4, results in self-reactive
CD8+ T cells and autoimmune vitiligo., *Autoimmunity* **44**, 599–606 (2011).

36. C. Jacquemin, C. Martins, F. Lucchese, D. Thiolat, A. Taieb, J. Seneschal, K. Boniface, NKG2D Defines a
Subset of Skin Effector Memory CD8 T Cells with Proinflammatory Functions in Vitiligo., *J. Invest. Dermatol.* **140**,
1143–1153.e5 (2020).

37. J. G. van den Boorn, D. Konijnenberg, T. A. M. Dellemijn, J. P. W. van der Veen, J. D. Bos, C. J. M. Melief, F.
A. Vyth-Dreese, R. M. Luiten, Autoimmune destruction of skin melanocytes by perilesional T cells from vitiligo
patients., *J. Invest. Dermatol.* **129**, 2220–2232 (2009).

38. C. T. Taylor, C. C. Scholz, The effect of HIF on metabolism and immunity., *Nat. Rev. Nephrol.* **18**, 573–587
(2022).

39. V. Schatz, P. Neubert, F. Rieger, J. Jantsch, Hypoxia, Hypoxia-Inducible Factor-1α, and Innate Antileishmanial

Immune Responses., *Front. Immunol.* **9**, 216 (2018).
40. T. Charpentier, A. Hammami, S. Stäger, Hypoxia inducible factor 1α: A critical factor for the immune response
to pathogens and Leishmania., *Cell Immunol.* **309**, 42–49 (2016).
41. A. Mahnke, R. J. Meier, V. Schatz, J. Hofmann, K. Castiglione, U. Schleicher, O. S. Wolfbeis, C. Bogdan, J.
Jantsch, Hypoxia in Leishmania major skin lesions impairs the NO-dependent leishmanicidal activity of
macrophages., *J. Invest. Dermatol.* **134**, 2339–2346 (2014).
42. A. Bowlin, H. Roys, H. Wanjala, M. Bettadapura, G. Venugopal, J. Surma, M. C. Simon, T. Weinkopff,
Hypoxia inducible factor signaling in macrophages promotes lymphangiogenesis in Leishmania major infection.,
Infect. Immun. (2021), doi:10.1128/IAI.00124-21.
43. A. Degrossoli, M. C. Colhone, W. W. Arrais-Silva, S. Giorgio, Hypoxia modulates expression of the 70-kD heat
shock protein and reduces Leishmania infection in macrophages., *J Biomed Sci* **11**, 847–854 (2004).
44. M. C. Colhone, W. W. Arrais-Silva, C. Picoli, S. Giorgio, Effect of hypoxia on macrophage infection by
Leishmania amazonensis., *J. Parasitol.* **90**, 510–515 (2004).
45. W. W. Arrais-Silva, V. A. Paffaro, A. T. Yamada, S. Giorgio, Expression of hypoxia-inducible factor-1alpha in
the cutaneous lesions of BALB/c mice infected with Leishmania amazonensis., *Exp Mol Pathol* **78**, 49–54 (2005).
46. A. Hammami, B. M. Abidin, K. M. Heinonen, S. Stäger, HIF-1α hampers dendritic cell function and Th1
generation during chronic visceral leishmaniasis., *Sci. Rep.* **8**, 3500 (2018).
47. A. Hammami, B. M. Abidin, T. Charpentier, A. Fabié, A.-P. Duguay, K. M. Heinonen, S. Stäger, HIF-1α is a
key regulator in potentiating suppressor activity and limiting the microbicidal capacity of MDSC-like cells during
visceral leishmaniasis., *PLoS Pathog.* **13**, e1006616 (2017).
48. A. Hammami, T. Charpentier, M. Smans, S. Stäger, IRF-5-Mediated Inflammation Limits CD8+ T Cell
Expansion by Inducing HIF-1α and Impairing Dendritic Cell Functions during Leishmania Infection., *PLoS Pathog.*
11, e1004938 (2015).
49. I. Mesquita, C. Ferreira, D. Moreira, G. E. G. Kluck, A. M. Barbosa, E. Torrado, R. J. Dinis-Oliveira, L. G.
Gonçalves, C.-J. Beauparlant, A. Droit, L. Berod, T. Sparwasser, N. Bodhale, B. Saha, F. Rodrigues, C. Cunha, A.
Carvalho, A. G. Castro, J. Estaquier, R. Silvestre, The Absence of HIF-1α Increases Susceptibility to Leishmania
donovani Infection via Activation of BNIP3/mTOR/SREBP-1c Axis., *Cell Rep.* **30**, 4052–4064.e7 (2020).
50. F. O. Novais, B. T. Nguyen, P. Scott, Granzyme B inhibition by tofacitinib blocks the pathology induced by
CD8 T cells in cutaneous leishmaniasis., *J. Invest. Dermatol.* **141**, 575–585 (2021).
51. C. Farias Amorim, V. M. Lovins, T. P. Singh, F. O. Novais, J. C. Harris, A. S. Lago, L. P. Carvalho, E. M.
Carvalho, D. P. Beiting, P. Scott, E. A. Grice, The skin microbiome enhances disease through IL-1b and delays
healing in cutaneous leishmaniasis patients., *medRxiv* (2023), doi:10.1101/2023.02.02.23285247.
52. F. O. Novais, L. P. Carvalho, S. Passos, D. S. Roos, E. M. Carvalho, P. Scott, D. P. Beiting, Genomic profiling
of human Leishmania braziliensis lesions identifies transcriptional modules associated with cutaneous
immunopathology., *J. Invest. Dermatol.* **135**, 94–101 (2015).
53. E. J. Crosby, M. H. Goldschmidt, E. J. Wherry, P. Scott, Engagement of NKG2D on bystander memory CD8 T
cells promotes increased immunopathology following Leishmania major infection., *PLoS Pathog.* **10**, e1003970
(2014).
54. C. da S. Santos, V. Boaventura, C. Ribeiro Cardoso, N. Tavares, M. J. Lordelo, A. Noronha, J. Costa, V. M.
Borges, C. I. de Oliveira, J. Van Weyenbergh, A. Barral, M. Barral-Netto, C. I. Brodskyn, CD8(+) granzyme B(+)-
mediated tissue injury vs. CD4(+)IFNγ(+)-mediated parasite killing in human cutaneous leishmaniasis., *J. Invest.*
Dermatol. **133**, 1533–1540 (2013).
55. Y. Chen, R. Zander, A. Khatun, D. M. Schauder, W. Cui, Transcriptional and epigenetic regulation of effector
and memory CD8 T cell differentiation., *Front. Immunol.* **9**, 2826 (2018).
56. J. T. Chang, E. J. Wherry, A. W. Goldrath, Molecular regulation of effector and memory T cell differentiation.,
Nat. Immunol. **15**, 1104–1115 (2014).
57. B. Yu, K. Zhang, J. J. Milner, C. Toma, R. Chen, J. P. Scott-Browne, R. M. Pereira, S. Crotty, J. T. Chang, M.
E. Pipkin, W. Wang, A. W. Goldrath, Epigenetic landscapes reveal transcription factors that regulate CD8+ T cell
differentiation., *Nat. Immunol.* **18**, 573–582 (2017).
58. A. L. Harris, Hypoxia--a key regulatory factor in tumour growth., *Nat. Rev. Cancer* **2**, 38–47 (2002).
59. S. M. Evans, A. E. Schrlau, A. A. Chalian, P. Zhang, C. J. Koch, Oxygen levels in normal and previously
irradiated human skin as assessed by EF5 binding., *J. Invest. Dermatol.* **126**, 2596–2606 (2006).
60. N. T. S. Evans, P. F. D. Naylor, The systemic oxygen supply to the surface of human skin, *Respir Physiol* **3**, 21–
37 (1967).
61. E. S. Friedman, K. Bittinger, T. V. Esipova, L. Hou, L. Chau, J. Jiang, C. Mesaros, P. J. Lund, X. Liang, G. A.
FitzGerald, M. Goulian, D. Lee, B. A. Garcia, I. A. Blair, S. A. Vinogradov, G. D. Wu, Microbes vs. chemistry in

the origin of the anaerobic gut lumen., *Proc. Natl. Acad. Sci. USA* **115**, 4170–4175 (2018).

62. H. Shin, S. D. Blackburn, A. M. Intlekofer, C. Kao, J. M. Angelosanto, S. L. Reiner, E. J. Wherry, A role for the
transcriptional repressor Blimp-1 in CD8(+) T cell exhaustion during chronic viral infection., *Immunity* **31**, 309–320
(2009).

63. Q. Lin, Y. Huang, F. J. Giordano, Z. Yun, Generation of a hypoxia-sensing mouse model., *Genesis* **58**, e23352
(2020).

64. S.-H. Fu, L.-T. Yeh, C.-C. Chu, B. L.-J. Yen, H.-K. Sytwu, New insights into Blimp-1 in T lymphocytes: a
divergent regulator of cell destiny and effector function., *J Biomed Sci* **24**, 49 (2017).

65. M. Montes de Oca, R. Kumar, F. de Labastida Rivera, F. H. Amante, M. Sheel, R. J. Faleiro, P. T. Bunn, S. E.
Best, L. Beattie, S. S. Ng, C. L. Edwards, W. Muller, E. Cretney, S. L. Nutt, M. J. Smyth, A. Haque, G. R. Hill, S.
Sundar, A. Kallies, C. R. Engwerda, Blimp-1-Dependent IL-10 Production by Tr1 Cells Regulates TNF-Mediated
Tissue Pathology., *PLoS Pathog.* **12**, e1005398 (2016).

66. Y. Belkaid, E. Von Stebut, S. Mendez, R. Lira, E. Caler, S. Bertholet, M. C. Udey, D. Sacks, CD8+ T cells are
required for primary immunity in C57BL/6 mice following low-dose, intradermal challenge with Leishmania major.,
J. Immunol. **168**, 3992–4000 (2002).

67. F. O. Novais, A. C. Wong, D. O. Villareal, D. P. Beiting, P. Scott, CD8+ T Cells Lack Local Signals To Produce
IFN-γ in the Skin during Leishmania Infection., *J. Immunol.* **200**, 1737–1745 (2018).

68. T. Weinkopff, H. Roys, A. Bowlin, P. Scott, Leishmania Infection Induces Macrophage Vascular Endothelial
Growth Factor A Production in an ARNT/HIF-Dependent Manner., *Infect. Immun.* **87** (2019),
doi:10.1128/IAI.00088-19.

69. N. C. Peters, J. G. Egen, N. Secundino, A. Debrabant, N. Kimblin, S. Kamhawi, P. Lawyer, M. P. Fay, R. N.
Germain, D. Sacks, In vivo imaging reveals an essential role for neutrophils in leishmaniasis transmitted by sand
flies., *Science* **321**, 970–974 (2008).

70. K. Passelli, O. Billion, F. Tacchini-Cottier, The impact of neutrophil recruitment to the skin on the pathology
induced by leishmania infection., *Front. Immunol.* **12**, 649348 (2021).

71. C. P. Pedraza-Zamora, J. Delgado-Domínguez, J. Zamora-Chimal, I. Becker, Th17 cells and neutrophils: Close
collaborators in chronic Leishmania mexicana infections leading to disease severity., *Parasite Immunol* **39** (2017),
doi:10.1111/pim.12420.

72. M. Charmoy, B. P. Hurrell, A. Romano, S. H. Lee, F. Ribeiro-Gomes, N. Riteau, K. Mayer-Barber, F. Tacchini-
Cottier, D. L. Sacks, The Nlrp3 inflammasome, IL-1β, and neutrophil recruitment are required for susceptibility to a
nonhealing strain of Leishmania major in C57BL/6 mice., *Eur. J. Immunol.* **46**, 897–911 (2016).

73. S. da C. Gonçalves-de-Albuquerque, R. Pessoa-E-Silva, L. A. M. Trajano-Silva, T. C. de Goes, R. C. S. de
Morais, C. N. da C Oliveira, V. M. B. de Lorena, M. de Paiva-Cavalcanti, The equivocal role of th17 cells and
neutrophils on immunopathogenesis of leishmaniasis., *Front. Immunol.* **8**, 1437 (2017).

74. C. Gimblet, J. S. Meisel, M. A. Loesche, S. D. Cole, J. Horwinski, F. O. Novais, A. M. Misic, C. W. Bradley, D.
P. Beiting, S. C. Rankin, L. P. Carvalho, E. M. Carvalho, P. Scott, E. A. Grice, Cutaneous Leishmaniasis Induces a
Transmissible Dysbiotic Skin Microbiota that Promotes Skin Inflammation., *Cell Host Microbe* **22**, 13–24.e4
(2017).

75. Z. M. Moghadam, P. Henneke, J. Kolter, From flies to men: ROS and the NADPH oxidase in phagocytes.,
Front. Cell Dev. Biol. **9**, 628991 (2021).

76. C. Farias Amorim, F. O Novais, B. T. Nguyen, M. T. Nascimento, J. Lago, A. S. Lago, L. P. Carvalho, D. P.
Beiting, P. Scott, Localized skin inflammation during cutaneous leishmaniasis drives a chronic, systemic IFN-γ
signature., *PLoS Negl. Trop. Dis.* **15**, e0009321 (2021).

77. C. T. Turner, M. R. Zeglinski, K. C. Richardson, S. Santacruz, S. Hiroyasu, C. Wang, H. Zhao, Y. Shen, R.
Sehmi, H. Lima, G. M. Gauvreau, D. J. Granville, Granzyme B Contributes to Barrier Dysfunction in Oxazolone-
Induced Skin Inflammation through E-Cadherin and FLG Cleavage., *J. Invest. Dermatol.* **141**, 36–47 (2021).

78. J. K. Leavey, R. L. Tarleton, Cutting edge: dysfunctional CD8+ T cells reside in nonlymphoid tissues during
chronic Trypanosoma cruzi infection., *J. Immunol.* **170**, 2264–2268 (2003).

79. S. Chouaib, M. Z. Noman, K. Kosmatopoulos, M. A. Curran, Hypoxic stress: obstacles and opportunities for
innovative immunotherapy of cancer., *Oncogene* **36**, 439–445 (2017).

80. D. K.-C. Chiu, A. P.-W. Tse, I. M.-J. Xu, J. Di Cui, R. K.-H. Lai, L. L. Li, H.-Y. Koh, F. H.-C. Tsang, L. L.
Wei, C.-M. Wong, I. O.-L. Ng, C. C.-L. Wong, Hypoxia inducible factor HIF-1 promotes myeloid-derived
suppressor cells accumulation through ENTPD2/CD39L1 in hepatocellular carcinoma., *Nat. Commun.* **8**, 517
(2017).

81. S.-H. Chiou, V. I. Risca, G. X. Wang, D. Yang, B. M. Grüner, A. S. Kathiria, R. K. Ma, D. Vaka, P. Chu, M.
Kozak, L. Castellini, E. E. Graves, G. E. Kim, P. Mourrain, A. C. Koong, A. J. Giaccia, M. M. Winslow, BLIMP1

induces transient metastatic heterogeneity in pancreatic cancer., *Cancer Discov.* **7**, 1184–1199 (2017).

82. I. Liikanen, C. Lauhan, S. Quon, K. Omilusik, A. T. Phan, L. B. Bartrolí, A. Ferry, J. Goulding, J. Chen, J. P.
Scott-Browne, J. T. Yustein, N. E. Scharping, D. A. Witherden, A. W. Goldrath, Hypoxia-inducible factor activity
promotes antitumor effector function and tissue residency by CD8+ T cells., *J. Clin. Invest.* **131** (2021),
doi:10.1172/JCI143729.

83. P. Veliça, P. P. Cunha, N. Vojnovic, I. P. Foskolou, D. Bargiela, M. Gojkovic, H. Rundqvist, R. S. Johnson,
Modified Hypoxia-Inducible Factor Expression in CD8+ T Cells Increases Antitumor Efficacy., *Cancer Immunol*
Res **9**, 401–414 (2021).

84. P. P. Cunha, E. Minogue, L. C. M. Krause, R. M. Hess, D. Bargiela, B. J. Wadsworth, L. Barbieri, C. Brombach,
I. P. Foskolou, I. Bogeski, P. Velica, R. S. Johnson, Oxygen levels at the time of activation determine T cell
persistence and immunotherapeutic efficacy., *Elife* **12** (2023), doi:10.7554/eLife.84280.

85. A. L. Doedens, A. T. Phan, M. H. Stradner, J. K. Fujimoto, J. V. Nguyen, E. Yang, R. S. Johnson, A. W.
Goldrath, Hypoxia-inducible factors enhance the effector responses of CD8(+) T cells to persistent antigen., *Nat.*
Immunol. **14**, 1173–1182 (2013).

86. D. K. Finlay, E. Rosenzweig, L. V. Sinclair, C. Feijoo-Carnero, J. L. Hukelmann, J. Rolf, A. A. Panteleyev, K.
Okkenhaug, D. A. Cantrell, PDK1 regulation of mTOR and hypoxia-inducible factor 1 integrate metabolism and
migration of CD8+ T cells., *J. Exp. Med.* **209**, 2441–2453 (2012).

87. Y. Gropper, T. Feferman, T. Shalit, T.-M. Salame, Z. Porat, G. Shakhar, Culturing CTLs under Hypoxic
Conditions Enhances Their Cytolysis and Improves Their Anti-tumor Function., *Cell Rep.* **20**, 2547–2555 (2017).

88. A. J. Little, P.-M. Chen, M. D. Vesely, R. N. Khan, J. Fiedler, J. Garritano, F. M. Islam, J. M. McNiff, J. E.
Craft, HIF-1 regulates pathogenic cytotoxic T cells in lupus skin disease, *JCI Insight* (2023).

89. F. Hasan, Y. Chiu, R. M. Shaw, J. Wang, C. Yee, Hypoxia acts as an environmental cue for the human tissue-
resident memory T cell differentiation program., *JCI Insight* **6** (2021), doi:10.1172/jci.insight.138970.

90. N. Bannoud, T. Dalotto-Moreno, L. Kindgard, P. A. García, A. G. Blidner, K. V. Mariño, G. A. Rabinovich, D.
O. Croci, Hypoxia supports differentiation of terminally exhausted CD8 T cells., *Front. Immunol.* **12**, 660944
(2021).

91. N. E. Scharping, D. B. Rivadeneira, A. V. Menk, P. D. A. Vignali, B. R. Ford, N. L. Rittenhouse, R. Peralta, Y.
Wang, Y. Wang, K. DePeaux, A. C. Poholek, G. M. Delgoffe, Mitochondrial stress induced by continuous
stimulation under hypoxia rapidly drives T cell exhaustion., *Nat. Immunol.* **22**, 205–215 (2021).

92. B. R. Ford, N. L. Rittenhouse, N. E. Scharping, P. D. A. Vignali, A. T. Frisch, R. Peralta, G. M. Delgoffe, A. C.
Poholek, Altered costimulatory signals and hypoxia support chromatin landscapes limiting the functional potential
of exhausted T cells in cancer, *BioRxiv* (2021), doi:10.1101/2021.07.11.451947.

93. F. M. Behr, N. A. M. Kragten, T. H. Wesselink, B. Nota, R. A. W. van Lier, D. Amsen, R. Stark, P. Hombrink,
K. P. J. M. van Gisbergen, Blimp-1 Rather Than Hobit Drives the Formation of Tissue-Resident Memory CD8+ T
Cells in the Lungs., *Front. Immunol.* **10**, 400 (2019).

94. L. K. Mackay, M. Minnich, N. A. M. Kragten, Y. Liao, B. Nota, C. Seillet, A. Zaid, K. Man, S. Preston, D.
Freestone, A. Braun, E. Wynne-Jones, F. M. Behr, R. Stark, D. G. Pellicci, D. I. Godfrey, G. T. Belz, M. Pellegrini,
T. Gebhardt, M. Busslinger, W. Shi, F. R. Carbone, R. A. W. van Lier, A. Kallies, K. P. J. M. van Gisbergen, Hobit
and Blimp1 instruct a universal transcriptional program of tissue residency in lymphocytes., *Science* **352**, 459–463
(2016).

95. N. A. M. Kragten, F. M. Behr, F. A. Vieira Braga, E. B. M. Remmerswaal, T. H. Wesselink, A. E. Oja, P.
Hombrink, A. Kallies, R. A. W. van Lier, R. Stark, K. P. J. M. van Gisbergen, Blimp-1 induces and Hobit maintains
the cytotoxic mediator granzyme B in CD8 T cells., *Eur. J. Immunol.* **48**, 1644–1662 (2018).

96. E. Doz, R. Lombard, F. Carreras, D. Buzoni-Gatel, N. Winter, Mycobacteria-infected dendritic cells attract
neutrophils that produce IL-10 and specifically shut down Th17 CD4 T cells through their IL-10 receptor., *J.*
Immunol. **191**, 3818–3826 (2013).

97. M. Rambault, É. Doz-Deblauwe, Y. Le Vern, F. Carreras, P. Cunha, P. Germon, P. Rainard, N. Winter, A.
Remot, Neutrophils encompass a regulatory subset suppressing T cells in apparently healthy cattle and mice., *Front.*
Immunol. **12**, 625244 (2021).

98. N. Glodde, T. Bald, D. van den Boorn-Konijnenberg, K. Nakamura, J. S. O'Donnell, S. Szczepanski, M.
Brandes, S. Eickhoff, I. Das, N. Shridhar, D. Hinze, M. Rogava, T. C. van der Sluis, J. J. Ruotsalainen, E. Gaffal, J.
Landsberg, K. U. Ludwig, C. Wilhelm, M. Riek-Burchardt, A. J. Müller, C. Gebhardt, R. A. Scolyer, G. V. Long, V.
Janzen, M. W. L. Teng, W. Kastenmüller, M. Mazzone, M. J. Smyth, T. Tüting, M. Hölzel, Reactive Neutrophil
Responses Dependent on the Receptor Tyrosine Kinase c-MET Limit Cancer Immunotherapy., *Immunity* **47**, 789–
802.e9 (2017).

99. S. B. Coffelt, K. Kersten, C. W. Doornebal, J. Weiden, K. Vrijland, C.-S. Hau, N. J. M. Verstegen, M.

Ciampricotti, L. J. A. C. Hawinkels, J. Jonkers, K. E. de Visser, IL-17-producing γδ T cells and neutrophils conspire
to promote breast cancer metastasis., *Nature* **522**, 345–348 (2015).
100. P. J. Volberding, G. Xin, M. Y. Kasmani, A. Khatun, A. K. Brown, C. Nguyen, J. S. Stancill, E. Martinez, J. A.
Corbett, W. Cui, Suppressive neutrophils require PIM1 for metabolic fitness and survival during chronic viral
infection., *Cell Rep.* **35**, 109160 (2021).
101. F. C. Weber, T. Németh, J. Z. Csepregi, A. Dudeck, A. Roers, B. Ozsvári, E. Oswald, L. G. Puskás, T. Jakob,
A. Mócsai, S. F. Martin, Neutrophils are required for both the sensitization and elicitation phase of contact
hypersensitivity., *J. Exp. Med.* **212**, 15–22 (2015).
102. F. O. Novais, R. C. Santiago, A. Báfica, R. Khouri, L. Afonso, V. M. Borges, C. Brodskyn, M. Barral-Netto, A.
Barral, C. I. de Oliveira, Neutrophils and macrophages cooperate in host resistance against Leishmania braziliensis
infection., *J. Immunol.* **183**, 8088–8098 (2009).
103. A. B. Guimarães-Costa, M. T. C. Nascimento, G. S. Froment, R. P. P. Soares, F. N. Morgado, F. Conceição-
Silva, E. M. Saraiva, Leishmania amazonensis promastigotes induce and are killed by neutrophil extracellular traps.,
Proc. Natl. Acad. Sci. USA **106**, 6748–6753 (2009).
104. F. L. Ribeiro-Gomes, N. C. Peters, A. Debrabant, D. L. Sacks, Efficient capture of infected neutrophils by
dendritic cells in the skin inhibits the early anti-leishmania response., *PLoS Pathog.* **8**, e1002536 (2012).
105. G. van Zandbergen, M. Klinger, A. Mueller, S. Dannenberg, A. Gebert, W. Solbach, T. Laskay, Cutting edge:
neutrophil granulocyte serves as a vector for Leishmania entry into macrophages., *J. Immunol.* **173**, 6521–6525
(2004).
106. C. Gonzalez-Lombana, C. Gimblet, O. Bacellar, W. W. Oliveira, S. Passos, L. P. Carvalho, M. Goldschmidt, E.
M. Carvalho, P. Scott, IL-17 mediates immunopathology in the absence of IL-10 following Leishmania major
infection., *PLoS Pathog.* **9**, e1003243 (2013).
107. P. Rao, S. Suvas, Development of inflammatory hypoxia and prevalence of glycolytic metabolism in
progressing herpes stromal keratitis lesions., *J. Immunol.* **202**, 514–526 (2019).
108. E. L. Campbell, W. J. Bruyninckx, C. J. Kelly, L. E. Glover, E. N. McNamee, B. E. Bowers, A. J. Bayless, M.
Scully, B. J. Saeedi, L. Golden-Mason, S. F. Ehrentraut, V. F. Curtis, A. Burgess, J. F. Garvey, A. Sorensen, R.
Nemenoff, P. Jedlicka, C. T. Taylor, D. J. Kominsky, S. P. Colgan, Transmigrating neutrophils shape the mucosal
microenvironment through localized oxygen depletion to influence resolution of inflammation., *Immunity* **40**, 66–77
(2014).
109. J. D. Pollock, D. A. Williams, M. A. Gifford, L. L. Li, X. Du, J. Fisherman, S. H. Orkin, C. M. Doerschuk, M.
C. Dinauer, Mouse model of X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide
production., *Nat. Genet.* **9**, 202–209 (1995).
110. G. F. Späth, S. M. Beverley, A lipophosphoglycan-independent method for isolation of infective Leishmania
metacyclic promastigotes by density gradient centrifugation., *Exp Parasitol* **99**, 97–103 (2001).
111. J. E. Uzonna, K. L. Joyce, P. Scott, Low dose Leishmania major promotes a transient T helper cell type 2
response that is down-regulated by interferon gamma-producing CD8+ T cells., *J. Exp. Med.* **199**, 1559–1566
(2004).
888
889

Acknowledgments: The authors would like to thank Dr. Juhi Bagaitkar for providing Cybb

deficient mice.

Funding:

National Institutes of Health grant R01AI162711 (FON)

National Institute of Health training program T32 AI165391 "Interdisciplinary Program

in Microbe-Host Biology" (EAF)

National Institute of Health training program T32 AI118684 (WZ)

Host-Microbial Analytic and Repository Core in the Center for Molecular Studies in

Digestive and Liver Diseases (P30DK050306) (GDW)

Author contributions:

Conceptualization: FON

Methodology: EAF, CFA, RAS, EDSH, KM, LAS, WZ, GDW, FON, PLC

Investigation: EAF, CFA, RAS, EDSH, KM, LAS, WZ, FON, PLC

Visualization: EAF, CFA, KAM, WZ

Funding acquisition: FON, CIO

Project administration: FON

Supervision: FON

Writing – original draft: EAF, CFA, FON

Writing – review & editing: PLC

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Raw sequence data from purified mouse CD8 T cells are

available on the NCBI GEO BioProject PRJNA794119. The complete RNA-seq data analysis, R

code, file inputs, and outputs used to perform the transcriptional analysis presented in this current

manuscript are available on the GitHub repository "Hypoxia_FON"

(https://github.com/camilafarias112/Hypoxia_FON).

Figure 1

Fig 1. CD8 T cell function is tissue-specific.

(a) Granzyme B (GzmB) expression in CD8 T cells from dLNs and lesions from C57BL/6 mice in week two post-
infection with *L. major*. Representative data from more than three experiments with at least four mice per
experiment. (b) Granzyme B expression in dLN CD8 T cells from *L. major* infected C57BL/6 mice treated with
FTY720 or vehicle daily 12 days post-infection for 10 days. Data from one independent experiment with five mice
per group. (c - left) Schematic representation of the transfer of CD45.2 dLN purified CD8 T cells into the lesions of
CD45.1 recipients, both infected for three weeks. (c - right) Granzyme B expression in donor CD8 T cells from
dLNs and lesions from recipient mice 48 hours post-transfer. Data from 2 experiments independent experiments
combined. (d-g) RNA-seq analyses from antigen-experienced (CD44high) CD8 T cells purified from dLNs and
lesions of *L. major*-infected mice in week five. (d) Principal component analysis showing principal component 1
(PC1) and PC2. (e) Heat map of top 20 differentially expressed genes (DEGs) between purified CD8 T cells from
the lesions compared to dLN (FDR>0.05 and FC=1.5). (f) Expression of genes encoding for transcription factors
associated with effector-like T cell functions is represented in a heat map, as well as fold-change (FC) between
lesional and dLN CD8 T cells. (g) Heatmap of hypoxia-related pathways enriched in lesional and dLN CD8 T cells.
*P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001 and ****P ≤ 0.0001. ns = non-significant. FDR = false discovery rates. FC =
fold change.

Figure 2

Fig 2. Hypoxia induces granzyme B and *Prdm1* expression in CD8 T cells.

(a-b) Immunofluorescence staining and confocal microscopy of horizontal sections from lesions of *L. major*
C57BL/6-infected mice for two weeks. Mice received pimonidazole one hour before euthanasia. The top panels
show contralateral ears (naïve skin), and the bottom panels are from infected ears (lesions). Pimonidazole (green)
and nuclear stain DAPI (blue). Representative image from two naïve and three infected ears. Bars (a) 200 μm and
(b) 100 μm. (c) Partial pressure of oxygen in mmHg in dLNs and lesions of C57BL/6 mice infected with *L. major*
for two weeks. Data from two independent experiments combined. (d) Granzyme B (GzmB) expression in CD8 T
cells from dLNs of C57BL/6 mice infected with *L. major* cultured with Dimethyloxalylglycine (DMOG) or vehicle.
(e) Fold-change (FC) of *Prdm1* mRNA over the average expression of vehicle-treated cells measured by qPCR in
DMOG or vehicle-treated purified CD8 T cells. **P ≤ 0.01 and ***P ≤ 0.001

Figure 3

Fig 3. Blimp-1 expression is induced by hypoxia within skin lesions.

(a) Blimp-1 YFP reporter mice expression in CD8 T cells from dLNs and lesions from Blimp-1 reporter mice infected with *L. major* for one week. Data from two independent experiments combined. (b) Granzyme B (GzmB) and Blimp-1 YFP expression in CD8 T cells from dLNs and lesions from Blimp-1 reporter mice infected with *L. major* for one week. (c) Schematic representation of OOD^{cre}Blimp-1$^{flox/flox}$ mice. (d) OOD^{cre}Blimp-1$^{flox/flox}$ mice were infected with *L. major* for one week when daily tamoxifen injections commenced. Mice were euthanized at week two post-infection. GzmB expression of lesional CD8 T cells in tamoxifen-treated and untreated mice. Data from two independent experiments combined. *P ≤ 0.05, **P ≤ 0.01, and ****P ≤ 0.0001.

Figure 4

Fig 4. Blimp-1 expression is required for CD8 T cell-mediated disease.

RAG$^{-/-}$ mice were infected with *L. braziliensis* and reconstituted with CD8 T cells purified from WT or Blimp-1^{cKO} mice. (a) Ear thickness and (b) parasite numbers in lesions at seven weeks of infection. (c) CD8 T cell frequency and (d) GzmB expression by CD8 T cells in lesions were assessed directly ex vivo by flow cytometry seven weeks after infection. Data represent 3 individual experiments with 3-5 mice per group. *P ≤ 0.05 and **P ≤ 0.01. ns = non-significant

Figure 5

Fig 5. Neutrophils consume oxygen from lesions and stimulate granzyme B expression in CD8 T cells.

(a-b) C57BL/6 mice infected with *L. major* received pimonidazole one hour before euthanasia at two hours, five hours, two weeks, and nine weeks. (a) Frequency of neutrophils in lesions and (b) pimonidazole expression in neutrophils (CD11b⁺Ly6Gᵖᵒˢ) and other myeloid cells (CD11b⁺Ly6Gⁿᵉᵍ) in lesions. (c) Confocal microscopy of one representative image of Ly6G (pink) and pimonidazole (green) staining in the skin infected for two weeks. Scale bar = 200 μm. (d) Pixel intensity of pimonidazole and Ly6G based on 27 regions (Fig. S3). (e) Confocal microscopy of two representative images of nuclear (DAPI) staining (blue), Ly6G (pink), and pimonidazole (green) in lesions infected for two weeks. Scale bar = 100 μm. (f-g) C57BL/6 mice infected with *L. major* were injected every three days with anti-Ly6G (clone 1A8) or isotype control until euthanasia at two weeks post-infection. Frequency of (f) neutrophils in lesions. (g) Granzyme B (GzmB) frequency and mean fluorescence intensity (MFI) in CD8 T cells in lesions. (h) Granzyme B frequency and MFI in CD8 T cells in the lesions of WT or Cybb⁻ᐟ⁻ mice infected with *L. major* for two weeks. *P ≤ 0.05, ***P ≤ 0.001 and ****P ≤ 0.0001.

Figure 6

987
Fig 6. The degree of hypoxia-related gene expression is associated with the presence of neutrophils in patients.

(a) Hypoxia score calculated in the blood from healthy subjects (n=14) and patients (n=50); (b) human intact skin
(n=6) and cutaneous leishmaniasis lesions from patients (n=51) based on the Harris et al. hypoxia-gene signature. (c)
Unsupervised hierarchical clustering classified human skin samples in 'baseline' and 'hypoxic' lesions according to
their hypoxia-gene phenotypes. (d) Principal Component Analysis (PCA) of baseline and hypoxic cutaneous
leishmaniasis lesions. (e) Neutrophil abundances estimated by Microenvironment Cell Populations (MCP) counter.
Wilcoxon was used for statistical analysis. (f) Scatter plot showing differential gene expression analysis between
hypoxic vs. baseline lesions (FDR>0.01 and FC=1.5). In blue are the top DGEs from this analysis. (g) Model
summarizing our findings. **P ≤ 0.01, and ***P ≤ 0.001. HS = Healthy subjects. CL = cutaneous leishmaniasis.
FDR = false discovery rates. FC = fold change.

Figure S1

Fig S1. Genes overexpressed in CD8 T cells purified from dLNs compared to lesions.

C57BL/6 mice were infected with *L. major*, and at the peak of the lesion, antigen-experienced (CD44high) CD8 T cells were purified by flow cytometry cell sorting and used for RNA-seq. Heat map of top 20 differentially expressed genes overexpressed in CD8 T cells purified from dLN compared to lesions.

Figure S2

Fig S2. Pimonidazole expression in *Leishmania*-infected skin.

C57BL/6 mice infected with *L. major* received pimonidazole one hour before euthanasia at two weeks. (a) Individual images from confocal microscopy (Fig. 2a). Nuclear (DAPI) staining (blue) and pimonidazole (green) in contralateral ears (naïve skin) and infected ears (lesions) for two weeks. Scale bar = 200 μm. (b) Individual images from confocal microscopy (Fig. 2b). Nuclear (DAPI) staining (blue) and pimonidazole (green) in contralateral ears (naïve skin) and infected ears (lesions) for two weeks. Scale bar = 100 μm.

Figure S3

Sample 1

Sample 1

Sample 1

Sample 2

1018
1019

1020   Fig S3. Neutrophil and pimonidazole expression in *Leishmania*-infected skin.
1021
C57BL/6 mice infected with *L. major* received pimonidazole one hour before euthanasia at two weeks. (a)
Individual images from confocal microscopy (Fig. 5c). Nuclear (DAPI) staining (blue), Ly6G (pink), and
pimonidazole (green) in the skin infected for two weeks. Scale bar = 200 μm. (b) Confocal microscopy image was
divided into 27 regions to measure the pixel intensity of Pimonidazole (green) and Ly6G (pink) shown in Fig. 5d.
Scale bar = 100 μm. (c) Individual images from confocal microscopy (Fig. 5e). Nuclear (DAPI) staining (blue),
Ly6G (pink), and pimonidazole (green) in the skin infected for two weeks. Scale bar = 100 μm.
1028