Refining the resolution of the yeast genotype-phenotype map using single-cell RNA-sequencing

Authors: Arnaud N'Guessan[1], Wen Yuan Tong[2,¥], Hamed Heydari[3,4], Alex N Nguyen Ba[1,2]

1. Department of Cell and Systems Biology, University of Toronto, Ramsay Wright Laboratories, 25 Harbord St, M5S3G5, Toronto, Ontario, Canada.

2. Department of Biology, University of Toronto at Mississauga, 3359 Mississauga Rd, L5L 1C5, Mississauga, Ontario, Canada

3. Department of Molecular Genetics, University of Toronto, Medical Science Building, Room 4386, 1 King's College Cir, M5S1A8, Toronto, Ontario, Canada.

4. Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, 160 College Street, Room 230, M5S3E1, Toronto, Ontario, Canada.

¥. Present address: Center of Molecular and Cellular Oncology, Yale University, 300 George Street Suite 6400, New Haven, CT

ABSTRACT

Genotype-phenotype mapping (GPM), or the association of trait variation to genetic variation, has been a long-lasting problem in biology. The existing approaches to this problem allowed researchers to partially understand within- and between-species variation as well as the emergence or evolution of phenotypes. However, traditional GPM methods typically ignore the transcriptome or have low statistical power due to challenges related to dataset scale. Thus, it is not clear to what extent selection modulates transcriptomes and whether cis- or trans-regulatory elements are more important. To overcome these challenges, we leveraged the cost efficiency and scalability of single-cell RNA sequencing (scRNA-seq) by collecting data from 18,233 yeast cells from 4,489 F2 segregants derived from an F1 cross between the laboratory strain BY4741 and the vineyard strain RM11-1a. More precisely, we performed eQTL mapping with the scRNA-seq data to identify single-cell eQTL (sc-eQTL) and transcriptome variation patterns associated with fitness variation inferred from the segregant bulk fitness assay. Due to the larger scale of our dataset and its multidimensionality, we could recapitulate results from decades of work in GPM from yeast bulk assays while revealing new associations between phenotypic and transcriptomic variations at a broad scale. We evaluated the strength of the association between phenotype variation and expression variation, revealed new hotspots of gene expression regulation associated with trait variation, revealed new gene functions with high expression heritability, and highlighted the larger aggregate effect of trans-regulation compared to cis-regulation. Altogether, these results suggest that integrating large-scale scRNA-seq data into GPM improves our understanding of trait variation in the context of transcriptomic regulation.

INTRODUCTION

The process by which DNA encodes proteins via transcription and translation has been studied for decades to make sense of organisms' phenotypes. However, being able to explain organisms' phenotypes from their genetic material, i.e. genotype-phenotype mapping (GPM), has been a long-lasting problem with important applications (1,2). Indeed, making sense of genetic variation at the phenotypic level enables the understanding of trait variation between and within species as well as the emergence and evolution of phenotypes (3). For instance, reverse genetics approaches, e.g. gene knockout or transgenic technologies, and forward genetics approaches like GWAS and QTL mapping helped in determining the function of multiple genes and the effects of mutations on growth in different environments (4). However, reverse genetics approaches typically fail to account for natural variation and forward genetics approaches like QTL mapping typically focus on genetic and phenotypic variation, so they cannot highlight selection on the transcriptome.

An essential characteristic of this problem is the multi-layered organization of the GPM. Indeed, GPM is not strictly restricted to the direct association between genotypes and phenotypes. This association is better resolved and complemented by understanding the intermediary transcriptome layer, e.g. cell mechanisms at the transcriptomic level are involved in diseases and pathogenicity (2,5–8). However, it is not clear to what extent transcriptomic changes relate to phenotypic changes or selection. Pioneering work from Mary-Claire King and Allan Charles Wilson set the tone for investigating this question by proposing that variations in morphological and behavioral traits arise more often through gene expression regulation than evolution at the protein-coding level (9). François Jacob then postulated an essay that stemmed from this theory in which he highlights how evolution acts as a tinkerer that works from already available material, i.e. through regulation of gene expression, to create new adaptations (10). This constituted the core of the evolutionary developmental biology, which matured into the still-debated claim that new adaptations mainly emerge through cis-regulation of gene expression, i.e. through noncoding DNA regulating a neighbor gene contrarily to trans-regulators acting on distant genes (11–14). This debate has been reinforced by the technical difficulties and complexity of assessing the evolution and outcome of mutations in non-coding regions (11,12). For example, although human genome-wide association studies (GWAS) loci are mostly identified in noncoding regions and have well-characterized cis-regulatory effects, their trans-regulatory effects are usually challenging to detect (15–17). Indeed, small effect sizes, multiple testing corrections and high false-positive rates for trans-regulation impose statistical and computational challenges for its detection (18,19). Advances in sequencing technologies helped to partially solve these issues, particularly in the context of transcriptome analyses of the model organism *Saccharomyces cerevisiae*. For instance, Brem et al (2002) used microarray technology to relate the gene expression profiles of 40 yeast segregants from a lab (BY) and natural vineyard strain (RM) to their genetic markers (20). They found that cis-acting modulation is the main mechanism for regulating gene expression. Nearly two decades later, by greatly increasing statistical power, Albert and collaborators (2018) found that most of the expression variation arises through trans-regulation using non-multiplexed RNA-seq to analyze 5,720 genes in 1,012 yeast segregants generated by a crossing between RM and BY (21). The analysis method they used, i.e. expression quantitative trait loci (eQTL) mapping,

consists in correlating allele frequencies to gene expression levels to find the loci modulating
expression.

Although eQTL mapping is a traditional GPM analysis that accounts for the transcriptomic
layer, it is typically realized through non-multiplexed RNA-seq which tends to have low statistical
power due to challenges with experimental scale and confounding factors (22,23). Thus, eQTL
mapping traditionally cannot identify significant low-effect regulatory mutations that are
important for understanding the genetic bases of complex traits and diseases (24,25). Furthermore,
most eQTL studies only assess the average transcriptomic profile of bulk populations without
being able to capture the profile of rare cell lineages within a population. This is a critical limitation
in heterogenous populations such as cancer or microbial populations where rare lineages can drive
relapse or drug resistance (26).

Here, we sought to circumvent the challenges of non-multiplexed bulk RNA-seq imposed
by the scale and population heterogeneity by performing eQTL mapping through single-cell RNA
sequencing (scRNA-seq) of a pool of ~4,500 well-characterized F2 segregants derived from a yeast
cross (21,27,28). In the same way that combinatorial indexing/barcoding and multiplexing enable
the collection of large-scale fitness and genotype data (29), we hypothesized that scRNA-seq could
help us collect both genotype and expression data on a large pool of segregants. We employ several
strategies to overcome previous obstacles of eQTL mapping studies: i) we pool cells from
thousands of F2 segregants during the growth step and perform a single scRNA-seq run on the
culture to account for environmental effects. The expression profiles of these lineages cannot be
captured cost-effectively with bulk RNA-seq as pooling segregants in a single bulk assay would
not capture individual strains' expression profiles. Moreover, ii) from the exome sequencing data
of single cells, we take advantage of the reference panel to validate that we accurately infer the
genotype of each cell from extremely low number of reads mapping to polymorphic sites per cell
(effectively ~0.2x coverage). Finally, iii) we leverage the presence of thousands of high-coverage
cells to train an unsupervised learning model to correct (denoise) and infer an imputed expression
profile of poorly covered cells.

Using this approach, we integrated the resulting transcriptomic data from growth in rich
media with a pre-existing yeast GPM to refine GPM features at a broad scale. For instance, we
estimated the heritability of the transcriptome and the extent to which the transcriptome is
associated with fitness. This allowed us to reveal that a negligible portion of expression variation
is related to the studied phenotype variation independently of genotypic variation. This implies
that almost all the phenotypic variation related to expression is also modulated by mutations. This
differentiates our work from previous bulk RNA-seq studies where expression is systematically
controlled for differences in lineage growth or phenotypes due to an assumption that a considerable
amount of phenotypic variation is related to expression independently of genomic mutations (21).
We also show that the increased scale of our scRNA-seq dataset enables single-cell eQTL mapping
(sc-eQTL) and identifies more hotspots of expression regulation containing previously identified
QTL. We also exploit the identified sc-eQTL to analyze the patterns of cis- and trans-regulation
in the GPM.

Our single-cell RNA-seq approach is consistent with yeast GPM results from non-multiplexed assays

We initially aimed to show that performing scRNA-seq at a large scale can generate data that are consistent with non-multiplexed DNA and RNA sequencing. For this purpose, we analyzed a dataset of thousands of yeast lineages generated by Nguyen Ba and collaborators (2022) (29). To understand the yeast GPM, they collected fitness and genotype data from ~100,000 F2 segregants derived from an F1 cross between a laboratory strain of yeast (BY) and a natural vineyard strain (RM) (**Figure 1A**).

Figure 1 F2 yeast segregants datasets. A) Reference panel and experiments from the barcoded bulk sequencing previously described in (29). The 99,995 F2 yeast segregants in the reference panel were derived from an F1 cross between a laboratory strain of yeast (BY) and a natural vineyard strain (RM). Thus, they only have 2 possible alleles at each of the 41,594 polymorphic sites. The lineage barcodes enabled fitness estimation from competition assays in 18 environments recapitulating the adaptation to temperature gradients, the ability to process different sources of carbon and the resistance to antifungal compounds. B) Pooled scRNA-seq dataset from a single batch. We performed scRNA-seq of the first batch of F2 segregants (n=4,489) to obtain genotypes that are similar to the reference panel and cell's expression profiles. The F2 segregant barcodes are typically not expressed, so single cells have their own cell barcode, which is included in all amplified reads of the same droplet. Non-covered sites, sequencing errors and the presence of reads in the wrong library (index swapping) are corrected for using the HMM described in Figure S1.

Using this approach named barcoded bulk QTL mapping or BB-QTL mapping, they revealed the
complex polygenic and pleiotropic nature of phenotypes as well as an unprecedented number of
pairwise epistatic interactions. To integrate transcriptomic data into this GPM, we performed
scRNA-seq using the 10X Genomics Chromium microfluidics platform (30). Other scRNA-seq
methods like Smart-seq2 offer a better breadth of coverage per cell (31). However, due to the
higher throughput or number of cells generated by 10x chromium and the different coverage
profiles across cells from the same F2 segregant, it is more advantageous to use 10x Chromium to
capture transcriptomic variation and reconstruct the reference genomes from single cells (31,32).
The throughput of 10x Genomics is typically in the order of 10^4 cells which limits us to a few
thousand lineages per library if we want to obtain a reasonable number of cells per lineage. This
is why we focus on a single batch of 4,489 F2 segregants. This method allowed us to obtain both
genotype and expression profiles from 18,233 cells of the first batch of F2 segregants (**Figure 1B**).
The F2 segregant barcodes are typically not expressed, so single cells have their own cell barcode,
which is included in all amplified reads of the same droplet. Although this short-read scRNA-seq
is a high-throughput approach (**Table 1**), it comes with challenges like gene dropouts and low
sequencing depth in some cells caused by technical artifacts of PCR amplification (25,26).

Metric	Value
Library size (number of reads)	479,781,081
Median number of covered genes per cell	1,311
Median number of UMIs per cell	5,158
Doublet rate	0.10
Average number of cell barcodes per F2 segregant	4.83
Polymorphic sites mean breadth of coverage	0.03

Table 1 scRNA-seq dataset features. The doublets were defined as scRNA-seq barcodes with
high coverage (>=third quartile) without significant genotype association to any F2 segregants and
for which the 2 most similar F2 segregants do not share genotypic similarity ($R^2 < 0.1$). The breadth
of coverage is defined as the proportion of BY/RM polymorphic sites covered in a single cell.

To overcome these challenges, the unique molecular identifiers (UMIs) of the 10X Genomics
platform provide a control for PCR amplification biases by quantifying gene expression from
unique transcribed molecule counts instead of sequencing read counts (31). However, UMIs still
do not correct for dropouts or missing data in the expression profile, which could obscure potential
associations between genotype and transcription (see below). We addressed this issue by fitting an
imputation model called DISCERN to the expression data (33). DISCERN is a neural network that
learns how to reconstruct the expression profile of high-coverage cells after embedding it in a
lower dimension. This eliminates the gene dropouts and denoises the expression profile. The fact
that the reconstructed expression is highly but not perfectly correlated to the original expression

(mean R^2 =50.0%, median R^2 =52.9%; R^2 s.d. = 10.1%) and the high variance explained by the
first principal component of the imputed expression (99.6%) is consistent with an effective
denoising. We take advantage of this denoised transcriptome for variance partitioning (as
described later).

In addition, Hidden Markov Models (HMMs) can infer accurate genotype data even at sequencing
depths as low as 0.1x (29). Nguyen Ba and collaborators (2022) designed an HMM to infer the
segregants genotypes from the observed reads at low depth of DNA sequencing by accounting for
sequencing error rate, recombination rate, and index swapping rate (29). As there are only two
ancestral lineages, there are only two possible alleles for the strains at each of the 41,594
polymorphic sites. Thus, the genotype of the segregants can be represented by the frequency of
only one of the parental alleles, which is RM in the dataset. Applying this model to low-coverage
segregants yielded genotypes that are significantly similar to high-coverage replicates (29). We
sought to use a similar model to infer genotypes from scRNA-seq data but we anticipated that
some of these parameters may differ due to increased error rate of the reverse-transcriptase,
increased index swapping due to pooled reaction, etc (**Figure S1**). In Nguyen Ba et al, those rates
were heuristically determined, but here we estimated these from the read mapping data and found
that re-estimated parameters from data increased the proportion of recovered strains in the single
cell data from 58.6% to 65.7%.

After adapting the HMM to the scRNA-seq data, we sought to validate that the resulting
cell genotypes relate well to their corresponding strain in the reference panel obtained by non-
multiplexed DNA sequencing strategies. Ideally, each single-cell barcode (from 10x Genomics
Chromium) should be associated with a single cell and a cell should have a clear match with a
unique strain in the reference panel. However, several factors can obscure these associations, e.g.
a single-cell droplet containing cells from 2 different strains, a low-coverage cell, uncertainty in
the allele of the reference genotype, etc. Thus, we designed an approach to assign cells to the
correct reference panel strain (see **Methods**). This approach relies on two metrics of similarity
between the cells and the strains' genotypes, i.e. the expected distance between them, which should
be minimized for the best match, and the relatedness (R^2). The statistical significance of the
relatedness between single cells and reference lineages was determined by a permutation test
(**Figure S2**). From the read mapping alone, we obtained a mean R^2 of 0.59 (σ = 0.19 and median
= 0.64), which was significantly improved after applying our HMM to correct for misidentified
alleles and imputing data in low-coverage sites using recombination probability. Indeed, the
single-cell HMM genotypes yield a mean R^2 of 0.73 (σ = 0.18 and median = 0.81; **Figure 2A**).
This approach was also successful in a recent eQTL mapping on segregants with a similar genetic
background (34). We found that the distribution of relatedness after HMM was still left-skewed,
with a minority of cells statistically significantly assigned to a reference genotype despite having
what appeared to be low relatedness. Upon investigation, it was found that these could be explained
by sequencing coverage and genotyping uncertainty in the single-cell and/or in the reference panel
genotype(**s**) (**Figure S3; Table S1**).

A

B

Figure 2 Single-cell RNA-seq data recapitulate bulk DNA and RNA assays results. A) Effect
of the HMM on the relatedness between single-cell genotypes and their closest reference lineage.
The single-cell original genotype represents the genotype of the cells before the correction with
the HMM. The relatedness to the closest lineage in batch 1 has been measured with the adjusted
R^2. To control for genotype uncertainty, only the 13,069 barcodes with a significant lineage

assignment (lineage-barcode genotype correlation FDR<0.05) and a reference lineage with lower uncertainty than the single cell HMM are selected, which represents 72.2% of the barcodes. We then rounded the genotypes to remove the uncertainty during the comparison. Wilcoxon signed test p-value is indicated above the violin plots. B) Narrow-sense heritability measured with scRNA-seq consensus genotypes and non-multiplexed DNA sequencing. Three datasets are compared through a t-test on the bootstrap narrow-sense heritabilities. The consensus genotypes of the batch 1 F2 segregants obtained from the associated scRNA-seq data are represented in blue, the batch 1 F2 segregant genotypes in the reference dataset (bulk sequencing) are represented in yellow and the whole reference dataset of F2 segregants is represented in grey. For each dataset, the narrow-sense heritability has been measured from 500 random subsamples of 1000 F2 segregants. The bars represent the mean narrow-sense heritability, and the error bar length represents the 95% CI. The results are illustrated for the 15 environmental conditions where a minority of batch 1 F2 segregants have missing fitness data. The 23C-30C represents the temperature for the competition assay in YPD media while the other phenotypes represent growth on YNB, molasses (mol), mannose (Mann) or raffinose (raff) and chemical resistance to copper sulfate (Cu), ethanol (eth), guanidinium chloride (gu), lithium acetate (Li), Sodium dodecyl sulfate (SDS) and suloctidil (suloc) (29).

To further establish that the genotyping obtained from scRNA-seq data was comparable to previous non-multiplexed genotyping of the reference genotype panel, we estimated the contribution of genetic variation to the phenotypic variation, i.e. fitness heritability. Nguyen Ba and collaborators (2022) estimated the narrow- and broad-sense heritabilities of complex phenotypes associated with temperature gradient, carbon source and chemical resistance for which RM and BY segregants exhibit a significant level of diversity (29). We used our lineage assignment to that panel to obtain fitness but used our single-cell genotyping to perform this association (**Methods**). Encouragingly, all GCTA-REML estimates of narrow-sense heritability in the scRNA-seq batch 1 dataset (blue in **Figure 2B**) are similar to the estimates from Nguyen Ba and collaborators (2022) whether or not we focus on batch 1 (yellow and grey rectangles in **Figure 2B**).

Although the variance partitioning is consistent with previous studies, it only provides a broad view of the genotype-phenotype map as it does not allow the identification of the loci that significantly explain phenotype variation. If the genotypes obtained by scRNA-seq were of high quality, then we would expect that a QTL mapping model from scRNA-seq would yield a model similar to that of non-multiplexed DNA sequencing data. To achieve this, we used a cross-validated stepwise forward linear regression on the strain fitness and consensus genotypes data from single cells that shared the same lineage assignment (**Methods**). Performing the QTL mapping on the batch 1 scRNA-seq dataset enabled the identification of 29 QTL compared to 31 QTL identified with the bulk barcoded approach (**Tables S2 and S3**) (29). These QTL were largely similar as shown by the non-significant difference between the effect sizes (Wilcoxon signed rank test $p = 0.29$) and by a model similarity metric (29) that considers the recombination distance between matched QTL, the similarity of the effect sizes and the allele frequencies (**Methods**).

Using this approach, we estimated that the similarity score between the batch 1 single cells QTL and the batch 1 BB-QTL is 86.2% while each model respectively had a similarity score of 78.7% and 78.2% with the full BB-QTL mapping performed on the 99,950 F2 segregants (29) (**Figure S4**). The QTL identified from the scRNA-seq dataset also recapitulated several important biological features of the reference panel such as an enrichment of non-synonymous and disordered region QTL (29) (**Figure S5**).

Integrating scRNA-seq data to an existing GPM highlights selection on the transcriptome

Having shown that our scRNA-seq allows us to relate the cells to the corresponding F2 segregant fitness data via the genotype relatedness, we next sought to highlight new associations within the BY/RM GPM. Selection is often highlighted at the genotype level through convergent evolution, an increase in allele frequency within a population, or population genetics metric (35–37). However, the central dogma of molecular biology and the evolution tinkering model both entail that phenotype variation should be linked to transcriptomic variation. As our dataset included all these variables, we sought to evaluate the expression-genotype association strength and provide a variance partitioning framework to evaluate the association between the transcriptome and trait variation (**Methods**). The GCTA-REML variance partitioning model used to estimate trait heritability in the previous section can also be modified to include gene expression as the response variable and cell genotypes as the only random effect (**Methods**). This enables the quantification of expression heritability, i.e. the variance of expression explained by genotype. Using this approach on our large-scale dataset, we estimated that genotype explains 75.7% of expression variance. This estimate is weighted for expression levels as genes that are more expressed account for more of the expression variation. At the individual gene level, the genotype explains, on average, 35.4% of transcript level variation (σ = 4.51%; min=31.0%; median = 33.4%; max=100%), which is slightly but significantly higher than previous estimates from non-multiplexed eQTL mapping studies (Wilcoxon signed rank p < 2.2e-16). Indeed, Albert and collaborators (2018) estimated that at the gene level, the genotype explains on average 30.0% of the expression variance using a lower-power dataset of 5,720 genes in 1,012 yeast segregants generated by the same parental strains (RM and BY).

In addition to expression heritability, our "one-pot" approach and the multidimensionality of our scRNA-seq dataset enable the evaluation of the association between traits and expression. This was previously challenging with large-scale bulk RNA-seq experiments as they require batching strategies. This results in batch effects, which are linked to environmental and technical variations that affect lineages' growth rates and expression. To eliminate this bias, Albert et al. (2018) controlled gene expression levels for variation in growth rates across lineages, making it impossible to correlate fitness to gene expression. Measuring the gene expression profiles of thousands of lineages simultaneously removes the need for batching strategies as it allows us to collect enough transcriptomes to understand the associations between different components of the genotype-phenotype map, which we demonstrate with the 30C phenotype (**Figure 3**).

Figure 3 Variance partitioning of the 30C phenotype from scRNA-seq. We fitted F2 segregant fitness data to their cells' consensus genotype and expression profiles using GTCA-GREML at different sample sizes. Because each segregant had multiple representatives/cells in the scRNA-seq data, the consensus profile of these lineages has been obtained from the cell with the highest transcript level. The bars represent the proportion of the fitness/phenotype variance across F2 segregants (whole rectangle area) explained by each model component. The size of the red genotype-exclusive component is obtained by subtracting the size of the shared component from the variance explained by the model in Equation 2. The red cell illustrates this effect as a mutation changes its fitness by modifying the structure of the transcript product. The size of the blue expression-exclusive component is obtained by subtracting the size of the shared component from the variance explained by the model in Equation 3. An epigenetic modification increasing the transcript level of the gene in the blue cell illustrates this effect. The purple area represents the effect of the interaction of mutation and expression on fitness variation observed across the segregants. In the example shown, a mutation in the promoter changes the fitness of the purple cell through a down-regulation of the gene expression. This component is obtained by subtracting the variance explained by the model in Equations 1 or 2 from the one in Equation 4, as there is a shared component of trait variation explained by genotype and expression variation. Finally, the black bar represents the model's residual, which includes any phenomenon that causes fitness variation that we did not explicitly measure and include in the model. The size of the residuals is obtained by subtracting the sum of the size of all the defined components (red + blue + purple) from 1.

The components of this variance partitioning all relate to at least one biological phenomenon. Indeed, the portion of trait variation explained exclusively by the genotype variation (red in **Figure 3**) represents the effect of mutations on fitness via several biological phenomena, such as protein stability, enzymatic function, etc, independent of expression level. For the 30C phenotype, this

component explains 20.9% of the fitness variation in the BY/RM background, which is lower than the 47.8% explained by the shared component between phenotype, genotype and expression variations (purple in **Figure 3**). The latter represents the association between selection (fitness) and the transcriptome either through loci influencing fitness via expression directly, e.g. a mutation in the promoter, or through loci affecting expression via an effect on cell fitness (indirectly) (38,39). Its considerable association with fitness variation thus supports the evolution and tinkering model for the 30C phenotype. As for the phenotype variation explained exclusively by gene expression (blue in **Figure 3**), it could represent epigenetics and stochastic gene expression, which weakly explain variations in the 30C phenotype (0.69%). Moreover, the total variance explained by the model is consistent across sample sizes, highlighting the robustness of the GREML method. However, increasing the sample size allowed an increase in the relative contribution of the genotype-expression interaction component at the expense of the other components, suggesting that increasing the scale of the dataset increases the confidence of the associations between expression and genotype changes that contribute to trait variation. We also note that denoising the expression data with DISCERN slightly increases the contribution of the genotype-expression interaction and decreases the residual of the model by up to 8% (**Figure S6**).

Although this model explains a large proportion of the 30C phenotype variance and its heritability, the residuals still represent ~30% of fitness variation (**Figure 3**). This could be explained by unmeasured factors like high-order epistasis. However, the broad-sense heritability of this phenotype is similar to the narrow-sense heritability, suggesting that the residuals are mostly not explained by genotype and expression (29). Nguyen Ba et al. (2022) also estimated that epistasis only explained around 5% of fitness (29). Other factors like mitochondrial mutations, post-transcriptional changes, protein-protein interactions, or other protein properties could play a role in explaining the residual of the model and are good candidates to extend this analysis. These results suggest that a single run of scRNA-seq on a single batch of F2 yeast segregants converges with bulk DNA sequencing results while refining broad-scale features of the GPM.

The increased scale of scRNA-seq enables the discovery of new eQTL

Our integrative scRNA-seq approach is not limited to enabling the quantification of the association between transcriptomic changes and trait variation. Indeed, the same approach we used to identify QTL can be used to detect loci regulating gene expression which can reveal the cell mechanisms underlying trait variation through transcriptomic changes. We thus modified the QTL mapping framework such that the response variable is the level of expression of a single gene in the single cells (**Methods**). Because our cells are not synchronized, differences in the cell cycle could create transcriptomic variation that could obscure the association with genotypic variation. To control for this, we averaged out the effect of cell cycle by reconstructing the consensus expression and genotype profiles of each segregant significantly associated to the cells. This approach is a cost-efficient way to perform eQTL mapping from the expression profile and genotype of cells from thousands of lineages in a multiplexed way (sc-eQTL mapping).

Consistent with yeast non-multiplexed eQTL results, the genes with the highest expression heritability are enriched in functions related to carbohydrate catabolic process (GO:0016052) and cellular biosynthetic process (translation GO:0006412, organelle assembly GO:0070925,

ribosome biogenesis GO:0042254 and gene expression GO:0010467) (Fisher's exact test
FDR<0.05; **Methods**). In both datasets, these genes are also highly expressed, which reflects the
positive correlation between expression heritability and expression levels ($R^2 = 0.81$ and $p < 2.2e$-
16). Although some of these highly expressed genes have important functions modulated by
mutations, their higher heritability could also be explained by higher statistical power to measure
heritability when expression counts are higher. Moreover, genes with the lowest expression
heritability observed in the RM/BY background, which we defined as the bottom 10% expression
heritability, are enriched in functions related to the cell cycle biological process (GO:0007049,
Fisher's exact test FDR<0.05) which is consistent with bulk RNA-seq eQTL (21,40). This does
not suggest that genes with this function are not important for variation in gene expression at the
whole-genome scale but rather that mutations observed in the RM/BY background are not
correlated with variation in the expression of these specific genes. This could reflect conservation
for genes involved in the cell cycle biological process (41,42).

Because of the increased scale of our collection, our approach is more powered to estimate
the gene expression heritability. Indeed, we detected a median of 21 eQTL per gene (μ=34.3,
σ=29.7) which is almost 4 times higher than what has been detected by the largest non-multiplexed
RNA-seq dataset in the same genetic background (21). This is expected as the power to detect trait
loci is positively correlated to the sample size (29). We validated this trend in our dataset by
performing a power analysis with the 10 most heritable genes (**Figure S7;** linear regression
adjusted $R^2 = 0.95$ and p<0.05), which had the following functions: glycolysis, gluconeogenesis,
pentose phosphate pathways, AMP metabolism and chaperone. These newly detected eQTL
probably have small effect sizes as they only allow to increase the expression heritability by 5%
on average. However, for the 3,730 out of 6,240 genes (59.8%) that have higher expression
heritability in our dataset, this increase is on average 15.7%. We were also able to detect new
overrepresented biological processes, i.e., the DNA metabolic process (GO:0006259) and the
response to nutrient levels (GO:0031667), for which the variation of expression levels is weakly
associated with the genetic variation observed across the F2 RM/BY segregants.

The functional enrichment analysis using scRNA-seq data revealed new associations
between expression heritability and biological processes in the RM/BY genetic background.
However, while it suggests that many eQTL are also QTL, it cannot accurately point to the specific
loci involved in trait variation and cannot address whether mutations on regulatory hubs have
stronger effects on traits. To investigate this, we mapped the QTL to hotspots of gene regulation
(or regulatory hubs), which we defined as 25 kb genomic windows that were repeatedly identified
in the eQTL mapping procedure (for different genes). This was done to acknowledge the
uncertainty in the exact position of the eQTL due to linkage disequilibrium and power. We then
ranked the 30C QTL identified by Nguyen Ba and collaborators (2022) based on their absolute
effect size and correlated it to the rank of the eQTL hotspots based on the number of regulated
genes. This resulted in a positive correlation (Spearman ρ = 0.33 and $p = 5.21e$-5), suggesting that
larger effects on the regulatory network translate into larger trait variation. Indeed, we observed
that some previously reported high-effect-size QTL genes are located in eQTL hotspots, e.g.
MKT1, HAP1, and IRA2 (**Figure 4A**).

Figure 4: eQTL features underlying trait variation across the BY/RM segregants. A) Mapping of the 30C QTL in the eQTL hotspots. We represent the hotspots of expression regulation as genomic windows (25 kb) to acknowledge the uncertainty around the real position of the eQTL

due to linkage disequilibrium. We annotated the 5 top eQTL hotspots and the eQTL hotspots in
which the top additive QTL identified by the BB-QTL mapping of the 30C phenotype are located.
In these regions, we represented the most affected trans-regulated genes in red, the most affected
cis-regulated gene in blue and the top QTL genes in black. The double quotation characters
represent the absence of such genes in the associated region. We also represented the rank of the
QTL in the set of 159 QTL of the 30C phenotype. B) Partitioning of the expression heritability or
explained variance (R^2) among cis- and trans-eQTL. Each pair of points connected by a line
represents a gene. Green lines represent the genes that only have trans-eQTL and orange lines
represent the genes that have both trans- and cis-eQTL. C) Comparison of the mean effect size
between cis- and trans-eQTL. Each pair of points connected by a line represents a gene. The line
color represents the ratio of the average effect size between cis- and trans-eQTL. The sample size
of each eQTL category is represented in the x-axis. This is the number of trans-eQTL and cis-
eQTL used for calculating the average effect sizes per gene not the number of points per
distribution.

Performing this rank test on individual genes also yielded the result that eQTL effect is correlated
with the fitness effect for 35.1% of the genes (permutation test $p < 0.05$, see **Methods**). Although
this correlation does not apply to most genes, it highlights potential regulatory mechanisms
explaining the importance of the strongest growth loci or QTL. For instance, MKT1, i.e. the
strongest growth loci, is part of a regulation hotspot affecting genes that are important for yeast
growth like ENP1 which is involved in RNA processing and HXT6 which is involved in glucose
uptake (43–45). Among the strongest growth loci, VPS70 is part of a hotspot of regulation that
strongly affects the expression of RSF2, a zinc-finger protein regulating glycerol-based growth
and respiration (46). Furthermore, the highest peaks for expression regulation contain important
growth loci on chromosome IV (13[th] and 23[rd] strongest QTL among 159 QTL; **Figure 4A**). This
hotspot is also responsible for regulating TDH3, which is involved in glycolysis and
gluconeogenesis, and can have an important effect on fitness (47). A recent eQTL mapping study
on segregants with a similar genetic background that also leverages one-pot scRNA-seq and
genotype inference also identified such hotspots of regulation and highlighted their role in the
trade-off between reproduction and cell cycle progression (34).

The presence of these hotspots suggests that expression differences in BY/RM would
predominantly be explained by mutations in trans-regulatory elements. To test this, we partitioned
the variation in gene expression between cis- and trans-regulatory loci for each gene (see
Methods). This analysis revealed that all the genes are affected by at least one polymorphic trans-
regulatory locus and that these polymorphic trans-regulatory loci explain most of that gene's
expression (**Figure 4B**). It is well known that mutations in promoters and nearby enhancers can
influence gene expression (49,50). Indeed, we identified many genes that contained an allele in a
cis-regulatory element that strongly explains that gene's expression variation (n=750 genes out of
6088, **Figure 4B**). As expected, mutations in cis-regulatory elements were of stronger effect size
than trans-eQTL individually, but the cumulative aggregate effect of all trans-eQTL acting on that
gene was comparable to the few cis-eQTL they had (**Figure 4C**). This can be explained by the fact
that there are more opportunities for mutations to arise in trans-regulatory elements. Finally, we
found that trans-eQTL have two times higher odds of affecting cell fitness than cis-eQTL (χ^2 p =

0.01). Furthermore, by repeating our eQTL hotspot analysis with Albert et al. (2018) data, we
observed a non-significant association between eQTL hotspot and QTL (χ^2 $p = 0.50$). In that bulk
RNA-seq dataset, QTL have less odds of being in eQTL hotspots compared to other loci (odd ratio
$= 0.72$) while in this scRNA-seq dataset, QTL odds of being in eQTL hotspots are two times higher
than other loci. This could be explained by the smaller sample size of that bulk RNA-seq dataset
and by the fact that they control for differences in growth across lineages. Taken together, these
results suggest that the link between the genetic basis of transcription variation across RM/BY
segregants and fitness could only be revealed by integrating large-scale transcriptomic data to an
existing GPM, which scRNA-seq facilitates.

CONCLUSION

By leveraging the scalability of scRNA-seq, we obtained thousands of transcriptomes from a
reference pool of strains in a single experiment. This enabled the analysis of the association
between genotype, transcriptome, and phenotype at an unprecedented scale. Questions
surrounding transcriptomic variation and phenotypic variation have been at the center of many
previous quantitative genetics studies (19–21,27,48,51–54). For instance, how strong is the
connection between the transcriptome and trait variation? Although most GWAS loci are located
at noncoding regions and are well-characterized for cis-regulation effects, is trans-regulation more
important for phenotypic and expression variation? What are the gene functions that are the most
heritable and modulable at the expression level? These ideas and discoveries all support the fact
that researchers can gain valuable insight into the evolution of traits by integrating the
transcriptome in GPM analyses, which can translate into fundamental knowledge or other
important applications where phenotypes evolve.

In this study, we took advantage of a previously characterized BY/RM cross where the genetic
basis of growth in various environments was examined in detail (29). By integrating transcriptomic
data in this genotype-phenotype map, we identified transcriptome components that are involved in
trait variation and evaluated their contribution. Similar to a previous study, which obtained
transcriptomes by individual strain sequencing, we found that gene expression is highly heritable.
Further, our study design also allowed us to conclude that gene expression contributes to a
significant portion of the phenotypic variation in this strain collection.

This finding is corroborated by our findings that most eQTL detected in our study were previously
shown to be QTL. This is perhaps not surprising given that QTL in this cross were previously
inferred to be in regulatory genes, but this provides a more mechanistic view of the effect of an
allele on phenotype. Indeed, we find a bias for trans-regulation for generating transcription
innovation where the cumulative effect of trans-eQTL on gene expression are significant. That is
not to say that cis-regulatory alleles are dispensable as cis-regulatory alleles often have a large
effect on gene expression. This is similar to studies on human transcriptomes where trans-eQTL
are cumulatively more impactful and enriched in sets of complex trait loci (19,52–54). This
genome-wide view of the genetic basis of transcriptional variation has consequences for the
evolution of phenotypes, as the target size afforded by trans-eQTL is far larger than cis-eQTL.

Thus, adaptation to small and fluctuating environmental changes may proceed preferentially
through allelic changes or recombination of many small-effect trans-eQTL, but large expression
changes are likely to require some cis-eQTL.

In this study, we leveraged the availability of genotype and phenotype data on our pool of strains.
This was obtained by liquid handling robotics and pooled competitive growth assay with barcode
sequencing. While this was performed on a very large scale, it was essentially obtained by brute-
force and through approaches that are not necessarily applicable to other systems. Although it is
clear from our results that single-cell sequencing can achieve the same genotype quality as single-
reaction genotyping, it is much harder to obtain phenotype data from scRNA-seq. Thus, our
framework might not be readily translatable to other systems where similar studies on the GPM
are desirable. However, two observations from this cross can be used to suggest an experimental
approach. First, while epistasis is important, it contributes to a relatively small portion of the
phenotypic variance. Second, transcriptomic variation contributes little to the missing heritability.
Thus, it may be possible to use predicted fitness instead of observed fitness and recapitulate
findings from this study. Indeed, the phenotype under study has a negligible component of its
variance explained by transcription independent of the genotype, but this feature was unknown
before the study. This alternative approach can only relate expression to the heritable component
of traits, which would have missed this variance explained by transcription independent of the
genotype and explains why we did not employ it. Predicted fitness could be obtained from bulk-
segregant analysis where the additive effect of loci can be inferred from whole-genome sequencing
(28,55). However, it is unclear whether the proposed experimental approach is generalizable in
other contexts, and collecting fitness data may not be necessary if a GPM analysis is already
available for the studied system.

Despite the study's limitation on generalizability, our scRNA-seq framework helps bridge the
understanding of how genetic variation influences transcriptomic variation at a broad scale. Our
framework relies on identifying the genome of single cells from the transcriptome, which is going
to be possible from low-coverage sequencing when genetic variation within the pool is high (such
as this cross, microbiome sequencing, or cancer cells with extensive copy number variation), and
from low cell diversity with sufficient transcriptomic variation such that aggregation of single-
cells with similar transcriptomes can afford pseudo-high coverage sequencing. Although we
restricted the scope of this work to broad-scale transcriptomic and genomic patterns related to trait
variation, we demonstrated that it is possible to integrate large-scale transcriptome data into an
existing GPM, which can be exploited in the future for more fine-scale analysis like co-expression
analysis and highlighting how specific trans-regulatory patterns within the cell network lead to
trait variation. For instance, identifying new eQTL could enable a better understanding of the gene
network by highlighting the downstream trans-regulatory effects of mutations and their association
with trait variation. For most of the genes, this newly discovered eQTL explained more than 15%
of their transcript level variation on average and could help characterize how cells react to
environmental changes. Moreover, the fact that it is an integrative approach allowed us to evaluate
the association between phenotypes, expression, and mutations. Although we did not validate the
strength of this association for multiple traits and biological systems, this approach is readily
available to help screen for traits that are evolving independently of eQTL if weak correlations

between phenotypic variation and genotype-expression interaction are observed, i.e. phenotypes
driven by epigenetics. Finally, the residuals of the model could help hypothesize on the
contribution of unmeasured phenomena like protein-protein interaction, gene content variation or
post-transcriptional modification, although it could be modest. Thus, integrating genotype,
transcriptome, and phenotype using scRNA-seq data can be particularly efficient for developing a
better understanding of other important traits or diseases.

MATERIAL AND METHODS

Yeast strains and segregants

We analyzed cells from a single batch (batch 1) of 4,489 F2 segregants obtained from a F1 cross between the yeast laboratory strain BY4741 and the vineyard strain RM11-1a generated in a previous study (29). These strains have been selected to generate this collection of segregants because they exhibit differences in multiple phenotypes including the adaptation to temperature, the ability to process different sources of carbon and the ability to resist antifungal compounds. Therefore, the genetic variation observed across the segregants can be correlated to the differences in growth rate observed in the 18 environments recapitulating these phenotypes in the Nguyen et al (2022) study (29). The selection of the batch is random and the fact that we performed the analyses on a single batch eliminates batch effects that could obscure variable associations. Genotypes and fitness data used were the same ones obtained in the previous study.

Yeast growth and single-cell RNA-sequencing protocol

To prepare strains for scRNA sequencing, we unfroze the batch of segregants and inoculated approximately $5*10^6$ cells in YPD (1% Yeast Extract, 2% Peptone, 2% Dextrose) to saturation. The next day, about 10^7 cells were passaged to 5 mL of fresh YPD and grew for 4 hours to bring cells to log-phase. We then pelleted 100 ul of cells and resuspended them in spheroplasting solution (5 mg/mL zymolyase 20T, 10 mM DTT, 1 M Sorbitol, 100 mM Sodium Phosphate pH 7.4) at a concentration of 10^7 cells/mL. The cells were incubated at 37 degrees Celcius for approximately 10 minutes at which point spheroplasting was verified by mixing a small aliquot of cells with detergent to observe lysis. The cells at this point were quantified using a hemocytometer and prepared using the standard 10x Genomics Gel Beads-in-emulsion (GEM) protocol. We used the Chromium Next GEM Single-cell 3' Reagent Kit to prepare the sequencing libraries and sequenced on a NextSeq 500 high-output flow cell.

We note that the cells analyzed here were grown in bulk and assayed for their transcriptome in log-phase. Our fitness data was obtained from competitive bulk fitness assays which include several whole growth cycles over multiple days and thus capture lag phase, exponential growth, and saturation. Nevertheless, previous experiments had shown that fitness was mostly determined by exponential growth which suggests that our analysis is adequate even if the cells were prepared for sequencing at a single time point.

Single-cell RNA-sequencing data parsing

From the scRNA-seq reads, we obtained gene expression levels and allele counts using the pipeline count from CellRanger version 3.1.0 (56). For each of the ancestral strains, i.e RM11-1a and BY4741, the pipeline mapped the scRNA-seq reads to the reference genome, filtered the barcodes by comparing the UMI count per barcode distribution to a background model of empty gel-bead in-emulsion, and counted the number of UMI per gene per barcode. The barcode filtering retained 18,233 barcodes. For each barcode, we then counted the number of RM and BY alleles at each polymorphic site by parsing the RM and BY bam files using a Python script (https://github.com/arnaud00013/sc-eQTL/tree/main/II_scRNA-seq_genotyping). This script only

573 keeps reads that mapped at the same loci on both reference genomes to increase the level of
574 confidence of the mapping.

Correction and imputation of single-cell genotypes with a Hidden Markov Model

Because there are only two possible alleles at each polymorphic sites of the F2 RM/BY segregants,
their genotype can be recapitulated by a quantitative variable measuring the proportion of reads
from one of the parental strains, which is RM in our dataset. The raw allele count data provides a
first estimate of this RM allele frequency at each polymorphic site. However, due to the low mean
depth of coverage of scRNA-seq data (0.2x), the absence of reads in some polymorphic sites and
the biases introduced during sequencing like index hopping/swapping, we expect that the raw data
can be imputed and corrected for errors and uncertainty in the observed alleles. Therefore, we
applied a Hidden Markov Model (HMM) on the observed allele count. Such model can infer
accurate genotype data at sequencing depths as low as 0.1x (29,31,57). Nguyen Ba and
collaborators (2022) designed an HMM to infer the segregants genotypes from bulk DNA
sequencing by accounting for sequencing error rate, recombination rate and index swapping rate
(29). Because scRNA-seq uses the reverse transcriptase, which has a higher error rate, and because
it is a pooled assay with higher chances of index swapping, we expected the HMM parameter to
differ for the single cell data. Therefore, we adapted the HMM to scRNA-seq data by measuring
its parameters in our dataset (**Figure S1**). The scripts are available on GitHub
(https://github.com/arnaud00013/sc-eQTL/tree/main/II_scRNA-seq_genotyping).

Assigning single cells to the reference panel strains

594 To evaluate the level of relatedness between the reference panel strains and the imputed single cell
595 genotypes, we used the expected distance to identify the strain that best relate to each single cell:

$$Expected\ distance(g_c, g_s) = \sum_{i=1}^{41594} g_c + g_s - 2g_c g_s \qquad (Eq.1)$$

597 where g_c is the cell genotype and g_s is the strain genotype. Next, we assigned the single cell to its
598 best match in the studied batch of 4,489 trains only if this match is better than the best match in
599 randomly generated batches of the same size (**Figure S2**). This procedure is implemented and
600 available at https://github.com/arnaud00013/sc-eQTL/tree/main/III_Genotype_analysis).

Partitioning the phenotypic variance into genetic and transcriptomic components

602 To analyze the yeast GPM at a broad scale and to evaluate the association between selection and
603 the transcriptome, we estimated the contribution of genetic and transcriptomic variations to
604 phenotypic variation from scRNA-seq data. More precisely, we performed a Genome-wide
605 Complex Trait Analysis (GCTA) by fitting a linear mixed model to the data using the restricted
606 maximum-likelihood (REML) method (58):

$$y = X\beta + W_g u_g + \varepsilon_g \qquad (Eq.2)$$

$$y = X\beta + W_e u_e + \varepsilon_e \qquad (Eq.3)$$

$$y = X\beta + W_g u_g + W_e u_e + \varepsilon \qquad\qquad \text{(Eq.4)}$$

where y is the fitness vector for the n segregants with a significant association to batch 1 cells, X is the nxk matrix of k fixed effects, β is the vector of k coefficients of the fixed effects, W_g is the nxp genotype matrix, u_g is the vector of p SNP effects, W_e is the nxm expression matrix, u_e is the vector of m gene expression effects and ε is the error term. To decrease the effect of gene dropouts and noise in the gene expression profile of each cell, we fitted an imputation model called DISCERN to the expression data (33). DISCERN is an autoencoder that learns how to reconstruct the expression profile of high-coverage cells after embedding it in a lower dimension. These operations eliminate the gene dropouts and denoises the expression profile. We also controlled for the effect of the cell cycle on gene expression by using the consensus F2 segregants expression and genotypes from the associated cell data. The consensus data has been obtained using the cell with the maximum read count. Because the dataset does not include fixed effects, we set the fixed effect to a vector of ones such that its coefficients represent the mean fitness while the genotype and expression data are the random effects that explain the fitness variance along with the error terms. The REML solution assumes that the data follow a Gaussian distribution, so the data are standardized before fitting the model. We also divided the standardized expression counts by the cell sum of expression counts to control for molecule count biases across cells. The cell fitness is based on the fitness of the closest segregant in batch 1 as measured by the expected distance. Because this model is linear and additive, it can be compared to the estimates of narrow-sense heritability obtained by Nguyen Ba and collaborators (2022) (29). The difference between the variance explained in equation 4 and equations 2 or 3 allow to infer the variance explained only by the genotype or the expression component of the model. The code for the variance partitioning is available on GitHub (https://github.com/arnaud00013/sc-eQTL/tree/main/IV_variance_partitioning).

Estimating the expression heritability from scRNA-seq

To obtain this estimate from scRNA-seq data, we considered the fact that GCTA-REML only takes a vector as a response variable while the gene expression matrix is multi-dimensional. To solve this, we orthogonalized the gene expression matrix using principal component analysis (PCA), and used its first PC, which explains 99.6% of the variance, as a response variable of the model. In case 99% of the expression matrix is explained by more than one PC, it is possible to use a weighted sum to obtain the expression heritability. Indeed, if the expression PCs recapitulate the total expression variance and are orthogonal or independent to each other, then the sum of the PCs variance explained by genotype should be the expression heritability. If "k" is the number of PCs explaining at least 99% of expression variance:

$$Expression\ heritability = \sum_{i=1}^{k} PC_i\ eigen\ value * "PC_i \sim genotype"\ model\ R^2 \qquad \text{(Eq.5)}$$

QTL mapping

To identify the loci that influence cell fitness, we performed a linear regression on the consensus genotypes of the strains from the scRNA-seq data and the strain fitness. We decided to use the consensus genotypes of the strains as they relate better to the bulk segregant genomes. To build the consensus genotypes, we defined cells from the same lineage as the ones that shared the same closest F2 segregant in batch 1. Next, we used the median to obtain cells' consensus genotypes as it is less sensitive to outliers and because it yields the best relatedness to the batch 1 reference genotypes (median R^2 = 87.0%; μ=79.5%; σ=18.2; **Figure S4**). We selected the QTL in the linear models using cross-validation on the scRNA-seq data. This analysis consists in dividing the dataset into 10 random partitions of similar sample sizes and running a cross-validated stepwise forward linear regression on each partition. For each partition, the model starts with no QTL and a linear model "Fitness ~ Genotype" is fitted using the genotype data at each polymorphic site, where the correlation coefficient represents the effect size of the SNP. Then, the forward search starts and at each iteration, a new locus with the minimum linear model residual sum of squares (RSS) is added to the QTL model, which is updated with new effect sizes after the addition of a new SNP. Because the order of addition of QTL matters in the forward search and because some QTL are linked or collinear, the model can be refined by exploring different QTL around the local optima. These steps are repeated until the model RSS cannot be improved anymore or until the number of QTL reaches an arbitrary maximum far from the cross-validated number of QTL. After the forward search is completed in each partition, the algorithm calculates the optimal λ values that minimizes the objective function F_o:

$$F_o(\beta) = RSS(\beta) + Lasso\ penalty(\beta)$$

$$\|Y - X\beta\|_2^2 + \lambda\|\beta\|_0 \text{ (Eq.6)}$$

where β is the vector of SNP effect sizes in the QTL model, $\|Y - X\beta\|_2^2$ is the RSS of the linear QTL model, λ defines the penalty for adding a new SNP to the model and $\|\beta\|_0$ is the number of SNPs in the QTL model. This objective function has the property to add sparsity in the QTL model and thus avoid overestimating the number of QTL while being consistent (29). The optimal λ has a minimum of log(n) which corresponds to the Bayesian Information Criterion (BIC), which is known to yield correct models asymptotically (59). This allows to consider the possibility that a sparser model than the one found using the BIC could yield better predictive power on a test set while avoiding overfitting. The optimal λ values found in all the partitions are then averaged and the resulting mean λ is used to solve the objective function in the full dataset, which yields the optimal QTL model. The cross-validation assumes that the partitions are independent, such that the variance explained by the model and the number of relevant QTL are unbiased estimates.

Highlighting hotspots of gene regulation through eQTL mapping

To identify the loci regulating gene expression regulation, we adapted the QTL mapping framework using expression as the predicted phenotype. Because this approach had to be repeated for each of the 6,240 genes, we needed to modify it so that the execution time was convenient. To do so, the parameter λ was not estimated using cross-validation but rather from the Bayesian Inference Criterion (BIC), i.e. λ = log(n) where n is the number of cells. We found that the BIC

was often selected by the cross-validation procedure when tested on a few genes and thus we do
not believe that this approach will significantly change our results.

To acknowledge the uncertainty around the exact position of eQTL due to linkage disequilibrium,
we define eQTL hotspots as 25 kb genomic windows that were repeatedly identified in the eQTL
mapping procedure. Finally, we defined hotspots of gene regulation as genomic windows in the
fourth quantile of the distribution of the number of regulated genes per window and the fourth
quantile of the number of eQTL per window. The code for the single cell eQTL mapping is
available on GitHub (https://github.com/arnaud00013/sc-eQTL/tree/main/V_sc_eQTL_mapping).

Functional enrichment analysis by gene ontology annotation

To highlight gene functions enriched at different levels of expression or expression heritability,
we performed the panther database binomial test for statistical overrepresentation of gene ontology
biological processes (40,60). A low level was defined as within the 25% bottom part of the
distribution (<Q1) while a high level was defined as within the top 25% part of the distribution
(>Q3). The *p*-values were corrected for multiple testing using the false discovery rate correction
(FDR).

Matching QTL to eQTL

To evaluate the contribution of gene expression regulation to fitness variation, we created a model
to match QTL and eQTL based on the similarity of loci and the similarity of predicted effect on
gene expression. More precisely, for each of the 6,088 genes for which we could detect eQTL, we
performed a new eQTL model by correlating the expression level of the gene to the genetic
variation at QTL positions. This allowed us to measure the predicted effect of the QTL on gene
expression. We then calculated the distance between the QTL and the real eQTL of the gene based
on recombination distance within each chromosome, which decreases exponentially with genetic
distance, and the difference in the predicted effect on the gene expression using the formulation
developed by Nguyen Ba et al (2022) (29). Next, we used the same Needleman-Wunsch algorithm
to find the most likely set of pairing between QTL and eQTL, where an unmatched QTL is also
possible but penalized. Finally, we determined the proportion of genes for which gene expression
regulation is associated with higher fitness. To do so, for each gene, we performed a permutation
test by comparing the average rank of the matched QTL of the gene to the average rank of 999
random subsets of unmatched QTL of the same size. The p-value is the proportion of random
subsets of unmatched QTL with a higher average QTL rank than the set of matched QTL.

Comparing cis- and trans-eQTL contribution to expression variation

719 We used the definition of local eQTL in Albert et al. (2018) to define cis-eQTL, i.e. any eQTL
720 between 1,000 bp upstream of the gene and 200 bp downstream of the gene. Thus, we defined
721 trans-eQTL as the eQTL that do not follow this criterion. For each gene, we then performed
722 variance partitioning using the GCTA:

$$y = X\beta + W_{g\_cis}u_{g\_cis} + \varepsilon_{cis} \tag{Eq.7}$$

$$y = X\beta + W_{g\_trans}u_{g\_trans} + \varepsilon_{trans} \tag{Eq.8}$$

$$725 \qquad\qquad y = X\beta + W_{g\_cis}u_{g\_cis} + W_{g\_trans}u_{g\_trans} + \varepsilon \qquad\qquad\qquad \text{(Eq.9)}$$

where y is the vector of expression level of the gene across the n cells, X is the nxk matrix of k
fixed effects, β is the vector of k coefficients of the fixed effects, W_{g_cis} is the nxp cis-eQTL
genotype matrix, u_{g_cis} is the vector of p cis-eQTL effects on expression, W_{g_trans} is the nxm
trans-eQTL expression matrix, u_e is the vector of m trans-eQTL effects on expression and ε
represent the error terms. Because the dataset does not include fixed effects, we set the fixed effect
to a vector of ones such that its coefficients represent the mean expression level while the cis-
eQTL and trans-eQTL genotypes are the random effects that explain the expression variance along
with the error terms. We can infer the variance explained by the cis-eQTL by the difference in
variance explained between the models in equations 9 and 8. Likewise, the difference of variance
explained by the models in equations 9 and 7 can help us estimate the variance explained by the
trans-eQTL. Finally, we estimate the effect sizes using the absolute value of the correlation
coefficients of each loci (**Figure S8**) and compare the median between the cis- and trans-eQTL
from the same gene (paired data) with a Wilcoxon signed rank test.

DATA AVAILABILITY

The code used for this study is available and explained at https://github.com/arnaud00013/sc-eQTL and the original single-cell reads from the pooled segregants scRNA-seq assay have been uploaded in the NCBI BioProject database with the accession number PRJNA1022775. The single-cell barcodes expression data are also available at https://github.com/arnaud00013/sc-eQTL as an archive file named Matrix_gene_expression_barcodes_1_to_9000.csv.tar.gz or Matrix_gene_expression_barcodes_9001_to_18233.csv.tar.gz.

ACKNOWLEDGEMENTS

ANNB acknowledges support from the Natural Science and Engineering Research Council of Canada (NSERC RGPIN-2021–02716 and DGECR-2021–00117) and AN acknowledges support from the Natural Science and Engineering Research Council (NSERC CGS-D App Id: 569340-2022, UTF Fellowship from the University of Toronto and Rustom H. Dasture Graduate Scholarship in Cell and Systems Biology). The computations in this paper were enabled by resources provided by Compute Canada. We also thank the Centre for the Analysis of Genome Evolution and Function (CAGEF) and TCAG at SickKids for sequencing support.

REFERENCES

1. Bartoli C, Roux F. Genome-Wide Association Studies In Plant Pathosystems: Toward an Ecological Genomics Approach. Front Plant Sci [Internet]. 2017 May 23;8. Available from: ://WOS:000402030200001

2. Ferreira MA, Gamazon ER, Al-Ejeh F, Aittomaki K, Andrulis IL, Anton-Culver H, et al. Genome-wide association and transcriptome studies identify target genes and risk loci for breast cancer. Nat Commun [Internet]. 2019 Apr 15;10. Available from: ://WOS:000464494100010

3. Aguet F, Alasoo K, Li YI, Battle A, Im HK, Montgomery SB, et al. Molecular quantitative trait loci. Nat Rev Methods Primer. 2023 Jan 25;3(1):1–22.

4. Tarantino LM, Eisener-Dorman AF. Forward Genetic Approaches to Understanding Complex Behaviors. Curr Top Behav Neurosci. 2012;12:25–58.

5. Casamassimi A, Federico A, Rienzo M, Esposito S, Ciccodicola A. Transcriptome Profiling in Human Diseases: New Advances and Perspectives. Int J Mol Sci. 2017 Jul 29;18(8):1652.

6. Wainberg M, Sinnott-Armstrong N, Mancuso N, Barbeira AN, Knowles DA, Golan D, et al. Opportunities and challenges for transcriptome-wide association studies. Nat Genet. 2019 Apr;51(4):592–9.

7. Transcriptome: Connecting the Genome to Gene Function | Learn Science at Scitable [Internet]. [cited 2023 Aug 31]. Available from: https://www.nature.com/scitable/topicpage/transcriptome-connecting-the-genome-to-gene-function-605/

8. Williams CG, Lee HJ, Asatsuma T, Vento-Tormo R, Haque A. An introduction to spatial transcriptomics for biomedical research. Genome Med. 2022 Jun 27;14(1):68.

9. King MC, Wilson AC. Evolution at Two Levels in Humans and Chimpanzees. Science. 1975 Apr 11;188(4184):107–16.

10. Jacob F. Evolution and Tinkering. Science. 1977 Jun 10;196(4295):1161–6.

11. Hoekstra HE, Coyne JA. The Locus of Evolution: Evo Devo and the Genetics of Adaptation. Evolution. 2007;61(5):995–1016.

12. Kratochwil CF, Meyer A. Evolution: Tinkering within Gene Regulatory Landscapes. Curr Biol. 2015 Mar 30;25(7):R285–8.

13. Primig M, Williams RM, Winzeler EA, Tevzadze GG, Conway AR, Hwang SY, et al. The core meiotic transcriptome in budding yeasts. Nat Genet. 2000 Dec;26(4):415–23.

14. Cavalieri D, Townsend JP, Hartl DL. Manifold anomalies in gene expression in a vineyard isolate of Saccharomyces cerevisiae revealed by DNA microarray analysis. Proc Natl Acad Sci U S A. 2000 Oct 24;97(22):12369–74.

15. Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al. Systematic Localization of Common Disease-Associated Variation in Regulatory DNA. Science. 2012 Sep 7;337(6099):1190–5.

16. Schaub MA, Boyle AP, Kundaje A, Batzoglou S, Snyder M. Linking disease associations with
regulatory information in the human genome. Genome Res. 2012 Sep;22(9):1748–59.

17. Nicolae DL, Gamazon E, Zhang W, Duan S, Dolan ME, Cox NJ. Trait-Associated SNPs Are More Likely
to Be eQTLs: Annotation to Enhance Discovery from GWAS. PLoS Genet. 2010 Apr 1;6(4):e1000888.

18. Dutta D, He Y, Saha A, Arvanitis M, Battle A, Chatterjee N. Aggregative trans-eQTL analysis detects
trait-specific target gene sets in whole blood. Nat Commun. 2022 Jul 26;13(1):4323.

19. Wang L, Babushkin N, Liu Z, Liu X. Trans-eQTL mapping in gene sets identifies network effects of
genetic variants. Cell Genomics. 2024 Apr 1;4(4):100538.

20. Brem RB, Yvert G, Clinton R, Kruglyak L. Genetic Dissection of Transcriptional Regulation in Budding
Yeast. Science [Internet]. 2002 Apr 26 [cited 2022 Apr 29]; Available from:
https://www.science.org/doi/pdf/10.1126/science.1069516

21. Albert FW, Bloom JS, Siegel J, Day L, Kruglyak L. Genetics of trans-regulatory variation in gene
expression. Wittkopp PJ, editor. eLife. 2018 Jul 17;7:e35471.

22. Schwarz T, Boltz T, Hou K, Bot M, Duan C, Loohuis LO, et al. Powerful eQTL mapping through low-
coverage RNA sequencing. Hum Genet Genomics Adv. 2022 Jul 14;3(3):100103.

23. Fan Y, Zhu H, Song Y, Peng Q, Zhou X. Efficient and effective control of confounding in eQTL mapping
studies through joint differential expression and Mendelian randomization analyses. Bioinformatics.
2020 Aug 13;37(3):296–302.

24. Bush WS, Moore JH. Chapter 11: Genome-Wide Association Studies. Plos Comput Biol [Internet].
2012 Dec;8(12). Available from: ://WOS:000312901500028

25. Lorenz K, Cohen BA. Small- and Large-Effect Quantitative Trait Locus Interactions Underlie Variation
in Yeast Sporulation Efficiency. Genetics. 2012 Nov;192(3):1123-+.

26. Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, et al. Rare cell variability and
drug- induced reprogramming as a mode of cancer drug resistance. Nature. 2017 Jun
15;546(7658):431-+.

27. Bloom JS, Ehrenreich IM, Loo WT, Lite TLV, Kruglyak L. Finding the sources of missing heritability in a
yeast cross. Nature. 2013 Feb;494(7436):234–7.

28. Ehrenreich IM, Torabi N, Jia Y, Kent J, Martis S, Shapiro JA, et al. Dissection of genetically complex
traits with extremely large pools of yeast segregants. Nature. 2010 Apr;464(7291):1039–42.

29. Nguyen Ba AN, Lawrence KR, Rego-Costa A, Gopalakrishnan S, Temko D, Michor F, et al. Barcoded
Bulk QTL mapping reveals highly polygenic and epistatic architecture of complex traits in yeast.
Verstrepen KJ, editor. eLife. 2022 février;11:e73983.

30. Vermeersch L, Jariani A, Helsen J, Heineike BM, Verstrepen KJ. Single-Cell RNA Sequencing in
YeastYeast Using the 10× Genomics Chromium Device. In: Devaux F, editor. Yeast Functional

Genomics: Methods and Protocols [Internet]. New York, NY: Springer US; 2022 [cited 2024 Jun 16].
p. 3–20. Available from: https://doi.org/10.1007/978-1-0716-2257-5_1

31. Hwang B, Lee JH, Bang D. Single-cell RNA sequencing technologies and bioinformatics pipelines. Exp
Mol Med [Internet]. 2018 Aug 7;50. Available from: ://WOS:000441266700002

32. Jariani A, Vermeersch L, Cerulus B, Perez-Samper G, Voordeckers K, Van Brussel T, et al. A new
protocol for single-cell RNA-seq reveals stochastic gene expression during lag phase in budding
yeast. Rokas A, Barkai N, editors. eLife. 2020 May 18;9:e55320.

33. Hausmann F, Ergen C, Khatri R, Marouf M, Hänzelmann S, Gagliani N, et al. DISCERN: deep single-cell
expression reconstruction for improved cell clustering and cell subtype and state detection.
Genome Biol. 2023 Sep 20;24(1):212.

34. Boocock J, Alexander N, Alamo Tapia L, Walter-McNeill L, Patel SP, Munugala C, et al. Single-cell
eQTL mapping in yeast reveals a tradeoff between growth and reproduction. Brem R, Moses AM,
editors. eLife. 2025 Mar 12;13:RP95566.

35. Bergström A, Simpson JT, Salinas F, Barré B, Parts L, Zia A, et al. A High-Definition View of Functional
Genetic Variation from Natural Yeast Genomes. Mol Biol Evol. 2014 Apr 1;31(4):872–88.

36. Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. The dynamics of molecular evolution over
60,000 generations. Nature. 2017/10/19 ed. 2017 Nov 2;551(7678):45–50.

37. Johnson MS, Gopalakrishnan S, Goyal J, Dillingham ME, Bakerlee CW, Humphrey PT, et al.
Phenotypic and molecular evolution across 10,000 generations in laboratory budding yeast
populations. Verstrepen KJ, Wittkopp PJ, Verstrepen KJ, Hodgins-Davis A, editors. eLife. 2021 Jan
19;10:e63910.

38. Sun XM, Bowman A, Priestman M, Bertaux F, Martinez-Segura A, Tang W, et al. Size-Dependent
Increase in RNA Polymerase II Initiation Rates Mediates Gene Expression Scaling with Cell Size. Curr
Biol. 2020 Apr 6;30(7):1217-1230.e7.

39. Marguerat S, Bähler J. Coordinating genome expression with cell size. Trends Genet. 2012 Nov
1;28(11):560–5.

40. Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, et al. Protocol Update for large-scale
genome and gene function analysis with the PANTHER classification system (v.14.0). Nat Protoc.
2019 Mar;14(3):703–21.

41. Bähler J. Cell-cycle control of gene expression in budding and fission yeast. Annu Rev Genet.
2005;39:69–94.

42. Yu L, Castillo LP, Mnaimneh S, Hughes TR, Brown GW. A Survey of Essential Gene Function in the
Yeast Cell Division Cycle. Mol Biol Cell. 2006 Nov;17(11):4736–47.

43. Roos J, Luz JM, Centoducati S, Sternglanz R, Lennarz WJ. ENP1, an essential gene encoding a nuclear
protein that is highly conserved from yeast to humans. Gene. 1997 Jan 31;185(1):137–46.

44. Chen W, Bucaria J, Band DA, Sutton A, Sternglanz R. Enp1, a yeast protein associated with U3 and
U14 snoRNAs, is required for pre-rRNA processing and 40S subunit synthesis. Nucleic Acids Res.
2003 Jan 15;31(2):690–9.

45. Roy A, Dement AD, Cho KH, Kim JH. Assessing Glucose Uptake through the Yeast Hexose Transporter
1 (Hxt1). PLOS ONE. 2015 Mar 27;10(3):e0121985.

46. Lu L, Roberts GG, Oszust C, Hudson AP. The YJR127C/ZMS1 gene product is involved in glycerol-
based respiratory growth of the yeast Saccharomyces cerevisiae. Curr Genet. 2005 Oct 1;48(4):235–
46.

47. Vande Zande P, Hill MS, Wittkopp PJ. Pleiotropic effects of trans-regulatory mutations on fitness and
gene expression. Science. 2022 Jul;377(6601):105–9.

48. Brem RB, Storey JD, Whittle J, Kruglyak L. Genetic interactions between polymorphisms that affect
gene expression in yeast. Nature. 2005 Aug;436(7051):701–3.

49. Mattioli K, Oliveros W, Gerhardinger C, Andergassen D, Maass PG, Rinn JL, et al. Cis and trans effects
differentially contribute to the evolution of promoters and enhancers. Genome Biol. 2020 Aug
20;21(1):210.

50. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution of gene
regulation. Nat Rev Genet. 2012 Jun 18;13(7):505–16.

51. Bloom JS, Boocock J, Treusch S, Sadhu MJ, Day L, Oates-Barker H, et al. Rare variants contribute
disproportionately to quantitative trait variation in yeast. Landry CR, Barkai N, editors. eLife. 2019
Oct 24;8:e49212.

52. Westra HJ, Peters MJ, Esko T, Yaghootkar H, Schurmann C, Kettunen J, et al. Systematic
identification of trans-eQTLs as putative drivers of known disease associations. Nat Genet. 2013
Oct;45(10):1238–43.

53. Yao C, Joehanes R, Johnson AD, Huan T, Liu C, Freedman JE, et al. Dynamic Role of trans Regulation
of Gene Expression in Relation to Complex Traits. Am J Hum Genet. 2017 Apr 6;100(4):571–80.

54. Liu X, Li YI, Pritchard JK. Trans Effects on Gene Expression Can Drive Omnigenic Inheritance. Cell.
2019 May 2;177(4):1022-1034.e6.

55. Brauer MJ, Christianson CM, Pai DA, Dunham MJ. Mapping novel traits by array-assisted bulk
segregant analysis in Saccharomyces cerevisiae. Genetics. 2006 Jul;173(3):1813–6.

56. Zheng GX, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel digital
transcriptional profiling of single cells. Nat Commun. 2017/01/17 ed. 2017 Jan 16;8:14049.

57. Stark R, Grzelak M, Hadfield J. RNA sequencing: the teenage years. Nat Rev Genet. 2019
Nov;20(11):631–56.

58. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. Am
J Hum Genet. 2010/12/21 ed. 2011 Jan 7;88(1):76–82.

59. Harrell FE. Regression Modeling Strategies: With Applications to Linear Models, Logistic Regression,
and Survival Analysis [Internet]. New York, NY: Springer; 2001 [cited 2023 Mar 29]. (Springer Series
in Statistics). Available from: http://link.springer.com/10.1007/978-1-4757-3462-1

60. PANTHER: Making genome-scale phylogenetics accessible to all [Internet]. [cited 2023 Sep 29].
Available from: https://onlinelibrary.wiley.com/doi/epdf/10.1002/pro.4218

900