Refining the resolution of the yeast genotype-phenotype map using single-cell RNA-sequencing

Authors: Arnaud N'Guessan[1], Wen Yuan Tong[2,¥], Hamed Heydari[3,4], Alex N Nguyen Ba[1,2]

1. Department of Cell and Systems Biology, University of Toronto, Ramsay Wright Laboratories, 25 Harbord St, M5S3G5, Toronto, Ontario, Canada.

2. Department of Biology, University of Toronto at Mississauga, 3359 Mississauga Rd, L5L 1C5, Mississauga, Ontario, Canada

3. Department of Molecular Genetics, University of Toronto, Medical Science Building, Room 4386, 1 King's College Cir, M5S1A8, Toronto, Ontario, Canada.

4. Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, 160 College Street, Room 230, M5S3E1, Toronto, Ontario, Canada.

¥. Present address: Center of Molecular and Cellular Oncology, Yale University, 300 George Street Suite 6400, New Haven, CT

ABSTRACT

Genotype-phenotype mapping (GPM) or the association of trait variation to genetic variation has been a long-lasting problem in biology. The existing approaches to this problem allowed researchers to partially understand within- and between-species variation as well as the emergence or evolution of phenotypes. However, traditional GPM methods typically ignore the transcriptome or have low statistical power due to challenges related to dataset scale. Thus, it is not clear to what extent selection modulates transcriptomes and whether cis- or trans-regulatory elements are more important. To overcome these challenges, we leveraged the cost efficiency and scalability of single-cell RNA sequencing (scRNA-seq) by collecting data from 18,233 yeast cells from 4,489 segregants of a cross between the laboratory strain BY4741 and the vineyard strain RM11-1a. More precisely, we performed eQTL mapping with the scRNA-seq data to identify single-cell eQTL (sc-eQTL) and transcriptome variation patterns associated to fitness variation inferred from the segregants' bulk fitness assay. Due to the larger scale of our dataset, we were able to recapitulate results from decades of work in GPM from yeast bulk assays while revealing new associations between phenotypic and transcriptomic variations. The multidimensionality of this dataset also allowed us to measure phenotype and expression heritability and partition the variance of cell fitness into genotype and expression components to highlight selective pressure at both levels. Altogether these results suggest that integrating large-scale scRNA-seq data into GPM improves our understanding of trait variation in the context of transcriptomic regulation.

INTRODUCTION

The process by which DNA encodes proteins via transcription and translation has been studied for decades to make sense of organisms' phenotypes. However, being able to explain organisms' phenotypes from their genetic material, i.e. genotype-phenotype mapping (GPM), has been a long-lasting problem with important applications (1,2). Indeed, making sense of genetic variation at the phenotypic level enables the understanding of trait variation between and within species as well as the emergence and evolution of phenotypes (3). For instance, reverse genetics approaches, e.g. gene knockout or transgenic technologies, and forward genetics approaches like GWAS and QTL mapping helped in determining the function of multiple genes and the effects of mutations on growth in different environments (4). However, reverse genetics approaches typically fail to account for natural variation and forward genetics approaches like QTL mapping typically focus on genetic and phenotypic variation so they cannot highlight selection on the transcriptome.

An essential characteristic of this problem is the multi-layered organization of the GPM. Indeed, GPM is not strictly restricted to the direct association between genotypes and phenotypes. This association is better resolved and complemented by understanding the intermediary transcriptome layer, e.g. cell mechanisms at the transcriptomic level are involved in diseases and pathogenicity (2,5–8). However, it is not clear to what extent transcriptomic changes relate to phenotypic changes or selection. Pioneering work from Mary-Claire King and Allan Charles Wilson set the tone for investigating this question by proposing that variations in morphological and behavioral traits arise more often through gene expression regulation than evolution at the protein-coding level (9). François Jacob then postulated an essay that stemmed from this theory in which he highlights how evolution acts as a tinkerer that works from already available material, i.e. through regulation of gene expression, to create new adaptations (10). This constituted the core of the evolutionary developmental biology which matured into the still-debated claim that new adaptations mainly emerge through cis-regulation of gene expression, i.e. through noncoding DNA regulating a neighbor gene contrarily to trans-regulators acting on distant genes (11–14). This debate has been reinforced by the technical difficulties and complexity of assessing the evolution and outcome of mutations in non-coding regions (11,12). Advances in sequencing technologies have clarified some of these hypotheses, particularly in the context of transcriptome analyses of the model organism *Saccharomyces cerevisiae*. For instance, Brem et al (2002) used microarray technology to relate the gene expression profiles of 40 yeast segregants from a lab (BY) and natural vineyard strain (RM) to their genetic markers (15). They found that cis-acting modulation is the main mechanism for regulating gene expression. Nearly two decades later, by greatly increasing statistical power, Albert and collaborators (2018) found that most of the expression variation arise through trans-regulation using non-multiplexed RNA-seq to analyze 5,720 genes in 1,012 yeast segregants generated by a crossing between RM and BY (16). The analysis method they used, i.e. expression quantitative trait loci (eQTL) mapping, consists in correlating allele frequencies to gene expression levels to find the loci modulating expression.

Although eQTL mapping is a traditional GPM analysis that accounts for the transcriptomic layer, it is typically realized through non-multiplexed RNA-seq which tends to have low statistical power due to challenges with experimental scale and confounding factors (17,18). Thus, eQTL

mapping traditionally cannot identify significant low-effect regulatory mutations that are important for understanding the genetic bases of complex traits and diseases (19,20). Furthermore, most eQTL studies only assess the average transcriptomic profile of bulk populations without being able to capture the profile of rare cell lineages within a population. This is a critical limitation in heterogenous populations such as cancer or microbial populations where rare lineages can drive relapse or drug resistance (21).

Here, we sought to circumvent the challenges of non-multiplexed bulk RNA-seq imposed by the scale and population heterogeneity by performing eQTL mapping through single-cell RNA sequencing (scRNA-seq) of a pool of ~4500 well-characterized F1 segregants of a yeast cross (16,22,23). In the same way that combinatorial indexing/barcoding and multiplexing enable the collection of large-scale fitness and genotype data (24), we hypothesized that scRNA-seq can help us collect both genotype and expression data on a large pool of segregants. We employ several strategies to overcome previous obstacles of eQTL mapping studies: i) we pool cells from thousands of segregants during the growth step and perform a single scRNA-seq run on the culture to account for environmental effects, and ii) from the exome sequencing data of single-cells we take advantage of the reference panel to validate that we accurately infer the genotype of each cell from extremely low number of reads mapping to polymorphic sites per cell (effectively ~0.2x coverage).

Using this approach, we integrated the resulting transcriptomic data from growth in rich media with a pre-existing yeast GPM. We estimated the heritability of the transcriptome and the extent at which transcriptome is associated with fitness. We show that this increased scale from scRNA-seq enables eQTL mapping directly without the use of a reference genotype panel, and relate identified single-cell eQTL (sc-eQTL) to previously identified QTL. We also exploit the identified sc-eQTL to analyze the patterns of cis- and trans- regulation in the GPM.

Our single-cell RNA-seq approach is consistent with yeast GPM results from non-multiplexed assays

We initially aimed to show that performing scRNA-seq at a large scale can generate data that are consistent with non-multiplexed DNA and RNA sequencing. To do so, we analyzed a dataset of thousands of yeast lineages generated by Nguyen Ba and collaborators (2022) (24). To understand the yeast GPM, they collected fitness and genotype data from ~100,000 segregants of an F1 cross between a laboratory strain of yeast (BY) and a natural vineyard strain (RM) (**Figure 1A**).

Figure 1 Yeast segregants datasets. A) Reference panel from the barcoded bulk sequencing. The
99,995 yeast segregants in the reference panel come from a F1 cross between a laboratory strain
of yeast (BY) and a natural vineyard strain (RM) (24). Thus, they only have 2 possible alleles at
each of the 41,594 polymorphic sites. The lineages barcodes enabled fitness estimation from
competition assays in 18 environments recapitulating the adaptation to temperature gradients, the
ability to process different sources of carbon and the resistance to antifungal compounds. B)
Pooled scRNA-seq dataset from a single batch. We performed scRNA-seq of the first batch of
segregants (n=4,489) to obtain genotypes that are similar to the reference panel and single cell's
expression profiles. Non-covered sites, sequencing errors and the presence of reads in the wrong
library (index swapping) are corrected for using the HMM described in Figure S1.

Using this approach named barcoded bulk QTL mapping or BB-QTL mapping, they revealed the
complex polygenic and pleiotropic nature of phenotypes as well as an unprecedented number of
pairwise epistatic interactions. To integrate transcriptomic data to that GPM, we performed
scRNA-seq using the 10X Genomics Chromium microfluidics platform and obtained both
genotype and expression profiles from 18,233 cells of the first batch of segregants (**Figure 1B**).
This short-read scRNA-seq method comes with challenges like low-coverage sites due to technical
sequencing biases and low sequencing depth in some cells (25,26). To overcome these challenges,
the unique molecular identifiers (UMIs) of the 10X Genomics platform provide a control for
technical biases by quantifying gene expression from unique transcribed molecule counts instead
of reads counts (25). In addition, Hidden Markov Models (HMMs) can infer accurate genotype

data even at sequencing depths as low as 0.1x (24). Nguyen Ba and collaborators (2022) designed
an HMM to infer the segregants genotypes from the observed reads at low depth of DNA
sequencing by accounting for sequencing error rate, recombination rate and index swapping rate
(24). As there are only two ancestral lineages, there are only two possible alleles for the strains at
each of the 41,594 polymorphic sites. Thus, the genotype of the segregants can be represented by
the frequency of only one of the parental alleles, which is RM in the dataset. Applying this model
to low-coverage segregants yielded genotypes that are significantly similar to high-coverage
replicates (24). We sought to use a similar model to infer genotypes from scRNA-seq data, but we
anticipated that some of these parameters may differ due to increased error rate of the reverse-
transcriptase, increased index swapping due to pooled-reaction, etc (**Figure S1**). In Nguyen Ba et
al, those rates were heuristically determined, but here we estimated these from the read mapping
data and found that re-estimated parameters from data increase the proportion of recovered strains
in the single cell data from 58.6% to 72.0%.

After adapting the HMM to the scRNA-seq data, we sought to validate that the resulting
cell genotypes relate well to their corresponding strain in the reference panel obtained by non-
multiplexed DNA sequencing strategies. Ideally, each single-cell barcode (from 10x Genomics
Chromium) should be associated with a single cell and a cell should have a clear match with a
unique strain in the reference panel. However, several factors can obscure these associations, e.g.
a single-cell droplet containing cells from 2 different strains, a low-coverage cell, uncertainty in
the allele of the reference genotype, etc. Thus, we designed an approach to clearly assign cells to
the correct reference panel strain (see **Methods**). This approach relies on two metrics of similarity
between the cells and the strains' genotypes, i.e. the expected distance between them, which should
be minimized for the best match, and the relatedness (R^2). The statistical significance of the
relatedness between single cells and reference lineages was determined by a permutation test
(**Figure S2**). From the read mapping alone, we obtained a mean R^2 of 0.59 ($\sigma = 0.19$ and median
= 0.64), which was significantly improved after applying our HMM to correct for mis-identified
alleles and imputing data in low-coverage sites using recombination probability. Indeed, the
single-cell HMM genotypes yield a mean R^2 of 0.73 ($\sigma = 0.18$ and median = 0.81; **Figure2A**). We
found that the distribution of relatedness after HMM was still left-skewed, with many cells
statistically significantly assigned to a reference genotype despite having what appeared to be low
relatedness. Upon investigation, it was found that these could be explained by genotyping
uncertainty either in the single-cell and/or in the reference panel genotype (**s**) (**Table S1**).

A

B

161
Figure 2 Single-cell RNA-seq data recapitulate bulk DNA and RNA assays results. A) Effect
of the HMM on the relatedness between single cell genotypes and their closest reference lineage.
The single-cell original genotype represents the genotype of the cells before the correction with
the HMM. The relatedness to the closest lineage in batch1 has been measured with the adjusted
R^2. To control for genotype uncertainty, only the 13,069 barcodes with a significant lineage
assignment (lineage-barcode genotype correlation FDR<0.05) and a reference lineage with a lower
uncertainty than the single cell HMM are selected, which represents 72.2% of the barcodes. We

then rounded the genotypes to remove the uncertainty during the comparison. Wilcoxon signed
test p-value is indicated above the violin plots. B) Narrow-sense heritability measured with non-
multiplexed DNA sequencing and scRNA-seq. The grey bars represent the scRNA-seq estimates
of narrow-sense heritability while the red dots represent the estimates from bulk DNA sequencing.
The interval of confidence of the bulk DNA sequencing is indicated by the red line around the red
dot and was obtained from genotype and phenotype measurement error in the BB-QTL paper (24).
The 23C-37C represents the temperature for the competition assay in YPD media while the other
phenotypes represent growth on YNB, molasses (mol), mannose (Mann) or raffinose (raff) and
chemical resistance to copper sulfate (Cu), ethanol (eth), guanidinium chloride (gu), lithium
acetate (Li), Sodium dodecyl sulfate (SDS) and suloctidil (suloc) (24).

To further establish that the genotyping obtained from scRNA-seq data was comparable to
previous non-multiplexed genotyping of the reference genotype panel, we estimated the
contribution of genetic variation to the phenotypic variation, i.e. fitness heritability. Nguyen Ba
and collaborators (2022) estimated the narrow- and broad-sense heritabilities of complex
phenotypes associated with temperature gradient, carbon source and chemical resistance for which
RM and BY segregants exhibit a significant level of diversity (24). We used our lineage assignment
to that panel to obtain fitness but used our single-cell genotyping to perform this association.
Encouragingly, most GCTA-REML estimates of narrow-sense heritability are within the
confidence intervals of Nguyen Ba and collaborators (2022) estimates (**Figure 2B**).

Although the variance partitioning is consistent with previous studies, it only provides a
broad view of the genotype-phenotype map as it does not allow to identify the loci that significantly
explain phenotype variation. If the genotypes obtained by scRNA-seq were of high-quality, then
we would expect that a QTL mapping model from scRNA-seq would yield a similar model than
non-multiplexed DNA sequencing data. To do so, we used a cross-validated stepwise forward
linear regression on the strain fitness and consensus genotypes data from single-cells that shared
the same lineage assignment (**Methods**). Performing the QTL mapping on the batch 1 scRNA-seq
dataset enabled the identification of 29 QTL compared to 31 QTL identified with the bulk barcoded
approach (**Tables S2 and S3**) (24). These QTL were largely similar as shown by the non-
significant difference between the effect sizes (Wilcoxon signed rank test $p = 0.29$) and by a model
similarity metric (24) that considers the recombination distance between matched QTL, the
similarity of the effect sizes and the allele frequencies (**Methods**). Using this approach, we
estimated that the similarity score between the batch 1 single cells QTL and the batch 1 BB-QTL
is 86.2% while each model respectively had a similarity score of 78.7% and 78.2% with the full
BB-QTL mapping performed on 99,950 segregants (24) (**Figure S4**). The QTL identified from the
scRNA-seq dataset also recapitulated several important biological features of the reference panel
such as an enrichment of non-synonymous and disordered region QTL (24) (**Figure S5**).

Finally, the variance partitioning model can also be modified to include gene expression as
the response variable and cell genotypes as the only random effect (**Methods**). This enables the
quantification of expression heritability, i.e. the variance of expression explained by genotype.
Using this approach, we estimated that genotype explains 72.3% of expression variance, which is

consistent with results from previous non-multiplexed eQTL mapping studies. Indeed, Albert and
collaborators (2018) estimated that genotype explains 70% of expression variance using a dataset
of 5720 genes in 1012 yeast segregants generated by the same parental strains (RM and BY).

Integrating scRNA-seq data to an existing GPM highlights selection on the transcriptome

Having shown that scRNA-seq is consistent with non-multiplexed assays while being more
scalable, we next sought to highlight new associations within the BY/RM GPM. Selection is often
highlighted at the genotype level through convergent evolution, increase in allele frequency within
a population or population genetics metric (26–28). However, the central dogma of molecular
biology and evolution tinkering entail that phenotype variation should be linked to transcriptomic
variation. As our dataset included all these variables, we sought to provide a variance partitioning
framework to evaluate the association between the transcriptome and trait variation (**Methods**)
with the 30C phenotype as an example (**Figure 3**).

222
Figure 3 Variance partitioning of the 30C phenotype from scRNA-seq data. The percentages
represent the proportion of fitness variance (whole rectangle area) explained by the components.
The ellipse area represents the phenotype variance explained by genotype variation and the circle
area represents the phenotype variance explained by expression variation. The black area of the
rectangle represents the residual of the model while the other colored areas represent the shared
and exclusive components explaining fitness variation.

The components of this variance partitioning all relate to at least one biological phenomenon.
Indeed, the portion of trait variation explained exclusively by the genotype variation (red in **Figure**
3) represents the effect of mutations on fitness via several biological phenomena such as protein
stability, enzymatic function etc, independent of expression level. For the 30C phenotype, this
component explains 31.2% of the fitness variation in the BY/RM background which is similar to
the 29.7% explained by the shared component between phenotype, genotype and expression
variations (purple in **Figure 3**). The latter represents the association between selection (fitness)
and the transcriptome either through loci influencing fitness via expression directly or through loci

affecting expression via an effect on cell fitness (indirectly) (29,30). Its considerable association
to fitness variation thus supports the evolution tinkering model. As for the phenotype variation
explained exclusively by gene expression (blue in **Figure 3**), it could represent epigenetics and
stochastic gene expression, which weakly explain variations in the 30C phenotype.

Although this model accurately estimates the narrow-sense heritability of 30C, the
residuals still represent 37.7% of fitness variation. This could be explained by unmeasured factors
like high-order epistasis, mitochondrial mutations or protein properties but the broad-sense
heritability of this phenotype is similar to the narrow-sense heritability, suggesting that the
residuals are mostly not explained by genotype and expression (24). Nguyen Ba et al. (2022) also
estimated that epistasis only explained around 5% of fitness (24). These results suggest that a
single run of scRNA-seq on a single batch of yeast segregants converge with bulk DNA sequencing
results while revealing previously hidden components of the GPM.

Revealing hidden components of the yeast GPM with scRNA-seq

Our integrative scRNA-seq approach is not limited to enabling the quantification of the association
between transcriptomic changes and trait variation. Indeed, the same approach we used to identify
QTL can be used to detect loci regulating gene expression which can reveal the cell mechanisms
underlying trait variation through transcriptomic changes. We thus modified the QTL mapping
framework such that the response variable is the level of expression of a single gene in the single
cells (**Methods**). This approach is a cost-efficient way to perform eQTL mapping from the
expression profile and genotype of cells from thousands of lineages in a multiplexed way (sc-
eQTL mapping).

Consistent with yeast non-multiplexed eQTL results, the genes with the highest expression
heritability are enriched in functions related to carbohydrate catabolic process (GO:0016052) and
cellular biosynthetic process (translation GO:0006412, organelle assembly GO:0070925,
ribosome biogenesis GO:0042254 and gene expression GO:0010467) (Fisher's exact test
FDR<0.05; **Methods**). In both datasets, these genes are also highly expressed, which reflects the
positive correlation between expression heritability and expression levels ($R^2 = 0.66$ and $p < 2.2e-16$).
Conversely, genes with the lowest expression heritability observed in the RM/BY background,
which we defined as the bottom 10% expression heritability, are enriched in functions related to
the cell cycle biological process (GO:0007049, Fisher's exact test FDR<0.05) (16,31).

Because of the increased scale of our collection, our approach is more powered to estimate
the gene heritability. We were thus able to detect new overrepresented biological processes, i.e.
DNA metabolic process (GO:0006259) and the response to nutrient levels (GO:0031667), for
which the variation of expression levels is weakly associated to the genetic variation observed
across the RM/BY segregants.

The functional enrichment analysis using scRNA-seq data revealed new associations
between expression heritability and biological processes in the RM/BY genetic background.
However, while it suggests that many eQTL are also QTL, it cannot accurately point to the specific

277 loci involved in trait variation and cannot address whether mutations on regulatory hubs have
278 stronger effects on traits. To investigate this, we mapped the QTL to hotspots of gene regulation
279 (or regulatory hubs), which we defined as 25 kb genomic windows that were repeatedly identified
280 in the eQTL mapping procedure (for different genes). This was done to acknowledge the
281 uncertainty in the exact position of the eQTL due to linkage disequilibrium and power. We then
282 ranked the 30C QTL identified by Nguyen Ba and collaborators (2022) based on their absolute
283 effect size and correlated it to the rank of the eQTL hotspots based on the number of regulated
284 genes. This resulted in a positive correlation (Spearman $\rho = 0.33$ and $p = 5.21e\text{-}5$), suggesting that
285 larger effects on the regulatory network translate into larger trait variation. Indeed, we observed
286 that some previously reported high-effect-size QTL genes are located in eQTL hotspots, eg MKT1,
287 HAP1, and IRA2 (**Figure 4A**).

Figure 4: eQTL features underlying trait variation across the BY/RM segregants. A) Mapping of the 30C QTL in the eQTL hotspots. We represent the hotspots of expression regulation as genomic windows (25 kb) to acknowledge the uncertainty around the real position of the eQTL due to linkage disequilibrium. We annotated the 5 top eQTL hotspots and the eQTL hotspots in which the top additive QTL identified by the BB-QTL mapping of the 30C phenotype are located. In these regions, we represented the most affected trans-regulated genes in red, the most affect cis-

regulated gene in blue and the genes of the top QTL in black. The double quotation characters
represent the absence of such genes in the associated region. We also represented the rank of the
QTL in the set of 159 QTL of the 30C phenotype. B) Partitioning of the expression heritability or
explained variance (R^2) among cis- and trans-eQTL. Each pair of points connected by a line
represents a gene. Green lines represent the genes that are only have trans-eQTL and orange lines
represent the genes that have both trans- and cis-eQTL. C) Comparison of the mean effect size
between cis- and trans-eQTL. Each pair of points connected by a line represents a gene. The ratio
of the average effect size between cis- and trans-eQTL is represented by the line color. The sample
size of each eQTL category is represented in the x axis. This is the number of trans-eQTL and cis-
eQTL used for calculating the average effect sizes per gene not the number of points per
distribution.

Performing this rank-test on individual genes also yielded the result that eQTL effect is correlated
with fitness effect for 35.1% of the genes (permutation test $p < 0.05$, see **Methods**). Although this
correlation does not apply to most genes, it reveals potential regulatory mechanisms explaining
the importance of the strongest growth loci or QTL. For instance, MKT1, i.e. the strongest growth
loci, is part of a regulation hotspot affecting genes that are important for yeast growth like ENP1
which is involved in RNA processing and HXT6 which is involved in glucose uptake (32–34).
Among the strongest growth loci, VPS70 is part of a hotspot of regulation that strongly affects the
expression of RSF2, a zinc-finger protein regulating glycerol-based growth and respiration (35).
Furthermore, the highest peak for expression regulation contains important growth loci in
chromosome IV around the mating type loci. This suggests the presence of cells with different
mating types in the dataset which we confirmed from the read mapping to Mat-a and Mat-α genes.
This is consistent with previous budding yeast eQTL mapping and is also expected because the
mating types in yeast express sets of genes that are "turned off" in other mating types (15,16,36).
This peak of expression regulation is also responsible for regulating TDH3 which is involved in
glycolysis and glucogenesis and can have important effect on fitness (37).

These hotspots suggest that expression differences in BY/RM would predominantly be due
to mutations in trans-regulatory elements. To test this, we partitioned the variation in gene
expression between cis- and trans- regulatory loci for each gene (see Methods). This analysis
revealed that all the genes are affected by at least one polymorphic trans-regulatory locus and that
these polymorphic trans-regulatory loci explain most of that gene's expression (**Figure 4B**). It is
well known that mutations in promoters and nearby enhancers can influence gene expression
(38,39). Indeed, we identified many genes that contained an allele in a cis-regulatory element that
strongly explain that gene's expression variation (n=750 genes out of 6088, **Figure 4B**). As
expected, mutations in cis-regulatory elements were of stronger effect size than trans-eQTL
individually, but the cumulative aggregate effect of all trans-eQTL acting on that gene was
comparable to the few cis-eQTL they had (**Figure 4C**). This can be explained by the fact that there
are more opportunities for mutations to arise in trans-regulatory elements. Finally, we found that
trans-eQTL have two times higher odds of affecting cell fitness than cis-eQTL (χ^2 $p = 0.01$).

Taken together, the link between the genetic basis of transcription variation across RM/BY segregants and fitness could only be revealed by integrating large-scale transcriptomic data to an existing GPM, which scRNA-seq facilitates.

CONCLUSION

By leveraging the scalability of scRNA-seq, we obtained thousands of transcriptomes from a reference pool of strains in a single experiment. This enabled the analysis of association between genotype, transcriptome, and phenotype at an unprecedented scale. Questions surrounding transcriptomic variation and phenotypic variation have been at the center of many previous quantitative genetics studies (15,16,22,36,40). These ideas and discoveries all support the fact that researchers can gain valuable insight about the evolution of traits by integrating the transcriptome in GPM analyses, which can translate into fundamental knowledge or other important applications where phenotypes evolve.

In this study, we took advantage of a previously characterized BY/RM cross where the genetic basis of growth in various environments was examined in detail (24). By integrating transcriptomic data in this genotype-phenotype map, we revealed how transcriptomic components are involved in trait variation. Similar to a previous study, which obtained transcriptomes by individual strain sequencing, we found that gene expression is highly heritable. Further, our study design also allowed us to conclude that gene expression contributes to a significant portion of the phenotypic variation in this strain collection.

This finding is corroborated by our findings that most eQTL detected in our study were previously shown to be QTL. This is perhaps not surprising given that QTL in this cross were previously inferred to be in regulatory genes, but this provides a more mechanistic view of the effect of an allele on phenotype. Indeed, we find a bias for trans-regulation for generating transcription innovation where the cumulative effect of trans-eQTL on gene expression are significant. That is not to say that cis-regulatory alleles are dispensable as cis-regulatory alleles often have large effect on gene expression. This genome-wide view of the genetic basis of transcriptional variation has consequences for the evolution of phenotypes, as the target size afforded by trans-eQTL is far larger than cis-eQTL. Thus, adaptation to small and fluctuation environmental changes may proceed preferentially through allelic changes or recombination of many small-effect trans-eQTL, but large expression changes are likely to require some cis-eQTL.

In this study, we leveraged the fact that our pool of strain was previously genotyped and phenotyped. This was obtained by liquid handling robotics and pooled competitive growth assay with barcode sequencing. While this was performed on a very large scale, it was essentially obtained by brute-force and through approaches that are not necessarily applicable to other systems. While it is clear from our results that genotyping single-cells can achieve the same genotype quality as single-reaction genotyping, it is much harder to obtain phenotyping data from scRNA-seq. Thus, our framework might not be readily translatable to other systems where similar studies on the GPM are desirable. However, two observations from this cross can be used to suggest an experimental approach. First, while epistasis is important, it contributes to a relatively

small portion of the phenotypic variance. Further, transcriptomic variation contributes little to the
missing heritability. Thus, it may be possible to use predicted fitness instead of observed fitness
and recapitulate essentially similar results as this study. Predicted fitness could be obtained from
bulk-segregant analysis where the additive effect of loci can be inferred from whole-genome
sequencing (23,41). While it is not clear if these observations are generalizable, it may be possible
to verify this for a study system of interest with some modest time-course single-cell based
sequencing where low-coverage genotyping is possible.

However, despite the study's limitation on generalizability, our scRNA-seq framework helps
bridge understanding of how genetic variation influences transcriptomic variation. Our framework
relies on identifying the genome of single-cells from the transcriptome, which is going to be
possible from low-coverage sequencing when genetic variation within the pool is high (such as
this cross, microbiome sequencing, or cancer cells with extensive copy number variation), and
from low cell diversity with sufficient transcriptomic variation such that aggregation of single-
cells with similar transcriptomes can afford pseudo-high coverage sequencing. Thus, integrating
genotype, transcriptome, and phenotype using scRNA-seq data can be particularly efficient for
developing a more fundamental understanding of other important traits or diseases.

MATERIAL AND METHODS

Yeast strains and segregants

We analyzed cells from a single batch (batch 1) of 4,489 segregants obtained from a F1 cross between the yeast laboratory strain BY4741 and the vineyard strain RM11-1a generated in a previous study (24). These strains have been selected to generate this collection of segregants because they exhibit differences in multiple phenotypes including the adaptation to temperature, the ability to process different sources of carbon and the ability to resist antifungal compounds. Therefore, the genetic variation observed across the segregants can be correlated to the differences in growth rate observed in the 18 environments recapitulating these phenotypes in the Nguyen et al (2022) study (24). The selection of the batch is random and the fact that we performed the analyses on a single batch eliminates batch effects that could obscure variable associations. Genotypes and fitness data used were the same ones obtained in the previous study.

Yeast growth and single-cell RNA-sequencing protocol

To prepare strains for scRNA sequencing, we unfroze the batch of segregants and inoculated approximately $5*10^6$ cells in YPD (1% Yeast Extract, 2% Peptone, 2% Dextrose) to saturation. The next day, about 10^7 cells were passaged to 5 mL of fresh YPD and grew for 4 hours to bring cells to log-phase. We then pelleted 100 ul of cells and resuspended them in spheroplasting solution (5 mg/mL zymolyase 20T, 10 mM DTT, 1 M Sorbitol, 100 mM Sodium Phosphate pH 7.4) at a concentration of 10^7 cells/mL. The cells were incubated at 37 degrees Celcius for approximately 10 minutes at which point spheroplasting was verified by mixing a small aliquot of cells with detergent to observe lysis. The cells at this point were quantified using a hemocytometer and prepared using the standard 10x Genomics Gel Beads-in-emulsion (GEM) protocol. We used the Chromium Next GEM Single-cell 3' Reagent Kit to prepare the sequencing libraries and sequenced on a NextSeq 500 high-output flow cell.

We note that the cells analyzed here were grown in bulk and assayed for their transcriptome in log-phase. Our fitness data was obtained from competitive bulk fitness assays which includes several whole growth cycle over multiple days and thus captures lag phase, exponential growth, and saturation. Nevertheless, previous experiments had shown that fitness was mostly determined by exponential growth which suggests that our analysis is adequate even if the cells were prepared for sequencing at a single time point.

Single-cell RNA-sequencing data parsing

From the scRNA-seq reads, we obtained gene expression levels and allele counts using the pipeline count from CellRanger version 3.1.0 (42). For each of the ancestral strain, i.e RM11-1a and BY4741, the pipeline mapped the scRNA-seq reads to the reference genome, filtered the barcodes by comparing the UMI count per barcode distribution to a background model of empty gel-bead in-emulsion, and counted the number of UMI per gene per barcode. The barcode filtering retained 18,233 barcodes. For each barcode, we then counted the number of RM and BY alleles at each polymorphic site by parsing the RM and BY bam files using a python script (https://github.com/arnaud00013/sc-eQTL/tree/main/II_scRNA-seq_genotyping). This script only

429　keeps reads that mapped at the same loci on both reference genomes to increase the level of
430　confidence of the mapping.

Correction and imputation of single-cell genotypes with a Hidden Markov Model

Because there are only two possible alleles at each polymorphic sites of the RM/BY segregants,
their genotype can be recapitulated by a quantitative variable measuring the proportion of reads
from one of the parental strains, which is RM in our dataset. The raw allele count data provides a
first estimate of this RM allele frequency at each polymorphic site. However, due to the low mean
depth of coverage of scRNA-seq data (0.2x), the absence of reads in some polymorphic sites and
the biases introduced during sequencing like index hopping/swapping, we expect that the raw data
can be imputed and corrected for errors and uncertainty in the observed alleles. Therefore, we
applied a Hidden Markov Model (HMM) on the observed allele count. Such model can infer
accurate genotype data at sequencing depths as low as 0.1x (24,25,43). Nguyen Ba and
collaborators (2022) designed an HMM to infer the segregants genotypes from bulk DNA
sequencing by accounting for sequencing error rate, recombination rate and index swapping rate
(24). Because scRNA-seq uses the reverse transcriptase, which has a higher error rate, and because
it is a pooled assay with higher chances of index swapping, we expected the HMM parameter to
differ for the single cell data. Therefore, we adapted the HMM to scRNA-seq data by measuring
its parameters in our dataset (**Figure S1**). The scripts are available on GitHub
(https://github.com/arnaud00013/sc-eQTL/tree/main/II_scRNA-seq_genotyping).

Assigning single cells to the reference panel strains

450　To evaluate the level of relatedness between the reference panel strains and the imputed single cell
451　genotypes, we used the expected distance to identify the strain that best relate to each single cell:

$$Expected\ distance(g_c, g_s) = \sum_{i=1}^{41594} g_c + g_s - 2g_c g_s \qquad \text{(Eq.1)}$$

453　where g_c is the cell genotype and g_s is the strain genotype. Next, we assigned the single cell to its
454　best match in the studied batch of 4,489 trains only if this match is better than the best match in
455　randomly generated batches of the same size (**Figure S2**). This procedure is implemented and
456　available at https://github.com/arnaud00013/sc-eQTL/tree/main/III_Genotype_analysis).

Partitioning the phenotypic variance into genetic and transcriptomic components

458　To analyze the yeast GPM at a broad scale and to evaluate the association between selection and
459　the transcriptome, we estimated the contribution of genetic and transcriptomic variations to
460　phenotypic variation from scRNA-seq data. More precisely, we performed a Genome-wide
461　Complex Trait Analysis (GCTA) by fitting a linear mixed model to the data using the restricted
462　maximum-likelihood (REML) method (44):

$$y = X\beta + W_g u_g + \varepsilon_g \qquad \text{(Eq.2)}$$

$$y = X\beta + W_e u_e + \varepsilon_e \qquad \text{(Eq.3)}$$

$$y = X\beta + W_g u_g + W_e u_e + \varepsilon \qquad\qquad\qquad \text{(Eq.4)}$$

where y is the fitness vector for the n cells, X is the nxk matrix of k fixed effects, β is the vector of
k coefficients of the fixed effects, W_g is the nxp genotype matrix, u_g is the vector of p SNP effects,
W_e is the nxm expression matrix, u_e is the vector of m gene expression effects and ε is the error
term. Because the dataset does not include fixed effects, we set the fixed effect to a vector of ones
such that its coefficients represent the mean fitness while the genotype and expression data are the
random effects that explain the fitness variance along with the error terms. The REML solution
assumes that the data follow a Gaussian distribution, so the data are standardized before fitting the
model. We also divided the standardized expression counts by the cell sum of expression counts
to control for molecule count biases across cells. The cell fitness is based on the fitness of the
closest segregant in batch 1 as measured by the expected distance. Because this model is linear
and additive, it can be compared to the estimates of narrow-sense heritability obtained by Nguyen
Ba and collaborators (2022) (24). The difference between the variance explained in equation 4 and
equations 2 or 3 allow to infer the variance explained only by the genotype or the expression
component of the model. The code for the variance partitioning is available on GitHub
(https://github.com/arnaud00013/sc-eQTL/tree/main/IV_variance_partitioning).

Estimating the expression heritability from scRNA-seq

To obtain this estimate from scRNA-seq data, we needed to consider the fact that GCTA-REML
only takes a vector as a response variable while the gene expression matrix is multi-dimensional.
To solve this, we orthogonalized the gene expression matrix using principal component analysis
(PCA), and used each of the PC one at a time as a response variable of the model. Indeed, if the
expression PCs recapitulate the total expression variance and are orthogonal or independent to
each other, then the sum of the PCs variance explained by genotype should be the expression
heritability. To save time, we only used the 898 expression PCs that explain 99% of expression
variance:

$Expression\ heritability = \sum_{i=1}^{898} PC_i\ eigen\ value * "PC_i \sim genotype"\ model\ R^2$ (Eq.5)

QTL mapping

To identify the loci that influence cell fitness, we performed a linear regression on the consensus
genotypes of the strains from the single cell data and the strain fitness. We decided to use the
consensus genotypes of the strains as they relate better to the bulk segregant genomes. To build
the consensus genotypes, we defined cells from the same lineage as the ones that shared the same
closest segregant in batch 1. Next, we used the median to obtain cells' consensus genotypes as it is
less sensitive to outliers and because it yields the best relatedness to the batch 1 reference
genotypes (median $R^2 = 87.0\%$; $\mu=79.5\%$; $\sigma=18.2$; **Figure S3**). We selected the QTL in the linear
models using cross-validation on the scRNA-seq data. This analysis consists in dividing the dataset
into 10 random partitions of similar sample sizes and running a cross-validated stepwise forward

linear regression on each partition. For each partition, the model starts with no QTL and a linear
model "Fitness ~ Genotype" is fitted using the genotype data at each polymorphic site, where the
correlation coefficient represents the effect size of the SNP. Then, the forward search starts and at
each iteration, a new locus with the minimum linear model residual sum of squares (RSS) is added
to the QTL model, which is updated with new effect sizes after the addition of a new SNP. Because
the order of addition of QTL matters in the forward search and because some QTL are linked or
collinear, the model can be refined by exploring different QTL around the local optima. These
steps are repeated until the model RSS cannot be improved anymore or until the number of QTL
reaches an arbitrary maximum far from the cross-validated number of QTL. After the forward
search is completed in each partition, the algorithm calculates the optimal λ values that minimizes
the objective function F_o:

$$F_o(\beta) = RSS(\beta) + Lasso\ penalty(\beta)$$

$$\|Y - X\beta\|_2^2 + \lambda\|\beta\|_0 \text{ (Eq.6)}$$

where β is the vector of SNP effect sizes in the QTL model, $\|Y - X\beta\|_2^2$ is the RSS of the linear
QTL model, λ defines the penalty for adding a new SNP to the model and $\|\beta\|_0$ is the number of
SNPs in the QTL model. This objective function has the property to add sparsity in the QTL model
and thus avoid overestimating the number of QTL while being consistent (24). The optimal λ has
a minimum of log(n) which corresponds to the Bayesian Information Criterion (BIC), which is
known to yield correct models asymptotically (45). This allows to consider the possibility that a
sparser model than the one found using the BIC could yield better predictive power on a test set
while avoiding overfitting. The optimal λ values found in all the partitions are then averaged and
the resulting mean λ is used to solve the objective function in the full dataset, which yields the
optimal QTL model. The cross-validation assumes that the partitions are independent, such that
the variance explained by the model and the number of relevant QTL are unbiased estimates.

Highlighting hotspots of gene regulation through eQTL mapping

To identify the loci regulating gene expression regulation, we adapted the QTL mapping
framework using expression as the predicted phenotype. Because this approach had to be repeated
for each of the 6,240 genes, we needed to modify it so that the execution time is convenient. To
do so, the parameter λ was not estimated using cross validation but rather from the Bayesian
Inference Criterion (BIC), i.e. $\lambda = log(n)$ where n is the number of cells. We found that the BIC
was often selected by the cross-validation procedure when tested on a few genes and thus we do
not believe that this approach will significantly change our results.

To acknowledge the uncertainty around the exact position of eQTL due to linkage disequilibrium,
we define eQTL hotspots as 25 kb genomic windows that were repeatedly identified in the eQTL
mapping procedure. The code for the single cell eQTL mapping is available on GitHub
(https://github.com/arnaud00013/sc-eQTL/tree/main/V_sc_eQTL_mapping).

Functional enrichment analysis by gene ontology annotation

To highlight gene functions enriched at different levels of expression or expression heritability,
we performed the panther database binomial test for statistical overrepresentation of gene ontology

biological processes (31,46). A low level was defined as within the 25% bottom part of the
distribution (<Q1) while a high level was defined as within the top 25% part of the distribution
(>Q3). The *p*-values were corrected for multiple testing using the false discovery rate correction
(FDR).

Matching QTL to eQTL

To evaluate the contribution of gene expression regulation to fitness variation, we created a model
to match QTL and eQTL based on the similarity of loci and the similarity of predicted effect on
gene expression. More precisely, for each of the 6,088 genes for which we could detect eQTL, we
performed a new eQTL model by correlating the expression level of the gene to the genetic
variation at QTL positions. This allowed us to measure the predicted effect of the QTL on gene
expression. We then calculated the distance between the QTL and the real eQTL of the gene based
on recombination distance within each chromosome, which decreases exponentially with genetic
distance, and the difference in the predicted effect on the gene expression using the formulation
developed by Nguyen Ba et al (2022) (24). Next, we used the same Needleman-Wunsch algorithm
to find the most likely set of pairing between QTL and eQTL, where an unmatched QTL is also
possible but penalized. Finally, we determined the proportion of genes for which gene expression
regulation is associated with higher fitness. To do so, for each gene, we performed a permutation
test by comparing the average rank of the matched QTL of the gene to the average rank of 999
random subsets of unmatched QTL of the same size. The p-value is the proportion of random
subsets of unmatched QTL with a higher average QTL rank than the set of matched QTL.

Comparing cis- and trans-eQTL contribution to expression variation

563 We used the definition of local eQTL in Albert et al. (2018) to define cis-eQTL, i.e. any eQTL
564 between 1,000 bp upstream of the gene and 200 bp downstream of the gene. Thus, we defined
565 trans-eQTL as the eQTL that do not follow this criterion. For each gene, we then performed
566 variance partitioning using the GCTA:

$$y = X\beta + W_{g\_cis}u_{g\_cis} + \varepsilon_{cis} \tag{Eq.7}$$

$$y = X\beta + W_{g\_trans}u_{g\_trans} + \varepsilon_{trans} \tag{Eq.8}$$

$$y = X\beta + W_{g\_cis}u_{g\_cis} + W_{g\_trans}u_{g\_trans} + \varepsilon \tag{Eq.9}$$

where *y* is the vector of expression level of the gene across the *n* cells, *X* is the *n*x*k* matrix of *k*
fixed effects, β is the vector of *k* coefficients of the fixed effects, W_{g_cis} is the *n*x*p* cis-eQTL
genotype matrix, u_{g_cis} is the vector of *p* cis-eQTL effects on expression, W_{g_trans} is the *n*x*m*
trans-eQTL expression matrix, u_e is the vector of *m* trans-eQTL effects on expression and ε
represent the error terms. Because the dataset does not include fixed effects, we set the fixed effect
to a vector of ones such that its coefficients represent the mean expression level while the cis-
eQTL and trans-eQTL genotypes are the random effects that explain the expression variance along
with the error terms. We can infer the variance explained by the cis-eQTL by the difference in
variance explained between the models in equations 9 and 8. Likewise, the difference of variance
explained by the models in equations 9 and 7 can help us estimate the variance explained by the

trans-eQTL. Finally, we estimate the effect sizes using the absolute value of the correlation
coefficients of each loci and compare the mean between the cis- and trans-eQTL from the same
gene (paired data) with a Wilcoxon signed rank test.

DATA AVAILABILITY

The code used for this study is available and explained at https://github.com/arnaud00013/sc-eQTL and the original single-cell reads from the pooled segregants scRNA-seq assay have been uploaded in the NCBI BioProject database with the accession number PRJNA1022775. The single-cell barcodes expression data are also available at https://github.com/arnaud00013/sc-eQTL as an archive file named Matrix_gene_expression_barcodes_1_to_9000.csv.tar.gz or Matrix_gene_expression_barcodes_9001_to_18233.csv.tar.gz.

ACKNOWLEDGEMENTS

ANNB acknowledges support from the Natural Science and Engineering Research Council of Canada (NSERC RGPIN-2021–02716 and DGECR-2021–00117) and AN acknowledges support from the Natural Science and Engineering Research Council (NSERC CGS-D App Id: 569340-2022, UTF Fellowship from the University of Toronto and Rustom H. Dasture Graduate Scholarship in Cell and Systems Biology). The computations in this paper were enabled by resources provided by Compute Canada. We also thank the Centre for the Analysis of Genome Evolution and Function (CAGEF) and TCAG at SickKids for sequencing support.

BIBLIOGRAPHY

1. Bartoli C, Roux F. Genome-Wide Association Studies In Plant Pathosystems: Toward an Ecological Genomics Approach. Front Plant Sci [Internet]. 2017 May 23;8. Available from: ://WOS:000402030200001

2. Ferreira MA, Gamazon ER, Al-Ejeh F, Aittomaki K, Andrulis IL, Anton-Culver H, et al. Genome-wide association and transcriptome studies identify target genes and risk loci for breast cancer. Nat Commun [Internet]. 2019 Apr 15;10. Available from: ://WOS:000464494100010

3. Aguet F, Alasoo K, Li YI, Battle A, Im HK, Montgomery SB, et al. Molecular quantitative trait loci. Nat Rev Methods Primer. 2023 Jan 25;3(1):1–22.

4. Tarantino LM, Eisener-Dorman AF. Forward Genetic Approaches to Understanding Complex Behaviors. Curr Top Behav Neurosci. 2012;12:25–58.

5. Casamassimi A, Federico A, Rienzo M, Esposito S, Ciccodicola A. Transcriptome Profiling in Human Diseases: New Advances and Perspectives. Int J Mol Sci. 2017 Jul 29;18(8):1652.

6. Wainberg M, Sinnott-Armstrong N, Mancuso N, Barbeira AN, Knowles DA, Golan D, et al. Opportunities and challenges for transcriptome-wide association studies. Nat Genet. 2019 Apr;51(4):592–9.

7. Transcriptome: Connecting the Genome to Gene Function | Learn Science at Scitable [Internet]. [cited 2023 Aug 31]. Available from: https://www.nature.com/scitable/topicpage/transcriptome-connecting-the-genome-to-gene-function-605/

8. Williams CG, Lee HJ, Asatsuma T, Vento-Tormo R, Haque A. An introduction to spatial transcriptomics for biomedical research. Genome Med. 2022 Jun 27;14(1):68.

9. King MC, Wilson AC. Evolution at Two Levels in Humans and Chimpanzees. Science. 1975 Apr 11;188(4184):107–16.

10. Jacob F. Evolution and Tinkering. Science. 1977 Jun 10;196(4295):1161–6.

11. Hoekstra HE, Coyne JA. The Locus of Evolution: Evo Devo and the Genetics of Adaptation. Evolution. 2007;61(5):995–1016.

12. Kratochwil CF, Meyer A. Evolution: Tinkering within Gene Regulatory Landscapes. Curr Biol. 2015 Mar 30;25(7):R285–8.

13. Primig M, Williams RM, Winzeler EA, Tevzadze GG, Conway AR, Hwang SY, et al. The core meiotic transcriptome in budding yeasts. Nat Genet. 2000 Dec;26(4):415–23.

14. Cavalieri D, Townsend JP, Hartl DL. Manifold anomalies in gene expression in a vineyard
isolate of Saccharomyces cerevisiae revealed by DNA microarray analysis. Proc Natl Acad
Sci U S A. 2000 Oct 24;97(22):12369–74.

15. Brem RB, Yvert G, Clinton R, Kruglyak L. Genetic Dissection of Transcriptional Regulation
in Budding Yeast. Science [Internet]. 2002 Apr 26 [cited 2022 Apr 29]; Available from:
https://www.science.org/doi/pdf/10.1126/science.1069516

16. Albert FW, Bloom JS, Siegel J, Day L, Kruglyak L. Genetics of trans-regulatory variation in
gene expression. Wittkopp PJ, editor. eLife. 2018 Jul 17;7:e35471.

17. Schwarz T, Boltz T, Hou K, Bot M, Duan C, Loohuis LO, et al. Powerful eQTL mapping
through low-coverage RNA sequencing. Hum Genet Genomics Adv. 2022 Jul
14;3(3):100103.

18. Fan Y, Zhu H, Song Y, Peng Q, Zhou X. Efficient and effective control of confounding in
eQTL mapping studies through joint differential expression and Mendelian randomization
analyses. Bioinformatics. 2020 Aug 13;37(3):296–302.

19. Bush WS, Moore JH. Chapter 11: Genome-Wide Association Studies. Plos Comput Biol
[Internet]. 2012 Dec;8(12). Available from: ://WOS:000312901500028

20. Lorenz K, Cohen BA. Small- and Large-Effect Quantitative Trait Locus Interactions
Underlie Variation in Yeast Sporulation Efficiency. Genetics. 2012 Nov;192(3):1123-+.

21. Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, et al. Rare cell
variability and drug- induced reprogramming as a mode of cancer drug resistance. Nature.
2017 Jun 15;546(7658):431-+.

22. Bloom JS, Ehrenreich IM, Loo WT, Lite TLV, Kruglyak L. Finding the sources of missing
heritability in a yeast cross. Nature. 2013 Feb;494(7436):234–7.

23. Ehrenreich IM, Torabi N, Jia Y, Kent J, Martis S, Shapiro JA, et al. Dissection of genetically
complex traits with extremely large pools of yeast segregants. Nature. 2010
Apr;464(7291):1039–42.

24. Nguyen Ba AN, Lawrence KR, Rego-Costa A, Gopalakrishnan S, Temko D, Michor F, et al.
Barcoded Bulk QTL mapping reveals highly polygenic and epistatic architecture of complex
traits in yeast. Verstrepen KJ, editor. eLife. 2022 février;11:e73983.

25. Hwang B, Lee JH, Bang D. Single-cell RNA sequencing technologies and bioinformatics
pipelines. Exp Mol Med [Internet]. 2018 Aug 7;50. Available from:
://WOS:000441266700002

26. Bergström A, Simpson JT, Salinas F, Barré B, Parts L, Zia A, et al. A High-Definition View
of Functional Genetic Variation from Natural Yeast Genomes. Mol Biol Evol. 2014 Apr
1;31(4):872–88.

27. Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. The dynamics of molecular
evolution over 60,000 generations. Nature. 2017/10/19 ed. 2017 Nov 2;551(7678):45–50.

28. Johnson MS, Gopalakrishnan S, Goyal J, Dillingham ME, Bakerlee CW, Humphrey PT, et
al. Phenotypic and molecular evolution across 10,000 generations in laboratory budding
yeast populations. Verstrepen KJ, Wittkopp PJ, Verstrepen KJ, Hodgins-Davis A, editors.
eLife. 2021 Jan 19;10:e63910.

29. Sun XM, Bowman A, Priestman M, Bertaux F, Martinez-Segura A, Tang W, et al. Size-
Dependent Increase in RNA Polymerase II Initiation Rates Mediates Gene Expression
Scaling with Cell Size. Curr Biol. 2020 Apr 6;30(7):1217-1230.e7.

30. Marguerat S, Bähler J. Coordinating genome expression with cell size. Trends Genet. 2012
Nov 1;28(11):560–5.

31. Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, et al. Protocol Update for large-
scale genome and gene function analysis with the PANTHER classification system (v.14.0).
Nat Protoc. 2019 Mar;14(3):703–21.

32. Roos J, Luz JM, Centoducati S, Sternglanz R, Lennarz WJ. ENP1, an essential gene
encoding a nuclear protein that is highly conserved from yeast to humans. Gene. 1997 Jan
31;185(1):137–46.

33. Chen W, Bucaria J, Band DA, Sutton A, Sternglanz R. Enp1, a yeast protein associated with
U3 and U14 snoRNAs, is required for pre-rRNA processing and 40S subunit synthesis.
Nucleic Acids Res. 2003 Jan 15;31(2):690–9.

34. Roy A, Dement AD, Cho KH, Kim JH. Assessing Glucose Uptake through the Yeast Hexose
Transporter 1 (Hxt1). PLOS ONE. 2015 Mar 27;10(3):e0121985.

35. Lu L, Roberts GG, Oszust C, Hudson AP. The YJR127C/ZMS1 gene product is involved in
glycerol-based respiratory growth of the yeast Saccharomyces cerevisiae. Curr Genet. 2005
Oct 1;48(4):235–46.

36. Brem RB, Storey JD, Whittle J, Kruglyak L. Genetic interactions between polymorphisms
that affect gene expression in yeast. Nature. 2005 Aug;436(7051):701–3.

37. Vande Zande P, Hill MS, Wittkopp PJ. Pleiotropic effects of trans-regulatory mutations on
fitness and gene expression. Science. 2022 Jul;377(6601):105–9.

38. Mattioli K, Oliveros W, Gerhardinger C, Andergassen D, Maass PG, Rinn JL, et al. Cis and
trans effects differentially contribute to the evolution of promoters and enhancers. Genome
Biol. 2020 Aug 20;21(1):210.

39. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution
of gene regulation. Nat Rev Genet. 2012 Jun 18;13(7):505–16.

40. Bloom JS, Boocock J, Treusch S, Sadhu MJ, Day L, Oates-Barker H, et al. Rare variants
contribute disproportionately to quantitative trait variation in yeast. Landry CR, Barkai N,
editors. eLife. 2019 Oct 24;8:e49212.

41. Brauer MJ, Christianson CM, Pai DA, Dunham MJ. Mapping novel traits by array-assisted
bulk segregant analysis in Saccharomyces cerevisiae. Genetics. 2006 Jul;173(3):1813–6.

42. Zheng GX, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel
digital transcriptional profiling of single cells. Nat Commun. 2017/01/17 ed. 2017 Jan
16;8:14049.

43. Stark R, Grzelak M, Hadfield J. RNA sequencing: the teenage years. Nat Rev Genet. 2019
Nov;20(11):631–56.

44. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait
analysis. Am J Hum Genet. 2010/12/21 ed. 2011 Jan 7;88(1):76–82.

45. Harrell FE. Regression Modeling Strategies: With Applications to Linear Models, Logistic
Regression, and Survival Analysis [Internet]. New York, NY: Springer; 2001 [cited 2023
Mar 29]. (Springer Series in Statistics). Available from:
http://link.springer.com/10.1007/978-1-4757-3462-1

46. PANTHER: Making genome-scale phylogenetics accessible to all [Internet]. [cited 2023 Sep
29]. Available from: https://onlinelibrary.wiley.com/doi/epdf/10.1002/pro.4218

47. Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Res. 2000 Jan 1;28(1):27–30.

721