1   **Title: Nutritional sex-specificity on bacterial metabolites during mosquito development**

2   **leads to adult sex-ratio distortion**

Running title: Metabolism and sex ratio distortion in mosquitoes

Authors: Ottavia Romoli*[1, 2], Javier Serrato-Salas*[1], Chloé Gapp[1], Yanouk Epelboin[1], Pol Figueras

Ivern[1], Frédéric Barras[3] and Mathilde Gendrin[1]

* These authors contributed equally and are listed by alphabetical order

Affiliations:

1. Microbiota of Insect Vectors group, Institut Pasteur de la Guyane, 97306 Cayenne

2. Institut Pasteur, Université Paris Cité, CNRS UMR3569, Viruses and RNAi Unit, F-75015 Paris,

France.

3. Université Paris Cité, Department of Microbiology, SAMe Unit, Institut Pasteur, 25-28 rue du

Dr Roux, 75015 Paris, France

1 Abstract (199 words)

Mosquitoes rely on their microbiota for B vitamin synthesis. We previously found that *Aedes aegypti*

third-instar larvae cleared of their microbiota were impaired in their development, notably due to

a lack of folic acid (vitamin B9). In this study, we investigated the effects of other B vitamins on the

development of germ-free mosquito larvae. We found that diet supplementation using a cocktail of

seven B vitamins did not improve mosquito developmental success, but rather had a significant

impact on the sex-ratio of the resulting adults, with an enrichment of female mosquitoes emerging

from B vitamin-treated larvae. A transcriptomic analysis of male and female larvae identified some

sex-specific regulated genes upon vitamin treatment. When treating germ-free larvae with high

doses of individual B vitamins, we detected a specific toxic effect related to biotin (vitamin B7)

exposure at high concentrations. We then provided germ-free larvae with varying biotin doses or

with varying bacterial counts, and showed that males are sensitive to biotin toxicity at a lower

concentration than females, and require less bacteria-derived nutrients than females. These

findings shed new light on sex-specific nutritional requirements and toxicity thresholds during the

development of insect larvae, which impact the sex ratio of adults.

16 Introduction

Aedes aegypti mosquitoes are vectors of pathogens responsible for various human diseases,

notably including yellow fever or dengue fever. They are predominantly concentrated in tropical

and neotropical regions, poising a significant threat to approximately 3.9 billion people who could

potentially contract dengue (WHO, 2023). Mosquitoes are holometabolous insects, undergoing a

complete metamorphic transformation from aquatic larvae to terrestrial adults. Consequently,

several adult physiological traits, such as size and lifespan, heavily hinge on the quality of the insect

larval development, specifically influenced by the nutritional status during this developmental

stage.

Mosquitoes depend heavily on their microbiota for vital nutrients crucial for their larval

development. In their natural habitat, mosquito larvae primarily feed on microorganisms,

particulate organic matter, and detritus present in their breeding sites (Clements 1992). In

controlled laboratory rearing environments, the main food sources for larvae typically comprise fish

food or commercial pet food designed for rodents, cats, or dogs. Mosquito larvae hatched from

microbe-free eggs and maintained in sterile conditions are halted in their development when

provided with a sterile conventional diet (Coon et al. 2014). However, their development can be

rescued when a live microbiota is introduced or when provided nutrient-rich diet and kept in the

dark, strongly implying that the microbiota plays a fundamental role in furnishing to its mosquito

host essential nutrients that are not present in the diet (Coon et al. 2014; Romoli et al. 2021; Correa

et al. 2018; Wang et al. 2021). These essential nutrients encompass critical elements such as

essential amino acids, nucleosides, and B vitamins, which are beyond the synthetic capabilities of

most insects (Wang et al. 2021; Romoli et al. 2021). The investigation of mosquito nutritional

requirements and how the microbiota affects larval development could unveil crucial metabolic

insights that might serve as the foundation for innovative vector control strategies.

To unravel the mechanisms behind the intricate interactions between mosquitoes and their

microbiota, we previously developed a method that enables the generation of germ-free

mosquitoes at virtually any stage of their development. Our approach consists in the colonisation

of *Ae. aegypti* mosquitoes with an *Escherichia coli* strain auxotrophic for two amino acids, d-Alanine

(D-Ala) and meso-diaminopimelic acid (*m*DAP), that are bacteria-specific and crucial for bacterial

cell wall synthesis. When bacteria are supplied alongside D-Ala and *m*DAP to germ-free larvae,

mosquito development is rescued. As soon as the larval rearing medium is changed to sterile water

deprived of bacteria and of D-Ala and *m*DAP, bacterial growth is arrested and germ-free larvae are

obtained. Using this method, we have been able to pinpoint folic acid as one of the critical

metabolites furnished by the microbiota and required during mosquito larval development (Romoli

et al. 2021).

Vitamins of the B group are cofactors carrying on numerous cellular processes,

encompassing essential functions in the electron transport chain (riboflavin, nicotinic acid), amino

acid metabolism (pyridoxine, folic acid), lipid metabolism (biotin, riboflavin, nicotinic acid), and

nucleic acid metabolism (folic acid, nicotinic acid)(Serrato-Salas and Gendrin 2023). They are

required in very low amounts, as they serve as coenzymes and are not consumed by the enzymes

for which they act as cofactors (Douglas 2017). Insects depend on their food and microbiota to

supply B vitamins as they lack the complete metabolic pathways to synthesize them (Dadd 1973).

The insect model *Drosophila melanogaster* requires all B vitamins for optimal larval development

(Piper 2017) and similar necessities have been hypothesized for mosquitoes. As a matter of fact, B

vitamins have been identified, along with sterols, aminoacids and nucelosides, as compounds in

synthetic diets designed for rearing mosquito larvae in the absence of a microbiota (Trager 1935;

Lea et al. 1956; Singh and Brown 1957). A more recent study has highlighted the critical roles played

by amino acids and B vitamins in mosquito larval growth. Notably, riboflavin has been shown to be

required throughout the entire larval development as it is light sensitive, and thiamine, pyridoxine,

and folic acid were found to be specifically required by *Ae. aegypti* larvae to initiate pupation (Wang

et al. 2021).

In our previous study, we found that folic acid provision partially rescued the development

success of mosquito larvae deprived of their microbiota during late instars (Romoli et al. 2021).

Here, we tested the effect of supplying different B vitamin doses on *Ae. aegypti* larval development.

Larvae cleared of their microbiota during their third instar and supplemented with increasing

amounts of B vitamins did not show any improvement in their developmental success when compared to germ-free non-supplemented counterparts. Surprisingly though, we observed an impact on the sex-ratio of the resulting adults, with a significant enrichment of female mosquitoes emerging from B vitamin-treated larvae. We further investigate potential mechanisms involved in this sex-specific effect of B vitamins via transcriptomics, diet supplementation and bacterial monocolonisation. We show that during larval development, males require less bacteria-derived products, notably biotin, than females and that they are sensitive to lower doses of biotin, resulting in the observed sex ratio distortion.

Results

Vitamin supplementation affects mosquito sex-ratio in axenic conditions

We previously reported that *Ae. aegypti* larvae cleared of their microbiota at the third instar showed lower developmental success compared to conventionally reared larvae, with only 10–20 % of individuals successfully completing the metamorphosis stage. The proportion of germ-free individuals reaching the adult stage increased to ~50–60 % if folic acid (vitamin B9) was supplemented to the larval diet (0.25–1.25 mg/mL), suggesting an important role of the microbiota in providing this B vitamin to larvae (Romoli et al. 2021). As the rescue was not complete, we wondered if other B vitamins potentially produced by the microbiota were involved in mosquito larval development and could further increase the proportion of larvae developing into adults. We produced germ-free third-instar larvae, following our transient colonization protocol (Romoli et al. 2021): after egg sterilization, we mono-colonized first instar larvae with the *E. coli* HA416 strain, auxotrophic for D-Ala and *m*-DAP, in the presence of these amino acids, and starved larvae from D-Ala and *m*-DAP shortly after reaching the third instar to turn them germ-free. We supplemented these larvae with a solution containing six B vitamins (biotin, folic acid, nicotinic acid, pyridoxine,

riboflavin, and thiamine) and choline (not classified as B vitamin anymore, but recognized as

essential nutrient), at concentrations previously reported to support *Ae. aegypti* larval development

in germ-free conditions (VIT1x (Singh and Brown 1957)). We did not observe any significant increase

in the percentage of adult mosquitoes (Supplemental Figure S1, proportion of adults: p = 0.55, not

significant (ns); see Supplemental Table S1 for details on statistics). Since the folic acid concentration

supplemented in our previous study was 5–25 times more concentrated that the concentration used

in Singh & Brown, 1957, we decided to test the effect of four and eight times higher B vitamin

concentrations (VIT4x and VIT8x) on larval development. Again, we did not observe an increase in

the proportion of adult mosquitoes, rather a marginally significant decrease with VIT8x (Figure 1A,

proportion of adults: p < 0.0001; AUX vs GF, VIT4x, or VIT8x: p < 0.0001; GF vs VIT4x: p = 0.73; GF vs

VIT8x: p = 0.058; Supplemental Table S1). However, while in germ-free conditions the small number

of mosquitoes completing their development was due to a similar proportion of larvae dying and

being stalled at the larval stage (~30 %), the vitamin treatment significantly doubled mortality rates

to ~60 % (Figure 1A, proportion of dead larvae: p < 0.0001; AUX vs GF, VIT4x, or VIT8x: p < 0.0001;

GF vs VIT4x: p = 0.010; GF vs VIT8x: p < 0.0001; Supplemental Table S1). Interestingly, we saw a

significant shift in the sex ratio of the fully developed mosquitoes, with more than 80 % of adult

mosquitoes being females when larvae were treated with VIT4x and VIT8x solutions (Figure 1B, p <

0.0001; AUX or GF vs VIT4x: p = 0.0010; AUX vs VIT8x: p = 0.0037; GF vs VIT8x: p = 0.0019;

Supplemental Table S1). This suggested a sex-specific effect of B vitamins on mosquito

development.

Transcriptomic analysis of germ-free larvae treated with B vitamins

We had two alternative hypotheses to interpret the observed impact of B vitamins on sex

ratio without significantly affecting overall development success. This effect can be attributed to a

concomitant increase and decrease in the fitness of female and male larvae respectively upon

vitamin supplementation, due to differential nutritional needs and to toxic effects of these

compounds at the tested concentrations. Alternatively, some feminization mechanism may

influence male trait maturation during larval development . To sort out which of the two possibilities

prevailed, we conducted a transcriptomic study on male and female larvae cleared of their

microbiota at the beginning of the third instar, kept in germ-free conditions (GF), and then

supplemented with VIT8x until sampling, 3-6 h after the beginning of the fourth larval instar (Figure

2A; this transcriptomic study is detailed in Supplemental data and Supplemental Figures S2, S3). We

observed on one hand that VIT 8x induced the heat-shock-protein related gene *AAEL017976* in

males, which would lend credence to the toxicity hypothesis (Figure 2B). In parallel, it regulated

AAEL012340 and *AAEL017067* in females, assigned to encode lipase 1 precursor and peritrophin-

48-like, which also may point to a better development in females (Figure 2D, E). Conversely, we also

observed a strong induction of *AAEL013606* in males, predicted to encode a SRY (Sex-determining

region of Y chromosome)-like protein, which is a sex-determination factor and may alternatively

point to a potential feminizing impact of B vitamins (Figure 2C). Hence, our transcriptomic analysis

did not provide a clear conclusion on the way vitamins affected the sex-ratio.

Effect of individual B vitamins on larval development and adult sex ratio

We reasoned that if vitamins had a toxic effect on males, we could detect it more clearly at

a higher concentration. For this reason, we treated decolonised *Ae. aegypti* larvae with vitamins at

high dose, 50x compared to the reference concentration (Singh and Brown 1957), using single

vitamins to narrow down which B vitamin was causing this sex-specific effect. We added a 16x dose

of folic acid to all tested conditions to obtain enough adult mosquitoes in the germ-free control

group and have a more reliable comparisons on adult mosquito sex-ratio. We observed a significant

decrease in larval development success in folic acid and biotin treatments (Figure 3A, proportion of

adults: $p < 0.0001$; GF vs folic acid: $p = 0.015$; GF vs biotin: $p < 0.0001$; all other comparisons $p >$

0.05, ns; Supplemental Table S1). In particular, while ~90 % larvae developed into adults in the germ-

free control, only ~60 % and ~5 % mosquitoes reached the adult stage in the folic acid and biotin

treatments, respectively. While the addition of folic acid induced a non-significant increase in the

proportion of mosquitoes that were blocked in their larval development (from ~2 % in the control

to ~20 %, Figure 3A, $p = 0.0002$; GF vs folic acid: $p = 0.062$, ns; Supplemental Table S1), biotin induced

a strong mortality on larvae (from ~10 % in the control to ~70 %, Figure 3A, $p < 0.0001$; GF vs biotin:

$p < 0.0001$; Supplemental Table S1). The sex-ratio of the resulting adults was not significantly

affected by any vitamin supplementation, although no viable male mosquito developed from biotin-

treated larvae (Figure 3B, $p = 0.73$, ns; Supplemental Table S1). This low statistical effect was

probably due to the small number of biotin-treated larvae that could complete their development

(adult mosquitoes per replicate (sex): 1/24 (non identified); 1/24 (female); 3/24 (females)),

suggesting that the biotin 50x dose was generally toxic to both sexes. However, we could not

determine if the absence of fully developed males after biotin supplementation was due to a male-

killing effect or to the feminization of phenotypic characters in male mosquitoes.

Biotin requirements and toxicity are sex-specific

The experimental set up used in previous experiments could not distinguish between biotin-

related male toxicity or feminization of male mosquitoes. In fact, our visual analysis of fully-

developed mosquitoes indicated that adult mosquitoes displayed female-specific phenotypic

characteristics, but this analysis did not yield information on the genotype of those mosquitoes and

of those that did not reach adulthood. We took advantage of a recently established *Ae. aegypti*

genetic sexing strain (Aaeg-M) characterized by the insertion of the *eGFP* transgene in the male-

specific M locus (Lutrat et al. 2023). This allowed us to sort first instar larvae by sex right after egg

sterilization and perform the full experiment on male and female larvae in parallel. The effect of

four increasing biotin concentrations (1x, 4x, 8x and 20x) was tested on decolonized third instar

larvae. As done previously, we added 16x folic acid to all tested solutions to increase the number of

adult mosquitoes.

When analysing the global results independently from the mosquito sex, we observed that

the addition of folic acid alone did not significantly increase the proportion of larvae completing

their development to adulthood in the Aaeg-M strain, with ~20 % of fully developed adults in both

germ-free controls, with or without folic acid (Figure 3C, $p < 0.0001$; GF vs GF+folic acid: $p = 1$, ns;

Supplemental Table S1). The proportion of fully developed mosquitoes at 1x biotin concentration

increased to ~30 %, while it decreased to ~20 % and ~10 % in 8x and 20x biotin treatments,

respectively (Figure 3C, see Supplemental Table S1 for individual comparisons). In all treatments,

the variation in the proportion of adult mosquitoes was due to an equivalent change in the

percentage of dead mosquitoes rather than of stunted larvae.

Data sorted by sex indicated a differential biotin requirement for male and female

mosquitoes: while the biotin 1x concentration was the best one to support the development of

males (~ 30 %, Figure 3, $p < 0.0001$, see Supplemental Table S1 for individual comparisons), the 4x

concentration resulted in the highest development success in females (~ 50 %, Figure 3E, $p < 0.0001$,

see Supplemental Table S1 for individual comparisons). In females, the effect of the sole addition of

biotin 4X was not statistically different ($p = 0.25$, ns; Supplemental Table S1), but the addition of

biotin 4x and folic acid was ($p = 0.0001$; Supplemental Table S1). The sole addition of folic acid did

not significantly improve the development of germ-free larvae in these conditions ($p = 1$, ns;

Supplemental Table S1).

When comparing male and female mosquitoes in their developmental rates to pupa (i.e., the

proportion of larvae starting metamorphosis by day), we observed similar proportions of

mosquitoes starting metamorphosis in all conditions, with a higher percentage of males going

through pupation in germ-free ($p < 0.0001$), biotin 1x ($p < 0.0001$) and 8x ($p = 0.02$) treatments

(Supplemental Figure S4A; Supplemental Table S1). However, a significant proportion of these

pupae were not able to complete the developmental stage and died before emergence

(Supplemental Figure S4B). Interestingly, the addition of 16x folic acid alone to germ-free larvae

differentially affected the pupation rates of male and female mosquitoes, suggesting that sex-

dependent requirements are not exclusive for biotin (p = 0.0005; Supplemental Figure S4B;

Supplemental Table S1). Taken together these data suggest that the clearance of the microbiota

during the third instar impacts mosquito development at the metamorphosis stage, and that the

addition of biotin at a 1x concentration improves male development while a 4x concentration

already causes toxicity; in females, the optimal concentration for development is 4x.

For each experiment, adult mosquitoes were visually inspected to confirm that their

genotype (GFP fluorescence status in L1) matched their adult phenotype. In parallel, an experiment

with New Orleans mosquitoes was performed only on decolonised larvae kept germ-free or

supplemented with 4x or 8x biotin. To compare mosquito phenotype and genotype, emerged

mosquitoes were visually analysed to confirm their sex. Their DNA was extracted and subjected to

dual PCR on *Nix* and *Actin*. PCR amplicons were analysed using a capillary electrophoresis system

that allowed to automatically detect gene-specific signals. Among the analysed mosquitoes (GF: n =

70, biotin 4x: n = 65, biotin 8x: n = 59), none showed discordant results between genotype and

phenotype, further confirming that biotin had a toxic effect but did not induce any male

feminization.

Males require less bacteria-derived metabolites than females for development

Based on these results, we hypothesized that males required lower quantities of bacteria-

derived metabolites than females. To test this, we provided larvae with a low concentration of

auxotrophic bacteria so that bacterial growth was required during larval development. Thus, a *m-*

DAP and D-Ala solution at a conventional (1x) or 100-fold-diluted (0.01x) concentration were

provided every three days until pupa appearance. This allowed to maintain a bacterial population

while ensuring that bacterial loads were 2.2-73 fold lower when provided the diluted solution

(Figure 4A, 1x vs 0.01x : $p < 0.0001$ – t = 24 h, 48 h, 72 h, 144 h, 168 h, 240 h, $p = 0.013$ – t = 196 h,

$p = 0.0014$ – t = 216 h; Supplemental Table S1). Larvae received the same amount of larval food in

both conditions. This reduction in bacterial counts led to a lower development success (76 % to 33

%, $p < 0.0001$, Figure 4B). Development was also slower (11.4 d to 12.4 d: $p = 0.012$; Figure 4D;

Supplemental Table S1). While females generally develop slower than males, this did not explain

slower development as larvae that developed in 15 days or more in the 0.01x condition were only

males. As expected, sex ratio was also distorted towards a higher proportion of males with lower

amounts of bacteria (Figure 4C; 59 % to 89 %, $p = 0.043$; Supplemental Table S1). When doing a 10-

fold dilution (0.1x), we observed similar trends, albeit with non-significant differences on sex ratio,

partly because less replicates were performed (Figure S5A-D; % adults – P = 0.0061; sex – $p = 0.32$;

time – 0.012; Supplemental Table S1).

We then wished to test whether such effects were specific to the provision of biotin. The use of a

Biotin synthase (BioB) auxotrophic mutant was impossible in the experiment condition set-up

because it would require the addition of biotin for its growth. We thus relied on a Δ*iscUA* mutant

that is deficient in the main Fe-S cluster biogenesis pathway. In this mutant, activity of BioB is sharply

decreased and the level of biotin synthesized much lower than in a wild type (Reyda et al. 2009).

We observed that Δ*iscUA* has a significantly lower ability to support larval development to

adulthood than the control (65 % respect to 83 %: $p = 0.0019$; Figure 4E; Supplemental Table S1)

and that it supported the development of a slightly higher proportion of males, albeit marginally

significantly ($p = 0.085$; Figure 4F; Supplemental Table S1). This result supported the view that biotin

level lowering bears consequence on larval growth and sex ratio. Another possibility was that the

reason the Δ*iscUA* mutant was less efficient in supporting larvae development was related to its

reduced growth rate and altered fitness. Therefore, to test this last possibility, we used a Δ*mnmA E.*

coli mutant known to show slow growth rate (Zhou et al. 2021). The gene *mnmA* encodes an enzyme

required for thiolation of a subset of tRNAs. Larvae fed with Δ*mnmA* had an intermediate

development success ($p = 0.15$; Figure 4E; Supplemental Table S1) and a significant increase in

proportion of males among adults ($p = 0.015$; Figure 4F; Supplemental Table S1). Altogether these

results indicated that microbial growth, and presumably associated richness in metabolite

production, rather than specifically biotin level, is important for sex ratio.

Discussion

The mosquito microbiota provides essential nutrients to its host. Among these nutrients, B

vitamins have been shown to be required by mosquitoes to complete their larval development

(Singh and Brown 1957; Romoli et al. 2021; Wang et al. 2021). Here, we found that the larval B

vitamins requirements, especially biotin, are sex-specific: males require less B vitamins than females

for a successful development. Together, the differences in nutritional requirements and in

thresholds of toxicity led to a sex-ratio distortion after supplementing diet of germ-free *Ae. aegypti*

larvae with increasing doses of B vitamins. Specifically, there was a notable reduction in the

proportion of male mosquitoes emerging from larvae that had been subjected to vitamin

supplementation. We conclude that males require less bacterial-derived metabolites, including

biotin, for development.

Evolutionarily speaking, favouring male development when nutritional conditions are scarce may be a way to maintain a progeny that will be successful if it spreads and finds mating partners elsewhere, while when nutritional conditions are optimal, having a high development success of both sexes may allow an efficient local colonization. While our experiments were performed with individuals kept separately, the impact on sex ratio may be exacerbated in a population as male and female mosquitoes have different developmental dynamics at the larval stage, where females emerge later than males. In line with this, female adult body sizes are impacted more significantly than males by the composition of larval diet (van Schoor et al. 2020). Female development may be longer because mosquito larvae must reach a critical mass to successfully commence metamorphosis, and this mass is higher for females than males (Chambers and Klowden, 1992). In species of scarab beetles where males are larger than females, instantaneous growth rate is not different between sexes but optimal growth lasts longer in males (Vendl et al. 2018). Contrary to what we observe in mosquitoes, total development time is however not necessarily correlated with size in these beetles (Vendl et al. 2018). Similarly in *Drosophila*, females are larger than males, yet they develop on average 4 hours earlier than males, indicating that final mass and development timing are not necessarily positively correlated. During fly metamorphosis, this protogyny phenotype is genetically controlled by *Sex lethal*, the master sex switch gene (Seong and Kang 2022).

Diet composition at the larval stage has already been shown to affect the sex ratio of the adult mosquito population, with higher proportions of males emerging in starvation or suboptimal feeding conditions (Souza et al. 2019). Our data now specifies that males and females have different requirements in bacteria-derived metabolites, notably biotin, while the amount of provided food did not vary in our experiments. Females may also need more energy storage than males for egg production, as egg production after the first blood meal has been found to be strongly correlated with richness of the larval food (Yan et al. 2023).

The observed male-killing effect appears to be a specific outcome resulting from exposure

to one or a combination of B vitamins when exceeding a concentration threshold. We observed that

the optimal concentration of biotin for the development of germ-free female larvae was four times

higher than what had been previously reported as sufficient to sustain mosquito larval development

in germ-free conditions (Singh and Brown 1957), while this concentration was already toxic to

males. Depending on the vitamin, our 4x and 8x concentrations were also 2–160-fold higher than

the concentrations used more recently to formulate a chemically defined medium for germ-free *Ae.*

aegypti larvae (Wang et al. 2021). These two studies identified osmotic pressure as a critical limiting

factor for artificial diets, describing a rapid larval mortality when the rearing medium contained

either 11.6 g/L (Singh and Brown 1957) or 113.7 g/L (Wang et al. 2021) of amino acids. The osmotic

pressure of these solutions is ~25 or ~256 times higher than that of the 8x vitamin B solution we

tested. Therefore, it is unlikely that the elevated mortality observed in our experiments could be

due to high solute concentration. Yet, our transcriptomic analysis revealed the upregulation of a

chloride channel encoding gene in both male and female larvae treated with the 8x vitamin solution,

which may reflect a response to an increase in solute concentrations.

To investigate the potential mechanisms behind the male-specific toxic effect of B vitamins,

we studied the transcriptomes of germ-free male and female larvae exposed to vitamin solutions.

We encountered several problems in the process of sample sex assignment via PCR and RNA

degradation, which led us to exclude one replicate and the entire vitamin 1x condition. This resulted

in a relatively low number of genes that exhibited sex-specific regulation when compared to a prior

transcriptomic study (Matthews et al. 2018). With the same sex assignment technique, Matthews

et al's reported approximately 3400 differentially regulated genes between male and female fourth

instar larvae. Between 35 % and 40 % of the genes identified in our transcriptomic study overlapped

with those identified by Matthews *et al*, suggesting a reasonable degree of consistency with our

data, considering that our larvae were germ-free. Most significantly, *Nix* was specifically enriched in male transcriptomes. These observations suggest that while we likely missed many regulated genes, those identified in our study are reliably true positives. The analysis of genes differentially regulated in vitamin 8x conditions identified different sets of genes in male and female larvae. Notably, the upregulation of genes linked to stress response, DNA binding, development or sexual reproduction was specific to male larvae. This observation suggests that the administration of high doses of vitamins interferes with critical cellular processes, as indicated by the upregulation of a heat-shock protein coding gene. The vitamin-induced stress response might explain the observed reduction in developmental rates among male larvae. However, it remains challenging to draw definitive conclusions regarding the precise mechanisms that underlie these effects.

When we administered individual B vitamins at elevated doses (50x), we observed that biotin, and to a lesser extent, folic acid, had lethal effects on germ-free larvae. However, only biotin was found to have an impact on sex ratio at this dose. Use the Aaeg-M GSS strain, we were able to distinguish the developmental success of male and female larvae independently and to validate the absence of a feminizing effect caused by biotin. To maintain consistent experimental conditions, we added folic acid to all tested conditions. Surprisingly, this addition in the absence of biotin did not significantly affect the proportion of fully developed adults in germ-free conditions, in contrast to what we typically observe ((Romoli et al. 2021), Figure 3A). This discrepancy could potentially be explained by differences in the mosquito strains used in the two sets of experiments (New Orleans and Aaeg-M GSS, which was created from mosquitoes sourced in Thailand (Lutrat et al. 2023)). However, we think that the variations in the timing between the two types of experiments were responsible for the different outcomes of folate supplementation. Due to the large number of larvae and to diet autofluorescence, first instar larvae of the Aaeg-M GSS strain had to be starved for 24 h while assigning the sex via GFP fluorescence before adding bacteria and food. Hence, larvae were

initially in a deprived metabolic status which may have carry-over effects on later development. We hypothesize that folic acid alone is not sufficient to complement their requirements. Furthermore, experiments showed a noteworthy degree of variability of development success between replicates. While experimental conditions were extremely controlled, replicates were conducted with different batches of eggs produced in conventional conditions. This variability underscores the influence of various factors such as the maternal gonotrophic cycle or the age of the eggs on the outcomes of larval development. The quality of the blood on which mothers have fed and the quantity of eggs laid by each female could potentially influence the amounts of vitamins in embryos, consistently with observations that arthropod development is influenced by mother's age and diet (Goos et al. 2018; Deas et al. 2019; Serrato-Salas and Gendrin 2023).

Using the GFP-sexing strain, we showed that male and female mosquito larvae cleared of their microbiota exhibit different biotin requirements for their development (0.5 µg/mL for males and 2 µg/mL for females). Importantly, both sexes show heightened mortality rates during metamorphosis when exposed to higher biotin concentrations. Although more than 50 % of the larvae successfully transitioned into pupae under all tested conditions, only a range of 10 to 40 % emerged as fully developed adults (see Supplemental Figure S4A). This observation suggests that exposure to elevated biotin levels during the third and fourth instars predominantly impacts pupae. Biotin has been found to be essential for intestinal stem cell mitosis in *Drosophila* (Neophytou and Pitsouli 2022), and is more generally involved in cell cycle progression (Velazquez-Arellano and Hernandez-Vazquez 2020), suggesting its importance during life stages with high cell proliferation rates. During metamorphosis, tissue remodelling increases cell proliferation, hence the number of biotin-responsive cells may be relatively high. Elevated biotin doses may specifically deregulate cell cycle at this stage and explain why we observe high pupal mortality. In line with our observation, previous studies underscored the essential role of biotin in *Ae. aegypti* larval development, but they

also showed that high biotin concentrations induced mortality in pupae when larvae were reared in

both axenic or conventional conditions (TRAGER 1948; Pillai and Madhukar 1969). Some degree of

biotin toxicity has been observed in pupae of the flour beetle *Tribolium confusum* as well (Fraenkel

and Blewett 1943). Furthermore, high biotin concentrations that still allowed *Ae. aegypti*

development were reported to reduce adult fertility; they notably caused follicle degeneration in

females during the post-ovipositional period (Pillai and Madhukar 1969). This effect of biotin on

fertility seems to be conserved in other insect species such as the Mexican fruit fly *Anastrepha*

ludens (Benschoter and Paniagua G. 1966), the house fly *Musca domestica* (Benschoter 1967) and

the beetle *Dermestes maculatus* (Cohen and Levinson 1968). In these insects, females tended to lay

fewer eggs when fed biotin-rich diets and eggs showed lower hatching rates. In *D. maculatus* the

impact of biotin on embryogenesis appeared to be related to the binding of this vitamin to insoluble

yolk proteins, resulting in reduced amino acid availability within the embryo (Levinson and Cohen

1973).

Mosquitoes are incapable of synthesising B vitamins, thus relying on their diet and

microbiota for the essential provision of these metabolites (Douglas 2017; Romoli et al. 2021; Wang

et al. 2021; Serrato-Salas and Gendrin 2023). Notably, germ-free mosquito larvae can only develop

if a rich diet is provided and if measures are taken to protect vitamins from light-induced

degradation (Correa et al. 2018; Wang et al. 2021). Several bacterial strains including *E. coli* are

capable of rescuing mosquito development when introduced to germ-free larvae, while others such

are *Microbacterium* are not (Coon et al. 2014; Romoli et al. 2021). This correlates with genomic data

showing that *E. coli* possess complete metabolic pathways for B vitamin synthesis (Serrato-Salas and

Gendrin 2023), while *Microbacterium* does not (Coon et al. 2014; KEGG website). Our findings

regarding biotin toxicity and previous data indicating biotin impact on insect fertility even raise the

possibility that bacteria might interfere with mosquito sex ratio or, more generally, with mosquito

physiology, by delivering high concentrations of some B vitamins. Intriguingly, genomic analyses of

Wolbachia genomes have identified a riboflavin transporter gene to be associated with cytoplasmic

incompatibility (Scholz et al. 2020). Taken together, these findings underscore the critical need to

explore bacterial-induced mechanisms that involve B vitamins because they influence the mosquito

host larval development, toxicity, fertility, and the alteration of the sex-ratio.

Materials and Methods

Mosquitoes and bacteria

Escherichia coli HA 416 was grown in lysogeny broth (LB) supplemented with 50 µg/mL kanamycin,

50 µg/mL *m*-DAP and 200 µg/mL D-Ala. For all experiments, *E. coli* cultures were inoculated from

single fresh colonies in liquid LB with appropriate supplementation and incubated at 30 °C, shaking

at 150 rpm for 16 h. *E. coli* FBE051 (wt), FBE356 (Δ*iscUA*) and FBE584 (Δ*mnmA*), used for Figure 4D-

E, were cultured in the same conditions in LB with (Δ*iscUA* and Δ*mnmA*) or without (wt) 50 µg/mL

kanamycin supplementation.

Aedes aegypti mosquitoes belonged either to the New Orleans strain or to the Aaeg-M genetic

sexing strain (GSS, (Lutrat et al. 2023)). Both colonies were maintained under standard insectary

conditions at 28-30 °C on a 12:12 h light/dark cycle. Mosquitoes are routinely blood-fed either on

beef blood provided by the local slaughterhouse (Abattoir Territorial de Rémire Montjoly) or on

anesthetized mice. The protocol of blood feeding on mice has been validated by the French

Direction générale de la recherche et de l'innovation, ethical board # 089, under the agreement #

973021. Gnotobiotic mosquitoes were maintained in a climatic chamber at 80 % relative humidity

on a 12:12 h light/dark 28 °C/25 °C cycle.

Vitamin solutions

B vitamins (Sigma-Aldrich) were dissolved in the appropriate solvent at the concentration indicated

in Table 1. Stock solutions were adjusted to pH 7.4-8.0 and filtered through a 0.22 µM membrane

filter and stored at -20 °C until use.

Generation of gnotobiotic larvae

Germ-free larvae were obtained as previously described (Romoli et al. 2021). Briefly, eggs were

placed on top of a filtration unit and surface sterilised by subsequent washes in 70 % ethanol for 5

min, 1 % bleach for 5 min, and 70 % ethanol for 5 min. After rinsing three times with sterile water,

eggs were transferred to a sterile 25-cm^2 cell-culture flask filled with ~20 mL of sterile water.

Approximately 20-30 eggs were inoculated in 3 mL of liquid LB and incubated for 48 h at 30 °C

shaking at 150-200 rpm to confirm sterility.

The following day (day two), a 16 h culture of auxotrophic *E. coli* was centrifuged, and the bacterial

pellet was resuspended in 5 times the initial culture volume of sterile water supplemented with *m*-

DAP (12.5 µg/mL) and D-Ala (50 µg/mL). Larvae were individually placed in 24-well plates together

with 2 mL of bacterial suspension and ~50 µL of autoclaved TetraMin Baby fish food suspension.

Larvae were kept in a climatic chamber at 80 % RH with 12:12 h light/dark 28 °C/25 °C cycle.

To achieve bacterial decolonisation, larvae that moulted to the third instar in a time window of 5 h

during day 4 were washed in sterile water and individually transferred in a 24-well plate filled with

1.5 mL of sterile medium and 50 µL of sterile fish food in each well. Larvae obtained with this

method were shown to be largely germ-free after 12 h since transfer (Romoli et al. 2021).

The Aaeg-M GSS required a sex-sorting step prior to be transiently colonised by auxotrophic *E. coli*.

This strain carries an *eGFP* transgene in the male-specific M locus, thus only male mosquitoes

express GFP. Axenic first instar larvae were individually placed in 96-well plates and checked for

green fluorescence under an EVOS FL Auto System (Thermo Scientific). After sex-sorting, larvae

were individually placed in 24-well plates. Bacteria were added the following day (day three)

therefore the experiment schedule was shifted by one day.

For experiments shown Figure 4 and Supplemental Figure S5, bacterial suspensions were diluted in

sterile water to 100 CFU/mL. In Figure 4A-D and Supplemental Figure S5, amino acids were provided

at a conventional concentration (1x, *m*-DAP – 12.5 µg/mL and D-Ala – 50 µg/mL) or after 1:100

dilution in sterile water. For each condition, the initial concentration was provided again every three

days until pupae appeared.

Vitamin treatment of axenic larvae

At the time of transfer, germ-free third instar larvae were randomly assigned to the different

treatments and kept individually in 24-well plates. The larval medium consisted either of sterile

water alone (germ-free condition) or sterile water supplemented with a single vitamin or with a mix

of B vitamins (see Table 1 for reference B vitamin concentration). Sterile fish food was supplied to

each larva. The 24-well plates were placed in a climatic chamber at 80 % RH with 12:12 h light/dark

28 °C/25 °C cycle into plastic boxes covered with aluminium foil to prevent B vitamins from light-

degradation (Wang et al. 2021).

Analysis of axenic larvae developmental success in different vitamin conditions

Larvae treated with different B vitamin conditions were observed daily to track their developmental

success. For each larva, the following parameters were recorded: day of moulting to the fourth

instar, day of pupation, day of adult emergence, sex; in case of failed development to adult, the day

of death was recorded. Larvae were monitored for 14 days after transfer in the germ-free medium.

CFU quantification

For each condition, 20 µL of breeding water from 6 different wells were sampled and each sample

was transferred into 200 µL of sterile water. For each time point, sampling was randomly carried

out among the wells whose larvae were at the most advanced stage. For CFU counts, bacterial

suspensions were serially diluted in sterile water and 10 µL was spotted on LB agar plates, incubated

at 30 °C and counted 24 h later.

Growth curves

Bacteria were grown overnight in LB medium at 30 °C then diluted in LB at 1:50. Growth curves were

produced by quantifying optical density at a wavelength of 600 nm every 30 min during 24 hours in

a FLUOStar Omega plate reader (BMG Labtech). Plates were shaken before each measurement. All

the conditions were performed in triplicates (i.e. starting from three distinct colonies) and technical

duplicates.

Statistical analyses

Graphs were created with GraphPad Prism (version 10.0.2). Statistical analyses on developmental

success and sex-ratio of vitamin-treated larvae were performed with generalised linear mixed

models (GLMM) using the lme4, lmerTest and lsmeans packages in R (version 4.3.0). The replicate

was set as a random effect and an ANOVA was performed on a logistic regression (glmer). For

developmental rate analyses, a Log-rank (Mantel-Cox) test was performed in GraphPad Prism

testing the incremental proportion of pupae or adults per day. Supplemental Table S1 details

statistical information and number of individuals analysed per replicate. Numbers were rounded to

two significant figures.

Competing Interest Statement

The authors declare no competing interests.

Acknowledgments

We would like to pay our gratitude to our colleague Jean-Géraud Issaly, who passed away in December 2022 and whose work was essential for mosquito rearing and colony maintenance in the lab. We thank Eric Marois (CNRS, INSERM, University of Strasbourg) for providing the Aaeg-M strain, and Siegfried Hapfelmeier (University of Bern) for providing the *Escherichia coli* HA 416 strain. We also thank Stencey Fontenelle (Institut Pasteur de la Guyane) for technical assistance during experiments and Emmanuel Sechet for strain handling. This study is funded by the French Government's Investissement d'Avenir program, Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (grant ANR10-LABX-62-IBEID), and by ANR JCJC MosMi funding to M.G. (grant ANR-18-CE15-0007).

Author Contributions

Conceptualization, Validation – O.R., J.S.S., and M.G. Methodology: O.R., J.S.S., Y.E., F.B. and M.G Software, Formal analysis– O.R. and J.S.S. Data Curation, O.R., J.S.S., Y.E. Investigation – O.R., J.S.S., C.G., Y.E., P.F.I., and M.G. Resources, F. B. Writing – Original Draft, Visualization – O.R., M.G. Writing – Review & Editing – O.R., J.S.S., C.G., F.B. and M.G. Supervision, Project administration, Funding acquisition – M.G.

References

Benschoter CA. 1967. Effect of Dietary Biotin on Reproduction of the House Fly. *J Econ Entomol* **60**: 1326–1328. https://dx.doi.org/10.1093/jee/60.5.1326 (Accessed September 4, 2023).

Benschoter CA, Paniagua G. R. 1966. Reproduction and Longevity of Mexican Fruit Flies, Anastrepha ludens (Diptera: Tephritidae), Fed Biotin in the Diet. *Ann Entomol Soc Am* **59**: 298–300. https://dx.doi.org/10.1093/aesa/59.2.298 (Accessed September 4, 2023).

Chambers GM, Klowden MJ. CORRELATION OF NUTRITIONAL RESERVES WITH A CRITICAL WEIGHT FOR PUPATION IN LARVAL AEDES AEGYPTI iVIOSqUITONS.

Clements AN. 1992. *The biology of mosquitoes. Volume 1. Development, nutrition and reproduction.* Eds. Chapman and Hall. New York.

Cohen E, Levinson HZ. 1968. Disrupted fertility of the hidebeetle Dermestes maculatus (Deg.) due to dietary overdosage of biotin. *Experientia* **24**: 367–368. https://link.springer.com/article/10.1007/BF02140826 (Accessed September 4, 2023).

Coon KL, Vogel KJ, Brown MR, Strand MR. 2014. Mosquitoes rely on their gut microbiota for development. *Mol Ecol* **23**: 2727. /pmc/articles/PMC4083365/ (Accessed May 26, 2023).

Correa MA, Matusovsky B, Brackney DE, Steven B. 2018. Generation of axenic Aedes aegypti demonstrate live bacteria are not required for mosquito development. *Nature Communications 2018 9:1* **9**: 1–10. https://www.nature.com/articles/s41467-018-07014-2 (Accessed December 17, 2022).

Dadd RH. 1973. Insect nutrition: current developments and metabolic implications. *Annu Rev Entomol* **18**: 381–420. https://pubmed.ncbi.nlm.nih.gov/4218469/ (Accessed September 13, 2023).

Deas JB, Blondel L, Extavour CG. 2019. Ancestral and offspring nutrition interact to affect life-history traits in Drosophila melanogaster. *Proceedings of the Royal Society B: Biological Sciences* **286**. /pmc/articles/PMC6408897/ (Accessed September 14, 2023).

Douglas AE. 2017. The B vitamin nutrition of insects: the contributions of diet, microbiome and horizontally acquired genes. *Curr Opin Insect Sci* **23**: 65–69.

Fraenkel G, Blewett M. 1943. The vitamin B-complex requirements of several insects. *Biochemical Journal* **37**: 686. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1257997/ (Accessed September 4, 2023).

Goos JM, Cameron |, Swain J, Munch SB, Walsh MR. 2018. Maternal diet and age alter direct and indirect relationships between life-history traits across multiple generations. https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-2435.13258.

Lea AO, Dimond JB, Delong DM. 1956. A Chemically Defined Medium for Rearing Aedes aegypti Larvae. *J Econ Entomol* **49**: 313–315. https://dx.doi.org/10.1093/jee/49.3.313 (Accessed September 4, 2023).

Levinson HZ, Cohen E. 1973. The action of overdosed biotin on reproduction of the hide beetle, Dermestes maculatus. *J Insect Physiol* **19**: 551–558.

Lutrat C, Burckbuchler M, Olmo RP, Beugnon R, Fontaine A, Akbari OS, Argilés-Herrero R, Baldet T, Bouyer J, Marois E. 2023. Combining two genetic sexing strains allows sorting of non-transgenic males for Aedes genetic control. *Communications Biology 2023 6:1* **6**: 1–12. https://www.nature.com/articles/s42003-023-05030-7 (Accessed September 4, 2023).

Matthews BJ, Dudchenko O, Kingan SB, Koren S, Antoshechkin I, Crawford JE, Glassford WJ, Herre M, Redmond SN, Rose NH, et al. 2018. Improved reference genome of

Aedes aegypti informs arbovirus vector control. *Nature* **563**: 501–507. https://pubmed.ncbi.nlm.nih.gov/30429615/ (Accessed May 17, 2023).

Neophytou C, Pitsouli C. 2022. Biotin controls intestinal stem cell mitosis and host-microbiome interactions. *Cell Rep* **38**. http://www.cell.com/article/S2211124722002418/fulltext (Accessed September 14, 2023).

Pillai MKK, Madhukar BVR. 1969. Effect of biotin on the fertility of the yellow fever mosquito, Aedes aegypti (L.). *Naturwissenschaften* **56**: 218–219. https://pubmed.ncbi.nlm.nih.gov/5360109/ (Accessed September 4, 2023).

Piper MD. 2017. Using artificial diets to understand the nutritional physiology of Drosophila melanogaster. *Curr Opin Insect Sci* **23**: 104–111.

Reyda MR, Fugate CJ, Jarrett JT. 2009. A Complex Between Biotin Synthase and The Iron-Sulfur Cluster Assembly Chaperone HscA That Enhances In Vivo Cluster Assembly. *Biochemistry* **48**: 10782. /pmc/articles/PMC2809015/ (Accessed December 23, 2023).

Romoli O, Schönbeck JC, Hapfelmeier S, Gendrin M. 2021. Production of germ-free mosquitoes via transient colonization allows stage-specific investigation of host–microbiota interactions. *Nat Commun* **12**.

Scholz M, Albanese D, Tuohy K, Donati C, Segata N, Rota-Stabelli O. 2020. Large scale genome reconstructions illuminate Wolbachia evolution. *Nat Commun* **11**. /pmc/articles/PMC7568565/ (Accessed September 4, 2023).

Seong KH, Kang S. 2022. Noncanonical function of the Sex lethal gene controls the protogyny phenotype in Drosophila melanogaster. *Scientific Reports 2022 12:1* **12**: 1–9. https://www.nature.com/articles/s41598-022-05147-5 (Accessed December 23, 2023).

Serrato-Salas J, Gendrin M. 2023. Involvement of Microbiota in Insect Physiology: Focus on B Vitamins. *mBio* **14**. /pmc/articles/PMC9972918/ (Accessed September 4, 2023).

Singh KRP, Brown AWA. 1957. Nutritional requirements of Aedes aegypti L. *J Insect Physiol* **1**: 199–220.

Souza RS, Virginio F, Riback TIS, Suesdek L, Barufi JB, Genta FA. 2019. Microorganism-based larval diets affect mosquito development, size and nutritional reserves in the yellow fever mosquito aedes aegypti (Diptera: Culicidae). *Front Physiol* **10**. /pmc/articles/PMC6465640/ (Accessed October 18, 2023).

TRAGER W. 1948. BIOTIN AND FAT-SOLUBLE MATERIALS WITH BIOTIN ACTIVITY IN THE NUTRITION OF MOSQUITO LARVAE. *Journal of Biological Chemistry* **176**: 1211–1223.

Trager WM. 1935. ON THE NUTRITIONAL REQUIREMENTS OF MOSQUITO LARVAE (AEDES AEGYPTI). *Am J Epidemiol* **22**: 475–493. https://dx.doi.org/10.1093/oxfordjournals.aje.a118187 (Accessed September 4, 2023).

Van Schoor T, Kelly ET, Tam N, Attardo GM. 2020. Impacts of Dietary Nutritional Composition on Larval Development and Adult Body Composition in the Yellow Fever Mosquito (Aedes aegypti). *Insects* **11**: 1–15. /pmc/articles/PMC7469193/ (Accessed October 9, 2023).

Velazquez-Arellano A, Hernandez-Vazquez ADJ. 2020. Vitamins as Cofactors for Energy Homeostasis and Their Genomic Control, With Special Reference to Biotin, Thiamine, and Pantothenic Acid. *Principles of Nutrigenetics and Nutrigenomics: Fundamentals of Individualized Nutrition* 271–277.

Vendl T, Šípek P, Kouklík O, Kratochvíl L. 2018. Hidden complexity in the ontogeny of sexual size dimorphism in male-larger beetles. *Scientific Reports 2018 8:1* **8**: 1–10. https://www.nature.com/articles/s41598-018-24047-1 (Accessed December 23, 2023).

Wang Y, Eum JH, Harrison RE, Valzania L, Yang X, Johnson JA, Huck DT, Brown MR,
Strand MR. 2021. Riboflavin instability is a key factor underlying the requirement of a
gut microbiota for mosquito development. *Proc Natl Acad Sci U S A* **118**:
e2101080118. https://www.pnas.org/doi/abs/10.1073/pnas.2101080118 (Accessed
September 4, 2023).
Yan J, Kim CH, Chesser L, Ramirez JL, Stone CM. 2023. Nutritional stress compromises
mosquito fitness and antiviral immunity, while enhancing dengue virus infection
susceptibility. *Communications Biology 2023 6:1* **6**: 1–15.
https://www.nature.com/articles/s42003-023-05516-4 (Accessed December 23, 2023).
Zhou J, Lénon M, Ravanat JL, Touati N, Velours C, Podskoczyj K, Leszczynska G,
Fontecave M, Barras F, Golinelli-Pimpaneau B. 2021. Iron–sulfur biology invades
tRNA modification: the case of U34 sulfuration. *Nucleic Acids Res* **49**: 3997.
/pmc/articles/PMC8053098/ (Accessed December 23, 2023).
KEGG PATHWAY: Biotin metabolism – Microbacterium sp. SGAir0570.
https://www.kegg.jp/pathway/mics00780 (Accessed September 4, 2023a).
Vector-borne diseases. https://www.who.int/news-room/fact-sheets/detail/vector-borne-
diseases (Accessed September 4, 2023b).
18

1 **Tables**

Vitamin	Reference concentration (Singh and Brown 1957)	Solvent	Stock concentration
Biotin (B7)	0.5 µg/mL	NaOH 0.1 M	0.4 mg/mL
Choline	25 µg/mL	Water	50 mg/mL
Folic Acid (B9)	30 µg/mL	NaOH 0.1 M	50 mg/mL
Nicotinic Acid (B3)	10 µg/mL	NaOH 0.1 M	50 mg/mL
Pyridoxine (B6)	10 µg/mL	Water	50 mg/mL
Riboflavin (B2)	20 µg/mL	NaOH 0.1 M	10 mg/mL
Thiamine (B1)	10 µg/mL	Water	50 mg/mL

Table 1. List of the B vitamins tested with their reference and stock concentrations, and the

solvent used for stock solutions.

Supplemental Table S1. Statistical analyses. For each figure, the type of analysis, response variable

(dependent variable), predictor (independent variable), random effect, sample size in each

replicate, test results, and results of multiple comparisons (if applicable) are shown. Glmm:

generalised linear mixed model. Lsmeans: least square means.

Supplemental Table S2. Genes differentially regulated between male and female larvae kept in

the germ-free condition. List of genes up- (red) and down-regulated (blue) in germ-free male vs

female larvae, in the absence of vitamin supplementation. For each gene, Vectorbase ID, $\log_2$ fold

value, *padj*, description, and transcript type are indicated. The last column indicates whether the

gene was also identified in (Matthews et al. 2018) as differentially regulated in male vs female

larvae.

Supplemental Table S3. Genes differentially regulated between male and female larvae treated

with the 8x vitamin solution. List of genes up- (red) and down-regulated (blue) in germ-free male

vs female larvae treated with the 8x B vitamin solution. For each gene, Vectorbase ID, $\log_2$ fold value,

padj, description, and transcript type are indicated. The last column indicates whether the gene was

also identified in (Matthews et al. 2018)as differentially regulated in male vs female larvae.

Supplemental Table S4. Sequences of *Nix* and *Actin* primers used in this study. For each gene, the

Vectorbase accession, 5'-3' sequence, and references are indicated.

Figures

Figure 1. Development success and sex-ratio of mosquitoes exposed to B vitamins during larval

development. (**A**) Proportion of individuals completing their development (dark grey), blocked at

the larval stage (white), or dying during development (light grey) when initially reared with the

auxotrophic *E. coli* strain (AUX), reversibly colonized at the beginning of the third instar and kept

thereafter in germ-free conditions (GF) or supplemented with a 4x (VIT4x) or 8x (VIT8x) B vitamin

solution. Bar charts represent the mean ± SEM of three independent replicates of 24 to 48

mosquitoes per condition. (**B**) Proportion of male adult mosquitoes emerging from the larvae

treated in (**A**). See Supplemental Table S1 for detailed statistical information.

Figure 2. Transcriptomic analysis of male and female larvae exposed to B vitamins. (**A**) Experimental set-up: mosquito larvae reared in gnotobiotic conditions with the auxotrophic *E. coli* strain were decolonized at the beginning of the third instar (L3) to obtain germ-free larvae. Larvae were either kept in germ-free conditions or treated with 8x B vitamin solutions. After three to six hours since moulting to the fourth instar (L4), larvae were individually collected. DNA was extracted from individual larvae for sex assignment using the *Nix* gene, while RNA was extracted for transcriptome sequencing on male and female larvae. A more precise version of the experimental set up can be found in Supplemental Figure S2. (**B-E**) Reads Per Kilobase per Million mapped read (RPKM) values of representative genes differentially regulated with B vitamins in males (**B, C**) and females (**D, E**). Genes are named according to their Vectorbase ID.

Figure 3. Development success of male and female mosquito larvae exposed to high doses of single B vitamins or to increasing doses of biotin. (**A**) Proportion of mosquitoes completing their development (dark grey), blocked at the larval stage (white), or dying (light grey) when reversibly colonized at the beginning of the third instar and kept in germ-free conditions (GF) or supplemented with a 50x solution of folic acid, choline, thiamine, nicotinic acid, biotin, riboflavin, or pyridoxine. A 16x folic acid solution was also added to all tested conditions. Bar charts represent the mean ± SEM of three independent replicates except for riboflavin (two replicates) of 24 mosquitoes per condition. (**B**) Proportion of male adult mosquitoes emerging from the larvae treated in (**A**). (**C-E**) Proportion of mosquitoes (Aaeg-M strain) completing their development (dark grey), blocked at the larval stage (white), or dying (light grey) when reversibly colonized at the beginning of the third instar and kept in germ-free conditions (GF) or provided with a 1x, 4x, 8x, or 20x biotin solution. A 16x folic acid solution was also added to all tested conditions. Bar charts represent the mean ± SEM

of four independent replicates except for GF+folic acid (three replicates) of 16 to 190 mosquitoes

per sex and condition. Data in (**C**) are not sorted by sex, while data in (**D**) and (**E**) represent results

for male and female mosquitoes, respectively. See Supplemental Table S1 for detailed statistical

information.

Figure 4. Development success and sex ratio of larvae with decreased support in bacterial

metabolites. In **A-D**, auxotrophic *E. coli*-colonized larvae were treated with 1x or 0.01x

concentrations of D-Ala and *m*-DAP and 5 replicates were performed. (**A**) Number of colony forming

units (CFU)/mL in wells containing an individual. Each dot represents a well, 6 wells/time

point/replicate were tested in 5 replicates (0 h, 24 h, 72 h, 144 h and 168 h), 4 replicates (48 h) or 3

replicates (196 h, 216 h, 240 h). (**B**) Proportion of larvae reaching adulthood, (Dev), blocked in

development (Non-dev) or dead prior to reaching adulthood), (**C**) Proportion of males amongst the

adults and (**D**) number of days until reaching adulthood. In **E-F**, larvae were colonised with wt *E. coli*,

growth deficient mutant Δ*mnmA* or the Δ*iscUA* mutant, that is deficient in a biotin production. (**E**)

Proportion of larvae reaching adulthood, (Dev), blocked in development (Non-dev) or dead prior to

reaching adulthood), (**C**) Proportion of males amongst the adults. In **D**, each dot represents one

individual.

Supplemental Figure S1. Development success of mosquitoes exposed to B vitamins at 1x

concentrations during larval development. Proportion of mosquitoes completing their

development (dark grey), blocked at the larval stage (white), or dying (light grey) when reversibly

colonized until the beginning of the third instar and kept in standard germ-free conditions (GF) or

supplemented with a 1x (VIT1x) B vitamin solution. Bar charts represent the mean ± SEM of three

independent replicates of 48 mosquitoes per condition.

Supplemental Figure S2. Experimental set-up and quality check of transcriptomic analysis of male and female larvae exposed to B vitamins. (**A**) Experimental set-up: mosquito larvae reared in gnotobiotic conditions with the auxotrophic *E. coli* strain were decolonized at the beginning of the third instar (L3) to obtain germ-free larvae. Larvae were either kept in germ-free conditions or treated with 1x or 8x B vitamin solutions. After three to six hours since moulting to the fourth instar (L4), up to 150 larvae per condition were individually collected. Three independent replicates were performed. DNA was extracted from individual larvae for sex assignment using the *Nix* gene, while RNA was extracted for transcriptome sequencing on male and female larvae. A quality control step on the proportion of *Nix* reads and of overall alignment rates and a second check of initial *Nix* PCR results, allowed to select samples for data analysis. (**B**) Raw counts of read mapping on the *Nix* gene in the sequenced transcriptomes. Total number of reads mapping on the *Nix* gene in the transcriptomes of female and male larvae reversibly colonized at the beginning of the third instar and kept in germ-free conditions (GF) or supplemented with a 1x (VIT1x) or 8x (VIT8x) B vitamin solution. Results from the three replicates (A, B, and C) are shown separately. (**C-G**) Principal Component Analysis (PCA) of transcriptomes passing the quality check and used for the final analysis. (**C**) PCA of transcriptomes of female (star) and male (circle) larvae kept in germ-free conditions (white) or treated with vitamins (VIT8x, yellow). (**D, E**) PCA of transcriptomes of female (star) and male (circle) larvae kept in germ-free conditions (**D**) or treated with vitamins (**E**). (**F, G**) PCA of transcriptomes of male (**F**) and female (**G**) larvae kept in germ-free conditions (white) or treated with vitamins (yellow).

Supplemental Figure S3. Genes differentially regulated in male and female larvae exposed to B vitamins. (**A**) Number of genes up- (red) and downregulated (blue) in male vs female mosquitoes

when larvae were kept in germ-free conditions or treated with the 8x B vitamin solution. The numbers in parenthesis represent the number of genes that are shared between our study and the Matthews *et al* transcriptome (Matthews et al. 2018). (**B**) Number of genes up- (red) and downregulated (blue) in B vitamin vs germ-free conditions in male and female larvae. (**C, D**) List of genes up- (red) and down-regulated (blue) in male (**C**) and female (**D**) larvae treated with the B vitamin solution. For each gene, Vectorbase ID, $\log_2$ fold value, *padj*, description, and transcript type (coding gene: yellow; noncoding RNA (ncRNA): green; pseudogene: blue) are indicated. The last column shows paralog or ortholog genes with the name of the species and a brief function description. Asterisks (*) indicate description copied from FlyBase.

Supplemental Figure S4. Developmental rates to pupa or adult of mosquitoes exposed to increasing doses of biotin during larval development. Proportion of male (black) and female (red) larvae becoming pupae (**A**) or adult (**B**) per day when reversibly colonized at the beginning of the third instar and kept in germ-free conditions (GF) or provided with a 1x, 4x, 8x, or 20x biotin solution. A 16x folic acid solution was also added to all tested conditions. Each dot represents the mean ± SEM of four independent replicates except for GF+folic acid (three replicates) of 16 to 190 mosquitoes per sex and condition. See Supplementary Table 1 for detailed statistical information.

Supplemental Figure S5. Development success and sex ratio of larvae with decreased number of bacteria. In **A-D**, auxotrophic *E. coli*-colonized larvae were treated with 1x or 0.1x concentrations of D-Ala and *m*-DAP and 3 replicates were performed. (**A**) Number of CFUs/mL in wells containing an individual. Each dot represents a well, 6 wells/time point/replicate were tested in each replicate. (**B**) Proportion of larvae reaching adulthood, (Dev), blocked in development (Non-dev) or dead prior

1 to reaching adulthood), (**C**) Proportion of males amongst the adults and (**D**) number of days until

2 reaching adulthood. In **D**, each dot represents one individual.

4 **Figure S6. Growth curves of wt *E. coli*, growth deficient mutant Δ*mnmA* or biotin-deficient mutant**

5 **Δ*iscUA*.** Optical density of individual wells of a 96-well plate, measured every 30 min during 23 h 30

6 min. For each bacterium, 3 independent colonies were used to start the pre-cultures (referred to as

7 A, B, C), and pre-cultures were diluted to 1:50 as duplicates to start the experiment. Data show the

8 average ± SEM of the six wells per strain per time point.

A **Development success** **B** **Proportion of male adults**

A Experimental set-up

Decolonised L3 larvae → Germ-free / Vitamin 8x → Sampling 3-6h after moulting to L4 + Molecular sexing + Transcriptome analysis

B *AAEL017976* HSP70

C *AAEL013606* SRY-like

D *AAEL012340* Lipase 1 precursor

E *AAEL017067* Peritrophin-48-like

1
2

A Bacterial load

B Development success

C Proportion of male adults

D Development duration

E Development success

F Proportion of male adults

1
2