Title: Sex differences in aggression and its neural substrate in a cichlid fish

Authors: Lillian R. Jackson[1]; Mariam Dumitrascu[1]; Beau A. Alward[1,2]*

Affiliations: [1]University of Houston, Department of Psychology; [2]University of Houston, Department of Biology and Biochemistry

*Corresponding author: balward@uh.edu

Abstract

Background: Aggression is ubiquitous among social species and functions to maintains social dominance hierarchies. The African cichlid fish *Astatotilapia burtoni* is an ideal study species for studying aggression due to their unique and flexible dominance hierarchy. However, female aggression in this species and the neural mechanisms of aggression in both sexes is not well understood.

Methods: To further understand the potential sex differences in aggression in this species, we characterized aggression in male and female *A. burtoni* in a mirror assay. We then quantified neural activation patterns in regions of the social behavior network (SBN) to investigate if differences in behavior are reflected in the brain with immunohistochemistry by detecting the phosphorylated ribosome marker phospho-S6 ribosomal protein (pS6), a marker for neural activation.

Results: We found that *A. burtoni* perform both identical and sex-specific aggressive behaviors in response to a mirror assay. We observed sex differences in pS6 immunoreactivity in the Vv, a homolog of the lateral septum in mammals. Males but not females had higher ps6 immunoreactivity in the ATn after the aggression assay. The ATn is a homolog of the ventromedial hypothalamus in mammals, which is strongly implicated in the regulation of aggression in males. Several regions also have higher pS6 immunoreactivity in negative controls than fish exposed to a mirror, implicating inhibitory neurons in suppressing aggression until a relevant stimulus is present.

Conclusions: Male and female *A. burtoni* display both similar and sexually dimorphic behavioral patterns in aggression in response to a mirror assay. There are also sex differences in the corresponding neural activation patterns in the SBN. These findings suggest that distinct neural circuitry underlie aggressive behavior in male and female *A. burtoni*, serving as a foundation for future work investigating the molecular and neural underpinnings of sexually dimorphic behaviors in this species to reveal fundamental insights into understanding aggression.

Keywords: aggression, sex differences, sexual dimorphism, neural activation, social behavior network

Introduction

Aggression is a complex social behavior defined as agonistic acts directed towards conspecifics and is used to resolve conflict, maintain social dominance hierarchies, gain access to mates, and defend resources like shelter and food [1,2]. Although aggression is metabolically costly and oftentimes risky, it exists in nearly all species [3]. Extensive knowledge has been gained on the molecular and neural mechanisms of aggression in both sexes; however, male intrasexual aggression has been studied more extensively in this regard [3–6]. Moreover, our most in-depth understanding of the mechanisms of aggression come from a few traditional model organisms such as mice, rats, and fruit flies [3,3,7]. To gain a comprehensive understanding of the molecular and neural control of sexually dimorphic behaviors, models are needed in which 1) both sexes perform aggression in similar contexts and, if possible, 2) both sexes perform the same behaviors in those similar contexts [8]. Achieving both goals allows for the control of environmental and social stimuli to isolate the fundamental molecular and neural substrates that underlie aggression. However, both goals have been challenging to achieve in current models of the molecular and neural control of aggression.

The African cichlid fish *Astatotilapia burtoni* may be an especially useful species in which to gain fundamental insights into the sex differences in aggression and its molecular and neural control [9–14]. *A. burtoni* exhibit dynamic social interactions and decades of field and laboratory studies have revealed novel insights into the mechanisms governing social behavior. Male *A. burtoni* exist in two dominance states that can be transitioned between depending on the social environment [15]. Dominant males are brightly colored, have high levels of circulating sex steroid hormones (androgens and estrogens), and perform reproductive and aggressive behaviors. Aggressive behaviors such as attacks and lateral displays are distinct and easily identifiable in behavioral assays (Figure 1). In contrast, subordinate males are drably colored, have low levels of circulating sex steroid hormones, and perform submissive behaviors [16,17]. Female *A. burtoni* do not naturally form a social hierarchy and typically shoal with other females. However, when in all-female communities, females acquire certain male-typical dominance traits including enhanced aggression [18], suggesting some mechanisms regulating male-typical dominance behaviors are present in females. Still, outside of all-female communities, aggressive behavioral patterns in *female A. burtoni* have not been characterized. To definitively determine which behaviors females use during aggressive interactions, and whether they are similar to male aggressive behaviors, requires an assay in which both sexes can be studied in similar contexts.

Here, we characterize aggressive behaviors in male and female *A. burtoni* using a mirror assay. In *A. burtoni* and other model organisms like zebrafish, mirrors elicit aggressive behavioral responses because the fish treats its reflection as another fish [19,20]. We reasoned that mirror assays would be especially useful for studying aggression in *A. burtoni* since males alter their aggression if an intruder male is larger, smaller, or the same size (within 5% of other males) in length [21]. Therefore, to successfully compare aggression in males and females requires the use of a paradigm like the mirror assay, which controls for size given that their reflection should be treated as a fish identical in size to it.

We find that males and females perform both identical and sexually dimorphic aggressive behaviors during a mirror assay. We found that males and females in this assay also exhibit sexually dimorphic neural activation patterns as measured by phosphorylated ribosomes throughout brain regions of the social behavior network (SBN), a conserved, interconnected circuit critical for integrating social cues and modulating social behaviors. Our findings show suites of convergent and divergent behavioral and neural sex differences in the control of aggression in *A. burtoni*, suggesting independent mechanisms mediate sexually dimorphic behaviors in this species. These results highlight the importance of studying sex differences in diverse organisms to illuminate the fundamental mechanisms underlying the generation of social behavior and suggest *A. burtoni* may be a useful model for understanding the molecular and neural control of sexually dimorphic aggression.

Methods

Animal Subjects

111
Adult *A. burtoni* were bred from a wild-caught stock that originated from Lake Tanganyika, Africa and
housed in environmental conditions that mimic their natural equatorial habitat (28 °C; pH 8.0; 12:12 h
light/dark cycle with full spectrum illumination; constant aeration). Aquaria contained gravel-covered bottoms
with terra cotta pots cut in half to serve as shelters. Fish were fed cichlid flakes daily. All experimental
procedures were approved by the University of Houston Institutional Animal Care and Use Committee
(Protocol #202000001).
118
119 *Mirror Assay behavior recording*
120
Male (n=8) and female (n=9) *Astatotilapia burtoni* were housed in a pre-assay tank for approximately
two weeks prior to being assayed in a mirror assay. The pre-assay tank consisted of a 60.5 L plexiglass tank
with a clear perforated divider that separated males and females on separate halves. Subjects were housed
in groups of 6 fish at a time, including 3 gravid females and 3 dominant males on each side. After two weeks
in the pre-assay tank, each subject's behaviors were recorded in a mirror assay. One dominant male and one
gravid female were assayed each day until all the subjects in a pre-assay group were assayed (Figure 2).
127
The mirror assay tank consisted of a white tub (Sterilite; 400 mm x 318 mm x 12 mm) filled 6 cm deep
with UV-sterilized aquaria water with a mirror replacing one side of the assay tank. An opaque cover was first
placed on the mirror for a 15-minute habituation period to allow subjects to acclimate to the assay tank. The
opaque cover was then removed, and 30 minutes of behavior was recorded. We also included negative
control males (n= 5) and females (n= 8) in the same exact pre-assay conditions except the mirror cover was
not lifted after the 15-minute habituation period. Immediately following the mirror assay, subjects were
euthanized in an ice bath for 1-2 minutes before rapid cervical transection followed immediately by the
harvesting of tissue.
136
137 *Behavioral analysis*
138
Behavior was recorded using a digital video camera and was quantified using the freely available
BORIS (Behavioral Observation Research Interactive Software). Multiple types of behavior were quantified:
behaviors considered not to be aggressive or "neutral" (tap, graze); subordinate behaviors (flee, retreat);
aggressive behaviors (attack, lateral display, rostral display); and a male-typical reproductive behavior
(quiver) (Figure 1). Taps were defined as making gentle mouth contact with the mirror. Grazing was defined
as swimming alongside the mirror for at least 3 seconds. Fleeing was defined as an abrupt turn and swim
away from the mirror. Retreats were defined as a rapid retreat swim from the mirror. Attacks were defined as
a rapid swim towards the mirror, making mouth contact, and vigorously wiggling the body. Lateral displays
were defined as the presentation of the side of the body towards the mirror, conforming the body at times in a
slightly convex shape while lateral to the mirror with erect fins. Lateral displays were further separated into
type 1 and type 2 lateral displays to distinguish between more and less vigorous lateral displays respectively.
Rostral displays were defined as flaring the opercula towards the mirror without making direct contact with the
mirror. Quivers were defined as a rapid vibration of the body towards the mirror while conforming in a convex
shape. A group of behaviors were summed together for their mouth movements, termed mouth contact
(summed attacks, taps, and rostral displays). Additionally, total behaviors were calculated as the sum of all
behaviors and total aggression was the sum of all aggressive behaviors (attacks, lateral displays, rostral
displays, and quivers). Furthermore, another behavior was identified that appears to be associated with
aggression that we have termed an s-curve display, defined as a curved body conformation facing the mirror
at a distance.
158
Tracking and activity data was collected on pre-mirror and mirror assay videos using Noldus
EthoVisionXT software. Center-point detection was used to track the subjects throughout the assay tank.
Distance moved (cm) was calculated as the accumulated distance the subject moved throughout the assay
duration. Velocity (cm/s) was calculated as distance over time. Activity was measured as the average % of
pixels changed over time. We defined a zone close to the mirror to measure the proximity of subjects near
the mirror, extending 2.4 cm out from the mirror. Zone duration (s) was calculated as the cumulative duration
of time spent in the zone. Zone frequency was calculated as the number of times the center-point of the

subject entered the zone. Zone latency was defined as the lapse of time until the zone is first entered. Total and mean distances to the zone (cm) were calculated as the cumulative and mean distances from the zone respectively.

A series of dyad assays similar to that used recently in male *A. burtoni* [21] were also performed to ensure that female aggressive behaviors observed in the mirror assay were also performed towards a live opponent. Gravid resident females (n = 5) were housed in a 20.8 L glass tank containing gravel, half of a terra cotta pot, and constant aeration for two days before being introduced to a gravid intruder female. The intruder female was an unfamiliar conspecific that was size-matched within 5% of the resident's standard length. After the intruder female was added to the tank, the following interaction was recorded for one hour and aggressive behaviors performed by the resident female were quantified.

Inter-behavioral Intervals

We also calculated Inter-behavioral intervals as described in [22]. The log files containing the results from BORIS were analyzed for inter-behavior intervals (IBI) using custom R software (available at https://github.com/AlwardLab). IBI were determined by calculating the time between successive behaviors and averaging across all IBIs within a given log file. Negative control fish performed almost no behavior so were not included in analyses of temporal patterns of behavior.

Morphological and steroid hormone analysis

Subjects were assessed for standard length (SL), body mass (BM), gonad mass, and gonadosomatic index [GSI = (gonad mass / body mass) $*$ 100]. Male GSI ranged from 0.324 - 1.813 (average GSI = 0.895) and female GSI ranged from 1.074 - 9.122 (average GSI = 5.903).Blood samples were collected with capillary tubes from the caudal vein, centrifuged for 10 min at 8 rpm, and the plasma was removed and stored at -80°C until assayed.

Plasma 11-ketotestosterone (11-KT) levels were measured using a commercially available enzyme immunoassay (EIA) kit (Cayman Chemical Company, Ann Arbor, MI, USA) as previously described and validated for this species [23]. For the 11-KT assay, a 2-μl sample of plasma from each subject was extracted two times using 200 μl of ethyl ether and evaporated under a fume hood before re-constitution in EIA assay buffer (1:40 dilution). EIA kit protocols were then strictly followed, plates were read at 405 nm using a microplate reader (Biorad), and steroid concentrations were determined based on standard curves. All samples were assayed in duplicate.

Immunohistochemistry for pS6

Following the cervical transection of subjects, brains were fixed in 4% paraformaldehyde in PBS (pH =7.4) for 1-2 hours and then transferred to 30% sucrose in PBS. Brains were kept in the sucrose solution at 4°C until they sunk and then embedded in mounting media Neg50 and stored at -80°C. Brains were sectioned at 30 μm. We analyzed neural activation in the brain following an aggression assay vs. a negative control condition using an immunohistochemistry protocol to detect the phosphorylated ribosome marker pS6, similar to that described in Butler et al., 2018.

We performed immunohistochemistry to detect ps6, a proxy of neural activation in the brain [25] that has been used previously in *A. burtoni* brains [24,26] After sectioning, the slides were dried in a desiccator for 48 hours, then stored at -80˚C. Slides were then dried again in the desiccator for 48 hours prior to the immunohistochemistry protocol. Slides were outlined with a hydrophobic barrier using a PAP pen and then immersed in boiling citric acid (10 mM, pH = 6) for 5 minutes twice. The slides were then rinsed with 1x PBS (pH = 7.4) for 5 minutes three times. Nonspecific binding was blocked by incubating slides in 1x PBS containing 5% donkey serum and 0.3% Triton X-100 (PBS-T) for 30 minutes at room temperature. Then slides were incubated in PBS-T containing 0.5% donkey serum with the pS6 primary antibody (1:300, Invitrogen™ Phospho-S6 (Ser244, Ser247) polyclonal antibody) overnight at room temperature. Following the primary antibody incubation, slides were rinsed with 1x PBS for 5 minutes three times. Slides were then

221 incubated in PBS-T containing 0.5% donkey serum with the donkey anti-rabbit secondary antibody (1:300,
222 Invitrogen Alexa Fluor™ 488) at room temperature for 1 hour. The slides were rinsed with DAPI (1:500,
223 Sigma-Aldrich) diluted in PBS for 5 minutes and then coverslipped with aqua-poly/mount and stored at 4˚C.
224
225 *Quantification of ps6 immunoreactivity*
226
Stained slides for the pS6 immunohistochemistry were imaged using a Nikon Eclipse 80i Microscope
MicroFire™ at 20x magnification in GFP and DAPI and quantified in regions of the social behavior network
(SBN). The social behavior network consists of interconnected nuclei that are implicated in the production of
social behavior [27,28]. We quantified pS6 immunoreactivity in the following brain regions of the SBN: the
ventral part of the ventral telencephalon (Vv) (lateral septum homolog), the supracomissural part of the
ventral pallium (Vs) (medial amygdala and bed nucleus of the stria terminalis partial homolog),
subpopulations of the preoptic area (POA): parvocellular preoptic nucleus, anterior part (nPPa),
magnocellular preoptic nucleus, magnocellular division (nMMp), magnocellular preoptic nucleus,
parvocellular division (nPMp); anterior tuberal nucleus (ATn) (ventromedial hypothalamus homolog), and the
periaqueductal gray / central gray (PAG/CG). We additionally quantified pS6 immunoreactivity in the vagal
lobe (VL), vagal motor nucleus (Xm), medial part of the dorsal telencephalon, subdivision 3 (Dm-3), and the
lateral nucleus of the dorsal telencephalon, granular region (Dl-g) due to the involvement of these regions in
either processing sensory information or the social decision-making network, which is thought to evaluate the
salience of social cues [28].
241
The number of pS6 positive cells were quantified for each ROI using ImageJ. A polygon was drawn
around the ROI in each hemisphere and each pS6 positive cell within the ROI was counted manually. The
cell density for each ROI was calculated per hemisphere (cells/μm^2) and then the cell density was summed
between the hemispheres. This total cell density measure was averaged across two consecutive sections for
each region per subject, and the consecutive sections were selected at random using a random number
generator. If the subject did not have two consecutive sections for a ROI, the cell density was taken from one
section at random.
249
250 *Statistical Analysis*
251
All statistical tests were performed using Graphpad Prism version 10. Two-way ANOVAs were used to
compare 1) male and female aggressive behaviors over time in the mirror assay and 2) effects of sex and
mirror condition on behaviors and pS6 immunoreactivity between negative control and mirror males and
females. Following a significant main effect for ANOVA, Šídák's test was used for multiple comparisons.
When normality and equality of variance assumptions were not met, a log transformation of the data was
performed before conducting the two-way ANOVA. If log transformations did not correct the data to meet
assumptions, we still conducted a two-way ANOVA on the log-transformed data for consistency; this only
occurred for two dependent variables (activity and ps6 immunoreactivity in the ATn). We also performed
Fisher's tests on the frequency of performance of rostral displays and quivers. Effects were considered
significant at $p \leq 0.05$. For several behaviors, log transformations were not feasible due to zeros in the
negative control groups. For these behaviors, two-way ANOVAs were conducted, and additional chi-square
tests were run.
264
265 **Results**
266
267 *Male and female A. burtoni perform the same and different aggressive behaviors towards a mirror*
268
269 Contour maps of fish movement over time that were generated in Noldus showed that before the
270 opaque cover was lifted fish swim around the edges of the assay arena (Fig. 3a). However, fish spent the
271 vast majority of their time in front of the mirror once the cover was removed.
272
273 There was a significant effect of time but not sex or an interaction on attacks (Fig. 3b; attacks, Two-
274 way ANOVA, sex: $p = 0.3885$, $F_{(1, 15)} = 0.7888$, time: $p < 0.0001$, $F_{(2.435, 36.53)} = 12.60$, interaction: $p = 0.3688$
275 $F_{(3, 45)} = 1.076$). There was a significant effect of sex and time but not an interaction on lateral displays (Fig.

3c; lateral displays, Two-way ANOVA, sex: p = 0.0399, $F_{(1, 15)}$ = 5.059, time: p = 0.0126, $F_{(2.353, 35.29)}$ = 4.618, interaction: p = 0.0805, $F_{(3, 45)}$ = 2.397), wherein males performed more lateral displays than females. We also observed a significant effect of sex on rostral displays (Fig. 3d; rostral displays, Two-way ANOVA, sex: p = 0.0075, $F_{(1, 15)}$ = 9.537, time: p = 0.1724, $F_{(1.529, 22.93)}$ = 1.946, interaction: p = 0.1357, $F_{(3, 45)}$ = 1.946), where males produced more rostral displays than females. Additionally, there was a significant effect of sex on quivers (Fig. 3e; quivers, Two-way ANOVA, sex: p = 0.0413, $F_{(1, 15)}$ = 4.982, time: p = 0.0882, $F_{(1.306, 19.59)}$ = 3.038, interaction: p = 0.0530, $F_{(3, 45)}$ = 2.760). Specifically, females performed more quivers than males. Fisher's exact tests showed more males than females performed rostral displays (6/8 males versus 1/9 females; p=0.0152) while more females than males performed quivers (2/8 males versus 8/9 females; p=0.0152). We were surprised to observe females perform a male-typical mating behavior, quivers, in the mirror assay, given the behavior is putatively used as an aggressive display in a mirror assay by females. To confirm females do these behaviors towards another female, we conducted resident intruder tests with focal resident females against intruder females [21]. Female residents performed quivers, lateral displays, and attacks towards intruder females and also chased them (Fig. S1), indicating quivers are a normal component of the behavioral repertoire of female-female aggression. There were no sex differences in median or mean IBI between mirror males and females (Fig. S2a; Median IBI, Unpaired t-test, p = 0.1724, t = 1.433, df = 15, Fig. S2b; Mean IBI, Mann-Whitney, p = 0.0927, U = 18).

We compared the behaviors of the NC fish to mirror fish and found across all behaviors NC males and females performed no or very few acts of any behavior types (Fig. S3). Overall, Two-way ANOVAs revealed consistent effects of condition and not sex or an interaction on behaviors performed during the mirror assay (See Supplementary Results for detailed statistical findings; Fig. S3). This shows the NC fish included in this assay are sufficient as a control group when comparing ps6 immunoreactivity to fish exposed to the mirror.

There was a significant effect of condition but not sex or an interaction on latency to behave (Fig. S3g; latency to behave, Two-way ANOVA, sex: p = 0.2149, $F_{(1, 26)}$ = 1.616, condition: p = 0.0408, $F_{(1, 26)}$ = 4.632, interaction: p = 0.0679, $F_{(1, 26)}$ = 3.630), wherein mirror males had a longer latency to behave than negative control males (Fig. S3g; latency to behave, Male Mirror – NC, p = 0.0259, t = 2.666, df = 26). There were no significant effects of sex, condition, or sex*condition on latency to aggression (Fig. S3h; latency to aggression, Two-way ANOVA, sex: p = 0.8680, $F_{(1, 17)}$ = 0.02848, condition: p = 0.7141, $F_{(1, 17)}$ = 0.1388, interaction: p = 0.1165, $F_{(1, 17)}$ = 2.735). Additionally, there were no significant effects of sex, condition, or sex*condition on distance moved and velocity (Fig. S4a; distance moved, Two-way ANOVA of log transformation, sex: p = 0.8537, $F_{(1, 26)}$ = 0.03466, condition: p = 0.7970, $F_{(1, 26)}$ = 0.06752, interaction: p = 0.8036, $F_{(1, 26)}$ = 0.06312, Fig. S4b; velocity, Two-way ANOVA of log transformation, sex: p = 0.7945, F (1, 26) = 0.06927, condition: p = 0.9439, F (1, 26) = 0.005043, interaction: p = 0.7528, F (1, 26) = 0.1013).

We observed a significant effect of condition but not sex or an interaction on activity (Fig. S4c; activity, Two-way ANOVA of log transformation, sex: p = 0.2757, $F_{(1, 25)}$ = 1.242, condition: p = 0.0013, $F_{(1, 25)}$ = 13.13, interaction: p = 0.5318, $F_{(1, 25)}$ = 0.4020). One mirror female outlier was removed. Mirror females have higher activity than negative control females and males follow this same pattern although not significant (Fig. S4c, activity, Šídák's test, Female Mirror – NC, p = 0.0072, t = 3.214, df = 25). No significant effects of sex, condition or sex*condition were observed on zone frequency or zone latency (Fig. S4d; zone frequency, Two-way ANOVA of log transformation, sex: p = 0.7472, $F_{(1, 26)}$ = 0.1061, condition: p = 0.8358, $F_{(1, 26)}$ = 0.04381, interaction: p = 0.3929, $F_{(1, 26)}$ = 0.7549, Fig. S4f; zone latency, Two-way ANOVA, sex: p = 0.1351, $F_{(1, 26)}$ = 2.378, condition: p = 0.1573, $F_{(1, 26)}$ = 2.121, interaction: p = 0.1452, $F_{(1, 26)}$ = 2.256). There was a significant effect of condition but not sex or an interaction on zone duration (Fig. S4e; zone duration, Two-way ANOVA, sex: p = 0.0907, $F_{(1, 26)}$ = 3.086, condition: p = 0.0369, $F_{(1, 26)}$ = 4.841, interaction: p = 0.2392, $F_{(1, 26)}$ = 1.451) and mirror females had a longer zone duration than negative control females while males did not differ by condition (Fig. S4e, zone duration, Šídák's test, Female Mirror – NC, p = 0.0285, t = 2.624, df = 26).

Condition—but not sex or an interaction—had an effect on on mean distance and total distance to the zone (Fig. S4g; mean distance to the zone, Two-way ANOVA of log transformation, sex: p = 0.2036, $F_{(1, 26)}$ = 1.701, condition: p = 0.0032, $F_{(1, 26)}$ = 10.53, interaction: p = 0.7699, $F_{(1, 26)}$ = 0.08735, Fig. S4h; total distance to the zone, Two-way ANOVA of log transformation, sex: p = 0.2308, $F_{(1, 26)}$ = 1.506, condition: p = 0.0030, $F_{(1, 26)}$ = 10.71, interaction: p = 0.7529, $F_{(1, 26)}$ = 0.1012). Mirror females had a higher mean and total distance

to the zone than negative control females and males followed this pattern but was not statistically significant (Fig. S4g, mean distance to the zone, Šídák's test, Female Mirror – NC, $p = 0.0223$, $t = 2.730$, df = 26, Fig. S4h; total distance to the zone, Šídák's test, Female Mirror – NC, $p = 0.0204$, $t = 2.768$, df = 26).

Sex differences and similarities in neural activation patterns of the SBN after aggression

No significant effects of sex, condition, or sex*condition on ps6 immunoreactivity were observed in the POA or subregions nPPa and nPMp, as well as in the Dl-g, Dm-3, and Xm (Fig. 4a; POA, Two-way ANOVA, sex: $p = 0.3719$, $F_{(1, 25)} = 0.8266$, condition: $p = 0.1268$, $F_{(1, 25)} = 2.494$, interaction: $p = 0.9556$, $F_{(1, 25)} = 0.003165$, Fig. S5a; nPPa Two-way ANOVA, sex: $p = 0.1497$, $F_{(1, 25)} = 2.210$, condition: $p = 0.1101$, $F_{(1, 25)} = 2.744$, interaction: $p = 0.8910$, $F_{(1, 25)} = 0.01917$, Fig. S5c; nPMp, Two-way ANOVA, sex: $p = 0.4148$, $F_{(1, 19)} = 0.6952$, condition: $p = 0.5210$, $F_{(1, 19)} = 0.4276$, interaction: $p = 0.5545$, $F_{(1, 19)} = 0.3621$, Fig. S5d; Dl-g, Two-way ANOVA, sex: $p = 0.1271$, $F_{(1, 24)} = 2.497$, condition: $p = 0.8580$, $F_{(1, 24)} = 0.03271$, interaction: $p = 0.2951$, $F_{(1, 24)} = 1.146$, Fig. S5e; Dm-3, Two-way ANOVA, sex: $p = 0.3922$, $F_{(1, 25)} = 0.7580$, condition: $p = 0.5113$, $F_{(1, 25)} = 0.4440$, interaction: $p = 0.0919$, $F_{(1, 25)} = 3.073$, Fig. S5f; Xm, Two-way ANOVA of log transformation, sex: $p = 0.1418$, $F_{(1, 23)} = 2.314$, condition: $p = 0.1078$, $F_{(1, 23)} = 2.800$, interaction: $p = 0.9505$, $F_{(1, 23)} = 0.003933$). However, there was a significant effect of condition, but not sex or an interaction, on pS6 immunoreactivity in one subregion of the POA, the nMMp (Fig. S5b; nMMp, Two-way ANOVA, sex: $p = 0.7492$, $F_{(1, 24)} = 0.1046$, condition: $p = 0.0400$, $F_{(1, 24)} = 4.718$, interaction: $p = 0.9390$, $F_{(1, 24)} = 0.005974$) in which negative controls had higher ps6 immunoreactivity in this region on average, but this was not statistically significant. There was a significant effect of sex on ps6 immunoreactivity in the Vv, where females had higher pS6 immunoreactivity compared to males (Fig. 4b; Vv, Two-way ANOVA, sex: $p = 0.0010$, $F_{(1, 23)} = 14.21$, condition: $p = 0.3178$, $F_{(1, 23)} = 1.043$, interaction: $p = 0.0624$, $F_{(1, 23)} = 3.834$).

We observed a significant effect of condition on ps6+ cells in the Vs, PAG/CG, and the VL (Fig. 4c; Vs, Two-way ANOVA, sex: $p = 0.0793$, $F_{(1, 25)} = 3.347$, condition: $p = 0.0040$, $F_{(1, 25)} = 10.03$, interaction: $p = 0.1571$, $F_{(1, 25)} = 2.127$, Fig. 4d; PAG, Two-way ANOVA of log transformation, sex: $p = 0.4585$, $F_{(1, 22)} = 0.5695$, condition: $p = 0.0316$, $F_{(1, 22)} = 5.272$, interaction: $p = 0.5344$, $F_{(1, 22)} = 0.3985$, Fig. 4e; CG, Two-way ANOVA, sex: $p = 0.2459$, $F_{(1, 25)} = 1.412$, condition: $p = 0.0009$, $F_{(1, 25)} = 14.06$, interaction: $p = 0.4308$, $F_{(1, 25)} = 0.6412$, Fig. 4f; VL, Two-way ANOVA, sex: $p = 0.3216$, $F_{(1, 23)} = 1.026$, condition: $p = 0.0021$, $F_{(1, 23)} = 11.96$, interaction: $p = 0.9752$, $F_{(1, 23)} = 0.0009858$). Specifically, in the Vs, PAG/CG, and the VL, female negative controls had higher pS6 immunoreactivity than mirror females while males did not differ by condition (Fig. 4c, Vs, Šídák's test, Female Mirror – NC, $p = 0.0020$; Fig. 4d, PAG, Šídák's test, Female Mirror – NC, $p = 0.0408$; Fig. 4e, CG, Šídák's test, Female Mirror – NC, $p = 0.0024$, df = 25, Fig. 4f, VL, Šídák's test, Female Mirror – NC, $p = 0.0181$). Mirror males and negative controls did not differ in ps6 immunoreactivity in the PAG and CG (Fig. 4d, PAG, Šídák's test, Male Mirror – NC, $p = 0.5310$; Fig. 4e, CG, Šídák's test, Male Mirror – NC, $p = 0.1378$). Although not statistically significant, male negative controls had higher pS6 immunoreactivity in the VL compared to mirror males (Fig. 4f, VL, Šídák's test, Male Mirror – NC, $p = 0.0773$, $t = 2.184$, df = 23), following the same trend as females in the VL.

A significant sex*condition interaction on ps6 immunoreactivity in the ATn was observed (Fig. 5a; ATn, Two-way ANOVA of log transformation, sex: $p = 0.5059$, $F_{(1, 24)} = 0.4562$, condition: $p = 0.4089$, $F_{(1, 24)} = 0.7065$, interaction: $p = 0.0040$, $F_{(1, 24)} = 10.14$). Specifically, males exposed to the mirror had more ps6+ cells in the ATn compared to negative control males, while this was not true for females (Fig. 5a, ATn, Šídák's test, Male Mirror – NC, $p = 0.0313$, Female Mirror – NC, $p = 0.1467$).

No effects of sex or condition on 11KT

There were no significant effects of sex, condition or sex*condition on 11-KT (Fig. S6a; 11-KT, Two-way ANOVA, sex: $p = 0.6283$, $F_{(1, 26)} = 0.2400$, condition: $p = 0.1882$, $F_{(1, 26)} = 1.826$, interaction: $p = 0.9053$, $F_{(1, 26)} = 0.01443$).

Discussion

We aimed to compare male and female intrasexual aggression in *A. burtoni*, a non-traditional model organism of social behavior that has the potential to yield novel insights into the sexually dimorphic control of social behavior. We determined that in an identical context that elicits aggression, male and female *A. burtoni* perform a mixture of identical behaviors and sex-specific behaviors. Interestingly, quivers, which are performed by males in reproductive contexts, were performed by females in the mirror assay, suggesting it is used as an aggressive signal by females. As in other species in which females adopt male-like mounting behaviors towards other females to assert dominance, perhaps in teleosts this behavior maintains female social status [29].

Intrasexual male aggression is thought to follow specific "rules of engagement" between opponents, assessing the resource holding power of the other to minimize risk in conflict [30]. Since *A. burtoni* males alter their aggression depending on the relative size of their opponent, we can safely assume that intrasexual male aggression possesses a level of opponent assessment. In our study, we find that males exposed to a mirror have a longer latency to behave compared to negative control males. Perhaps this longer duration of time to investigate the mirror serves as a lengthy assessment period since the male is faced with an identically-sized opponent. However, we don't see this difference in female *A. burtoni* exposed to a mirror vs. negative controls. Perhaps intrasexual female aggression in *A. burtoni* does not follow the same opponent assessment as it does for *A. burtoni* males, supported by a study in Texas cichlids that suggests that males and females use a different set of rules to settle conflict [31].

We also revealed a significant difference in condition in activity and zone duration in which mirror females have higher activity and spend more time in the zone near the mirror compared to negative control females. Although not statistically significant, mirror males also have higher activity levels on average than negative controls. Since the change in the environment between the pre-mirror and mirror assay is more pronounced in trials of fish exposed to a mirror compared to the negative controls (opaque cover removed vs. remained on), we expect to see mirror subjects behaving more actively in their assay. Additionally, the mirror allows subjects to be more engaging with their reflection compared to an opaque cover which may be manifesting here in activity levels. Mirror females spending more time near the mirror may reflect an aspect of exploration or investigation that is not necessary for males in this context. In fact, female rats show more exploratory activity than males in a novel environment and have shorter approach latencies to novelty [32]. Perhaps this aspect of elevated exploration in females is consistent with a longer zone duration in mirror females that we have shown here.

We found that 11-KT did not differ between mirror and negative control conditions, which is in line with findings in other teleost species [33,34] but contrasts with previous findings in male *A. burtoni* where mirror assays were also used [20]. However, as argued by Oliveira and Canario [35], the previous mirror assay study in male *A. burtoni* [20] has a few issues that could explain the higher 11-KT in mirror fish compared to negative controls, including the confound of exposure to a competing male's chemical cues in the same territory directly before exposure to a mirror. Indeed, the lack of a rise in androgens in a mirror assay that we observed in both males and females has been explained by the fact that there is no clear winner of the fight as is seen against a live opponent because all aggressive acts are perceived to elicit an equal counter response [14,33–36].

In multiple brain regions we observed impacts of sex and exposure to the mirror on neural activation as measured by ps6 immunoreactivity. For several brain regions no differences in pS6 activation between negative control and mirror fish, including the POA (nPPa and nPMp), Dl-g, Dm-3, and Xm, were observed. The POA plays a role in aggressive and reproductive behaviors across vertebrates [28]. In *A. burtoni*, there is an increase in *egr1* expression in the nMMp in males after fighting but not after courting, emphasizing the role of this POA subregion in modulating aggression [37]. When we investigated the pS6 immunoreactivity in the POA we expected to see higher pS6 immunoreactivity in the POA in fish exposed to the mirror compared to the negative control condition. Surprisingly, we see no differences in pS6 immunoreactivity as a function of sex or condition for two of the subregions of the POA (nPPa and nPMp), and although we find a significant effect of condition on pS6 immunoreactivity in the nMMp, it was higher in negative controls. Another study in *A. burtoni* found no effect of social stimulus on c-Fos expression in the POA when males were exposed to reproductive or aggressive contexts [38]. Interestingly, the males in the aforementioned study were not

engaged in full contact with stimulus fish and therefore not receiving all sensory signals. Hence, it may be that the POA neurons activate in response to integrating several multisensory components of social information, but solely the visual input and behavioral output we have observed in this study is not sufficient for the activation of these neurons.

The Dl-g (homolog of the medial pallium; hippocampus) is involved in processing sensory input and receiving and processing socially relevant hydrodynamic cues in *A. burtoni* [39]. The Dm (homologous to the basolateral amygdala, blAMY) also receives sensory input including lateral line information [40–42]. pS6 immunoreactivity in the Dl-g, Dm-3, and Xm did not differ by sex or condition and may indicate that mechanosensory input is similar in the negative control condition and mirror conditions despite distinct differences in behavioral output. Some behavioral measures (distance moved (cm) and velocity (cm/s)) do not differ by sex or condition and may provide evidence of similar mechanosensory input between these groups. It is important to note that the lack of socially relevant hydrodynamic cues that accompany a fight with a real opponent are not present in the mirror assay, and thus regions like the Dl-g, Dm-3, and Xm that process sensory information may not be relevant in an aggressive assay lacking this dynamic sensory input.

One region that did display a sex difference in pS6 immunoreactivity is the Vv. The Vv, homologous to the LS, is thought to integrate internal state with the social environment since this region is shared between the social behavior network and the mesolimbic reward system [28]. The Vv has also been implicated in processing social information in anxiety-like contexts. In previous work, female *A. burtoni* had higher *cfos* expression in the Vv after watching a preferred male lose a fight [43]. In this study, we found higher pS6 immunoreactivity in the Vv of females compared to males regardless of condition, which may indicate that the mirror assay is more anxiogenic to females, or that the Vv is more active overall in females than it is in males. Future experiments aimed at testing the role of the Vv in explicitly anxiogenic behavior paradigms in both males and females are thus warranted.

The ventromedial hypothalamus (VMH) has been implicated in mediating aggressive behaviors across several species [6]. In mammals, *cfos* expression in the VMH increases after aggressive interactions in males, and specifically the causal role of the ventrolateral portion of the VMH (VMHvl) in aggression in male mice is well established [3,44–46]. In *A. burtoni*, the ATn (homologous to the VMH) is likely involved in processing mechanosensory information in male-male aggressive contexts [39]. In our study, pS6 immunoreactivity in the ATn is higher in males exposed to a mirror than in negative controls, supporting the importance of activity in the ATn in mediating male-male aggression. While ps6 immunoreactivity was not higher in mirror versus control females, this does not preclude the ATn potentially being involved in female aggression in a environment-specific manner. For example, ATn neurons may become activated during female aggression in particular contexts and/or a specific sub-population of neurons in the ATn that remains to be identified regulates female aggression as the ventrolateral portion of the VMH does in male mice. Investigations into whether a specific locus within the fish VMH regulates aggression in both male and female *A. burtoni* will be key in disentangling the neural basis of aggression in this species.

Some regions of interest had higher neural activation in negative controls than in mirror fish. These results suggest the potential involvement of inhibitory neurons in suppressing activity in specific aggression circuits. Glutamic acid decarboxylase (Gad) is the rate-limiting enzyme that converts glutamate (an excitatory neurotransmitter) into GABA (an inhibitory neurotransmitter). Gad exists in two highly conserved isoforms, *gad1* and *gad2*. In *A. burtoni*, the Vs, CG, and VL express both *gad1* and *gad2* [47]. The Vs, PAG/CG and VL show higher pS6 immunoreactivity in negative controls and since *gad1* and *gad2* are both expressed in these regions, we postulate that the inhibitory neurons in these regions are important for suppressing aggressive behaviors. We hypothesize that inhibitory neurons containing GABA are activated in negative control fish whereas lower activation of these neurons permits aggressive behaviors in mirror fish.

Perspectives and significance

Our results revealed both similar and sexually dimorphic behavioral patterns underlying aggressive responses to a mirror in male and female *A. burtoni.* We also observed extensive sex differences in neural activation throughout regions of the SBN in response to our aggression assay, suggesting distinct neurons

and neural circuitry underlie aggressive behavior in male and female *A. burtoni.* Interestingly, neural activation in the ATn reflected aggressive state in males but not females, suggesting this region controls aggression in male *A. burtoni* as it does in mice [3,7,45,48]. We also provide evidence that the inhibition of neural activity in a variety of brain regions in the SBN is associated with aggression in *A. burtoni* as well. These findings overall lay a foundation of testable hypotheses for future studies to understand the molecular and neural regulation of sexually dimorphic behaviors in *A. burtoni*.

Gaining a fundamental understanding of the molecular and neural basis of aggression requires studies in both sexes in which the conditions and behaviors performed by either sex are similar. In our mirror assay we were able to stimulate aggressive responses in both males and females. The behaviors performed by males were the same as the suite of behaviors they perform in dyadic social interactions and we confirmed that this is the case for females as well. A significant benefit of assessing aggression in a setup that controls for the environmental and social conditions in males and females is it is presumed that while 1) both sexes are experiencing an elevated motivation to behave aggressively, they are 2) also performing distinct and similar aggressive behavioral acts. Thus, a benefit for the use of *A. burtoni* as a model organism for elucidating the molecular and neural mechanisms governing sexually dimorphic aggression is that the molecular and neural mechanisms of the motivation versus sensorimotor aspects of aggression in both sexes can be studied in the same environment. Therefore, studying sex differences in aggression in *A. burtoni* may lead to new ideas on the fundamental mechanisms controlling social behavior generally, yielding novel hypotheses that can be tested across species.

Supplementary Information

Supplementary results and figures are in the Supplementary Information file.

Availability of data and materials

The datasets generated and/or analyzed during the current study are available in the manuscript itself, the Supplementary Information file, and online at https://github.com/AlwardLab.

Declarations

Ethics approval and consent to participate

Experimental procedures were conducted according to the ethical guidelines for the care and use of laboratory animals. Experiments were approved by the University of Houston Institutional Animal Care and Use Committee (Protocol #202000001).

Consent for publication

Not applicable

Competing interests

The authors declare no competing or financial interests.

Funding

This research was supported by a Beckman Young Investigator Award from the Arnold and Mabel Beckman Foundation, an NIH grant R35GM142799, and a University of Houston-National Research University Fund startup R0503962 to B.A.A.

Acknowledgements

549   The authors thank Andrew Hoadley for procedural assistance as well as Mariana Lopez for helpful
550   discussions on interpreting findings.

Authors' contributions

552
553

554   B.A.A. and L.R.J. designed the study; L.R.J. performed the experiments; L.R.J. and M.D. collected
555   the data; B.A.A. and L.R.J. analyzed the data; L.R.J. made the figures. B.A.A. and L.R.J. wrote and
556   edited the manuscript.

References

558
559

1. Adkins-Regan E. Hormones and Animal Social Behavior. Princeton University Press; 2005.

2. Nelson R, Kriegsfeld LJ. An Introduction to Behavioral Endocrinology [Internet]. Sinauer; 2017. Available
from: https://books.google.com/books?id=pBiBjgEACAAJ

3. Lischinsky JE, Lin D. Neural mechanisms of aggression across species. Nat Neurosci. 2020;23.

4. Bayless DW, Shah NM. Genetic dissection of neural circuits underlying sexually dimorphic social
behaviours. Philos Trans R Soc B Biol Sci. 2016;371:1–10.

5. Been LE, Gibbons AB, Meisel RL. Towards a neurobiology of female aggression. Neuropharmacology.
2019;156:107451.

6. Nelson RJ, Trainor BC. Neural mechanisms of aggression. Nat Rev Neurosci. 2007;8:536–46.

7. Anderson DJ. Circuit modules linking internal states and social behaviour in flies and mice. Nat Rev
Neurosci. 2016;17:692–704.

8. McCarthy MM, Arnold AP, Ball GF, Blaustein JD, De Vries GeertJ. Sex Differences in the Brain: The Not
So Inconvenient Truth. J Neurosci. 2012;32:2241–7.

9. Alward BA, Hoadley AP, Jackson LR, Lopez MS. Genetic dissection of steroid-hormone modulated social
behavior: Novel paralogous genes are a boon for discovery. Horm Behav. 2023;147:105295.

10. Fernald RD. Social control of the brain. Annu Rev Neurosci. 2012;35:133–51.

11. Fernald RD, Maruska KP. Social information changes the brain. Proc Natl Acad Sci. 2012;109:17194–9.

12. Jackson LR, Lopez MS, Alward B. Breaking Through the Bottleneck: Krogh's Principle in Behavioral
Neuroendocrinology and the Potential of Gene Editing. Integr Comp Biol. 2023;63:428–43.

13. Maruska KP, Fernald RD. Astatotilapia burtoni: A Model System for Analyzing the Neurobiology of
Behavior. ACS Chem Neurosci. 2018;9:1951–62.

14. Munley KM, Alward BA. Control of social status by sex steroids: insights from teleost fishes. Mol Psychol
Brain Behav Soc. 2023;2:21.

15. Maruska KP, Fernald RD. Social Regulation of Male Reproductive Plasticity in an African Cichlid Fish.
Integr Comp Biol. 2013;53:938–50.

16. Maruska KP. Social Transitions Cause Rapid Behavioral and Neuroendocrine Changes. Integr Comp
Biol. 2015;55:1–13.

17. Maruska KP, Butler JM. Reproductive- and Social-State Plasticity of Multiple Sensory Systems in a
Cichlid Fish. Integr Comp Biol. 2021;61:249–68.

18. Renn SCP, Fraser EJ, Aubin-Horth N, Trainor BC, Hofmann HA. Females of an African cichlid fish display
male-typical social dominance behavior and elevated androgens in the absence of males. Horm Behav.
2012;61.

19. Ariyomo TO, Watt PJ. Aggression and sex differences in lateralization in the zebrafish. Anim Behav.
2013;86:617–22.

20. Desjardins JK, Fernald RD. What do fish make of mirror images? Biol Lett. 2010;6.

21. Alward BA, Cathers PH, Blakkan DM, Hoadley AP, Fernald RD. A behavioral logic underlying aggression
in an African cichlid fish. Ethology. 2021;127:572–81.

22. Alward BA, Hilliard AT, York RA, Fernald RD. Hormonal regulation of social ascent and temporal patterns
of behavior in an African cichlid. Horm Behav. 2019;107:83–95.

23. Maruska KP, Fernald RD. Steroid receptor expression in the fish inner earvaries with sex, social status,
and reproductive state. BMC Neurosci. 2010;11.

24. Butler JM, Whitlow SM, Roberts DA, Maruska KP. Neural and behavioural correlates of repeated social
defeat. Sci Rep. 2018;8:6818.

25. Knight ZA, Tan K, Birsoy K, Schmidt S, Garrison JL, Wysocki RW, et al. Resource Molecular Profiling of
Activated Neurons by Phosphorylated Ribosome Capture. Cell. 2012;151:1126–37.

26. Maruska KP, Butler JM, Field KE, Forester C, Augustus A. Neural Activation Patterns Associated with
Maternal Mouthbrooding and Energetic State in an African Cichlid Fish. Neuroscience [Internet]. 2020;
Available from: https://doi.org/10.1016/j.neuroscience.2020.07.025

27. Newman SW. The Medial Extended Amygdala in Male Reproductive Behavior A Node in the Mammalian
Social Behavior Network. Ann N Y Acad Sci. 1999;877:242–57.

28. O'Connell LA, Hofmann HA. The Vertebrate mesolimbic reward system and social behavior network: A
comparative synthesis. J Comp Neurol. 2011;519.

29. FANG J, CLEMENS LG. Contextual determinants of female–female mounting in laboratory rats. Anim
Behav. 1999;57:545–55.

30. Parker GA. Assessment strategy and the evolution of fighting behaviour. J Theor Biol. 1974;47:223–43.

31. Draud M. Female and male Texas cichlids (Herichthys cyanoguttatum) do not fight by the same rules.
Behav Ecol. 2004;15:102–8.

32. Williams, DI, Wells PA. Differences in home-cage-emergence in the rat in relation to infantile handling.
Psychon Sci. 1970;18:168–9.

33. Oliveira RF, Carneiro LA, Canário AVM. No hormonal response in tied fights. Nature. 2005;437:207–8.

34. Oliveira RF. Social behavior in context: Hormonal modulation of behavioral plasticity and social
competence. Integr Comp Biol. 2009;49:423–40.

35. Oliveira RF, Canário AVM. *Nemo* through the looking-glass: a commentary on Desjardins & Fernald. Biol
Lett. 2011;7:487–8.

36. Hirschenhauser K, Wittek M, Johnston P, Möstl E. Social context rather than behavioral output or winning
modulates post-conflict testosterone responses in Japanese quail (Coturnix japonica). Physiol Behav.
2008;95:457–63.

37. Loveland JL, Fernald RD. Differential activation of vasotocin neurons in contexts that elicit aggression
and courtship. Behav Brain Res. 2017;317:188–203.

38. O'Connell LA, Rigney MM, Dykstra DW, Hofmann HA. Neuroendocrine Mechanisms Underlying Sensory
Integration of Social Signals. J Neuroendocrinol. 2013;25:644–54.

39. Butler JM, Maruska KP. The Mechanosensory Lateral Line System Mediates Activation of Socially-
Relevant Brain Regions during Territorial Interactions. Front Behav Neurosci [Internet]. 2016 [cited 2023 Sep
28];10. Available from: http://journal.frontiersin.org/Article/10.3389/fnbeh.2016.00093/abstract

40. Folgueira M, Anadón R, Yáñez J. Experimental study of the connections of the telencephalon in the
rainbow trout (*Oncorhynchus mykiss*). II: Dorsal area and preoptic region: Telencephalic Connections in
Trout. J Comp Neurol. 2004;480:204–33.

41. Yamamoto N, Ito H. Fiber connections of the anterior preglomerular nucleus in cyprinids with notes on
telencephalic connections of the preglomerular complex. J Comp Neurol. 2005;491:212–33.

42. Yamamoto N, Ito H. Visual, lateral line, and auditory ascending pathways to the dorsal telencephalic area
through the rostrolateral region of the lateral preglomerular nucleus in cyprinids. J Comp Neurol.
2008;508:615–47.

43. Desjardins JK, Klausner JQ, Fernald RD. Female genomic response to mate information. Proc Natl Acad
Sci. 2010;107.

44. Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, et al. Functional identification of an aggression
locus in the mouse hypothalamus. Nature. 2011;470:221–6.

45. Yang T, Yang CF, Chizari MD, Maheswaranathan N, Burke KJ, Borius M, et al. Social Control of
Hypothalamus-Mediated Male Aggression. Neuron. 2017;95:955-970.e4.

46. Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, et al. Functional identification of an aggression
locus in the mouse hypothalamus. Nature. 2011;470:221–6.

47. Maruska KP, Butler JM, Field KE, Porter DT. Localization of glutamatergic, GABAergic, and cholinergic
neurons in the brain of the African cichlid fish, *Astatotilapia burtoni*. J Comp Neurol. 2017;525:610–38.

48. Lee H, Kim D, Remedios R, Anthony TE, Chang A, Madisen L, et al. Scalable control of mounting and
attack by Esr1+ neurons in the ventromedial hypothalamus. Nature. 2014;509:627–32.

Figure 1. Typical behaviors characterized in *A. burtoni*. Aggressive and reproductive behaviors are typically performed by dominant males. Submissive behaviors are performed by subordinate males.

Pre-assay: 2 weeks

Day 1 | Day 2 | Day 3

Mirror Assay

Assay 1 ♂ and 1 ♀ each day (alternating order)

Figure 2. Pre-assay schematic for negative control and mirror fish. Males and females were housed in a tank separated by a perforated divider for 2 weeks. Following this pre-assay period, males and females were assayed in pairs on consecutive days in a mirror assay.

666
667
668
669
670 Figure 3. Male and female *A. burtoni* perform different aggressive behaviors. (a) Representative track
671 visualization of male and female subjects before (PRE) and throughout (Mirror) a mirror assay trial. (b) Males
672 and females both perform attacks in a mirror assay. (c)(d) Males perform more lateral displays and rostral
673 displays than females. (e) Females perform more quivers than males.

680
681
682 Figure 4. Sex differences and similarities in neural activation patterns. (a) pS6 cell density does not differ by
683 condition or sex in the POA. (b) Females have higher pS6 immunoreactivity in the Vv than males. (c)(d)(e)(f)
684 pS6 cell density is higher in negative control fish than in mirror fish in the Vs, PAG, CG, and the VL. Multiple
685 comparisons reveal that pS6 cell density is higher in negative control females compared to mirror females for
686 these regions. NC, negative control. ** $p < 0.01$, * $p < 0.05$. ns, not statistically significant.
687

688
689
690 Figure 5. Males and females have opposite neural activation patterns in the ATn. (a) pS6 cell density in the
691 ATn has an interaction effect between sex and condition. (b) Representative section of the ATn. (c)
692 Representative sections of the ATn stained with pS6 antibody to detect neural activation show lower signal in
693 a male in the control condition compared to a male exposed to a mirror. NC, negative control. ** p < 0.01
694
695