Tumor-infiltrating nerves functionally alter brain circuits and modulate behavior in a mouse model of head-and-neck cancer

Jeffrey Barr [1†], Austin Walz [1†], Anthony C. Restaino[1,3], Moran Amit[4], Sarah M. Barclay[1],

Elisabeth G. Vichaya[2], William C. Spanos[1,3], Robert Dantzer[4], Sebastien Talbot[5], Paola D.

Vermeer[1,3,*]

[1]Sanford Research, Cancer Biology and Immunotherapies Group, Sioux Falls, South Dakota,

USA.

[2]Baylor University, Department of Psychology and Neuroscience, Waco, TX, USA.

[3]University of South Dakota, Sanford School of Medicine, Vermillion, SD, USA.

[4]University of Texas, MD Anderson Cancer Center, Houston, TX, USA.

[5]Queen's University, Department of Biomedical and Molecular Sciences, Kingston, Ontario,

Canada.

[†] equal first authors

*Corresponding author   Email: Paola.Vermeer@sanfordhealth.org

Conflict of interest: RD is a consultant for GoodCap, Toronto; the consultancy is not related to

the present work.

Abstract: Cancer patients often experience changes in mental health, prompting an exploration into whether nerves infiltrating tumors contribute to these alterations by impacting brain functions. Using a mouse model for head and neck cancer and neuronal tracing we show that tumor-infiltrating nerves connect to distinct brain areas. The activation of this neuronal circuitry altered behaviors (decreased nest-building, increased latency to eat a cookie, and reduced wheel running). Tumor-infiltrating nociceptor neurons exhibited heightened calcium activity and brain regions receiving these neural projections showed elevated Fos as well as increased calcium responses compared to non-tumor-bearing counterparts.

The genetic elimination of nociceptor neurons decreased brain Fos expression and mitigated the behavioral alterations induced by the presence of the tumor. While analgesic treatment restored nesting and cookie test behaviors, it did not fully restore voluntary wheel running indicating that pain is not the exclusive driver of such behavioral shifts. Unraveling the interaction between the tumor, infiltrating nerves, and the brain is pivotal to developing targeted interventions to alleviate the mental health burdens associated with cancer.

Significance Statement: Head and neck cancers are infiltrated by sensory nerves which connect to a pre-existing circuit that includes areas in the brain. Neurons within this circuit are altered and mediate modifications in behavior.

Introduction:

The prevalence of mental health disorders (e.g., depression, anxiety, suicide) in cancer patients is significantly greater than in the general population [1-3]. For those patients with no prior psychiatric history, a cancer diagnosis increases the risk of mental health decline [4]. The presence of cancer introduces many stressors (physical, financial, relational) into the lives of patients, thus a negative impact on mental health may not be surprising. However, these changes persist even in long-term cancer survivors. For instance, decade-long cancer survivors maintain an increased incidence of depression (approximately 12%, depending on the cancer type) as compared to the population at large (3-5%) [5-7]. While the intensity and prevalence of psychological symptoms in cancer patients fluctuate before, during, and after treatment for a given type of cancer, and between cancer types, it remains higher than in the general population [8-10]. The association of cancer with impaired mental health is directly mediated by the disease, its treatment or both; these findings suggest that the development of a tumor alters brain functions.

We have demonstrated the presence of TRPV1-expressing nociceptor neurons in head and neck squamous cell carcinomas (HNSCC) [11], melanoma, cervical [12], and ovarian cancers [13]. Nerve recruitment to the tumor bed is an active process that involves the release of soluble factors, including neurotrophins [14-16] and neuropeptides [17]. Tumor-released small extracellular vesicles (sEVs) also recruit loco-regional nerves to the tumor bed [11,18]. While these and other studies establish that solid peripheral tumors engage with the peripheral nervous system [11,13,19-22], it raises the possibility of a direct neuronal connection from the tumor to the brain.

A recent study used pseudorabies virus and mapped a connection from tumor-infiltrating nerves in an orthotopic model of murine lung cancer to areas in the brain [23]. We expand these findings and demonstrate that HNSCC-associated nerves are transcriptionally and functionally altered and project to discrete regions in the brain. The brain neurons connected to the tumor manifest

increased activity, which is associated with behavioral alterations in tumor-bearing animals.

Consistent with this, newly diagnosed HNSCC patients suffer high rates of depression and

anxiety and lower quality of life [24]. As most cancer patients face cancer-related pain [25-27], we

tested whether treating pain could restore normal behavior. While pain treatment restored normal

function at the tumor site, such as nesting behavior, it only partially restored normal voluntary

running wheel behavior. Our findings suggest that, in addition to pain, tumor-infiltrating nerves

communicate signals to the brain that lead to cancer-associated changes in behavior.

Results

Tumor innervation begins early in disease. MOC2-7 cells are HNSCC cancer cells derived

from a CXCR3 null mouse on a C57BL/6 background [28]. Their implantation in male mice results

in dense innervation of tumors with nociceptor nerves[29,30]. To define the timing of tumor

innervation, C57BL/6 wildtype male mice were orthotopically implanted (oral cavity) with

MOC2-7 cells, and tumors were collected on days 4-, 10-, and 20- post-implantation. Western

blot analysis of whole tumor lysate indicated expression of Tau, a neuronal marker [31], as early as

day four post-tumor implantation, which increased over time (**Fig. 1A**; full westerns in

Supplemental Fig. 1A-B; densitometric quantification of westerns, **Supplemental Fig. 1E**). A

similar increase in the neuronal marker doublecortin (DCX) was noted (**Fig. 1B**; full western

Supplemental Fig. 1C-D; densitometric quantification of westerns, **Supplemental Fig. 1F**).

While DCX is well-known for its expression in immature neurons, it is also expressed in adult

peripheral neurons, including dorsal root ganglia [32], and in non-neuronal tissues[33]. The DCX

signal did not originate from the tumors, as MOC2-7 cell lysate was negative for this protein

(**Fig. 1B**, Full western **Supplemental Fig 1C**).   Consistent with this, immunohistochemical

staining of these tumors with the neuronal marker β-III tubulin (β3T), demonstrated the

increasing presence of nerves beginning on day four post-tumor implantation (**Fig. 1C-E**).

Tumor-infiltrating nerves map to the ipsilateral trigeminal ganglion and into the CNS. To

map the origin of tumor-infiltrating nerves and define the circuits they converge upon, mice with

palpable oral MOC2-7 tumors (approximately day 15 post-tumor implantation) were intra-

tumorally injected with wheat germ agglutinin, WGA, a neural tracer conjugated to a

fluorophore (n= 10 mice). WGA is a lectin molecule that specifically binds to sialic acid residues

present ubiquitously on neuronal membranes. It has been used extensively to map neuronal

circuits centrally and in the periphery [34,35] and is a known transganglionic and transsynaptic

neuronal tracer [36-38] making it ideal for mapping neural circuits [39-43]. Following tracer injection,

tumor growth was permitted for an additional 3-7 days to allow time for tracer labeling to occur.

Animals were then euthanized, and tumors, trigeminal (TGM) ganglia, and brains were harvested

and analyzed by microscopy. Microscopic examination of tumors revealed nerves with robust

WGA signals (**Fig 1F**) as well as the V3 branch of the ipsilateral, but not the contralateral, TGM

ganglion (**Fig 1G,H**). Consistent with the restricted labeling of tumor-infiltrating nerves, the

tracer did not diffuse outside the tumor mass (**Fig. 1I**).

Examination of brains from MOC2-7 tumor-bearing animals revealed tracer[+] neurons in specific

regions including spinal nucleus of the trigeminal (SpVc), parabrachial nucleus (PBN), and

central amygdala (CeA) (**Fig 1J**). Sections of brain from these regions show the presence of

tracer-positive neurons (**Fig 1K**). Tracer injections of equivalent volume and concentration into

the oral cavities of control non-tumor-bearing animals did not label the TGM ganglia nor areas in

the brain (data not shown). A larger volume (10 μl) of tracer injected into non-tumor bearing

mice resulted in tracer labeling of ipsilateral TGM neurons and brain (**Fig. 1L**). These control

studies indicate that the nerve density and distribution present in the tumor bed are higher than in

control mice, with the consequence that a small volume (2 µl instead of 10 µl) of tracer is

sufficient to result in nerve labeling. The mapped circuit encompasses pre-existing connections

to brain areas that regulate pain and affect [44-47]. These data indicate that the nerve infiltration into

the tumor extends this circuit.

Tumor-infiltrating neurons have altered transcript levels. The observation of nerves

infiltrating the tumor mass prompted us to ask whether they undergo alterations by their mere

presence within this "foreign" environment. Consistent with this, we previously found that

melanoma-infiltrating neurons have a unique transcriptome [48]. To test whether this is also the

case in HNSCC, we orthotopically implanted MOC2-7 cells into the oral cavity of male wildtype

mice. After 14 days, TGM neurons were harvested and analyzed by qPCR using a commercial

array for neural transmission and membrane trafficking genes. Compared to control TGM (non-

tumoral) neurons, ipsilateral TGM neurons from tumor-bearing animals harbored increased

expression (>4-fold) in genes involved in neuronal signaling/receptors (*Gabrg1, Gabra4,*

Grin2c, Grm3) and synaptic transmission (*Gria2*) (**Fig. 2A; Supplemental Table 1**) [49]. The gene

demonstrating the highest increase in expression, *Fus*, was of particular interest; it increases in

expression within DRG neurons following nerve injury and contributes to injury-induced pain

[50,51]. Of note, we purposefully used whole trigeminal ganglia rather than FACS-sorted tracer-

positive dissociated neurons to avoid artificially imposing injury and altering the transcript levels

of these cells [52,53]. Thus, significantly elevated expression of *Fus* by ipsilateral TGM neurons

from tumor-bearing animals suggests the presence of neuronal injury induced by the malignancy.

This is consistent with our previous findings [54] and those of others [55] showing that tumor-

infiltrating nerves harbor higher expression of nerve-injury transcripts and neuronal sensitization.

Tumor-infiltrating neurons are functionally changed. Given this transcriptomic alteration, we

tested whether tumor-infiltrating neurons exhibit functional alterations. Thus, MOC2-7 tumor-

bearing mice were intra-tumorally injected with the fluorophore-conjugated WGA tracer. Five

days later, tracer[+] TGM ganglia were cultured, and responsiveness to noxious stimuli was

analyzed by calcium microscopy. Neuronal Ca^{+2} responses to capsaicin (300 nM), which binds

and activates TRPV1 channels, were measured by Fluo-4AM fluorescence microscopy. We

found that tracer[+] tumor-infiltrating ipsilateral TGM neurons show increased responsiveness to

capsaicin compared to tracer negative contralateral neurons (amplitude, **Fig. 2B-D;** the area

under the curve**, Fig. 2E**). Similar to contralateral TGM neurons, non-tumor-bearing TGM

neurons elicited a normal response to capsaicin (**Fig. 2F**).

This heightened sensitivity could reflect increased TRPV1 expression and/or its phosphorylation.

We tested whether this was the case using western blotting and discovered an overexpression of

the sigma 1 receptor (σ1R) and increased TRPV1 phosphorylation in the TGM ganglia from

tumor-bearing animals (**Fig. 2G-I,** full westerns **Supplemental Fig 2A-C**). Of note, the σ1R is

an endoplasmic reticulum chaperone protein that directly interacts with TRPV1 and regulates its

membrane expression, [56] while protein kinase C phosphorylation of serine 502 and 800 of

TRPV1 reduces the receptor's activation threshold [57-60]. Taken together, our data indicate that

HNSCC-infiltrating nerves have a unique transcriptome and a heightened sensitivity to noxious

stimuli characterized by an increased TRPV1 expression and phosphorylation consistent with

TRPV1 sensitization secondary to oral cancer [61].

Tumor-brain circuit neurons harbor elevated activity. The transcriptional and functional

changes evident in tracer[+] tumor-infiltrating neurons could result in alterations in central target

neurons. To assess this, brain sections from MOC2-7 tumor-bearing and non-tumor-bearing

animals were immunofluorescently stained for cFos and ΔFosB, two markers of neuronal activity

with different courses of expression [62-64]. ΔFosB expression was increased in several brain

regions of tumor-bearing animals, while cFos expression was also increased in the PBN (**Fig.**

3A-F). Predominant differences in the long-lived ΔFosB were expected [65], as those with the

short-lived cFos changes are more challenging to capture using single time point assessment.

Next, we assessed the neuronal activity of tumor-bearing animal brains using stereotaxically

injected AAV1-Syn-GCaMP6f, a viral vector encoding a neuron-specific synapsin-driven

calcium sensor [66], into the PBN. Two weeks after intra-cranial injection of the virus, mice were

orally implanted with MOC2-7 cells. Approximately 18-days post-tumor inoculation, the animals

were euthanized, and the neuronal calcium activity was recorded in *ex vivo* brain slices using a

mini scope (**Supplemental Fig. 3A, B**). While baseline fluorescence was similar between tumor-

bearing and control animals, that recorded upon stimulation (KCl; 30 mM) was significantly

higher in neurons of tumor-bearing animals (**Fig. 3G-J; Supplemental Fig 3A, B**). These

functional and Fos staining data indicate that central neurons within the tumor-brain circuit are

functionally altered compared to their healthy brain counterparts.

Ablation of tumor-infiltrating neurons attenuates cancer-induced brain alterations. Cancer

patients, and even more notably, survivors, experienced poor mental health [1,67-69]. The neural

connection between tumor and brain, together with our finding that TRPV1-expressing

nociceptor neurons within this circuit become functionally altered, might contribute to these

changes. To test whether this is the case, we genetically engineered mice with ablated nociceptor

neurons (TRPV1-Cre::Floxed-DTA). TRPV1-Cre::Floxed-DTA animals lack TRPV1-expressing

cells (including neurons) as well as many other nociceptive neurons. These nociceptor-neuron-

ablated mice have been previously characterized and lack the expected sensitivity to temperature

as well as itch and pain reactions to chemical mediators such as capsaicin [70]. Fos immunostaining

of brains from nociceptor ablated and control (C57BL/6) mice show no significant differences

(**Supplemental Fig 3C-E**) indicating that, in the absence of a malignancy, the neurons in these

regions are not differentially activated. First, we confirmed the absence of TRPV1$^+$ neurons in

the TGM of these ablated animals (**Fig. 4A**). We then analyzed Fos expression in the brain of

MOC2-7-bearing nociceptor intact and ablated mice. While Fos expression was similar between

non-tumor bearing mice of the two genotypes (**Supplemental Fig. 3C-E**), the absence of

nociceptor neurons in tumor-bearing animals decreases cFos and ΔFosB in the PBN, and ΔFosB

in the SpVc (**Fig. 4B, C**). Other tested regions were not impacted (**Fig. 4D**). These data suggest

that the tumor-brain communication was disrupted in the absence of nociceptor neurons.

Alleviation of pain does not always restore behavior. To evaluate the effects of this disruption

on cancer-induced behavioral changes, we assessed the animals' general well-being through

nesting behavior [71] and anhedonia using the cookie test [72,73], as well as body weight and food

disappearance as surrogates for oral pain and/or loss of appetite. Nociceptor-neuron-ablated mice

showed increased nesting performance (**Fig. 4E**) and decreased anhedonia (**Fig. 4F**) compared to

intact mice. This was accompanied by smaller tumor growth (**Fig. 4G**) and increased survival

(**Fig. 4H**). Given the impact of nociceptor neuron ablation on tumor growth, we wondered

whether differences in tumor volume contributed to the behavioral differences we noted. Thus,

the behavior data were graphed as a function of tumor volume (**Supplemental Fig 4A, B**). A

simple linear regression model was used to fit the data. In the case of nesting scores, the linear

regression did not fit the data points very well making it difficult to assess nesting scores at a

given tumor volume (**Supplemental Fig 4A**). However, the linear regression model fit the time

to interact data better. Here, the data suggest that tumor volume did not influence behavior as at

any given tumor volume the time to interact with the cookie is generally smaller in TRPV1-

Cre::Floxed-DTA animals as compared to C57BL/6 animals (**Supplemental Fig 4B**). While both

groups showed similar body weight loss (**Fig. 4I**), nociceptor ablated animals show a transient

(Wk 1, 4, 5) decrease in food disappearance (**Fig. 4J**). Significant effects for all statistical

analyses in Figure 4 are presented in Supplementary Table 2.

The developmental ablation of TRPV1 neurons can lead to unintended effects on tumor

innervation and behavior. Thus, we chemoablated TRPV1 neurons in 4-week-old juvenile

C57BL/6 mice using resiniferatoxin (RTX) [74,75]. We examined nesting behavior in RTX and

vehicle-treated C57BL/6 male mice orthotopically implanted with MOC2-7 cells. An additional

control group consisted of age-matched C57BL/6 mice without tumors. Although all groups

displayed similar nesting scores initially, the scores in vehicle-treated mice declined significantly

faster compared to RTX-treated mice (**Fig 5A**). Body weight decreased in all tumor-bearing

mice, but the extent of weight loss was similar between vehicle and RTX-treated mice (**Fig 5B**).

Similar to the developmental ablation of TRPV1 neurons, their chemoablation post-development

results in a similar decrease in tumor growth compared to controls (**Fig 5C**). Since HNSCC

tumors cause oral pain and both nesting and the cookie test require the use of the mouth, we also

evaluated the effect of nociceptor ablation on voluntary wheel running to gain additional insights

into cancer-associated fatigue, a surrogate for depressive-like behaviors in tumor-bearing mice [76-

80]. To assess the potential influence of cancer-associated pain, 1-week post tumor inoculation,

groups of mice were treated with carprofen (a non-steroidal anti-inflammatory), extended-release

buprenorphine (opioid), or vehicle. We found that wheel running decreased as tumors

progressed, an effect partially alleviated by carprofen and buprenorphine (**Fig 6A, B**). The

nesting behavior (**Fig 6C**), body weight (**Fig 6D**), and food disappearance (**Fig 6E**) were also

alleviated by the painkillers. Neither carprofen nor buprenorphine impacted MOC2-7 tumor

growth compared to vehicle-treated animals (**Fig 6F**), but buprenorphine increased tumor growth

compared to carprofen-treated mice (significant effects for all statistical analyses in Figure 5 are

presented in Supplementary Table 3). These findings suggest that choosing pain management

drugs in the context of cancer can potentially influence tumor growth adversely. Furthermore,

the data reveal that behaviors associated with the tumor site are adversely affected by cancer-

associated pain. Pain-induced anhedonia is mediated by changes in the reward pathway.

Specifically, in the context of pain, dopaminergic neurons in the ventral tegmental area (VTA)

become less responsive to pain and release less serotonin. This decreased serotonin results in

disinhibition of GABA release; the resulting increased GABA promotes an increased inhibitory

drive leading to anhedonia [81] and, when extreme, anorexia. Carprofen and buprenorphine

treatments completely reversed nesting behavior and significantly improved eating.

Inflammation [82] and opioids [83] directly influence reward processing and though our tracing

studies did not indicate that the tumor-brain circuit includes the VTA, this brain region may be

indirectly impacted by tumor-induced pain in the oral cavity. Thus, an alternative interpretation

of the data is that the effects of carprofen and buprenorphine treatments on nesting and food

consumption may be due to inhibition of anhedonia (and anorexia) rather than, or in addition to,

relieving oral pain. However, merely alleviating pain (or inhibiting anhedonia) does not fully

restore all behaviors, as evidenced by persistent issues in activities like wheel running.

Discussion

We have previously demonstrated that head and neck squamous cell carcinoma is associated

with innervation by sensory neurons and that substance P, one of the principal neuropeptides

these fibers release, drives malignant cell proliferation and migration [29]. Significantly, patients

suffering from HNSCC often experience depressive disorders, which are poorly managed

clinically [84-86]. Here, we found that nociceptor neurons infiltrating HNSCC connect to a pre-

existing brain-projecting circuit [87,88] that includes the trigeminal ganglia, the spinal nucleus of the

trigeminal (SpVc), as well as the parabrachial nucleus (PBN), and central amygdala (CeA)[89-92].

This connection occurs independently of pain, and this cancer-brain circuit drives behavioral

alterations in tumor-bearing mice. While the elimination of TRPV1-expressing nociceptor

neurons reduces tumor growth and enhances survival rates, it also highlights a potential

therapeutic target for mitigating depressive behaviors in cancer patients. This underscores the

complex interplay between sensory neurons, cancer progression, and mental health, uncovering a

two-pronged approach to improving HNSCC patients' physical and psychological health.

In melanoma [48] and HNSCC, we demonstrated significant alterations in transcript levels within

tumor-infiltrating nerves. Several mechanisms can account for these changes. For instance,

TRPV1 stimulation is sufficient to trigger the activation of the AP-1 transcription factor within

neurons [93]. Such TRPV1 activation could occur in response to low pH [94] or hypoxia [95-97], which

are both prevalent within the tumor microenvironment. Alternatively, we previously

demonstrated that tumor-released sEVs, which transport microRNAs (miRNAs), induced

sprouting of loco-regional nerves to the tumor bed [11,98]. These sEV-transported miRNAs could

also modulate various neuronal transcription factors and, in turn, drive the transcriptomic

changes we observed in these neurons.

Our data indicate that these peripheral neuronal changes also influence the functioning of the

brain areas to which they connect. Although we did not make the complete cartography of all

brain regions used by tumor-infiltrating nerves, we identified a few critical nuclei. Subsequent

work will use emerging circuit mapping techniques for whole brain profiling [99]. Nevertheless,

our tracing in tumor-bearing nociceptor-neuron-ablated and intact animals was sufficient to

functionally implicate TRPV1-expressing neurons in this communication.

Trigeminal ganglia neurons are composed of ~80% of $Na_V1.8^+$ nociceptor neurons, with one-half

being peptidergic ($TRPV1^+$, $TRPA1^+$) neurons [100,101,70]. In the TRPV1-Cre::Floxed-DTA mouse,

these peptidergic neurons are eliminated, leaving intact ~50% of pain-sensing neurons. The

remaining presence of these non-peptidergic, largely $MrgD^+$, neurons might explain why nesting

behavior in TRPV1-Cre::Floxed-DTA tumor-bearing animals is only partially restored.

Inflammation contributes to pain in cancer [102]. Consistent with this, treatment of tumor-bearing

mice with analgesic drugs (carprofen, buprenorphine) resulted in complete restoration of nest

building and performance in the cookie test. These data show that the constraint imposed by the

tumor on behavioral activities that require the use of the oral cavity is mainly due to pain and not

to the physical interference of the tumor mass with the pattern of oral activities that need to take

place for mice to perform the corresponding behavior.

Cancer pain often interferes with activities that don't involve oral movements, a phenomenon

evident in the impaired voluntary wheel running observed in tumor-bearing mice. Although

analgesic drugs can partially mitigate this deficit, they fail to restore this behavior fully, despite

their efficacy in completely restoring oral activity behaviors like nesting and the cookie test.

Moreover, given that carprofen and buprenorphine decrease inflammation [103], their ability to

restore normal nesting and cookie test behaviors (which require the use of the oral cavity where

the tumor is located) suggests that inflammation at the tumor site contributed to the decline in

these behaviors in vehicle-treated animals. Since both drugs were given systemically and each

only partially restored wheel running, it suggests that systemic inflammation alone cannot fully

account for the decline in wheel running seen in vehicle-treated animals. We posit that the

inflammation- and pain-independent component of this behavioral decline is mediated via the

transcriptional and functional alterations in the cancer-brain circuit.

This suggests that factors beyond pain contribute to the observed behavioral changes. These

alterations could be attributed to depression [104] or represent a competition between the energy

demands of the HNSCC malignant cells and the host's skeletal muscles [105-107].

Neuro-immune interactions have been studied in the context of a variety of conditions including,

but not limited to infection [108], inflammation [109,110], homeostasis in the gut [111-113], as well as

neurological diseases[114,115]. Neuro-immune communications in the context of cancer and

behavior have also been studied (e.g., sickness behavior, depression) [116-118] however, these

studies did not assess these interactions at the tumor bed. Investigations into neuro-immune

interactions occurring within primary malignancies which harbor nerves have shed light on these

critical communications. In the context of melanoma, which is innervated by sensory nerves, we

identified that release of the neuropeptide calcitonin gene related peptide (CGRP) induces

immune suppression. This effect is mediated by CGRP binding to its receptor, RAMP1, which is

expressed on CD8+ T cells [48]. A study utilizing a different syngeneic model of oral cancer

similarly found an immune suppressive role for CGRP [119-121]. These studies demonstrate that

neuro-immune interactions occur at the tumor bed. Our current findings indicating that tumor-

infiltrating nerves connect to a circuit that includes regions within the brain suggest that neuro-

immune interactions within the peripheral malignancy may contribute to the behavioral

alterations we studied.

Our current findings suggest that interrupting tumor-to-brain communication can mitigate the

mental health decline often associated with cancer. Yet, a higher prevalence of depression

persists among long-term cancer survivors compared to individuals without a cancer history.

Having received treatment and showing no evidence of disease, these survivors continue to battle

mental health issues [6,7]. This phenomenon prompts a critical question - why does this

discrepancy exist? It is conceivable that these patients, through treatments like surgery, radiation,

or chemotherapy, have disrupted the tumor-brain connection and have sustained this

disconnection over the years. However, unlike animal models, their mental health decline is not

reversed. Several speculations, grounded in known factors, can be considered.

The human experience with cancer is distinct from that of mouse models; individuals can live

with malignant growths for extended periods, often unaware of their existence due to

asymptomatic or nonspecific symptoms. During this undiagnosed period, the tumor-brain circuit

might induce irreversible changes in central neurons through activity-dependent transcriptional

modifications [122,123]. These alterations, anchored in neuronal plasticity, adaptation, and behavior,

may become permanent even after the cancer is diagnosed and treated.

Moving forward, to reverse these entrenched brain changes, understanding the specific

transcriptional alterations incurred by central neurons is vital. Unraveling this complexity could

pave the way for therapies that reverse the neuronal activity impacts, fostering improved mental

health. Although in its early stages, our insight into the role of nerves in cancer is growing, and

highlighting neuronal targets for pharmacological interventions.

Our study is not exempt from limitations. First, our exploration of the tumor-to-brain pathway

was confined to mice with HNSCC and used a single cancer cell line. Hence, the universality of

our conclusions warrants verification through additional HNSCC cell lines and diverse cancer

models. Furthermore, the exclusive focus on HNSCC, particularly considering the nerve-rich

oral region, raises questions about the pathway's establishment and behavioral impact in other

contexts. Our behavioral assessment was not exhaustive; a comprehensive analysis of nerve-

dependent and independent behaviors remains forthcoming. Behaviors could also be influenced

by elements like soluble factors or the energy competition between the tumor and behavioral

activities.

Head and neck cancer is predominantly a cancer in males; it occurs in males three times more

often than in females [124], this disparity increases in certain parts of the world. While smoking

cigarettes and drinking alcohol are risk factors for HPV negative head and neck squamous cell

carcinoma, even males that do not smoke and drink are have a higher susceptibility for this

cancer than females [125,126]. Thus, our studies used only male mice. However, we do recognize

that females also get this cancer. In fact, female patients with head and neck cancer, particularly

oral cancer, report more pain than their male counterparts [127,128]. These findings suggest that

differences in tumor innervation exist in males and females. Our studies have also solely

involved male mice, presenting a clear gap in understanding the potential sex differences in the

development of the tumor-to-brain communication pathway and its subsequent influence on

behavior. The inclusion of female subjects in future research is essential to provide a

comprehensive insight into these processes.

Second, our nerve tracing methodology, where we injected 2 µl of WGA into the tumor bed, is

designed to label only the tumor-infiltrating nerves and their connections. However, this

approach fails to label *all* tumor-infiltrating nerves, predominantly those not close to the WGA-

injected region and nerves that are in contact with or influenced by the tumor but are not directly

infiltrating it. This limitation is a calculated one, reflecting a technical trade-off. We aimed to

ensure the specificity of capturing only tumor-infiltrating nerves, which meant sacrificing the

comprehensiveness of labeling all nerves associated with the tumor. This constraint extends to

our calcium imaging studies and brain tracing. Although we can confidently assert that all

labeled neurons are indeed tumor-infiltrating or connected, we cannot conclusively state that all

tracer-negative ipsilateral nerves are unconnected to the tumor-brain circuit.

Even with these constraints, our findings are important. We have demonstrated that tumor-

infiltrating nerves are integrated into a pre-established neuronal circuit. This circuit extends from

the tumor bed to the TGM ganglion, connects to the SpVc, and projects into the brain,

influencing behavior in both pain-dependent and independent ways. This revelation paves the

way for in-depth exploration into the mechanistic underpinnings of cancer-associated alterations

in well-being and mental health, shedding light on potential therapeutic interventions to alleviate

these profound effects.

MATERIALS AND METHODS

Study approval. All animal studies were performed with approval from the Institutional Animal Care and Use Committee at Sanford Research and were within institutional guidelines and complied with all relevant ethical regulations. Sanford Research has an Animal Welfare Assurance on file with the Office of Laboratory Animal Welfare (assurance number: A-4568-01) and is accredited by AAALAC, Intl. Sanford Health is also a licensed research facility under the authority of the United States Department of Agriculture (USDA) with USDA certificate number 46-R-011.

Inclusion & Ethics Statement. All studies utilizing animals were performed with approval from the appropriate ethical bodies to ensure sufficient protection was in place. This study was a collaborative, multi-disciplinary, multi-institutional effort with contributions from researchers in academic positions.

Cell lines. MOC2-7 cells (RRID:CVCL_ZD34) [28] (previously known as MOC7) [129] were a kind gift from Dr. Ravindra Uppaluri (Dana-Farber Cancer Institute, Boston, MA). They were maintained in DMEM with 10% fetal calf serum and cultured at 37°C and 5% CO_2; with culture medium refreshed every three days. MOC2-7 tumors have been characterized as non-inflamed and poorly immunogenic [130-132]. These cells can be obtained through Kerafest (www.kerafest.com).

Animal studies. All animal experiments were performed in the Sanford Research Animal

Resource Center which is a specific pathogen free facility. All mice were maintained in IVC

Tecniplast Green line Seal Safe Plus cages which were opened only under aseptic conditions in

an animal transfer station. All cages were changed every other week using aseptic technique. All

cages had individual HEPA filtered air. Animal rooms were maintained at $75°$F, 30-70%

humidity, with a minimum of 15 air changes per hour, and a 14:10 light/dark cycle. Corncob

bedding, which was autoclaved prior to use, was maintained in all cages. Irradiated, sterile food

(Envigo) and acidified water (pH 2.8-3.0) were available *ad libitum*. There was a maximum of 5

mice/cage. All animals were observed daily for abnormal behavior, signs of illness or distress,

the availability of food and water and proper husbandry. Animals injected with murine tumor

cells were 10-week-old C57BL/6 or TRPV1-Cre::Floxed-DTA mice (The Jackson Laboratory)

weighing approximately 24 g. Investigators were blinded to the groups when assessing animals

(e.g., measuring tumors). Animals were numbered by ear punch and cage number.

Nociceptor-neuron-ablated TRPV1-Cre::Floxed-DTA animals were generated by crossing

ROSA26-DTA (diphtheria toxin A, B6.129P2-*Gt(ROSA)26Sor* [tm1(DTA)Lky]/J)(Jax#009669) and

TRPV1-Cre (B6.129-*Trpv1*[tm1(cre)Bbm] /J) [133](Jax# 017769) mice; the progeny (TRPV1-

Cre::Floxed-DTA) express DTA (diphtheria toxin fragment A) under control of the TRPV1

promoter, thereby genetically ablating all TRPV1 expressing cells (including TRPV1-expressing

neurons) throughout development. Absence of TRPV1-expressing sensory neurons was validated

by IHC staining of TGM ganglia for TRPV1.

Chemoablation of TRPV1-expressing neurons post-development was accomplished with

resiniferatoxin (RTX), a naturally occurring potent TRPV1 agonist. RTX treatment results in the

specific ablation of TRPV1 neurons (post-development), while sparing other TRPV1 expressing

(non-neuronal) cells. RTX treatment in mice consisted of three injections of RTX given 24 hours

apart. The injections are given in the flank and were: 30 µg/kg on day 1, 70 µg/kg on day 2 and

100 µg/kg on day 3. Just prior to the first RTX injection, mice are treated with 0.1 mg/kg of

buprenorphine, given intraperitoneally, as the first injection causes pain. Subsequent injections

do not require buprenorphine. It takes 4 weeks for TRPV1 nerve ablation to occur. Control mice

received injections with vehicle solution (DMSO with Tween80 in PBS). The tail flick test was

used to confirm nerve ablation.

Tail flick test. Prior to testing, mice are acclimated to the room for 30 minutes. A water bath is

pre-warmed to 52°$\square$C and a thermometer is used to constantly monitor the temperature of the

water. The mice are picked up using a fresh paper towel so that only their tail is outside the paper

towel. Once the mouse is relaxed in the tester's hand, the mouse's tail is lowered into the water

so that a quarter of the tail is submerged. The time for the tail to be flicked out of the water is

recorded. If no response has occurred by 20 seconds, the mouse's tail is removed from the water

to prevent tissue damage. A response time latency greater than 10 seconds signifies denervation.

Tumor implantation. The cells injected into animals were cultured in media containing 10%

fetal calf serum. When cells are harvested for tumor injections, they are first washed two times

with PBS and then trypsinized to detach the cells from the plate. Cells are collected, washed

again with PBS and resuspended with DMEM without serum; this is what is injected into

animals. We harvest cells in this way in order to eliminate any serum being injected into mice.

Ketamine (87.5 mg/kg)/xylazine (10 mg/kg) were used to anesthetize mice prior to oral cavity

tumor implantation. Tumors were initiated as follows: using a 25-gauge needle, cells (5×10^4

cells in a total of 50 µl) were implanted orthotopically in the oral cavity of male C57BL/6 mice.

Specifically, the needle was placed proximal to the crease of the mouse cheek, inserted at

approximately a 45° angle, bevel deep, and tumor cells injected into this location of the cheek

pouch. Mice were allowed to recover from anesthesia on a heating blanket and returned to their

home cage. Control (non-tumor) mice were treated the same way only their injection contained

media alone (no cells). Tumor growth was monitored weekly by caliper measurements of

isoflurane anesthetized animals and tumor volume calculated was using the formula: ½

(length*(width^2)). Non-tumor control mice were also anesthetized with isoflurane each time

that tumor-bearing animals were anesthetized for tumor measurements.

Euthanasia criteria: Criteria for euthanasia in our IACUC approved protocol include maximum

tumor volume of 1000mm^3, edema, extended period of weight loss progressing to emaciation,

impaired mobility or lesions interfering with eating, drinking or ambulation, rapid weight loss

(>20% in 1 week), as well as weight loss at or more than 20% of baseline. In addition to tumor

size and weight loss, we use the body condition score to evaluate the state of animals and to

determine euthanasia.

Drug treatments. Carprofen (Pivetal) was provided in the drinking water (10 mg/kg which is

equivalent to 0.067 mg/ml) which was refreshed every 7 days. Buprenorphine-ER (Fidelis

Animal Health), a 72-hour extended-release formulation, was given by subcutaneous injection

(3.25 mg/kg) every 72 hours. Vehicle-treated animals were treated with subcutaneous injection

of saline. Drug treatments commenced on day 7 post-tumor implantation in all groups. Given

that buprenorphine treatment requires a subcutaneous injection every 3 days, mice in the other

groups (vehicle and carprofen) received a subcutaneous injection of saline at the same volume

(60 µl) as the buprenorphine-treated animals. In this way, all animal stress from handling and

injection was the same.

Nest building. Nest building is an innate behavior performed in rodents of both sexes and is a

general indication of well-being [71]; it also measures the motivation to perform a goal-directed

task. To assess nesting behavior, individually housed mice were given a nesting square (Ancare)

overnight. The following morning, nests were scored by two independent scorers who were

blinded to the conditions [134,135]. The scoring system is 0-4 and based on the percentage of the

nestlet that is shredded and the height of the nest. Two baseline measurements were conducted

prior to tumor implantation followed by weekly testing. Exclusion criteria of an average baseline

score of 3 or greater was used to ensure all mice were good nest builders at baseline (n=3

C57BL/6 mice and n=7 TRPV1-Cre::Floxed-DTA mice were excluded for this reason).

Cookie test: The cookie test is a variation of the sucrose preference test, a test of anhedonia (the

inability to experience enjoyment or interest in previously rewarding activities). These test the

animal's endogenous inclination for sweet tastes and the notion that it derives pleasure from

consuming sweets. One hour prior to testing, mouse food is removed and the animals are

acclimated to the brightly lit testing room. Following the acclimation period, mice were given a

piece of cookie (approximately 1.5 g). The time it took the mouse to bite the cookie from its

placement in the cage was measured to determine time-to-interact. Each mouse was acclimated

to the cookie every other day for 2 weeks prior to tumor implantation with a baseline

measurement followed by weekly testing. Mice were individually housed. One mouse from the

C57BL/6 group was excluded due to incomplete data (mouse died before completion of the

study).

Voluntary wheel running: Prior to tumor implantation, mice were singly housed with a running

wheel maintained in the home cage continuously. The only time that the wheel was removed was

when mice were undergoing nesting. The mice were acclimated to the wheel for 2 weeks prior to

tumor implantation. This acclimatization period provided the time to stabilize their running

performance. Running wheels were maintained in the home cage for the duration of the study.

As mice are most active during the dark phase, nighttime running (8pm to 8am) was assessed.

Running data were collected in 1-minute bins continuously throughout the duration of the

experiment.

Food disappearance. Animals were provided 100 g of solid food per cage each week. At the

end of each week, the remaining food was weighed, and this value was used to calculate the

amount of food that disappeared in that week per cage. For all behavioral experiments, mice

were singly housed.

Neural tracing of orthotopic HNSCC tumors. When oral tumors reached approximately 5 x 5

mm in size, neural tracer was injected into the tumor as described below. Ketamine (87.5

mg/kg)/xylazine (10 mg/kg) were used to anesthetize the mice. A 10 µL Hamilton syringe with

30-G needle was loaded with 1% Wheat Germ Agglutinin (WGA) conjugated to either

AlexaFluor 594 (Invitrogen) or CF568 (Biotium) in PBS and 2 µl was slowly injected intra-

tumorally with the bevel side up. The needle was inserted approximately midway through the

tumor then pulled back slightly to reduce pressure and leakage of tracer following injection. The

injection of tracer occurred slowly over the course of 10 minutes. After the tracer was injected,

the needle was kept in place for 2 additional minutes before being slowly removed. Mice were

placed on a heating pad until recovery from anesthesia. Five to seven days later, mice were

deeply anesthetized and transcardially perfused with ice cold PBS followed by 4%

paraformaldehyde and trigeminal ganglia (TGM) were carefully removed and placed in HBSS in

a 24-well plate kept on ice. TGM harvesting was completed as follows: euthanized animals were

subjected to a midline incision while in the prone position to expose the crown of the skull. The

brainstem was separated from the spinal cord by a transverse cut and the top of the skull was

removed, exposing the brainstem and TGM. All tissues (tumor, ganglia and brain) were fixed,

sectioned, and imaged for WGA labeled nerve fibers and somas under confocal microscopy.

WGA injection into control (non-tumor bearing animals) was performed the same way and

injection was into the same oral region where tumors were implanted. Volumes of 2 or 10 µl of

WGA were used.

Immunohistochemical (IHC) staining. Tissues were formalin fixed, paraffin-embedded and cut

into 5-µm sections. The BenchMark® XT automated slide staining system (Ventana Medical

Systems, Inc.) was used to optimize antibody dilutions and staining. The Ventana CC1 solution

was used to perform the antigen retrieval step (basic pH tris base buffer). Tissues were incubated

in primary antibody for 1 hour. The Ventana iView DAB detection kit was used as the

chromogen and the slides were counterstained with hematoxylin.

Antibody used for immunohistochemistry (IHC). Anti-β-III Tubulin (Abcam, Cat# ab78078,

1:250, RRID: AB 2256751), Anti-TRPV1 (Alomone Labs Cat# ACC-030, 1:400,

RRID:AB_2313819).

Antibodies used for immunofluorescence (IF). cFos (Cell Signaling, Cat# 2250, 1:10,000,

RRID: AB 2247211), ΔFosB (Abcam, Ab11959, 1:5,000, RRID:AB_298732). The following

secondary antibodies were used: Alexa568 anti-rabbit (Thermo, Cat# A-11011, 1:500,

RRID:AB_143157), Alexa488 anti-mouse (Thermo, Cat# A-11001, 1:500, RRID:AB_2534069).

Antibodies used for western blot. β-actin (Sigma Life Science, Cat#A2228, 1:1000, RRID:

AB_476697), Tau (Abcam, Cat# ab75714, 1:500, RRID:AB_1310734), phosphorylated TRPV1

(Thermo Fisher Scientific, Cat# PA5-64860, 1:500, RRID:AB_2663797), Sigma-1R

(ProteinTech, Cat# 15168-1-AP, 1:1,000, RRID:AB_2301712), Doublecortin (DCX) (Santa

cruz, Cat# sc-271390, 1:1000, RRID:AB_10610966).

Brain immunostaining and analysis. Brains were post-fixed for twenty-four hours. Coronal or

sagittal sections were cut at 40 µm using a vibratome. Sequential fluorescent immunolabeling

was performed for cFos or ΔFosB on free-floating sections. All sections were blocked with 10%

goat serum in 0.3% Tx-100. All antibodies were diluted in PBS containing 0.03% Triton X-100

and 2% normal goat serum. Sections were incubated in Fos antibody for 72 hours. Following

washes in 0.1 M phosphate buffered saline, sections were incubated in secondary antibody for 3

hours. Following additional washes, sections were mounted onto glass slides, and cover slipped

with cytoseal prior to imaging. The total numbers of Fos-labeled nuclei were counted on $10 \times 2D$

images acquired using a laser-scanning confocal microscope (Nikon A1R) and verified as

positive if the signal filled the nucleus and stood out clearly compared to surrounding tissue [136].

Nuclei were counterstained with DAPI. Quantification of Fos-labeled cells was performed using

digital thresholding of Fos-immunoreactive nuclei. The threshold for detection was set at a level

where dark Fos-immunoreactive nuclei were counted, but nuclei with light labeling, similar to

background staining, were not. For each animal, 3-4 sections were selected at 120 μm intervals.

ImageJ cell counter (v. 1.52) was used to quantify Fos+ nuclei per region / section.

Analysis of transcriptional changes in tumor-infiltrating neurons. To assess transcriptional

changes in tumor-infiltrating neurons we utilize a qPCR array (ScienCell, GeneQuery Neural

Transmission and Membrane Trafficking, #MGK008). These qPCR-ready 96 well plates enable

rapid profiling of 88 key genes important for neuronal functions (listed in Table S1). Mice

bearing MOC2-7 oral tumors were injected with tracer (as described above); TGM ganglia were

isolated on day 28 post-tumor implantation, their RNA harvested, converted into cDNA and then

assayed with the array as per manufacturer's instructions (ScienCell, GeneQuery arrays). We

used N=4 TGM/group from n=4 mice/group; n=4 plates/condition. Control RNA was isolated

from TGM ganglia from age-matched non-tumor bearing mice (n=4 mice). Relative gene

expression was calculated as per manufacturer's recommendations.

Western blot analysis of whole tumor lysate. Tumors were excised from euthanized mice,

taking care to eliminate as much non-tumor tissue as possible. Tumors were then placed in

approximately 500μl lysis buffer (50mM Tris HCl pH 7.4, 100mM NaCl, 100mM NaF, 10mM

NaPPi, 2mM Na_3VO_4, 10% glycerol, HALT protease inhibitor cocktail) with 1% TX-100 and

kept on ice for 10 minutes. The homogenate was sonicated 3x for 15 secs and then incubated on

ice for 15 minutes. Samples were then centrifuged at 2000 g for 5 minutes at 4°C, supernatant

was collected and further centrifuged at 12,000 g for 10 minutes at 4°C. Protein concentrations

of the supernatants were determined by BCA protein assay.

Western blot analysis of ganglia lysate. Trigeminal ganglia were harvested as described above.

Ganglia were lysed using a homogenizer in lysis buffer at 4°C. Homogenates were centrifuged at

10,000 g for 20 minutes and protein concentrations determined from the supernatant. Protein

concentration was measured by BCA protein assay (Pierce, Cat#23225) and 40 µg of total

protein were separated by SDS-PAGE, transferred to PDVF membranes which were then

blocked for 30 minutes at room temperature (RT) in 5% milk in PBS. Membranes were

incubated with primary antibody overnight at 4°C. Following 1xTBST washes, membranes were

probed with an HRP-conjugated secondary antibody (1:10,000 dilution) for 1 hour at RT,

washed and imaged on a Li-COR Odyssey FC imaging system. Densitometric quantification was

used to assess changes in protein expression.

Western blot densitometry analysis. Raw images of western blots were analyzed using ImageJ.

Briefly, images were opened and bands of interested were selected by gating with the rectangle

selection tool. Densitometry was measured based on grey scale analysis of selected area.

Analysis was conducted on proteins of interest as well as the loading control band (β-actin).

Relative expression of proteins of interest was determined as a fraction of β-actin densitometry.

Neuron culture. Trigeminal ganglia were harvested from MOC2-7 tumor-bearing or non-tumor

animals and enzyme digested in papain, and then collagenase II/dispase at 37 □ for 15 min. After

washing and trituration, cells were plated onto a thin layer of Matrigel® in glass bottom dishes

and cultured with HamsF12 supplemented with 10% FBS. The cells were maintained in an

incubator (5% CO_2, 37 °C) for 24 h before they were used for calcium imaging experiments.

Ca2+ imaging. TGM neurons from non-tumor and tumor-bearing animals (n=4-6

mice/condition) were imaged on the same day. Neurons were incubated with the calcium

indicator, Fluo-4AM, at 37°C for 20 min. After dye loading, the cells were washed, and Live

Cell Imaging Solution (Thermo-Fisher) with 20 mM glucose was added. Calcium imaging was

conducted at room temperature. Changes in intracellular Ca^{2+} were measured using a Nikon

scanning confocal microscope with a□10x objective. Fluo-4AM was excited at 488 nm using an

argon laser with intensity attenuated to 1%. The fluorescence images were acquired in the

confocal frame (1024□×□1024 pixels) scan mode. After 1 min of baseline measure, capsaicin

(300nM final concentration) was added. Ca^{2+} images were recorded before, during and after

capsaicin application. Image acquisition and analysis were achieved using NIS-Elements

imaging software. Fluo-4AM responses were standardized and shown as percent change from the

initial frame. Data are presented as the relative change in fluorescence ($\Delta F/F_0$), where F_0 is the

basal fluorescence and $\Delta F = F - F_0$ with F being the measured intensity recorded during the

experiment. Calcium responses were analyzed only for neurons responding to ionomycin (10

µM, positive control) to ensure neuronal health. Treatment with the cell permeable Ca^{2+} chelator,

BAPTA (200 µM), served as a negative control.

Stereotaxic AAV injection. Male mice were anesthetized with 2% isoflurane and placed in a

stereotaxic head frame on a heating pad. A midline incision was made down the scalp and a

craniotomy was made using a micro drill. A 10-μl Hamilton syringe was used to infuse 1 μl of

AAV1/Syn-GCaMP6f-WPRESV40 (titer 4.65×10^{13} GC per ml, via Addgene) into the

parabrachial nucleus (−5.3 mm anteroposterior, -1.3 mm mediolateral, −3 mm dorsoventral) via

a microsyringe pump. After infusion, the needle was kept at the injection site for 5 min and then

slowly withdrawn. Two weeks following stereotaxic surgeries, tumors were introduced by

implanting MOC2-7 cells orthotopically into the oral cavity as described above.

Ex vivo **Ca2+ imaging of brain slices.** Two weeks post-tumor (or sham) implantation, wildtype

and tumor-bearing mice (n=3 mice/group) were anesthetized under isoflurane and perfused

intracardially with 10 ml of ice-cold N-methyl-d-glucamine (NMDG) solution [92 mM NMDG,

30 mM NaHCO$_3$, 25 mM glucose, 20 mM Hepes, 10 mM MgSO$_4$, 5 mM sodium ascorbate, 3

mM sodium pyruvate, 2.5 mM KCl, 2 mM thiourea, 1.25 mM NaH$_2$PO$_4$, and 0.5 mM CaCl$_2$ (pH

7.3, 300 mOsm, bubbled with 95% O2 and 5% CO$_2$)] [137]. The brains were quickly removed and

placed into additional ice-cold NMDG solution for slicing. Coronal slices (150 μm) were cut

using a vibratome (n=3 slices/brain). Slices were transferred to Hepes holding solution and

warmed to 37°C (bubbled with 95% O$_2$ and 5% CO$_2$) for 1 hour. After incubation, slices were

transferred to the recording chamber with RT (22° to 25°C) recording solution. A miniaturized

microscope (Miniscope V4) imaged the GCaMP6f signal from the slices. Each video was

processed with motion correction and ΔF/F calculation. Regions of interest (ROIs; considered as

a single cell soma) were manually selected. ROIs that exhibited short bursts of ΔF/F changes or

fluctuations during the recording were analyzed. The ΔF/F changes were then aligned with the

time window of treatment. Movement correction was performed using the motion correction

module in the EZcalcium toolbox employed in MATLAB [138]. The fluorescence intensity trace of

each neuron was extracted, and ΔF/F was calculated (CNMF).

Statistical analysis. GraphPad Prism (version 10.0.3, 2023) was used for all statistical analyses.

Gene expression: For qRT-PCR analysis, Ct values for each gene were normalized to that of the

reference gene. Statistical analysis by multiple student's t-test.

Ca^{2+} imaging: Statistical analysis by unpaired student's t-test. Fluo-4AM/GCaMP6f responses

were standardized and shown as percent change from the initial frame. Data are presented as the

relative change in fluorescence ($\Delta F/F_0$), where F_0 is the basal fluorescence and $\Delta F = F - F_0$ with F

being the peak response.

Western blot: Membranes were visualized, and proteins were quantified using the Odyssey

infrared imaging system and software (Li-COR). Densitometric quantification of western blots

was performed by normalizing the signal from tumor or non-tumor ganglia to the β-actin

(loading control) signal and differences assessed by one-way ANOVA with post-hoc Tukey test.

Fos brain immunostaining: The numbers of Fos (cFos, ΔFosB) immune-positive neurons in brain

sections from tumor-bearing or non-tumor animals were quantified as described above and

statistically analyzed by 2-way ANOVA with multiple comparisons.

Nesting scores: The variations over time of the nesting scores for each group were statistically

analyzed by repeated measures ANOVA. Exclusion criteria of an average baseline score of 3 or

greater was used to ensure all mice were good nest builders.

Time to interact scores (cookie test): Statistical analysis by repeated measures ANOVA.

Voluntary wheel running: The data were collected in 1-minute bins continuously. When

compiling the data, the sum of bins between 8pm to 8am is taken for each individual animal for

voluntary nightly running. Baseline is calculated by taking the final 3 days prior to tumor

implantation and averaging the values. The percent change is then calculated for each day from

baseline for each mouse. The data were then analyzed using a repeated measures 2-way

ANOVA.

Food disappearance: Statistical analysis by repeated measures ANOVA.

Weight: Statistical analysis by repeated measures ANOVA.

Tumor growth curves: Statistical analysis of tumor growth curves by repeated measures ANOVA

Kaplan Meier survival: Statistical analysis of survival by Log-rank (Mantel-Cox) test.

700
701
702
703

References

1   Carreira, H. *et al.* Associations Between Breast Cancer Survivorship and Adverse Mental
Health Outcomes: A Systematic Review. *Journal of the National Cancer Institute* **110**,
1311-1327 (2018). https://doi.org:10.1093/jnci/djy177
2   Pitman, A., Suleman, S., Hyde, N. & Hodgkiss, A. Depression and anxiety in patients
with cancer. *Bmj* **361**, k1415 (2018). https://doi.org:10.1136/bmj.k1415
3   Walker, J. *et al.* Prevalence of depression in adults with cancer: a systematic review.
Annals of oncology : official journal of the European Society for Medical Oncology **24**,
895-900 (2013). https://doi.org:10.1093/annonc/mds575
4   Zhu, J. *et al.* First-onset mental disorders after cancer diagnosis and cancer-specific
mortality: a nationwide cohort study. *Annals of oncology : official journal of the*
European Society for Medical Oncology **28**, 1964-1969 (2017).
https://doi.org:10.1093/annonc/mdx265

5   Steel, Z. *et al.* The global prevalence of common mental disorders: a systematic review
and meta-analysis 1980-2013. *Int J Epidemiol* **43**, 476-493 (2014).
https://doi.org:10.1093/ije/dyu038
6   Gotze, H. *et al.* Depression and anxiety in long-term survivors 5 and 10 years after cancer
diagnosis. *Supportive care in cancer : official journal of the Multinational Association of*
Supportive Care in Cancer **28**, 211-220 (2020). https://doi.org:10.1007/s00520-019-
04805-1
7   Kuba, K. *et al.* Risk for depression and anxiety in long-term survivors of hematologic
cancer. *Health Psychol* **38**, 187-195 (2019). https://doi.org:10.1037/hea0000713
8   Wang, Y. H. *et al.* Depression and anxiety in relation to cancer incidence and mortality: a
systematic review and meta-analysis of cohort studies. *Mol Psychiatry* **25**, 1487-1499
(2020). https://doi.org:10.1038/s41380-019-0595-x
9   Naser, A. Y. *et al.* Depression and Anxiety in Patients With Cancer: A Cross-Sectional
Study. *Front Psychol* **12**, 585534 (2021). https://doi.org:10.3389/fpsyg.2021.585534
10   Tsaras, K. *et al.* Assessment of Depression and Anxiety in Breast Cancer Patients:
Prevalence and Associated Factors. *Asian Pacific journal of cancer prevention : APJCP*
19, 1661-1669 (2018). https://doi.org:10.22034/APJCP.2018.19.6.1661
11   Madeo, M. *et al.* Cancer exosomes induce tumor innervation. *Nature communications* **9**,
4284 (2018). https://doi.org:10.1038/s41467-018-06640-0
12   Lucido, C. T. *et al.* Innervation of cervical carcinoma is mediated by cancer-derived
exosomes. *Gynecologic oncology* (2019). https://doi.org:10.1016/j.ygyno.2019.04.651
13   Barr, J. L. *et al.* Intra-Tumoral Nerve-Tracing in a Novel Syngeneic Model of High-
Grade Serous Ovarian Carcinoma. *Cells* **10** (2021).
https://doi.org:10.3390/cells10123491
14   Renz, B. W. *et al.* beta2 Adrenergic-Neurotrophin Feedforward Loop Promotes
Pancreatic Cancer. *Cancer cell* **33**, 75-90 e77 (2018).
https://doi.org:10.1016/j.ccell.2017.11.007
15   Hayakawa, Y. *et al.* Nerve Growth Factor Promotes Gastric Tumorigenesis through
Aberrant Cholinergic Signaling. *Cancer cell* **31**, 21-34 (2017).
https://doi.org:10.1016/j.ccell.2016.11.005
16   Wang, K. *et al.* The neurotrophic factor neurturin contributes toward an aggressive
cancer cell phenotype, neuropathic pain and neuronal plasticity in pancreatic cancer.
Carcinogenesis **35**, 103-113 (2014). https://doi.org:10.1093/carcin/bgt312
17   Kasprzak, A. & Adamek, A. The Neuropeptide System and Colorectal Cancer Liver
Metastases: Mechanisms and Management. *International journal of molecular sciences*
21 (2020). https://doi.org:10.3390/ijms21103494
18   Silverman, D. A. *et al.* Cancer-Associated Neurogenesis and Nerve-Cancer Cross-talk.
Cancer research **81**, 1431-1440 (2021). https://doi.org:10.1158/0008-5472.CAN-20-2793
19   Cole, S. W., Nagaraja, A. S., Lutgendorf, S. K., Green, P. A. & Sood, A. K. Sympathetic
nervous system regulation of the tumour microenvironment. *Nat Rev Cancer* **15**, 563-572
(2015). https://doi.org:10.1038/nrc3978
20   Magnon, C. *et al.* Autonomic nerve development contributes to prostate cancer
progression. *Science* **341**, 1236361 (2013). https://doi.org:10.1126/science.1236361
21   Sadighparvar, S., Darband, S. G., Ghaderi-Pakdel, F., Mihanfar, A. & Majidinia, M.
Parasympathetic, but not sympathetic denervation, suppressed colorectal cancer

progression. *Eur J Pharmacol* **913**, 174626 (2021).
https://doi.org:10.1016/j.ejphar.2021.174626

22   Chen, D. & Ayala, G. E. Innervating Prostate Cancer. *N Engl J Med* **378**, 675-677
(2018). https://doi.org:10.1056/NEJMcibr1714003

23   Chen, M. *et al.* The Rostral Ventromedial and Lateral Medulla Are the Major Areas
Responsive to Lung Cancer Progression among Brainstem Lung-Innervating Nuclei.
Brain Sci **12** (2022). https://doi.org:10.3390/brainsci12111486

24   Henry, M. *et al.* Psychosocial outcomes of human papillomavirus (HPV)- and non-HPV-
related head and neck cancers: A longitudinal study. *Psychooncology* **31**, 185-197 (2022).
https://doi.org:10.1002/pon.5803

25   Virgen, C. G. *et al.* Pharmacological management of cancer pain: Novel therapeutics.
Biomed Pharmacother **156**, 113871 (2022). https://doi.org:10.1016/j.biopha.2022.113871

26   Hjermstad, M. J., Fainsinger, R., Kaasa, S. & European Palliative Care Research, C.
Assessment and classification of cancer pain. *Curr Opin Support Palliat Care* **3**, 24-30
(2009). https://doi.org:10.1097/SPC.0b013e3283260644

27   van den Beuken-van Everdingen, M. H. *et al.* Prevalence of pain in patients with cancer:
a systematic review of the past 40 years. *Annals of oncology : official journal of the
European Society for Medical Oncology* **18**, 1437-1449 (2007).
https://doi.org:10.1093/annonc/mdm056

28   Judd, N. P. *et al.* ERK1/2 regulation of CD44 modulates oral cancer aggressiveness.
Cancer research **72**, 365-374 (2012). https://doi.org:10.1158/0008-5472.CAN-11-1831

29   Restaino, A. C. *et al.* Functional neuronal circuits promote disease progression in cancer.
Sci Adv **9**, eade4443 (2023). https://doi.org:10.1126/sciadv.ade4443

30   Menezes, F. D. S., Fernandes, G. A., Antunes, J. L. F., Villa, L. L. & Toporcov, T. N.
Global incidence trends in head and neck cancer for HPV-related and -unrelated subsites:
A systematic review of population-based studies. *Oral oncology* **115**, 105177 (2021).
https://doi.org:10.1016/j.oraloncology.2020.105177

31   Georgieff, I. S. *et al.* Expression of high molecular weight tau in the central and
peripheral nervous systems. *Journal of cell science* **105 (Pt 3)**, 729-737 (1993).
https://doi.org:10.1242/jcs.105.3.729

32   Dellarole, A. & Grilli, M. Adult dorsal root ganglia sensory neurons express the early
neuronal fate marker doublecortin. *J Comp Neurol* **511**, 318-328 (2008).
https://doi.org:10.1002/cne.21845

33   Bernreuther, C., Salein, N., Matschke, J. & Hagel, C. Expression of doublecortin in
tumours of the central and peripheral nervous system and in human non-neuronal tissues.
Acta Neuropathol **111**, 247-254 (2006). https://doi.org:10.1007/s00401-006-0038-z

34   Borges, L. F. & Sidman, R. L. Axonal transport of lectins in the peripheral nervous
system. *The Journal of neuroscience : the official journal of the Society for Neuroscience*
2, 647-653 (1982).

35   van der Want, J. J., Klooster, J., Cardozo, B. N., de Weerd, H. & Liem, R. S. Tract-
tracing in the nervous system of vertebrates using horseradish peroxidase and its
conjugates: tracers, chromogens and stabilization for light and electron microscopy.
Brain Res Brain Res Protoc **1**, 269-279 (1997). https://doi.org:10.1016/s1385-
299x(96)00042-6

36 McNicholas, K. & Michael, M. Z. Immuno-characterization of Exosomes Using
Nanoparticle Tracking Analysis. *Methods in molecular biology* **1545**, 35-42 (2017).
https://doi.org:10.1007/978-1-4939-6728-5_3
37 Dumas, M., Schwab, M. E. & Thoenen, H. Retrograde axonal transport of specific
macromolecules as a tool for characterizing nerve terminal membranes. *Journal of*
neurobiology **10**, 179-197 (1979). https://doi.org:10.1002/neu.480100207
38 Sillitoe, R. V. Mossy Fibers Terminate Directly Within Purkinje Cell Zones During
Mouse Development. *Cerebellum* **15**, 14-17 (2016). https://doi.org:10.1007/s12311-015-
0712-6
39 Tabuchi, K. *et al.* GAL4/UAS-WGA system as a powerful tool for tracing Drosophila
transsynaptic neural pathways. *Journal of neuroscience research* **59**, 94-99 (2000).
40 Levy, S. L., White, J. J., Lackey, E. P., Schwartz, L. & Sillitoe, R. V. WGA-Alexa
Conjugates for Axonal Tracing. *Current protocols in neuroscience* **79**, 1 28 21-21 28 24
(2017). https://doi.org:10.1002/cpns.28
41 Itaya, S. K. Anterograde transsynaptic transport of WGA-HRP in rat olfactory pathways.
Brain research **409**, 205-214 (1987). https://doi.org:10.1016/0006-8993(87)90703-7
42 Itaya, S. K., Van Hoesen, G. W. & Barnes, C. L. Anterograde transsynaptic transport of
WGA-HRP in the limbic system of rat and monkey. *Brain research* **398**, 397-402 (1986).
https://doi.org:10.1016/0006-8993(86)91504-0
43 Carson, K. A. & Mesulam, M. M. Ultrastructural evidence in mice that
transganglionically transported horseradish peroxidase-wheat germ agglutinin conjugate
reaches the intraspinal terminations of sensory neurons. *Neuroscience letters* **29**, 201-206
(1982). https://doi.org:10.1016/0304-3940(82)90317-2
44 Yeh, L. F., Watanabe, M., Sulkes-Cuevas, J. & Johansen, J. P. Dysregulation of aversive
signaling pathways: a novel circuit endophenotype for pain and anxiety disorders. *Curr*
Opin Neurobiol **48**, 37-44 (2018). https://doi.org:10.1016/j.conb.2017.09.006
45 Hu, H. Reward and Aversion. *Annual review of neuroscience* **39**, 297-324 (2016).
https://doi.org:10.1146/annurev-neuro-070815-014106
46 Zhu, Y. B. *et al.* PBN-PVT projections modulate negative affective states in mice. *eLife*
11 (2022). https://doi.org:10.7554/eLife.68372
47 Russo, S. J. & Nestler, E. J. The brain reward circuitry in mood disorders. *Nature*
reviews. Neuroscience **14**, 609-625 (2013). https://doi.org:10.1038/nrn3381
48 Balood, M. *et al.* Nociceptor neurons affect cancer immunosurveillance. *Nature* (2022).
https://doi.org:10.1038/s41586-022-05374-w
49 Michetti, F. *et al.* The S100B story: from biomarker to active factor in neural injury.
Journal of neurochemistry **148**, 168-187 (2019). https://doi.org:10.1111/jnc.14574
50 Han, G. *et al.* FUS Contributes to Nerve Injury-Induced Nociceptive Hypersensitivity by
Activating NF-kappaB Pathway in Primary Sensory Neurons. *J Neurosci* **43**, 1267-1278
(2023). https://doi.org:10.1523/JNEUROSCI.2082-22.2022
51 Du, S. *et al.* A nerve injury-specific long noncoding RNA promotes neuropathic pain by
increasing Ccl2 expression. *J Clin Invest* **132** (2022). https://doi.org:10.1172/JCI153563
52 Liu, Q. *et al.* Transcriptional Alterations of Mouse Trigeminal Ganglion Neurons
Following Orofacial Inflammation Revealed by Single-Cell Analysis. *Frontiers in*
cellular neuroscience **16**, 885569 (2022). https://doi.org:10.3389/fncel.2022.885569

53 Malin, S. A., Davis, B. M. & Molliver, D. C. Production of dissociated sensory neuron
cultures and considerations for their use in studying neuronal function and plasticity.
Nature protocols **2**, 152-160 (2007). https://doi.org:10.1038/nprot.2006.461
54 Inyang, K. E. *et al.* HPV+ head and neck cancer-derived small extracellular vesicles
communicate with TRPV1+ neurons to mediate cancer pain. *Pain* (2023).
https://doi.org:10.1097/j.pain.0000000000003045
55 Horan, N. L. *et al.* The impact of tumor immunogenicity on cancer pain phenotype using
syngeneic oral cancer mouse models. *Front Pain Res (Lausanne)* **3**, 991725 (2022).
https://doi.org:10.3389/fpain.2022.991725
56 Ortiz-Renteria, M. *et al.* TRPV1 channels and the progesterone receptor Sig-1R interact
to regulate pain. *Proceedings of the National Academy of Sciences of the United States of
America* **115**, E1657-E1666 (2018). https://doi.org:10.1073/pnas.1715972115
57 Bhave, G. *et al.* Protein kinase C phosphorylation sensitizes but does not activate the
capsaicin receptor transient receptor potential vanilloid 1 (TRPV1). *Proceedings of the
National Academy of Sciences of the United States of America* **100**, 12480-12485 (2003).
https://doi.org:10.1073/pnas.2032100100
58 Numazaki, M. *et al.* Structural determinant of TRPV1 desensitization interacts with
calmodulin. *Proceedings of the National Academy of Sciences of the United States of
America* **100**, 8002-8006 (2003). https://doi.org:10.1073/pnas.1337252100
59 Numazaki, M., Tominaga, T., Toyooka, H. & Tominaga, M. Direct phosphorylation of
capsaicin receptor VR1 by protein kinase Cepsilon and identification of two target serine
residues. *The Journal of biological chemistry* **277**, 13375-13378 (2002).
https://doi.org:10.1074/jbc.C200104200
60 Bhave, G. *et al.* cAMP-dependent protein kinase regulates desensitization of the
capsaicin receptor (VR1) by direct phosphorylation. *Neuron* **35**, 721-731 (2002).
https://doi.org:10.1016/s0896-6273(02)00802-4
61 Scheff, N. N. *et al.* Oral cancer induced TRPV1 sensitization is mediated by PAR(2)
signaling in primary afferent neurons innervating the cancer microenvironment. *Scientific
reports* **12**, 4121 (2022). https://doi.org:10.1038/s41598-022-08005-6
62 Chung, L. A Brief Introduction to the Transduction of Neural Activity into Fos Signal.
Dev Reprod **19**, 61-67 (2015). https://doi.org:10.12717/DR.2015.19.2.061
63 Hudson, A. E. Genetic Reporters of Neuronal Activity: c-Fos and G-CaMP6. *Methods in
enzymology* **603**, 197-220 (2018). https://doi.org:10.1016/bs.mie.2018.01.023
64 Nestler, E. J., Barrot, M. & Self, D. W. DeltaFosB: a sustained molecular switch for
addiction. *Proceedings of the National Academy of Sciences of the United States of
America* **98**, 11042-11046 (2001). https://doi.org:10.1073/pnas.191352698
65 Nestler, E. J., Kelz, M. B. & Chen, J. DeltaFosB: a molecular mediator of long-term
neural and behavioral plasticity. *Brain research* **835**, 10-17 (1999).
https://doi.org:10.1016/s0006-8993(98)01191-3
66 Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity.
Nature **499**, 295-300 (2013). https://doi.org:10.1038/nature12354
67 Niedzwiedz, C. L., Knifton, L., Robb, K. A., Katikireddi, S. V. & Smith, D. J. Depression
and anxiety among people living with and beyond cancer: a growing clinical and research
priority. *BMC cancer* **19**, 943 (2019). https://doi.org:10.1186/s12885-019-6181-4
68 Utley, M., Adeyanju, T., Bernardo, B., Paskett, E. D. & Krok-Schoen, J. L. The
association between mental health, social support and physical health outcomes among

older female cancer survivors. *J Geriatr Oncol* **13**, 834-838 (2022).
https://doi.org:10.1016/j.jgo.2022.04.001
69 Ji, X. *et al.* National estimates of mental health service use and unmet needs among adult
cancer survivors. *Cancer* **127**, 4296-4305 (2021). https://doi.org:10.1002/cncr.33827
70 Mishra, S. K., Tisel, S. M., Orestes, P., Bhangoo, S. K. & Hoon, M. A. TRPV1-lineage
neurons are required for thermal sensation. *The EMBO journal* **30**, 582-593 (2011).
https://doi.org:10.1038/emboj.2010.325
71 Neely, C. L. C., Pedemonte, K. A., Boggs, K. N. & Flinn, J. M. Nest Building Behavior
as an Early Indicator of Behavioral Deficits in Mice. *Journal of visualized experiments :*
JoVE (2019). https://doi.org:10.3791/60139
72 Liu, M. Y. *et al.* Sucrose preference test for measurement of stress-induced anhedonia in
mice. *Nature protocols* **13**, 1686-1698 (2018). https://doi.org:10.1038/s41596-018-0011-
z
73 Eliwa, H. *et al.* Adult neurogenesis augmentation attenuates anhedonia and HPA axis
dysregulation in a mouse model of chronic stress and depression.
Psychoneuroendocrinology **124**, 105097 (2021).
https://doi.org:10.1016/j.psyneuen.2020.105097
74 Liu, J. *et al.* TRPV1(+) sensory nerves modulate corneal inflammation after epithelial
abrasion via RAMP1 and SSTR5 signaling. *Mucosal immunology* **15**, 867-881 (2022).
https://doi.org:10.1038/s41385-022-00533-8
75 Mishra, S. K. & Hoon, M. A. Ablation of TrpV1 neurons reveals their selective role in
thermal pain sensation. *Mol Cell Neurosci* **43**, 157-163 (2010).
https://doi.org:10.1016/j.mcn.2009.10.006
76 Norden, D. M. *et al.* Tumor growth increases neuroinflammation, fatigue and depressive-
like behavior prior to alterations in muscle function. *Brain, behavior, and immunity* **43**,
76-85 (2015). https://doi.org:10.1016/j.bbi.2014.07.013
77 Zombeck, J. A., Fey, E. G., Lyng, G. D. & Sonis, S. T. A clinically translatable mouse
model for chemotherapy-related fatigue. *Comp Med* **63**, 491-497 (2013).
78 Grossberg, A. J. *et al.* Tumor-Associated Fatigue in Cancer Patients Develops
Independently of IL1 Signaling. *Cancer research* **78**, 695-705 (2018).
https://doi.org:10.1158/0008-5472.CAN-17-2168
79 Bower, J. E. *et al.* Inflammation and behavioral symptoms after breast cancer treatment:
do fatigue, depression, and sleep disturbance share a common underlying mechanism?
Journal of clinical oncology : official journal of the American Society of Clinical
Oncology **29**, 3517-3522 (2011). https://doi.org:10.1200/JCO.2011.36.1154
80 Norden, D. M. *et al.* Fluoxetine prevents the development of depressive-like behavior in
a mouse model of cancer related fatigue. *Physiol Behav* **140**, 230-235 (2015).
https://doi.org:10.1016/j.physbeh.2014.12.045
81 Markovic, T. *et al.* Pain induces adaptations in ventral tegmental area dopamine neurons
to drive anhedonia-like behavior. *Nat Neurosci* **24**, 1601-1613 (2021).
https://doi.org:10.1038/s41593-021-00924-3
82 Bekhbat, M., Treadway, M. T. & Felger, J. C. Inflammation as a Pathophysiologic
Pathway to Anhedonia: Mechanisms and Therapeutic Implications. *Curr Top Behav*
Neurosci **58**, 397-419 (2022). https://doi.org:10.1007/7854_2021_294

Le Merrer, J., Becker, J. A., Befort, K. & Kieffer, B. L. Reward processing by the opioid system in the brain. *Physiological reviews* **89**, 1379-1412 (2009). https://doi.org:10.1152/physrev.00005.2009

Chen, A. M. *et al.* Depression among long-term survivors of head and neck cancer treated with radiation therapy. *JAMA Otolaryngol Head Neck Surg* **139**, 885-889 (2013). https://doi.org:10.1001/jamaoto.2013.4072

Lazure, K. E., Lydiatt, W. M., Denman, D. & Burke, W. J. Association between depression and survival or disease recurrence in patients with head and neck cancer enrolled in a depression prevention trial. *Head & neck* **31**, 888-892 (2009). https://doi.org:10.1002/hed.21046

Elaldi, R. *et al.* Correlations between long-term quality of life and patient needs and concerns following head and neck cancer treatment and the impact of psychological distress. A multicentric cross-sectional study. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies* **278**, 2437-2445 (2021). https://doi.org:10.1007/s00405-020-06326-8

Navratilova, E. *et al.* Positive emotions and brain reward circuits in chronic pain. *J Comp Neurol* **524**, 1646-1652 (2016). https://doi.org:10.1002/cne.23968

Mercer Lindsay, N., Chen, C., Gilam, G., Mackey, S. & Scherrer, G. Brain circuits for pain and its treatment. *Science translational medicine* **13**, eabj7360 (2021). https://doi.org:10.1126/scitranslmed.abj7360

Ito, M. *et al.* The parabrachial-to-amygdala pathway provides aversive information to induce avoidance behavior in mice. *Mol Brain* **14**, 94 (2021). https://doi.org:10.1186/s13041-021-00807-5

Jaramillo, A. A., Brown, J. A. & Winder, D. G. Danger and distress: Parabrachial-extended amygdala circuits. *Neuropharmacology* **198**, 108757 (2021). https://doi.org:10.1016/j.neuropharm.2021.108757

Ge, J., Cai, Y. & Pan, Z. Z. Synaptic plasticity in two cell types of central amygdala for regulation of emotion and pain. *Front Cell Neurosci* **16**, 997360 (2022). https://doi.org:10.3389/fncel.2022.997360

Torres-Rodriguez, J. M. *et al.* The parabrachial to central amygdala pathway is critical to injury-induced pain sensitization in mice. *Neuropsychopharmacology* **49**, 508-520 (2024). https://doi.org:10.1038/s41386-023-01673-6

Backes, T. M. *et al.* Stimulation of TRPV1 channels activates the AP-1 transcription factor. *Biochemical pharmacology* **150**, 160-169 (2018). https://doi.org:10.1016/j.bcp.2018.02.008

Boedtkjer, E. & Pedersen, S. F. The Acidic Tumor Microenvironment as a Driver of Cancer. *Annual review of physiology* (2019). https://doi.org:10.1146/annurev-physiol-021119-034627

Pocock, R. & Hobert, O. Oxygen levels affect axon guidance and neuronal migration in Caenorhabditis elegans. *Nature neuroscience* **11**, 894-900 (2008). https://doi.org:10.1038/nn.2152

Pocock, R. & Hobert, O. Hypoxia activates a latent circuit for processing gustatory information in C. elegans. *Nature neuroscience* **13**, 610-614 (2010). https://doi.org:10.1038/nn.2537

97   Stevenson, T. J. *et al.* Hypoxia disruption of vertebrate CNS pathfinding through
ephrinB2 Is rescued by magnesium. *PLoS Genet* **8**, e1002638 (2012).
https://doi.org:10.1371/journal.pgen.1002638
98   Amit, M. *et al.* Loss of p53 drives neuron reprogramming in head and neck cancer.
Nature **578**, 449-454 (2020). https://doi.org:10.1038/s41586-020-1996-3
99   Ueda, H. R. *et al.* Whole-Brain Profiling of Cells and Circuits in Mammals by Tissue
Clearing and Light-Sheet Microscopy. *Neuron* **106**, 369-387 (2020).
https://doi.org:10.1016/j.neuron.2020.03.004
100   Kobayashi, K. *et al.* Distinct expression of TRPM8, TRPA1, and TRPV1 mRNAs in rat
primary afferent neurons with adelta/c-fibers and colocalization with trk receptors. *J*
Comp Neurol **493**, 596-606 (2005). https://doi.org:10.1002/cne.20794
101   Mickle, A. D., Shepherd, A. J. & Mohapatra, D. P. Sensory TRP channels: the key
transducers of nociception and pain. *Prog Mol Biol Transl Sci* **131**, 73-118 (2015).
https://doi.org:10.1016/bs.pmbts.2015.01.002
102   Renna, M. E. *et al.* Depression and anxiety in colorectal cancer patients: Ties to pain,
fatigue, and inflammation. *Psychooncology* **31**, 1536-1544 (2022).
https://doi.org:10.1002/pon.5986
103   Shepherd, A. J. & Mohapatra, D. P. Pharmacological validation of voluntary gait and
mechanical sensitivity assays associated with inflammatory and neuropathic pain in mice.
Neuropharmacology **130**, 18-29 (2018).
https://doi.org:10.1016/j.neuropharm.2017.11.036
104   Walker, A. K., Kavelaars, A., Heijnen, C. J. & Dantzer, R. Neuroinflammation and
comorbidity of pain and depression. *Pharmacological reviews* **66**, 80-101 (2014).
https://doi.org:10.1124/pr.113.008144
105   Grossberg, A. J. *et al.* Interleukin 6-independent metabolic reprogramming as a driver of
cancer-related fatigue. *Brain, behavior, and immunity* **88**, 230-241 (2020).
https://doi.org:10.1016/j.bbi.2020.05.043
106   Thaker, P. H. *et al.* Chronic stress promotes tumor growth and angiogenesis in a mouse
model of ovarian carcinoma. *Nature medicine* **12**, 939-944 (2006).
https://doi.org:10.1038/nm1447
107   Kim-Fuchs, C. *et al.* Chronic stress accelerates pancreatic cancer growth and invasion: a
critical role for beta-adrenergic signaling in the pancreatic microenvironment. *Brain,*
behavior, and immunity **40**, 40-47 (2014). https://doi.org:10.1016/j.bbi.2014.02.019
108   Camp, B., Stegemann-Koniszewski, S. & Schreiber, J. Infection-Associated Mechanisms
of Neuro-Inflammation and Neuro-Immune Crosstalk in Chronic Respiratory Diseases.
Int J Mol Sci **22** (2021). https://doi.org:10.3390/ijms22115699
109   Kabata, H. & Artis, D. Neuro-immune crosstalk and allergic inflammation. *J Clin Invest*
129, 1475-1482 (2019). https://doi.org:10.1172/JCI124609
110   Chen, O., Donnelly, C. R. & Ji, R. R. Regulation of pain by neuro-immune interactions
between macrophages and nociceptor sensory neurons. *Curr Opin Neurobiol* **62**, 17-25
(2020). https://doi.org:10.1016/j.conb.2019.11.006
111   van Baarle, L., Stakenborg, M. & Matteoli, G. Enteric neuro-immune interactions in
intestinal health and disease. *Semin Immunol* **70**, 101819 (2023).
https://doi.org:10.1016/j.smim.2023.101819

Jacobson, A., Yang, D., Vella, M. & Chiu, I. M. The intestinal neuro-immune axis: crosstalk between neurons, immune cells, and microbes. *Mucosal immunology* **14**, 555-565 (2021). https://doi.org/10.1038/s41385-020-00368-1

Jakob, M. O., Murugan, S. & Klose, C. S. N. Neuro-Immune Circuits Regulate Immune Responses in Tissues and Organ Homeostasis. *Front Immunol* **11**, 308 (2020). https://doi.org/10.3389/fimmu.2020.00308

Tan, E. K. *et al.* Parkinson disease and the immune system - associations, mechanisms and therapeutics. *Nat Rev Neurol* **16**, 303-318 (2020). https://doi.org/10.1038/s41582-020-0344-4

Jorfi, M. *et al.* Infiltrating CD8(+) T cells exacerbate Alzheimer's disease pathology in a 3D human neuroimmune axis model. *Nat Neurosci* **26**, 1489-1504 (2023). https://doi.org/10.1038/s41593-023-01415-3

Dantzer, R., O'Connor, J. C., Freund, G. G., Johnson, R. W. & Kelley, K. W. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* **9**, 46-56 (2008). https://doi.org/10.1038/nrn2297

Kopec, A. M., Smith, C. J. & Bilbo, S. D. Neuro-Immune Mechanisms Regulating Social Behavior: Dopamine as Mediator? *Trends Neurosci* **42**, 337-348 (2019). https://doi.org/10.1016/j.tins.2019.02.005

Dantzer, R. Neuroimmune Interactions: From the Brain to the Immune System and Vice Versa. *Physiological reviews* **98**, 477-504 (2018). https://doi.org/10.1152/physrev.00039.2016

McIlvried, L. A., Atherton, M. A., Horan, N. L., Goch, T. N. & Scheff, N. N. Sensory Neurotransmitter Calcitonin Gene-Related Peptide Modulates Tumor Growth and Lymphocyte Infiltration in Oral Squamous Cell Carcinoma. *Adv Biol (Weinh)*, e2200019 (2022). https://doi.org/10.1002/adbi.202200019

Zhang, Y. *et al.* Cancer cells co-opt nociceptive nerves to thrive in nutrient-poor environments and upon nutrient-starvation therapies. *Cell Metab* **34**, 1999-2017 e1910 (2022). https://doi.org/10.1016/j.cmet.2022.10.012

Campos, C. A. *et al.* Cancer-induced anorexia and malaise are mediated by CGRP neurons in the parabrachial nucleus. *Nat Neurosci* **20**, 934-942 (2017). https://doi.org/10.1038/nn.4574

Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018). https://doi.org/10.1016/j.neuron.2018.10.013

Takayanagi-Kiya, S. & Kiya, T. Activity-dependent visualization and control of neural circuits for courtship behavior in the fly Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 5715-5720 (2019). https://doi.org/10.1073/pnas.1814628116

Gaeta, A. *et al.* Are sex and gender considered in head and neck cancer clinical studies? *NPJ Precis Oncol* **7**, 84 (2023). https://doi.org/10.1038/s41698-023-00439-z

Park, J. O. *et al.* Sex Differences in the Prevalence of Head and Neck Cancers: A 10-Year Follow-Up Study of 10 Million Healthy People. *Cancers (Basel)* **14** (2022). https://doi.org/10.3390/cancers14102521

Dittberner, A. *et al.* Gender Disparities in Epidemiology, Treatment, and Outcome for Head and Neck Cancer in Germany: A Population-Based Long-Term Analysis from 1996

1074   to 2016 of the Thuringian Cancer Registry. *Cancers (Basel)* **12** (2020). https://doi.org:10.3390/cancers12113418

1076   127   Bianchini, C. *et al.* Pain in head and neck cancer patients: the role of gender. *J BUON* **24**, 2220-2226 (2019).

1078   128   Reyes-Gibby, C. C. *et al.* Survival patterns in squamous cell carcinoma of the head and neck: pain as an independent prognostic factor for survival. *J Pain* **15**, 1015-1022 (2014). https://doi.org:10.1016/j.jpain.2014.07.003

1081   129   Onken, M. D. *et al.* A surprising cross-species conservation in the genomic landscape of mouse and human oral cancer identifies a transcriptional signature predicting metastatic disease. *Clinical cancer research : an official journal of the American Association for Cancer Research* **20**, 2873-2884 (2014). https://doi.org:10.1158/1078-0432.CCR-14-0205

130   Moore, E. *et al.* Established T Cell-Inflamed Tumors Rejected after Adaptive Resistance Was Reversed by Combination STING Activation and PD-1 Pathway Blockade. *Cancer Immunol Res* **4**, 1061-1071 (2016). https://doi.org:10.1158/2326-6066.CIR-16-0104

131   Moore, E. C. *et al.* Enhanced Tumor Control with Combination mTOR and PD-L1 Inhibition in Syngeneic Oral Cavity Cancers. *Cancer Immunol Res* **4**, 611-620 (2016). https://doi.org:10.1158/2326-6066.CIR-15-0252

132   Shah, S., Caruso, A., Cash, H., Waes, C. V. & Allen, C. T. Pools of programmed death-ligand within the oral cavity tumor microenvironment: Variable alteration by targeted therapies. *Head Neck* **38**, 1176-1186 (2016). https://doi.org:10.1002/hed.24269

133   Cavanaugh, D. J. *et al.* Trpv1 reporter mice reveal highly restricted brain distribution and functional expression in arteriolar smooth muscle cells. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 5067-5077 (2011). https://doi.org:10.1523/JNEUROSCI.6451-10.2011

134   Deacon, R. M. Assessing nest building in mice. *Nature protocols* **1**, 1117-1119 (2006). https://doi.org:10.1038/nprot.2006.170

135   Kraeuter, A. K., Guest, P. C. & Sarnyai, Z. The Nest Building Test in Mice for Assessment of General Well-Being. *Methods in molecular biology* **1916**, 87-91 (2019). https://doi.org:10.1007/978-1-4939-8994-2_7

136   Kee, N., Teixeira, C. M., Wang, A. H. & Frankland, P. W. Imaging activation of adult-generated granule cells in spatial memory. *Nature protocols* **2**, 3033-3044 (2007). https://doi.org:10.1038/nprot.2007.415

137   Ting, J. T. *et al.* Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine Protective Recovery Method. *Journal of visualized experiments : JoVE* (2018). https://doi.org:10.3791/53825

138   Cantu, D. A. *et al.* EZcalcium: Open-Source Toolbox for Analysis of Calcium Imaging Data. *Front Neural Circuits* **14**, 25 (2020). https://doi.org:10.3389/fncir.2020.00025

1112

Acknowledgments: We thank the Histology and Imaging Core (Sanford Research, supported by

National Institutes of Health, National Institute of General Medical Sciences, Center of

Biomedical Research Excellence 5P20GM103548 and P30GM145398), specifically Claire

Evans who provided her services and expertise towards this project. We also thank the following

funding sources for their critical contributions to this work: National Institutes of Health,

National Institute of Dental and Craniofacial Research grant 1R01DE032712-01 (PDV),

Institutional Development Award (IDeA) from the National Institute of General Medical

Sciences of the National Institutes of Health grant 5P20GM103548 (PDV). DaCCoTA Scholar

Award supported by the National Institute of General Medical Sciences of the National Institutes

of Health grant U54 GM128729 (PDV). National Institute of Health, National Cancer Institute

grant R37 1R37CA242006-01A1 (MA). Stiefel family Discovery award (MA). Institutional

Research Grant (MA). Disruptive Science Moonshot award, MDACC (MA). Canadian Institutes

of Health Research grants 162211, 461274, 461275 (ST). National Institutes of Health (R01

CA193522; R21 NS130712) (RD).

1127

1128   **Materials & Correspondence:** Correspondence and material requests should be addressed to

1129   Paola D. Vermeer: Paola.Vermeer@sanfordhealth.org

1130   **Data availability statement:** Data will be available upon reasonable request.
1131
1132   **Figure legends**
1133
Figure 1. Tumor-infiltrating nerves form a circuit connecting to the brain. (**A**) Western blot
of MOC2-7 whole tumor lysates for Tau (**A**) and Doublecortin, DCX (**B**). Tumors were
harvested at different time points (as indicated) post-tumor implantation (n=3 mice/time point).
Mouse brain, positive control; MOC2-7, whole cell lysate control; β-actin, loading control.
Immunohistochemical staining for β-III tubulin (β3T, brown, arrows) of tumors harvested on
days 4 (**C**), 10 (**D**) and 20 (**E**) post tumor implantation. Scale bar, 100 µm. Dotted boxes are
shown at higher magnification below each panel; scale bar, 50µm. Arrows highlight positive
nerves staining. Representative *en face* confocal image of WGA positive (red) neurites within
tumor (**F**) and the ipsilateral trigeminal ganglion (**G**) from a mouse orthotopically implanted with
a MOC2-7 tumor and intra-tumorally injected with the tracer. Scale bars, 200µm. (**H**) Low
magnification section of ipsilateral and contralateral TGM ganglia from a MOC2-7 tumor-
bearing mouse that was intra-tumorally injected with WGA. The V3 branch of the TGM is
marked with an arrow. WGA (red) positive neurons found only in the ipsilateral TGM. Scale bar,
500 µm. (**I**) Compressed Z-stack of confocal images demonstrating restricted WGA (red) within

the tumor bed (cytokeratin, green). Vertical and horizontal cross-sections highlight that WGA
(red) localizes strictly within the tumor (green) and does not leak out. Yellow shows the merged
file. (**J**) Diagram showing the location of the trigeminal (TGM) ganglion, extension of its
neurites into the tumor bed and the existing circuit which includes the spinal nucleus of the TGM
(SpVc), the parabrachial nucleus (PBN) and the central amygdala (CeA) which get labeled
following WGA injection into tumor. (**K**) Representative confocal images of the SpVc, PBN and
CeA following injection of 2 µl of WGA into an oral MOC2-7 tumor (n=10 mice).
Representative confocal images of SpVc, PBN and CeA following injection of (**L**) 10 µl (n=3
mice) of WGA into the oral cavity of a non-tumor bearing control animal. Scale bar, 50 µm.
WGA (red looks pink due to DAPI), DAPI nuclear counterstain (blue).

Figure 2. Tumor-infiltrating neurons become transcriptionally and functionally altered.
Trigeminal (TGM) ganglia from tumor bearing (red) or non-tumor bearing (blue) mice were
analyzed by (**A**) quantitative PCR array. N=4 TGM/group from n=4 mice/group; n=4
arrays/group. Ct values for each gene normalized to that of a housekeeping gene. Statistical
analysis by multiple students t-test. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$, ****, $p<0.00001$. Data
are expressed as means ± standard deviation. (**B**) Representative fluorescent images of
dissociated neurons from the ipsilateral or contralateral TGM ganglia from a MOC2-7 tumor-
bearing mouse. N=6 of each ganglia harvested from 6 tumor-bearing animals with n= 4-8
neurons analyzed/ganglia. The color palette reflects the strength of the calcium signal. Images
are taken pre- and post-treatment with capsaicin. (**C**) Graph of average change in fluorescence
for all neurons analyzed in each group (ipsilateral and contralateral). Statistical analysis by
student's t-test. Arrow, time of capsaicin (300 nM) stimulation. Graph of average peak of the
curve (**D**) and area under the curve (**E**) from panel C. Statistical analysis by student's t-test. **,
$p<0.01$. (**F**) Quantification of Ca^{+2} responses to capsaicin (300 nM) from non-tumoral and
tumor-infiltrating, tracer^{+} (Tumor-infiltrating) neurons. N= 4-8 neurons/group from n=4 control
and n=7 tumor-bearing mice. Statistical analysis by paired students t-test, *, $p<0.05$. Data
expressed as means ± SEM. (**G**) Representative western blot of trigeminal ganglia from non-
tumor (N) or MOC2-7 tumor-bearing (T) animals for phosphorylated TRPV1 (pTRPV1, Ser502,
Ser800), σ1 receptor (σ1R) and β-actin (loading control). Densitometric quantification of
western blots for pTRPV1 (**H**) and σ1R (**I**). Statistical analysis by student's t-test. *, $p< 0.05$.
Data are expressed as means□±□SEM. (n=3-4 mice/group).

Figure 3. Elevated CNS neuronal activity in tumor bearing mice. Representative fluorescent
photomicrographs of the spinal nucleus of the TGM (SpVc) (**A**), parabrachial nucleus (PBN) (**B**)
and central amygdala (CeA) (**C**) from a control (-tumor) and MOC2-7 tumor-bearing (+tumor)
mouse brains immuonfluorescently stained for cFos (red) or ΔFosB (green). Nuclei
counterstained with DAPI (blue). Dotted circle denotes each brain region. Scale bar, 100 µm.
Brain landmarks: Spc, superior cerebellar peduncle; VL, lateral ventricle; IA, intercalated
amygdalar nucleus. Quantification of cFos and ΔFosB staining in SpVc (**D**), PBN (**E**) and CeA
(**F**) from control (-tumor) and tumor-bearing (+tumor) mice. N= 4 mice/group with n=2-4
sections analyzed/brain region/mouse/group. Data are expressed as the mean ± SEM. Statistical
analysis by one-way ANOVA*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ns, not significant. Scale bar,
100 µm. Representative heat maps of regions of interest (ROI), each of which represent one
neuron, within an *ex vivo* brain slice from non-tumor (control) (**G**) and tumor-bearing (**H**) mice.
The color palette reflects the strength of the calcium signal. Arrow denotes the time of KCl (30

mM) application. N= 3 mice/group with n=3 slices/brain. Quantification of calcium imaging data (n= 5 mice/group with n=2-4 sections analyzed/mouse) analyzing the peak of the curve (**I**) and the area under the curve (**J**). Statistical analysis by student's t-test. ****, p< 0.0001; ***, p< 001.

Figure 4. Intra-tumoral neurons impact behavior. (**A**) Bright field photomicrograph of TGM ganglion from C57BL/6 or TRPV1-Cre::Floxed-DTA mice IHC stained for TRPV1 (brown). Scale bar, 100μm. Quantification of cFos and ΔFosB from the brains of C57BL/6 (C57) or TRPV1-Cre::Floxed-DTA (DTA) animals with MOC2-7 oral tumors in the Spinal nucleus of the trigeminal (SpVc) (**B**), the parabrachial nucleus (PBN) (**C**) and the central amygdala (CeA) (**D**). N=4 mice/group with n=2-4 sections analyzed/brain region/mouse. Statistical analysis by one-way ANOVA. **, p<0.01; ***, p<0.001. Data displayed as mean ± SEM. C57BL/6 (black, n=15) and TRPV1-Cre::Floxed-DTA (blue, n=14) mice were orthotopically implanted with MOC2-7 tumor and behavior assessed weekly and statistically analyzed by repeated measures ANOVA. (**E**) Graph of nesting scores over time. There is a main effect of time such that there is a decline in nesting performance over time. We also see a time by strain interaction in that C57BL/6 mice show a greater decline than the nociceptor neuron ablated TRPV1-Cre::Floxed-DTA mice. Post hoc testing shows a significant difference in nesting on day 27 (p=0.0315). (**F**) Graph of time to interact with the cookie in the cookie test. There is a main effect of time such that the time to interact with the cookie declines over time. There is a main effect of strain such that nociceptor neuron ablated TRPV1-Cre::Floxed-DTA mice were faster to interact with the cookie. Finally, there is a time by strain interaction such that nociceptor neuron ablated TRPV1-Cre::Floxed-DTA mice showed progressively faster task performance, while in C57BL/6 mice interaction time plateaued. Post hoc testing demonstrates a significant difference between the groups on days 3, 22, and 27 (p<0.05). (**G**) Tumor growth curves for mice that underwent behavioral testing. Statistical analysis by repeated measures ANOVA. There is a main effect of time such that there is an increase in tumor volume over time. We also see a time by strain interaction in that C57BL/6 mice have larger tumors than nociceptor ablated TRPV1-Cre::Floxed-DTA mice. (**H**) Kaplan-Meier survival curve for mice in panel I. Statistical analysis by Log-rank (Mantel-Cox) test. ****, p<0.0001. (**I**) Graph of % change in weight from baseline. Dotted line represents baseline. Statistical analysis by repeated measures ANOVA. There was a main effect of time such that both groups showed an initial increase in weight followed by a decrease. Post hoc testing shows a significant difference between the groups on days 5, 10, and 20. (**J**) Graph of % change from baseline in food consumption. Dotted line represents baseline. Statistical analysis by repeated measures ANOVA. There is a main effect of time such that there is a decline in % food disappearance over time. There is a time by strain interaction such that C57BL/6 mice show a greater decline in % food disappearance compared to nociceptor neuron ablated TRPV1-Cre::Floxed-DTA mice. In all panels, error bars are SEM.

Figure 5. Denervation of tumors improves behavior, attenuates body weight loss, and reduces tumor growth. (**A**) Graph of nesting scores over time. Statistical analysis by repeated measures ANOVA (group x time): The time factor and its interaction with the group factor are significant with a significant difference between RTX- and vehicle-treated mice on the last time point (p<0.001). (**B**) Graph of % change in weight from baseline. Statistical analysis by repeated measures ANOVA (group x time): The time factor and its interaction with the group factor are significant with a significant difference between groups on the last time point (p<0.01). (**C**) Tumor volume according to treatment and time: Statistical analysis by repeated measures

1240 ANOVA (treatment x time): The time factor and its interaction with the group factor are
1241 significant with a significant difference between treatments on the last two time points
1242 (p<0.001). N=20 mice/group.
1243
Figure 6. Treating for pain does not restore all behaviors. C57BL/6 male mice were
orthotopically implanted with MOC2-7 tumors and separated into three groups: carprofen treated
(10mg/kg, blue, n=10), buprenorphine treated (3.25mg/kg, red, n=10) or vehicle treated (black,
n=9). Mice underwent behavioral testing. All analyses are by repeated measures ANOVA. (**A**)
Data are graphed as % of baseline for overnight wheel running. The dotted line represents the
baseline. There is a main effect of time such that there is a decline in voluntary wheel running
over time. We also see a time by drug (carprofen) interaction in that vehicle-treated mice show a
greater decline in wheel running than carprofen-treated mice. Post hoc testing shows a significant
difference in wheel running (p<0.0001). (**B**) There is a main effect of time such that there is a
decline in voluntary wheel running over time. We also see a time by drug (buprenorphine)
interaction in that vehicle-treated mice show a greater decline in wheel running than the
buprenorphine-treated mice. Post hoc testing shows a significant difference in wheel running
(p<0.0001). (**C**) Graph of nesting scores over time. When comparing vehicle vs carprofen
groups, there is a main effect of time such that there is a decline in nesting performance over
time. We also see a time by drug (carprofen) interaction in that vehicle-treated mice show a
greater decline than carprofen-treated mice. Similarly, we see a main effect of time when
comparing the vehicle and buprenorphine groups. There is a time by drug (buprenorphine)
interaction in that vehicle-treated mice show a greater decline than buprenorphine-treated mice.
(**D**) Graph of % change in weight from baseline. The dotted line represents the baseline. There
was a main effect of time. There is also a drug (carprofen) by time interaction in that vehicle-
treated mice show a greater decline in weight than the carprofen-treated mice. In fact, carprofen
treated mice gain weight. Similarly, when comparing the vehicle and buprenorphine groups,
there is a main effect of time. There is also an interaction of drug (buprenorphine) by time in that
vehicle-treated mice show a greater decline in weight than the buprenorphine-treated mice. In
fact, the buprenorphine treated mice gain weight. When carprofen and buprenorphine-treated
animals are compared, there is an interaction of time with treatment such that buprenorphine
treated animals gain more weight than carprofen-treated animals. (**E**) Graph of % change in food
disappearance from baseline. When comparing the vehicle vs carprofen group, there is a main
effect of time such that there is a decline in food disappearance over time. We also see a time by
treatment interaction such that carprofen-treated animals do not decline in food disappearance as
much as vehicle treated animals. When comparing the vehicle vs buprenorphine group, there is a
main effect of time. In the buprenorphine treated group, there is a time by treatment interaction
such that the buprenorphine-treated animals demonstrate the smallest decline in food
disappearance. Finally, comparison of the carprofen and buprenorphine groups shows that there
is an interaction between time and treatment such that buprenorphine-treated animals show the
least reduction in food disappearance. (**F**) Tumor volume was monitored weekly for mice in all
groups. There is a main effect of treatment such that mice treated with carprofen show a greater
decline in tumor volume than buprenorphine-treated mice.
1282
1283
1284
1285

1286