Tumor-infiltrating nerves functionally alter brain circuits and modulate behavior in a mouse model of head-and-neck cancer

Jeffrey Barr [1†], Austin Walz [1†], Anthony C. Restaino[1,3], Moran Amit[4], Sarah M. Barclay[1],

Elisabeth G. Vichaya[2], William C. Spanos[1,3], Robert Dantzer[4], Sebastien Talbot[5], Paola D.

Vermeer[1,3,*]

[1]Sanford Research, Cancer Biology and Immunotherapies Group, Sioux Falls, South Dakota,

USA.

[2]Baylor University, Department of Psychology and Neuroscience, Waco, TX, USA.

[3]University of South Dakota, Sanford School of Medicine, Vermillion, SD, USA.

[4]University of Texas, MD Anderson Cancer Center, Houston, TX, USA.

[5]Queen's University, Department of Biomedical and Molecular Sciences, Kingston, Ontario,

Canada.

[†] equal first authors

*Corresponding author   Email: Paola.Vermeer@sanfordhealth.org

Conflict of interest: RD is a consultant for GoodCap, Toronto; the consultancy is not related to

the present work.

Abstract: Cancer patients often experience changes in mental health, prompting an exploration into whether nerves infiltrating tumors contribute to these alterations by impacting brain functions. Using a male mouse model for head and neck cancer, we utilized neuronal tracing techniques and show that tumor-infiltrating nerves indeed connect to distinct brain areas via the ipsilateral trigeminal ganglion. The activation of this neuronal circuitry led to behavioral alterations represented by decreased nest-building, increased latency to eat a cookie, and reduced wheel running. Tumor-infiltrating nociceptor neurons exhibited heightened activity, as indicated by increased calcium mobilization. Correspondingly, the specific brain regions receiving these neural projections showed elevated cFos and delta FosB expression in tumor-bearing mice, alongside markedly intensified calcium responses compared to non-tumor-bearing counterparts.

The genetic elimination of nociceptor neurons in tumor-bearing mice led to decreased brain Fos expression and mitigated the behavioral alterations induced by the presence of the tumor. While analgesic treatment successfully restored behaviors involving oral movements to normalcy in tumor-bearing mice, it did not have a similar therapeutic effect on voluntary wheel running. This discrepancy points towards an intricate relationship, where pain is not the exclusive driver of such behavioral shifts. Unraveling the interaction between the tumor, infiltrating nerves, and the brain is pivotal to developing targeted interventions to alleviate the mental health burdens associated with cancer.

Significance Statement: Head and neck cancers are infiltrated by sensory nerves which connect to a pre-existing circuit that includes areas in the brain. Neurons within this circuit are altered and mediate modifications in behavior.

Introduction:

The prevalence of mental health disorders (e.g., depression, anxiety, suicide) in cancer patients is

significantly greater than in the general population [1-3]. For those patients with no prior

psychiatric history, a cancer diagnosis increases the risk of mental health decline [4]. The presence

of cancer introduces many stressors (physical, financial, relational) into the lives of patients, thus

a negative impact on mental health may not be surprising. However, these changes persist even

in long-term cancer survivors. For instance, decade-long cancer survivors maintain an increased

incidence of depression (approximately 12%, depending on the cancer type) as compared to the

population at large (3-5%) [5-7]. While the intensity and prevalence of psychological symptoms in

cancer patients fluctuate before, during, and after treatment for a given type of cancer, and

between cancer types, it remains higher than in the general population [8-10]. Whether the

association of cancer with impaired mental health is directly mediated by the disease, its

treatment or both remains unclear, yet it suggests that the development of a tumor may alter

brain functions.

We have demonstrated the presence of TRPV1-expressing nociceptor neurons in head and neck

squamous cell carcinomas (HNSCC) [11], melanoma, cervical [12], and ovarian cancers [13]. Nerve

recruitment to the tumor bed is an active process that involves the release of soluble factors,

including neurotrophins [14-16] and neuropeptides [17]. Tumor-released small extracellular vesicles

(sEVs) also recruit loco-regional nerves to the tumor bed [11,18]. While these and other studies

establish that solid peripheral tumors engage with the peripheral nervous system [11,13,19-23], it

raises the possibility of a direct neuronal connection from the tumor to the brain.

A recent study used pseudorabies virus and mapped a connection from tumor-infiltrating nerves

in an orthotopic model of murine lung cancer to areas in the brain [24]. We expand these findings

and demonstrate that HNSCC-associated nerves are transcriptionally and functionally altered and

project to discrete regions in the brain. The brain neurons connected to the tumor harbor

increased activity, which is associated with behavioral alterations in tumor-bearing animals.

Consistent with this, newly diagnosed HNSCC patients suffer high rates of depression and

anxiety and lower quality of life [25]. As most cancer patients face cancer-related pain [26-28], we

tested whether treating pain could restore normal behavior. While pain treatment restored normal

function at the tumor site, such as nesting behavior, it only partially restored normal voluntary

running wheel behavior. Our findings suggest that, in addition to pain, tumor-infiltrating nerves

communicate signals to the brain that lead to cancer-associated changes in behavior.

Results

Tumor innervation begins early in disease. MOC2-7 cells are HNSCC cancer cells derived

from a CXCR3 null mouse under a C57BL/6 background [29]. Their implantation in male mice

results in dense innervation of tumors with nociceptor nerves[38,39]. To define the timing of tumor

innervation, wildtype male mice were orthotopically implanted (oral cavity) with MOC2-7 cells,

and tumors were collected on days 4-, 10-, and 20- post-implantation. Western blot analysis of

whole tumor lysate indicated expression of Tau, a neuronal marker [40], as early as day four post-

tumor implantation, which increased over time (**Fig. 1A**; full westerns in **Supplemental Fig. 1A-**

B; densitometric quantification of westerns, **Supplemental Fig. 1E**). A similar increase in the

neuronal marker doublecortin (DCX) was noted (**Fig. 1B**; full western **Supplemental Fig. 1C-D**;

densitometric quantification of westerns, **Supplemental Fig. 1F**). While DCX is well-known for

its expression in immature neurons, it is also expressed in adult peripheral neurons, including

dorsal root ganglia [41], and in non-neuronal tissues[42]. The DCX signal did not originate from the

tumors, as MOC2-7 cell lysate was negative for this protein (**Fig. 1B**, Full western

Supplemental Fig 1C). Consistent with this, immunohistochemical staining of these tumors

with the neuronal marker β-III tubulin (β3T), demonstrated the increasing presence of nerves

beginning on day four post-tumor implantation (**Fig. 1C-E**).

Tumor-infiltrating nerves map to the ipsilateral trigeminal ganglion and into the CNS. To

map the origin of tumor-infiltrating nerves and define the circuits they converge upon, mice with

palpable oral MOC2-7 tumors (approximately day 15 post-tumor implantation) were intra-

tumorally injected with wheat germ agglutinin, WGA, a neural tracer conjugated to a

fluorophore (n= 10 mice). WGA is a lectin molecule that specifically binds to sialic acid residues

present ubiquitously on neuronal membranes. It has been utilized extensively to map neuronal

circuits centrally and in the periphery [43,44] and is a known transganglionic and transynaptic

neuronal tracer [45-47] making it ideal for mapping neural circuits [48-52]. Following tracer injection,

tumor growth was permitted for an additional 3-7 days to allow time for tracer labeling to occur.

Animals were then euthanized, and tumors, trigeminal (TGM) ganglia, and brains were harvested

and analyzed by microscopy. Microscopic examination of tumors revealed nerves with robust

WGA signals (**Fig 1F**) as well as the V3 branch of the ipsilateral, but not the contralateral, TGM

ganglion (**Fig 1G,H**). Consistent with the restricted labeling of tumor-infiltrating nerves, the

tracer did not diffuse outside the tumor mass (**Fig. 1I**).

Examination of brains from MOC2-7 tumor-bearing animals revealed tracer[+] neurons in specific

regions [including spinal nucleus of the trigeminal (SpVc), parabrachial nucleus (PBN), and

central amygdala (CeA)] (**Fig 1J**). Sections of brain from these regions show the presence of

tracer-positive neurons (**Fig 1K**). Tracer injections of equivalent volume and concentration into

the oral cavities of control non-tumor-bearing animals did not label the TGM ganglia nor areas in

the brain (data not shown). A larger volume (10µl) of tracer injected into non-tumor bearing

mice resulted in tracer labeling of ipsilateral TGM neurons and brain (**Fig. 1L**). These control

studies indicate that the nerve density and distribution present in the tumor bed are higher than in

control mice, with the consequence that a small volume (2 µl instead of 10 µl) of tracer is

sufficient to result in nerve labeling. The mapped circuit encompasses pre-existing connections

to brain areas that regulate pain and affect [53-56]. These data indicate that the nerve infiltration into

the tumor extends this circuit.

Tumor-infiltrating neurons are transcriptionally modified. The observation of nerves

infiltrating the tumor mass prompted us to ask whether they undergo alterations by their mere

presence within this "foreign" environment. Consistent with this, we previously found that

melanoma-infiltrating neurons have a unique transcriptome [57]. To test whether this is also the

case in HNSCC, we orthotopically implanted MOC2-7 cells into the oral cavity of male wildtype

mice. After 14 days, TGM neurons were harvested and analyzed by qPCR using a commercial

array for neural transmission and membrane trafficking genes. Compared to control TGM (non-

tumoral) neurons, ipsilateral TGM neurons from tumor-bearing animals harbored increased

expression (<4-fold) in genes involved in neuronal signaling/receptors (*Gabrg1, Gabra4,*

Grin2c, Grm3) and synaptic transmission (*Gria2*) (**Fig. 2A; Supplemental Table 1**) [58]. Of note,

we purposefully used whole trigeminal ganglia rather than FACS-sorted tracer-positive

dissociated neurons to avoid artificially imposing injury and altering the transcriptome of these

cells [59,60].

Tumor-infiltrating neurons are functionally changed. Given this transcriptomic alteration, we tested whether tumor-infiltrating neurons exhibit functional alterations. Thus, MOC2-7 tumor-bearing mice were intra-tumorally injected with the fluorophore-conjugated WGA tracer. Five days later, tracer[+] TGM ganglia were cultured, and responsiveness to noxious stimuli was analyzed by calcium microscopy. Neuronal Ca^{+2} responses to capsaicin (300 nM), which binds and activates TRPV1 channels, were measured by Fluo-4AM fluorescence microscopy. We found that tracer[+] tumor-infiltrating ipsilateral TGM neurons show increased responsiveness to capsaicin compared to tracer negative contralateral neurons (amplitude, **Fig. 2B-D;** the area under the curve**, Fig. 2E**). Similar to contralateral TGM neurons, non-tumor-bearing TGM neurons elicited a normal response to capsaicin (**Fig. 2F**).

This heightened sensitivity could reflect increased TRPV1 expression and/or its phosphorylation. We tested whether this was the case using western blotting and discovered an overexpression of the sigma 1 receptor (σ1R) and increased TRPV1 phosphorylation in the TGM ganglia from tumor-bearing animals (**Fig. 2G-I,** full westerns **Supplemental Fig 2A-C**). Of note, the σ1R is an endoplasmic reticulum chaperone protein that directly interacts with TRPV1 and regulates its membrane expression, [61] while protein kinase C phosphorylation of serine 502 and 800 of TRPV1 reduces the receptor's activation threshold [62-65]. Taken together, our data indicate that HNSCC-infiltrating nerves have a unique transcriptome and a heightened sensitivity to noxious stimuli characterized by an increased TRPV1 expression and phosphorylation consistent with TRPV1 sensitization secondary to oral cancer [66].

Tumor-brain circuit neurons harbor elevated activity. The transcriptional and functional changes evident in tracer[+] tumor-infiltrating neurons could result in alterations in central target neurons. To assess this, brain sections from MOC2-7 tumor-bearing and non-tumor-bearing

animals were immunofluorescently stained for cFos and ΔFosB, two markers of neuronal activity

with different courses of expression [67-69]. ΔFosB expression was increased in several brain

regions of tumor-bearing animals, while cFos expression was also increased in the PBN (**Fig.**

3A-F). Predominant differences in the long-lived ΔFosB were expected [70], as those with the

short-lived cFos changes are more challenging to capture using single time point assessment.

Next, we assessed the neuronal activity of tumor-bearing animal brains using stereotaxically

injected AAV1-Syn-GCaMP6f, a viral vector encoding a neuron-specific synapsin-driven

calcium sensor [71], into the PBN. Two weeks after intra-cranial injection of the virus, mice were

orally implanted with MOC2-7 cells. Approximately 18 days post-tumor inoculation, the animals

were euthanized, and the neuronal calcium activity was recorded in *ex vivo* brain slices using a

mini scope (**Suppl Fig. 3A,B**). While baseline fluorescence was similar between tumor-bearing

and control animals, that recorded upon stimulation (KCl; 30 mM) was significantly higher in

neurons of tumor-bearing animals (**Fig. 3G-J; Supplemental Fig 3A,B**). These functional and

Fos staining data indicate that central neurons within the tumor-brain circuit are functionally

altered compared to their healthy brain counterparts.

Ablation of tumor-infiltrating neurons attenuates cancer-induced brain alterations. Cancer

patients, and even more notably, survivors, experienced poor mental health [1,72-74]. The neural

connection between tumor and brain, together with our finding that TRPV1-expressing

nociceptor neurons within this circuit become functionally altered, might contribute to these

changes. To test whether this is the case, we genetically engineered mice with ablated nociceptor

neurons (TRPV1^{cre}::DTA$^{fl/wt}$). TRPV1^{cre}::DTA$^{fl/wt}$ animals lack TRPV1-expressing cells

(including neurons) as well as many other nociceptive neurons. These nociceptor neuron ablated

mice have been previously characterized and lack the expected sensitivity to temperature as well

as itch and pain reactions to chemical mediators such as capsaicin [75]. cFos and ΔFosB

immunostaining of brains from nociceptor ablated and control (C57BL/6) mice show no

significant differences (**Supplemental Fig 3C-E**) indicating that, in the absence of a malignancy,

the neurons in these regions are not differentially activated. First, we confirmed the absence of

TRPV1$^+$ neurons in the TGM of these ablated animals (**Fig. 4A**). We then analyzed cFos and

ΔFosB expression in the brain of MOC2-7-bearing nociceptor intact and ablated mice. While Fos

expression was similar between non-tumor bearing mice of the two genotypes (**Supplemental**

Fig. 3C-E), the absence of nociceptor neurons decreases cFos and ΔFosB in the PBN, and

ΔFosB in the SpVc (**Fig. 4B, C**). Other tested regions were not impacted (**Fig. 4D**). These data

suggest that the tumor-brain communication was disrupted in the absence of nociceptor neurons.

Alleviation of pain does not always restore behavior. To evaluate the effects of this disruption

on cancer-induced behavioral changes, we assessed the animals' general well-being through

nesting behavior [32] and anhedonia using the cookie test [76,77], as well as body weight and the

speed of food disappearance as a surrogate for oral pain. Nociceptor neuron ablated mice showed

increased nesting performance (**Fig. 4E**) and decreased anhedonia (**Fig. 4F**) compared to intact

mice. This was accompanied by smaller tumor growth (**Fig. 4G**) and increased survival (**Fig.**

4H). While both groups showed similar body weight loss (**Fig. 4I**), nociceptor ablated animals

show a transient (Wk 1, 4, 5) decrease in food disappearance (**Fig. 4J**). Significant effects for all

statistical analyses in Figure 4 are presented in Supplementary Table 2.

The developmental ablation of TRPV1 neurons can lead to unintended effects on tumor

innervation and behavior. Thus, we chemoablated TRVP1 neurons in 4-week-old juvenile

C57BL/6 mice using resiniferatoxin (RTX) [39,40]. We examined nesting behavior in RTX and

vehicle-treated C57BL/6 male mice orthotopically implanted with MOC2-7 cells. An additional

control group consisted of age-matched C57BL/6 mice without tumors. Although all groups

displayed similar nesting scores initially, the scores in vehicle-treated mice declined significantly

faster compared to RTX-treated mice (**Fig 5A**). Body weight decreased in all tumor-bearing

mice, but the extent of weight loss was similar between vehicle and RTX-treated mice (**Fig 5B**).

Similar to the developmental ablation of TRPV1 neurons, their chemoablation post-development

results in a similar decrease in tumor growth compared to controls (**Fig 5C**).Since HNSCC

tumors cause oral pain and both nesting and the cookie test require the use of the mouth, we also

evaluated the effect of nociceptor ablation on voluntary wheel running to gain additional insights

into cancer-associated fatigue, a surrogate for depressive-like behaviors in tumor-bearing mice [78-

82]. To assess the potential influence of cancer-associated pain, 1-week post tumor inoculation,

groups of mice were treated with carprofen (a non-steroidal anti-inflammatory), extended-release

buprenorphine (opioid), or vehicle. We found that wheel running decreased as tumors

progressed, an effect partially alleviated by carprofen and buprenorphine (**Fig 6A,B**). The

nesting behavior (**Fig 6C**), body weight (**Fig 6D**), and food disappearance (**Fig 6E**) were also

alleviated by the painkillers. Neither carprofen nor buprenorphine impacted MOC2-7 tumor

growth compared to vehicle-treated animals (**Fig 6F**), but buprenorphine increased tumor growth

compared to carprofen-treated mice (significant effects for all statistical analyses in Figure 5 are

presented in Supplementary Table 3). These findings suggest that choosing pain management

drugs in the context of cancer can potentially influence tumor growth adversely. Furthermore,

the data reveals that behaviors associated with the tumor site are adversely affected by cancer-

associated pain. However, merely alleviating pain does not fully restore all behaviors, as

evidenced by persistent issues in activities like wheel running.

Discussion

We have previously demonstrated that head and neck squamous cell carcinoma is associated with innervation by sensory neurons and that substance P, one of the principal neuropeptides these fibers release, drives malignant cell proliferation and migration [38]. Significantly, patients suffering from HNSCC often experience depressive disorders, which are poorly managed clinically [83-85]. Here, we now found that nociceptor neurons infiltrating HNSCC connect to a pre-existing brain-projecting circuit [86,87] that includes the trigeminal ganglia, the spinal nucleus of the trigeminal (SpVc), as well as the parabrachial nucleus (PBN), and central amygdala (CeA). This connection occurs independently of pain, and this cancer-brain circuit drives behavioral alterations in tumor-bearing mice. While the elimination of TRPV1-expressing nociceptor neurons reduces tumor growth and enhances survival rates, it also highlights a potential therapeutic target for mitigating depressive behaviors in cancer patients. This underscores the complex interplay between sensory neurons, cancer progression, and mental health, uncovering a two-pronged approach to improving HNSCC patients' physical and psychological health.

In melanoma [57] and HNSCC, we demonstrated significant transcriptional alterations within tumor-infiltrating nerves. Several mechanisms can account for these changes. For instance, TRPV1 stimulation is sufficient to trigger the activation of the AP-1 transcription factor within neurons [88]. Such TRPV1 activation could occur in response to low pH [89] or hypoxia [90-92], which are both prevalent within the tumor microenvironment. Alternatively, we previously demonstrated that tumor-released sEVs, which transport microRNAs (miRNAs), induced sprouting of loco-regional nerves to the tumor bed [11,93]. These sEV-transported miRNAs could

also modulate various neuronal transcription factors and, in turn, drive the transcriptomic

changes we observed in these neurons.

Our data indicate that these peripheral neuronal changes also influence the functioning of the

brain areas to which they connect. Although we did not make the complete cartography of all

brain regions utilized by tumor-infiltrating nerves, we identified a few critical nuclei. Subsequent

work will use emerging circuit mapping techniques for whole brain profiling [94]. Nevertheless,

our tracing in tumor-bearing nociceptor neuron ablated, and intact animals was sufficient to

functionally implicate TRPV1-expressing neurons in this communication.

Trigeminal ganglia neurons are composed of ~80% of $Na_V1.8^+$ nociceptor neurons, with one-half

being peptidergic (TRPV1$^+$, TRPA1$^+$) neurons [95,96,75]. In the TRPV1^{cre}::DTA$^{fl/wt}$ mouse, these

peptidergic neurons are eliminated, leaving intact ~50% of pain-sensing neurons. The remaining

presence of these non-peptidergic, largely MrgD$^+$, neurons might explain why nesting behavior

in TRPV1^{cre}::DTA$^{fl/wt}$ tumor-bearing animals is only partially restored.

Inflammation contributes to pain in cancer [97]. Consistent with this, treatment of tumor-bearing

mice with analgesic drugs (carprofen, buprenorphine) resulted in complete restoration of nest

building and performance in the cookie test. These data show that the constraint imposed by the

tumor on behavioral activities that require the use of the oral cavity is mainly due to pain and not

to the physical interference of the tumor mass with the pattern of oral activities that need to take

place for mice to perform the corresponding behavior.

Cancer pain often interferes with activities that don't involve oral movements, a phenomenon

evident in the impaired voluntary wheel running observed in tumor-bearing mice. Although

analgesic drugs can partially mitigate this deficit, they fail to restore this behavior fully, despite

their efficacy in completely restoring oral activity behaviors like nesting and the cookie test. This

suggests that factors beyond pain contribute to the observed behavioral changes. These

alterations could be attributed to depression [98] or represent a competition between the energy

demands of the HNSCC malignant cells and the host's skeletal muscles [99-101].

Our current findings suggest that interrupting tumor-to-brain communication can mitigate the

mental health decline often associated with cancer. Yet, a higher prevalence of depression

persists among long-term cancer survivors compared to individuals without a cancer history.

Having received treatment and showing no evidence of disease, these survivors continue to battle

mental health issues [6,7]. This phenomenon prompts a critical question - why does this

discrepancy exist? It is conceivable that these patients, through treatments like surgery, radiation,

or chemotherapy, have disrupted the tumor-brain connection and have sustained this

disconnection over the years. However, unlike animal models, their mental health decline is not

reversed. Several speculations, grounded in known factors, can be considered.

The human experience with cancer is distinct from that of mouse models; individuals can live

with malignant growths for extended periods, often unaware of their existence due to

asymptomatic or nonspecific symptoms. During this undiagnosed period, the tumor-brain circuit

might induce irreversible changes in central neurons through activity-dependent transcriptional

modifications [102,103]. These alterations, anchored in neuronal plasticity, adaptation, and behavior,

may become permanent even after the cancer is diagnosed and treated.

Moving forward, to reverse these entrenched brain changes, understanding the specific

transcriptional alterations incurred by central neurons is vital. Unraveling this complexity could

pave the way for therapies that reverse the neuronal activity impacts, fostering improved mental

health. Although in its early stages, our insight into the role of nerves in cancer is growing, and

highlighting neuronal targets for pharmacological interventions.

Our study is not exempt from limitations. First, our exploration of the tumor-to-brain pathway

was confined to mice with HNSCC and utilized a single cancer cell line. Hence, the universality

of our conclusions warrants verification through additional HNSCC cell lines and diverse cancer

models. Furthermore, the exclusive focus on HNSCC, particularly considering the nerve-rich

oral region, raises questions about the pathway's establishment and behavioral impact in other

contexts. Our behavioral assessment was not exhaustive; a comprehensive analysis of nerve-

dependent and independent behaviors remains forthcoming. Behaviors could also be influenced

by elements like soluble factors or the energy competition between the tumor and behavioral

activities. Our studies have also solely involved male mice, presenting a clear gap in

understanding the potential sex differences in the development of the tumor-to-brain

communication pathway and its subsequent influence on behavior. The inclusion of female

subjects in future research is essential to provide a comprehensive insight into these processes.

Second, our nerve tracing methodology, where we injected 2 µl of WGA into the tumor bed, is

designed to label only the tumor-infiltrating nerves and their connections. However, this

approach fails to label *all* tumor-infiltrating nerves, predominantly those not close to the WGA-

injected region and nerves that are in contact with or influenced by the tumor but are not directly

infiltrating it. This limitation is a calculated one, reflecting a technical trade-off. We aimed to

ensure the specificity of capturing only tumor-infiltrating nerves, which meant sacrificing the

comprehensiveness of labeling all nerves associated with the tumor. This constraint extends to

our calcium imaging studies and brain tracing. Although we can confidently assert that all

labeled neurons are indeed tumor-infiltrating or connected, we cannot conclusively state that all

tracer-negative ipsilateral nerves are unconnected to the tumor-brain circuit.

Even with these constraints, our findings are important. We have demonstrated that tumor-

infiltrating nerves are integrated into a pre-established neuronal circuit. This circuit extends from

the tumor bed to the TGM ganglion, connects to the SpVc, and projects into the brain,

influencing behavior in both pain-dependent and independent ways. This revelation paves the

way for in-depth exploration into the mechanistic underpinnings of cancer-associated alterations

in well-being and mental health, shedding light on potential therapeutic interventions to alleviate

these profound effects.

MATERIALS AND METHODS

Study approval. All animal studies were performed with approval from the Institutional Animal

Care and Use Committee at Sanford Research and were within institutional guidelines and

complied with all relevant ethical regulations. Sanford Research has an Animal Welfare

Assurance on file with the Office of Laboratory Animal Welfare (assurance number: A-4568-01)

and is accredited by AAALAC, Intl. Sanford Health is also a licensed research facility under the

authority of the United States Department of Agriculture (USDA) with USDA certificate number

46-R-011.

Inclusion & Ethics Statement. All studies utilizing animals were performed with approval from

the appropriate ethical bodies to ensure sufficient protection was in place. This study was a

collaborative, multi-disciplinary, multi-institutional effort with contributions from researchers in

academic positions.

Cell lines. MOC2-7 cells (RRID:CVCL_ZD34) [29] (previously known as MOC7) [30] were a kind

gift from Dr. Ravindra Uppaluri (Dana-Farber Cancer Institute, Boston, MA). They were

maintained in DMEM with 10% fetal calf serum and cultured at 37°C and 5% CO_2, with culture

medium refreshed every three days. These cells can be obtained through Kerafest

(www.kerafest.com).

Animal studies. All animal experiments were performed in the Sanford Research Animal

Resource Center which is a specific pathogen free facility. All mice were maintained in IVC

Tecniplast Green line Seal Safe Plus cages which were opened only under aseptic conditions in

an animal transfer station. All cages were changed every other week using aseptic technique. All

cages had individual HEPA filtered air. Animal rooms were maintained at 75°F, 30-70%

humidity, with a minimum of 15 air changes per hour, and a 14:10 light/dark cycle. Corncob

bedding, which was autoclaved prior to use, was maintained in all cages. Irradiated, sterile food

(Envigo) and acidified water (pH 2.8-3.0) were available *ad libitum*. There was a maximum of 5

mice/cage. All animals were observed daily for abnormal behavior, signs of illness or distress,

the availability of food and water and proper husbandry. Animals injected with murine tumor

cells were 10-week-old C57BL/6 or TRPV1^{cre}::DTA$^{fl/wt}$ mice (The Jackson Laboratory)

weighing approximately 24 gm. Investigators were blinded to the groups when assessing animals

(e.g., measuring tumors). Animals were numbered by ear punch and cage number.

Nociceptor neuron ablated TRPV1^{cre}::DTA$^{fl/wt}$ animals were generated by crossing ROSA26-

DTA (diphtheria toxin A, B6.129P2-*Gt(ROSA)26Sor $^{tm1(DTA)Lky}$*/J) and TRPV1-Cre (B6.129-

Trpv1$^{tm1(cre)Bbm}$ /J) [31] mice; the progeny (TRPV1^{cre}::DTA$^{fl/wt}$) express DTA (diphtheria toxin

fragment A) under control of the TRPV1 promoter, thereby genetically ablating all TRPV1

expressing cells (including TRPV1-expressing neurons) throughout development. Absence of

TRPV1-expressing sensory neurons was validated by IHC staining of TGM ganglia for TRPV1.

Chemoablation of TRPV1-expressing neurons post-development was accomplished with

resiniferatoxin (RTX), a naturally occurring potent TRPV1 agonist. RTX treatment results in the

specific ablation of TRPV1 neurons (post-development), while sparing other TRPV1 expressing

(non-neuronal) cells. RTX treatment in mice consisted of three injections of RTX given 24 hours

apart. The injections are given in the flank and were: 30ug/kg on day 1, 70ug/kg on day 2 and

100ug/kg on day 3. Just prior to the first RTX injection, mice are treated with 0.1 mg/kg of

buprenorphine, given intraperitoneally, as the first injection causes pain. Subsequent injections

do not require buprenorphine. It takes 4 weeks for TRPV1 nerve ablation to occur. Control mice

received injections with vehicle solution (DMSO with Tween80 in PBS). The tail flick test was

used to confirm nerve ablation.

Tail flick test. Prior to testing, mice are acclimated to the room for 30 minutes. A water bath is

pre-warmed to 52□C and a thermometer is used to constantly monitor the temperature of the

water. The mice are picked up using a fresh paper towel so that only their tail is outside the paper

towel. Once the mouse is relaxed in the tester's hand, the mouse's tail is lowered into the water

so that a quarter of the tail is submerged. The time for the tail to be flicked out of the water is

recorded. If no response has occurred by 20 seconds, the mouse's tail is removed from the water

to prevent tissue damage. A response time latency greater than 10 seconds signifies denervation.

Tumor implantation. Ketamine (87.5 mg/kg)/xylazine (10 mg/kg) were used to anesthetize

mice prior to oral cavity tumor implantation. Tumors were initiated as follows: using a 25-gauge

needle, cells (5×10^4 cells in a total of 50 μl) were implanted orthotopically in the oral cavity of

male C57BL/6 mice. Specifically, the needle was placed proximal to the crease of the mouse

cheek, inserted at approximately a 45° angle, bevel deep, and tumor cells injected into this

location of the cheek pouch. Mice were allowed to recover from anesthesia on a heating blanket

and returned to their home cage. Control (non-tumor) mice were treated the same way only their

injection contained media alone (no cells). Tumor growth was monitored weekly by caliper

measurements of isoflurane anesthetized animals and tumor volume calculated was using the

formula: ½ (length*(width^2)).

Drug treatments. Carprofen (Pivetal) was provided in the drinking water (10mg/kg which is

equivalent to 0.067 mg/ml) which was refreshed every 7 days. Buprenorphine-ER (Fidelis

Animal Health), a 72-hour extended-release formulation, was given by subcutaneous injection

(3.25 mg/kg) every 72 hours. Vehicle-treated animals were treated with subcutaneous injection

of saline. Drug treatments commenced on day 7 post-tumor implantation in all groups. Given

that buprenorphine treatment requires a subcutaneous injection every 3 days, mice in the other

groups (vehicle and carprofen) received a subcutaneous injection of saline at the same volume

(60 µl) as the buprenorphine-treated animals. In this way, all animal stress from handling and

injection was the same.

Nest building. Nest building is an innate behavior performed in rodents of both sexes and is a

general indication of well-being [32]; it also measures the motivation to perform a goal-directed

task. To assess nesting behavior, individually housed mice were given a nesting square (Ancare)

overnight. The following morning, nests were scored by two independent scorers who were

blinded to the conditions [33,34]. The scoring system is 0-4 and based on the percentage of the

nestlet that is shredded and the height of the nest. Two baseline measurements were conducted

prior to tumor implantation followed by weekly testing. Exclusion criteria of an average baseline

score of 3 or greater was used to ensure all mice were good nest builders at baseline (n=3

C57BL/6 mice and n=7 TRPV1^{cre}::DTA$^{fl/wt}$ mice were excluded for this reason).

Cookie test: The cookie test is a variation of the sucrose preference test, a test of anhedonia (the

inability to experience enjoyment or interest in previously rewarding activities). These test the

animal's endogenous inclination for sweet tastes and the notion that it derives pleasure from

consuming sweets. One hour prior to testing, the mice were fasted and acclimated to the brightly

lit testing room. Following the acclimation period, mice were given a piece of cookie

(approximately 1.5 gm). The time it took the mouse to bite the cookie from its placement in the

cage was measured to determine time-to-interact. Each mouse was acclimated to the cookie

every other day for 2 weeks prior to tumor implantation with a baseline measurement followed

by weekly testing. Mice were individually housed. One mouse from the C57BL/6 group was

excluded due to incomplete data (mouse died before completion of the study).

Voluntary wheel running: Prior to tumor implantation, mice were singly housed with a running

wheel maintained in the home cage continuously. The only time that the wheel was removed was

when mice were undergoing nesting. The mice were acclimated to the wheel for 2 weeks prior to

tumor implantation. This acclimatization period provided the time to stabilize their running

performance. Running wheels were maintained in the home cage for the duration of the study.

As mice are most active during the dark phase, nighttime running (8pm to 8am) was assessed.

Running data were collected in 1-minute bins continuously throughout the duration of the

experiment.

Food disappearance. Animals were provided 100 g of solid food per cage each week. At the

end of each week, the remaining food was weighed, and this value was used to calculate the

amount of food that disappeared in that week per cage. For all behavioral experiments, mice

were singly housed.

Neural tracing of orthotopic HNSCC tumors. When oral tumors reached approximately 5 x 5

mm in size, neural tracer was injected into the tumor as described below. Ketamine (87.5

mg/kg)/xylazine (10 mg/kg) were used to anesthetize the mice. A 10 µL Hamilton syringe with

457 30-G needle was loaded with 1% Wheat Germ Agglutinin (WGA) conjugated to either

458 AlexaFluor 594 (Invitrogen) or CF568 (Biotium) in PBS and 2μl was slowly injected intra-

459 tumorally with the bevel side up. The needle was inserted approximately midway through the

460 tumor then pulled back slightly to reduce pressure and leakage of tracer following injection. The

461 injection of tracer occurred slowly over the course of 10 minutes. After the tracer was injected,

462 the needle was kept in place for 2 additional minutes before being slowly removed. Mice were

463 placed on a heating pad until recovery from anesthesia. Five to seven days later, mice were

464 deeply anesthetized and transcardially perfused with ice cold PBS followed by 4%

465 paraformaldehyde and trigeminal ganglia (TGM) were carefully removed and placed in HBSS in

466 a 24-well plate kept on ice. TGM harvesting was completed as follows: euthanized animals were

467 subjected to a midline incision while in the prone position to expose the crown of the skull. The

468 brainstem was separated from the spinal cord by a transverse cut and the top of the skull was

469 removed, exposing the brainstem and TGM. All tissues (tumor, ganglia and brain) were fixed,

470 sectioned, and imaged for WGA labeled nerve fibers and somas under confocal microscopy.

471 WGA injection into control (non-tumor bearing animals) was performed the same way and

472 injection was into the same oral region where tumors were implanted. Volumes of 2 or 10 μl of

473 WGA were used.

475 **Immunohistochemical (IHC) staining.** Tissues were formalin fixed, paraffin-embedded and cut

476 into 5 µm sections. The BenchMark® XT automated slide staining system (Ventana Medical

477 Systems, Inc.) was used to optimize antibody dilutions and staining. The Ventana CC1 solution

478 was used to perform the antigen retrieval step (basic pH tris base buffer). Tissues were incubated

in primary antibody for 1 hour. The Ventana iView DAB detection kit was used as the

chromogen and the slides were counterstained with hematoxylin.

Antibody utilized for immunohistochemistry (IHC). Anti-β-III Tubulin (Abcam, Cat#

ab78078, 1:250, RRID: AB 2256751), Anti-TRPV1 (Alomone Labs Cat# ACC-030, 1:400,

RRID:AB_2313819).

Antibodies utilized for immunofluorescence (IF). cFos (Cell Signaling, Cat# 2250, 1:10,000,

RRID: AB 2247211), ΔFosB (Abcam, Ab11959, 1:5,000, RRID:AB_298732). The following

secondary antibodies were used: Alexa568 anti-rabbit (Thermo, Cat# A-11011, 1:500,

RRID:AB_143157), Alexa488 anti-mouse (Thermo, Cat# A-11001, 1:500, RRID:AB_2534069).

Antibodies utilized for western blot. β-actin (Sigma Life Science, Cat#A2228, 1:1000, RRID:

AB_476697), Tau (Abcam, Cat# ab75714, 1:500, RRID:AB_1310734), phosphorylated TRPV1

(Thermo Fisher Scientific, Cat# PA5-64860, 1:500, RRID:AB_2663797), Sigma-1R

(ProteinTech, Cat# 15168-1-AP, 1:1,000, RRID:AB_2301712), Doublecortin (DCX) (Santa

cruz, Cat# sc-271390, 1:1000, RRID:AB_10610966).

Brain immunostaining and analysis. Brains were post-fixed for twenty-four hours. Coronal or

sagittal sections were cut at 40 µm using a vibratome. Sequential fluorescent immunolabeling

was performed for cFos or ΔFosB on free-floating sections. All sections were blocked with 10%

goat serum in 0.3% Tx-100. All antibodies were diluted in PBS containing 0.03% Triton X-100

and 2% normal goat serum. Sections were incubated in Fos antibody for 72 hours. Following

washes in 0.1M phosphate buffered saline, sections were incubated in secondary antibody for 3

hours. Following additional washes, sections were mounted onto glass slides, and cover slipped

with cytoseal prior to imaging. The total numbers of cFos or ΔFosB-labeled nuclei were counted

on 10 × 2D images acquired using a laser-scanning confocal microscope (Nikon A1R) and

verified as positive if the signal filled the nucleus and stood out clearly compared to surrounding

tissue [35]. Nuclei were counterstained with DAPI. Quantification of Fos-labeled cells was

performed using digital thresholding of Fos-immunoreactive nuclei. The threshold for detection

was set at a level where dark Fos-immunoreactive nuclei were counted, but nuclei with light

labeling, similar to background staining, were not. For each animal, 3-4 sections were selected at

120 μm intervals. ImageJ cell counter (v. 1.52) was used to quantify Fos+ nuclei per region /

section.

Analysis of transcriptional changes in tumor-infiltrating neurons. To assess transcriptional

changes in tumor-infiltrating neurons we utilize a qPCR array (ScienCell, GeneQuery Neural

Transmission and Membrane Trafficking, #MGK008). These qPCR-ready 96 well plates enable

rapid profiling of 88 key genes important for neuronal functions (listed in Table S1). Mice

bearing MOC2-7 oral tumors were injected with tracer (as described above); TGM ganglia were

isolated on day 28 post-tumor implantation, their RNA harvested, converted into cDNA and then

assayed with the array as per manufacturer's instructions (ScienCell, GeneQuery arrays). We

utilized N=4 TGM/group from n=4 mice/group; n=4 plates/condition. Control RNA was isolated

from TGM ganglia from age-matched non-tumor bearing mice (n=4 mice). Relative gene

expression was calculated as per manufacturer's recommendations.

Western blot analysis of whole tumor lysate. Tumors were excised from euthanized mice,

taking care to eliminate as much non-tumor tissue as possible. Tumors were then placed in

approximately 500µl lysis buffer (50mM Tris HCl pH 7.4, 100mM NaCl, 100mM NaF, 10mM

NaPPi, 2mM Na_3VO_4, 10% glycerol, HALT protease inhibitor cocktail) with 1% TX-100 and

kept on ice for 10 minutes. The homogenate was sonicated 3x for 15 secs and then incubated on

ice for 15 minutes. Samples were then centrifuged at 2000 g for 5 minutes at 4°C, supernatant

was collected and further centrifuged at 12,000 g for 10 minutes at 4°C. Protein concentrations

of the supernatants were determined by BCA protein assay.

Western blot analysis of ganglia lysate. Trigeminal ganglia were harvested as described above.

Ganglia were lysed using a homogenizer in lysis buffer at 4°C. Homogenates were centrifuged at

10,000 g for 20 minutes and protein concentrations determined from the supernatant. Protein

concentration was measured by BCA protein assay (Pierce, Cat#23225) and 40 ug of total

protein were separated by SDS-PAGE, transferred to PDVF membranes which were then

blocked for 30 minutes at room temperature (RT) in 5% milk in PBS. Membranes were

incubated with primary antibody overnight at 4°C. Following 1xTBST washes, membranes were

probed with an HRP-conjugated secondary antibody (1:10,000 dilution) for 1 hour at RT,

washed and imaged on a Li-COR Odyssey FC imaging system. Densitometric quantification was

used to assess changes in protein expression.

Western blot densitometry analysis. Raw images of western blots were analyzed using ImageJ.

Briefly, images were opened and bands of interested were selected by gating with the rectangle

selection tool. Densitometry was measured based on grey scale analysis of selected area.

Analysis was conducted on proteins of interest as well as the loading control band (β-actin).

Relative expression of proteins of interest was determined as a fraction of β-actin densitometry.

Neuron culture. Trigeminal ganglia were harvested from MOC2-7 tumor-bearing or non-tumor

animals and enzyme digested in papain, and then collagenase II/dispase at 37 □ for 15 min. After

washing and trituration, cells were plated onto a thin layer of Matrigel® in glass bottom dishes

and cultured with HamsF12 supplemented with 10% FBS. The cells were maintained in an

incubator (5% CO_2, 37 °C) for 24 h before they were used for calcium imaging experiments.

Ca2+ imaging. TGM neurons from non-tumor and tumor-bearing animals (n=4-6

mice/condition) were imaged on the same day. Neurons were incubated with the calcium

indicator, Fluo-4AM, at 37°C for 20 min. After dye loading, the cells were washed, and Live

Cell Imaging Solution (Thermo-Fisher) with 20 mM glucose was added. Calcium imaging was

conducted at room temperature. Changes in intracellular Ca^{2+} were measured using a Nikon

scanning confocal microscope with a□10x objective. Fluo-4AM was excited at 488 nm using an

argon laser with intensity attenuated to 1%. The fluorescence images were acquired in the

confocal frame (1024□×□1024 pixels) scan mode. After 1 min of baseline measure, capsaicin

(300nM final concentration) was added. Ca^{2+} images were recorded before, during and after

capsaicin application. Image acquisition and analysis were achieved using NIS-Elements

imaging software. Fluo-4AM responses were standardized and shown as percent change from the

initial frame. Data are presented as the relative change in fluorescence ($\Delta F/F_0$), where F_0 is the

basal fluorescence and $\Delta F=F-F_0$ with F being the measured intensity recorded during the

experiment. Calcium responses were analyzed only for neurons responding to ionomycin (10

µM, positive control) to ensure neuronal health. Treatment with the cell permeable Ca^{2+} chelator,

BAPTA (200 µM), served as a negative control.

Stereotaxic AAV injection. Male mice were anesthetized with 2% isoflurane and placed in a

stereotaxic head frame on a heating pad. A midline incision was made down the scalp and a

craniotomy was made using a micro drill. A 10-µl Hamilton syringe was used to infuse 1 µl of

AAV1/Syn-GCaMP6f-WPRESV40 (titer 4.65×10^{13} GC per ml, via Addgene) into the

parabrachial nucleus (−5.3 mm anteroposterior, -1.3 mm mediolateral, −3 mm dorsoventral) via

a microsyringe pump. After infusion, the needle was kept at the injection site for 5 min and then

slowly withdrawn. Two weeks following stereotaxic surgeries, tumors were introduced by

implanting MOC2-7 cells orthotopically into the oral cavity as described above.

Ex vivo **Ca2+ imaging of brain slices.** Two weeks post-tumor (or sham) implantation, wildtype

and tumor-bearing mice (n=3 mice/group) were anesthetized under isoflurane and perfused

intracardially with 10 ml of ice-cold N-methyl-d-glucamine (NMDG) solution [92 mM NMDG,

30 mM $NaHCO_3$, 25 mM glucose, 20 mM Hepes, 10 mM $MgSO_4$, 5 mM sodium ascorbate, 3

mM sodium pyruvate, 2.5 mM KCl, 2 mM thiourea, 1.25 mM NaH_2PO_4, and 0.5 mM $CaCl_2$ (pH

7.3, 300 mOsm, bubbled with 95% O2 and 5% CO_2)] [36]. The brains were quickly removed and

placed into additional ice-cold NMDG solution for slicing. Coronal slices (150µm) were cut

using a vibratome (n=3 slices/brain). Slices were transferred to Hepes holding solution and

warmed to 37°C (bubbled with 95% O_2 and 5% CO_2) for 1 hour. After incubation, slices were

transferred to the recording chamber with RT (22° to 25°C) recording solution. A miniaturized

microscope (Miniscope V4) imaged the GCaMP6f signal from the slices. Each video was

processed with motion correction and ΔF/F calculation. Regions of interest (ROIs; considered as

a single cell soma) were manually selected. ROIs that exhibited short bursts of ΔF/F changes or

fluctuations during the recording were analyzed. The ΔF/F changes were then aligned with the

time window of treatment. Movement correction was performed using the motion correction

module in the EZcalcium toolbox employed in MATLAB [37]. The fluorescence intensity trace of

each neuron was extracted, and ΔF/F was calculated (CNMF).

Statistical analysis. GraphPad Prism (version 10.0.3, 2023) was utilized for all statistical

analyses.

Gene expression: For qRT-PCR analysis, Ct values for each gene were normalized to that of the

reference gene. Statistical analysis by multiple student's t-test.

Ca^{2+} imaging: Statistical analysis by unpaired student's t-test. Fluo-4AM/GCaMP6f responses

were standardized and shown as percent change from the initial frame. Data are presented as the

relative change in fluorescence ($\Delta F/F_0$), where F_0 is the basal fluorescence and $\Delta F = F - F_0$ with F

being the peak response.

Western blot: Membranes were visualized, and proteins were quantified using the Odyssey

infrared imaging system and software (Li-COR). Densitometric quantification of western blots

was performed by normalizing the signal from tumor or non-tumor ganglia to the β-actin

(loading control) signal and differences assessed by one-way ANOVA with post-hoc Tukey test.

Fos brain immunostaining: The numbers of Fos (cFos, ΔFosB) immune-positive neurons in brain

sections from tumor-bearing or non-tumor animals were quantified as described above and

statistically analyzed by 2-way ANOVA with multiple comparisons.

Nesting scores: The variations over time of the nesting scores for each group were statistically analyzed by repeated measures ANOVA. Exclusion criteria of an average baseline score of 3 or greater was used to ensure all mice were good nest builders.

Time to interact scores (cookie test): Statistical analysis by repeated measures ANOVA.

Voluntary wheel running: The data were collected in 1-minute bins continuously. When compiling the data, the sum of bins between 8pm to 8am is taken for each individual animal for voluntary nightly running. Baseline is calculated by taking the final 3 days prior to tumor implantation and averaging the values. The percent change is then calculated for each day from baseline for each mouse. The data were then analyzed using a repeated measures 2-way ANOVA.

Food disappearance: Statistical analysis by repeated measures ANOVA.

Weight: Statistical analysis by repeated measures ANOVA.

Tumor growth curves: Statistical analysis of tumor growth curves by repeated measures ANOVA

Kaplan Meier survival: Statistical analysis of survival by Log-rank (Mantel-Cox) test.

References

1   Carreira, H. *et al.* Associations Between Breast Cancer Survivorship and Adverse Mental Health Outcomes: A Systematic Review. *Journal of the National Cancer Institute* **110**, 1311-1327 (2018). https://doi.org:10.1093/jnci/djy177

2   Pitman, A., Suleman, S., Hyde, N. & Hodgkiss, A. Depression and anxiety in patients
with cancer. *Bmj* **361**, k1415 (2018). https://doi.org:10.1136/bmj.k1415
3   Walker, J. *et al.* Prevalence of depression in adults with cancer: a systematic review.
Annals of oncology : official journal of the European Society for Medical Oncology **24**,
895-900 (2013). https://doi.org:10.1093/annonc/mds575
4   Zhu, J. *et al.* First-onset mental disorders after cancer diagnosis and cancer-specific
mortality: a nationwide cohort study. *Annals of oncology : official journal of the*
European Society for Medical Oncology **28**, 1964-1969 (2017).
https://doi.org:10.1093/annonc/mdx265
5   Steel, Z. *et al.* The global prevalence of common mental disorders: a systematic review
and meta-analysis 1980-2013. *Int J Epidemiol* **43**, 476-493 (2014).
https://doi.org:10.1093/ije/dyu038
6   Gotze, H. *et al.* Depression and anxiety in long-term survivors 5 and 10 years after cancer
diagnosis. *Supportive care in cancer : official journal of the Multinational Association of*
Supportive Care in Cancer **28**, 211-220 (2020). https://doi.org:10.1007/s00520-019-
04805-1
7   Kuba, K. *et al.* Risk for depression and anxiety in long-term survivors of hematologic
cancer. *Health Psychol* **38**, 187-195 (2019). https://doi.org:10.1037/hea0000713
8   Wang, Y. H. *et al.* Depression and anxiety in relation to cancer incidence and mortality: a
systematic review and meta-analysis of cohort studies. *Mol Psychiatry* **25**, 1487-1499
(2020). https://doi.org:10.1038/s41380-019-0595-x
9   Naser, A. Y. *et al.* Depression and Anxiety in Patients With Cancer: A Cross-Sectional
Study. *Front Psychol* **12**, 585534 (2021). https://doi.org:10.3389/fpsyg.2021.585534
10   Tsaras, K. *et al.* Assessment of Depression and Anxiety in Breast Cancer Patients:
Prevalence and Associated Factors. *Asian Pacific journal of cancer prevention : APJCP*
19, 1661-1669 (2018). https://doi.org:10.22034/APJCP.2018.19.6.1661
11   Madeo, M. *et al.* Cancer exosomes induce tumor innervation. *Nature communications* **9**,
4284 (2018). https://doi.org:10.1038/s41467-018-06640-0
12   Lucido, C. T. *et al.* Innervation of cervical carcinoma is mediated by cancer-derived
exosomes. *Gynecologic oncology* (2019). https://doi.org:10.1016/j.ygyno.2019.04.651
13   Barr, J. L. *et al.* Intra-Tumoral Nerve-Tracing in a Novel Syngeneic Model of High-
Grade Serous Ovarian Carcinoma. *Cells* **10** (2021).
https://doi.org:10.3390/cells10123491
14   Renz, B. W. *et al.* beta2 Adrenergic-Neurotrophin Feedforward Loop Promotes
Pancreatic Cancer. *Cancer cell* **33**, 75-90 e77 (2018).
https://doi.org:10.1016/j.ccell.2017.11.007
15   Hayakawa, Y. *et al.* Nerve Growth Factor Promotes Gastric Tumorigenesis through
Aberrant Cholinergic Signaling. *Cancer cell* **31**, 21-34 (2017).
https://doi.org:10.1016/j.ccell.2016.11.005
16   Wang, K. *et al.* The neurotrophic factor neurturin contributes toward an aggressive
cancer cell phenotype, neuropathic pain and neuronal plasticity in pancreatic cancer.
Carcinogenesis **35**, 103-113 (2014). https://doi.org:10.1093/carcin/bgt312
17   Kasprzak, A. & Adamek, A. The Neuropeptide System and Colorectal Cancer Liver
Metastases: Mechanisms and Management. *International journal of molecular sciences*
21 (2020). https://doi.org:10.3390/ijms21103494

688	18	Silverman, D. A. *et al.* Cancer-Associated Neurogenesis and Nerve-Cancer Cross-talk.
689		*Cancer research* **81**, 1431-1440 (2021). https://doi.org:10.1158/0008-5472.CAN-20-2793
690	19	Cole, S. W., Nagaraja, A. S., Lutgendorf, S. K., Green, P. A. & Sood, A. K. Sympathetic
691		nervous system regulation of the tumour microenvironment. *Nature reviews. Cancer* **15**,
692		563-572 (2015). https://doi.org:10.1038/nrc3978
693	20	Magnon, C. *et al.* Autonomic nerve development contributes to prostate cancer
694		progression. *Science* **341**, 1236361 (2013). https://doi.org:10.1126/science.1236361
695	21	Sadighparvar, S., Darband, S. G., Ghaderi-Pakdel, F., Mihanfar, A. & Majidinia, M.
696		Parasympathetic, but not sympathetic denervation, suppressed colorectal cancer
697		progression. *Eur J Pharmacol* **913**, 174626 (2021).
698		https://doi.org:10.1016/j.ejphar.2021.174626
699	22	Chen, D. & Ayala, G. E. Innervating Prostate Cancer. *The New England journal of*
700		*medicine* **378**, 675-677 (2018). https://doi.org:10.1056/NEJMcibr1714003
701	23	Kamiya, A. *et al.* Genetic manipulation of autonomic nerve fiber innervation and activity
702		and its effect on breast cancer progression. *Nature neuroscience* **22**, 1289-1305 (2019).
703		https://doi.org:10.1038/s41593-019-0430-3
704	24	Chen, M. *et al.* The Rostral Ventromedial and Lateral Medulla Are the Major Areas
705		Responsive to Lung Cancer Progression among Brainstem Lung-Innervating Nuclei.
706		*Brain Sci* **12** (2022). https://doi.org:10.3390/brainsci12111486
707	25	Henry, M. *et al.* Psychosocial outcomes of human papillomavirus (HPV)- and non-HPV-
708		related head and neck cancers: A longitudinal study. *Psychooncology* **31**, 185-197 (2022).
709		https://doi.org:10.1002/pon.5803
710	26	Virgen, C. G. *et al.* Pharmacological management of cancer pain: Novel therapeutics.
711		*Biomed Pharmacother* **156**, 113871 (2022). https://doi.org:10.1016/j.biopha.2022.113871
712	27	Hjermstad, M. J., Fainsinger, R., Kaasa, S. & European Palliative Care Research, C.
713		Assessment and classification of cancer pain. *Curr Opin Support Palliat Care* **3**, 24-30
714		(2009). https://doi.org:10.1097/SPC.0b013e3283260644
715	28	van den Beuken-van Everdingen, M. H. *et al.* Prevalence of pain in patients with cancer:
716		a systematic review of the past 40 years. *Annals of oncology : official journal of the*
717		*European Society for Medical Oncology* **18**, 1437-1449 (2007).
718		https://doi.org:10.1093/annonc/mdm056
719	29	Judd, N. P. *et al.* ERK1/2 regulation of CD44 modulates oral cancer aggressiveness.
720		*Cancer research* **72**, 365-374 (2012). https://doi.org:10.1158/0008-5472.CAN-11-1831
721	30	Onken, M. D. *et al.* A surprising cross-species conservation in the genomic landscape of
722		mouse and human oral cancer identifies a transcriptional signature predicting metastatic
723		disease. *Clinical cancer research : an official journal of the American Association for*
724		*Cancer Research* **20**, 2873-2884 (2014). https://doi.org:10.1158/1078-0432.CCR-14-
725		0205
726	31	Cavanaugh, D. J. *et al.* Trpv1 reporter mice reveal highly restricted brain distribution and
727		functional expression in arteriolar smooth muscle cells. *The Journal of neuroscience : the*
728		*official journal of the Society for Neuroscience* **31**, 5067-5077 (2011).
729		https://doi.org:10.1523/JNEUROSCI.6451-10.2011
730	32	Neely, C. L. C., Pedemonte, K. A., Boggs, K. N. & Flinn, J. M. Nest Building Behavior
731		as an Early Indicator of Behavioral Deficits in Mice. *Journal of visualized experiments :*
732		*JoVE* (2019). https://doi.org:10.3791/60139

33 Deacon, R. M. Assessing nest building in mice. *Nature protocols* **1**, 1117-1119 (2006).
https://doi.org:10.1038/nprot.2006.170
34 Kraeuter, A. K., Guest, P. C. & Sarnyai, Z. The Nest Building Test in Mice for
Assessment of General Well-Being. *Methods in molecular biology* **1916**, 87-91 (2019).
https://doi.org:10.1007/978-1-4939-8994-2_7
35 Kee, N., Teixeira, C. M., Wang, A. H. & Frankland, P. W. Imaging activation of adult-
generated granule cells in spatial memory. *Nature protocols* **2**, 3033-3044 (2007).
https://doi.org:10.1038/nprot.2007.415
36 Ting, J. T. *et al.* Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-
glucamine Protective Recovery Method. *Journal of visualized experiments : JoVE*
(2018). https://doi.org:10.3791/53825
37 Cantu, D. A. *et al.* EZcalcium: Open-Source Toolbox for Analysis of Calcium Imaging
Data. *Front Neural Circuits* **14**, 25 (2020). https://doi.org:10.3389/fncir.2020.00025
38 Restaino, A. C. *et al.* Functional neuronal circuits promote disease progression in cancer.
Sci Adv **9**, eade4443 (2023). https://doi.org:10.1126/sciadv.ade4443
39 Menezes, F. D. S., Fernandes, G. A., Antunes, J. L. F., Villa, L. L. & Toporcov, T. N.
Global incidence trends in head and neck cancer for HPV-related and -unrelated subsites:
A systematic review of population-based studies. *Oral oncology* **115**, 105177 (2021).
https://doi.org:10.1016/j.oraloncology.2020.105177
40 Georgieff, I. S. *et al.* Expression of high molecular weight tau in the central and
peripheral nervous systems. *Journal of cell science* **105 (Pt 3)**, 729-737 (1993).
https://doi.org:10.1242/jcs.105.3.729
41 Dellarole, A. & Grilli, M. Adult dorsal root ganglia sensory neurons express the early
neuronal fate marker doublecortin. *J Comp Neurol* **511**, 318-328 (2008).
https://doi.org:10.1002/cne.21845
42 Bernreuther, C., Salein, N., Matschke, J. & Hagel, C. Expression of doublecortin in
tumours of the central and peripheral nervous system and in human non-neuronal tissues.
Acta Neuropathol **111**, 247-254 (2006). https://doi.org:10.1007/s00401-006-0038-z
43 Borges, L. F. & Sidman, R. L. Axonal transport of lectins in the peripheral nervous
system. *The Journal of neuroscience : the official journal of the Society for Neuroscience*
2, 647-653 (1982).
44 van der Want, J. J., Klooster, J., Cardozo, B. N., de Weerd, H. & Liem, R. S. Tract-
tracing in the nervous system of vertebrates using horseradish peroxidase and its
conjugates: tracers, chromogens and stabilization for light and electron microscopy.
Brain Res Brain Res Protoc **1**, 269-279 (1997). https://doi.org:10.1016/s1385-
299x(96)00042-6
45 McNicholas, K. & Michael, M. Z. Immuno-characterization of Exosomes Using
Nanoparticle Tracking Analysis. *Methods in molecular biology* **1545**, 35-42 (2017).
https://doi.org:10.1007/978-1-4939-6728-5_3
46 Dumas, M., Schwab, M. E. & Thoenen, H. Retrograde axonal transport of specific
macromolecules as a tool for characterizing nerve terminal membranes. *Journal of*
neurobiology **10**, 179-197 (1979). https://doi.org:10.1002/neu.480100207
47 Sillitoe, R. V. Mossy Fibers Terminate Directly Within Purkinje Cell Zones During
Mouse Development. *Cerebellum* **15**, 14-17 (2016). https://doi.org:10.1007/s12311-015-
0712-6

Tabuchi, K. *et al.* GAL4/UAS-WGA system as a powerful tool for tracing Drosophila transsynaptic neural pathways. *Journal of neuroscience research* **59**, 94-99 (2000).

Levy, S. L., White, J. J., Lackey, E. P., Schwartz, L. & Sillitoe, R. V. WGA-Alexa Conjugates for Axonal Tracing. *Current protocols in neuroscience* **79**, 1 28 21-21 28 24 (2017). https://doi.org:10.1002/cpns.28

Itaya, S. K. Anterograde transsynaptic transport of WGA-HRP in rat olfactory pathways. *Brain research* **409**, 205-214 (1987). https://doi.org:10.1016/0006-8993(87)90703-7

Itaya, S. K., Van Hoesen, G. W. & Barnes, C. L. Anterograde transsynaptic transport of WGA-HRP in the limbic system of rat and monkey. *Brain research* **398**, 397-402 (1986). https://doi.org:10.1016/0006-8993(86)91504-0

Carson, K. A. & Mesulam, M. M. Ultrastructural evidence in mice that transganglionically transported horseradish peroxidase-wheat germ agglutinin conjugate reaches the intraspinal terminations of sensory neurons. *Neuroscience letters* **29**, 201-206 (1982). https://doi.org:10.1016/0304-3940(82)90317-2

Yeh, L. F., Watanabe, M., Sulkes-Cuevas, J. & Johansen, J. P. Dysregulation of aversive signaling pathways: a novel circuit endophenotype for pain and anxiety disorders. *Curr Opin Neurobiol* **48**, 37-44 (2018). https://doi.org:10.1016/j.conb.2017.09.006

Hu, H. Reward and Aversion. *Annual review of neuroscience* **39**, 297-324 (2016). https://doi.org:10.1146/annurev-neuro-070815-014106

Zhu, Y. B. *et al.* PBN-PVT projections modulate negative affective states in mice. *eLife* **11** (2022). https://doi.org:10.7554/eLife.68372

Russo, S. J. & Nestler, E. J. The brain reward circuitry in mood disorders. *Nature reviews. Neuroscience* **14**, 609-625 (2013). https://doi.org:10.1038/nrn3381

Balood, M. *et al.* Nociceptor neurons affect cancer immunosurveillance. *Nature* (2022). https://doi.org:10.1038/s41586-022-05374-w

Michetti, F. *et al.* The S100B story: from biomarker to active factor in neural injury. *Journal of neurochemistry* **148**, 168-187 (2019). https://doi.org:10.1111/jnc.14574

Liu, Q. *et al.* Transcriptional Alterations of Mouse Trigeminal Ganglion Neurons Following Orofacial Inflammation Revealed by Single-Cell Analysis. *Frontiers in cellular neuroscience* **16**, 885569 (2022). https://doi.org:10.3389/fncel.2022.885569

Malin, S. A., Davis, B. M. & Molliver, D. C. Production of dissociated sensory neuron cultures and considerations for their use in studying neuronal function and plasticity. *Nature protocols* **2**, 152-160 (2007). https://doi.org:10.1038/nprot.2006.461

Ortiz-Renteria, M. *et al.* TRPV1 channels and the progesterone receptor Sig-1R interact to regulate pain. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E1657-E1666 (2018). https://doi.org:10.1073/pnas.1715972115

Bhave, G. *et al.* Protein kinase C phosphorylation sensitizes but does not activate the capsaicin receptor transient receptor potential vanilloid 1 (TRPV1). *Proceedings of the National Academy of Sciences of the United States of America* **100**, 12480-12485 (2003). https://doi.org:10.1073/pnas.2032100100

Numazaki, M. *et al.* Structural determinant of TRPV1 desensitization interacts with calmodulin. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 8002-8006 (2003). https://doi.org:10.1073/pnas.1337252100

Numazaki, M., Tominaga, T., Toyooka, H. & Tominaga, M. Direct phosphorylation of capsaicin receptor VR1 by protein kinase Cepsilon and identification of two target serine

residues. *The Journal of biological chemistry* **277**, 13375-13378 (2002).
https://doi.org:10.1074/jbc.C200104200

65   Bhave, G. *et al.* cAMP-dependent protein kinase regulates desensitization of the
capsaicin receptor (VR1) by direct phosphorylation. *Neuron* **35**, 721-731 (2002).
https://doi.org:10.1016/s0896-6273(02)00802-4

66   Scheff, N. N. *et al.* Oral cancer induced TRPV1 sensitization is mediated by PAR(2)
signaling in primary afferent neurons innervating the cancer microenvironment. *Scientific*
reports **12**, 4121 (2022). https://doi.org:10.1038/s41598-022-08005-6

67   Chung, L. A Brief Introduction to the Transduction of Neural Activity into Fos Signal.
Dev Reprod **19**, 61-67 (2015). https://doi.org:10.12717/DR.2015.19.2.061

68   Hudson, A. E. Genetic Reporters of Neuronal Activity: c-Fos and G-CaMP6. *Methods in*
enzymology **603**, 197-220 (2018). https://doi.org:10.1016/bs.mie.2018.01.023

69   Nestler, E. J., Barrot, M. & Self, D. W. DeltaFosB: a sustained molecular switch for
addiction. *Proceedings of the National Academy of Sciences of the United States of*
America **98**, 11042-11046 (2001). https://doi.org:10.1073/pnas.191352698

70   Nestler, E. J., Kelz, M. B. & Chen, J. DeltaFosB: a molecular mediator of long-term
neural and behavioral plasticity. *Brain research* **835**, 10-17 (1999).
https://doi.org:10.1016/s0006-8993(98)01191-3

71   Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity.
Nature **499**, 295-300 (2013). https://doi.org:10.1038/nature12354

72   Niedzwiedz, C. L., Knifton, L., Robb, K. A., Katikireddi, S. V. & Smith, D. J. Depression
and anxiety among people living with and beyond cancer: a growing clinical and research
priority. *BMC cancer* **19**, 943 (2019). https://doi.org:10.1186/s12885-019-6181-4

73   Utley, M., Adeyanju, T., Bernardo, B., Paskett, E. D. & Krok-Schoen, J. L. The
association between mental health, social support and physical health outcomes among
older female cancer survivors. *J Geriatr Oncol* **13**, 834-838 (2022).
https://doi.org:10.1016/j.jgo.2022.04.001

74   Ji, X. *et al.* National estimates of mental health service use and unmet needs among adult
cancer survivors. *Cancer* **127**, 4296-4305 (2021). https://doi.org:10.1002/cncr.33827

75   Mishra, S. K., Tisel, S. M., Orestes, P., Bhangoo, S. K. & Hoon, M. A. TRPV1-lineage
neurons are required for thermal sensation. *The EMBO journal* **30**, 582-593 (2011).
https://doi.org:10.1038/emboj.2010.325

76   Liu, M. Y. *et al.* Sucrose preference test for measurement of stress-induced anhedonia in
mice. *Nature protocols* **13**, 1686-1698 (2018). https://doi.org:10.1038/s41596-018-0011-
z

77   Eliwa, H. *et al.* Adult neurogenesis augmentation attenuates anhedonia and HPA axis
dysregulation in a mouse model of chronic stress and depression.
Psychoneuroendocrinology **124**, 105097 (2021).
https://doi.org:10.1016/j.psyneuen.2020.105097

78   Norden, D. M. *et al.* Tumor growth increases neuroinflammation, fatigue and depressive-
like behavior prior to alterations in muscle function. *Brain, behavior, and immunity* **43**,
76-85 (2015). https://doi.org:10.1016/j.bbi.2014.07.013

79   Zombeck, J. A., Fey, E. G., Lyng, G. D. & Sonis, S. T. A clinically translatable mouse
model for chemotherapy-related fatigue. *Comp Med* **63**, 491-497 (2013).

80   Grossberg, A. J. *et al.* Tumor-Associated Fatigue in Cancer Patients Develops
Independently of IL1 Signaling. *Cancer research* **78**, 695-705 (2018).
https://doi.org:10.1158/0008-5472.CAN-17-2168
81   Bower, J. E. *et al.* Inflammation and behavioral symptoms after breast cancer treatment:
do fatigue, depression, and sleep disturbance share a common underlying mechanism?
Journal of clinical oncology : official journal of the American Society of Clinical
Oncology **29**, 3517-3522 (2011). https://doi.org:10.1200/JCO.2011.36.1154
82   Norden, D. M. *et al.* Fluoxetine prevents the development of depressive-like behavior in
a mouse model of cancer related fatigue. *Physiol Behav* **140**, 230-235 (2015).
https://doi.org:10.1016/j.physbeh.2014.12.045
83   Chen, A. M. *et al.* Depression among long-term survivors of head and neck cancer treated
with radiation therapy. *JAMA Otolaryngol Head Neck Surg* **139**, 885-889 (2013).
https://doi.org:10.1001/jamaoto.2013.4072
84   Lazure, K. E., Lydiatt, W. M., Denman, D. & Burke, W. J. Association between
depression and survival or disease recurrence in patients with head and neck cancer
enrolled in a depression prevention trial. *Head & neck* **31**, 888-892 (2009).
https://doi.org:10.1002/hed.21046
85   Elaldi, R. *et al.* Correlations between long-term quality of life and patient needs and
concerns following head and neck cancer treatment and the impact of psychological
distress. A multicentric cross-sectional study. *European archives of oto-rhino-*
laryngology : official journal of the European Federation of Oto-Rhino-Laryngological
Societies **278**, 2437-2445 (2021). https://doi.org:10.1007/s00405-020-06326-8
86   Navratilova, E. *et al.* Positive emotions and brain reward circuits in chronic pain. *J Comp*
Neurol **524**, 1646-1652 (2016). https://doi.org:10.1002/cne.23968
87   Mercer Lindsay, N., Chen, C., Gilam, G., Mackey, S. & Scherrer, G. Brain circuits for
pain and its treatment. *Science translational medicine* **13**, eabj7360 (2021).
https://doi.org:10.1126/scitranslmed.abj7360
88   Backes, T. M. *et al.* Stimulation of TRPV1 channels activates the AP-1 transcription
factor. *Biochemical pharmacology* **150**, 160-169 (2018).
https://doi.org:10.1016/j.bcp.2018.02.008
89   Boedtkjer, E. & Pedersen, S. F. The Acidic Tumor Microenvironment as a Driver of
Cancer. *Annual review of physiology* (2019). https://doi.org:10.1146/annurev-physiol-
021119-034627
90   Pocock, R. & Hobert, O. Oxygen levels affect axon guidance and neuronal migration in
Caenorhabditis elegans. *Nature neuroscience* **11**, 894-900 (2008).
https://doi.org:10.1038/nn.2152
91   Pocock, R. & Hobert, O. Hypoxia activates a latent circuit for processing gustatory
information in C. elegans. *Nature neuroscience* **13**, 610-614 (2010).
https://doi.org:10.1038/nn.2537
92   Stevenson, T. J. *et al.* Hypoxia disruption of vertebrate CNS pathfinding through
ephrinB2 Is rescued by magnesium. *PLoS Genet* **8**, e1002638 (2012).
https://doi.org:10.1371/journal.pgen.1002638
93   Amit, M. *et al.* Loss of p53 drives neuron reprogramming in head and neck cancer.
Nature **578**, 449-454 (2020). https://doi.org:10.1038/s41586-020-1996-3

Ueda, H. R. *et al.* Whole-Brain Profiling of Cells and Circuits in Mammals by Tissue Clearing and Light-Sheet Microscopy. *Neuron* **106**, 369-387 (2020). https://doi.org:10.1016/j.neuron.2020.03.004

Kobayashi, K. *et al.* Distinct expression of TRPM8, TRPA1, and TRPV1 mRNAs in rat primary afferent neurons with adelta/c-fibers and colocalization with trk receptors. *J Comp Neurol* **493**, 596-606 (2005). https://doi.org:10.1002/cne.20794

Mickle, A. D., Shepherd, A. J. & Mohapatra, D. P. Sensory TRP channels: the key transducers of nociception and pain. *Prog Mol Biol Transl Sci* **131**, 73-118 (2015). https://doi.org:10.1016/bs.pmbts.2015.01.002

Renna, M. E. *et al.* Depression and anxiety in colorectal cancer patients: Ties to pain, fatigue, and inflammation. *Psychooncology* **31**, 1536-1544 (2022). https://doi.org:10.1002/pon.5986

Walker, A. K., Kavelaars, A., Heijnen, C. J. & Dantzer, R. Neuroinflammation and comorbidity of pain and depression. *Pharmacological reviews* **66**, 80-101 (2014). https://doi.org:10.1124/pr.113.008144

Grossberg, A. J. *et al.* Interleukin 6-independent metabolic reprogramming as a driver of cancer-related fatigue. *Brain, behavior, and immunity* **88**, 230-241 (2020). https://doi.org:10.1016/j.bbi.2020.05.043

Thaker, P. H. *et al.* Chronic stress promotes tumor growth and angiogenesis in a mouse model of ovarian carcinoma. *Nature medicine* **12**, 939-944 (2006). https://doi.org:10.1038/nm1447

Kim-Fuchs, C. *et al.* Chronic stress accelerates pancreatic cancer growth and invasion: a critical role for beta-adrenergic signaling in the pancreatic microenvironment. *Brain, behavior, and immunity* **40**, 40-47 (2014). https://doi.org:10.1016/j.bbi.2014.02.019

Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018). https://doi.org:10.1016/j.neuron.2018.10.013

Takayanagi-Kiya, S. & Kiya, T. Activity-dependent visualization and control of neural circuits for courtship behavior in the fly Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 5715-5720 (2019). https://doi.org:10.1073/pnas.1814628116

Acknowledgments: We thank the Histology and Imaging Core (Sanford Research, supported by National Institutes of Health, National Institute of General Medical Sciences, Center of Biomedical Research Excellence 5P20GM103548 and P30GM145398), specifically Claire Evans who provided her services and expertise towards this project. We also thank the following funding sources for their critical contributions to this work: National Institutes of Health, National Institute of Dental and Craniofacial Research grant 1R01DE032712-01 (PDV),

Institutional Development Award (IDeA) from the National Institute of General Medical

Sciences of the National Institutes of Health grant 5P20GM103548 (PDV). DaCCoTA Scholar

Award supported by the National Institute of General Medical Sciences of the National Institutes

of Health grant U54 GM128729 (PDV). National Institute of Health, National Cancer Institute

grant R37 1R37CA242006-01A1 (MA). Stiefel family Discovery award (MA). Institutional

Research Grant (MA). Disruptive Science Moonshot award, MDACC (MA). Canadian Institutes

of Health Research grants 162211, 461274, 461275 (ST). National Institutes of Health (R01

CA193522; R21 NS130712) (RD).

Materials & Correspondence: Correspondence and material requests should be addressed to

Paola D. Vermeer: Paola.Vermeer@sanfordhealth.org

Data availability statement: Data will be available upon reasonable request.
961
962
963

Fig 1. Tumor-infiltrating nerves form a circuit connecting to the brain. (**A**) Western blot of

MOC2-7 whole tumor lysates for Tau (**A**) and Doublecortin, DCX (**B**). Tumors were harvested

at different time points (as indicated) post-tumor implantation (n=3 mice/time point). Mouse

brain, positive control; MOC2-7, whole cell lysate control; β-actin, loading control.

Immunohistochemical staining for β-III tubulin (β3T, brown, arrows) of tumors harvested on

days 4 (**C**), 10 (**D**) and 20 (**E**) post tumor implantation. Scale bar, 100 µm. Dotted boxes are

shown at higher magnification below each panel; scale bar, 50µm. Arrows highlight positive

nerves staining. Representative *en face* confocal image of WGA positive (red) neurites within

tumor (**F**) and the ipsilateral trigeminal ganglion (**G**) from a mouse orthotopically implanted with

a MOC2-7 tumor and intra-tumorally injected with the tracer. Scale bars, 200µm. (**H**) Low

magnification section of ipsilateral and contralateral TGM ganglia from a MOC2-7 tumor-

bearing mouse that was intra-tumorally injected with WGA. The V3 branch of the TGM is

marked with an arrow. WGA (red) positive neurons found only in the ipsilateral TGM. Scale bar,

500 µm. (**I**) Compressed Z-stack of confocal images demonstrating restricted WGA (red) within

the tumor bed (cytokeratin, green). Vertical and horizontal cross-sections highlight that WGA

(red) localizes strictly within the tumor (green) and does not leak out. Yellow shows the merged

file. (**J**) Diagram showing the location of the trigeminal (TGM) ganglion, extension of its

neurites into the tumor bed and the existing circuit which includes the spinal nucleus of the TGM

(SpVc), the parabrachial nucleus (PBN) and the central amygdala (CeA) which get labeled

following WGA injection into tumor. (**K**) Representative confocal images of the SpVc, PBN and

CeA following injection of 2 µl of WGA into an oral MOC2-7 tumor (n=10 mice).

Representative confocal images of SpVc, PBN and CeA following injection of (**L**) 10 µl (n=3

mice) of WGA into the oral cavity of a non-tumor bearing control animal. Scale bar, 50 μm.

WGA (red looks pink due to DaPi), DaPi nuclear counterstain (blue).

Fig 2. Tumor-infiltrating neurons become transcriptionally and functionally altered.

Trigeminal (TGM) ganglia from tumor bearing (red) or non-tumor bearing (blue) mice were

analyzed by (**A**) quantitative PCR array. N=4 TGM/group from n=4 mice/group; n=4

arrays/group. Ct values for each gene normalized to that of a housekeeping gene. Statistical

analysis by multiple students t-test. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$, ****, $p<0.00001$. Data

are expressed as means ± standard deviation. (**B**) Representative fluorescent images of

dissociated neurons from the ipsilateral or contralateral TGM ganglia from a MOC2-7 tumor-

bearing mouse. N=6 of each ganglia harvested from 6 tumor-bearing animals with n= 4-8

neurons analyzed/ganglia. The color palette reflects the strength of the calcium signal. Images

are taken pre- and post-treatment with capsaicin. (**C**) Graph of average change in fluorescence

for all neurons analyzed in each group (ipsilateral and contralateral). Statistical analysis by

student's t-test. Arrow, time of capsaicin (300 nM) stimulation. Graph of average peak of the

curve (**D**) and area under the curve (**E**) from panel C. Statistical analysis by student's t-test. **,

$p<0.01$. (**F**) Quantification of Ca^{+2} responses to capsaicin (300 nM) from non-tumoral (Control)

and tumor-infiltrating, tracer$^+$ (Tumor-infiltrating) neurons. N= 4-8 neurons/group from n=4

control and n=7 tumor-bearing mice. Statistical analysis by paired students t-test, *, $p<0.05$.

Data expressed as means ± SEM. (**G**) Representative western blot of trigeminal ganglia from

non-tumor (N) or MOC2-7 tumor-bearing (T) animals for phosphorylated TRPV1 (pTRPV1,

Ser502, Ser800), σ1 receptor (σ1R) and β-actin (loading control). Densitometric quantification

of western blots for pTRPV1 (**H**) and σ1R (**I**). Statistical analysis by student's t-test. *, $p< 0.05$.

Data are expressed as means□±□SEM. (n=3-4 mice/group).

1010

1011

Fig 3. Elevated CNS neuronal activity in tumor bearing mice. Representative fluorescent

photomicrographs of the spinal nucleus of the TGM (SpVc) (**A**), parabrachial nucleus (PBN) (**B**)

and central amygdala (CeA) (**C**) from a control (-tumor) and MOC2-7 tumor-bearing (+tumor)

mouse brains immuonfluorescently stained for cFos (red) or ΔFosB (green). Nuclei

counterstained with DaPi (blue). Dotted circle denotes each brain region. Scale bar, 100 μm.

Brain landmarks: Spc, superior cerebellar peduncle; VL, lateral ventricle; IA, intercalated

amygdalar nucleus. Quantification of cFos and ΔFosB staining in SpVc (**D**), PBN (**E**) and CeA

(**F**) from control (-tumor) and tumor-bearing (+tumor) mice. N= 4 mice/group with n=2-4

sections analyzed/brain region/mouse/group. Data are expressed as the mean ± SEM. Statistical

analysis by one-way ANOVA*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ns, not significant. Scale bar,

100 μm. Representative heat maps of regions of interest (ROI), each of which represent one

neuron, within an *ex vivo* brain slice from non-tumor (control) (**G**) and tumor-bearing (**H**) mice.

The color palette reflects the strength of the calcium signal. Arrow denotes the time of KCl (30

mM) application. N= 3 mice/group with n=3 slices/brain. Quantification of calcium imaging data

from (n= 5 mice/group with n=2-4 sections analyzed/mouse) analyzing the peak of the curve (**I**)

and the area under the curve (**J**). Statistical analysis by student's t-test. ****, $p< 0.0001$; ***, $p<$

001.

1029

Fig 4. Intra-tumoral neurons impact behavior. (**A**) Bright field photomicrograph of TGM

ganglion from C57BL/6 or TRPV1[cre]::DTA[fl/wt] mice IHC stained for TRPV1 (brown). Scale bar,

100μm. Quantification of cFos and ΔFosB from the brains of C57Bl/6 (C57) or

TRPV1[cre]::DTA[fl/wt] (DTA) animals with MOC2-7 oral tumors in the Spinal nucleus of the

trigeminal (SpVc) (**B**), the parabrachial nucleus (PBN) (**C**) and the central amygdala (CeA) (**D**).

N=4 mice/group with n=2-4 sections analyzed/brain region/mouse. Statistical analysis by one-

way ANOVA. **, $p<0.01$; ***, $p<0.001$. Data displayed as mean $\pm$ SEM. C57BL/6 (black,

n=15) and TRPV1^{cre}::DTA$^{fl/wt}$ (blue, n=14) mice were orthotopically implanted with MOC2-7

tumor and behavior assessed weekly and statistically analyzed by repeated measures ANOVA.

(**E**) Graph of nesting scores over time. There is a main effect of time such that there is a decline

in nesting performance over time. We also see a time by strain interaction in that C57BL/6 mice

show a greater decline than the nociceptor neuron ablated TRPV1^{cre}::DTA$^{fl/wt}$ mice. Post hoc

testing shows a significant difference in nesting on day 27 ($p=0.0315$). (**F**) Graph of time to

interact with the cookie in the cookie test. There is a main effect of time such that the time to

interact with the cookie declines over time. There is a main effect of strain such that nociceptor

neuron ablated TRPV1^{cre}::DTA$^{fl/wt}$ mice were faster to interact with the cookie. Finally, there is a

time by strain interaction such that nociceptor neuron ablated TRPV1^{cre}::DTA$^{fl/wt}$ mice showed

progressively faster task performance, while in C57BL/6 mice interaction time plateaued. Post

hoc testing demonstrates a significant difference between the groups on days 3, 22, and 27

($p<0.05$). (**G**) Tumor growth curves for mice that underwent behavioral testing. Statistical

analysis by repeated measures ANOVA. There is a main effect of time such that there is an

increase in tumor volume over time. We also see a time by strain interaction in that C57BL/6

mice have larger tumors than nociceptor ablated TRPV1^{cre}::DTA$^{fl/wt}$ mice. (**H**) Kaplan-Meier

survival curve for mice in panel I. Statistical analysis by Log-rank (Mantel-Cox) test. ****,

$p<0.0001$. (**I**) Graph of % change in weight from baseline. Dotted line represents baseline.

Statistical analysis by repeated measures ANOVA. There was a main effect of time such that

both groups showed an initial increase in weight followed by a decrease. Post hoc testing shows

a significant difference between the groups on days 5, 10, and 20. (**J**) Graph of % change from

baseline in food consumption. Dotted line represents baseline. Statistical analysis by repeated

measures ANOVA. There is a main effect of time such that there is a decline in % food

disappearance over time. There is a time by strain interaction such that C57BL/6 mice show a

greater decline in % food disappearance compared to nociceptor neuron ablated

TRPV1^{cre}::DTA$^{fl/wt}$ mice. In all panels, error bars are SEM.

1063
Fig 5. Denervation of tumors improves behavior, attenuates body weight loss, and reduces

tumor growth. (**A**) Graph of nesting scores over time. Statistical analysis by repeated measures

ANOVA (group x time): The time factor and its interaction with the group factor are significant

with a significant difference between RTX- and vehicle-treated mice on the last time point

(p<0.001). (**B**) Graph of % change in weight from baseline. Statistical analysis by repeated

measures ANOVA (group x time): The time factor and its interaction with the group factor are

significant with a significant difference between groups on the last time point (p<0.01). (**C**)

Tumor volume according to treatment and time: Statistical analysis by repeated measures

ANOVA (treatment x time): The time factor and its interaction with the group factor are

significant with a significant difference between treatments on the last two time points

(p<0.001). N=20 mice/group.

1075
Fig 6. Treating for pain does not restore all behaviors. C57BL/6 male mice were

orthotopically implanted with MOC2-7 tumors and separated into three groups: carprofen treated

(10mg/kg, blue, n=10), buprenorphine treated (3.25mg/kg, red, n=10) or vehicle treated (black,

n=9). Mice underwent behavioral testing. All analyses are by repeated measures ANOVA. (**A**)

Data are graphed as % of baseline for overnight wheel running. The dotted line represents the

baseline. There is a main effect of time such that there is a decline in voluntary wheel running

over time. We also see a time by drug (carprofen) interaction in that vehicle-treated mice show a

greater decline in wheel running than carprofen-treated mice. Post hoc testing shows a significant

difference in wheel running (p<0.0001). (**B**) There is a main effect of time such that there is a

decline in voluntary wheel running over time. We also see a time by drug (buprenorphine)

interaction in that vehicle-treated mice show a greater decline in wheel running than the

buprenorphine-treated mice. Post hoc testing shows a significant difference in wheel running

(p<0.0001). (**C**) Graph of nesting scores over time. When comparing vehicle vs carprofen

groups, there is a main effect of time such that there is a decline in nesting performance over

time. We also see a time by drug (carprofen) interaction in that vehicle-treated mice show a

greater decline than carprofen-treated mice. Similarly, we see a main effect of time when

comparing the vehicle and buprenorphine groups. There is a time by drug (buprenorphine)

interaction in that vehicle-treated mice show a greater decline than buprenorphine-treated mice.

(**D**) Graph of % change in weight from baseline. The dotted line represents the baseline. There

was a main effect of time. There is also a drug (carprofen) by time interaction in that vehicle-

treated mice show a greater decline in weight than the carprofen-treated mice. In fact, carprofen

treated mice gain weight. Similarly, when comparing the vehicle and buprenorphine groups,

there is a main effect of time. There is also an interaction of drug (buprenorphine) by time in that

vehicle-treated mice show a greater decline in weight than the buprenorphine-treated mice. In

fact, the buprenorphine treated mice gain weight. When carprofen and buprenorphine-treated

animals are compared, there is an interaction of time with treatment such that buprenorphine

treated animals gain more weight than carprofen-treated animals. (**E**) Graph of % change in food

disappearance from baseline. When comparing the vehicle vs carprofen group, there is a main

effect of time such that there is a decline in food disappearance over time. We also see a time by

treatment interaction such that carprofen-treated animals do not decline in food disappearance as

much as vehicle treated animals. When comparing the vehicle vs buprenorphine group, there is a

main effect of time. In the buprenorphine treated group, there is a time by treatment interaction

such that the buprenorphine-treated animals demonstrate the smallest decline in food

disappearance. Finally, comparison of the carprofen and buprenorphine groups shows that there

is an interaction between time and treatment such that buprenorphine-treated animals show the

least reduction in food disappearance. (**F**) Tumor volume was monitored weekly for mice in all

groups. There is a main effect of treatment such that mice treated with carprofen show a greater

decline in tumor volume than buprenorphine-treated mice.

Significant effects of statistics in Supplementary Table S3.

1115

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6