The genetic consequences of range expansion and its influence on diploidization in polyploids

William W. Booker[1*] and Daniel R. Schrider[1]

[1]Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27514-2916, United States of America

* Corresponding author

E-mail: wbooker14@gmail.com

Keywords: Polyploid, population genetics, expansion load, allele surfing, diploidization, range expansion

Despite newly formed polyploids being subjected to myriad fitness consequences, the relative prevalence of polyploidy both contemporarily and in ancestral branches of the tree of life suggests alternative advantages that outweigh these consequences. One proposed advantage is that polyploids have an elevated adaptive potential that enables them to colonize novel habitats such as previously glaciated areas. However, previous research conducted in diploids suggests that range expansion comes with a fitness cost as deleterious mutations may fix rapidly on the expansion front. Here, we interrogate the potential consequences of expansion in polyploids by conducting spatially explicit forward-in-time simulations of autopolyploids, allopolyploids, and diploids to investigate how ploidy and inheritance patterns impact the relative ability of polyploids to expand their range. We show that under realistic dominance models, autopolyploids suffer greater fitness reductions than diploids as a result of range expansion due to the fixation of increased mutational load that is masked in the range core. Alternatively, the disomic inheritance of allopolyploids provides a shield to this fixation resulting in minimal fitness consequences under an empirically estimated DFE. In light of this advantage provided by disomy, we investigate how range expansion may influence cytogenetic diploidization through the reversion to disomy in autotetraploids. We show that under both a model of where the mode of inheritance is determined by a small number of loci and a model where inheritance is regulated by chromosomal similarity, disomy evolves more rapidly on the expansion front than in the range core, and that this dynamic inheritance model has additional effects on fitness. Together our results point to a complex interaction between dominance, ploidy, inheritance, and recombination on fitness as a population spreads across a geographic range.

INTRODUCTION

Polyploidy as a mechanism of evolution has had a profound influence on shaping the tree of life (Otto and Whitton 2000; Gregory and Mable 2005; Albertin and Marullo 2012; Barker et al. 2016; Zhan et al. 2016; Van de Peer et al. 2017; Li et al. 2018; One Thousand Plant Transcriptomes Initiative 2019). However, In many ways this observation does not reconcile with the known negative fitness consequences imbued by polyploidization. Once formed, polyploids often suffer meiotic abnormalities and genomic instability that result in reduced fertility and offspring viability as well as phenotypic instability (Mayer and Aguilera 1990; Comai et al. 2000; Ramsey and Schemske 2002; Morgan et al. 2020). Should a polyploid organism get past these hurdles, the minority cytotype exclusion principle suggests they have a very low probability of reproductive success by virtue of being surrounded by their more numerous non-polyploid relatives (Levin 1975; Husband 2000). Newly formed polyploids are also likely to be ecologically similar to their progenitor species, introducing additional pressures through the competition for resources (Rodriguez 1996; Yamauchi et al. 2004).

Despite these disadvantages, the prevalence of polyploid species suggests there are inherent advantages that polyploidy provides. One prevailing theory is that polyploidy provides an advantage in times of stress (Van de Peer et al. 2021), and they may more rapidly expand into novel or deglaciated habitats (Brochmann et al. 2004; Comai 2005; David 2022). The precise mechanisms that provide polyploids with these advantages aren't entirely clear. One hypothesis proposes that polyploids simply have a greater adaptive potential resultant from a greater effective genic mutation rate (Otto 2007) and that duplicate genes allow for more specialized functionalization of one or both copies (Lynch and Conery 2000; Lynch and Force 2000; Adams and Wendel 2005; Gout and Lynch 2015). Similarly, more rapid adaptation may come from a multiplication of the gene regulatory network, as whole genome duplication creates a level of redundancy that frees parts of the network to reconfigure (Freeling and Thomas 2006; Hegarty and Hiscock 2007; Fusco et al. 2010; Yao et al. 2019), which can be particularly beneficial in times of environmental change (Ebadi et al. 2023). Alternatively, glacial cycles may play a more direct role in driving rates of polyploid formation by initiating secondary contact of diploid species (Stebbins 1984) potentially creating allopolyploids with fixed heterozygosity (Stebbins 1985; Brochmann et al. 2004).

The adaptive advantages provided by polyploidy for colonization of new habitats are not uncontroversial. Firstly, polyploids are not always more robust in harsher conditions and can have more restricted ranges (Levin 2002; Parisod et al. 2010). Although polyploids have a greater effective mutation rate that can produce novel adaptive mutations, this process also means polyploids will carry a greater mutational load that must be purged

(Otto 2007). Additionally, the idea that mutations on duplicate genes are neutral because the non-mutated copy retains its function is not necessarily true (Mable and Otto 2001). Perhaps most glaring is the immediate reduction in effective population size following polyploidization (Stebbins 1950), limiting the variation upon which selection can act for any adaptation.

Though there is a long history of discussion on the adaptive advantages polyploids may have in expanding into new environments, little work has been done to understand the genetic consequences of those expansions. As a species expands its range, populations along the expansion front are colonized by a small number of individuals that are more likely to colonize subsequent populations (Moreau et al. 2011). This phenomenon has a compounding effect, increasing an allele's probability of fixation due to the elevated force of drift along the expansion front (Peischl et al. 2013; Peischl et al. 2015). This effect is particularly pronounced for recessive weakly deleterious mutations which can be masked from the effects of selection in larger core populations (Peischl and Excoffier 2015). As a result, populations on the edge of an expansion should have reduced fitness compared to populations in the core. Although the primary consequence of this process is reduced fitness, range expansions can alter the landscape of neutral genetic diversity across the genome (Schlichta et al. 2022) and can promote the hybridization of species coming into secondary contact (MacPherson et al. 2022).

The ability for polyploids to mask deleterious alleles through whole genome duplication is often invoked to describe a relative advantage of polyploidy. However, theoretical work has shown that this masking results in a higher mutational load in polyploids (Haldane 1932; Hill 1970; Bever and Felber 1992) resulting in greater inbreeding depression in small populations and self-fertilizers (Bennett 1976). Because the mutational load present prior to expansion becomes a primary driver of expansion load (Peischl and Excoffier 2015), polyploids may be more severely impacted by the consequences of expanding their range. The effects of range expansion are also heavily influenced by recombination (Peischl et al. 2015)–a force that acts qualitatively differently during diploid and tetraploid meiosis (Grandont et al. 2013; Stenberg and Saura 2013), altering the ability of expanding migrants to purge the accumulating load. In total, these observations suggest that range expansion should have an inherently different effect on polyploid genomes.

Just as all polyploids must expand their range to ensure successful establishment, the ultimate fate of all polyploids is returning into a functionally diploid state (Wolfe 2001; Conant et al. 2014; Robertson et al. 2017; Mandáková and Lysak 2018; Li et al. 2021). While diploidization may refer to the process of fractionation (sometimes referred to as genic diploidization), where multiple copies of genes are generally lost or silenced (Freeling et al. 2012), diploidization first occurs cytogenetically to restore normal meiotic

behavior. Because multivalent formation often results in meiotic errors, and therefore has large deleterious fitness effects, bivalent pairing is often restored relatively quickly (Feldman and Levy 2012; Feldman et al. 2012; Tayalé and Parisod 2013; Morgan et al. 2021). However, the fitness consequences of tetrasomic inheritance itself are less clear, and the restoration of disomy is often prolonged over millions of years while lineages continue to diversify (Lien et al. 2016; Scott et al. 2016; Robertson et al. 2017; Parey et al. 2022; Redmond et al. 2022). Because expansion load is driven by the fixation of deleterious recessive mutations, the evolution of disomy might then provide a fitness advantage to recently formed species by preventing the fixation of alleles across subgenomes.

Here, we explore both the genetic consequences of range expansion in polyploid species, and the role that these consequences play in driving cytogenetic diploidization and the evolution of disomy. Specifically, we develop a framework for spatially explicit forward-in-time simulations of autotetraploid, allotetraploid, and diploid species, simulating the expansion of species out of their core range. We investigate expansion load as it relates to the pattern of inheritance, ploidy, and dominance to better understand how the range expansion of polyploids contributes to their fitness in comparison to diploids. In addition, we consider how selection for disomic inheritance may be influenced by expansion load by developing two models of diploidization and investigating the parameters in which diploidization is favored or disfavored in a geographically expanding species. Ultimately, we aim to identify the scenarios in which polyploidy and diploidization provide advantages or disadvantages for expansion and therefore contribute to their probability of establishment in novel regions.

METHODS

Simulation of Tetraploids

All simulations were conducted in SLiM version 4.0.1 (Haller and Messer 2023). Although SLiM does not support polyploid species natively, modeling polyploids is possible through some modifications. In general, these modifications were to the behavior of recombination and segregation and to fitness calculations in polyploid individuals.

To modify recombination for tetraploids, we added an additional population to SLiM that was reproductively isolated from all other populations and that served as storage for the additional two chromosomes present in each tetraploid individual. During initialization, each individual in the storage population is assigned, or "tagged", to an individual in the focal population. For both auto- and allopolyploids we assume all chromosomes form bivalents and there is no double reduction. During reproduction, for autopolyploids we model tetrasomic inheritance by choosing, for a given parent, two random

chromosomes from the four total chromosomes in the focal and storage populations to recombine. These two recombinant chromosomes are then passed on to a new offspring individual with one chromosome randomly chosen to be passed to the offspring individual in the focal population, and the other to the corresponding member of the storage population that is tagged to the offspring individual. This same process is repeated for the other parent of this individual. For allopolyploids, disomic inheritance was modeled by generating recombinants of the focal and storage chromosomes separately, thereby ensuring the two subgenomes remained distinct.

To calculate fitness for tetraploids, first the count of derived alleles across all four chromosomes was obtained. This allele count was then multiplied by the product of the selection and dominance coefficients. Although diploids only have a single dominance coefficient given a single heterozygous state, tetraploids have three coefficients, corresponding to the possible heterozygote copy numbers of the derived allele (one, two, or three). As such, the vector of dominance coefficients depends on the model of dominance under consideration. All dominance coefficient vectors, excluding those obtained from the empirical distribution of fitness effects (DFE) model (described below), can be found in Table 1. To determine the effect of expansion by itself, all fitnesses were rescaled to one by dividing each population's fitness by the fitness of the outermost burn-in population at the start of expansion.

Demographic Model

We used a stepping stone model where the total population consisted of a series of demes along a single dimension, and migration occurred between adjacent demes. Prior to expansion, demes at both edges were reflecting, with no migrants expanding beyond either edge. During expansion, reflection at one edge was removed, and migration beyond this deme resulted in the colonization of an additional deme.

Within each deme, we allowed population size to grow according to the logistic growth model defined in Eq. 1:

Eq. 1 $\quad N' \; = \dfrac{e^r N_t}{1 + \dfrac{(e^r - 1)N_t}{K}}$

Here, N_t is the number of individuals at time t, and N' is the rate parameter of a poisson distribution representing the number of individuals at time $t + 1$. K is the carrying capacity of the deme and r is the intrinsic growth rate. For all simulations, generations are discrete and non-overlapping. At the start of each generation, individuals are killed as a function of their fitness, and reproduction immediately follows this step. Individuals are dioecious, and mates are chosen at random with replacement until N_{t+1} ~

Poisson(*N'*) offspring are generated. Individual fitness has no effect on reproductive success among survivors of the death phase. Following reproduction, individuals migrated to adjacent populations at rate *m*, with the direction of migration chosen at random, provided individuals were not in the first deme, or the fifth deme during burn-in.

<u>Model Parameters</u>

To start, 5 demes were initialized at carrying capacity and allowed to evolve for a burn-in of 2500 generations, and migration was limited to between adjacent demes among these 5 initial demes only. Although 2500 generations represents 5N generations (5 demes at N=100) which is shorter than the time required for all alleles to coalesce, we limited the burn-in to 2500 generations because modeling all or a majority of the mutations as deleterious eventually leads to a population crash under some parameters for certain models. Following this burn-in, individuals were allowed to migrate into new demes from the edge deme, and at this point migration was allowed between all pairs of adjacent demes. For all simulations, the carrying capacity *K* was set to 100, *r*=log(2), *m*=0.05. For the genomic parameters, each individual had a 1 Mb genome, and deleterious mutations occurred at a rate of 2.5×10^{-8} for the fixed-s model and 5×10^{-8} for our empirical DFE model. For all simulations the recombination rate (ρ) was set equal to the deleterious mutation rate.

Fitness was calculated multiplicatively as shown in Eq. 2, where *w* is the individual's fitness and *s* is the selection coefficient for the given mutation *i*, and h_j is the dominance coefficient for mutation *i* given that it is present at copy number *j* in the individual under consideration:

Eq. 2 $$w = \prod_{i}^{n}(1 + h_j * s_i)$$

For all simulations with our fixed-*s* model, *s* was set to -0.005. For simulations based on an empirically estimated DFE, *s* was distributed as described below. To calculate the effect of mutations outside of the homozygous state, we multiplied *s* by the dominance coefficient for the individual's copy number of the derived allele as specified by the dominance model being simulated (Table 1). Here, the additive model describes a DFE where *s* is scaled linearly by the proportion of alleles in the mutated state. The recessive model describes a DFE where there is no selective effect as long as one wild-type allele is present at the locus, and the duplex model describes a DFE where selection is dosage dependent and acts only when the wild-type allele is lower than 50% in frequency (see Hill 1970).

Table 1. Dominance coefficients (*h*) for diploids and tetraploids for each model and allelic state.

Model	Diploid *h* vector (h_0,h_1,h_2)	Tetraploid *h* vector (h_0,h_1,h_2,h_3,h_4)
Additive	(0.0, 0.5, 1.0)	(0.0, 0.25, 0.50, 0.75, 1.0)
Recessive	(0.0, 0.0, 1.0)	(0.0, 0.0, 0.0, 0.0, 1.0)
Recessive (duplex)	(0.0, 0.0, 1.0)	(0.0, 0.0, 0.0, 1.0, 1.0)

<u>Empirical DFE</u>

In an attempt to model the effects of expansion load on more realistic populations, we modeled an expanding population experiencing mutations whose fitness effects were drawn from an empirically estimated DFE. To do so, we used the genome wide *Arabidopsis lyrata* DFE from Huber et al. (2018). In addition to having a distribution of selection coefficients, the model from Huber et al. (2018) also modeled the relationship between dominance coefficient *h* and *s* (Eq. 3). Here, $\theta_{intercept}$ defines the value of *h* at s = 0, and θ_{rate} defines the rate at which *h* approaches 0 with a declining negative *s*. To broadly summarize the distributions of *s* and *h* defined by this model, all mutations are deleterious, with most being weakly deleterious, and there is a negative relationship between *h* and *s*. For our simulations, the DFE was parameterized with *s* as a gamma distribution with α = 0.16 and β = 0.0092, and *h* was parameterized with $\theta_{intercept}$ = 0.978 θ_{rate} = 50328.

Eq. 3 $h = \dfrac{1}{\frac{1}{\theta_{intercept}} - \theta_{rate} s}$

To translate the dominance coefficient from the diploid DFE to tetraploids, we translated the flux model of dominance from Kascer and Burns (1981) by using the estimated *h* from Eq. 3 as the point on the flux curve at 0.5 wild-type frequency. Eq. 4 is adapted from Boucher et al. (2016) and models the shape of the flux curve given *h*. Here, *W* is the fitness effect for that mutation given the percentage of ancestral or wild-type alleles W^P, and a value *B* that determines the shape of the curve and that is in turn determined by the value of *h* as defined in Eq. 5. Combining the two equations, we can then solve for the fitness effect of any mutation given the *h* value calculated from Eq. 3 and the percentage of ancestral alleles (Eq. 6). Importantly, Eq. 5 and Eq. 6 are not solvable for

$h = 0.5$, however the curve at this value represents the additive model therefore the desired fitness value ($W_{tetraploid}$) in such cases can be trivially obtained from W^P.

Eq. 4 $W = \dfrac{(1+B)W^P}{W^P + B}$

Eq. 5 $B = \dfrac{h}{1-2h}$

Eq. 6 $W_{tetraploid} = \dfrac{(1+\frac{h}{1-2h})W^P}{W^P + \frac{h}{1-2h}}$

<u>Cytogenetic Diploidization</u>

To model the evolution of cytogenetic diploidization in range-expanding populations, we ran our simulations as described previously, but allowed for additional mutations that altered the inheritance pattern. All individuals start as autopolyploids with bivalent pairing but free recombination between all four chromosomes. As diploidization mutations are added, the pairing behavior of chromosomes is altered depending on the model of diploidization being simulated (see below). Importantly, for these mutations we did not assume an infinite sites model, and allowed diploidization mutations to revert to the ancestral state.

Although the mechanisms for diploidization are not entirely clear, evidence suggests two distinct processes are largely responsible (see Li et al. 2021). One process involves mutations that regulate the meiotic machinery which fully or partially restore normal meiotic pairing behavior (Sánchez-Morán et al. 2001; Jenczewski et al. 2003; Liu et al. 2006; Henry et al. 2014; Gonzalo et al. 2019; Morgan et al. 2020). We refer to this as the dominant meiotic model, and we define its parameters in Eq. 7. Here, d is the number of unique derived diploidization alleles in an individual, and λ is the number of mutations required to restore disomic inheritance. Under this model, subgenomes are predefined but all chromosomes freely recombine. As diploidization mutations occur, each mutation has a fixed effect in weighting the probability of recombination between chromosomes of the same subgenome, until $\frac{d}{\lambda} \geq 1$, at which point the subgenomes are differentiated and never recombine.

Eq. 7 $P(disomy) = max(\frac{d}{\lambda}, 1)$

Alternatively, diploidization may occur as a process of chromosomal differentiation, where large rearrangements, duplications, deletions, or other structural changes drive differences between chromosomes such that they are no longer able to pair. Le Comber et al. (2010) showed that drift of neutral mutations resulting in chromosomal differences alongside fitness consequences for pairing failures was sufficient to drive the evolution of diploidization. Because expansion is akin to accelerating the force of drift, it is possible expansion may more rapidly result in the evolution of diploidization.

To model this behavior, we adapted the model from Le Comber et al. (2010) and applied it to our simulations. Here, we allowed for λ diploidization loci randomly distributed across the chromosome to evolve at a rate of μ_{dip} mutations per locus per generation. Diploidization mutations have no fitness effects themselves and only affect the pairing efficiency. At the start of reproduction, the percent sequence identity is calculated for every combination of chromosome pairs. Percent sequence identity is then used to calculate the pairing efficiency of each combination as shown in Eq. 8 adapted from Le Comber et al. (2010). The nature of this equation is such that there is a sharp decline in pairing efficiency at around 90% similarity, with similarity below 80% resulting in poor pairing efficiency (approximately 0) and above 90% having perfect efficiency (approximately 100). Once pairing efficiency is calculated for each combination of chromosomes, the average efficiency for each possible set of pairs (i.e. ab-cd, ac-bd, ad-bc) is calculated, and the set of pairs is chosen by random sample weighted by their pairing efficiency. This process is completed for each parent, and the probability of successful reproduction is calculated based on the average pairing efficiency of the two parents–mimicking errors in meiosis resulting in inviable offspring. Should the offspring not survive, another random set of parents is chosen to reproduce, and the process is repeated until a successful offspring is produced.

To determine if disomy had evolved, we calculated the diploidization index as the difference in pairing efficiency between the top two possible sets of pairs, with the idea that under disomy one possible pair should be at near 100% pairing efficiency and all other pairs at 0%. Because disomy in this model is less specific and there is no pairing associated fitness cost when pairing efficiency is below 90%, we defined a population as disomic if it had an average diploidization index equal to or above 0.9.

Eq. 8 $P(pairing_{ij}) = \dfrac{8}{0.8 + 80e^{78 - (\frac{(\lambda - d_{ij}) \times 100}{\lambda})}}$

RESULTS

Expansion load has a similar effect across tetraploids and diploids for additive models

To examine the consequences of expansion load in polyploids without the effects of dominance, we first modeled populations expanding their range with mutations having an additive effect on fitness relative to their frequency. Under this additive model, the reduction in fitness in the edge of the species range as a result of geographic expansion is remarkably similar across all tetraploid and diploid models, although autotetraploids may have slightly higher fitness along the edge (Fig. 1a; Fig. 2a). Interestingly, allotetraploids have a much worse starting unscaled fitness but experience a linear decline in edge fitness as the expansion proceeds that has a very similar slope to that of autopolyploids and diploids (Supp. Fig. 1). Core fitness declines fastest in allotetraploids and slowest in autotetraploids, with diploids more similar to but ultimately less fit than autotetraploids (Fig. 2b). Allopolyploids have a greater effective mutation rate than diploids, but do not benefit from a greater effective recombination rate–making deleterious mutations appear more rapidly but without any concomitant improvement in the ability to purge them. Autopolyploids also have a greater effective mutation rate, but also recombination between all possible pairs of four chromosomes which reduces the probability that mutations reach high frequency. As such, autotetraploids start with a generally higher number of mutations per individual than diploids (Fig. 2d), but fewer mutations per genome (Fig. 2c), suggesting that many of these mutations are at low frequency. This in turn results in a large fraction of mutations in tetraploids to be lost quickly at the start of expansion before later increasing linearly over time (Fig. 2d).

Figure 1. Mean fitness in each deme at each generation under additive (a), recessive (b), and empirical DFE (c) dominance models. Fitnesses for each deme are scaled by the initial fitness of the core at the onset of the range expansion, and then averaged across all replicates. Note that the scales are different for each dominance model, and that under the empirical DFE model the scale for allotetraploids is different from autotetraploids and diploids.

Figure 2. Statistics summarizing fitness and variation within spatially expanding populations under the additive dominance model. (a) Mean fitness (scaled) of the edge

deme. (b) Mean fitness (scaled) of the innermost core deme. Note the different *y*-axis scales in panels a and b. (c) Mean number of deleterious mutations per genome in the outermost deme (i.e. the edge). (d) Mean number of deleterious mutations per individual in the edge deme. (e) Number of fixed mutations in the edge deme. (f) proportion of the mutational load contributed by mutations originating along the edge deme. Lines show the average value of each statistic across 25 replicates, and dashed lines show ± 1 standard deviation.

The masking effect of tetraploidy in core populations results in rapid fitness decline during expansion under recessive models

We next examined the consequences of expansion load in polyploids under a more realistic model where all deleterious mutations are recessive. Under both the fully recessive and duplex models, where selection acts if mutations are homozygous or have fewer than two copies of the wild-type allele, respectively, fitness in autotetraploids decreases rapidly on the expansion front (Fig. 1b). The magnitude of this fitness decline in the early stages of expansion in autotetraploids is significantly greater than that observed in diploids, although the decline eventually steadies out to a similar slope in both ploidies (Fig. 3a). A further look at the dynamics driving this pattern demonstrate that autopolyploid genomes and individuals have more deleterious mutations at the start of the expansion (Fig. 3c,d) resulting in a much greater number of deleterious mutations that fix along the expansion front (Fig. 3e). In contrast to the edge, however, fitness in the core is maintained at high levels in autotetraploids under a recessive model, whereas diploid fitness steadily decreases (Fig. 3b). Although in a single panmictic population fitness in autotetraploids does decrease, fitness is still considerably greater at all generations in autopolyploids when the recombination rate is equal to the mutation rate (Supp. Fig. 2). When ϱ is increased fitness between tetraploids and diploids becomes more similar, but fitness is still higher in autotetraploids. As such, although autotetraploid fitness in the core will eventually decline, it should be maintained at higher levels than diploids outside of extreme recombination rates.

Figure 3. Statistics summarizing fitness and variation within spatially expanding populations under the recessive model. (a) Mean fitness (scaled) of the edge deme. (b)

Mean fitness (scaled) of the innermost core deme. Note the different y-axis scales in panels a and b. (c) Mean number of deleterious mutations per genome in the edge deme. (d) Mean number of deleterious mutations per individual in the edge deme. (e) Number of fixed mutations in the edge deme. (f) Proportion of the mutational load contributed by mutations originating along the edge deme. Lines show the average value of each statistic across 25 replicates, and dashed lines show ± 1 standard deviation.

Autotetraploids behave similarly under an empirical DFE as in the recessive model

We next examined a DFE that was adapted from one estimated from *Arabidopsis lyrata* (Huber et al. 2018) that models the relationship between the dominance and selection coefficients of deleterious mutations (Methods). Under this empirical DFE model, autotetraploids and diploids behave generally similarly to the fully recessive model (Fig. 1c, Fig. 4a-f). Again, autotetraploids accumulate more deleterious mutations during the burn-in because most mutations in the DFE are weakly deleterious and therefore proliferate in an autotetraploid where they are generally masked from the effects of selection, and therefore, like in the recessive model, exhibit an initial rapid drop prior to linear growth in the number of deleterious mutations. Also similar to the fully recessive model, autotetraploid fitness is largely stable in the core populations under this model, while diploid core fitness slowly declines (Fig. 4b). Although the incomplete recessiveness of the mutations in this model results in some fitness decline in allotetraploids, the magnitude of the fitness effects at 50% or lower frequency (the maximum frequency they can reach in allotetraploids) are miniscule and result in little difference between core and edge populations, and little fitness consequences generally.

Figure 4. Empirical DFE model statistics. (a) Mean fitness (scaled) of the edge deme. (b) Mean fitness (scaled) of the innermost core deme. Note the different *y*-axis scales in panels a and b. (c) Mean number of deleterious mutations per genome in the edge

deme. (d) Mean number of deleterious mutations per individual in the edge deme. (e) Number of fixed mutations in the edge deme. For allopolyploids, fixed mutations are those fixed within each subgenome. (f) proportion of the mutational load contributed by mutations originating along the edge deme. Lines show the average value of each statistic across 25 replicates, and dashed lines show ± 1 standard deviation.

Cytogenetic diploidization is more likely on the expansion front when disomy is restored by dominant mutations at one or two loci

Because recessive deleterious mutations fix rapidly in a spatially expanding autopolyploid with tetrasomic inheritance, we incorporated the ability to evolve disomy into our models to investigate if expansion increases the rate of cytogenetic diploidization in autopolyploids. Under a model in which disomy is restored by dominant mutations, the average rate at which populations diverge from random chromosomal associations (tetrasomy, diploidization index = 0) to complete subgenomic association (disomy, diploidization index = 1) is faster in core populations (Fig. 5a, Supp. Fig. 3). However, the average rate doesn't accurately capture the stochastic dynamics of disomy evolution in expanding populations, where rapid fixation of diploidization alleles along the expansion front can lead to near instantaneous evolution of disomy along the edge. To investigate this, we considered the proportion of simulations where complete disomy (population diploidization index = 1) had evolved in core and edge populations. Here, where 1 or 2 diploidization mutations are required for full disomy to evolve, the proportion of simulations where the entire population is disomic is greater for edge rather than core populations (Fig. 5b).

Evidence that the evolution of disomy in edge populations under this model is largely stochastic comes from the observation that the selection coefficient has little if any effect on the rate of diploidization (Fig. 5a,b). Although our simulations of expansion load under a recessive model show diploidization should be favored, because the selection coefficient has little effect on the edge, the evolution of disomy under this model is likely constrained by the number of mutations required to restore disomy (λ), the mutation rate of diploidization mutations (μ_{dip}), and the migration rate (m). Interestingly, although s has little effect on edge populations, there does appear to be a small increase in the rate of diploidization at higher selective coefficients in core populations (Fig. 5a,b). Because fitness is stable in the core in a recessive model (Fig. 3b), it's unlikely that the fitness of the core itself is driving this process. More likely, this pattern is driven by migration from nearby midrange populations where disomy provides a selective benefit and thus diploidization is more prevalent.

Additional evidence for the stochastic dynamics in these models can be seen investigating individual simulations (Supp. Fig. 4). For non-edge populations, the diploidization index fluctuates and may steadily evolve towards disomy. For edge populations however, changes to diploidization index happen quickly and are generally maintained in that and all subsequently colonized demes. Another interesting phenomenon individual simulation results show is that even if disomy doesn't evolve on the edge more quickly, the stochastic dynamics occurring along the edge can have lasting effects long after those populations have established (e.g. maintained higher diploidization index in additional generations at $\lambda = 5$).

Figure 5. Diploidization of core and edge populations under the dominant meiotic model for polyploids with a fully recessive model of dominance. (a) Average diploidization index of core and edge populations under varying selection coefficients. (b) Proportion of simulations where disomy is evolved (population average diploidization index = 1) under varying selective coefficients for λ of 1 and 2. Dashed lines for (a) represent ± 1 standard deviation. For $\lambda = 1$ and 2 the number of simulations ran was 100 rather than 25 for other models due to the large variance.

Diploidization occurs much more rapidly on the expansion front if pairing behavior is regulated by chromosomal similarity

While mutations of large effect have been previously discovered to restore disomy in several systems, evidence also suggests cytogenetic diploidization may proceed through chromosomal differentiation. Using a model adapted from Le Comber et al. (2010) where chromosomal similarity regulates pairing behavior with additional fitness costs, we find that diploidization is more likely along the expansion front for all parameters investigated (Fig. 6). Unlike the dominant meiotic model, this was true for both the average diploidization index (Fig. 6a) and the proportion of simulations where disomy had evolved (Fig. 6b). Using the original diploidization mutation rate of 1×10^{-3} from Le Comber et al. (2010), no simulations in the core population had evolved disomy by 1000 generations. However, even at s = -0.0025, disomy had evolved along the edge for a significant proportion of simulations, with disomy in edge populations evolving by 1k generations in all simulations with s = -0.01. Doubling the rate of diploidization mutations results in some core simulations restoring disomy within 1000 generations, but edge populations still typically restore disomy much sooner (Fig. 6b).

In contrast to the dominant meiotic model, the selection coefficient of deleterious mutations has a clear effect in the pairing efficiency model for both core and edge populations, with the rate of diploidization increasing as the selection coefficient becomes more negative. Although our simulations of expansion load show a clear selective benefit of evolving disomy early along the edge (Fig. 3), the cause for diploidization in the core is less direct. However, as described in the Methods, the nature of this model is that there is a fitness cost if chromosomes are not either homogeneous in a unimodal distribution (tetrasomic) or in a bimodal distribution of two homogeneous groups (disomic). As such, because the model includes pairing-associated fitness costs (hereafter abbreviated as PAFC) there should be a fitness cost to the formation of chromosome pairs with high nucleotide diversity, thereby increasing the likelihood that a deleterious mutation will be found in the homozygous state. Indeed, in these simulations we observed a reduction in nucleotide diversity at deleterious mutations (Supp. Fig. 5a) and a correlated reduction in fitness (Supp. Fig. 5b) in the core. However this effect is delayed, with an initial increase in nucleotide diversity before the reduction. Because this initial increase follows the trajectory of the model without the PAFC, it is likely this initial increase is due to the preferential pairing increasing nucleotide diversity by initially trending towards a bimodal distribution of diversity prior to hitting any fitness cost when chromosomes become too differentiated.

In edge populations along the expansion front we observed a similar phenomenon as in the core–that pairing-associated fitness costs exacerbate the genetic load incurred from expansion (Supp. Fig. 5b) due to an initial additional loss of nucleotide diversity (Supp.

Fig. 5a). Although the magnitude of this additional load is dependent upon the degree to which pairing affects fitness from Eq. 8, the additional loss incurred when the model incorporates a PAFC significantly affects both the total fitness cost as well as the rate of fitness loss during the initial expansion. However, as expansion continues and disomy is restored, diploidization slows the rate of fitness decline, and in later generations fitness along the range edge is higher under this model of diploidization than without diploidization (Supp. Fig. 5b).

Pairing associated fitness costs generate barriers to gene flow between disomic edge demes and tetrasomic interior demes

We further explored the dynamics of the pairing efficiency model by investigating models without PAFCs at varying selection coefficients including complete neutrality. Interestingly, although the PAFCs are necessary for the evolution of disomy in core populations (Figs. 5 and 6), PAFCs actually decrease the rate at which disomy evolves along the expansion front (Supp. Fig. 6). For all selection coefficients investigated, both the diploidization index and the proportion of simulations with restored disomy are higher in both the core and edge demes when PAFCs are removed (although this state is transient, see below). In core populations, however, the diploidization index increases initially before returning to a lower baseline (Supp. Fig. 6b). Even with model parameters where disomy evolved in most core populations by 1000 generations ($s = -0.01$, $\mu_{dip} = 2\times10^{-3}$), the diploidization index remained low in non-PAFC simulations (Supp. Fig. 7a) as the lack of fitness constraints results in pairing efficiency immediately falling towards 0 (Supp. Fig. 7b).

In contrast to core populations, pairing efficiency along the range edge is maintained near 1 throughout expansion with occasional sharp drops due to drift that are quickly recovered (Supp. Fig. 8). These drops generally coincide with an increase in the diploidization index, and occasionally are followed by the restoration of disomy. It is not clear why some but not all drops in pairing efficiency in tetrasomic populations result in the restoration of disomy. However, following the evolution of disomy, pairing efficiency is generally more stable around 1, indicating minimal losses due to gamete failure once disomy is restored.

When disomy evolves in edge populations, this state is generally maintained at that and any additionally colonized demes, indicating a fixation of this state on one side of the range edge (Supp. Fig. 9). Additionally, there is little spread of the disomic state to interior populations. When disomy evolves in interior populations, the spread is generally slow. One cause for this phenomenon may additionally be due to the PAFCs limiting migration, as F2 generations of offspring from tetrasomic and disomic parents may incur a fitness cost due to low pairing efficiency. If this is the case, we should

expect a higher F_{ST} between the populations along the tetrasomy-disomy border compared to other population comparisons. Indeed, we observe elevated F_{ST} along that border in simulations with PAFCs (Supp. Fig. 10a), but no such elevated F_{ST} when PAFCs are not included (Supp. Fig. 10b). Interestingly, we also observe higher F_{ST} between adjacent demes that are disomic as opposed to between adjacent demes that are tetrasomic (Supp. Fig. 10a) indicating additional downstream effects once disomy evolves. When PAFCs are removed from the model, F_{ST} between adjacent demes is minimal, including along the tetrasomy-disomy border (Supp. Fig. 10b). One result of this increased effective rate of gene flow is that, without PAFCs, disomy along the edge is transient, although it is also possible that reversion to tetrasomy is expected without PAFCs even in an isolated population–a scenario untested here.

Figure 6. Diploidization of core and edge populations under the pairing efficiency model for polyploids with a fully recessive model of dominance. (a) Average diploidization index of core and edge populations under varying selection coefficients for μ_{dip} = 1×10⁻³ and μ_{dip}=2×10⁻³. (b) Proportion of simulations where disomy is evolved (population average diploidization index ≥ 0.9) under varying selective coefficients for μ_{dip} = 1×10⁻³ and μ_{dip}=2×10⁻³. Dashed lines in (a) represent ± 1 standard deviation.

DISCUSSION

The genetic consequences of range expansion in polyploids

The results of this study have several implications for our understanding of polyploid evolution and establishment. Perhaps most immediately is the relationship between inheritance and expansion in polyploids, introducing an additional layer of complexity to the longstanding hypothesis that polyploidy confers an advantage to colonizing novel and deglaciated habitats. More specifically, outside of an unlikely model of additive dominance, the masking advantage provided by whole genome duplication when mutations are partially or fully recessive results in an accumulation of deleterious mutations that in turn leads to a protracted drop in the fitness of polyploids with tetrasomic inheritance during expansion relative to their diploid counterparts. Abundant evidence has demonstrated a link between polyploidy and previously glaciated areas (Brochmann et al. 2004; Paun et al. 2006; Novikova et al. 2018; Sutherland and Galloway 2018; Rice et al. 2019; David 2022; Booker et al. 2023). While support for this observation has many adaptive explanations–such as fixed heterozygosity (Stebbins 1985; Brochmann et al. 2004), increased adaptability of gene regulatory networks (Freeling and Thomas 2006; Hegarty and Hiscock 2007; Fusco et al. 2010; Yao et al. 2019; Ebadi et al. 2023), or the relaxed constraint on gene duplicates enabling more rapid adaptation (Lynch and Conery 2000; Lynch and Force 2000; Gout and Lynch 2015)–these arguments too rely on some level of subgenomic differentiation, and therefore disomy, to maintain their validity. As such, the results herein suggest polyploids with tetrasomic inheritance should be uniquely disadvantaged in colonizing new areas as polyploidy confers significant consequences as a result of expansion load–not to mention additional disadvantages due to meiotic errors as a result of polyploidy (Mayer and Aguilera 1990; Comai et al. 2000; Ramsey and Schemske 2002; Morgan et al. 2020). However, more work extending the scenarios examined here to include other potential advantages of polyploidy would be required to determine whether or not the adaptive benefits to polyploidy may in some cases offset the disadvantages that we show here.

One potential way to offset the disadvantages inherent to autopolyploids expanding their range is through interploid hybridization. Despite the fitness consequences of hybridizing across ploidies, numerous studies have demonstrated pervasive gene flow between autopolyploids and their diploid relatives when their ranges overlap (Arnold et al. 2016; Marburger et al. 2019; Monnahan et al. 2019; Bogart et al. 2020; Novikova et al. 2020; Shastry et al. 2021; Booker et al. 2022). Although hybridization across ploidies during formation is often considered a way for neo-polyploids to overcome the massive population bottleneck during formation (Stebbins 1980), admixture following their establishment should be generally disfavored as mixed-ploidy offspring suffer extensive fitness consequences (Ramsey and Schemske 1998)–though this may be mitigated in higher-level ploidies (Peskoller et al. 2021; Sutherland and Galloway 2021). However, hybridization to relieve the deleterious load accumulated through range expansion may help to explain why this phenomenon is so pervasive, and why the evolutionary histories

of autopolyploids can be so reticulate (e.g. Booker et al. 2022). Importantly, interploid hybridization has also been documented in numerous allopolyploids (Ma et al. 2010; Sutherland and Galloway 2018; Kryvokhyzha et al. 2019b; Kryvokhyzha et al. 2019a; Zhao et al. 2019). While disomic neo-allopolyploids should be shielded from the effects of expansion, some level of tetrasomy (as is often observed in allopolyploids, see Li et al. 2021) or fractionation should make some degree of post-expansion interploid hybridization more likely.

While autopolyploids may be uniquely disadvantaged in expanding their range under realistic models of dominance, the results of this study show a markedly different pattern in allopolyploids. Allopolyploids may be at a disadvantage under an additive model, though this is largely due to fitness consequences incurred in isolation. However the restriction of recombination across subgenomes prevents the fixation of alleles as a result of expansion (unless the same mutation occurs on both subgenomes), therefore providing a benefit if deleterious mutations are fully or partially recessive. Importantly, the model investigated here is of a neo-allopolyploid where no fractionation or genic diploidization, wherein gene duplicates are gradually lost or silenced leading to dynamics more and more similar to diploids, has occurred. In reality, allopolyploids exist somewhere on the continuum between the neo-allopolyploids investigated here and a functionally diploid paleo-polyploid. Anywhere outside of the functionally diploid paleo-polyploid extreme, allopolyploids should suffer fewer consequences of expansion load relative to diploids. However, because disomic inheritance results in the accumulation of fixed deleterious mutations within each subgenome (Fig. 4e), these dynamics may have consequences for the process of fractionation (gene silencing, subfunctionalization, and loss) in later stages of polyploid evolution. Furthermore, because genes on one subgenome may have a high number of deleterious mutations, this may hinder the adaptability of allopolyploids through functional redundancy.

Although the present study focuses on the consequences of expansion for polyploids already formed, the work here and elsewhere provokes a reconsideration of the hypotheses purported by Stebbins (1984, 1985) and evidenced by Brochmann et al. (2004): that allopolyploid formation is promoted in the arctic as glacial cycles promote secondary contact which can fix heterozygosity in newly formed polyploids. Macpherson et al. (2022) showed that range expansion promotes hybridization of species when they come in contact by providing immediate relief to the genetic load built up by expansion, and under some parameters the benefit provided by hybridization is enough to counteract the negative effects of Bateson-Dobzhansky-Muller incompatibilities. Allopolyploid formation rather than diploid hybridization provides these same benefits, while also introducing strong reproductive isolation. As demonstrated here, should these newly formed allopolyploids have largely disomic inheritance, their formation here would not only provide immediate relief to the already fixed load incurred by the

range-expanding diploids, but they would also be protected from the build up of additional load as expansion continues. Although the establishment and persistence of a new polyploid under these circumstances is still a strong evolutionary hurdle to overcome, the benefits relative to diploids regarding expansion load should make this more likely.

The terms we use here as autopolyploid and allopolyploid provide shorthand for tetrasomically and disomically inheriting polyploids. The definition of these terms are not exact, often referring to the nature of their formation or behavior in meiosis. In reality, nearly half of all identified allopolyploids have some level of multivalent formation (Li et al. 2021) and therefore deviate from the disomic inheritance modeled in allopolyploids here. As such, the particular consequences of expansion prescribed to autotetraploids are generally applicable across polyploids to the degree which their inheritance allows deleterious mutations to fix across all chromosomal copies. Similarly, the benefits described here to allotetraploids translate to the extent in which recombination across homoeologs is restricted, and many tetraploids may experience dynamics somewhere in between those of the fully tetrasomic and disomic tetraploids examined here. Additionally, although this work focuses on tetraploids, the masking potential of polyploidy–and the consequences of that masking when those species expand their range–should be generally applicable to polyploids that are not tetraploids.

Cytogenetic diploidization of range-expanding polyploids

We modeled the dynamics that regulate the inheritance pattern of polyploid meiosis during a range expansion and observed several interesting behaviors related to both the nature of diploidization and the fitness consequences of expansion. Regardless of the method by which cytogenetic diploidization proceeds, results from this study show that expansion increases the rate at which disomy is restored in a polyploid. Although the models of diploidization explored here are fairly simple, they represent two empirically identified modes of cytogenetic diploidization in nature. The better-understood of these two models is the dominant meiotic model, which incorporates dominant mutations that alter the meiotic machinery itself. Among these *Ph1*, a gene identified in grasses that when present suppresses pairing of homoeologs and facilitates pairing of homologs by affecting the interactions between microtubules and kinases (Vega and Feldman 1998), has been the most extensively studied (Riley and Chapman 1958; Riley 1960; Sears 1976; Sánchez-Morán et al. 2001; Feldman and Levy 2012; Li et al. 2021). A similar gene that experiences mutations of large effect which regulate pairing behavior is *PrBn* in *Brassica napus* (Jenczewski et al. 2003), and an additional QTL of large effect has also been found in *Arabidopsis suecica* (Henry et al. 2014). However, other identified pathways to the genetic control of meiosis have also been observed, such as *MHS4* copy number regulation that may be preserved in angiosperms generally (Gonzalo et al.

2019) and a complex landscape of epistatic interactions in *Arabidopsis arenosa* and *A. thaliana* neo-polyploids (Morgan et al. 2020; Morgan et al. 2022; Gonzalo et al. 2023).

Under the dominant meiotic model, the evolution of disomy is largely stochastic and minimally dependent on the fitness consequences of expansion (Fig. 5b,c). Instead, the rate of evolution of disomy is determined by the number of mutations required to restore disomy, and the chance that any already-segregating mutations fix is higher on the expansion edge. Still, there is some evidence that the genetic load incurred by expansion that remains in previously colonized areas influences the rate of diploidization under this model as diploidization mutations expand out from these populations (i.e. elevated core diploidization rates; s = -0.01, Fig. 5a,b). As well, even if disomy is not completely restored along the edge when $\lambda > 1$, any mutations that influence inheritance and that happened to fix along the edge may persist as the range continues to expand and the former edge populations become interior populations (Supp. Fig. 4). Thus, the stochastic fixation of disomy-restoring mutations along the range edge may increase the rate at which disomy evolves in the species across a broad portion of its range.

While our best-known examples of meiotic regulation in polyploids come from genetically controlled mechanisms such as those described above, it is not necessarily the case that this is the predominant mechanism of cytogenetic diploidization as identifying the relative contribution of chromosomal differentiation to pairing behavior (i.e. the pairing efficiency model) is inherently more difficult. Still, competition experiments have shown that pairing preferences are linked to some aspect of sequence divergence in certain systems (Bomblies 2022). Indeed, both large structural changes, such as inversions (Parra-Nunez et al. 2018), and the more uniform accumulation of sequence divergence along the chromosome (Braz et al. 2021) have been shown to result in preferential pairing.

We found the influence of expansion on the evolution of disomy through the pairing efficiency model to be patently different from that of the dominant meiotic model. Under this model, cytogenetic diploidization is more deterministic (Supp. Fig. 9) and the rate of disomy restoration is linked to the selection coefficient in both core and edge populations (Fig. 6b,c). Although this was not the original purpose of the present study, the result that adding deleterious recessive mutations to the model of Le Comber et al. (2010) increases the speed in which disomy is restored lends additional credence to the validity of this model as a pathway to diploidization. Furthermore, this result also makes this model less reliant on subfunctionalization and neofunctionalization as a necessary accelerator of diploidization.

Deleterious recessive mutations influence the rate of disomy restoration because during the tetrasomic phase of a population there is a fitness consequence (PAFCs) to heterozygosity, and selection acts to increase homozygosity generally–increasing the probability recessive mutations will be found in the homozygous state as well. While this acts to reduce fitness due to deleterious variation generally, it also introduces a selective pressure to transition to disomy, where high levels of nucleotide diversity are no longer disadvantageous. The result of this behavior on edge populations is drastic: the reduction in nucleotide diversity as a result of drift due to expansion acts in tandem with the similar reduction due to pairing behavior (Supp. Fig. 5a). As a consequence, fitness drops rapidly along the expansion front (Supp. Fig. 5b). This adds a strong selective pressure to transition to disomy, providing compelling evidence that expansion may be an important driver of cytogenetic diploidization in polyploids. Furthermore, the consequences of this behavior also suggest that autopolyploids transitioning to disomy through preferential pairing of chromosomes may suffer additional fitness losses during expansion (Supp. Fig. 5d).

Importantly, the empirical examples above discuss pairing behavior as it relates to both chiasma formation (e.g. bivalents or multivalents) and inheritance. While the pairing efficiency model indirectly includes some consequences of multivalent formation (PAFCs), we do not explicitly model multivalent formation as part of these models. In large part this is to simplify the models and identify the influence of range expansion on disomy restoration as a potential conduit of diploidization. Evolving disomy can be viewed as an extreme alteration to the recombination landscape of an organism, and additional work that might identify other routes of recombination modification such as modeling multivalent formation, more realistic recombination maps, and dynamic recombination rates and landscapes as chromosomes diverge may provide additional routes through which an expansion influences diploidization of polyploids.

Conclusions

The present work was conducted to examine aspects of polyploid evolution and establishment under two processes all polyploids proceed through: range expansion and diploidization. To do so, we have introduced more realistic models that allowed us to observe behaviors that do not follow additively from these extensions in complexity. In one case the behavior renders an otherwise advantageous element disadvantageous: in autopolyploids, the duplicated genome provides a fitness benefit to an isolated population through masking recessive mutations, but this same mechanism in turn results in severe fitness consequences as deleterious mutations accumulate along the range edge as the population expands. In other instances we made observations that were wholly unexpected. For example, while the transition from tetrasomy to disomy should provide a fitness benefit to an expanding autopolyploid, the

process of this transition may result in significant additional fitness consequences that could delay any such benefit (i.e. if disomy evolves through the accumulation of differences between subgenomes, preferential pairing increases the likelihood deleterious mutations are found in the homozygous state). Additionally, by modeling cytogenetic diploidization spatially, we found that asynchronous evolution of disomy across the landscape can create barriers to gene flow–a behavior that could result in future speciation. Spatial dynamics play an important role in the evolutionary trajectory of all species, but the conditions that regulate polyploid formation and establishment suggest an elevated importance of this role in polyploid evolution. We anticipate that future work incorporating these dynamics into population genetic models of polyploids will continue to help answer remaining questions and paradoxes regarding polyploid evolution.

ACKNOWLEDGEMENTS

We would first like to thank Ben Haller and Peter Ralph for providing prompt and necessary feedback while developing the polyploid models for these simulations. We would also like to thank the attendants of the 2023 Polyploidy Tree of Life Conference for their helpful comments and feedback of this work as it was ongoing, Tyler Kent for providing valuable feedback on this manuscript and discussion as the project was ongoing, and the remainder of the Schrider lab for their input throughout this project. This work was supported by NIH grants HG010774 and GM138286.

DATA ACCESSIBILITY

All scripts used for this study can be found in the github repository
https://github.com/wbooker/polyploid_expansion_load

REFERENCES

Adams, K. L., and J. F. Wendel. 2005. Polyploidy and genome evolution in plants. Current Opinion in Plant Biology 8:135–141.

Albertin, W., and P. Marullo. 2012. Polyploidy in fungi: evolution after whole-genome duplication. Proceedings. Biological Sciences / the Royal Society 279:2497–2509.

Arnold, B. J., B. Lahner, J. M. DaCosta, C. M. Weisman, J. D. Hollister, D. E. Salt, K. Bomblies, et al. 2016. Borrowed alleles and convergence in serpentine adaptation. Proceedings of the National Academy of Sciences of the United States of America 113:8320–8325.

Barker, M. S., B. C. Husband, and J. C. Pires. 2016. Spreading Winge and flying high: The evolutionary importance of polyploidy after a century of study. American Journal of Botany 103:1139–1145.

Bennett, J. H. 1976. Expectations for inbreeding depression on self-fertilization of tetraploids. Biometrics 32:449–452.

Bever, J. D., and F. R. Felber. 1992. The theoretical population genetics of autopolyploidy. Oxford surveys in evolutionary biology 8:185–185.

Bogart, J. P., P. Burgess, and J. Fu. 2020. Revisiting the evolution of the North American tetraploid treefrog (Hyla versicolor). Genome 63:547–560.

Bomblies, K. 2022. Learning to tango with four (or more): the molecular basis of adaptation to polyploid meiosis. Plant reproduction.

Booker, W. W., H. C. Gerhardt, A. R. Lemmon, M. B. Ptacek, A. T. B. Hassinger, J. Schul, and E. M. Lemmon. 2022. The complex history of genome duplication and

hybridization in north american gray treefrogs. Molecular Biology and Evolution 39.

Booker, W. W., E. M. Lemmon, A. R. Lemmon, M. B. Ptacek, A. T. B. Hassinger, J. Schul, and H. C. Gerhardt. 2023. Biogeography and the evolution of acoustic communication in the polyploid North American grey treefrog complex. Molecular Ecology 32:4863–4879.

Braz, G. T., F. Yu, H. Zhao, Z. Deng, J. A. Birchler, and J. Jiang. 2021. Preferential meiotic chromosome pairing among homologous chromosomes with cryptic sequence variation in tetraploid maize. The New Phytologist 229:3294–3302.

Brochmann, C., A. K. Brysting, I. G. Alsos, L. Borgen, H. H. Grundt, A. C. Scheen, and R. Elven. 2004. Polyploidy in arctic plants. Biological Journal of the Linnean Society 82:521–536.

Comai, L., A. P. Tyagi, K. Winter, R. Holmes-Davis, S. H. Reynolds, Y. Stevens, and B. Byers. 2000. Phenotypic instability and rapid gene silencing in newly formed arabidopsis allotetraploids. The Plant Cell 12:1551–1568.

Comai, L. 2005. The advantages and disadvantages of being polyploid. Nature Reviews. Genetics 6:836–846.

Conant, G. C., J. A. Birchler, and J. C. Pires. 2014. Dosage, duplication, and diploidization: clarifying the interplay of multiple models for duplicate gene evolution over time. Current Opinion in Plant Biology 19:91–98.

David, K. T. 2022. Global gradients in the distribution of animal polyploids. Proceedings of the National Academy of Sciences of the United States of America 119:e2214070119.

Ebadi, M., Q. Bafort, E. Mizrachi, P. Audenaert, P. Simoens, M. Van Montagu, D. Bonte, et al. 2023. The duplication of genomes and genetic networks and its potential for evolutionary adaptation and survival during environmental turmoil. Proceedings of the National Academy of Sciences of the United States of America 120:e2307289120.

Feldman, M., A. Levy, B. Chalhoub, and K. Kashkush. 2012. Genomic plasticity in polyploid wheat. In P. S. Soltis & D. E. Soltis, eds., Polyploidy and genome evolution (pp. 109–135). Springer Berlin Heidelberg, Berlin, Heidelberg.

Feldman, M., and A. A. Levy. 2012. Genome evolution due to allopolyploidization in wheat. Genetics 192:763–774.

Freeling, M., and B. C. Thomas. 2006. Gene-balanced duplications, like tetraploidy, provide predictable drive to increase morphological complexity. Genome Research 16:805–814.

Freeling, M., M. R. Woodhouse, S. Subramaniam, G. Turco, D. Lisch, and J. C. Schnable. 2012. Fractionation mutagenesis and similar consequences of mechanisms removing dispensable or less-expressed DNA in plants. Current Opinion in Plant Biology 15:131–139.

Fusco, D., L. Grassi, B. Bassetti, M. Caselle, and M. Cosentino Lagomarsino. 2010. Ordered structure of the transcription network inherited from the yeast whole-genome duplication. BMC Systems Biology 4:77.

Gonzalo, A., M.-O. Lucas, C. Charpentier, G. Sandmann, A. Lloyd, and E. Jenczewski. 2019. Reducing MSH4 copy number prevents meiotic crossovers between non-homologous chromosomes in Brassica napus. Nature Communications

10:2354.

Gonzalo, A., P. Parra-Nunez, A. L. Bachmann, E. Sanchez-Moran, and K. Bomblies. 2023. Partial cytological diploidization of neoautotetraploid meiosis by induced cross-over rate reduction. Proceedings of the National Academy of Sciences of the United States of America 120:e2305002120.

Gout, J.-F., and M. Lynch. 2015. Maintenance and loss of duplicated genes by dosage subfunctionalization. Molecular Biology and Evolution 32:2141–2148.

Grandont, L., E. Jenczewski, and A. Lloyd. 2013. Meiosis and its deviations in polyploid plants. Cytogenetic and Genome Research 140:171–184.

Gregory, T. R., and B. K. Mable. 2005. Polyploidy in Animals. The evolution of the genome (pp. 427–517). Elsevier.

Haldane, J. B. S. 1932. The Causes of Evolution (1st ed.). Longmans, Green and Co., New York.

Haller, B. C., and P. W. Messer. 2023. SLiM 4: Multispecies Eco-Evolutionary Modeling. The American Naturalist 201:E127–E139.

Hegarty, M., and S. Hiscock. 2007. Polyploidy: doubling up for evolutionary success. Current Biology 17:R927-9.

Henry, I. M., B. P. Dilkes, A. Tyagi, J. Gao, B. Christensen, and L. Comai. 2014. The BOY NAMED SUE quantitative trait locus confers increased meiotic stability to an adapted natural allopolyploid of Arabidopsis. The Plant Cell 26:181–194.

Hill, R. R. 1970. Selection in autotetraploids. TAG. Theoretical and Applied Genetics. Theoretische und Angewandte Genetik 41:181–186.

Huber, C. D., A. Durvasula, A. M. Hancock, and K. E. Lohmueller. 2018. Gene

expression drives the evolution of dominance. Nature Communications 9:2750.

Husband, B. C. 2000. Constraints on polyploid evolution: a test of the minority cytotype exclusion principle. Proceedings. Biological Sciences / the Royal Society 267:217–223.

Jenczewski, E., F. Eber, A. Grimaud, S. Huet, M. O. Lucas, H. Monod, and A. M. Chèvre. 2003. PrBn, a major gene controlling homeologous pairing in oilseed rape (Brassica napus) haploids. Genetics 164:645–653.

Kacser, H., and J. A. Burns. 1981. The molecular basis of dominance. Genetics 97:639–666.

Kryvokhyzha, D., P. Milesi, T. Duan, M. Orsucci, S. I. Wright, S. Glémin, and M. Lascoux. 2019a. Towards the new normal: Transcriptomic convergence and genomic legacy of the two subgenomes of an allopolyploid weed (Capsella bursa-pastoris). PLoS Genetics 15:e1008131.

Kryvokhyzha, D., A. Salcedo, M. C. Eriksson, T. Duan, N. Tawari, J. Chen, M. Guerrina, et al. 2019b. Parental legacy, demography, and admixture influenced the evolution of the two subgenomes of the tetraploid Capsella bursa-pastoris (Brassicaceae). PLoS Genetics 15:e1007949.

Levin, D. A. 1975. Minority Cytotype Exclusion in Local Plant Populations. Taxon 24:35.

Levin, D. A. 2002. The Role of Chromosomal Change in Plant Evolution (illustrated ed.). Oxford University Press.

Le Comber, S. C., M. L. Ainouche, A. Kovarik, and A. R. Leitch. 2010. Making a functional diploid: from polysomic to disomic inheritance. The New Phytologist 186:113–122.

Lien, S., B. F. Koop, S. R. Sandve, J. R. Miller, M. P. Kent, T. Nome, T. R. Hvidsten, et al. 2016. The Atlantic salmon genome provides insights into rediploidization. Nature 533:200–205.

Li, Z., M. T. W. McKibben, G. S. Finch, P. D. Blischak, B. L. Sutherland, and M. S. Barker. 2021. Patterns and processes of diploidization in land plants. Annual review of plant biology 72:387–410.

Li, Z., G. P. Tiley, S. R. Galuska, C. R. Reardon, T. I. Kidder, R. J. Rundell, and M. S. Barker. 2018. Multiple large-scale gene and genome duplications during the evolution of hexapods. Proceedings of the National Academy of Sciences of the United States of America 115:4713–4718.

Liu, Z., K. Adamczyk, M. Manzanares-Dauleux, F. Eber, M.-O. Lucas, R. Delourme, A. M. Chèvre, et al. 2006. Mapping PrBn and other quantitative trait loci responsible for the control of homeologous chromosome pairing in oilseed rape (Brassica napus L.) haploids. Genetics 174:1583–1596.

Lynch, M., and J. S. Conery. 2000. The evolutionary fate and consequences of duplicate genes. Science 290:1151–1155.

Lynch, M., and A. Force. 2000. The probability of duplicate gene preservation by subfunctionalization. Genetics 154:459–473.

Mable, B. K., and S. P. Otto. 2001. Masking and purging mutations following EMS treatment in haploid, diploid and tetraploid yeast (Saccharomyces cerevisiae). Genetical Research 77:9–26.

MacPherson, A., S. Wang, R. Yamaguchi, L. Rieseberg, and S. Otto. 2022. Parental population range expansion before secondary contact promotes heterosis. The

American Naturalist.

Ma, J.-X., Y.-N. Li, C. Vogl, F. Ehrendorfer, and Y.-P. Guo. 2010. Allopolyploid speciation and ongoing backcrossing between diploid progenitor and tetraploid progeny lineages in the Achillea millefolium species complex: analyses of single-copy nuclear genes and genomic AFLP. BMC Evolutionary Biology 10:100.

Mandáková, T., and M. A. Lysak. 2018. Post-polyploid diploidization and diversification through dysploid changes. Current Opinion in Plant Biology 42:55–65.

Marburger, S., P. Monnahan, P. J. Seear, S. H. Martin, J. Koch, P. Paajanen, M. Bohutínská, et al. 2019. Interspecific introgression mediates adaptation to whole genome duplication. Nature Communications 10:5218.

Mayer, V. W., and A. Aguilera. 1990. High levels of chromosome instability in polyploids of Saccharomyces cerevisiae. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis 231:177–186.

Monnahan, P., F. Kolář, P. Baduel, C. Sailer, J. Koch, R. Horvath, B. Laenen, et al. 2019. Pervasive population genomic consequences of genome duplication in Arabidopsis arenosa. Nature Ecology & Evolution 3:457–468.

Moreau, C., C. Bhérer, H. Vézina, M. Jomphe, D. Labuda, and L. Excoffier. 2011. Deep human genealogies reveal a selective advantage to be on an expanding wave front. Science 334:1148–1150.

Morgan, C., E. Knight, and K. Bomblies. 2022. The meiotic cohesin subunit REC8 contributes to multigenic adaptive evolution of autopolyploid meiosis in Arabidopsis arenosa. PLoS Genetics 18:e1010304.

Morgan, C., M. A. White, F. C. H. Franklin, D. Zickler, N. Kleckner, and K. Bomblies.

2021. Evolution of crossover interference enables stable autopolyploidy by ensuring pairwise partner connections in Arabidopsis arenosa. Current Biology 31:4713-4726.e4.

Morgan, C., H. Zhang, C. E. Henry, F. C. H. Franklin, and K. Bomblies. 2020. Derived alleles of two axis proteins affect meiotic traits in autotetraploid Arabidopsis arenosa. Proceedings of the National Academy of Sciences of the United States of America 117:8980–8988.

Novikova, P. Y., I. G. Brennan, W. Booker, M. Mahony, P. Doughty, A. R. Lemmon, E. Moriarty Lemmon, et al. 2020. Polyploidy breaks speciation barriers in Australian burrowing frogs Neobatrachus. PLoS Genetics 16:e1008769.

Novikova, P. Y., N. Hohmann, and Y. Van de Peer. 2018. Polyploid Arabidopsis species originated around recent glaciation maxima. Current Opinion in Plant Biology 42:8–15.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and the phylogenomics of green plants. Nature 574:679–685.

Otto, S. P., and J. Whitton. 2000. Polyploid Incidence and Evolution. Annual Review of Genetics 34:401–437.

Otto, S. P. 2007. The evolutionary consequences of polyploidy. Cell 131:452–462.

Parey, E., A. Louis, J. Montfort, Y. Guiguen, H. Roest Crollius, and C. Berthelot. 2022. An atlas of fish genome evolution reveals delayed rediploidization following the teleost whole-genome duplication. Genome Research 32:1685–1697.

Parisod, C., R. Holderegger, and C. Brochmann. 2010. Evolutionary consequences of autopolyploidy. The New Phytologist 186:5–17.

Parra-Nunez, P., M. Pradillo, and J. L. Santos. 2018. Competition for Chiasma
    Formation Between Identical and Homologous (But Not Identical) Chromosomes
    in Synthetic Autotetraploids of Arabidopsis thaliana. Frontiers in Plant Science
    9:1924.

Paun, O., T. F. Stuessy, and E. Hörandl. 2006. The role of hybridization, polyploidization
    and glaciation in the origin and evolution of the apomictic Ranunculus cassubicus
    complex. The New Phytologist 171:223–236.

Peischl, S., I. Dupanloup, M. Kirkpatrick, and L. Excoffier. 2013. On the accumulation of
    deleterious mutations during range expansions. Molecular Ecology
    22:5972–5982.

Peischl, S., and L. Excoffier. 2015. Expansion load: recessive mutations and the role of
    standing genetic variation. Molecular Ecology 24:2084–2094.

Peischl, S., M. Kirkpatrick, and L. Excoffier. 2015. Expansion load and the evolutionary
    dynamics of a species range. The American Naturalist 185:E81-93.

Peskoller, A., L. Silbernagl, K. Hülber, M. Sonnleitner, and P. Schönswetter. 2021. Do
    pentaploid hybrids mediate gene flow between tetraploid Senecio disjunctus and
    hexaploid S. carniolicus s. str. (S. carniolicus aggregate, Asteraceae)? Alpine
    botany.

Ramsey, J., and D. W. Schemske. 1998. Pathways, mechanisms, and rates of polyploid
    formation in flowering plants. Annual review of ecology and systematics
    29:467–501.

Ramsey, J., and D. W. Schemske. 2002. NEOPOLYPLOIDY IN FLOWERING PLANTS.
    Annual review of ecology and systematics 33:589–639.

Redmond, A. K., M. K. Gundappa, D. J. Macqueen, and A. McLysaght. 2022. Extensive lineage-specific rediploidisation masks shared whole genome duplication in the sturgeon-paddlefish ancestor. bioRxiv.

Rice, A., P. Šmarda, M. Novosolov, M. Drori, L. Glick, N. Sabath, S. Meiri, et al. 2019. The global biogeography of polyploid plants. Nature Ecology & Evolution 3:265–273.

Riley, R., and V. Chapman. 1958. Genetic Control of the Cytologically Diploid Behaviour of Hexaploid Wheat. Nature 182:713–715.

Riley, R. 1960. The diploidisation of polyploid wheat. Heredity 15:407–429.

Robertson, F. M., M. K. Gundappa, F. Grammes, T. R. Hvidsten, A. K. Redmond, S. Lien, S. A. M. Martin, et al. 2017. Lineage-specific rediploidization is a mechanism to explain time-lags between genome duplication and evolutionary diversification. Genome Biology 18:111.

Rodriguez, D. J. 1996. A Model for the Establishment of Polyploidy in Plants. The American Naturalist 147:33–46.

Sánchez-Morán, E., E. Benavente, and J. Orellana. 2001. Analysis of karyotypic stability of homoeologous-pairing (ph) mutants in allopolyploid wheats. Chromosoma 110:371–377.

Schlichta, F., S. Peischl, and L. Excoffier. 2022. The impact of genetic surfing on neutral genomic diversity. Molecular Biology and Evolution 39.

Scott, A. D., N. W. M. Stenz, P. K. Ingvarsson, and D. A. Baum. 2016. Whole genome duplication in coast redwood (Sequoia sempervirens) and its implications for explaining the rarity of polyploidy in conifers. The New Phytologist 211:186–193.

Sears, E. R. 1976. Genetic control of chromosome pairing in wheat. Annual Review of Genetics 10:31–51.

Shastry, V., P. E. Adams, D. Lindtke, E. G. Mandeville, T. L. Parchman, Z. Gompert, and C. A. Buerkle. 2021. Model-based genotype and ancestry estimation for potential hybrids with mixed-ploidy. Molecular Ecology Resources 21:1434–1451.

Stebbins, G. L. 1950. Variation and evolution in plants. Columbia University Press.

Stebbins, G. L. 1980. Polyploidy in plants: unsolved problems and prospects. Basic life sciences 13:495–520.

Stebbins, G. L. 1984. Polyploidy and the distribution of the arctic-alpine flora : new evidence and a new approach. Birkhäuser.

Stebbins, G. L. 1985. Polyploidy, hybridization, and the invasion of new habitats. Annals of the Missouri Botanical Garden 72:824.

Stenberg, P., and A. Saura. 2013. Meiosis and its deviations in polyploid animals. Cytogenetic and Genome Research 140:185–203.

Sutherland, B. L., and L. F. Galloway. 2018. Effects of glaciation and whole genome duplication on the distribution of the Campanula rotundifolia polyploid complex. American Journal of Botany 105:1760–1770.

Sutherland, B. L., and L. F. Galloway. 2021. Variation in heteroploid reproduction and gene flow across a polyploid complex: One size does not fit all. Ecology and Evolution 11:9676–9688.

Tayalé, A., and C. Parisod. 2013. Natural pathways to polyploidy in plants and consequences for genome reorganization. Cytogenetic and Genome Research 140:79–96.

Van de Peer, Y., T.-L. Ashman, P. S. Soltis, and D. E. Soltis. 2021. Polyploidy: an
evolutionary and ecological force in stressful times. The Plant Cell 33:11–26.

Van de Peer, Y., E. Mizrachi, and K. Marchal. 2017. The evolutionary significance of
polyploidy. Nature Reviews. Genetics 18:411–424.

Vega, J. M., and M. Feldman. 1998. Effect of the pairing gene Ph1 on centromere
misdivision in common wheat. Genetics 148:1285–1294.

Wolfe, K. H. 2001. Yesterday's polyploids and the mystery of diploidization. Nature
Reviews. Genetics 2:333–341.

Yamauchi, A., A. Hosokawa, H. Nagata, and M. Shimoda. 2004. Triploid bridge and role
of parthenogenesis in the evolution of autopolyploidy. The American Naturalist
164:101–112.

Yao, Y., L. Carretero-Paulet, and Y. Van de Peer. 2019. Using digital organisms to study
the evolutionary consequences of whole genome duplication and polyploidy. Plos
One 14:e0220257.

Zhan, S. H., M. Drori, E. E. Goldberg, S. P. Otto, and I. Mayrose. 2016. Phylogenetic
evidence for cladogenetic polyploidization in land plants. American Journal of
Botany 103:1252–1258.

Zhao, N., C. E. Grover, Z. Chen, J. F. Wendel, and J. Hua. 2019. Intergenomic gene
transfer in diploid and allopolyploid Gossypium. BMC Plant Biology 19:492.

Supplemental Figure 1. Unscaled mean fitness of the edge deme for additive models with no beneficial mutations.

Supplemental Figure 2. Fitness trajectory of a single population of diploids or autotetraploids under a recessive dominance model. Solid lines represent the average fitness from 100 simulations and dashed lines represent 1 standard deviation. For all simulations, K = 500, μ_{del} = 2.5×10^{-8}.

Supplemental Figure 3 . Diploidization of core and edge populations under the dominant meiotic model for polyploids with a fully recessive model of dominance. (a) Average diploidization index of core and edge populations (s = -0.01). (b) Proportion of simulations where disomy is evolved (population average diploidization index = 1) (s = -0.01). Dashed lines for (a) represent ± 1 standard deviation. For λ = 1 and 2 the number of simulations ran was 100 rather than 25 for other models due to the large variance.

Supplemental Figure 4. Diploidization index of example simulations under the dominant meiotic diploidization model with 1-5 mutations required to restore disomy. All simulations were run under a fully recessive dominance model with *s* = -0.005.

Supplemental Figure 5. The cost of pairing efficiency regulated diploidization on fitness and nucleotide diversity. a) Nucleotide diversity (π) in each generation of core and edge populations when the model includes no diploidization, diploidization under the pairing efficiency model with pairing associated fitness costs (PAFC), and diploidization under the pairing efficiency model without PAFC. b) Mean fitness (scaled at the start of expansion) in each generation of core and edge populations when the model includes no diploidization, diploidization under the pairing efficiency model with pairing associated fitness costs (PAFC), and diploidization under the pairing efficiency model without PAFC. All models were run with *s* = -0.005 and μ_{dip} = 1×10^{-3}. Note the different *y*-axis scales for the edge and core populations.

Supplemental Figure 6. Interaction of selection coefficient of standard deleterious mutations and the fitness of gametes determined by pairing efficiency on the evolution of disomy under the pairing efficiency model of diploidization. a) The proportion of simulations where disomy (diploidization index ≥ 0.9) has evolved if pairing efficiency affects (right) or does not (left) affect gamete fitness. b) Average diploidization index for each simulation if pairing efficiency affects (right) or does not affect (left) gamete fitness. Dashed line represents ± 1 standard deviation. For each parameter 25 simulations were run with μ_{dip} = 1×10⁻³.

Supplemental Figure 7. Effect of pairing efficiency affecting gamete fitness in core populations where disomy does evolve. For all figures, *s* = -0.01 and μ_{dip} = 2×10^{-3}. a) Average diploidization index for each simulation if pairing efficiency affects (right) or does not affect (left) gamete fitness. Dashed line represents ± 1 standard deviation, averages and standard deviation calculated from 25 independent simulations. b) Pairing efficiency of core populations if pairing efficiency affects (right) or does not affect (left) gamete fitness. Each line represents a single simulation.

Supplemental Figure 8. Pairing efficiency (black) and diploidization index (gray) of edge populations when disomy did not (a) or did (b) evolve within 1000 generations. Each figure represents a single simulation under a model where s = -0.005 and μ_{dip} = 1×10^{-3}.

Supplemental Figure 9. Diploidization index of example simulations under the pairing efficiency diploidization model. All simulations were run under a fully recessive dominance model with *s* = -0.005.

Supplemental Figure 10. Pairing associated fitness costs maintain disomy once evolved. a) Diploidization index (top) and F_{ST} between focal and previous deme (bottom) when diploidization involves pairing associated fitness costs. b) Diploidization index (top) and F_{ST} between focal and previous deme (bottom) when pairing associated fitness costs are not included in the diploidization model.