Phylogenomics, biogeography, and trait evolution of the Boletaceae (Boletales,

Agaricomycetes, Basidiomycota)

Keaton Tremble[a]*, Terry Henkel[b], Alexander Bradshaw[a], Colin Domnauer[a], Lyda

Brown[a], Lê Xuân Thám[c], Guliana Furci[d], Cathie Aime[e], Jean-Marc Moncalvo[f], Bryn

Dentinger[a]*

[a]Natural History Museum of Utah and School of Biological Sciences, University of

Utah, Salt Lake City, Utah 84108

[b]Department of Biological Sciences, California State Polytechnic University, Humboldt,

Arcata, California 95521

[c]Van Lang University, Ho Chi Minh City, Viet Nam

[d]Fungi Foundation, Brooklyn, New York 11222

[e]Department of Botany and Plant Pathology, Purdue University, West Lafayette,

Indiana 47907

[f]Department of Natural History, Royal Ontario Museum & Department of Ecology and

Evolutionary Biology, University of Toronto, Toronto M5S 2C6, Canada

ABSTRACT

The species-rich porcini mushroom family Boletaceae is a widespread and well-known

group of ectomycorrhizal (ECM) mushroom-forming fungi that has eluded intrafamilial

phylogenetic resolution despite many attempts using morphological traits and multi-

locus molecular datasets. In this study, we present a genome-wide molecular dataset of

1764 single-copy gene families from a global sampling of 418 Boletaceae specimens.

The resulting phylogenetic analysis has strong statistical support for most branches of

the tree, including the first statistically robust backbone. The enigmatic *Phylloboletellus*

chloephorus from non-ECM Argentinian subtropical forests was recovered as an early

diverging lineage within the Boletaceae. Time-calibrated branch lengths estimate that

the family first arose in the early- to mid-Cretaceous and underwent a rapid radiation in

the Eocene, possibly when the ECM nutritional mode arose with the emergence and

diversification of ECM angiosperms. Biogeographic reconstructions reveal a complex

history of vicariance and episodic long-distance dispersal correlated with historical

geologic events, including Gondwanan origins and cladogenesis patterns that parallel its

fragmentation. Ancestral state reconstruction of sporocarp morphological traits predicts

that the ancestor of the Boletaceae was lamellate with ornamented basidiospores,

contrary to most contemporary "bolete" morphologies. Transition rates indicated that

the lamellate hymenophore and sequestrate sporocarp are reversible traits. Together,

this study represents the most comprehensively sampled, data-rich molecular phylogeny

of the Boletaceae to date, enabling robust inferences of trait evolution and biogeography

in the group.

KEYWORDS: phylogenomics; Boletaceae; porcini; evolutionary radiation; Gondwana

INTRODUCTION

Evolutionary radiations result from short bursts of relatively rapid diversification and

demonstrate the creative power of evolution. Adaptive and non-adaptive radiations may

be the most common macroevolutionary patterns and are fundamental to the origins of

biodiversity (Simoes et al. 2016; Futuyma 1998; Schluter and McPeek 2000). Yet, the

causes of evolutionary radiations are poorly known for most groups, as most studies

have focused on animals and plants (Soulebeau et al. 2015). The other major

multicellular eukaryotic group, the Fungi, have largely been neglected (Varga et al.

2019).

The porcini mushroom family Boletaceae (FIG. 1) is an example of an

evolutionary radiation in Fungi (Bruns et al. 1992). The family is exceptionally diverse

(>2000 currently accepted species) and globally distributed, but poorly documented for

many regions. Yet, boletoid fungi are prevalent ectomycorrhizal (ECM) mutualists in

ecosystems dominated by ECM plants (Peay et al. 2010) and at least eight species are

traded globally as wild-collected edible mushrooms (Arora 2008; Sitta and Floriani

2008; Dentinger et al. 2010; Dentinger and Suz 2014). Despite their conspicuous

sporocarps, ecological dominance, and cultural importance, new species of Boletaceae

are regularly described from poorly explored habitats around the globe (e.g. Halling et

al. 2006, 2023; Fulgenzi et al. 2007, 2008, 2010; Neves et al. 2010; Husbands et al.

2013; Castellano et al. 2016; Chakraborty et al. 2015; Das et al. 2015, 2016; Henkel et

al. 2016; Magnago et al. 2017). New species have also recently been described from

wild-collected foods in markets (e.g. Das et al. 2015; Dentinger and Suz 2014; Halling

et al. 2014). While new Boletaceae species are increasingly understood in a global

phylogenetic context, shedding light on their origin, diversification, and migration, over

20 years of molecular phylogenetic studies using legacy loci have made little progress

towards resolving the deepest nodes ("backbones") in Boletaceae phylogenies

(Grubisha et al. 2001; Binder and Hibbett 2006; Drehmel et al. 2008; Dentinger et al.

2010; Nuhn et al. 2016; Wu et al. 2014).

A prominent consequence of this lack of phylogenetic resolution is the recent

explosion of new generic names to accommodate newly discovered species, or species

that are included in molecular phylogenetic analyses for the first time and recovered on

long branches with no supported affinity to existing named genera (e.g., Castellano et

al. 2016; Henkel et al. 2016; Badou et al. 2022; Halling et al. 2023). Few of these

studies have followed recommended best practices for naming new genera (Vellinga et

al. 2015). Moreover, many of these new Boletaceae genera are monotypic and require

identification to recognize, an impractical solution to the problem. Taken together, the

current situation is perhaps best described as a quagmire of nomenclatural, taxonomic,

ecological, and evolutionary speculation.

Beyond taxonomic concerns, the Boletacaeae presents a unique system to

identify the genetic mechanisms that contribute to diversification. The Boletaceae

appear to have underwent an early evolutionary radiation between 60-100 mya (Bruns

and Palmer 1989; Binder et al. 2006; Dentinger et al. 2010; Wu et al. 2014, 2016; Sato

et al. 2019). This early radiation has been correlated with the convergent evolution of

morphological traits, such as the lamellate hymenophore and gasteromycetization

(Castellano et al. 2016), suggesting that emergence of morphological diversity is

constrained by relatively few changes in developmental pathways.

Many factors have contributed to difficulties in generating robust phylogenetic

reconstructions for the Boletaceae. While phenomena such as incomplete lineage

sorting and hybridization may obscure historical phylogenetic signal, previous datasets

for the Boletaceae had patchy taxonomic and geographic sampling. These factors

impact accurate phylogenetic reconstruction, possibly exacerbated by the

aforementioned rapid radiation event (Bruns et al. 1992; Sato et al. 2017). Without a

phylogeny that is based on globally representative taxon sampling and statistically well-

supported resolution at all depths of the tree, it is impossible to name, classify, and

understand evolutionary history in the Boletaceae. For example, only a few recent

studies have included representatives of the exceptionally rich Australian boletoid funga

(Halling, Fechner, et al. 2015, 2023; Halling, Nuhn, et al. 2012). Boletoid fungi from

the Neotropics and Afrotropics are rarely represented in family-level analyses despite

their exceptional species richness (e.g. Heinemann 1951; Henkel et al. 2012).

Recent fieldwork has resulted in many new collections from under-sampled

regions (B. Dentinger, T. Henkel, R. Halling; unpublished data), and these are now

available to include in phylogenetic datasets aiming to achieve the first globally

representative sampling of the Boletaceae. Collections-based phylogenomics is also

effective for resolving ancient relationships among mushroom-forming fungi (Dentinger

et al. 2015; Liimatainen et al. 2021; Tremble et al. 2020). However, no one has yet

applied these methods to the Boletaceae. Moreover, whole genome sequencing of

mushroom forming fungi provides opportunities to go beyond phylogenetic

reconstruction. For example, whole genome sequencing can exceed legacy loci in

identifying population processes that generate biodiversity (e.g. Tremble et al. 2022).

For this study, we generated the first phylogeny of the Boletaceae that utilized a

very large molecular dataset comprised of 1764 genome-wide loci from 418 taxa across

the family. Specimens were collected in many tropical and temperate geographic

regions. To obtain broad geographic and taxon coverage we included type fungarium

specimens and recent new collections. We utilized specimens from previously under-

sampled regions including tropical Africa, southern South America, lowland tropical

South America, and Australia. Type species were sampled to facilitate future taxonomic

revisions of genera. Using our highly resolved genome-based phylogeny, we also

performed the first inclusive biogeographic reconstruction of the Boletaceae and

assessed morphological trait evolution. Overall, we provide new insights into the broad

patterns of evolution of this enigmatic fungal group.

MATERIALS AND METHODS

Sampling.—Taxon selection focused on obtaining representatives of all currently

accepted genera, selecting type species whenever possible. Because the current

understanding of genera is incomplete and rapidly changing, we could not include

representatives of all currently accepted genera that were published during the course of

this study. Specimens from geographic regions unrepresented in prior studies were also

included. A total of 418 Boletaceae specimens were gathered from a global distribution

using collections made by the authors, those borrowed from four institutions, and

donations from citizen scientists (SUPPLEMENTAL TABLE 1). In addition, we

utilized genome data publicly available from the JGI Mycocosm Portal (Grigoriev et al.

2014) for *Boletus coccyginus*, *B. reticuloceps*, *Butyriboletus roseoflavus*, *Chiua virens*,

Lanmaoa asiatica, and *Imleria badia* (Miyauchi et al. 2020, Wu et al. 2022, Kohler et

al. 2015). *Paxillus involutus*, *Paxillus adelphus*, and *Hydnomerulius pinastri* genomes

from JGI were used for outgroups (Kohler et al. 2015).

DNA extraction and sequencing. —Genomic DNA was extracted in one of three ways.

1) 10 mg of hymenophore tissue from each specimen was homogenized in 2.0 ml

screw-cap tubes containing a single 3.0 mm and 8 x 1.5 mm stainless steel beads using a

BeadBug™ microtube homogenizer (Sigma-Aldrich, #Z763713) for 120 seconds at a

speed setting of 3500 rpm. After physical disruption, DNA was extracted using the

Monarch® Genomic DNA Purification Kit (New England Biolabs, Massachusetts;

#T3010) with the Monarch® gDNA Tissue Lysis Buffer (#T3011) using double the

volume of lysis buffer, one hour lysis incubation at 56 C , and 550 µl of wash buffer

during each of the wash steps. 2) an in-house 96-well plate protocol where tissue is

physically homogenized, as above, after which 1000 µL of lysis buffer (1% sodium

dodecyl sulfate, 10 mM Tris, 10 mM EDTA, 5mM NaCl, 50mM dithiothreitol, pH 8.0)

is added. To this solution is added 4 µL of RNAse A (20 mg/mL), the solution is

vortexed, and then incubated at 37 C for 10 min. Next, 10 µL of proteinase K (20

mg/mL) is added, the solution vortexed, and then incubated at 56 C overnight on an

Eppendorf ThermoMixer® with agitation at 400 rpm. After lysis, the tubes are

centrifuged at max speed (17,000 x g) to pellet the cellular debris. 700 µL of

supernatant is removed to a new 1.7 mL microcentrifuge tube with hinged cap to which

162.5 µL 3.0 M potassium acetate (pH 5.5) is added. The solution is mixed briefly and

then put on ice for five min, followed by a second centrifugation, as above. Avoiding

the pellet, the supernatant is removed to a well of a 96-well 10 µM filter plate

(Enzymax, #EZ96FTP) set in a 2 mL MASTERBLOCK® collection plate (Grainer,

#780271). Filtration is achieved through centrifugation at 1500 x g for 2 min. The flow-

through is transferred to a new 1.7 ml microcentrifuge tube with hinged cap and

centrifuged, as above. Without disturbing the pellet, the supernatant is removed to a

new 2.0 mL microcentrifuge tupe with hinged cap and 1000 µL of binding buffer (5M

guanidium hydrochloride, 40% isopropanol) is added and the solution homogenized by

pipetting. The binding solution is then transferred to a well of a 96-well long-tip

AcroPrep™ Plate (PALL, #8133) that was pre-conditioned by pulling 400 µl Tris-HCl

buffer (pH 8.0) through using a vacuum manifold. DNA is bound to the filter by

centrifugation at 1500 x g for 2 min or using a vacuum manifold. The filter is washed

twice with 700 µl of wash buffer (20% solution of 80mM NaCl, 8mM Tris-HCl, pH 7.5

and 80% ethanol) using centrifugation or vacuum, and then the filter is dried with

centrifugation at 1500 x g for 15 min. Residual ethanol is removed by incubating the

filter plate at room temperature for 30 min. To elute the DNA from the filter, 50 µl of

elution buffer (0.1x Tris-EDTA buffer, pH 8-9) prewarmed to 60 C is added directly to

the filter, incubated for 2 min at room temperature, and eluted into a new 2 ml

MASTERBLOCK® collection plate with centrifugation at 1500 rpm for 2 min. The

elution step is repeated once. 3) a phenol-chloroform DNA extraction protocol where

tissue is physically homogenized, as above, and lysed using the Tissue Lysis buffer

from the Monarch® Genomic DNA Purification Kit (NEB, #T3010S) with double the

volume of lysis buffer and a 1 h incubation at 56 C. Then, total lysate was placed in

Phase Lock Gel™ Light tubes (QuantaBio, #2302820) along with an equal volume of

OmniPur® Phenol:Chloroform:Isoamyl Alcohol (25:24:1, TE-saturated, pH 8.0)

solution (MilliporeSigma, Calibiochem #D05686) and then mixed by gentle inversion

for 15 minutes using a fixed speed tube rotator. After mixing, tubes were centrifuged at

maximum speed (14,000 x g) for 10 minutes, then the aqueous (top) layer was

transferred to a new phase-lock gel tube and the process repeated. DNA precipitation of

the aqueous phase was performed by adding 5M NaCl to a final concentration of 0.3M

and two volumes of room temperature absolute ethanol, inverting the tubes 20x for

thorough mixing followed by an overnight incubation at -20C. DNA was pelleted by

centrifugation at 14,000 x g for 5 min, washed twice with freshly prepared, ice cold

70% ethanol, air-dried for 15 min at room temperature, and then resuspended in 150 µl

of Elution Buffer from the Monarch® Genomic DNA kit.

DNA extract quality was assessed for quality using a NanoDrop 1000 (Thermo

Scientific) and fragment integrity using agarose gel electrophoresis. Genomic DNAs

were sequenced using a combination of paired-end sequencing on the Illumina MiSeq,

HiSeq, and Novaseq sequencing platforms (SUPPLEMENTARY TABLE 2). All raw

reads and whole genome assemblies are deposited in the Short Read Archive

(Bioproject#PRJNA1022813).

Genome assembly, ortholog extraction and phylogenetic analysis. —Raw sequencing

reads were quality-filtered and adapter-trimmed using fastP v0.20.1 (Chen et al. 2018)

with default settings. Genome assemblies were produced from quality-filtered reads

using SPAdes v3.15.0 (Bankevich et al. 2012) with five k-mer values

(k=77,85,99,111,127). From each genome, we identified 1764 highly conserved single

copy orthologs using BUSCO with the "basidiomycota odb 10" dataset. Orthogroups

that were present in less than 75% of taxa and taxa with less than 20% ortholog

recovery were removed. Retained orthologs were aligned using MAFFT v7.397 (Katoh,

Rozewicki, and Yamada 2017) with the "L-INS-I" algorithm, and maximum- likelihood

gene-trees were inferred using IQ-TREE v2.0.3 (Minh et al. 2020) with automatic

model selection in ModelFinder (Kalyaanamoorthy et al. 2017) and ultrafast

bootstrapping (BS, (Hoang et al. 2018)) with 1000 replicates. A summary coalescent

species tree was constructed from the resulting gene trees using hybrid- ASTRAL

implemented in ASTER (v1.15) (Zhang and Mirarab 2022). Branch lengths in

substitutions/site were estimated under maximum likelihood on the species tree using

the "-te" option in IQ-TREE, with a partitioned concatenated alignment of all BUSCO

genes used in species tree construction.

Gene tree comparison. —To evaluate discordance, individual gene trees were

compared using six metrics calculated in SortaDate (average bootstrap support,

clocklike branch lengths, tree length; Smith et al. 2018) and the R package 'TreeDist'

(generalized Robinson-Foulds metrics; Smith 2020, 2022). In addition to data matrix

summaries (number of taxa, alignment length), Pearson's correlations were calculated

to determine relationships between metrics.

Divergence dating. —A timetree was inferred by applying the RelTime method

(Tamura et al. 2012, 2018) conducted in MEGA11 (Stecher et al. 2020, Tamura et al.

2021) to the species tree with ML-estimated branch lengths. To reduce computational

burden, time-calibrated branch lengths were calculated using the Maximum Likelihood

(ML) method and the General Time Reversible substitution model (Nei and Kumar

2000) from two sets of 100 genes: 1) the top 100 genes with well-supported clock-like

trees determined using SortaDate (Smith et al. 2018) and 2) the top 100 genes with the

smallest generalized Robinson-Foulds ('gRF') distances to the species tree calculated

using the R package 'TreeDist.' The timetree was computed using two sets of

calibration constraints. The first included two calibrations: 1) a secondary calibration

for the stem age of the Boletaceae from 50-150 mya (Varga et al. 2019, Wu et al. 2022)

and 2) a secondary calibration for the stem age of *Boletus edulis* from 5-13 mya

(Tremble et al. 2022). The second set included the former two calibrations plus four of

the five internal calibrations using the highly supported core shifts from Varga et al.

(2019). Because many of the clades in Varga et al. were incongruent with our topology,

calibrations were selected using the most inclusive node, except for *Aureoboletus* which

could not be reconciled with our results. The Tao et al. (2020) method was used to set

minimum and maximum time boundaries on nodes for which calibration densities were

provided, and to compute confidence intervals. Outgroup node ages were not estimated

because the RelTime method uses evolutionary rates from the ingroup to calculate

divergence times and does not assume that evolutionary rates in the ingroup clade apply

to the outgroup.

Ancestral state reconstruction. — Morphological/macrochemical traits were coded

according to original descriptions and verified with microscopic analysis when traits

were ambiguous. Four traits were scored as binary or multistate characters: 1)

hymenophore anatomy (straight tubes = 0, tubular with cross walls = 1, lamellate = 2,

not applicable = 3), 2) color changes from damage (none = 0, blue = 1, black/brown = 2,

not applicable = 3), and sporocarp morphology (pileate-stipitate with exposed

hymenophore = 0, secotioid = 1, gasteroid = 2), 4) spore ornamentation (smooth = 0,

not smooth = 1) (SUPPLEMENTARY TABLE 4). The state of hymenophore anatomy

defined as "tubular with cross walls" refers to hymenophores that have tubes at two or

more lengths, giving the appearance of a primary long tube with shorter internal cross-

walls. A "not applicable" category was scored for hymenophore anatomy and color

changes from damage to accommodate secotioid/gasteroid taxa and taxa with unknown

changes, respectively. An alternative coding scheme for sporocarp morphology was also

used in an attempt to disentangle the transition between pileate-stipitate and gasteroid

morphologies from the loss of ballistospory. In this alternative coding scheme, two

binary traits were scored: stipe (present = 0, absent = 1) and ballistospory (no = 0, yes

=1). Ancestral state reconstruction of the root node of the Boletaceae was implemented

with BayesTraits V4.0.0, using the MCMC approach over 1,100,000 iterations, with a

"burn-in" of 100,000 iterations (Mead and Pagel 2022). Model convergence was

assessed with the program Tracer (v1.7.1, Rambaut et al. 2018), and determined as an

effective sample size (ESS) of >300 for all variables.

Ancestral Range Reconstruction. —Numerous analytical methods for reconstructing

historical biogeography exist, accounting for processes such as vicariance, dispersal,

and cladogenesis (Ronquist 1994; Ree et al. 2005; Landis et al. 2013). To account for

these macroevolutionary processes in our ancestral state reconstruction in the

Boletaceae, we utilized BioGeography with Bayesian (and likelihood) Evolutionary

Analysis with R Scripts ("BioGeoBEARS"; Matzke 2018). Samples were coded in two

ways: 1) Paleotropical (consisting of Africa and tropical Asia), Neotropical (South and

Central America), South Temperate (temperate Australia and New Zealand), or North

Temperate (North America, Europe, northern temperate Asia), and 2) by these floristic

regions: Holarctic including Central America, Neotropical, Chilean-Patagonian,

African, Indo-Malesian, Australian, and Novo-Zealandic (Liu et al 2023)

(SUPPLEMENTARY TABLE 3). Central America was combined with the Holarctic

region as Central American ECM fungi are mostly derived from North American

ancestors (Halling 1996). The most likely model was chosen according to AIC and

weighted AIC score calculated in BioGeoBEARS.

RESULTS

DNA sequencing, genome assembly, and ortholog extraction. —Whole genome

sequencing of 418 specimens resulted in 13,794,532 paired-end reads per specimen on

average (SUPPLEMENTAL TABLE 2). On average, genome assemblies possessed an

assembly N50 of 12.9 Kbp (thousand base-pairs), total assembly length of 61.6 Mbp

(million base-pairs), 53,972 scaffolds, and a BUSCO score of 74.7%. 34 of 418

specimens possessed BUSCO scores less than 20%, and 175 specimens possessed

BUSCO scores greater than 90% (SUPPLEMENTARY TABLE 2). After removing

specimens with poor BUSCO recovery, our final dataset included 383 Boletaceae

specimens, three outgroup taxa, and 1461 single-copy orthologs.

Phylogenetic analysis. — The summary coalescent tree resolved most nodes with

strong support (FIG. 2). Many of the groups recovered are consistent with previous

studies but now with statistical support (Dentinger et al. 2010, Nuhn et al. 2013, Wu et

al. 2014). We recognized the six subfamilies following previous authors (Wu et al.

2014), which led us to formally recognize two new subfamilies (Tremble et al. 2023).

Many of the currently accepted genera that are not mono- or oligo-typic are

polyphyletic. One notable pattern is the phylogenetic placements of the endemic

Chilean taxa, all of which were recovered as ancient lineages of similar age within four

of the subfamilies: *Gastroboletus valdivianus* in Xerocomoidae, *Boletus loyita* in

Austroboletoidae, *Boletus loyo* in Suillelloidae, and *Boletus putidus* in Boletoidae.

Gene tree comparison. —Average bootstrap support had the highest positive

correlation with the number of taxa present (Pearson's coefficient = 0.76), and weak to

moderate negative correlations with alignment length (Pearson's coefficient = -0.17),

clocklike branch lengths (Pearson's coefficient = -0.13), and total tree length (Pearson's

coefficient = -0.20). Generalized Robinson-Foulds distances were weakly to moderately

negatively correlated with number of taxa (Pearson's coefficient = -0.33), total length

(Pearson's coefficient = -0.30), and clocklike branch lengths (Pearson's coefficient = -

0.13), and weakly to moderately positively correlated with alignment length (Pearson's

coefficient= 0.29) and average bootstrap support (Pearson's coefficient = 0.17).

Clocklike branch lengths and total tree length were weakly positively correlated

(Pearson's coefficient= 0.15).

Divergence dating. — Using the two- and six-calibration sets the following ages were

estimated (FIGS. 2, 3). Stem ages for the Boletaceae were estimated at 138-139 mya

and 77 mya. The crown age of the Boletaceae and origin of Chalciporoideae was

estimated at 103-105 mya and 63 mya (49-77 mya). The stem age of the

Phylloboletelloideae was estimated at 83-87 mya and 58 mya (49-77 mya)., The

radiation of the remaining subfamilies was estimated to have occurred between 61 and

51 mya. The origin of *Boletus* sensu stricto (i.e. "true porcini") was estimated at 38 mya

and 35 mya (29-42 mya), and its diversification was estimated at 29-30 mya and 26 mya

(20-34 mya).

Ancestral state reconstruction. —Ancestral state reconstruction with Bayestraits

achieved strong convergence (all variables with ESS > 300) and recovered the most

likely ancestor of the Boletaceae to have a pileate-stipitate, lamellate sporocarp with

ornamented basidiospores. The highest transition rates were observed in changes from

secotioid to gasteroid and from secotioid to pileate-stipitate morphologies. Secotioid

morphology appeared to be a relatively transient evolutionary state (TABLE 1). High

transition rates were also found to occur from gasteroid to pileate-stipitate morphology,

supporting evidence for the reversibility of gasteromycetization.

Ancestral range reconstruction. —Ancestral distribution reconstruction recovered a

likely Paleotropical origin of the Boletaceae (DEC+J model chosen with lowest AIC

and AICc for both coding sets), with two major descendant radiations originating in the

Paleotropics (Africa and Asia) and Neotropics (FIG. 3). In addition, we found evidence

for multiple diversification events spurred by the separation of Gondawana (FIG. 3).

Gondwanan separation occurred in two predominant phases: Phase 1, which involved

the separation of Southern South America, Southern Africa, Australia-Antarctica, and

Madagascar-India, beginning approximately 180 mya and largely completed by 120

mya (Jokat et al. 2003) and Phase 2, involving the separation of South America and

Africa, which was completed 80 mya (Reguero and Goin 2021). At the split between

the Austroboletoidae and Suillelloidae (FIG. 3), we estimated a putative Phase 1

Gondwanan separation to have occurred 150-120 mya, which led to rapid formation of

South Temperate, Neotropical, and Paleotropical lineages. Later, around 90-70 mya, at

least five putative Phase 2 separation events occurred, splitting the Paleotropical and

Neotropical lineages.

In our four category paleo-region coding set, the Boletaceae ancestor was

equally likely to be Neotropical or Paleotropical. However, the subsequent node that

leads to the rest of the Boletaceae (excluding Phyllobolletoidae and Chalciporoideae)

was well-supported as Paleotropical, as were all immediate descendent nodes. Our

coding of the Chalciporoideae and the single *Phylloboletellus* specimen likely had a

strong influence on deep-node ancestral range reconstructions. The backbone nodes of

the Boletaceae excluding Phylloboletoidae and Chalciporoideae were estimated as

Asian in origin, corroborating the four-category analysis, though with less confidence.

Migrations between phytogeographic regions were dominated by dispersals between the

Indo-Malesian and Holarctic regions (FIGS. 3,4).

DISCUSSION

The fully resolved phylogeny supports the recognition of eight subfamilies, including

the newly defined Phylloboletelloideae and Suillelloideae (Tremble et al. 2023). The

subfamilial relationships were resolved for the first time. The Chalciporoideae was

recovered as the earliest diverging group in the Boletaceae, a relationship previously

noted by Wu et al. (2014). The enigmatic *P. chloephorus* (Singer and Diglio 1952) was

the next lineage to branch off before the radiation that gave rise to the six additional

subfamilies. Previous studies have placed *Pseudoboletus parasiticus* in a position

similar to that of *P. chloephorus* in our study (Nuhn et al. 2013, Wu et al. 2014, Sato

and Toju 2019, Caiafa and Smith 2022). We were, however, unable to include a

representative of *P. parasiticus* in our study so cannot corroborate its putative

phylogenetic position.

The tree topology has intriguing implications for the role of ecological

transitions in Boletaceae diversification. Members of the earliest-diverging

Chalciporoideae species can be facultatively ECM, saprotrophic or mycoparasitic

(Caiafa and Smith 2022) and *P. chloephorus* may not be ECM given its occurrence in

non-ectotrophic forests (Singer and Diglio 1952). Moreover, *P. parasiticus* and other

Pseudoboletus spp. produce sporocarps directly attached to gasteroid *Scleroderma* and

Astraeus and are assumed to be mycoparasites (Raidl 1997; Binder and Hibbet, 2006;

Nuhn et al. 2013). Altogether, the basal position of these early diverging groups

suggests that the ancestor of the Boletaceae was likely saprotrophic and not ECM. This

possibility is in line with the results of Sato and Toju (2019) which indicated that the

ECM habit emerged with the origin of the six derived Boletaceae subfamilies. Genomic

changes coinciding with the emergence of an obligate ECM habit further support the

view that this nutritional shift has profoundly impacted Boletaceae (Wu et al. 2022).

Many taxonomic changes in the Boletaceae have been proposed in recent years.

In particular, new genera have been erected for phylogenetically unresolved lineages.

Many of these new genera are mono- or oligo-typic (composed of one or few species)

(e.g. Castellano et al. 2016; Chakraborty et al. 2015; Henkel et al. 2016; Halling et al.

2023). This proliferation of new genera has made it difficult or impossible to recognize

inclusive groupings without requiring knowledge of the species. On the bright side, the

strong nodal support throughout our phylogeny sets the stage for a new, comprehensive

and stable generic-level taxonomy. We will address this in subsequent works when all

recently described genera are represented.

Our biogeographic and divergence dating analyses support a Gondwanan origin

of the Boletaceae. Subsequent divergence was likely facilitated by continental drift-

based vicariance events and possible long-distance dispersals. Other recent studies have

shown that lineages of ECM fungi originated in Gondwana or more recently in

paleotropical regions (Hosaka et al. 2008, Matheny et al. 2009, Dentinger et al. 2010,

Ryberg and Matheny 2011, Kennedy et al. 2012, Sanchez-Ramirez et al. 2015, Han et

al. 2018; Hackel et al. 2022; Codjia et al. 2023). Phytogeographic endemism was

implied with our biogeographic reconstruction. The strongest migrations occurred

recently over the past 50 my between the Indo-Malesian and Holarctic regions. We

acknowledge the difficulty to determining origins and dispersal events in the absence of

fossils or other corroborating evidence. Nonetheless our study and others suggest that

vicariance may have played a strong role in the distribution of ECM fungal taxa, despite

the long-distance dispersal capacity of airborne spores (Matheny et al. 2009, Peay et al.

2010, Peay and Matheny 2016). Conversely, pure vicariance cannot explain the close

phylogenetic relationships seen between distantly disjunct taxa. Long-distance

dispersals may have occurred, albeit rarely. While long-distance dispersal is

demonstrably possible in the Boletaceae and other ECM lineages the likelihood of its

frequent occurrence is low (Geml et al. 2012; Hackel et al. 2022; Tremble et al. 2022).

Most basidiospores do not travel far from the parental sporocarp (Galante et al. 2011),

and the probability of two airborne basidiospores landing in close-enough proximity to

mate is negatively correlated with increasing distance from sporocarps (Peay et al.

2012; Golan and Pringle 2017). Such improbabilities notwithstanding, our

biogeographic patterns are consistent with episodic long-distance dispersal, possibly by

aerial dispersal of basidiospores, spores vectored by migrating animals (e.g., Elliot et al.

2019) or somatic mycelia on rafting vegetation (Thiel et al. 2005).

Our biogeographic reconstructions are consistent with the "Southern Route to

Asia" hypothesis (Wilf et al. 2019). This idea proposes that ECM Fagaceae and their

symbiotic fungi originated in Gondwana and were carried on Australia north to Asia. In

this scenario the Gondwanan ECM habitat tracked climatic niches on the desertifying

continent northward as the Australian plate collided with the Pacific plate. A relictual

ECM community remained in a newly isolated New Guinea and subsequently spread

northwest along the montane Australasian archipelago, followed by dispersal into

continental Asia. Many of the dispersal events we found between Indo-Malesia and

other regions, especially the Holarctic, are inferred within the last 20 my, coincident

with the late-Oligocene collision of Australia with the Pacific plate (Hall 2011). As

suggested by Halling et al. (2011) recent Boletaceae migrations likely occurred across

the Australasian archipelago and are corroborated by our inferred recent regional

dispersal events.

Biogeographic reconstructions are highly sensitive to taxon sampling and our

dataset is not immune to equivocal reconstructions. For example, in both coding

schemes the ancestral node of the Chalciporoideae had the highest probabilities of a

North Temperate and North American origin, respectively. However, with no

Chalciporoideae samples from Asia, Africa or Australia/New Zealand in our study, their

potential impacts on the reconstruction are unknown. Such sampling gaps

notwithstanding, we have the most geographically comprehensive sampling for

Boletaceae ever compiled and provide the first opportunity to examine global-scale

biogeographic patterns. Insights into the evolution of the Boletaceae are revealed for the

first time, despite mild uncertainty at a minority of nodes.

The evolutionary origins of distinctive regional Boletaceae assemblages have

long been a mystery (Horak 1977). For example, the endemic Boletaceae of Chile and

Argentina have not been included in previous phylogenetic studies, and their

morphology-based affinities have been inconclusive (Horak 1977). The recovery of

several Chilean species as ancient lineages in four of the subfamilies implies that they

have survived in isolation without speciating for millions of years. The closest relatives

of these Chilean boletes occurred in geographic regions as disjunct as North America,

lowland tropical northern South America, and Australia. *Boletus loyita* and *G.*

valdivianus were most closely related to extant Australian taxa, suggesting an origin

prior to final Gondwanan disarticulation (Reguero and Goin 2021). Close relationships

between southern Gondwanan Australian and southern South American taxa have been

documented elsewhere (Feng et al. 2017). In all likelihood Chilean boletes arose in

Gondwana, separated from their sister lineages during Gondwanan disarticulation, and

underwent no subsequent speciation for tens of millions of years.

Ancestral range reconstruction recovered an Asian origin of the core, "true

porcini" genus *Boletus* s. str. as previously suggested (Feng et al. 2012). However, we

cannot entirely rule out an African origin. The Central African endemic *Boletus*

alliaceus was recovered here as a sister taxon to *Boletus* s. str., and a similar

relationship was found for the recently described *Paxilloboletus africanus* (Badou et al.

2022). Furthermore, we estimated the origin of *Boletus* s. str to be 40 mya, which may

indicate why the sister lineages to *Boletus* s. str. are endemic to Africa. India had

separated from Africa and Madagascar ~120 mya (Reguero and Goin 2021), and at 40

mya was already colliding with Asia (Aitchison et al. 2007; Hu et al. 2016). If *B.*

alliaceus and *P. africanus* are indeed sister lineages of *Boletus* s. str., then the arrival

and subsequent diversification of true porcini in Asia must have been a dispersal event,

because the separation of India from mainland Africa ~180-170 mya (Hankel 1994)

occurred long before our estimated age of the *Boletus* s.s. ancestor (~40 mya). Even if a

more recent ancestor existed in Madagascar or the Seychelles, the separation of India

from these landmasses at ~90 mya (Storey et al. 1995) and ~64 mya (Norton and Sclater

1979), respectively, is still much older than our current age estimates for true porcini.

Furthermore, most or all ECM fungi in Madagascar appear to have arrived on the island

through dispersal its separation from Africa (Rivas-Ferreiro et al. 2023), so dispersal is

the most plausible mechanism unless ancient Malagasy relict taxa are discovered. In the

current study currently undescribed species of *Boletus* s. str. were recovered from

Taiwan, Malaysian Borneo, and the Gulf Coast of the US, indicating that much more

diversity exists in the genus. To sort out the origins and full diversity of *Boletus* s. str.

more mycological exploration and whole genome sequencing are needed. In particular,

discovery and analysis of true porcini basal lineages from India and Africa could shed

further light on the origin of this charismatic group.

Divergence dating estimated the origin of the Boletaceae at 138-139 mya, and

ancestral range reconstructions suggested it may be even older. Ranking genes with

different metrics had little impact on divergence date estimates. However, the two

calibrations sets gave very different estimates for most nodes. The estimated origin

dates of the Boletaceae using the Varga et al. (2019) calibrations were almost half those

of the two-calibration set. It is difficult to interpret these wildly different divergence

dates given the lack of fossil evidence. However, the dates estimated using the Varga et

al. (2019) calibrations are suspect due to extensive topological incongruence of their

phylogenetic trees with ours. Our older divergence estimate corroborates the results of

He et al. (2020) and our internal dates correspond with other results, such as the ~48

mya origin of the *Strobilomyces* group (Han et al. 2018). Our older divergence estimate

is also in line with the origin of ECM Pinaceae in the early Cretaceous (Brundrett and

Tedersoo 2018). Therefore, we consider the older estimate to be more accurate.

In our ancestral range reconstruction analysis, we found evidence of multiple

diversification events that may have been initiated by Gondwanan breakup. The first

phase of the Gondwanan separation postulated by Jokat et al. (2003) correlates well

with our estimated origin of the Boletaceae and indicates that the family was diverse

and widely distributed by 120 mya, substantially older than the estimated age from our

divergence dating analysis but within the 95% confidence interval. Our dates are at best

coarse estimates based on fossil-free secondary calibrations. However, the phylogenetic

pattern of vicariance that parallels the breakup of Gondwana is compelling and offers

corroborating evidence that our estimated ages may in fact be too young. In any case,

our divergence estimates suggest that the Boletaceae originated and diversified within

the early to late Cretaceous period. During this time global climate was warm and wet

(Hay and Floegal 2012), gymnosperms and subsequently angiosperms diversified (Crisp

and Cook 2011), and the supercontinental land masses broke apart (Jokat et al. 2003).

The Boletaceae mostly consists of species that form ECM assocations, but

emerging evidence suggests the ancestor may have had mycoparasitic capacity. The

gain of obligate ECM ecology in the six most-derived subfamilies likely occurred after

their divergence from the Phylloboletelloidae. *Phylloboletellus chloephorus* is from

non-ECM dominated habitats. The Chalciporoideae have not been definitively shown to

form ECM associations but do have saprotrophic or mycoparasitic capacities (Caiafa et

al. 2022). While mycoparasitic *Pseudoboletus* species have also been recovered as

early-diverging Boletaceae lineages (Nuhn et al. 2013, Sato and Toju 2019, Cortes-

Perez et al. 2023) and may have close affinities with lamellate *Phylloboletellus*, we

were not able to evaluate *Pseudoboletus* in this study. A more thorough investigation of

the ecology of early-diverging Boletaceae is needed to test this "mycoparasitic origin"

hypothesis.

A longest standing debate among Boletaceae systematists has centered on the

utility of morphological traits for defining natural genera. Basidiospore color,

ornamentation, and hymenophore arrangement were long emphasized in this respect.

These features have been used as evidence for dividing the Boletaceae into multiple

genera (e.g., Singer 1945a,b; 1947; Pegler and Young 1981) or treating nearly all

Boletaceae as a single genus (Corner 1972). More recently, hyphal anatomies of

sporocarp structures and pigment chemistry have been emphasized (e.g. Binder et al.

2002; Šutara 2005). Yet, despite the morphological and chemical variability in the

Boletaceae, the family is typified by the 'bolete' macromorphology of fleshy, pileate-

stipitate sporocarps with tubular hymenophores. In addition, type of basidiospore

ornamentation has been long considered to be a genus-unifying trait in genera such as

Strobilomyces, *Boletellus*, and *Austroboletus* (Berkeley 1851, Murrill 1909, Corner

1972, Pegler and Young 1981, Wolfe 1980). Despite such traditional views, our

ancestral state reconstructions suggest that the ancestor of the Boletaceae had a pileate-

stipitate sporocarp with a lamellate hymenophore and ornamented basidiospores. This

likely resulted from the basal phylogenetic position of the lamellate *P. chloephorus*

(Singer and Diglio 1952).

The evolution of sequestrate morphologies has long been thought to be

irreversible (Thiers 1984). The transition from pileate-stipitate to secotioid to gasteroid

morphology involves enclosure of the hymenophore and loss of ballistospory, which are

unlikely to be regained once lost (Hibbett et al. 1997, Hibbett 2004, Sanchez-Garcia et

al. 2020). However, prior studies have not fully rejected the hypothesis that

gasteromycetization is irreversible (Hibbett 2004; Wilson et al. 2011; Sanchez-Garcia et

al. 2020). Our ancestral state reconstructions under two coding schemes strongly

supported transition to and from gasteroid forms. This suggests that gains or losses of

ballistospory may also be reversible conditions. In our analyses, transitions between

pileate-stipitate with exposed hymenophore, secotioid, and gasteroid forms suggest that

the secotioid condition is intermediate, as indicated by the equal rates of transition to

gasteroid and pileate-stipitate forms. Moreover, inferred transition rates from pileate-

stipitate to both secotioid and gasteroid forms were almost zero, suggesting that the

secotioid condition is evolutionarily unstable. This result contrasts with the stability of

Cortinarius secotioid taxa suggested by Peinter et al. (2001). A plausible hypothesis is

that a membranous partial veil that covers the hymenophore at early stages of sporocarp

development may predispose it to gasteromycetization. However, although we did not

test this explicitly, based on the phylogenetic distribution of taxa with membranous

veils (e.g., *Pulveroboletus*, *Veloporphyrellus*), it is clear that the membranous partial

veil is a convergent trait not closely associated with secotioid or gasteroid

morphologies. A possible exception to this is *Veloboletus limbatus*, which is most

closely related to the gasteroid *Gastroboletus valdivianus*, although their common

ancestor was estimated at ~40 mya.

Transitions between hymenophore organization parallels the sporocarp

morphologies. The "tubular with crosswalls" hymenophore condition appears to be

intermediate between lamellate and tubulate forms, with transitions to tubes being 1.3-

5.8 times greater than the opposite transitions. However, the greatest transition rate was

from lamellate directly to tubulate forms, indicating that an intermediate morphology

may not be necessary or, like the secotioid condition, may be evolutionarily unstable.

Our data provide compelling evidence that lamellate hymenophores, color changes, and

gasteromycetization have evolved multiple times in the Boletaceae and are reversible.

SUMMARY

The Boletaceae underwent a rapid radiation and subsequent long period of phylogenetic

instability. These issues had long prevented accurate assessment and analysis of trait

evolution and conclusive generic-level taxonomic frameworks in the family. Previous

molecular phylogenetic studies were based on a limited taxon sampling and a few loci

and generated trees with many short branches and little deep-node support. This study

provides the first Boletaceae phylogeny with strong support at deep nodes, based on a

massive dataset of 1461 single copy genes from 383 genomes sampled from

taxonomically and geographically comprehensive specimens. Our analyses indicated

that the Boletaceae likely arose before Gondwanan breakup and that present-day

distributions are partly due to vicariance. Long-distance dispersal could not be ruled out

for some current distributions. Morphological traits resulted from convergence and

frequent reversals, which, along with rapid radiation, have long confounded attempts to

achieve a natural intrafamilial classification. This study provides new genomic data and

a solid phylogenetic framework that will enable a renewed foundational taxonomy as

well as deeper analysis of trait evolution.

AKNOWLEDGEMENTS

This work was supported by NSF-DEB awards DEB-2114785 to BD and DEB-0918591

to TWH and DEB-1556338 to TWH and BD; National Geographic Society's

Committee for Research and Exploration grants (6679-99, 7435-03, and 8481-08 to

TWH). JMM was supported by grants from the ROM Governors and the Natural

Sciences and Engineering Research Council of Canada, Discovery Program. We would

like to thank Dr. Roy Halling for the numerous specimens provided.

CONFLICTS OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

LITERATURE CITED

Aitchison JC, Ali JR, and Davis AM. 2007. When and where did India and Asia

collide? Journal of Geophysical Research: Solid Earth. 112:B5

Arora D. 2008. California porcini: three new taxa, observations on their harvest, and the

tragedy of no commons. Economic Botany. 62:356–375.

Badou SA, Furneaux B, De Kesel A, Khan FK, Houdanon RD, Ryberg M, Yorou NS.

2022. *Paxilloboletus* gen. nov., a new lamellate bolete genus from tropical Africa.

Mycological Progress. 21:243–256.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, and Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and

its applications to single-cell sequencing. Journal of Computational Biology. 19:455–

477.

Berkeley MJ. 1851. Decades of fungi. Decade XXXIV. Sikkim-Himalayan fungi

collected by Dr. Hooker. Hooker's Journal of Botany and Kew Gardens Miscellany.

3:77—84.

Binder M, Bresinsky A. 2002. *Retiboletus*, a new genus for a species-complex in the

Boletaceae producing retipolides. Feddes Repertorium. 113:30–40.

Binder M, Hibbet DS. 2006. Molecular systematics and biological diversification of

Boletales. Mycologia. 98:971–981.

Brundrett MC, Tedersoo L. Evolutionary history of mycorrhizal symbioses and global

host plant diversity. New Phytologist. 220:1108–1115.

Bruns TD, Palmer JD. 1989. Evolution of mushroom mitochondrial DNA: *Suillus* and

related genera. Journal of Molecular Evolution. 28:349–362.

Bruns TD, Vilgalys R, Barns SM, Gonzalez D, Hibbett DS, Lane DJ, Simon L, Stickel

S, Szaro TM, Weisburg WG, Sogin ML. 1992. Evolutionary relationships within the

fungi: analyses of nuclear small subunit rRNA sequences. Molecular Phylogenetics and

Evolution. 1:231—241.

Caiafa MV, Smith ME. 2022. Polyphyly, asexual reproduction and dual trophic mode in

Buchwaldoboletus. Fungal Ecology. 56:101—141.

Castellano MA, Elliott TF, Truong F, Séné O, Dentinger BTM, Henkel TW. 2016.

Kombocles bakaiana gen. sp. nov. (Boletaceae), a new sequestrate fungus from

Cameroon. IMA Fungus. 7: 239–245.

Chakraborty D, Das K. 2015. A new generic record of Boletaceae for Indian mycobiota.

Current Research in Environmental & Applied Mycology. 5:138–144.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics. 34:884–890.

Codjia JEI, Sánchez-Ramírez S, Ndolo Ebika ST, Wu G, Margaritescu S, Komura DL,

Oliveira JJS, Ryberg M, Tulloss RE, Yorou NS, Moncalvo JM, Yang ZL. 2023.

Historical biogeography and diversification of ringless *Amanita* (section *Vaginatae*)

support an African origin and suggest niche conservatism in the Americas. Molecular

Phylogenetics and Evolution. 178:107644.

Corner EJH. 1972. Boletus in Malaysia. The Botanic Gardens Singapore. The

Government Printer. 263pp

Cortés-Pérez A, Ramirez-Guillen F, Garcia-Jimenez J, Ramirez-Cruz V, Villalobos-

Arambula AR, Reynaga DMB, Guzman-Davalos L. 2023. *Pseudoboletus silvaticus*

(Boletaceae, Basidiomycota), a new species from Mexico. Phytotaxa. 589:27–38.

Crisp MD and Cook LG. 2011. Cenozoic extinctions account for the low diversity of

extant gymnosperms compared with angiosperms. New Phytologist. 192:997—1009.

Das K, Chakraborty D, Baghela A, Singh SK, Dentinger BTM. 2015. *Boletus*

lakhanpalii, a new species in Boletaceae from Sikkim India with uncertain phylogenetic

placement. Sydowia. 67:11–19.

Das K, Chakraborty D, Baghela A, Singh SK, Dentinger BTM. 2016. New species of

xerocomoid boletes (Boletaceae) from Himalayan India based on morphological and

molecular evidence. Mycologia. 108:753–764.

Dentinger BTM, Ammirati JF, Both EE, Desjardin DE, Halling RE, Henkel TW,

Moreau PA, Nagasawa E, Soytong K, Taylor AF, Watling R, Moncalvo JM,

McLaughlin DJ. 2010. Molecular phylogenetics of porcini mushrooms (*Boletus* section

Boletus). Molecular Phylogenetics and Evolution. 57:1276–1292.

Dentinger BTM, Gaya E, O'Brien H, Suz LM, Lachlan R, Diaz-Valderrama JR, Koch

RA, Aime MC. 2016. Tales from the crypt: genome mining from fungarium specimens

improves resolution of the mushroom tree of life. Biological Journal of the Linnean

Society. 117:11–32.

Dentinger BTM, Suz LM. 2014. What's for dinner? Undescribed species of porcini in a

commercial packet. PeerJ. 2:e570.

Drehmel D, James T, Vilgalys R. 2008. Molecular phylogeny and biodiversity of the

boletes. Fungi. 1:17—23.

Elliott TF, Jusino MA, Trappe JM, Lepp H, Ballard GA, Bruhl JJ, Verns K. 2019. A

global review of the ecological significance of symbiotic associations between birds and

fungi. Fungal Diversity. 98:161—194.

Feng F, Xu J, Wu G, Zeng NK, Li YC, Tolgor B, Kost GW, Yang ZL. 2012. DNA

sequence analyses reveal abundant diversity, endemism and evidence for Asian origin

of the porcini mushrooms. PLOS ONE. 7:e37567.

Feng YJ, Blackburn DC, Liang D, Hillis DM, Wake DB, Zhang P. 2017.

Phylogenomics reveals rapid, simultaneous diversification of three major clades of

Gondwanan frogs at the Cretaceous–Paleogene boundary. Proceedings of the National

Academy of Sciences. 114:E5864—E5870.

Fulgenzi TD, Halling RE, Henkel TW. 2010. *Fistulinella cinereoalba* sp. nov. and new

distribution records for *Austroboletus* from Guyana. Mycologia. 102:224–232.

Fulgenzi TD, Henkel TW, Halling RE. 2007. *Tylopilus orsonianus* sp. nov. and

Tylopilus eximius from Guyana. Mycologia. 99:622–627.

Fulgenzi TD., Mayor JR, Henkel TW, Halling RE. 2008. New species of *Boletellus*

from Guyana. Mycologia. 100:490–495.

Futuyma DJ. 1998. Evolutionary Biology. Quarterly Review of Biology. 73.

Galante TE, Horton TR, Swaney DP. 2011. 95% of basidiospores fall within 1 m of the

cap: a field-and modeling-based study. Mycologia. 103:1175–1183.

Geml , Timling I, Robinson CH, Lennon N, Nusbaum CH, Brochmann C, Noordeloos

ME, Taylor DL. 2012. An arctic community of symbiotic fungi assembled by long-

distance dispersers: phylogenetic diversity of ectomycorrhizal basidiomycetes in

Svalbard based on soil and sporocarp DNA. Journal of Biogeography. 39:74—88.

Golan JJ and Pringle A. 2017. Long-distance dispersal of fungi. Microbiology

Spectrum. 5:5—4.

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A,

Zhao X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2013.

MycoCosm portal: gearing up for 1000 fungal genomes. Nucleic Acids Research.

42:699–704

Grubisha LC, Trappe JM, Molina R, Spatafora JW. 2001. Biology of the

ectomycorrhizal genus *Rhizopogon*. V. Phylogenetic relationships in the Boletales

inferred from LSU rDNA sequences. Mycologia. 93:82–89.

Hackel J, Henkel TW, Moreau PA, De Crop E, Verbeken A, Sa M, Buyck B, Neves

MA, Vasco-Palacios A, Wartchow F, Schimann H, Carriconde F, Garnica S,

Courtecuisse R, Gardes M, Manzi S, Louisanna E, Roy M. 2022. Biogeographic history

of a large of ectomycorrhizal fungi, the Russulaceae, in the Neotropics and adjacent

regions. New Phytologist. 236:698—713.

Halling RE. 1996. Boletaceae (Agaricales): Latitudinal biodiversity and biological

interactions in Costa Rica and Colombia. Revista de Biología Tropical. 44:111–114.

Halling RE, Desjardin DE, Fechner N, Arora D, Soytong K, Dentinger BTM. 2014.

New Porcini (Boletus sect. Boletus) from Australia and Thailand. Mycologia. 106:830–

834.

Halling RE, Fechner NA, Holmes G, Davoodian N. 2023. *Kagaria* (Boletaceae,

Boletoideae) gen. nov. in Australia: Neither a *Tylopilus* nor a *Porphyrellus*. Fungal

Systematics and Evolution. 12:31—45.

Halling RE, Fechner N, Nuhn M, Osmundson T, Soytong K, Arora D, Binder M,

Hibbett D. 2015. Evolutionary relationships of *Heimioporus* and *Boletellus* (Boletales),

with an emphasis on Australian taxa including new species and new combinations in

Aureoboletus, *Hemileccinum* and *Xerocomus*. Australian Systematic Botany. 28:1–22.

Halling RE, Osmundson TW, Neves MA. 2006. *Austroboletus mutabilis* from northern

Queensland. Muelleria. 24:31–36.

Halling RE, Nuhn M, Osmundson T, Fechner N, Trappe JM, Soytong K, Arora D,

Hibbett DS, Binder M. 2012. Affinities of the *Boletus chromapes* group to *Royoungia*

and the description of two new genera, *Harrya* and *Australopilus*. Australian Systematic

Botany. 25:418–431.

Halling RE, Osmundson TE, Neves MA. 2008. Pacific boletes: implications for

biogeographic relationships. Mycological Research. 112:437–447.

He MQ, Zhao RL, Hyde KD, Begerow D, Kemler M, Yurkov A, McKenzie EHC,

Raspé O, Kakishima M, Sánchez-Ramírez S, Vellinga EC, Halling RE, Papp V,

Zmitrovich IV, Buyck B, Ertz D, Wijayawardene NN, Cui BK, Schoutteten N, Liu XZ,

Li TH, Yao YJ, Zhu XY, Liu AQ, Li GJ, Zhang MZ, Ling ZL, Cao B, Antonín V,

Boekhout T, da Silva BDB, De Crop E, Decock C, Dima B, Dutta AK, Fell JW, Geml J,

Ghobad-Nejhad M, Giachini AJ, Gibertoni TB, Gorjón SP, Haelewaters D, He SH,

Hodkinson BP, Horak E, Hoshino T, Justo A, Lim YW, Menolli N, Mešić A,

Moncalvo JM, Mueller GM, Nagy LG, Nilsson RH, Noordeloos M, Nuytinck J, Orihara

T, Ratchadawan C, Rajchenberg M, Silva-Filho AGS, Sulzbacher MA, Tkalčec Z,

Valenzuela R, Verbeken A, Vizzini A, Wartchow F, Wei TZ, Weiß M, Zhao CL, Kirk

PM. 2020. Notes, outline and divergence times of Basidiomycota. Fungal Diversity.

99:105—367.

Han LH, Feng B, Wu G, Halling RE, Buyck B, Yorou NS, Ebika STN, Yang ZL. 2018.

African origin and global distribution patterns: Evidence inferred from phylogenetic and

biogeographical analyses of ectomycorrhizal fungal genus *Strobilomyces*. Journal of

Biogeography. 45:201—212.

Hay WW and Floegel S. 2012. New thoughts about the Cretaceous climate and oceans.

Earth-Science Reviews. 115:262—272.

Heinemann P. 1951. Champignons recoltes au Congo belge par Madame M. Goossens-

Fontana I. Boletineae. Bulletin du Jardin botanique de l'État a Bruxelles. 21:223–346.

Henkel TW, Aime MC, Chin M, Miller SL, Vigalys R, Smith ME. 2012.

Ectomycorrhizal fungal sporocarp diversity and discovery of new taxa in *Dicymbe*

monodominant forests of the Guiana Shield. Biodiversity and Conservation. 21:2195—

2220.

Henkel TW, Obase K, Husbands D, Uehling JK, Bonito G, Aime MC, Smith ME. 2016.

New Boletaceae taxa from Guyana: *Binderoboletus* segoi gen. and sp. nov.,

Guyanaporus albipodus gen. and sp. nov., *Singerocomus rubriflavus* gen. and sp. nov.,

and a new combination for *Xerocomus inundabilis*. Mycologia. 108:157–173.

Hibbett DS. 2004. Trends in morphological evolution in Homobasidiomycetes inferred

using maximum likelihood: a comparison of binary and multistate approaches.

Systematic Biology. 53:889–903.

Hibbett DS, Pine EM, Langer E, Langer G, Donoghue MJ. 1997. Evolution of gilled

mushrooms and puffballs inferred from ribosomal DNA sequences. Proceedings of the

National Academy of Sciences. 94:12002–12006.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2:

Improving the ultrafast bootstrap approximation. Molecular Biology and Evolution.

35:518–522.

Horak E. 1977. New and rare boletes from Chile, *Nothofagus chilenos*, Ga*stroboletus*

valdivians, *Boletus loyita*, *Boletus putidus*. Boletín de la Sociedad Argentina de

Botánica.

Hosaka K, Castellano MA, Spatafora JW. 2008. Biogeography of Hysterangiales

(Phallomycetidae, Basidiomycota. Mycological Research. Phylogeography and

Biogeography of Fungi. 112:448–462.

Hu X, Garxanti E, Wang J, Huang W, An W, Webb A. 2016. The timing of India-Asia

collision onset–facts, theories, controversies. Earth Science Reviews. 160:264—299.

Husbands DR, Henkel TW, Bonito G, Vilgalys R, Smith ME. 2013. New species of

Xerocomus (Boletales) from the Guiana Shield, with notes on their mycorrhizal status

and fruiting occurrence. Mycologia. 105:422–435.

Jokat W, Boebel T, Konig M, Meyer U. 2003. Timing and geometry of early Gondwana

breakup. Journal of Geophysical Research: Solid Earth. 108:2156—2202.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods.

14:587–589.

Katoh K, Rozewicki J, Yamada KD. 2017. MAFFT online service: multiple sequence

alignment, interactive sequence choice and visualization. Briefings in Bioinformatics.

20:1160—1166.

Kennedy PG, Matheny PB, Ryberg KM, Henkel TW, Uehling JK, Smith ME. 2012.

Scaling up: examining the macroecology of ectomycorrhizal fungi. Molecular Ecology.

21:4151–4154.

Kohler A, Kuo A, Naga LG, Morin E, Berry KW, Buscot F, Canback B, Choi C,

Cichocki N, Clum A, Colpaert J, Copeland A, Costa MD, Dore J, Floudas D, Gay G,

Girlanda M, Henrissat B, Herrmann S, Hess J, Hogberg N, Johansson T, Khouja HR,

LaButti K, Lahrmann U, Levasseur A, Lindquist EA, Lipzen A, Marmeisse R, Martino

E, Murat C, Ngan CY, Nehls U, Plett JM, Pringle A, Ohm RA, Perotto S, Peter M,

Riley R, Rineau F, Ruytinx J, Salamov A, Shah F, Sun H, Tarkka M, Tritt A, Veneault-

Fourrey C, Zuccaro A, Mycorrhizal Genomics Initiative Consortium, Tunlid A,

Grigoriev IG, Hibbett DS, Martin F. 2015. Convergent losses of decay mechanisms and

rapid turnover of symbiosis genes in mycorrhizal mutualists. Nature Genetics. 47:410–

415.

Landis MJ, Matzke NJ, Moore BR, Huelsenbeck JP. 2013. Bayesian analysis of

biogeography when the number of areas is large. Systematic Biology. 62:789–804.

Liimatainen K, Kim JT, Pokorny L, Kirk PM, Dentinger BTM, Niskanen T. 2022.

Taming the beast: a revised classification of Cortinariaceae based on genomic data.

Fungal Diversity. 112:89–170.

Liu Y, Xu X, Dimitrov D, Pellissier L, Borregaard MK, Shrestha N, Su X, Luo A,

Zimmermann NE, Rahbek C, Wang Z. 2023. An updated floristic map of the world.

Nature Communications. 14:2990.

Magnago AC, Neves MA, da Silveira RMB. 2017. *Fistulinella ruschii*, sp. nov., and a

new record of *Fistulinella campinaranae var. scrobiculata* for the Atlantic Forest,

Brazil. Mycologia. 109:1003—1013.

Matheny PB, Aime MC, Bougher NL, Buyck B, Desjardin DE, Horak E, Kropp BR,

Lodge DJ, Soytong K, Trappe JM, Hibbett DS. 2009. Out of the Palaeotropics?

Historical biogeography and diversification of the cosmopolitan ectomycorrhizal

mushroom family Inocybaceae. Journal of Biogeography. 36:577–592.

Meade A and Pagel M. 2022. Ancestral state reconstruction using BayesTraits.

Environmental Microbial Evolution: Methods and Protocols. New York, NY: Springer

US. 255-266.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. Molecular Biology and Evolution. 37:1530–1534.

Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, Morin E,

Andreopoulos B, Barry KW, Bonito G, Buée M, Carver A, Chen C, Cichocki N, Clum

A, Culley D, Crous PW, Fauchery L, Girlanda M, Hayes RD, Kéri Z, LaButti K, Lipzen

A, Lombard V, Magnuson J, Maillard F, Murat C, Nolan M, Ohm RA, Pangilinan J,

Pereira MF, Perotto S, Peter M, Pfister S, Riley R, Sitrit Y, Stielow JB, Szöllősi G,

Žifčáková L, Štursová M, Spatafora JW, Tedersoo L, Vaario LM, Yamada A, Yan M,

Wang P, Xu J, Bruns T, Baldrian P, Vilgalys R, Dunand C, Henrissa Bt, Grigoriev IV,

Hibbett D, Nagy LG, Martin FM. 2020. Large-scale genome sequencing of mycorrhizal

fungi provides insights into the early evolution of symbiotic traits. Nature

Communications. 11:5125.

Murrill WA. 1909. The Boletaceae of North America—I. Mycologia. 1:4–18.

Nei M, Kumar S. 2000. Molecular evolution and phylogenetics. Oxford, New York:

Oxford University Press; p. 352.

Neves MA, Halling RE. 2010. Study on species of *Phylloporus* I: Neotropics and North

America. Mycologia. 102:923–943.

Norton IO, Sclater, JG. 1979. A model for the evolution of the Indian Ocean and the

breakup of Gondwanaland. Journal of Geophysical Research. 84:6803–6830.

Nuhn ME, Binder M, Taylor AFS, Halling RE, Hibbett DS. 2013. Phylogenetic

overview of the Boletineae. Fungal Biology. 117:479–511.

Peay KG, Kennedy PG, Davies SJ, Tan S, Bruns TD. 2010. Potential link between plant

and fungal distributions in a dipterocarp rainforest: community and phylogenetic

structure of tropical ectomycorrhizal fungi across a plant and soil ecotone. New

Phytologist. 185:529—542

Peay KG, Schubert MG, Nguyen NH, Bruns TD. 2012. Measuring ectomycorrhizal

fungal dispersal: macroecological patterns driven by microscopic propagules. Molecular

Ecology. 21:4122—4136.

Pegler DN, Young TWK. 1981. A natural arrangement of the Boletales, with reference

to spore morphology. Transactions of the British Mycological Society. 76:103–146.

Peintner U, Bougher NL, Castellano MA, Moncalvo JM, Moser MM, Trappe JM,

Vigalys R. 2001. Multiple origins of sequestrate fungi related to *Cortinarius*

(Cortinariaceae). American Journal of Botany. 88:2168-2179.

Raidl S. 1997. Studien zur ontogenese rhizomorphen von ektomykorrhiza. Bibliotheca

Mycologica 169:1–184.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior

summarization in bayesian phylogenetics ssing Tracer 1.7. Systematic Biology. 67:901–

904.

Ree RH, Moore BR, Webb CO, Donoghue MJ. 2005. A likelihood framework for

inferring the evolution of geographic range on phylogenetic trees. Evolution. 59:2299–

2311.

Reguero MA, Goin FJ. 2021. Paleogeography and biogeography of the Gondwanan

final breakup and its terrestrial vertebrates: New insights from southern South America

and the "double Noah's Ark" Antarctic Peninsula. Journal of South American Earth

Sciences. 108:103358.

Rivas-Ferreiro M, Skarha SM, Rakotonasolo F, Suz LM, Dentinger BTM. 2023. DNA-

based fungal diversity in Madagascar and arrival of the ectomycorrhizal fungi to the

island. Biotropica. 55:954-968.

Ronquist F. 1994. Ancestral areas and parsimony. Systematic Biology. 43:0267– 274.

Ryberg M and Matheny PB. 2011. Dealing with incomplete taxon sampling and

diversification of a large clade of mushroom-forming Fungi. Evolution. 65:1862–1878.

Sánchez-García M, Ryberg M, Khan FK, Hibbett DS. 2020. Fruiting body form, not

nutritional mode, is the major driver of diversification in mushroom-forming fungi.

Proceedings of the National Academy of Sciences. 117:32528–32534.

Sánchez-Ramírez S, Ryberg M, Khan FK, Varga T, Nagy LG, Hibbett DS. 2015. High

speciation rate at temperate latitudes explains unusual diversity gradients in a clade of

ectomycorrhizal fungi. Evolution. 69:2196–2209.

Sato H, Tanabe AS, Toju H. 2017. Host shifts enhance diversification of

ectomycorrhizal fungi: diversification rate analysis of the ectomycorrhizal fungal genera

Strobilomyces and *Afroboletus* with an 80-gene phylogeny. New Phytologist. 214: 443–

454.

Sato H, Toju H. 2019. Timing of evolutionary innovation: scenarios of evolutionary

diversification in a species-rich fungal clade, Boletales. New Phytologist. 222:1924–

1935.

Schluter D. 2000. Ecological character displacement in adaptive radiation. The

American Naturalist. 156:4–16.

Simões M, Breitkreuz L, Alvarado M, Baca S, Cooper JC, Heins L, Herzog K,

Lieberman BS. 2016. The evolving theory of evolutionary radiations. Trends in Ecology

& Evolution, 31:27–34.

Singer R. 1945a. The Boletineae of Florida with notes on extralimital species I. The

Strobilomycetaceae. Farlowia. 2:97–141.

Singer R. 1945b. The Boletineae of Florida with notes on extralimital species. II. The

Boletaceae. Farlowia. 2:223–303

Singer R. 1947. The Boletoideae of Florida with notes on extralimital Species III. The

American Midland Naturalist. 37:1–135.

Singer R and Digilio AP. 1951. Pródromo de la flora agaricina argentina. Lilloa. 5–461.

Sitta N, Floriani M. 2008. Nationalization and globalization trends in the wild

mushroom commerce of Italy with emphasis on porcini (*Boletus edulis* and allied

species. Economic Botany. 62:307.

Smith MR. 2020. Information theoretic generalized Robinson–Foulds metrics for

comparing phylogenetic trees. Bioinformatics. 36:5007–5013.

Smith MR. 2022. Robust analysis of phylogenetic tree space. Systematic Biology.

71:1255–1270.

Smith SA, Brown JW, Walker JF. 2018. So many genes, so little time: A practical

approach to divergence-time estimation in the genomic era. PLOS ONE. 13:e0197433.

Soulebeau A, Aubriot X, Gaudeul M, Rougan G, Hennequin S, Haevermans T,

Dubuisson JY, Jabbour F. 2015. The hypothesis of adaptive radiation in evolutionary

biology: hard facts about a hazy concept. Organisms Diversity & Evolution. 15:747–

761.

Stecher G, Tamura K, Kumar S. 2020. Molecular evolutionary genetics analysis

(MEGA) for macOS. Molecular Biology and Evolution. 37:1237–1239.

Storey B. 1995. The role of mantle plumes in continental breakup: case histories from

Gondwanaland. Nature. 377:301–308.

Šutara J. 2005. Central European genera of the Boletaceae and Suillaceae, with notes on

their anatomical characters. Czech Mycology. 57:1–50.

Šutara J. 2014. Anatomical structure of pores in European species of genera *Boletus*

s.str. and *Butyriboletus* (Boletaceae). Czech Mycology. 66:157–170.

Tamura K, Battistuzzi FU, Billing Ross P, Murillo O, Filipski A, Kumar S. Estimating

divergence times in large molecular phylogenies. Proceedings of the National Academy

of Sciences. 109:19333–19338.

Tamura K, Stecher G, Kumar S. 2021. MEGA11: molecular evolutionary genetics

analysis Version 11. Molecular Biology and Evolution. 38:3022–3027.

Tamura K, Tao Q, Kumar S. 2018. Theoretical foundation of the RelTime method for

estimating divergence times from variable evolutionary rates. Molecular Biology and

Evolution. 35:1770–1782.

1002

1003   Tao Q, Tamura K, Kumar S. 2020. Efficient methods for dating evolutionary

1004   divergences. Pp. 197– 219 in Ho SYW (Ed.) The molecular evolutionary clock: theory

1005   and practice. Cham, Switzerland: Springer Nature Switzerland AG.

1006

1007   Thiel M and Gutow L. 2005. The ecology of rafting in the marine environment. II. The

1008   rafting organisms and community. Oceanography and Marine Biology. 43:279—418.

1009

1010   Thiers HD. 1984. The secotioid syndrome. Mycologia, 76:1–8.

1011

1012   Tremble K, Hoffman JI, Dentinger BTM. 2022. Contrasting continental patterns of

1013   adaptive population divergence in the holarctic ectomycorrhizal fungus *Boletus edulis*.

1014   New Phytologist. 237:295—309.

1015

1016   Tremble K, Suz LM, Dentinger BTM. 2020. Lost in translation: population genomics

1017   and long-read sequencing reveals relaxation of concerted evolution of the ribosomal

1018   DNA cistron. Molecular Phylogenetics and Evolution. 148:106804.

1019

1020   Tremble K, Halling RE, Henkel TW, Moncalvo JM, Dentinger BTM. 2023.

1021   Nomenclatural novelties. Index Fungorum no. 555.

1022   https://www.indexfungorum.org/Publications/Index%20Fungorum%20no.555.pdf

1023

Varga T, Krizsán K, Földi C, Dima B, Sánchez-García M, Sánchez-Ramírez S, Szöllősi

GJ, Szarkándi JG, Papp V, Albert L, Andreopoulos W, Angelini C, Antonín V, Barry

KW, Bougher NL, Buchanan P, Buyck B, Bense V, Catcheside P, Chovatia M, Cooper

J, Dämon W, Desjardin D, Finy P, Geml J, Haridas S, Hughes K, Justo A, Karasiński D,

Kautmanova I, Kiss B, Kocsubé S, Kotiranta H, LaButti KM, Lechner BE, Liimatainen

K, Lipzen A, Lukács Z, Mihaltcheva S, Morgado LN, Niskanen T, Noordeloos ME,

Ohm RA, Ortiz-Santana B, Ovrebo C, Rácz N, Riley R, Savchenko A, Shiryaev A,

Soop K, Spirin V, Szebenyi C, Tomšovský M, Tulloss RE, Uehling J, Grigoriev IV,

Vágvölgyi C, PappT, Martin FM, Miettinen O, Hibbett DS, Nagy LG. 2019.

Megaphylogeny resolves global patterns of mushroom evolution. Nature Ecology &

Evolution. 3:668–678.

1035

1036   Vellinga EC, Kuyper TW, Ammirati J, Desjardin DE, Halling RE, Justo A, Laessoe T,

1037   Lebel T, Lodge DJ, Matheny PB, Methven AS, Moreau PA, Mueller GM, Noordeloos

1038   ME, Nuytinck J, Ovrebo CL, Verbeken A. 2015. Six simple guidelines for introducing

1039   new genera of fungi. IMA Fungus. 6:A65–A68.

1040

1041   Wilf P, Nixon KC, Gandolfo MA, Cuneo NR. 2019. Eocene Fagaceae from Patagonia

1042   and Gondwanan legacy in Asian rainforests. Science. 364:5139.

1043

1044   Wilson AW, Binder M, Hibbett DS. 2011. Effects of gasteroid fruiting body

1045   morphology on diversification rates in three independent clades of fungi estimated using

1046   binary state speciation and extinction analysis. Evolution. 65:1305—1322.

1047

1048   Wu G, Feng B, Xu J, Zhu XT, Li YC, Zeng NK, Hosen MI, Yang ZL. 2014. Molecular

1049   phylogenetic analyses redefine seven major clades and reveal 22 new generic clades in

1050   the fungal family Boletaceae. Fungal Diversity. 69:93–115.

1051

Wu G, Miyauchi S, Morin E, Kuo A, Drula E, Varga T, Kohler A, Feng B, Cao Y,

Lipzen A, Daum C, Hundley H, Pangilinan J, Johnson J, Barry K, LaButti K, Ng V,

Ahrendt S, Min B, Choi IG, Park H, Plett JM, Magnuson J, Spatafora JW, Nagy LG,

Henrissat B, Grigoriev IV, Yang ZL, Xu J, Martin FM. 2022. Evolutionary innovations

through gain and loss of genes in the ectomycorrhizal Boletales. New Phytologist.

233:1383–1400.

LEGENDS

Table 1. Morphological transition rates estimated with Bayestraits.

Figure 1. Selected Boletaceae collections from two of the most species-rich regions that

were newly sequenced in this study. A) *Boletus cervinococcineus*, Singapore (BD616);

B) *Heimioporus punctisporus*, Sarawak (BAKO2); C) unidenti- fied Boletaceae,

Vietnam (CTN-08-0007); D) unidentified Boletaceae (CTN-08-0029); E) *Spongiforma*

sp., Sarawak (BTNG10); F) *Leccinum* sp., Sarawak (SWK246); G) *Crocinoboletus*

laetissimus, Sarawak (SWK335); H) unidentified Boletaceae sp., Viet- nam (DLT-08-

0127); I) *Boletellus* sp., Sarawak (SWK356); J) unidentified Boletaceae, Vietnam

(CTN-08-0051); K) *Tylopilus* sp., Cameroon (BD655); L) *Xerocomus* sp. 9. Cameroon

(BD773); M) *Fistulinella staudtii*, Cameroon (BD848); N) *Boletellus* sp., Cameroon

(BD714); O) *Phylloporus cf. tubipes*, Cameroon (BD719); P) *Tylopilus s*p. 8, Cameroon

(BD816); Q) *Xerocomus* sp. 8, Cameroon (BD695); R) *Boletus alliaceus*, Cameroon

(BD697); S) *Tubosaeta brunneosetosa*, Cameroon (BD686); T) *Tylopilus* sp.,

Cameroon (BD716). Not to scale.

Figure 2. Time-calibrated phylogeny of Boletaceae using 1764 BUSCO genes.

Topology is a summary coalescent of individual best ML gene trees using astral-hybrid.

Numbers on branches are quartet probabilities. Branch lengths were converted to time

using the top 100 best gene trees estimated using SortaDate in RelTime. Inset a) map of

specimen origins with numbers of specimens from each geographic area. Inset

b) lineages-through-time plot calculated with the "ltt.plot" function in the R pack- age

"ape." The dashed line represents a constant birth-death rate. The shaded box indicates a

period of significant divergent increase from a constant birth-death rate indicative of a

rapid radiation.

Figure 3. Biogeographic reconstruction using BioGeoBEARS. Left-hand tree depicts 4-

state coding scheme (light blue=Neotropical, blue=Paleotropical, yellow=North

Temperate, red=South Temperate) and right-hand tree depicts floristic region coding

scheme (pink=Chilean-Patagonian, green=Indo-Malesian, blue=African,

yellow=Holarctic, red=Novo-Zealandic, light blue=Neotropical, orange=Australian).

Pie charts indicate the proportional likelihood of each state at a node. Red and black

boxes indicate Phase I and Phase II Gondwanan diversification events, respectively.

Figure 4. Dispersal events and rates inferred from BioGeoBEARS. Top: Map depicting

phytogeographic zones that are colored and labeled on the map. Map was ren- dered

using the 'imago' R code (https://github.com/hrbrmstr/imago) to reproduce the

AuthaGraph world map projection (http://www.authagraph.com/top/?lang=en). Curved

arrows indicate inferred directional dispersal events and are colored by rate values

following the table (Bottom). The stroke weight of the arrows has been scaled to percent

of the maximum rate value following the values in the table. Bottom: Table of dispersal

rates inferred under a DIVAlike+j model in BioGeoBEARS. Source regions are at left

and destination regions are along the top. Values are colored along a scale from cool to

warm (red being maximum).

Figure 5. Distribution of likelihood frequencies of morphological traits. A-C)

Illustrations of coded traits and inferred transition rates. Arrows indicate direction of

transition with stroke weights scaled to maximum rate value. D-I) Frequency

distributions of state probabilities for six trait codings. Traits and codes are labeled on

the x-axis.

1104

1105   *Corresponding author: Email: keaton.tremble@gmail.com. Current address:

1106   Department of Biology, Duke University, Durham, North Carolina 27708.

Trait Transition	Mean rate	Median rate
Ballistospory		
Yes to No	1.2303	1.0565
No to Yes	39.8088	35.7478
Spore Ornamentation		
Smooth to Not Smooth	0.7747	0.7498
Not Smooth to Smooth	1.741	1.464
Stipe		
Absent to Present	33.685	27.67
Present to Absent	0.8415	0.6788
Sporocarp Habit		
Secoitiod to Pileate-stipitate	51.64	51.47
Pileate-stipitate to Secoitiod	0.8178	0.7172
Secoitiod to Gasteroid	53.44	53.90
Gasteroid to Secoitiod	34.63	31.05
Pileate-stipitate to Gasteroid	0.8393	0.6749
Gasteroid to Pileate-stipitate	40.78	36.85
Hymenophore Bruising		
None to Blue	1.3515	1.2632
None to Black	0.6921	0.6921
Blue to None	5.3730	5.3540
Blue to Black	0.1732	0.1205
Black to None	3.2329	2.7385
Black to Blue	1.0294	0.7281
Hymenophore Anatomy		
Straight to Cross	0.6398	0.6092
Straight to Lamellete	0.2054	0.1715
Cross to Straight	3.694	2.960
Cross to Lamellate	2.4131	1.6755
Lamellate to Straight	5.483	5.004
Lamellate to Cross	1.858	1.275

Southeast Asia

Africa

	DESTINATION						
	Holarctic	Neotropical	Indo-Malesian	African	Australian	Novo-zealandic	Chilean-Patagonian
Holarctic	0	4.28	17.42	4.66	2.72	2.22	1.76
Neotropical	1.26	0	1.16	2.34	0.22	0.6	0.34
Indo-Malesian	20.52	3.12	0	8	4.32	2.54	0.64
African	6.46	6.1	5.56	0	1.24	0	0.2
Australian	2.38	1.34	2.16	0.5	0	0.66	1.06
Novozealandic	0.28	0.48	0.12	0	1.4	0	0
Chile-Patagonian	0.74	0.32	0.04	0.04	0.84	0	0