Phylogenomics, biogeography, and trait evolution of the Boletaceae (Boletales,

Agaricomycetes, Basidiomycota)

Keaton Tremble[a]*,Terry Henkel[b], Alexander Bradshaw[a], Colin Domnauer[a], Lê Xuân

Thám[c], Guliana Furci[d], Cathie Aime[e], Jean-Marc Moncalvo[f], Bryn Dentinger[a]*

[a]Natural History Museum of Utah and School of Biological Sciences, University of

Utah, Salt Lake City, Utah 84108

[b]Department of Biological Sciences, California State Polytechnic University, Humboldt,

Arcata, California 95521

[c]Van Lang University, Ho Chi Minh City, Viet Nam

[d]Fungi Foundation, Brooklyn, New York 11222, United States

[e]Department of Botany and Plant Pathology, Purdue University, West Lafayette,

Indiana 47907

[f]Department of Natural History, Royal Ontario Museum & Department of Ecology and

Evolutionary Biology, University of Toronto, Toronto M5S 2C6, Canada

ABSTRACT

The species-rich porcini mushroom family Boletaceae is a widespread and well-known

group of ectomycorrhizal (ECM) mushroom-forming fungi that has eluded intrafamilial

phylogenetic resolution despite many attempts using morphological traits and multi-

locus molecular datasets. In this study, we present a genome-wide molecular dataset of

1764 single-copy gene families from a global sampling of 418 Boletaceae specimens.

The resulting phylogenetic analysis has strong statistical support for most branches of

the tree, including the first statistically robust backbone. The enigmatic Phylloboletellus

chloephorus from non-ECM subtropical forests in northwestern Argentina is recovered

as an early diverging lineage within the Boletaceae. Time-calibrated branch lengths

estimate that the family first arose in the early-to-mid Cretaceous and underwent a rapid

radiation in the Eocene, possibly when the ECM nutritional mode arose with the

emergence and diversification of ECM angiosperms. Biogeographic reconstructions

reveal a complex history of vicariance and episodic long-distance dispersal correlated

with historical geologic events, including Gondwanan origins and cladogenesis patterns

that parallel its fragmentation. Ancestral state reconstruction of morphological traits

predicts that the ancestor of the Boletaceae was lamellate with ornamented

basidiospores, contrary to the most common "bolete" sporocarp morphology. Transition

rates indicate that the lamellate hymenophore and sequestrate sporocarp are reversible

traits. Together, this study represents the most comprehensively sampled, data-rich

molecular phylogeny of the Boletaceae to date, enabling robust inferences of trait

evolution and biogeography in the group.

KEYWORDS:

phylogenomics; Boletaceae; porcini; evolutionary radiation; Gondwana

INTRODUCTION

Evolutionary radiations result from short bursts of relatively rapid diversification and

demonstrate the creative power of evolution. Adaptive and non-adaptive radiations may

be the most common macroevolutionary patterns, and are fundamental to the origins of

biodiversity (Simoes et al. 2016; Futuyma 1998; Schluter and McPeek 2000). Yet, the

underlying causes of evolutionary radiations are poorly known for most groups, as most

studies have focused on animals and plants (Soulebeau et al. 2015). The other major

multicellular eukaryotic group, the Fungi, have largely neglected (Varga et al. 2019).

The porcini mushroom family Boletaceae (Fig. 1) is an example of an

evolutionary radiation in Fungi (Bruns et al. 1992). The family is exceptionally diverse

(>2000 currently accepted species) and globally distributed, but poorly documented for

many regions. Yet, they are prevalent ectomycorrhizal (ECM) mutualists in ecosystems

dominated by ECM plants (Peay et al. 2010) and at least eight species are traded

globally as wild-collected edible mushrooms (Arora 2008; Sitta and Floriani 2008;

Dentinger et al. 2010; Dentinger and Suz 2014). Despite their conspicuous sporocarps

(=mushrooms), ecological dominance, and cultural importance, new species of

Boletaceae are regularly described from poorly explored habitats around the globe(e.g.

Halling et al. 2006, 2023; Fulgenzi et al. 2007, 2008, 2010; Neves et al. 2010; Husbands

et al. 2013; Castellano et al. 2016; Chakraborty et al. 2015; Das et al. 2015, 2016;

Henkel et al. 2016; Magnago et al. 2017), as well as from wild-collected foods in local

markets and even from supermarkets, e.g., (e.g. Das et al. 2015; Dentinger and Suz

2014; Halling et al. 2014). While these new species are increasingly understood in a

global phylogenetic context, shedding light on their origin, diversification, and

migration in time and space, there has been little progress made towards resolving the

backbones in the phylogenies for Boletaceae using legacy loci (Grubisha et al. 2001;

Binder & Hibbett 2006; Drehmel et al. 2008; Dentinger et al. 2010; Nuhn et al. 2016;

Wu et al. 2014).

One of the most prominent consequences of this lack of phylogenetic resolution

is the recent explosion of new generic names to accommodate newly discovered

species, or species that are included in molecular phylogenetic analyses for the first time

and that are recovered on long branches with no supported affinity to existing generic

names (e.g., Castellano et al. 2016; Henkel et al. 2016; Badou et al. 2022; Halling et al.

2023). Although this is a predictable pattern for evolutionary radiations, it has resulted

in a recent proliferation of new genera, many of which are monotypic or require

identification of species to recognize, an impractical solution to the problem. Taken

together, the current situation is perhaps best described as a quagmire of nomenclatural,

taxonomic, ecological, and evolutionary speculation.

Beyond taxonomic concerns, the Boletacaeae presents a unique system to

identify the genetic mechanisms that contribute to diversification. Abundant evidence

indicates that the Boletaceae underwent an evolutionary radiation 60-100 million years

ago (Bruns and Palmer 1989; Binder et al. 2006; Dentinger et al. 2010; Wu et al. 2014,

2016; Sato et al. 2019). This early radiation has been correlated with the convergence of

morphological traits, such as lamellate ("gilled") mushrooms and gasteromycetization

(Castellano et al. 2016), which suggests that the development of novel morphological

diversity is constrained by relatively few changes in developmental pathways.

Sampling fungi for comparative studies is often a difficult and time-consuming

process due to their cryptic somatic lifestyles and the ephemeral, unpredictable

sporocarps. This can result in ad hoc taxon sampling that suffers from heavy sampling

bias.

Although progress has been made in describing new diversity and construction

of molecular phylogenetic datasets, the phylogeny for Boletaceae has been frustratingly

difficult to resolve. This is the consequence of molecular phylogenetic datasets that

included only one to four loci (Grubisha et al. 2001; Binder & Hibbett 2006; Drehmel et

al. 2008; Dentinger et al. 2010; Nuhn et al. 2013; Wu et al. 2014, 2016). While

phenomena such as incomplete lineage sorting and hybridization may obscure historical

phylogenetic signal, previous datasets for the Boletaceae have consisted of patchy

taxonomic and geographic sampling. All these factors' impacts on phylogenetic

reconstruction were intensified by the aforementioned rapid radiation event (Bruns et al.

1992; Sato et al. 2017). Without a robust phylogeny, it is impossible to name, classify,

reproduce ecological studies, and understand evolutionary history in the Boletaceae. For

example, only a few recent studies have included representatives of the exceptionally

rich Australian macrofunga (Halling, Fechner, et al. 2015, 2023; Halling, Nuhn, et al.

2012), and the Neotropics and Afrotropics have been conspicuously underrepresented in

family-level analyses despite their exceptional species richness (Heinemann 1951;

Henkel et al. 2012).

Recent fieldwork has resulted in many new collections from under-sampled

regions (B. Dentinger and T. Henkel, unpublished data), and these are now available to

include in phylogenetic datasets aiming to achieve the first globally representative

sampling of the Boletaceae. Collections-based phylogenomics is also effective for

resolving ancient relationships among mushroom-forming fungi (Dentinger et al. 2015,

Liimatainen et al. 2021, Tremble et al 2020). No one, however, has yet applied these

methods to the Boletaceae. Moreover, whole genome sequencing of mushroom forming

fungi provides opportunities to go beyond phylogeny reconstruction. For example,

whole genome sequencing can facilitate analyses of the population processes that

generate biodiversity (e.g. Tremble et al. 2022), resulting in a much richer insight into

biology than only a few molecular loci can provide on their own.

For this study, we generated the first robust phylogeny of the Boletaceae that

utilized a large molecular dataset comprised of 1764 genome-wide loci from 418 taxa

across the family collected in many tropical and temperate geographic regions. To

obtain sufficiently representative sampling, we targeted fungarium collections including

type specimens whenever possible, augmented by recent new collections to fill in

geographic and taxonomic gaps. We included samples from regions that have been

neglected in the past (e.g., Africa, southern South America, lowland tropical South

America, Australia), and wherever possible type species to provide an anchor for a

future taxonomic revisions of genera. We also performed the first inclusive

biogeographic reconstruction of the Boletaceae and assessment of morphological trait

evolution using our highly resolved genomic phylogeny. Overall, we provide

unprecedented insight into broad patterns of evolution in this enigmatic fungal group.

MATERIALS AND METHODS

Sampling

Taxon selection focused on obtaining representatives of all currently accepted genera,

selecting type species whenever possible. Because the current understanding of genera

is incomplete and rapidly changing, we could not include representatives of all currently

accepted genera that were published during the course of this study. Specimens from

geographic regions unrepresented in prior studies were also included. A total of 418

Boletaceae specimens were gathered from a global distribution using collections made

by the authors, those borrowed from four institutions, and donations from citizen

scientists (Supplementary Table 1). In addition, we utilized genome data publicly

available from the JGI Mycocosm Portal (Grigoriev et al. 2014) for *Boletus coccyginus*,

B. reticuloceps, *Butyriboletus roseoflavus*, *Chiua virens*, *Lanmaoa asiatica*, and *Imleria*

badia (Miyauchi et al. 2020, Wu et al. 2022, Kohler et al. 2015). Genomes of *Paxillus*

involutus, *Paxillus adelphus*, and *Hydnomerulius pinastri* (Kohler et al. 2015) from JGI

were used for outgroups.

DNA extraction and sequencing

Genomic DNA was extracted from 10mg of hymenophore tissue from each specimen.

Tissues was homogenized in 2.0 mL screw-cap tubes containing a single 3.0 mm and 8

x 1.5 mm stainless steel beads by bashing them in a BeadBug microtube homogenizer

(Sigma-Aldrich, #Z763713) for 120 seconds at a speed setting of 3500 rpm. After

physical disruption, DNA was extracted using the Monarch® Genomic DNA

Purification Kit (New England Biolabs, Massachusetts; #T3010) with the Monarch®

gDNA Tissue Lysis Buffer (#T3011) using double the volume of lysis buffer, 1 hour

lysis incubation at 56 C , and 550 µl of wash buffer during each of the wash steps. DNA

extract quality was assessed for quality using a NanoDrop 1000 (Thermo Scientific) and

fragment integrity using agarose gel electrophoresis. Genomic DNAs were sequenced

using a combination of paired-end sequencing on the Illumina MiSeq, HiSeq, and

Novaseq sequencing platforms (Supplementary Table 2). All raw reads and whole

genome assemblies are deposited in the Short Read Archive

(Bioproject#PRJNA1022813).

Genome assembly and ortholog extraction

Raw sequencing reads were quality-filtered and adapter-trimmed using fastP v0.20.1

(Chen et al. 2018) with default settings. Genome assemblies were produced from

quality-filtered reads using SPAdes v3.15.0 (Bankevich et al. 2012) with five k-mer

values (k=77,85,99,111,127). From each genome, we identified 1764 highly-conserved

single copy orthologs using BUSCO with the "basidiomycota odb 10" dataset.

Orthogroups that were present in less than 75% of taxa and taxa with less than 20%

ortholog recovery were removed. Retained orthologs were aligned using MAFFT

v7.397 (Katoh, Rozewicki, and Yamada 2017) with the "L-INS-I" algorithm, and

maximum- likelihood gene-trees were inferred using IQ-TREE v2.0.3 (Minh et al.

2020) with automatic model selection in ModelFinder (Kalyaanamoorthy et al. 2017)

and ultra- fast bootstrapping (BS, (Hoang et al. 2018)) with 1000 replicates. A summary

coalescent species tree was constructed from the resulting gene trees using hybrid-

ASTRAL implemented in ASTER (v1.15) (Zhang & Mirarab 2022). Branch lengths in

substitutions/site were estimated under maximum likelihood on the species tree using

the "-te" option in IQ-TREE, with a partitioned concatenated alignment of all BUSCO

genes used in species tree construction.

Gene tree comparison

To evaluate gene tree discordance, individual trees were compared using six metrics

calculated in SortaDate (average bootstrap support, clocklike branch lengths, tree

length; Smith et al. 2018) and the R package 'TreeDist' (generalized Robinson-Foulds

metrics; Smith 2020, 2022). In addition to data matrix summaries (number of taxa,

alignment length), Pearson's correlations were calculated to determine correlations

between metrics.

Divergence dating

A timetree was inferred by applying the RelTime method (Tamura et al. 2012, 2018)

conducted in MEGA11 (Stecher et al. 2020, Tamura et al. 2021) to the species tree with

ML-estimated branch lengths. To reduce computational burden, time-calibrated branch

lengths were calculated using the Maximum Likelihood (ML) method and the General

Time Reversible substitution model (Nei & Kumar 2000) from two sets of 100 genes: 1)

the top 100 genes with well-supported clock-like trees determined using SortaDate

(Smith et al. 2018) and 2) the top 100 genes with the smallest generalized Robinson-

Foulds ('gRF') distances to the species tree calculated using the R package 'TreeDist.'

The timetree was computed using two sets of calibration constraints. The first included

two calibrations: 1) a secondary calibration for the stem age of the Boletaceae from 50-

150 mya (Varga et al. 2019, Wu et al. 2022) and 2) a secondary calibration for the stem

age of Boletus edulis from 5-13 mya (Tremble et al. 2022). The second set included the

former two calibrations plus four of the five internal calibrations using the highly

supported core shifts from Varga et al. (2019). Because many of the clades in Varga et

al. were incongruent with our topology, calibrations were selected using the most

inclusive node, except for *Aureoboletus* which could not be reconciled with our results.

The Tao et al. (2020) method was used to set minimum and maximum time boundaries

on nodes for which calibration densities were provided. Confidence intervals were

computed using the Tao et al. (2020) method. Times were not estimated for outgroup

nodes because the RelTime method uses evolutionary rates from the ingroup to

calculate divergence times and does not assume that evolutionary rates in the ingroup

clade apply to the outgroup.

Ancestral Range Reconstruction

Numerous analytical methods for reconstructing historical biogeography exist,

accounting for processes such as vicariance, dispersal, and cladogenesis (Ronquist

1994; Ree et al. 2005; Landis et al. 2013). To account for these macroevolutionary

processes in our ancestral state reconstruction in the Boletaceae, we utilized

BioGeography with Bayesian (and likelihood) Evolutionary Analysis with R Scripts

("BioGeoBEARS"; Matzke 2018), which has unified the most prevalent biogeographic

models. Samples were coded in two ways: 1) Paleotropical (consisting of Africa and

Tropical Asia), Neotropical (South and Central America), South Temperate (temperate

Australia and New Zealand), or North Temperate (North America, Europe, Northern

tem- perate Asia), and 2) by floristic regions of the world, Holarctic including Central

America, Neotropical, Chilean-Patagonian, African, Indo-Malesian, Australian, and

Novo-Zealandic (Liu et al 2023) (Supplementary Table 3). Central America was

combined with the Holarctic region because prior studies have shown that Central

American ECM fungi are mostly derived from North American ancestors (Halling

1996). The most likely model was chosen according to AIC and weighted AIC score

calculated in BioGeoBEARS.

RESULTS

DNA sequencing, genome assembly, and ortholog extraction

Whole genome sequencing of 418 specimens resulted in 13,794,532 paired-end reads

per specimen on average (Supplementary Table 2). On average, genome assemblies

possessed an assembly N50 of 12.9 Kbp (thousand base-pairs), total assembly length of

61.6 Mbp (million base-pairs), 53,972 scaffolds, and a BUSCO score of 74.7%. 34 of

418 specimens possessed BUSCO scores less than 20%, and 175 specimens possessed

BUSCO scores greater than 90% (Supplementary Table 2). After removing specimens

with poor BUSCO recovery, our final dataset included 383 Boletaceae specimens, three

outgroup taxa, and 1461 single-copy orthologs.

Gene tree comparison

Average bootstrap support had the highest positive correlation with the number of taxa

present (Pearson's coefficient = 0.76), and weak to moderate negative correlations with

alignment length (Pearson's coefficient = -0.17), clocklike branch lengths (Pearson's

coefficient = -0.13), and total tree length (Pearson's coefficient = -0.20). Generalized

Robinson-Foulds distances were weakly to moderately negatively correlated with

number of taxa (Pearson's coefficient = -0.33), total length (Pearson's coefficient = -

0.30), and clocklike branch lengths (Pearson's coefficient = -0.13), and weakly to

moderately positively correlated with alignment length (Pearson's coefficient= 0.29)

and average bootstrap support (Pearson's coefficient = 0.17). Clocklike branch lengths

and total tree length were weakly positively correlated (Pearson's coefficient= 0.15).

Divergence Dating

Stem ages for the Boletaceae were estimated at 138-139 mya and 77 mya using the

two- and six-calibration sets, respectively. The crown age of the Boletaceae was

estimated at 103-105 mya and 63 mya (49-77 mya) using the two- and six-calibration

sets, respectively. The diversification of the Boletineae was estimated at 83-87 mya and

58 mya (49-77 mya) using the two- and six-calibration sets, respectively. The origin of

Boletus sensu stricto (i.e. "porcini) was estimated at 38 mya and 35 mya (29-42 mya),

and its diversification was estimated at 29-30 mya and 26 mya (20-34 mya), using the

two- and six-calibration sets, respectively.

Ancestral state reconstruction

Ancestral state reconstruction recovered the most likely ancestor of the Boletaceae to be

a pileate-stipitate, lamellate mushroom with ornamented spores. The highest transition

rates were observed in phenotypic changes from secotioid to gasteroid and from

secotioid to pileate-stipitate morphologies, recovering secotioid morphology as being a

relatively transient evolutionary state (Table 1). High transition rates were also found to

occur from gasteroid to pileate-stipitate morphology, supporting evidence for the

reversibility of gasteromycetation.

Ancestral Range Reconstruction

Ancestral distribution reconstruction recovered a likely Paleotropical origin of the

Boletaceae (DEC+J model chosen with lowest AIC and AICc for both coding sets),

with two major descendant radiations originating in the Paleotropics (Africa and Asia)

and Neotropics (Fig. 3). In addition, we found evidence for multiple diversification

events spurred by the separation of Gondawana (Fig. 3). Gondwanan separation

occurred in two predominant phases: Phase 1, involved the separation of Southern

South America, Southern Africa, Australia-Antarctica, and Madagascar-India, which

began approximately 180 MYA and was largely completed by 120 MYA (Jokat et al.

2003) and Phase 2, involving the separation of South America and Africa, which was

completed 80 MYA (Reguero and Goin 2021). At the split between the

Austroboletoidae and Suilelloidae (Fig. 3), we found a putative Phase 1 Gondwanan

separation event estimated to have occurred 150-120 MYA, which led to the rapid

formation of South Temperate, Neotropical, and Paleotropical lineages. Later, around

90-70 MYA, we found at least five putative Phase 2 separation events, leading to the

separation of Paleotropical and Neotropical lineages.

286   In our four-category paleo-region coding set, the Boletaceae ancestor was

equally likely to be Neotropical or Paleotropical. However, the subsequent node that

leads to the rest of the Boletaceae (excluding Phylloboletoidae and Chalciporoideae)

was recovered as Paleotropical with high likelihood, as were all subsequent daughter

nodes. The coding of the Chalciporoideae and the single *Phylloboletellus* specimen in

our dataset likely had a strong influence on ancestral range reconstructions at the

deepest nodes. The backbone nodes of the Boletaceae excluding Phylloboletoidae and

Chalciporoideae were estimated as Asian in origin, corroborating the four-category

analysis, though with less confidence. Migrations between phytogeographic regions

were dominated by dispersals between the Indo-Malesian and Holarctic regions (Fig.

3,4).

DISCUSSION

Phylogeny

The fully resolved backbone supports the recognition of eight subfamilies, including

two newly defined as Phylloboletelloideae and Suillelloideae(Tremble et al. 2023) with

their interrelationships resolved for the first time. The Chalciporoideae was recovered as

the earliest diverging group in the Boletaceae, a relationship previously noted by Wu et

al. (2014). The enigmatic *P. chloephorus* (Singer & Diglio 1952) is the next lineage to

branch off before the radiation that gave rise to the six additional subfamilies. Previous

studies have placed *Pseudoboletus parasiticus* in a position similar to where we

recovered *P. chloephorus* (e.g. Wu et al. 2014), but we were not able to include a

representative of *P. parasiticus* in this study and cannot corroborate this. Nonetheless,

the tree topology has intriguing implications for the role of ecological transitions in

Boletaceae diversification. Members of the earliest-diverging Chalciporoideae are

ecologically labile can be facultatively ECM or mycoparasitic (Caiafa and Smith 2022)

and *P. chloephorus,* the next lineage to branch off, is primarily known from non-

ectotrophic forests which suggests a non-ECM ecology (Singer & Diglio 1952).

Moreover, *P. parasiticus* is a parasite of *Scleroderma* spp. pseudotruffles (Raidl 1997;

Binder and Hibbet, 2006; Nuhn et al. 2013). Altogether, the basal position of these early

diverging groups suggests that the ancestor of the Boletaceae was non-ECM, and is in

line with a previous study which indicated that the ECM habit emerged with the origin

of the six derived subfamilies that subsequently radiated (Sato & Toju 2019). A number

of genomic changes coinciding with the emergence of obligate ectomycorrhizality (Wu

et al. 2022) further support the view that this shift in nutritional mode has profoundly

impacted evolution of the Boletaceae. Our phylogeny also recovered previously

undocumented sister relationships between the Zangioideae and Xerocomoideae, and

between Austroboletoideae and Suillelloideae. The phylogeny also recovered a new

clade composed of the Austroboletoideae, Suillelloideae and Boletoideae.

Taxonomic implications

Many taxonomic changes in the Boletaceae have been proposed in recent years.

Primarily these have involved the erection of new genera for phylogenetically

unresolved lineages. Many of these new genera are monotypic or composed of only a

few species (Castellano et al. 2016; Chakraborty et al. 2015; Henkel et al. 2016; Halling

et al. 2023). This burdensome proliferation of new genera has made it difficult or

impossible to recognize inclusive groupings without requiring knowledge of the species.

On the bright side, the strong nodal support throughout our phylogeny sets the stage for

a new, comprehensive and stable generic-level taxonomy which will be addressed in

subsequent work when all recently described genera are sampled. Nonetheless, we have

delimited higher-level groups into subfamilies following previous authors (Wu et al.

2014), which precipitated the need to formally recognize two new subfamilies (Tremble

et al. 2023). The subfamily Boletoideae is also expanded with taxa not previously

included.

Biogeography and chronology

Our biogeographic and divergence dating analyses support a Gondwanan origin of the

Boletaceae. Subsequent diversification was likely facilitated by vicariance events

correlated with continental drift and possible long-distance dispersal. These results

support the emerging consensus that numerous lineages of ECM fungi originated in

Gondwana or more recently in paleotropical regions (Hosaka et al. 2008, Matheny et al.

2009, Dentinger et al. 2010, Ryberg & Matheny 2011, Kennedy et al. 2012, Sanchez-

Ramirez et al. 2015, Han et al. 2018; Hackel et al. 2022; Codjia et al. 2023).

Phytogeographic endemism was implied with our biogeographic reconstruction. The

strongest migrations occurred recently over the past 50 mya between the Indo-Malesian

and Holarctic regions (Figs. 3, 4). We acknowledge the difficulty to determining origins

and dispersal events with certainty in the absence of a fossil record or other

corroborating evidence. Nonetheless it is clear from our results and those of other

studies that vicariance may have played a strong role in the distribution of ECM fungal

taxa, despite the long-distance dispersal capacity of airborne spores (Matheny et al.

2009, Peay et al. 2010, Peay & Matheny 2016). Conversely, pure vicariance cannot

explain the close phylogenetic relationships seen between distant disjunct taxa. It

follows that long-distance dispersals may have occurred, albeit rarely. While long-

distance dispersal is demonstrably possible in the Boletaceae (Tremble et al. 2022), and

has been documented in diverse ECM lineages (Gelm et al. 2012; Hackel et al. 2022), it

is obvious that successful long-distance dispersal is difficult. Most basidiospores do not

travel far from the sporocarp (Galante et al. 2011), and the probability of two

basidiospores landing in proximity to mate becomes highly unlikely with increasing

distance from sporocarps (Peay et al. 2012; Golan and Pringle 2017). Such

improbabilities notwithstanding, our biogeographic patterns are consistent with episodic

long-distance dispersal of basidiospores or even vegetative mycelium vectored by wind,

migrating birds (Elliot et al. 2019), or even rafting vegetation (Thiel et al. 2005).

Our biogeographic reconstructions are consistent with the "Southern Route to

Asia" hypothesis (Wilf et al. 2019). This idea proposes that ECM Fagaceae, and their

symbiont fungi, originated in Gondwana and drifted on Australia north to Asia. In this

scenario the Gondwana ECM habitat tracked climatic niches northward as the

Australian plate collided with the Pacific plate. A resulting relictual ECM community

remained in an emergent New Guinea and subsequently spread northwest along the

montane Australasian archipelago and dispersed onto continental Asia. Many of the

inferred dispersal events between Indo-Malesia and other regions, especially the

Holarctic, are inferred within the last 20 my (Figs. 3,4), coincident with the collision of

Australia with the Pacific plate in the later Oligocene (Hall 2011). More recent

migrations across the Australasian archipelago are likely (Halling et al. 2008),

consistent with the many recent dispersal events (Fig. 3).

Biogeographic reconstructions are highly sensitive to taxon sampling and

despite our best efforts this dataset is not immune to misleading reconstructions. For

example, in both coding schemes the ancestral node of the Chalciporoideae had the

highest probability of a North Temperate and North American origin. However, we did

not have access to samples of this clade from Asia, Africa or Australia/New Zealand,

and this could have had impacts on the reconstruction. Nonetheless we have the most

geographically comprehensive sampling for Boletaceae ever compiled, and the first

opportunity to examine global-scale biogeographic patterns. Thus, insights into the

evolution of the Boletaceae are revealed for the first time, despite mild uncertainty at a

minority of nodes.

The origin and evolution of distinctive Boletaceae funga in specific regions have

long been a mystery (Horak 1977). For example, the endemic Boletaceae of Chile and

Argentina have not been included in previous phylogenetic studies, and their

morphology based affinities have been inconclusive (Horak 1977). Four sampled

Chilean species were recovered as ancient lineages of similar age within four of the

subfamilies (*Gastroboletus valdivianus* in Xerocomoideae, *Boletus loyita* in

Austroboletoidae, *Boletus loyo* in Suillelloidae, *Boletus putidus* in Boletoidae).

Moreover, the closest relatives of the Chilean boletes occurred in geographic regions as

disjunct as North America, lowland tropical northern South America, and Australia.

Furthermore, *B. loyita* and *G. valdivianus* are most closely related to extant Australian

taxa and potentially originated prior to final Gondwanan disarticulation (Reguero and

Goin 2021). Close relationships between Australian and South American taxa is a

common pattern produced by Gondwanan separation (Feng et al. 2017). Altogether,

Chilean boletes likely arose in Gondwana, separated from their Australian sister

lineages during Gondwanan disarticulation, and experienced no subsequent speciation

for tens of millions of years.

Ancestral range reconstruction recovered an Asian origin of the core genus

Boletus s. str. (a.k.a "true porcini") as previously suggested (Feng et al. 2012).

However, we cannot entirely rule out an African origin. The Central African endemic

Boletus alliaceus was recovered here as a sister taxon to *Boletus* s. str., and another

study found a similar relationship for the recently described African *Paxilloboletus*

africanus (Badou et al. 2022). Furthermore, we estimated the origin of *Boletus* s. str to

be 40 mya, which may indicate why the sister lineages to *Boletus* s. str. are endemic to

Africa. At 40 mya the Indian subcontinent, which had separated from Africa and

Madagascar ~120 mya (Reguero and Goin 2021), was already colliding with Asia

(Aitchison et al. 2007; Hu et al. 2016). If *B. alliaceus* and *P. africanus* are indeed sister

lineages of *Boletus* s. str., then it is possible that the ancestor of true porcini is also

African, and arrived in Asia via the Indian subcontinent with subsequent diversification.

However, even more diversity of *Boletus* s. str. is still to be uncovered, including the

currently undescribed species from Taiwan, Malaysian Borneo, and the Gulf Coast of

the US reported here. Clearly more mycological exploration and specimen collecting

are needed to fully document global diversity of *Boletus* s. str. In particular, searches in

India and Africa for potential relictual lineages could shed light on the origin of this

important charismatic group.

Divergence dating estimated the origin of the Boletaceae to be 138-139 mya,

and ancestral range reconstructions suggest it may be even older. Ranking genes with

different metrics had little impact on divergence date estimates. However, the two sets

of calibrations gave very different estimates for most nodes. The calibration set that

included estimates from Varga et al. (2019) gave younger divergence ages than the

calibration set with only two calibrations. In fact, the estimated origin dates of the

Boletaceae using the Varga et al. (2019) calibrations were almost half those of the two-

calibration set. It is difficult to interpret these divergence dates with any certainty due to

a lack of fossil evidence. However, the dates estimated using the Varga et al. (2019)

internal calibrations may be viewed with extra skepticism due to extensive topological

incongruence of the Varga et al. trees with our topology. The older divergence estimates

are corroborated by other studies (e.g., He et al. 2020), and internal dates of this tree

largely correspond with other work, such as the ~48 MYA origin of the Stobilomyces

group reported by Han et al. (2018). These older divergence estimates also correspond

with the origin of ECM Pinaceae in the early Cretaceous (Brundrett & Tedersoo 2018).

Therefore, we consider these older estimates be more accurate.

In our ancestral range reconstruction analysis, we found evidence of multiple

diversification events that may have been initiated by the breakup of the Gondwanan

supercontinent. The putative phase 1 Gondwanan separation event identified within the

Boletoidae suggests that the Boletaceae was diverse and widely distributed by at least

120 mya (Jokat et al. 2003), substantially older than the estimated age from our

divergence dating analysis but within the 95% confidence interval. Our dates are at best

coarse estimates based on secondary calibrations and lack fossil-based calibrations.

However, the phylogenetic pattern of vicariance that parallels the breakup of Gondwana

is compelling and offers corroborating evidence that our estimated ages are in fact too

young. In any case, our divergence estimates suggest that the Boletaceae originated and

diversified within the early to late Cretaceous period, when the climate was warmer and

wetter than it is now (Hay and Floegal, 2012), when gymnosperms and subsequently

angiosperms diversified (Crisp and Cook, 2011), and when the large supercontinental

land masses broke apart (Jokat et al. 2003).

Trait evolution

The longest standing debates of Boletaceae systematics have centred on the evolution of

morphological traits and their utility for defining natural generic-level groups.

Basidiospore color, ornamentation, and hymenophore arrangement were long

emphasized in this debate, on the one hand used as evidence for dividing the Boletaceae

into multiple genera (Singer, Pegler & Young, etc.) or treating nearly all Boletaceae as a

single genus (Corner 1972). More recently, anatomy of the pileal surface and pigment

chemistry have been emphasized (e.g. Binder et al. 2002, Sutara 2014). Yet, despite the

tremendous morphological and chemical variability in the Boletaceae, the family is

typified by the 'bolete' macro-morphology (i.e. pileate-stipitate mushrooms with tubular

hymenophores). In addition, spore ornamentation has been considered to be a derived

trait, unifying species into readily recognizable genera such as *Strobilomyces*,

Boletellus, and *Austroboletus* (Berkeley 1851, Murrill 1909, Corner 1972, Pegler &

Young 1981, Wolfe 1980). Contrary to these iconic morphotypes, our ancestral state

reconstructions suggest that the ancestor of the Boletaceae was probably a pileate-

stipitate mushroom with lamellate hymenophore and ornamented spores. This is almost

certainly due in large part to the phylogenetic position of the enigmatic *Phylloboletellus*

chloephorus (Singer & Diglio 1952). It is also possible that the obligately ECM habit,

characterising most extant Boletaceae, was gained after its origin as evidenced by the

habitat of *Phylloboletellus* and the saprotrophic and mycoparasitic abilities of

Chalciporoideae (Caiafa et al. 2022). The mycoparasitic *Pseudoboletus parasiticus* and

P. silvaticus have also been recovered as early-diverging lineages of Boletaceae (Binder

& Hibbett 2006, Cortes- Perez et al. 2023) and may have a close affinity with

Phylloboletellus, but we were not able to evaluate their phylogenetic positions in this

study. A more thorough investigation of the ecology of early-diverging Boletaceae is

needed to test the mycoparasitic hypothesis.

The evolution of gasteroid morphologies has long been thought to be an

evolutionary dead-end (Thiers 1984). The transition from pileate-stipitate to gasteroid

morphology involves the loss of ballistospory, and regaining this complex mechanism

once lost is likely difficult (Hibbett et al. 1997, Hibbett 2004, Sanchez-Garcia et al.

2020). While prior studies have not fully rejected the hypothesis gasteroidization is

irreversible (Hibbett 2004; Wilson et al. 2011; Sanchez-Garcia et al. 2020), there is no

previous evidence that suggests otherwise. However, our ancestral state reconstructions

under two coding schemes strongly supported transition to and from gasteroid forms.

This suggests that gain or loss of ballistospory may be a reversible condition. In our

analyses, transitions between pileate-stipitate, secotioid, and gasteroid forms indicate

that the secotioid condition is intermediate, as suggested by equal rates of transitions to

both gasteroid forms and pileate-stipitate forms. Moreover, inferred rates from pileate-

stipitate to both secotioid and gasteroid forms was almost zero, suggesting that the

secotioid condition is evolutionarily unstable. This result conflicts with previous

investigations of secotioid stability, particularly those of *Cortinarius* secotioid taxa,

where evidence suggests long term stability (Peinter et al. 2001).

Transitions between hymenophore organization parallels the sporocarp

morphologies. The "tubes with crosswalls" condition appears to be an intermediate

between lamellate and tubulate forms, with transitions to the latter being 1.3-5.8x

greater than in the reverse. However, the greatest transition rate was from lamellate to

tubulate forms, indicating that an intermediate morphology may not be necessary or,

like the secotioid condition, may be evolutionarily unstable and rarely persists.

Our data provide compelling evidence that lamellate hymenophores, color

changes, and gasteromycetization have evolved multiple times in the Boletaceae and are

reversible conditions. Based on the phylogenetic distribution of taxa, it is also clear that

membranous partial veils have evolved independently many times, although we did not

code this explicitly.

CONCLUSION

The Boletaceae underwent a rapid radiation and subsequent long period of phylogenetic

instability. These issues had long limited accurate assessment and analysis of trait

evolution and conclusive generic-level taxonomic frameworks. Prior molecular

phylogenetic studies based on a limited taxon sampling and a few loci generated trees

with poor deep-node support and many short branches. This study provides the first

statistically well-supported phylogeny for the Boletaceae using 1461 single copy genes

from 383 genomes sampled from specimens representing the most taxonomically and

geographically comprehensive dataset compiled to date. Our analyses show that the

Boletaceae likely arose in Gondwana prior to its breakup and present-day distributions

are in part due to vicariance. Episodic long-distance dispersal could not, however, be

ruled out for some current distributions. Morphological traits have resulted from of

convergent evolution and frequent, reversible recycling of traits that, together with the

rapid radiation, has long confounded attempts to achieve a natural classification within

the family. This study provides new genomic data and a stable phylogenetic framework

that will enable a robust, foundational taxonomy and provide a resource for further

analysis of trait evolution.

AKNOWLEDGEMENTS

We would like to think Dr. Roy Halling for the numerous specimens provided. This

work was supported by NSF-DEB awards DEB-2114785 to BD and DEB-0918591 to

TWH and DEB-1556338 to TWH and BD; National Geographic Society's Committee

for Research and Exploration grants (6679-99, 7435-03, and 8481-08 to TWH). JMM

was supported by grants from the ROM Governors and the Natural Sciences and

Engineering Research Council of Canada, Discovery Program.

CONFLICTS OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

LITERATURE CITED

Aitchison, Jonathan C., Jason R. Ali, and Aileen M. Davis. 2007. When and where did

India and Asia collide? Journal of Geophysical Research: Solid Earth. 112:B5.

Arora, David. 2008. California Porcini: Three New Taxa, Observations on Their

Harvest, and the Tragedy of No Commons. Economic Botany. 62:356–375.

Badou, Sylvestre A. et al. 2022. *Paxilloboletus* gen. nov., a new lamellate bolete genus

from tropical Africa. Mycological Progress. 21:243–256.

Bankevich, Anton et al. 2012. SPAdes: A New Genome Assembly Algorithm and Its

Applications to Single-Cell Sequencing. Journal of Computational Biology. 19:455–

477.

Berkeley, M. J. 1851. Decades of fungi. Decade XXXIV. Sikkim-Himalayan fungi

collected by Dr. Hooker.

Binder, M. et al. 2002. *Retiboletus*, a new genus for a species-complex in the Boletaceae

producing retipolides. Feddes Repertorium. 113:30–40.

Binder, Manfred et al. 2006. Molecular systematics and biological diversification of

Boletales. Mycologia. 98:971–981.

Brundrett, Mark C. et al. 2018. Evolutionary history of mycorrhizal symbioses and

global host plant diversity. New Phytologist. 220:1108–1115.

Bruns, Thomas D. et al. 1989. Evolution of mushroom mitochondrial DNA: *Suillus* and

related genera. Journal of Molecular Evolution. 28:349–362.

Bruns, Thomas D., et al. 1992. Evolutionary relationships within the fungi: analyses of

nuclear small subunit rRNA sequences. Molecular phylogenetics and Evolution. 1:231-

241.

Caiafa, Marcos V. et al. 2022. Polyphyly, asexual reproduction and dual trophic mode

in *Buchwaldoboletus*. Fungal Ecology. 56:101-141.

Castellano, Michael A. et al. 2016. *Kombocles bakaiana* gen. sp. nov. (Boletaceae), a

new sequestrate fungus from Cameroon. IMA Fungus. 7: 239–245.

Chakraborty, Dyutiparna et al. (June 2015). "A new generic record of Boletaceae for

Indian mycobiota". In: Current Research in Environmental & Applied Mycology 5, pp.

138–144.

Chen, Shifu et al. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 34:884–890.

Codjia, Jean Evans I. et al. 2023. Historical biogeography and diversification of ringless

Amanita (section *Vaginatae*) support an African origin and suggest niche conservatism

in the Americas. Molecular Phylogenetics and Evolution. 178:107644.

Corner, E. J. H. (1972). Boletus in Malaysia.

Cortés-Pérez, Alonso et al. 2023. *Pseudoboletus silvaticus* (Boletaceae, Basidiomycota),

a new species from Mexico. Phytotaxa. 589:27–38.

Crisp, Michael D., and Lyn G. Cook. 2011. Cenozoic extinctions account for the low

diversity of extant gymnosperms compared with angiosperms. New Phytologist.

192:997-1009.

Das, K. et al. (2015). *Boletus lakhanpalii*, a new species in Boletaceae from Sikkim

India with uncertain phylogenetic placement. Sydowia. 67:11–19.

Das, Kanad et al. 2016. New species of xerocomoid boletes (Boletaceae) from

Himalayan India based on morphological and molecular evidence. Mycologia. 108:753–

764.

Dentinger, Bryn T. M., Joseph F. Ammirati, et al. 2010. Molecular phylogenetics of

porcini mushrooms (*Boletus* section *Boletus*). Molecular Phylogenetics and Evolution.

57:1276– 1292.

Dentinger, Bryn T. M., Ester Gaya, et al. 2016. Tales from the crypt: genome mining

from fungarium specimens improves resolution of the mushroom tree of life. Biological

Journal of the Linnean Society. 117:11–32.

Dentinger, Bryn T. M. and Laura M. Suz. 2014. What's for dinner? Undescribed species

of porcini in a commercial packet. PeerJ 2:e570.

Drehmel, D. et al. 2008. Molecular Phylogeny and Biodiversity of the Boletes. FUNGI

1:4.

Elliott, Todd F., et al. 2019. A global review of the ecological significance of symbiotic

associations between birds and fungi. Fungal Diversity. 98:161-194.

Feng, Bang et al. 2012. DNA Sequence Analyses Reveal Abundant Diversity,

Endemism and Evidence for Asian Origin of the Porcini Mushrooms. PLOS ONE

7:e37567.

Feng, Yan-Jie, et al. 2017. Phylogenomics reveals rapid, simultaneous diversification of

three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary.

Proceedings of the national Academy of Sciences 114:E5864-E5870.

Fulgenzi, Tara D., Roy E. Halling, et al. 2010. *Fistulinella cinereoalba* sp. nov. and new

distribution records for *Austroboletus* from Guyana. Mycologia 102:224–232.

Fulgenzi, Tara D., Terry W. Henkel, et al. 2007. "*Tylopilus orsonianus* sp. nov. and

Tylopilus eximius from Guyana. Mycologia 99:622–627.

Fulgenzi, Tara D., Jordan R. Mayor, et al. 2008. "New species of *Boletellus* from

Guyana. Mycologia 100:490–495.

Futuyma, Douglas J. 1998. Evolutionary Biology. Quarterly Review of Biology 73.

Galante, Tera E. et al. 2011. 95% of basidiospores fall within 1 m of the cap: a field-and

modeling-based study. Mycologia 103:1175–1183.

Geml, József, et al. 2012. An arctic community of symbiotic fungi assembled by long-

distance dispersers: phylogenetic diversity of ectomycorrhizal basidiomycetes in

Svalbard based on soil and sporocarp DNA. Journal of Biogeography 39:74-88.

Golan, Jacob J., and Anne Pringle. 2017. Long-distance dispersal of fungi.

Microbiology Spectrum 5:5-4.

Grigoriev, Igor V. et al. 2014. MycoCosm portal: gearing up for 1000 fungal genomes.

Nucleic Acids Research 42:699–704

Grubisha, Lisa C. et al. 2001. Biology of the ectomycorrhizal genus *Rhizopogon*. V.

Phylogenetic relationships in the Boletales inferred from LSU rDNA sequences.

Mycologia 93:82–89.

Hackel, J., Henkel, T.W., et al. 2022. Biogeographic history of a large of

ectomycorrhizal fungi, the Russulaceae, in the Neotropics and adjacent regions. New

Phytologist 236:698-713.

Halling, Roy E. 1996. Boletaceae (Agaricales): Latitudinal biodiversity and biological

interactions in Costa Rica and Colombia". In: Revista de Biología Tropical 44:111–114.

Halling, Roy E., Dennis E. Desjardin, et al. 2014. New Porcini (Boletus sect. Boletus)

from Australia and Thailand. Mycologia 106:830–834.

Halling, Roy E., N.A. Fechner, G. Holmes, N. Davoodian. 2023. *Kagaria* (Boletaceae,

Boletoideae) gen. nov. in Australia: Neither a *Tylopilus* nor a *Porphyrellus*. Fungal

Systematics and Evolution 12:31-45.

Halling, Roy E., Nigel Fechner, et al. 2015. Evolutionary relationships of *Heimioporus*

and *Boletellus* (Boletales), with an emphasis on Australian taxa including new species

and new combinations in *Aureoboletus*, *Hemileccinum* and *Xerocomus*. Australian

Systematic Botany 28:1–22.

Halling, Roy E. et al. 2006. *Austroboletus mutabilis* from northern Queensland.

Muelleria: An Australian Journal of Botany 24:31–36.

Halling, Roy E., Mitchell Nuhn, et al. 2012. Affinities of the *Boletus chromapes* group

to *Royoungia* and the description of two new genera, *Harrya* and *Australopilus*.

Australian Systematic Botany 25:418–431.

Halling, Roy E., Todd W. Osmundson, et al. 2008. Pacific boletes: implications for

biogeographic relationships. Mycological Research 112:437–447.

He, Mao-Qiang et al. 2020. Notes, outline and divergence times of Basidiomycota.

Fungal Diversity 99:105-367

Han, Li H., et al. 2018. African origin and global distribution patterns: Evidence

inferred from phylogenetic and biogeographical analyses of ectomycorrhizal fungal

genus *Strobilomyces*. Journal of Biogeography 45:201-212.

Hay, William W., and Sascha Floegel. 2012. New thoughts about the Cretaceous

climate and oceans. Earth-Science Reviews 115:262-272.

Heinemann, P. 1951. Champignons recoltes au Congo belge par Madame M. Goossens-

Fontana I. Boletineae. Bulletin du Jardin botanique de l'État a Bruxelles 21:223– 346.

Henkel, Terry W, Aime MC, Chin M, Miller SL, Vigalys R, Smith ME. 2012.

Ectomycorrhizal fungal sporocarp diversity and discovery of new taxa in Dicymbe

monodominant forests of the Guiana Shield. Biodiversity and Conservation 21:2195-

2220

Henkel, Terry W. et al. 2016. New Boletaceae taxa from Guyana: *Binderoboletus* segoi

gen. and sp. nov., *Guyanaporus albipodus* gen. and sp. nov., *Singerocomus rubriflavus*

gen. and sp. nov., and a new combination for *Xerocomus inundabilis*. Mycologia

108:157–173.

Hibbett, David S. 2004. Trends in Morphological Evolution in Homobasidiomycetes

Inferred Using Maximum Likelihood: A Comparison of Binary and Multistate

Approaches. Systematic Biology 53:889–903.

Hibbett, David S. et al. 1997. Evolution of gilled mushrooms and puffballs inferred

from ribosomal DNA sequences. Proceedings of the National Academy of Sciences

94:12002–12006.

Hoang, Diep Thi et al. 2018. UFBoot2: Improving the Ultrafast Bootstrap

Approximation. Molecular Biology and Evolution 35:518–522.

Horak, E. 1977. New and rare boletes from Chile, *Nothofagus chilenos*, Ga*stroboletus*

valdivians, *Boletus loyita*, *Boletus putidus*. Boletín de la Sociedad Argentina de

Botánica.

Hosaka, Kentaro et al. 2008. Biogeography of Hysterangiales (Phallomycetidae,

Basidiomycota. Mycological Research. Phylogeography and Biogeography of Fungi

112:448–462.

Hu, Xiumian, et al. 2016. The timing of India-Asia collision onset–Facts, theories,

controversies. Earth-Science Reviews 160:264-299.

Husbands, Dillon R. et al. 2013. New species of *Xerocomus* (Boletales) from the Guiana

Shield, with notes on their mycorrhizal status and fruiting occurrence. Mycologia

105:422–435.

Jokat, Wilfried et al. 2003. Timing and geometry of early Gondwana breakup. Journal

of Geophysical Research: Solid Earth 108:2156-2202.

Kalyaanamoorthy, Subha et al. 2017. ModelFinder: fast model selection for accurate

phylogenetic estimates. Nature Methods 14:587–589.

Katoh, Kazutaka et al. 2017. MAFFT online service: multiple sequence alignment,

interactive sequence choice and visualization. Briefings in Bioinformatics 20:1160-

1166.

Kennedy, Peter G. et al. 2012. Scaling up: examining the macroecology of

ectomycorrhizal fungi. Molecular Ecology 21:4151–4154.

Kohler, Annegret et al. 2015. Convergent losses of decay mechanisms and rapid

turnover of symbiosis genes in mycorrhizal mutualists. Nature Genetics 47:410–415.

Landis, Michael J. et al. 2013. Bayesian Analysis of Biogeography when the Number of

Areas is Large. Systematic Biology 62:789–804.

Liimatainen, Kare et al. 2022. Taming the beast: a revised classification of

Cortinariaceae based on genomic data. Fungal Diversity 112:89–170.

Liu, Yunpeng et al. 2023. An updated floristic map of the world. Nature

Communications 14:2990.

Magnago, A.C, et al. 2017. *Fistulinella ruschii*, sp. nov., and a new record of

Fistulinella campinaranae var. scrobiculata for the Atlantic Forest, Brazil. Mycologia,

109:1003-1013

Matheny, P. Brandon et al. 2009. Out of the Palaeotropics? Historical biogeography and

diversification of the cosmopolitan ectomycorrhizal mushroom family Inocybaceae.

Journal of Biogeography 36:577–592.

Minh, Bui Quang et al. 2020. IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution 37:1530–

1534.

Miyauchi, Shingo et al. 2020. Large-scale genome sequencing of mycorrhizal fungi

provides insights into the early evolution of symbiotic traits. Nature Communications

11:5125.

Murrill, William A. 1909. The Boletaceae of North America—I". Mycologia 1:4–18.

Nei, Masatoshi et al. 2000. Molecular Evolution and Phylogenetics. Oxford, New York:

Oxford University Press.

Neves, Maria Alice et al. 2010. Study on species of *Phylloporus* I: Neotropics and

North America. Mycologia 102:923–943.

Nuhn, Mitchell E. et al. 2013. Phylogenetic overview of the Boletineae. Fungal Biology

117:479–511.

Peay, Kabir G., et al. 2010. Potential link between plant and fungal distributions in a

dipterocarp rainforest: community and phylogenetic structure of tropical

ectomycorrhizal fungi across a plant and soil ecotone. New phytologist 185:529-542

Peay, Kabir G., et al. 2012. Measuring ectomycorrhizal fungal dispersal:

macroecological patterns driven by microscopic propagules. Molecular Ecology

21:4122-4136.

Pegler, D. N. et al. 1981. A natural arrangement of the Boletales, with reference to spore

morphology. Transactions of the British Mycological Society 76:103–146.

Peintner, Ursula, et al. 2001. Multiple origins of sequestrate fungi related to *Cortinarius*

(Cortinariaceae). American Journal of Botany 88:2168-2179.

Raidl S. 1997. Studien zur Ontogenese Rhizomorphen Von Ektomykorrhiza. Bibl

Mycol 169:1–184.

Rambaut, Andrew et al. 2018. Posterior Summarization in Bayesian Phylogenetics

Using Tracer 1.7. Systematic Biology 67:901–904.

Ree, Richard H. et al. 2005. A likelihood framework for inferring the evolution of

geographic range on phylogenetic trees. Evolution 59:2299–2311.

Reguero, M. A., Goin, F. J. 2021. Paleogeography and biogeography of the Gondwanan

final breakup and its terrestrial vertebrates: New insights from southern South America

and the "double Noah's Ark" Antarctic Peninsula. Journal of South American Earth

Sciences 108:103358.

Ronquist, Fredrik. 1994. Ancestral Areas and Parsimony. Systematic Biology 43:0267–

274.

Ryberg, Martin et al. 2011. Dealing with incomplete taxon sampling and diversification

of a large clade of mushroom-forming Fungi. Evolution 65:1862–1878.

Sánchez-García, Marisol et al. 2020. Fruiting body form, not nutritional mode, is the

major driver of diversification in mushroom-forming fungi. Proceedings of the National

Academy of Sciences 117:32528–32534.

Sánchez-Ramírez, Santiago et al. 2015. High speciation rate at temperate latitudes

explains unusual diversity gradients in a clade of ectomycorrhizal fungi. Evolution

69:2196–2209.

Sato, Hirotoshi, Akifumi S. Tanabe, et al. 2017. Host shifts enhance diversification of

ectomycorrhizal fungi: diversification rate analysis of the ectomycorrhizal fungal genera

Strobilomyces and *Afroboletus* with an 80-gene phylogeny. New Phytologist 214: 443–

454.

Sato, Hirotoshi and Hirokazu Toju. 2019. Timing of evolutionary innovation: scenarios

of evolutionary diversification in a species-rich fungal clade, Boletales. New

Phytologist 222:1924–1935.

Schluter, Dolph et al. 2000. Ecological Character Displacement in Adaptive Radiation.

The American Naturalist 156:4–16.

Simões, M. et al. 2016. The Evolving Theory of Evolutionary Radiations. Trends in

Ecology & Evolution 31:27–34.

Singer, Rolf 1947. The Boletoideae of Florida with Notes on Extralimital Species III.

The American Midland Naturalist 37:1–135.

Singer, Rolf and Antonio P. Digilio. 1951. Pródromo de la flora agaricina argentina.

Lilloa 5–461.

Sitta, Nicola et al. 2008. Nationalization and Globalization Trends in the Wild

Mushroom Commerce of Italy with Emphasis on Porcini (*Boletus edulis* and Allied

Species. Economic Botany 62:307.

Smith, Martin R. 2020. Information theoretic generalized Robinson–Foulds metrics for

comparing phylogenetic trees. Bioinformatics 36:5007–5013.

Smith, Martin R. 2022. Robust Analysis of Phylogenetic Tree Space. Systematic

Biology 71:1255–1270.

Smith, Stephen A. et al. 2018. So many genes, so little time: A practical approach to

divergence-time estimation in the genomic era. PLOS ONE 13:e0197433.

Soulebeau, Anaëlle et al. 2015. The hypothesis of adaptive radiation in evolutionary

biology: hard facts about a hazy concept. Organisms Diversity & Evolution 15:747–

761.

Stecher, Glen et al. 2020. Molecular Evolutionary Genetics Analysis (MEGA) for

macOS. Molecular Biology and Evolution 37:1237–1239.

Šutara, Josef. 2014. Anatomical structure of pores in European species of genera

Boletus s.str. and *Butyriboletus* (Boletaceae)". Czech Mycology 66:157–170.

Tamura, Koichiro, Fabia Ursula Battistuzzi, et al. 2012. Estimating divergence times in

large molecular phylogenies. Proceedings of the National Academy of Sciences

109:19333–19338.

Tamura, Koichiro, Glen Stecher, et al. 2021. MEGA11: Molecular Evolutionary

Genetics Analysis Version 11. Molecular Biology and Evolution 38:3022–3027.

Tamura, Koichiro, Qiqing Tao, et al. 2018. Theoretical Foundation of the RelTime

Method for Estimating Divergence Times from Variable Evolutionary Rates. Molecular

Biology and Evolution 35:1770–1782.

Tao, Qiqing et al. 2020. Efficient Methods for Dating Evolutionary Divergences. The

Molecular Evolutionary Clock: Theory and Practice 197– 219.

Thiel, Martin, and Lars Gutow. 2005. The ecology of rafting in the marine environment.

II. The rafting organisms and community. Oceanography and marine biology 43:279-

418.

Thiers, Harry D. 1984. The Secotioid Syndrome. Mycologia 76:1–8.

Tremble, Keaton, et al. 2022. Contrasting continental patterns of adaptive population

divergence in the holarctic ectomycorrhizal fungus *Boletus edulis*". New Phytologist

237:295-309.

Tremble, Keaton, Laura M. Suz, et al. 2020. Lost in translation: Population genomics

and long-read sequencing reveals relaxation of concerted evolution of the ribosomal

DNA cistron. Molecular Phylogenetics and Evolution 148:106804.

Tremble K, R.E. Halling, T.W. Henkel, J.-M. Moncalvo & B.T.M. Dentinger. 2023.

Nomenclatural novelties. Index Fungorum no. 555.

https://www.indexfungorum.org/Publications/Index%20Fungorum%20no.555.pdf

Varga, Torda et al. 2019. Megaphylogeny resolves global patterns of mushroom

evolution. Nature Ecology & Evolution 3:668–678.

Wilf, Peter et al. 2019. Eocene Fagaceae from Patagonia and Gondwanan legacy in

Asian rainforests. Science 364:5139.

Wilson, Andrew W., Manfred Binder, and David S. Hibbett. 2011. Effects of gasteroid

fruiting body morphology on diversification rates in three independent clades of fungi

estimated using binary state speciation and extinction analysis. Evolution 65:1305-1322.

Wu, Gang, Bang Feng, et al. 2014. Molecular phylogenetic analyses redefine seven

major clades and reveal 22 new generic clades in the fungal family Boletaceae. Fungal

Diversity 69.1, pp. 93–115. ISSN: 1560-2745, 1878-9129.

Wu, Gang, Shingo Miyauchi, et al. 2022. Evolutionary innovations through gain and

loss of genes in the ectomycorrhizal Boletales. New Phytologist 233:1383–1400.

LEGENDS

Table 1. Morphological transition rates estimated with Bayestraits.

Figure 1. Selected photographs of Boletaceae collections from two of the most species-

rich regions that are newly sequenced in this study. A) *Boletus cervinococcineus*,

Singapore (BD616); B) Heimioporus punctisporus, Sarawak (BAKO2); C) unidenti-

fied Boletaceae, Vietnam (CTN-08-0007); D) unidentified Boletaceae (CTN-08-0029);

E) Spongiforma sp., Sarawak (BTNG10); F) Leccinum sp., Sarawak (SWK246); G)

Crocinoboletus laetissimus, Sarawak (SWK335); H) unidentified Boletaceae sp., Viet-

nam (DLT-08-0127); I) Boletellus sp., Sarawak (SWK356); J) unidentified Boletaceae,

Vietnam (CTN-08-0051); K) Tylopilus sp., Cameroon (BD655); L) Xerocomus sp. 9.

Cameroon (BD773); M) Fistulinella staudtii, Cameroon (BD848); N) Boletellus sp.,

Cameroon (BD714); O) Phylloporus cf. tubipes, Cameroon (BD719); P) Tylopilus sp.

8, Cameroon (BD816); Q) Xerocomus sp. 8, Cameroon (BD695); R) Boletus alliaceus,

Cameroon (BD697); S) Tubosaeta brunneosetosa, Cameroon (BD686); T) Tylopilus sp.,

Cameroon (BD716). Not to scale. All photos used with permission from Bryn

Dentinger.

Figure 2: Time-calibrated phylogeny of Boletaceae using 1764 BUSCO genes.

Topology is a summary coalescent of individual best ML gene trees using astral-hybrid.

Num- bers on branches are quartet probabilities. Branch lengths were converted to time

using the top 100 best gene trees estimated using SortaDate in RelTime. Inset a) map of

specimen origins with numbers of specimens from each geographic area. Inset

b) lineages-through-time plot calculated with the "ltt.plot" function in the R pack- age

"ape." The dashed line represents a constant birth-death rate. The shaded box indicates a

period of significant divergent increase from a constant birth-death rate indicative of a

rapid radiation.

Figure 3. Biogeographic reconstruction using BioGeoBEARS. Left-hand tree depicts 4-

state coding scheme (lig2h2t blue=Neotropical, blue=Paleotropical, yellow=North

Temperate, red=South Temperate) and right-hand tree depicts floristic region coding

scheme (pink=Chilean-Patagonian, green=Indo-Malesian, blue=African,

yellow=Holarctic, red=Novo-Zealandic, light blue=Neotropical, orange=Australian).

Pie charts indicate the proportional likelihood of each state at a node. Red and black

boxes indicate Phase I and Phase II Gondwanan diversification events, respectively.

Figure 4. Dispersal events and rates inferred from BioGeoBEARS. Top: Map depicting

phytogeographic zones that are colored and labeled on the map. Map was ren- dered

using the 'imago' R code (https://github.com/hrbrmstr/imago) to reproduce the

AuthaGraph world map projection (http://www.authagraph.com/top/?lang=en). Curved

arrows indicate inferred directional dispersal events and are colored by rate values

following the table (Bottom). The stroke weight of the arrows has been scaled to percent

of the maximum rate value following the values in the table. Bottom: Table of dispersal

rates inferred under a DIVAlike+j model in BioGeoBEARS. Source regions are at left

and destination regions are along the top. Values are colored along a scale from cool to

warm (red being maximum).

Figure 5. Distribution of likelihood frequencies of morphological traits. A-C)

Illustrations of coded traits and inferred transition rates. Arrows indicate direction of

transition with stroke weights scaled to maximum rate value. D-I) Frequency

distributions of state probabilities for six trait codings. Traits and codes are labeled on

the x-axis.

*Corresponding author(s): keaton.tremble@gmail.com

Trait	Transition	Mean rate	Median rate
Ballistospory			
	Yes to No	1.2303	1.0565
	No to Yes	39.8088	35.7478
Spore Ornamentation			
	Smooth to Not Smooth	0.7747	0.7498
	Not Smooth to Smooth	1.741	1.464
Stipe			
	Absent to Present	33.685	27.67
	Present to Absent	0.8415	0.6788
Sporocarp Habit			
	Secoitiod to Pileate-stipitate	51.64	51.47
	Pileate-stipitate to Secoitiod	0.8178	0.7172
	Secoitiod to Gasteroid	53.44	53.90
	Gasteroid to Secoitiod	34.63	31.05
	Pileate-stipitate to Gasteroid	0.8393	0.6749
	Gasteroid to Pileate-stipitate	40.78	36.85
Hymenophore Bruising			
	None to Blue	1.3515	1.2632
	None to Black	0.6921	0.6921
	Blue to None	5.3730	5.3540
	Blue to Black	0.1732	0.1205
	Black to None	3.2329	2.7385
	Black to Blue	1.0294	0.7281
Hymenophore Anatomy			
	Straight to Cross	0.6398	0.6092
	Straight to Lamellete	0.2054	0.1715
	Cross to Straight	3.694	2.960
	Cross to Lamellate	2.4131	1.6755
	Lamellate to Straight	5.483	5.004
	Lamellate to Cross	1.858	1.275

Southeast Asia

Africa

Range reconstuction of Boletaceae using paleo-coding
model = DEC + J, LnL = -373.98, Weighted AIC = 9.961182e-01

Range reconstuction of Boletaceae using floristic region-coding
model = DIVALIKE + J, LnL = -464, Weighted AIC = 0.54

	DESTINATION						
SOURCE	Holarctic	Neotropical	Indo-Malesian	African	Australian	Novo-zealandic	Chilean-Patagonian
Holarctic	0	4.28	17.42	4.66	2.72	2.22	1.76
Neotropical	1.26	0	1.16	2.34	0.22	0.6	0.34
Indo-Malesian	20.52	3.12	0	8	4.32	2.54	0.64
African	6.46	6.1	5.56	0	1.24	0	0.2
Australian	2.38	1.34	2.16	0.5	0	0.66	1.06
Novozealandic	0.28	0.48	0.12	0	1.4	0	0
Chile-Patagonian	0.74	0.32	0.04	0.04	0.84	0	0

A)

Smooth spores → 0.7747 → Ornamented spores

Ornamented spores → 1.741 → Smooth spores

B)

Straight tubes → 0.6398 → Tubes with cross walls

Tubes with cross walls → 3.694 → Straight tubes

Straight tubes → 0.2054 → Lamellate

Lamellate → 5.483 → Straight tubes

Tubes with cross walls → 2.413 → Lamellate

Lamellate → 1.858 → Tubes with cross walls

C)

Secotioid → 51.64 → Pileate-stipitate

Pileate-stipitate → 0.8178 → Secotioid

Secotioid → 53.44 → Gasteroid

Gasteroid → 34.63 → Secotioid

Pileate-stipitate → 0.8393 → Gasteroid

Gasteroid → 40.78 → Pileate-stipitate

D) Spore Ornamentation, A=Smooth, B=Not Smooth — Boletaceae.P.A., Boletaceae.P.B., ANOVA, P < 0.001

E) Hymenophore Anatomy, A=Straight Tubes, B=Tubes with Cross Walls C=Lamellate — Boletaceae.P.A., Boletaceae.P.B., Boletaceae.P.C., ANOVA, P < 0.001

F) Sporocarp Habit, A=Pileate-Stipitate, B=Secotioid C=Gasteroid — Boletaceae.P.A., Boletaceae.P.B., Boletaceae.P.C., ANOVA, P < 0.001

G) Boletaceae.P.0., Boletaceae.P.1., ANOVA, P < 0.001

H) Boletaceae.P.0., Boletaceae.P.1., ANOVA, P < 0.001

I) Boletaceae.P.A., Boletaceae.P.B., Boletaceae.P.C., ANOVA, P < 0.001