**Increased stromal densities of B cells, CD103+ cells, and CD163+ M2-like macrophages associate with poor clinical outcomes in BCG treated non-muscle invasive bladder cancer**

Benjamin Ravenscroft[1], Priyanka Yolmo[2,3], Stephen Chenard[2,3], Sadaf Rahimi[2,3], Keiran Pace[2, 3], Kartik Sachdeva[2,3], Tamara Jamaspishvilli[1], Hamid Ghaedi[1,3], Andrew Garven[1,3], Kathrin Tyryshkin[1,3], David M. Berman[1,3], Chelsea Jackson[1,4], D. Robert Siemens[3,5], Madhuri Koti*[2, 3]

[1]Pathology and Molecular Medicine, Queen's University, Kingston, ON, Canada;[2]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON, Canada; [3]Cancer Research Institute, Queen's University, Kingston, ON, Canada; [4]Department of Pathology, University of Manitoba; [5]Department of Urology, Queen's University, Kingston, ON, Canada.

**Key words:** B cells, NMIBC, BCG immunotherapy

**\*To whom correspondence should be addressed**

 e-mail: kotim@queensu.ca

**Abstract**

Non-muscle invasive bladder cancer (NMIBC) constitutes a significant clinical challenge, with over 50% of patients experiencing poor clinical outcomes in the form of early recurrence or progression following treatment with Bacillus Calmette-Guerin (BCG) immunotherapy. The pre-treatment tumor immune microenvironment (TIME) is an established determinant of response to BCG. This study explores the spatial profiles of CD79a+ B cells, CD163+ M2-like macrophages, proliferating and tissue-resident phenotypes of T cells, along with PD-1/PD-L1 checkpoint expression in pre-BCG treatment tumors of 173 patients (139 males, 34 females). Multiplex immunofluorescence staining of a tumor tissue microarray, revealed elevated infiltration of CD79a+ B cells, CD163+ M2-like macrophages, CD103+ cells, and CD8+ T cells at the tumor invasive margins. Increased epithelial PD-L1 immune-checkpoint expression in tumors was observed in female and male patients who exhibited significantly shorter recurrence-free survival (RFS). Importantly, high CD79a+ B cell density in BCG-treated females in both stromal and epithelial compartments exhibited significantly shorter RFS and progression-free survival compared to males. Stromal CD79a+ B cell density was positively correlated with M2-like macrophages, CD8+ T cells, CD103+ cells and PD-1 expressing cells. CD79a+ B cells, CD103+ cells, and M2-like macrophage density were associated with higher grade and enriched in basal subtype tumor. This study highlights the significance of an understudied role of B cells and their cellular neighborhoods in the pre-treatment TIME and BCG-therapy response. Overall, findings from this study underscore the importance of considering sex-related immunobiological differences in the stromal compartments of bladder tumors towards the development of optimal therapeutic targeting strategies.

## Introduction

Approximately 75% of newly diagnosed bladder cancer patients present with NMIBC. While the incidence of NMIBC is four times higher in males, female patients with NMIBC often present with a more advanced stage and display earlier recurrence [1-4]. Patient age, tobacco smoking, and occupational exposure to chemicals are some of the major risk factors associated with NMIBC [5-7]. Bacillus Calmette-Guérin (BCG), a live attenuated strain of *Mycobacterium bovis*, is the adjuvant treatment for patients with intermediate or high-risk NMIBC [8, 9]. Intravesical BCG therapy entails 6 weekly instillations after transurethral resection of bladder tumor (TURBT), followed by a maintenance schedule of 3 weekly instillations every 3-6 months for 1-3 years depending on the risk stratification as defined by the American and European Urological Association [9]. Despite its proven efficacy, over 50% of the patients receiving BCG immunotherapy experience recurrence or become unresponsive to treatment [5, 6]. Second line treatments such as programmed cell death protein-1 receptor/ligand (PD-1/PD-L1) immune checkpoint axis inhibitors and replication deficient adeno viral gene-delivery vectors (Adstiladrin) are under trials for BCG unresponsive/recurrent disease [10-12]. While some have demonstrated modest benefits, these newer treatment approaches do not increase primary response to BCG and are not indicated in the BCG naïve setting [13-15].

The pre-treatment TIME is a major determinant of response to immunotherapy across various cancers [11, 16, 17]. Recent studies have established that the pre-treatment TIME plays an important role in determining BCG therapy response in patients with NMIBC [18-21]. This is especially important when considering BCG therapy, which involves a live bacterial preparation and its efficacy is dependent on multiple factors including the dose, viability of bacteria at the time of instillation, immune physiological status of the host, bacterial contact with the urothelium and uptake by cells in the bladder mucosal immune environment. Based on the infiltration profiles of CD8+ cytotoxic T cells, FoxP3+ $T_{reg}$ cells, and the expression patterns of PD-1/PD-L1+ immune checkpoints, both in pre-BCG treatment and recurrent tumors, a potential role of adaptive immune resistance has also been suggested [19]. Additionally, elevated T cell exhaustion has been reported as a major factor underlying poor response to BCG [20]. In our previous report, conducted on the whole transcriptomic profiles of the UROMOL

cohort (n=460) and spatial immunophenotyping of tumors from a large cohort assembled at the

Kingston Health Sciences Center (KHSC, n=332), we showed increased infiltration of B cells, M2-

like macrophages, and higher expression of PD-L1 in the pre-treatment TIME of both female

and male patients experiencing early recurrence, irrespective of the treatment type [18].

Notably, female patients with high-grade disease in both cohorts exhibited significantly shorter

progression free survival (PFS) and had higher infiltration of B cells and M2-like macrophages in

their pre-treatment tumors compared to their male counterparts. In the current study, we

characterized the association between immune cell spatial profiles in the BCG-naïve TURBT

specimens of female and male patients exhibiting differential response to BCG immunotherapy,

from our previously reported KHSC cohort [18]. Given the established differences in sex-

associated B cell immunobiology and the need for improved patient stratification for increased

benefits from BCG immunotherapy, we further evaluated patient sex-associated pre-BCG tumor

immune profiles and clinical outcomes.

**METHODS**

**Ethics statement**

This study was approved by the Queen's University Health Sciences and Affiliated Teaching

Hospitals Research Ethics Board.

***In silico* analysis of *CD79A, CD103* and *CD163* gene expression profiles in transcriptomic**

**profiles of tumors from BCG naïve high-risk patients**

Based on our previous findings [18] on the associations between tumor infiltrating CD79a+ B

cells and M2-like macrophage densities and overall poor clinical outcomes in patients with

NMIBC, irrespective of the treatment type, we first evaluated the publicly available normalized

bulk RNA-sequencing based tumor whole transcriptome profiles of 283 patients with NMIBC

that was recently generated by Jong *et al.,* [21]. The patients in this cohort were stratified into

different quartiles according to their *CD79a* and *CD163* gene expression levels. Patient clinical

demographics and outcome data was extracted from the supplementary information provided

by de Jong *et al.*, [21] These clinical attributes were then combined with the gene expression

profiles of the primary tumors. Subsequently, an analysis was conducted using the time to

disease progression and the progression status in R version 4.2.0. To visualize the relationship

between target *gene* expression and survival, Kaplan-Meier survival curves were generated by

employing a log-rank statistical analysis method and utilizing the survival and survminer

packages in R.

**KHSC-BCG Patient Cohort and Clinical Data**

Given their critical roles in mucosal immunobiology we then asked whether the spatial profiles

of CD79a+ B cells, CD163+ M2-like macrophages and other immune cells of interest (see below)

associate with variable response to BCG in female and male patients with NMIBC. Of the 332

patients reported in our previous study [18], a total of 173 patients (KHSC-BCG:139 males and

34 females) who received at least one induction round of BCG, defined as at least 5/6

instillations, and had no BCG treatment prior to entering the cohort, were selected for inclusion

in this study (**Table 1**). Adequate BCG treatment was defined as an initial induction round of at

least 5/6 instillations, followed by a maintenance round of at least 2/3 instillations [9]. Patients

were assessed for BCG therapy response and classified as either BCG refractory, BCG

responders, or an in-between state of therapy response. The BCG refractory state was defined

according to the recent consensus guidelines put forth by the European Association of Urology

[6], and BCG responders were patients without any high-grade recurrence within the two years,

following their first adequate BCG induction (**Table 1**)[8]. Due to our cohort size limitations and

the differences in clinical treatment versus ideal clinical trial treatment pathways, we used high-

grade recurrence as an endpoint for time-to-event analysis. The initiation point for time-to-

event analyses is the diagnosis of TURBT immediately prior to the initiation of BCG treatment.

Progression events were defined as recurrence to stage T2 or higher.

Baseline clinical data and NMIBC tissue were collected for eligible patients (**Table 2**). All the

tumor tissue specimens included in the study were index TURBT specimens. The median follow-

up from surgical resection (TURBT) was 54.3 months (95% CI 50.2 to 59.5 months) for the

cohort. 124 (71.7%) of the cohort were BCG responders and 37 (21.4%) were BCG refractory.

The entire cohort had a 24-month high-grade recurrence free survival (RFS) rate of 70% (95% CI

63.1% to 77.5%). The median time to HG recurrence was 7.6 months (95% CI 6.3 to 11.8

months), and the median follow-up time for BCG responders was 53.4 months (95% CI 47.8 to

59.5 months).

**133    Multiplex Immunofluorescence Staining**

Three panels of antibodies were used to stain the tissue microarrays (TMAs; **Supplementary**

**data, Figure S1**). The first panel consisted of antibodies identifying FoxP3+, CD8+, CD3+, and

Ki67+ cells. The second panel PD-1+, PD-L1+, and CK5+ cells and the third panel included

antibodies identifying GATA3+, CD103+, CD163+, and CD79a+ cells. The following cell types

were evaluated for expression in both the epithelium and stroma: proliferating and non-

proliferating cytotoxic T cells (CD3+CD8+Ki67+/-), cytotoxic regulatory and non-regulatory T

cells (CD3+FoxP3+/-), immune checkpoints (PD-1+, PD-L1+, PD-L1+ expressed on CK5+ cancer

cells and immune cells), CD103+ cells, CD163+ M2-like macrophages and CD79a+ B cells.

Staining and scanning of the TMA was performed using multiplex immunofluorescence (mIF)

staining at the Molecular and Cellular Immunology Core (MCIC) facility, BC Cancer Agency as

described earlier [18]. The antibodies utilized were supplied by Biocare Medical (Pacheco, CA,

USA) and distributed by Inter Medico (Markham, ON, CAN).

**147    Automated scoring and manual validation of immune cell density and distribution in mIF**

**148    stained tumors**

The stained TMA sections were scanned using the Vectra 3.0.5 multispectral imaging system

(PerkinElmer, MA, USA) and subsequently analyzed in InForm 2.4.0 (PerkinElmer, MA, USA).

The scanned TMAs underwent tissue segmentation into epithelium and stroma utilizing

InForm's tissue segmentation algorithm, specifically examining nuclear DAPI. Phenotyping was

performed by training three independent algorithms on ten randomly selected cores from the
cohort. The software identified positive pixels for all selected immune markers using each of
the three independent algorithms. The average was taken across the three algorithms for all
three sets of immune markers in each core. Cores containing ≤ 25% of the tissue were excluded
from subsequent analyses. Standard deviation between the three algorithms was calculated for
each of the immune markers within both the epithelial and stromal compartments. Outliers
were identified and cross-referenced with the composite image of the corresponding TMA core
to verify the discrepancy before excluding the outlying data points from further analysis. Any
quantifications that included misidentification of histological artifacts as immune cells by any of
the three algorithms were also excluded from analysis. Algorithms that consistently over-called
or under-called the immune marker quantification were excluded from analysis. Further visual
validation of the automated scoring was performed for randomly selected TMA cores.

**Statistical Analysis**

All analyses were conducted using either R version 4.1.2 or Python 3.9.12. Analysis of cell
population composition was performed using unpaired non-parametric Mann-Whitney tests.
The corrplot package was utilized to calculate and visualize the Spearman rank correlations of
various cell type, due to its non-parametric nature making it better suited for data lacking
normality. The Kaplan-Meier method and log-rank test were performed to determine the
association between time to high-grade recurrence and dichotomized infiltration of immune
cell phenotypes.

Differences in infiltration levels of immune cells between BCG refractory patients and BCG
responders were tested and presented with violin plots. Prior to statistical analysis, the
normality condition was checked for the density of each phenotype across the whole cohort
using the Shapiro-Wilk test. Failing to satisfy this condition, the Wilcox-rank sum test was used
to test for differences in immune cell infiltration in BCG responders and non-responders as it
does not assume normality.

To dichotomize infiltration of specific immune cell phenotypes, a combination of graphical
diagnostic plot methods as presented by Williams *et al.* and bootstrapping were used
([Microsoft Word - Technical report 79 cover mandrekar 6-06.doc (mayo.edu)](). Unique densities
of each immune phenotype were used to create indicator variables that could fit Cox models,
from which log-likelihood values could be extracted. These values were plotted, and LOESS
curve fitting was performed using the stats package. The discrete infiltration values closest to
the five most visually unique peaks were selected. Bootstrapping was performed using a
univariable cox model that compared the high/low infiltration groups for each of the six
possible cut point values previously identified [18]. The validated cut point was the
dichotomization value with the greatest frequency of maximum log-likelihood after
bootstrapping. Valid cut points were used to create dichotomized Kaplan-Meier curves
comparing RFS and PFS between high and low infiltration groups for various immune cell
phenotypes. The dichotomized Kaplan-Meier curves were typically truncated prior to the
number at risk in any group dropping below 5.

The distribution of clinically prognostic variables previously evaluated [18], including subtype,
grade, stage, AUA risk score, sex, and age, was rigorously examined for each dichotomized
immune cell phenotype. Statistical analysis involved employing Pearson's $\chi^2$ test or Fisher's
Exact test in cases where all assumptions were not met. P-values were calculated both without
adjustment for multiple testing and with adjustment for multiple testing using the Bonferroni
method.

The clinical impact of immune cell and immune checkpoint populations were analyzed using a
univariable Cox proportional hazards (CPH) model. The proportional hazards (PH) assumption
was verified using the Schoenfeld test, computed using the survival package. A nonsignificant (p
> 0.05) test statistic indicates that the PH assumption is met. Multivariable models were
created for each immune cell phenotype with age, sex, AUA risk score, and BCG treatment
amount, dichotomized into induction only and adequate BCG treatment groups, as covariates.
P-values were calculated with and without adjusting for multiple testing using the Bonferroni
method for correction of multiple testing. Results are reported graphically in a forest plot and
hazard ratios (HR) for immune cell density variables are scaled by the one standard deviation of

the given immune cell population when graphically reported. Scaling the hazard ratios by each
immune cell population's standard deviation in the cohort was performed to enable clearer
understanding of relative association between each cell type and RFS/PFS, as well as
comparison of each different cell type's effect size. Some cell types have a very low SD in the
cohort, such as stromal CK5+PD-L1+ (SD = 5.06), while others, such as stromal CD79a+ B cells
(SD = 137.13), have a very large standard deviation. Subsequently, reporting HR scaled by SD
(hereafter denoted $HR_{SD}$) shows effect size relative to expected magnitude of change in cell
infiltration for a given cell type [22].

**RESULTS**

***Increased expression of CD79A and CD163 genes in pre-treatment tumors is associated with***
***shorter progression free survival following treatment with BCG***
We first determined the association between expression levels of *CD79A* and *CD163* transcripts
and clinical outcomes using the publicly available and recently published large independent
cohort of BCG treated patients [21]. In this cohort of 283 patients, 20% of the patients were
female and the entire cohort was BCG-naive high-risk NMIBC with 34% of the patients
exhibiting progression after treatment with BCG. Kaplan-Meier survival analysis revealed that
high expression of *CD79A and CD163 are* associated with significantly shorter PFS following BCG
therapy (**Supplementary data, Figure S2**). In this cohort, this association was observed in both
male and female patients (not shown).

***High CD79a+ B cell infiltration in stromal and epithelial regions is associated with early***
***recurrence and progression following BCG therapy***

The patient clinical demographics of KHSC-BCG cohort are presented in **Table 2**. Of the 173
patients in the KHSC-BCG cohort, 72% patients had a diagnosis of high-grade disease. The
cohort reflected a female to male ratio of 1:4. Approximately 47% of patients in this cohort
received adequate BCG and risk stratification was performed as per the recent IBCG guidelines

(**Supplementary data, Figure S3**). High-grade recurrence was observed in 33% patients in this

cohort with 37 patients classified as BCG refractory and 124 as BCG responders. In this study,

we used CD79a as the pan B cell marker. Kaplan-Meier survival analysis revealed that both

female and male patients with high epithelial and stromal CD79a+ B cells in their pre-treatment

tumors had significantly shorter RFS and PFS (**Figure 1A, 1B** and **S4**). Furthermore, comparison

of the survival profiles specific to male and female patients revealed that females with high

stromal density of CD79a+ B cells in their pre-BCG tumors experienced worse RFS and PFS

(**Supplementary data, Figure S5**) compared to males. Quantification of immune cells in the

stromal regions showed a significantly higher density of CD79a+ B cells (p=0.015) and CD163+

M2 macrophages (p=0.027) in BCG refractory patients compared to the responders (**Figure 1C**

**and 1D**). Similar analysis of CD163+ cell densities within the epithelial compartment did not

show any significant differences (**Supplementary data, Figure S6**).

***Stromal CD79a+ B cell density is positively correlated with M2-like macrophages, CD8+***

***cytotoxic T cells, CD103+ cells and PD-1 immune checkpoint expression***

We then evaluated the correlations between cell types and immune checkpoint expression in

both the stromal and epithelial compartments. Spearman correlation analysis revealed

significantly positive correlations between CD79a+ B cells and CD163+ M2-like macrophages

(r=0.74, p=1x10$^{-7}$), CD3+ CD8+ cytotoxic T cell (r=0.55, p=4x10$^{-4}$), and PD-1 immune checkpoint

expressing cells (r=0.55, p=4x10$^{-4}$) in the stromal compartment of the BCG refractory group

(**Figure 2A**). Similar correlations were observed in the BCG responder group in which CD79a+ B

cells were significantly positively correlated with CD163+ M2 macrophages (r=0.61, p=3x10$^{-14}$),

CD3+CD8+ cytotoxic T cells (r=0.68, p=2x10$^{-10}$), and both PD-1 (r=0.67, p=2x10$^{-16}$) and PD-L1

(r=0.54, p=7x10$^{-11}$) immune checkpoints (**Figure 2B**). Importantly, within the BCG refractory

group, correlations between immune cell types in the epithelial compartments were not

significant as observed within the stromal region (**Figures 2C and 2D**). In the stromal regions of

tumors from BCG responder and refractory groups, CD163+ macrophages and CD8+ cytotoxic T

cells also depicted moderate positive correlation with each other (r=0.67-0.68). Cytotoxic T cell

density was highly correlated with the expression of PD-1 immune checkpoint (r=0.87-0.88) in
both the groups.

*Increased stromal density of CD8+ T cells, CD103+ cells and M2-like macrophages is*
*associated with early recurrence post-BCG*

Investigation of other immune cells in the stromal and epithelial compartment of each tumor
core demonstrated a higher density of CD163+ M2-like macrophages, CD3+CD8+ cytotoxic T
cells, and CD103+ cells in the BCG non-responder group. Kaplan-Meier survival analysis
revealed a significant association between high stromal density of immune cells (M2-like
macrophages, cytotoxic T cells and CD103+ cells) and overall shorter RFS (HR=2.87, p < 0.001,
**Figure 3**).

Spearman correlation analysis revealed that PD-1+ cell density is highly and significantly
correlated with CD103+ (r=0.64-0.83), CD163+ M2-like macrophages (r=0.62-0.65), and both
Ki67+ and Ki67- CD8+ T cells (r=0.8-0.85). Furthermore, significant correlations were observed
between PD-1 and CD103+ cells (r=0.64-0.83), in the stromal compartment indicate heightened
immune checkpoint expression in these cells. Similarly, a strong correlation (**Figure 2A and 2B**)
was found between PD-1 expression and both proliferating (Ki67+) and non-proliferating (Ki67-)
cytotoxic CD8 + T cells (r=0.8-0.85). High expression of PD-1 could potentially suggest T cell
activation in BCG responders and a state of immune exhaustion in BCG refractory patients.
These findings underscore the considerable variability in immune responses to BCG
immunotherapy.

*Elevated density of stromal CD79a+ B cells, CD8+ T cells, CD103+ cells, and M2-like*
*macrophages is associated with increased disease stage and heightened AUA risk score*
To determine the stage and risk associated immune cell profiles, we performed a Pearson's $\chi^2$
analysis (**Supplementary data**, **Table S1**). Among the patients who received BCG treatment, a
higher density of CD79a+ B cells, CD103+ cells, and M2-like macrophages in the stromal
compartment of their tumors was associated with higher grade ($\chi^2$(1, N=173) = 10.23, p = 0.001,

$\chi^2(1, N=173) = 5.54$, p = 0.019, $\chi^2(1, N=173) = 5.95$, p = 0.015, respectively) and stage ($\chi^2(1,$

N=173) = 6.11, p = 0.013, $\chi^2(1, N=173) = 9.55$, p = 0.002, respectively), except for the higher

CD79+ B cell density. The basal subtype was notably more prevalent in patients with an

elevated stromal density of CD79a+ B cells and CD103+ cells, as confirmed by a Fisher's Exact

test (p = 0.016 and 0.025, respectively).

***Univariable and multivariable analysis reveals high pre-BCG tumor immune cell density is***

***associated with a higher risk of recurrence and progression***

A CPH based analysis revealed significant associations between immune cell infiltration profiles

and clinical outcomes (**Figure 4**). Specifically, in the epithelium PD-1+ ($HR_{SD}$=1.223, p=0.026),

CD79a+ B cells ($HR_{SD}$=1.198, p=0.024), and CK5+ PD-L1+ cancer ($HR_{SD}$=1.241, p=0.019) cell

density was associated with worse RFS (**Supplementary data Table S2**). CD79a+ B cells

($HR_{SD}$=1.324, p=0.022), CD163+ M2-like macrophages ($HR_{SD}$=1.578, p < 0.001), CK5+PD-L1+

cancer cells ($HR_{SD}$=1.391, p=0.01), and PD-1+ immune checkpoint expressing cells ($HR_{SD}$=1.352,

p=0.017), in the epithelial compartment, were also associated with worse PFS (**Supplementary**

**data Table S3**).

In the stroma, CD3+CD8+ cytotoxic T cells ($HR_{SD}$=1.258, p=0.008), CD79a+ B cells ($HR_{SD}$=1.315, p

< 0.001), CD163+ M2-like macrophages ($HR_{SD}$=1.274, p=0.009), CD103+ cells ($HR_{SD}$=1.322,

p=0.014), and PD-1+ immune checkpoint expressing cells ($HR_{SD}$=1.415, p=0.006) were

associated with worse RFS. Only CD79a+ B cells ($HR_{SD}$=1.508, p=0.002), and CD163+ M2-like

macrophages ($HR_{SD}$=1.438, p=0.029), in the stromal compartment, were significantly associated

with PFS.

In a multivariable model considering sex, age, AUA risk score, and BCG treatment adequacy as

covariates, none of the immune cells in either the stromal or epithelial compartment were

significantly associated with a change in RFS after adjustment for multiple hypothesis testing.

CD79a+ B cells, in the stromal compartment, were however significantly associated with worse

PFS ($HR_{SD}$=1.663, p=0.03), with and without Bonferroni correction.

**Discussion**

Recurrence or progression to muscle invasive disease following BCG immunotherapy remain the major challenges in the management of patients with NMIBC. To determine the factors that lead to a poor response, studies have extensively examined the treatment-naïve TIME in tumors collected at TURBT from patients with NMIBC. Specifically, these prior investigations have focused on T cells and myeloid cells and have significantly advanced the current state of knowledge on the complex nature of the TIME in NMIBC tumors [19, 20, 23, 24]. However, clinical translation to actionable biomarkers is still unavailable. Additionally, B cells that constitute the key cellular mediators of mucosal immunity, via their dual functions as antigen presenting and antibody producing cells, remain understudied in the context of response to BCG immunotherapy.

A significant and novel observation from the current study is the association between a high density of stromal and intra-epithelial B cells in the BCG naïve tumors from patients who experienced early recurrence or progression to muscle invasive disease. Our findings demonstrate a robust association between CD79a+ B cell infiltration in both stromal and epithelial compartments, which is more pronounced in high-grade tumors from female patients who experienced early recurrence and disease progression following BCG therapy. These observations align with current literature highlighting the importance of CD79a+ B cells in driving tumor progression and fostering resistance to BCG therapy, particularly within the stromal milieu [21].

A higher stromal B cell density and a positive correlation between densities of CD79a+ B cells, M2-like macrophages, CD103+ cells, PD-1 immune checkpoint expression, compared to that of the epithelial compartment, is indicative of chronic inflammation and potential exhaustion within or outside of either stromal immune cell aggregates or bladder mucosa associated lymphoid tissue referred to as tertiary lymphoid structures (TLSs). Mucosa associated TLSs evolve as a result of chronic inflammation such as carcinogen exposure or persistent antigenic stimuli including recurrent urinary tract infection or autoimmune mediated Hunner's lesion interstitial cystitis [25]. Supporting this, our previous report demonstrated the compounding

effects of aging and chronic carcinogenesis associated inflammation on increased B cell
recruitment and TLS formation in the urinary bladder [26, 27]. We also showed the higher
expression of multiple immune exhaustion associated proteins within TLSs and their peri-
tumoral location associated with higher stage and poor response to BCG [26, 28]. A bladder
tumor associated chronic inflammatory state also suggests an exhausted pre-BCG host immune
system. Consequent to such systemic immune exhaustion, exposure to BCG bacteria at weekly
intervals during the induction phase of treatment potentially intensifies the chronic systemic
inflammatory state, specifically in patients with higher stromal B cell profiles in their pre-BCG
tumors. This is supported by a recent study conducted by Strangaard *et al.,* [20] which
examined pre- and post- BCG paired tumor samples and showed T cell exhaustion as the factor
underlying a BCG unresponsive disease. Overall, these findings suggest that in a subset of
patients who do not benefit from BCG, signatures of innate and adaptive immune exhaustion
within the pre-BCG tumor immune profiles could inform patient selection for
immunomodulatory BCG treatment.

Most recently, de Jong *et al.,* reported the presence of three BCG response subtypes (BRS1,
BRS2, and BRS3) of pre-treatment tumors that show a distinct association with progression of
NMIBC to muscle invasive disease [21]. Specifically, in this study based on bulk RNA-sequencing
of tumors, the BRS3 subtype enriched with immune gene signatures was shown to associate
with poor outcomes following BCG treatment. Tumors with a BRS3 subtype (increased
expression of genes associated with B cell, M1/M2 macrophage, CD8+ T cell, $T_{reg}$ function and
immune exhaustion) exhibited the most aggressive behavior and early progression. Notably,
luminal subtype tumors that are known to exhibit a low immune infiltration profile, were not
enriched for a BRS3 gene signature. Unsurprisingly, the post-BCG recurrent tumors also carried
the BRS3 phenotype and a basal tumor subtype [21]. In the current study, we used the pan B
cell marker - CD79a, to identify tumor infiltrating B cells. Evaluation of bulk RNA-Seq profiles
from the publicly available tumor transcriptomic profiles of 283 patients generated by de Jong
*et al.,* validated our findings on the increased expression of *CD79A* and *CD163* transcripts in
tumors from patients who experienced early progression to muscle invasive disease following
BCG therapy.

Our counterintuitive findings showing inverse correlations between higher density of cytotoxic

T cells and poor response to BCG immunotherapy, are also in alignment with previous literature

suggesting the development of adaptive immune resistance following treatment in BCG non-

responders [19, 21]. Previous studies have highlighted the involvement of regulatory B cell

subsets in fostering an immunosuppressive environment conducive to immune evasion and

tumor progression [29-31]. Remarkably, our study corroborates these prior findings,

establishing a positive correlation between CD79a+ B cells and CD163+ M2-like macrophages,

CD8+ Ki67+ proliferating cytotoxic T-cells, and PD-1 expression within the stromal compartment

of BCG-refractory patients.

The notable correlation between CD79a+ B-cells, CD8+ T cells and the PD-1 immune checkpoint

protein suggests a potential regulatory role exerted by immune checkpoint mechanisms in

modulating the functional activity of cytotoxic T cells in response to BCG therapy. Similarly, the

presence of CD8+ T cells and CD103+ cells implies an actively mounting immune response

against the tumor. However, their correlation with shorter RFS strongly suggests the possibility

of dysfunction or exhaustion, potentially arising from chronic antigen exposure or inhibitory

signals from other immune cell subsets. Immune cells expressing the $\alpha_E$ integrin CD103 (mainly

$T_{RM}$ cells) are widely distributed across mucosal surfaces [32]. In this study we did not use a T

cell phenotypic marker to define CD103+ cell, however, their significantly high correlation with

CD8+ cells and previous reports confirming the T cell specific intra-tumoral expression pattern

of CD103 and the chronic nature of the disease, is suggestive of their $T_{RM}$ phenotype [33, 34].

Indeed, the expression of CD103 has been reported on multiple cell types (Tregs, tissue resident

memory T cells) within the stromal compartment across various cancer types [35]. Future

studies should further characterize their effector memory and central memory functional

states.

Overall, these findings from multiple independent cohorts confirm that spatial organization and

density of B cells within the pre-BCG TIME could be developed into a robust clinically feasible

assay for application on TURBT specimens prior to initiation of BCG treatment.

An important and novel finding from our study is the significantly worse RFS and PFS in BCG
treated female patients with high stromal and intra-epithelial B cell density compared to that of
males. As reported by us and others, the physiologically higher B cell frequency in females
compared to that of males is likely a key factor underlying the higher degree of their trafficking
to the bladder TIME upon chronic antigenic stimulation. Previously, we have reported a
significantly shorter PFS in females based on evaluation of two large independent cohorts of
patients with NMIBC. We also showed that irrespective of the treatment type, a higher pre-
treatment intra-tumoral B cell density is associated with early RFS and PFS in patients with
NMIBC [18].

While these findings indicate that B cell proportions increase with disease progression and due
to the immune stimulatory effects of BCG bacteria, we speculate that upon treatment with
cytotoxic chemotherapy such as gemcitabine-docetaxel chemotherapy or immune checkpoint
inhibition, such responses may lead to a favorable response via immunogenic cell death
mediated events. Such associations have been reported in other cancers and in muscle invasive
bladder cancer, where higher pre-treatment B cell density associated with positive outcomes
following treatment [36-38]. Finally, in this study, we observed a significant association
between increased density of CDK5+ PDL-1+ cells indicative of an aggressive cancer cell
phenotype and shorter PFS. Constitutive expression of PD-L1 on cancer cells could result from
mutations in DNA damage repair genes or could be induced via the phenomenon of adaptive
immune resistance during disease progression [39]. While the mutational profiles of tumors
used in our study were not available, previous studies have reported an enrichment of DNA
damage repair gene mutations in high-grade NMIBC tumors [40, 41].

This study is not without limitations. We acknowledge that tumor cores of 0.6 - 1 mm diameter
in a TMA do not capture the heterogeneity within the larger tumor landscape. Recently,
Lindskrog *et al.*, [23] reported the presence of multiple tumor subtypes within a single tumor
specimen. This finding supports a potential caveat of TMA based studies and the need for
surveying whole tumor sections to derive a comprehensive spatial organization. However, with
regard to our previous report confirming stromal B cell infiltration profiles, and TLSs in whole
tumor sections of BCG non-responders, we posit that a careful selection of tissue cores

capturing tumor invasive margins is required for larger cohort-based biomarker studies. A

deeper characterization of the B cell subsets and their cellular neighborhoods using whole

tumor sections is also needed to gain further insights into their antigen presenting and/or

antibody producing functions. Finally, exploration of the finer spatial relationships between the

several immune cell populations and tumor invasive margins within pre- and post-BCG

treatment tumors is critical to determine if the localization of specific cellular phenotypes

associate with BCG outcome.

To conclude, the findings presented in this study advance the current understanding of the

immune landscape characterizing the NMIBC microenvironment and its association with patient

responses to BCG immunotherapy. Ultimately, these findings have implications not only for the

development of novel therapeutic strategies targeting B cell specific exhaustion but also to

inform approaches for precise sequencing of drugs targeting the immune checkpoint pathways

to improve the efficacy of treatments for NMIBC. The results also contribute to a growing body

of literature on sex differences in the immune cell landscape and the importance of sex

considerations in BCG immunotherapy and the need for a sex-disaggregated data analyses

approach in biomarker studies to address the interpretations associated with inconsistent

findings across different cohorts. Given the advent of newer immunotherapies as well as

gemcitabine-docetaxel chemotherapy in the BCG naïve setting, early identification of patients

who may not benefit from BCG, using information from pre-treatment immune landscape, will

allow precise incorporation of newer therapeutics in NMIBC.

**Conflict of interest statement**

All authors have no conflicts of interest to disclose.

**Author contributions and acknowledgements**

MK conceptualized the overall study and contributed to study design, manuscript writing and

revision. DRS and BR contributed to the study design. SC performed manual validation of mIF

staining and automated scoring. BR performed all data analysis relating to the KHSC cohort and

contributed to manuscript writing and revision. PY, SR and KP contributed to manuscript

writing and revision. KS, HG and AG performed additional analysis included in this study, such

as analysis for the independent validation cohort and subtyping and contributed to manuscript

writing and reviewing. CJ and DMB provided the KHSC tissue cohort and associated clinical data.

KT guided all data analysis. DRS and CJ helped with clinical classifications of BCG treated patient

specimens. All authors reviewed the manuscript. PY is supported by Robert Wilson Fellowship

for PhD training. This study was supported by research operating grants from Bladder Cancer

Canada (MK, DRS and DMB), Cancer Research Society (MK, DRS and DMB), Ontario Institute for

Cancer Research/Ontario Molecular Pathology Research Network (DMB); Ontario Ministry of

Research Innovation and Science: Early Researcher Award and Canada Foundation for

Innovation (MK). We thank Shakeel Virk at the QLMP for his assistance with imaging and Halo

software and Katy Milne at BC Cancer Agency's Molecular and Cellular Immunology Core for

immunofluorescence staining of TMAs.

**References**

1.      Koti, M., et al., *Sex Differences in Bladder Cancer Immunobiology and Outcomes: A Collaborative*
476             *Review with Implications for Treatment.* Eur Urol Oncol, 2020.
477     2.      Uhlig, A., et al., *Gender-specific Differences in Recurrence of Non–muscle-invasive Bladder*
478             *Cancer: A Systematic Review and Meta-analysis.* European Urology Focus, 2018. **4**(6): p. 924-
479             936.
480     3.      Marks, P., et al., *Female with bladder cancer: what and why is there a difference?* Transl Androl
481             Urol, 2016. **5**(5): p. 668-682.
482     4.      Krimphove, M.J., et al., *Sex-specific Differences in the Quality of Treatment of Muscle-invasive*
483             *Bladder Cancer Do Not Explain the Overall Survival Discrepancy.* Eur Urol Focus, 2019.
484     5.      Roumiguie, M., et al., *International Bladder Cancer Group Consensus Statement on Clinical Trial*
485             *Design for Patients with Bacillus Calmette-Guerin-exposed High-risk Non-muscle-invasive*
486             *Bladder Cancer.* Eur Urol, 2021.
487     6.      Lobo, N., et al., *Updated European Association of Urology (EAU) Prognostic Factor Risk Groups*
488             *Overestimate the Risk of Progression in Patients with Non-muscle-invasive Bladder Cancer*
489             *Treated with Bacillus Calmette-Guerin.* Eur Urol Oncol, 2021.
490     7.      Matulay, J.T., et al., *Contemporary Outcomes of Patients with Nonmuscle-Invasive Bladder*
491             *Cancer Treated with bacillus Calmette-Guerin: Implications for Clinical Trial Design.* J Urol, 2021.
492             **205**(6): p. 1612-1621.
493     8.      Babjuk, M., et al., *European Association of Urology Guidelines on Non-muscle-invasive Bladder*
494             *Cancer (Ta, T1, and Carcinoma in Situ).* Eur Urol, 2022. **81**(1): p. 75-94.
495     9.      Kamat, A.M., et al., *Definitions, End Points, and Clinical Trial Designs for Bladder Cancer:*
496             *Recommendations From the Society for Immunotherapy of Cancer and the International Bladder*
497             *Cancer Group.* J Clin Oncol, 2023: p. JCO2300307.
498     10.     Goldberg, I.P., et al., *Pharmacologic Therapies for Non-Muscle Invasive Bladder Cancer: Current*
499             *and Future Treatments.* Arch Pharmacol Ther, 2022. **4**(1): p. 13-22.
500     11.     Koti, M., et al., *Adaptive Immunity in Genitourinary Cancers.* Eur Urol Oncol, 2023.
501     12.     Nazmifar, M., et al., *Clinical and Preclinical Therapies for Bladder Cancer Following Bacillus*
502             *Calmette-Guerin Failure.* J Urol, 2023. **209**(1): p. 32-48.
503     13.     Packiam, V.T., et al., *The current landscape of salvage therapies for patients with bacillus*
504             *Calmette-Guerin unresponsive nonmuscle invasive bladder cancer.* Curr Opin Urol, 2021. **31**(3):
505             p. 178-187.
506     14.     Balar, A.V., et al., *Pembrolizumab monotherapy for the treatment of high-risk non-muscle-*
507             *invasive bladder cancer unresponsive to BCG (KEYNOTE-057): an open-label, single-arm,*
508             *multicentre, phase 2 study.* Lancet Oncol, 2021. **22**(7): p. 919-930.
509     15.     Flaig, T.W., et al., *NCCN Guidelines(R) Insights: Bladder Cancer, Version 2.2022.* J Natl Compr
510             Canc Netw, 2022. **20**(8): p. 866-878.
511     16.     Jiang, Y., et al., *Non-invasive tumor microenvironment evaluation and treatment response*
512             *prediction in gastric cancer using deep learning radiomics.* Cell Rep Med, 2023. **4**(8): p. 101146.
513     17.     Goto, W., et al., *Predictive value of improvement in the immune tumour microenvironment in*
514             *patients with breast cancer treated with neoadjuvant chemotherapy.* ESMO Open, 2018. **3**(6): p.
515             e000305.
516     18.     Chenard, S., et al., *Sexual Dimorphism in Outcomes of Non-muscle-invasive Bladder Cancer: A*
517             *Role of CD163+ Macrophages, B cells, and PD-L1 Immune Checkpoint.* Eur Urol Open Sci, 2021.
518             **29**: p. 50-58.

19.  Kates, M., et al., *Adaptive Immune Resistance to Intravesical BCG in Non-Muscle Invasive Bladder*
*Cancer: Implications for Prospective BCG-Unresponsive Trials.* Clin Cancer Res, 2020. **26**(4): p.
882-891.
20.  Strandgaard, T., et al., *Elevated T-cell Exhaustion and Urinary Tumor DNA Levels Are Associated*
*with Bacillus Calmette-Guerin Failure in Patients with Non-muscle-invasive Bladder Cancer.* Eur
Urol, 2022. **82**(6): p. 646-656.
21.  de Jong, F.C., et al., *Non-muscle-invasive bladder cancer molecular subtypes predict differential*
*response to intravesical Bacillus Calmette-Guerin.* Sci Transl Med, 2023. **15**(697): p. eabn4118.
22.  Newman, T.B. and W.S. Browner, *In defense of standardized regression coefficients.*
Epidemiology, 1991. **2**(5): p. 383-6.
23.  Lindskrog, S.V., et al., *Single-nucleus and Spatially Resolved Intratumor Subtype Heterogeneity in*
*Bladder Cancer.* Eur Urol Open Sci, 2023. **51**: p. 78-88.
24.  Sun, M., et al., *Infiltration and Polarization of Tumor-associated Macrophages Predict Prognosis*
*and Therapeutic Benefit in Muscle-Invasive Bladder Cancer.* Cancer Immunol Immunother, 2022.
**71**(6): p. 1497-1506.
25.  Koti, M. and D.R. Siemens, *The Evolving Landscape of Bladder Tertiary Lymphoid Structures.* J
Urol, 2023. **210**(1): p. 3-4.
26.  Koti, M., et al., *Tertiary Lymphoid Structures Associate with Tumour Stage in Urothelial Bladder*
*Cancer.* Bladder Cancer, 2017. **3**(4): p. 259-267.
27.  Hamade, A., et al., *Sex differences in the aging murine urinary bladder and influence on the*
*tumor immune microenvironment of a carcinogen-induced model of bladder cancer.* Biol Sex
Differ, 2022. **13**(1): p. 19.
28.  Priyanka Yolmo, S.R., Stephen Chenard, Gwenaëlle Conseil, Danielle Jenkins, Kartik Sachdeva,
Isaac Emon, Jake Hamilton, Minqi Xu, Manu Rangachari, Eva Michaud, Jose Mansure, Wassim
Kassouf, David M. Berman, D. Robert Siemens, Madhuri Kot, *Atypical B cells mediate poor*
*response to Bacillus Calmette Guérin immunotherapy in non-muscle invasive bladder cancer.*
bioRxiv, 2022(522127).
29.  Zirakzadeh, A.A., et al., *Tumour-associated B cells in urothelial urinary bladder cancer.* Scand J
Immunol, 2020. **91**(2): p. e12830.
30.  Schwartz, M., Y. Zhang, and J.D. Rosenblatt, *B cell regulation of the anti-tumor response and role*
*in carcinogenesis.* J Immunother Cancer, 2016. **4**: p. 40.
31.  Fehres, C.M., et al., *APRIL Induces a Novel Subset of IgA(+) Regulatory B Cells That Suppress*
*Inflammation via Expression of IL-10 and PD-L1.* Front Immunol, 2019. **10**: p. 1368.
32.  Sheridan, B.S. and L. Lefrancois, *Regional and mucosal memory T cells.* Nat Immunol, 2011.
**12**(6): p. 485-91.
33.  Wang, Z.Q., et al., *CD103 and Intratumoral Immune Response in Breast Cancer.* Clin Cancer Res,
2016. **22**(24): p. 6290-6297.
34.  Webb, J.R., K. Milne, and B.H. Nelson, *Location, location, location: CD103 demarcates*
*intraepithelial, prognostically favorable CD8(+) tumor-infiltrating lymphocytes in ovarian cancer.*
Oncoimmunology, 2014. **3**: p. e27668.
35.  Kim, Y., Y. Shin, and G.H. Kang, *Prognostic significance of CD103+ immune cells in solid tumor: a*
*systemic review and meta-analysis.* Sci Rep, 2019. **9**(1): p. 3808.
36.  Wahlin, S., et al., *Clinical impact of T cells, B cells and the PD-1/PD-L1 pathway in muscle invasive*
*bladder cancer: a comparative study of transurethral resection and cystectomy specimens.*
Oncoimmunology, 2019. **8**(11): p. e1644108.
37.  Fu, H., et al., *Identification and Validation of Stromal Immunotype Predict Survival and Benefit*
*from Adjuvant Chemotherapy in Patients with Muscle-Invasive Bladder Cancer.* Clin Cancer Res,
2018. **24**(13): p. 3069-3078.

38.    Vidotto, T., et al., *DNA damage repair gene mutations and their association with tumor immune*
*regulatory gene expression in muscle invasive bladder cancer subtypes.* J Immunother Cancer,
2019. **7**(1): p. 148.
39.    Miao, Y., et al., *Adaptive Immune Resistance Emerges from Tumor-Initiating Stem Cells.* Cell,
2019. **177**(5): p. 1172-1186 e14.
40.    Pietzak, E.J., et al., *Inherited Germline Cancer Susceptibility Gene Variants in Individuals with*
*Non-Muscle-Invasive Bladder Cancer.* Clin Cancer Res, 2022. **28**(19): p. 4267-4277.
41.    Lindskrog, S.V., et al., *An integrated multi-omics analysis identifies prognostic molecular*
*subtypes of non-muscle-invasive bladder cancer.* Nat Commun, 2021. **12**(1): p. 2301.

*Table 1: Categories of BCG therapy response.*

| BCG-refractory tumour (as per standard definitions)[8] |
|---|
| 1. If T1 G3/HG tumour is present at 3 mo. |
| 2. If TaG3/HG tumour is present after 3 months and/or at 6 mo., after either re-induction or first course of maintenance. |
| 3. If CIS (without concomitant papillary tumour) is present at 3 mo. and persists at 6 mo. after either reinduction or a first course of maintenance. For patients with CIS present at 3 mo., an additional BCG course can achieve a complete response in >50% of cases. |
| 4. If HG tumour appears during BCG maintenance therapy. |
| **BCG responsive** |
| 1. No HG tumour recurrence within 2 years following initial adequate BCG induction. |

581   *Table 2: Patient cohort clinical characteristics.*

| | BCG Responsive (N=124) | BCG Refractory (N=37) | BCG relapsing (N=12) | Overall (N=173) |
|---|---|---|---|---|
| **Age** | | | | |
| Mean | 71.5 | 68.3 | 72.3 | 70.8 |
| Median | 71.5 | 69.0 | 73.0 | 71.0 |
| Range | 35-93 | 45-85 | 62-82 | 35-93 |
| **Sex** | | | | |
| Male | 102 (82.3%) | 27 (73.0%) | 10 (83.3%) | 139 (80.3%) |
| Female | 22 (17.7%) | 10 (27.0%) | 2 (16.7%) | 34 (19.7%) |
| **AUA Risk Score** | | | | |
| High | 74 (59.7%) | 36 (97.3%) | 12 (100.0%) | 122 (70.5%) |
| Intermediate | 41 (33.1%) | 1 (2.7%) | 0 (0%) | 42 (24.3%) |
| Low | 9 (7.3%) | 0 (0%) | 0 (0%) | 9 (5.2%) |
| **Diagnosis TURBT Stage and Grade** | | | | |
| CIS | 11 (8.9%) | 1 (2.7%) | 1 (8.3%) | 13 (7.5%) |
| PUNLMP | 1 (0.8%) | 0 (0%) | 0 (0%) | 1 (0.6%) |
| T1HG | 32 (25.8%) | 15 (40.5%) | 4 (33.3%) | 51 (29.5%) |
| TaHG | 51 (41.1%) | 17 (45.9%) | 5 (41.7%) | 73 (42.2%) |
| TaLG | 29 (23.4%) | 2 (5.4%) | 2 (16.7%) | 33 (19.1%) |
| Unknown | 0 (0%) | 2 (5.4%) | 0 (0%) | 2 (1.2%) |
| **Time to Follow-up** | | | | |
| Median time to follow-up (months) | 53.4 | 57.9 | 65.4 | 54.3 |
| **BCG Adequacy** | | | | |
| Adequate BCG | 47 (37.9%) | 23 (62.2%) | 11 (91.7%) | 81 (46.8%) |
| Induction only | 77 (62.1%) | 14 (37.8%) | 1 (8.3%) | 92 (53.2%) |
| **Subtype** | | | | |
| Genomically Unstable | 13 (10.3%) | 3 (7.7%) | 1 (8.3%) | 17 (9.6%) |
| URO-KRT5+ | 11 (8.7%) | 2 (5.1%) | 2 (16.7%) | 15 (8.5%) |
| Urothelial-like | 9 (7.1%) | 7 (17.9%) | 1 (8.3%) | 17 (9.6%) |
| Basal | 1 (0.8%) | 1 (2.6%) | 0 (0.0%) | 2 (1.1%) |
| Not Subtyped | 92 (73.0%) | 26 (66.7%) | 8 (66.7%) | 126 |
| **Subtype Grade** | | | | |
| 1 | 27 (21.4%) | 7 (17.9%) | 2 (16.7%) | 36 (20.3%) |
| 2 | 7 (5.6%) | 6 (15.4%) | 2 (16.7%) | 15 (8.5%) |
| Not Subtyped | 92 (73.0%) | 26 (66.7%) | 8 (66.7%) | 126 |

**Figure 1. High intra-tumoral CD79a+ B cell density associates with early recurrence and progression post BCG immunotherapy. A)** Patients with high stromal CD79a+ B cell density have significantly shorter RFS compared to patients with low density (Log-rank p=5.83e-6, IRR=0.29, HR=3.18, p=1.75e-5). **B)** Patients with high stromal CD79a+ B cell density have significantly shorter PFS compared to patients with low density (Log-rank p=2.07e-2, IRR=0.32, HR=3.00, p=2.8e-2), **C)** BCG refractory patients have significantly higher stromal CD79a+ cell density compared to BCG responders (W$_{Mann-Whitney}$=1744.5, p=0.027), and **D)** BCG refractory patients have significantly higher stromal CD163 cell density compared to BCG responders (W$_{Mann-Whitney}$=1688.00, p=0.015).

**Figure 2. Stromal CD79a+ B cells are strongly correlated with CD163+ M2-like macrophages, CD8+ cytotoxic T cells, CD103+ cells and PD-1 immune checkpoint expression.** Spearman correlation plot for immune cell populations in the stromal (A and B) and epithelial (C and D) compartments of tumors from BCG responders and refractory. Correlation plots showing the correlations between immune cells in tumor from patients categorized as BCG refractory and responders.

**Figure 3. High tumor infiltrating M2-like macrophage, CD8+ and CD103+ T cell density associates with early recurrence post BCG immunotherapy. A)** Patients with high stromal CD163+ macrophage density have significantly worse RFS compared to patients with low density. Log-rank p=1.98e-6, IRR=0.31, HR=2.87, p=3.73e-4. **B)** Patients with high stromal CD3+CD8+ cell density have significantly worse RFS compared to patients with low density. Log-rank p=2.59e-4, IRR=0.34, HR=2.69, p=4.45e-2. **C)** Patients with high stromal CD103+ cell density have significantly worse RFS compared to patients with low density. Log-rank p=1.51e-3, IRR=0.40, HR=2.35, p=2.07e-3

**Figure 4. Forest plots demonstrating patient demographics and immune markers with clinical outcomes post BCG immunotherapy.** Forest plot based showing univariate Cox Proportional Hazards (CPH) regression analyses, with recurrence highlighted in black and progression in blue. Dots represent hazard ratios and lines represent the 95% confidence intervals. Hazard ratios for immune cell types are displayed graphically in terms of risk per standard deviation (σ) increment in cell infiltration. Multivariable CPH regression analyses were performed for each immune phenotype separately, including sex, age, AUA risk score, and amount of BCG treatment received after the analyzed tumor. P values were calculated using the Wald test, and multivariable p values were adjusted using the Bonferroni correction.