# Modulation by NPYR underlies experience-dependent, sexually dimorphic learning

Sonu Peedikayil-Kurien[1,2], Rizwanul Haque[1,2], Asaf Gat[1,2] and Meital Oren-Suissa[1,2*]

[1] Department of Brain Sciences, Weizmann Institute of Science; Rehovot, 7610001, Israel.

[2] Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot 7610001,

Israel

*Corresponding author. Email: meital.oren@weizmann.ac.il

**Abstract:** The evolutionary paths taken by each sex within a given species sometimes diverge, resulting in behavioral differences. Given their distinct needs, the mechanism by which each sex learns from a shared experience is still an open question. Here, we reveal sexual dimorphism in learning: *C. elegans* males do not learn to avoid the pathogenic bacteria PA14 as efficiently and rapidly as hermaphrodites. Notably, neuronal activity following pathogen exposure was dimorphic: hermaphrodites generate robust representations, while males, in line with their behavior, exhibit contrasting representations. Transcriptomic and behavioral analysis revealed that the neuropeptide receptor *npr-5*, an ortholog of the mammalian NPY receptor, regulates male learning by modulating neuronal activity. Furthermore, we show the dependency of the males' decision-making on their sexual status and demonstrate the pivotal role of *npr-5* as a modulator of incoming sensory cues. Taken together, we portray sex-specific plasticity in behavior toward a shared experience by modulating learning.

## Introduction

Learning facilitates the organism's ability to cope with a changing environment, making it an inherent property of biological neural systems[1]. To adapt to the environment, the organism needs to retain the learning experience and then use the stored information to alter future responses[2]. The synaptic basis of learning and memory has been well established and provides a solid foundation to assess the relationship between experience-dependent learning and survival[3].

Since sexual selection predicts that males and females are subjected to different selection pressures leading to dimorphisms[4,5], it stands to reason that the same would occur for an essential ability such as learning. Consistent with this prediction, several sex-shared traits such as perception and learning have been previously demonstrated to be modulated by the genetic sex[6–12]. It has also been proposed that learning can be a driving force for evolutionary change[13]. Thus, sex-specific evolutionary drives might be in tension with decision-making processes when facing an environmental cue[14,15]. While a significant portion of sex differences in behavior could be attributed to the neurochemical pathway of circulating sex hormones, it has been suggested that behavior is also influenced by other factors, such as life history and sex-specific gene expression[16]. This indicates a bi-directional relationship between sexual identity and the environment, with far-reaching consequences on cognitive abilities, including learning and memory. How sexual identity drives molecular and circuit changes toward a shared experience that intersects with learning and survival is still an open question.

The nematode *C. elegans* has been a useful model for the study of learning and memory at the molecular, cellular, neural circuitry, and behavioral levels[17]. Numerous studies have demonstrated that *C. elegans* is capable of non-associative learning and short-term memory, in addition to associative learning, long-term memory, and sensory integration of conflicting cues[18–22].

The two sexes of *C. elegans* display sexually dimorphic behaviors, such as mechanosensory responses to tail touch, dosage-dependent differences in nociceptive behaviors, and chemoattraction to odors and chemicals. Some of these differences have been mapped to corresponding network changes [8,23,24].

Here, we investigated if and how the genetic sex affects context- and experience-dependent behavioral plasticity when learning a shared environmental cue. We harnessed the well-established model of short-term pathogenic aversive conditioning in *C. elegans* [21,25,26], where hermaphrodites shift their preference from naïve attraction to aversion upon exposure to the pathogenic bacterium

*Pseudomonas aeruginosa* (PA14)[25,27]. We show that males are unable to avoid PA14 as efficiently

as hermaphrodites after training. This dimorphic avoidance, however, is not due to differences in

the immune responses, but due to layered differences in the sensory and subsequent downstream

processing. We observed context-dependent learning when exposing the males to hermaphrodites,

which altered their behavior towards PA14. Transcriptomic analyses indicate neuromodulatory

regulation, with one candidate, *npr-5*, regulating both the learned as well as context-dependent

avoidance by modulating sensory responses. We also unravel the importance of tissue-specific

sexual identity and decentralized molecular signaling in regulating male responses. Taken

together, our work underscores the diverse levels at which sexual identity modulates dimorphic

behavioral responses in each sex towards a shared experience.

**Aversive olfactory learning is sexually dimorphic**

To test whether short-term aversive learning is sexually dimorphic, we separated virgin males and

hermaphrodites and subjected them to an acute training paradigm of pathogenic bacterial exposure

(PA14 or *Escherichia coli* strain OP50 as control) that lasted 6 hours, followed by a choice-based

test (Fig. 1a). We found that while hermaphrodites efficiently avoided PA14 at the end of the 6-

hour mark, males did not display learned aversion (Fig. 1b-c). A similar dimorphic response was

observed when we tested the learning abilities of the sexes against another pathogenic bacterium,

*Staphylococcus aureus*[28,29] (Supplementary Fig. 1a-b). Analogously, when we tested the naïve

avoidance by exposing either sex to a lawn of PA14 and scored lawn occupancy after 6-8 hours,

males remained within the PA14 lawn while the hermaphrodites' occupancy was minimal

(Supplementary Fig. 1c). We speculated that these observed dimorphisms in short-term learning

could arise from differences at checkpoints of information processing within the sexes, such as i)

immune responses, ii) integration of sensory information, or iii) a deliberately chosen dimorphic

outcome to favor a sex-specific need, in a temporal manner.

To determine whether the immune system is activated in both sexes, we examined candidate genes

from multiple pathways activated during bacterial infection using either real-time qPCR analysis

or reporter expression levels[30]. The CUB-domain containing gene F08G5.6[31] and ShK-like toxin

gene F49F1.6, mainly activated by the PMK-1 p38 MAPK pathway[32] were elevated in both sexes

after exposure to PA14 (Supplementary Fig. 1d-e). The *pmk-1* pathway-independent immune

responder *fshr-1*[33,34] and its effector, F01D5.5 [35], another ShK-like toxin gene, were also elevated

in both sexes after training (Supplementary Fig. 1f-g). The *pmk-1* independent gene *irg-3* [30,35] was

also elevated in both sexes, but to a lesser extent in males (Supplementary Fig. 1h). In addition,

we imaged the activation of the infection responder gene *irg-1*, which has been shown to be

dependent on *zip-2*, a transcription factor specifically activated upon PA14 exposure and is

independent of *pmk-1* [30]. We found the *irg-1* levels in the anterior gut region of both sexes to be

similar (Fig. 1d-e). The similar pathogenic sensitivity was reflected in killing assays conducted to

measure PA14 virulence, where both males and hermaphrodites died with similar kinetics (Fig. 1f,

Supplementary Fig. 1i). This is in line with the observation that males do indeed avoid PA14, but

only after much more extended training periods (Supplementary Fig. 1j-k). Thus, even though the

immune response of the males was activated, they were unable to integrate the pathogenicity of

PA14 towards learned aversion as rapidly as hermaphrodites.

To evaluate if the avoidance of PA14 is either developmentally timed or inherently dimorphic, we

subjected juvenile animals to the same training paradigm as the adult animals (See Methods). We

observed that juveniles of both sexes failed to learn and actively preferred PA14 even after training

(Supplementary Fig. 1l-n). This suggests that the dimorphic behavior observed in adulthood results

from a developmentally mature nervous system[24].

To characterize the impact of sexual identity on the learning axis, we feminized or masculinized

specific tissues in hermaphrodites and males, respectively, and subjected them to the learning

paradigm (Fig. 1g). When only the nervous system is sex-reversed, hermaphrodites exhibited a

dramatic reduction in their learning ability (Fig. 1h-i). Contrary to our expectations, pan-

neuronally feminized males failed to learn (Fig. 1j-k), even though the sex reversal shifted the

olfactory preferences of feminized males for certain odors[24] (Supplementary Fig. 1o), suggesting

that the substrate of learned avoidance is restricted mostly to the nervous system in

hermaphrodites, but involves additional systems in males. In the absence of a direct link between

neuronal regulation and learning in males, we hypothesized that non-neuronal tissues such as the

gut and the germline might affect male learning. Indeed, previous studies have shown the

importance of the gut in regulating behavior[36,37], yet feminizing the gut of males was also

insufficient to enhance learning in males (Fig. 1j-k). Similar results were obtained when analyzing

males with a mutated germline, although germline-mutated hermaphrodites seemed to display

dampened avoidance (Supplementary Fig. 1p-q). However, when both the nervous system and the

gut were feminized, these males learned to avoid PA14 significantly better than wild-type males

(Fig. 1j-k). Thus, the default male drive is orchestrated by several tissues, and their
intercommunication is critical for the evaluation and subsequent response to external stimuli.

Figure 1. **Distinct tissues govern sexually dimorphic short-term learning of PA14 avoidance**

**a** Schematic representation of the training paradigm. A group of *C. elegans* was separated into males or hermaphrodites after late L4, exposed to the pathogenic bacterium *Pseudomonas aeruginosa* (PA14, cyan, "trained") or OP50 (control, magenta, "naïve") for 6 hours, and then assayed for their choice index against OP50 and PA14 (See Methods). In all figures, n represents the number of animals tested, and N represents the number of biological repeats. **b** The Choice Index of hermaphrodites (OP50, PA14: N = 9) and males (OP50, PA14: N = 7) subjected to the training paradigm in A shows dimorphic responses after training. **c** The learning indices of hermaphrodites (N = 9) and males (N = 7) are dimorphic. **d** Representative confocal micrographs of *irg-1*::GFP hermaphrodites and males trained on OP50 or PA14 after 6 hours of training. Scale bar = 50μm. **e** Quantification of the anterior region of the gut for *irg-1*::GFP in hermaphrodites (OP50: n = 13; PA14: n = 11; n = individual animals) and males (OP50: n = 13; PA14: n = 10; n = individual animals) after exposure to OP50 or PA14. **f** The percentage of live hermaphrodites (N = 6) and males (N = 6) in slow-killing assay. **g** Schematic representation of the tissue sex-reversal experiments (the results of the equivalent germline experiments are presented in Supplementary Fig. 1). **h-i** Choice **(h)** and Learning **(i)** Indices of wild-type (OP50, PA14: N = 6) and pan-neuronally masculinized (p*rab-3*::*fem-3*) hermaphrodites (OP50, PA14: N = 6). **j-k** Choice **(j)** and Learning **(k)** indices of wild-type (control) males (OP50, PA14: N = 5), pan-neuronally feminized (p*rab-3*::*tra-2(ic)*) males (OP50, PA14: N = 5), gut-feminized (p*elt-2*::*tra-2(ic)*) males (OP50, PA14: N = 8), and pan-neuronally and gut feminized (p*rab-3*::*tra-2(ic)*);p*elt-2*::*tra-2*) males (OP50, PA14, N = 5). Statistical test used are: in (**b**, **e**, **f**, **h**, **j**), a two-way ANOVA with Šídák's multiple comparisons test, in (**c**, **i**), a Mann Whitney test. In (**k**), a Brown-Forsythe and Welch's ANOVA with Dunnett's T3 multiple comparisons test. Box and Whisker plots with mean and entire data range. P values are indicated in the figure.

## Sexually dimorphic neuronal representations of PA14 exposure

We previously demonstrated that while nociceptive sensory neurons in *C. elegans* respond similarly in the two sexes to a noxious cue (glycerol), the network topology diverges, shaping sexually dimorphic behaviors[8]. The neurons regulating PA14 avoidance have been previously mapped[38,39]. The sensory neuron AWC is required for odor-mediated attraction, while AWB is required for odor-mediated repulsion[40–42]. In addition, these neurons have been recently

established as important in recognizing the odor bouquet presented by PA14 and for regulating

both adult and larval learning[35,38,39]. The dimorphic behavior we observed could then arise from

distinct computing of environmental cues (i.e., dimorphic neural states lead to dimorphic

behavior). To evaluate this, we used calcium imaging to measure the neuronal activity of sensory-

and inter-neurons of naïve and trained hermaphrodites and males and quantified the resulting

amplitudes during exposure to a sequence of OP50-PA14-OP50 supernatants (see methods). First,

we found higher calcium activity in naïve males, upon presentation of PA14, in the sensory neuron

AWC (OFF response) (Fig. 2a-b, i, Supplementary Fig. 2a, c, q) and AWB (ON and OFF response)

(Fig. 2c-d, j, Supplementary Fig. 2e, g, r-t), compared to naïve hermaphrodites. Next, we examined

the change in response after PA14 training. In both sexes, AWC exhibited no change in OFF

response after training, compared to naïve animals (Fig. 2a-b, i, Supplementary Fig. 2a-d, q).

However, the repellent odor-sensitive AWB displayed distinct dynamics. While there was no

change in the ON response of either sex after training (Fig. 2c-d, Supplementary Fig. 2r-s), the

OFF response was higher in hermaphrodites (Fig. 2c-d, j, Supplementary Fig. 2t). Taken together,

these results suggest that naïve responses in males are stronger in both AWB and AWC, while

hermaphrodites showed stronger trained response in AWB (Fig. 2m).

The sensory information from AWC and AWB is passed on to a downstream interneuron AIY,

which has been implicated as an information checkpoint that regulates memory retrieval [39,43–45] as

well as hermaphrodite training and forgetting states of PA14 aversive learning [26,38]. We observed

a reduction in calcium activity after training at the soma of the AIY neuron in both sexes (Fig. 2e-

f, k, Supplementary Fig. 2i-l, u). Yet, within the axonal regions of AIY, we observed a decrease in

calcium activity after training only in hermaphrodites but not in males (Fig. 2g-h, l, Supplementary

Fig. 2x). Interestingly, the latency of calcium responses in AIY soma (Supplementary Fig. 2v) and

axonal regions (Supplementary Fig. 2y) of trained males was smaller when compared to trained

hermaphrodites, while the time the neuron was active was unchanged in all the groups examined

(Supplementary Fig. 2w, z). Taken together, our results show an increased sensory response to

PA14 in AWB, and a subsequent decrease in the interneuron AIY only in hermaphrodites.

However, the neuronal response landscape was dimorphic with naïve male sensory responses

higher and the interneuron AIY displaying compartmentalized (soma vs axonal) dimorphic

responses (Fig. 2m).

To determine whether the observed dimorphisms in neuronal activity also extend to the synaptic vesicle organization of AIY, i.e., a readout of the learning process, we used a fluorescent readout of RAB-3, which regulates docking of synaptic vesicles[46–48]. The axonal regions of AIY can be distinguished according to their respective synaptic partners as Zone 2 and 3 (Fig. 2n) [48]. We quantified the mCherry expression levels of RAB-3 after training with PA14 and observed a non-dimorphic decrease in RAB-3 levels in both Zones 2 and 3 (Fig. 2o-p), suggesting that the sensory stimulus is processed in both sexes. However, we noticed that the extent of difference in RAB-3 expression between the naïve and trained males is lower than that of hermaphrodites (Fig. 2q), suggesting that dimorphic perceptive information is being communicated downstream.

Figure 2. **Distinct dimorphic compartmentalized information processing in response to PA14 stimulus. a-h** From top to bottom, SEM traces of naïve (magenta) and trained (cyan) animals of

both sexes in select neurons. The shaded region in the SEM traces corresponds to the exposure

window to PA14 supernatant in both naïve and trained animals. OP50 supernatant was used as the

buffer (See Methods). **a** AWC calcium levels in trained (n = 12) and naïve (n = 12) hermaphrodites.

**b** AWC calcium levels in trained (n = 13) and naïve (n = 11) males. **c** AWB calcium levels in

trained (n = 36) and naïve (n = 34) hermaphrodites. **d** AWB calcium levels in trained (n = 35) and

naïve (n = 35) males. **e** AIY soma calcium levels in trained (n = 36) and naïve (n = 34)

hermaphrodites. **f** AIY soma calcium levels in trained (n = 35) and naïve (n = 35) males. **g** AIY

axonal calcium levels in trained (n = 35) and naïve (n = 35) hermaphrodites. **h** AIY axonal calcium

levels in trained (n = 27) and naïve (n = 29) males. **i** Quantification of OFF response of AWC. **j**

Quantification of OFF response of AWB. **k** Quantification of ON response of AIY. **l**

Quantification of ON response of AIY Axon. **m** Table summarizing the neuronal responses

observed. Purple, significant response in hermaphrodites; Blue, Significant response in males;

Grey, no difference between the sexes. The sensory neurons AWC and AWB showed higher

calcium responses in the naïve males compared to naïve hermaphrodites. After training,

hermaphrodites exhibit dimorphic OFF responses in AWB, and ON response in AIY axon. **n**

Representative confocal micrographs of p*ttx-3*::*rab-3*::mCherry in naïve and trained

hermaphrodites and males showing expression in distinct zones of AIY, namely, a synapse-devoid

Zone 1 and synapse-rich Zones 2 and 3. Scale bar = 12μm. **o-p** Synaptic RAB-3 mCherry

fluorescence in AIY Zone 2 and Zone 3 axonal regions, respectively. Each dot represents a puncta

from 26-47 animals per group. **q** Normalized delta between naïve and trained RAB-3 expression

of both sexes in Zone 2 and Zone 3. Hermaphrodites in magenta, and Males in blue. Data is

normalized to hermaphrodites. In (**a-h**) change in fluorescence ($\Delta$F) was normalized to the average

GCaMP signal in 10 seconds before stimulus start ($F_0$). ON response was defined as the difference

in average $\Delta$F/ $F_0$ during 10 seconds after stimulus start to 5 seconds before stimulus start. OFF

response was calculated as the difference in average $\Delta$F/ $F_0$ during 10 seconds after stimulus

removal to 5 seconds before stimulus start. In (**i-l**, **o-p**), the statistical test used is Kruskal-Wallis

test with Dunn's multiple comparisons test was performed. Box and Whisker plots with mean and

entire data range. P values are indicated in the figure.

**Dimorphic molecular landscape of naïve and trained animals**

To characterize the molecular landscape of short-term dimorphic learning, we performed whole-
animal transcriptomics on sex-separated naïve and trained animals of both sexes (4 groups with 4
biological repeats), and the resultant RNA was sequenced using a bulk optimized protocol of the
single-cell MAR-seq protocol[49]. This protocol has been previously used to establish the molecular
profile of both sexes in all developmental stages[50]. To visualize the relationship between training
and sexual identity across all experimental groups, we performed a principal component analysis
(PCA) on the ~ 12,000 identified genes. The PCA revealed a clear separation by sex (Fig. 3a),
validated by the dimorphic expression of *lsy-27, tbx-9* and *tra-2* (higher in hermaphrodites[50,51])
and *mab-3, pkd-2* and *ins-39* (higher in males[50,52]) (Supplementary Fig. 3a). The training status of
the animals was also separated in the PCA (Fig. 3b). We found 869 and 846 differentially
expressed genes in hermaphrodites and males, respectively, after training (p value < 0.05;
Supplementary Data 1). Unlike the behavioral results, where both naïve and trained males
displayed similar attraction towards PA14 (Fig. 1b-c), their molecular landscape differed and
reflected the training experience (Fig. 3b). Volcano plots and functional GO term analysis revealed
similar changes in immune regulation, fatty acid metabolism, and other stress-related pathways in
both sexes, an expected outcome given the gustatory and pathogenic nature of the training
bacterium (Fig. 3c-d, Supplementary Fig. 3c, Supplementary Data 1, Supplementary Data 2). Of
the entire transcriptomics dataset, 70 genes were previously reported to be involved in PA14-
related behaviors (Supplementary Data 2)[26,27,37,39,53–64]. Consistent with our survival assays, most
of the genes implicated in PA14 survival, such as *daf-16* and *daf-7*, were non-dimorphic. Genes
involved in training and forgetting, such as *gst-38* and *gst-26* [26], were similarly regulated in trained
animals of both sexes (Supplementary Fig. 3d, Supplementary Data 2). qPCR analysis of metabolic
gene *acdh-1* [65], implicated in PA14 survival, was reduced non-dimorphically after PA14 exposure
as indicated by our transcriptomics dataset (Supplementary Fig. 3b, d).
To identify transient changes that regulate male learning, we focused on differentially expressed
neuro-specific gene classes, such as receptors and neuropeptide families (Fig. 3e, Supplementary
Data 2). We uncovered changes in transcription levels for several transcription factors. The
homeodomain *ceh-37*, which specifies the identity of the sensory neuron AWB[66]  and is involved
in regulating immunity from the gut[67] was upregulated in both sexes (Fig. 3e). *mxl-3*, involved in
lipolysis and expressed in sensory neurons including ASE, AWC and AWB[68,69] was
downregulated in males, while the H3K4 demethylase *rbr-2*, involved in axon guidance and

transmission of stress resistance across generations[70,71] was dimorphically regulated (Fig. 3e).

Interestingly, the levels of neuropeptide *ins-11*[61], a negative regulator of PA14 avoidance, were

increased in trained animals of both sexes (Fig. 3e), as confirmed by qPCR analysis

(Supplementary Fig. 3b, d). *unc-13*, a gene involved in synaptic vesicle clustering, was observed

to be dimorphically regulated after training in the sexes, along with the ligand-gated channel *lgc-*

*21* and the neuropeptide receptor *npr-5*. qPCR analysis confirmed the dimorphic expression of

*npr-5,* which is down-regulated in males after training (Fig. 3e, Supplementary Fig. 3b), but not

of *unc-13* (Supplementary Fig. 3b). Overall, our transcriptomic data suggest that while both sexes

respond to PA14 exposure with a similar molecular signature, additional modulatory mechanisms

may have evolved to regulate the observed male behavior.

**Sexually dimorphic modulation of the NPY receptor expression**

Recent evidence suggests the importance of integration between the nervous and the immune

systems in generating appropriate responses towards pathogenic bacterium[72]. Of note,

neuropeptides and their receptors play a major role in PA14 avoidance behavior[73]. One such

example is the G-protein coupled receptor *npr-1,* which has been shown to inhibit AQR, PQR, and

URX neurons to confer resistance to PA14[74,75]. Given the importance of the neuropeptidergic

system in modulating diverse behaviors, including learning[76], the neuropeptide receptor *npr-5*

emerged as a likely candidate. Notably, NPR-5 is an ortholog of the mammalian NPY receptor, a

key regulator of various behaviors, including learning and memory[77–81]. In *C. elegans*, *npr-5* is

involved in regulating foraging and fatty acid metabolism in hermaphrodites[82,83]. In addition, *npr-*

*5* was shown to modulate feeding by regulating the secretion of serotonin from ADF neurons in

an NMDAR-dependent manner[84], both of which have independently been implicated in PA14

aversive learning [27,85]. Thus, we sought to examine its role in male learning behavior.

To determine the change in expression post-training, we endogenously tagged *npr-5* using

CRISPR-Cas9 gene-editing and quantified its expression levels in head sensory neurons,

previously described to regulate behavior[82,83] (Fig. 3f-g). Consistent with previous studies[82], *npr-*

*5* was observed in body wall muscles and multiple neurons, including sensory neurons (Fig. 3f).

After training, *npr-5* levels rose in hermaphrodite AWB neurons but were reduced in male AWB

neurons (Fig. 3g). Intriguingly, this change in *npr-5* expression resembles the changes in activity

levels of AWB (Fig. 2c-d). Although ASJ has been previously described to be involved in

processing PA14 experience[37,64,86], and AIY displayed dimorphic response (Fig 2g, h, l), no

significant changes were observed in *npr-5* expression after training in ASJ and AIY (Fig. 3g).

These results support the transcriptomic data, suggesting *npr-5* is down-regulated in males upon

PA14 exposure (Fig. 3e), pointing to a critical role of *npr-5* in modulating learning, perhaps at the

sensory level.

npr-5::SL2::GFP::H2B

Figure 3. **Sexually dimorphic learning triggers sex-specific changes in NPYR expression. a-b**

Principle Component Analysis (PCA) of the transcriptomes of hermaphrodites and males (n = 4

independent samples from 600-800 animals/replicate) after training, ordered according to sex (**a**)

and training status (**b**), showing the validity of the data in separating the sexes and training states

by transcriptomics. **c-d** Volcano plots of naïve and trained hermaphrodites (**c**) and males (**d**),

depicting similar changes in transcripts **e** Heatmap of $\log_2$ fold-change of neuron-relevant genes

from naïve vs. trained males (FDR < 0.05) and hermaphrodites. Red indicates enrichment in naïve

animals (OP50), while blue indicates enrichment in trained animals (PA14). See also Supplemental

tables 1-2. **f** Representative confocal micrographs of *npr-5::SL2::GFP::H2B* CRISPR knock-in in

naïve and trained hermaphrodites and males showing expression in body wall muscles and sensory

cells. The inset shows the heatmap expression of *npr-5::SL2::GFP::H2B* in the selected region

with AWB and ASJ. Scale bar = 10µm. **g** Quantification of mean fluorescent signals of *npr*-5 in

ASJ, AIY and AWB, showing dimorphic expression changes in AWB after training (ASJ

(hermaphrodites: OP50, PA14 n =15; males: OP50, PA14 n =16); AIY (hermaphrodites: OP50 n

=13, PA14 n =15; males: OP50 n =15, PA14 n =13); AWB (hermaphrodites: OP50, PA14 n =15;

males: OP50 n =16, PA14 n =17). In (**g**), statistical test used is two-way ANOVA with Šídák's

multiple comparisons test. Box and Whisker plots with mean and entire data range. P values are

indicated in the figure.

**308    Neuropeptidergic modulation of sex-specific learning**

To evaluate the extent of neuropeptidergic regulation in male-specific PA14 aversive learning, we

subjected *npr-5* mutants to the training paradigm. Strikingly, males with a partial or full *npr-5*

deletion acquired the ability to learn quickly in a similar fashion as wild-type hermaphrodites (Fig.

4a-d). Although mutant *npr-5* hermaphrodites still avoided PA14, no significant changes were

observed between the naïve and trained responses (Fig. 4a, c). Previous studies[82], along with our

calcium imaging (Fig. 2c-d, j) and *npr-5* expression results (Fig. 3f-g) suggest a sensory-specific

site-of-action of *npr-5*. Therefore, we rescued *npr-5* expression specifically in AWB (using the

*str-1* promoter) and in ciliated neurons (using *tax-4* promoter) in an *npr-5* mutant background.

While both manipulations sufficed to impede the decision-making ability as indicated by the

choice index (Fig. 4a), reverting the animal back to the wild-type male phenotype, the *tax-4* rescue

strain displayed a stronger phenotype in the learning index (Fig. 4b). Thus, *npr-5* is required for

suppressing learned avoidance, and its activity, possibly in AWB and other sensory neurons,

suggests a sensory gating mechanism in males, in line with the observations made in the neuronal

activity of trained males (Fig. 2).

Previous studies have also established the link between various neuropeptides and the regulation
of behavioral states[86,87]. Given the dimorphic neuropeptidergic landscape and the decentralized
learning processing we observed in males (Fig. 1j-k), we postulated that the learning behavior in
males might be regulated by multiple signaling molecules in a molecular network (Supplementary
Fig. 4a). Indeed, when we screened select candidate neuropeptides using mutants or RNAi, we
observed that several neuropeptides such as *flp-33*, *flp-3* and *flp-18*, a cognate ligand of *npr-5,*
were involved (Fig. 4e, Supplementary Fig. 4b-c). The regulation of male PA14 aversive learning
by *npr-5* and multiple neuropeptides demonstrates that male learning is an amalgamation of
complex internal processing.

Figure 4. **NPR-5 regulates male-specific behavior. a-b** Choice (**a**) and Learning (**b**) Indices of
naïve (magenta) and trained (cyan) wild-type and *npr-5* mutant hermaphrodites and males and
male rescue strains where *npr-5* cDNA was expressed either under *str-1* promoter (AWB specific
expression [88]) or *tax-4* promoter (expression in ciliated neurons [89]); data shows sex-specific
behavioral effects in males (Choice Index: wild-type hermaphrodites: OP50 (N = 12), PA14 (N =
13); *npr-5* hermaphrodites: OP50 (N =10), PA14 (N = 12); wild-type males: OP50, PA14 (N =
10); *npr-5* males: OP50, PA14 (N = 9); p*str-1*::*npr-5*cDNA; *npr-5* males: OP50, PA14 (N = 5);
p*tax-4*::*npr-5*cDNA;*npr-5*  males:  OP50,  PA14  (N  =  4)).  Learning  Index:  wild-type

hermaphrodites: (N = 12); *npr-5* hermaphrodites: (N = 11); wild-type males: (N = 10); *npr-5* males:

(N = 9); p*str-1*::*npr-5*cDNA;*npr-5* males: (N = 5); p*tax-4*::*npr-5*cDNA;*npr-5* males: (N = 4)). **c-d**

Choice (**c**) and Learning (**d**) indices of naive (magenta) and trained (cyan) wild-type and *npr-5*

CRISPR mutant hermaphrodites and males, showing sex-specific behavioral effects in males

(hermaphrodites: OP50, PA14, Learning Index (n = 9); *npr-5* hermaphrodites: OP50, PA14,

Learning Index (N = 6); males: OP50, PA14, (N = 8), Learning index (N = 7); *npr-5* males: OP50,

PA14, Learning Index (N = 7)). **e** Learning indices of wild-type (N = 17), *flp-32* (*gk3692*) mutant,

*flp-33* (*gk1058*) mutant, *flp-18* (*db99*) mutant and *flp-18 (gk3063)* males (N = 6), showing

peptidergic regulation of male learning. Statistical test used are: in (**a, c**), a two-way ANOVA with

Tukey's multiple comparisons test, in (**b, d, e**), a Kruskal-Wallis test with Dunn's multiple

comparisons test. Box and Whisker plots with mean and entire data range. P values are indicated

in the figure.

***npr-5* functions in AWB to modulate PA14 responses**

Based on its sensory responses to PA14 (Fig. 2c-d, j), the observed decrease in *npr-5* expression

following training (Fig. 3g), and the change in behavior upon rescue (Fig. 4a), AWB is a likely

target of *npr-5* action. We, therefore, measured AWB activity in *npr-5* mutants and observed a

striking change in calcium response patterns, namely, a heightened OFF response (Fig. 5a-d,

Supplementary Fig. 5a-b), compared to wild-type animals (Fig. 2c-d). In addition, only *npr-5*

mutant male AWB neurons exhibited a stronger response to the removal of the PA14 stimulus

after training (Fig. 5c-d), similar to wild-type hermaphrodites (Fig. 2c, j). The *npr-5*

hermaphrodites showed a decrease in both ON and OFF responses after training (Fig. 5a, c,

Supplementary Fig. 5a, c-d). Intriguingly, when stimulated with diluted 2-nonanone, an aversive

odor recognized by AWB[90], wild-type animals displayed a non-dimorphic response

(Supplementary Fig. 5e, g-h). However, *npr-5* mutants showed higher activation in both sexes

when compared to their wild-type counterparts, with *npr-5* males displaying dimorphic heightened

activation (Supplementary Fig. 5e-h). Together, our results support the dimorphic inhibitory role

of *npr-5* in the proper integration of PA14 signals.

Even though we observe no change in *npr-5* expression in AIY (Fig. 3g), we found a stark decline

in the AIY soma calcium activity in only naïve, but not trained, *npr-5* mutants (Fig. 5e-g,

Supplementary Fig. 6a-b, e) compared to wild-type animals (Fig. 2e, f, k, Supplementary Fig. 2i-

l), suggesting that *npr-5* is involved in the recognition and processing of PA14 in the naïve state.

Similar observations were made in the axonal responses of AIY to PA14 odor in naïve mutant

hermaphrodites and males (Fig. 5h-j, Supplementary Fig. 6c-d, h). In contrast to wild-type males,

which retained high activation after training (Fig. 2h, l), mutant males showed reduced activation

(Fig. 5h-j, Supplementary Fig. 6c-d, h). The lack of change in responses between the naïve and

trained states is also reflected in the latency and the activation time of AIY in mutant animals

(Supplementary Fig. 6f-g, i-j). Taken together, these results suggest that *npr-5* plays a crucial role

in the integration of PA14 naïve cues in both sexes, but also the trained responses of males. This

is highlighted also at the synaptic level where the RAB-3 expression of *npr-5* mutant males

displayed an increase in mean fluorescence after training (Fig. 5k), unlike wild-type animals (Fig.

2o) and *npr-5* mutant hermaphrodites (Fig. 5k). Thus, exposure to PA14 is processed

dimorphically between the sexes, altered partly through modulation by *npr-5* in males.

Figure 5. **NPR-5 modulates state and sex-specific neuronal activity. a-b, e-f, h-i** From top to

bottom, SEM traces of GCaMP signals of select neurons of *npr-5* hermaphrodites and males in

naïve (magenta) and trained (cyan) animals. The shaded region corresponds to the exposure

window of PA14 supernatant. OP50 supernatant was used as the buffer (See Methods). AWB soma

calcium response of naïve and trained **(a)** *npr-5* mutant hermaphrodites (OP50 n = 28, PA14 n =

29) and **(b)** *npr-5* mutant males to PA14 supernatant (OP50 n = 30, PA14 n = 28). **c-d**

Quantification of **c** the OFF response and **d** AUC of OFF response of AWB. **e** AIY soma calcium

levels in naïve (n = 21) and trained (n = 23) *npr-5* mutant hermaphrodites. **f** AIY soma calcium

levels in naïve (n = 20) and trained (n = 21) *npr-5* mutant males. **g** Quantification of the ON

response of AIY soma. **h** AIY axon calcium levels in naïve (n = 20) and trained (n = 19) *npr-5*

mutant hermaphrodites. **i** AIY axon calcium levels in naïve (n = 19) and trained (n = 20) *npr-5*

mutant males. **j** Quantification of the ON response of AIY axon. **k** Mean quantification of synaptic

*rab-3::mCherry* fluorescence in naïve and trained *npr-5* mutant hermaphrodite and male AIY

axon. Each dot represents a puncta from n = 10-17 animals per group. In **(a-b, e-f, h-i)** change in

fluorescence was calculated as for Figure 2 (see Methods). In **(c-d, g, j, k)**, statistical test used is

Kruskal-Wallis test with Dunn's multiple comparisons test was performed. Box and Whisker plots

with mean and entire data range. P values are indicated in the figure.

**NPR-5 overexpression curbs state-specific male behavior**

Our pathogen-sensitivity and calcium-imaging results suggest that while males possess the ability

to recognize and potentially process the PA14 experience (Fig. 1c-d, Fig. 2), learning is repressed.

*C. elegans* males will abandon food in search of mates[24] and switch from avoidance to attraction

of aversive stimuli if hermaphrodites have been experienced[18,91]. Thus, we hypothesized that

perhaps repression in our paradigm is a result of the inherent evolutionary drive for mate search.

Therefore, diminishing males' sexual drive could lead to PA14 avoidance. To assess this, we

trained animals with the other sex ("sex-exposed") or without ("virgin") in the presence of either

OP50 or PA14 (Fig. 6a) and checked their choice indices. Sexual conditioning would predict that

in the presence of hermaphrodites, the males would be highly attracted to the coupled stimulus, as

evidenced by the retention of attraction to OP50 by the naïve hermaphrodite-exposed males (Fig.

6b). Surprisingly, we observed that, as opposed to virgin-trained males, which retain their

attraction to PA14, trained hermaphrodite-exposed males show decreased attraction to PA14 (Fig.

6c). Our results suggest that reducing the males' sexual or mate-searching drive might allow the

trained decision-making process to flow uninterrupted. To assess the potential role of *npr-5* in

regulating the integration of the internal state into the learning circuit, we overexpressed *npr-5* in

sensory neurons of wild-type males to simulate a perpetual 'naïve' state (Fig. 6b). These males

failed to avoid PA14, even after training, regardless of their sexual status (Fig. 6c). Taken together,

these results suggest that sex-specific needs act as a regulator of male learning, dictated and

processed by the inherent neuromodulatory state, where *npr-5* in the sensory neurons of males

halts potential learning by modulating the diverse variables that affect male decision-making

(Supplementary Fig. 7).

Figure 6. **The sexual status determines the behavioral outcomes. a** Schematic representation of

the conditioning training paradigm. A group of *C. elegans* was separated into males or

hermaphrodites after late L4, exposed to OP50 (magenta) or PA14 (cyan) for 6 hours either

separately ("virgin") or both sexes together ("sex-exposed"), and then assayed for their choice

index against OP50 and PA14 (See Methods). **b** Choice Index of OP50-exposed virgin and sex-

exposed wild-type hermaphrodites (virgin N = 5, sex-exposed N = 6), wild-type males (virgin,

sex-exposed N = 7) and *npr-5*-overexpression mutant (p*tax-4*::*npr-5*cDNA) males (virgin N = 4,

sex-exposed N = 6). **c** Choice Index of PA14-exposed virgin and sex-exposed wild-type

hermaphrodites (virgin N = 5, mated N = 5), wild-type males (virgin N = 7, mated N = 7) and *npr-*

*5*-overexpression males (p*tax-4*::*npr-5*cDNA virgin N = 4, mated N = 6). N refers to number of

independent assays with each dot representing a single population assay of 50-200 animals/assay.

In (**b**, **c**), statistical test used is two-way ANOVA with Tukey's multiple comparisons test. Box

and Whisker plots with mean and entire data range. P values are indicated in the figure.

**Discussion**

Sex-specific behaviors have long been recognized as both products and drivers of sexual selection,

however learning, a sex-shared trait, would naïvely be expected to withstand the forces of sexual

selection, as it is crucial to an organism's survival. Notably, a recent study demonstrated that in a

limited information paradigm, female rats outperform males[92]. These findings underscore the

importance of evaluating learning within the context of survival, where information is rapid,

limited, and temporal. In our study, we investigated the learning performance of both *C. elegans*

sexes in a multi-factorial, ethologically relevant paradigm, specifically focusing on pathogen

avoidance. Our results reveal an intriguing behavioral dimorphism: Males do not learn to avoid

the pathogenic bacteria PA14 as efficiently and rapidly as hermaphrodites.

The exposure to PA14 generates a neural representation of that experience[26,38,39]. Indeed, we

observe that two major olfactory neurons, AWC and AWB, and a cholinergic interneuron AIY,

reflect that training experience. Consistent with our notion that males can process the training

experience, we observe that the soma of AIY reduce its responses to PA14 supernatant, in a manner

similar to that of hermaphrodites, while there was no change in AWC after training in either of the

sexes. AWB displayed a biphasic response, with naïve males showing higher ON response than

naïve hermaphrodites and remained unaffected by training, while the OFF responses of AWB were

sexually dimorphic after training. The ON response in AWB could stem from a possible integration

mechanism of odors of both OP50 and PA14, a mechanism previously described by Yang et al[90]

or it is a response to the removal of OP50[38]. The OFF response of AWB increased in

hermaphrodites but reduced in males. Parallelly, the axonal regions of male AIY remain unaffected

but that of hermaphrodites showed a reduction analogous with their AIY somas. Although

compartmentalization of information processing has been shown before[93], we uncover a possible
sexually dimorphic component. The neuropeptide receptor *npr-5*, maintains these neural states as
its mutation caused a switch in male AWB OFF responses, which heightened after training, and a
complete silencing of naïve AIY responses. These point to the existence of a sensory gating
mechanism in the males where information is diverted, partly through *npr-5*, so as to continue
choosing PA14 over OP50 until the repression is alleviated.

Thus, in our investigation of the underlying molecular mechanisms that drive these behavioral
disparities, we identified several notable divergences in how males and hermaphrodites process
shared information during the learning process (Supplementary Fig. 7). First, learning in
hermaphrodites appears to be contingent on their nervous system, whereas males require the
combined processing of both their nervous system and gut to make critical choices between benign
and pathogenic bacteria. This indicates the existence of a complex regulatory mechanism in males
that overrides the immediate threat of infection, a finding in line with recent evidence highlighting
the importance of the gut-brain axis in behavior regulation[37,94]. Our study further suggests the
presence of a sexually dimorphic component. A potential explanation is the dispersed network of
neuropeptides we uncovered that could partially function from the gut. Second, our investigation
of neuronal correlates of learning revealed distinct sex-specific activation levels and responses to
PA14 stimuli in the sensory neurons and interneurons examined. This observation adds another
facet to the developing proposal of the sex differences in synaptic mechanisms that underlie
plasticity, learning, and memory[7]. Sex hormones have been shown to play a role in learning[95,96].
Importantly, our study highlights the role genetic sex plays in neuronal function independent of
sex hormones. Lastly, the MAR-seq analysis of naïve and trained animals points to transcriptomic
differences between males and hermaphrodites, particularly within the underlying
neuromodulatory network that potentially contributes to the observed behavioral differences.
Similar findings have been reported in butterflies, where a genomic analysis revealed a tissue and
gene family-specific bias between the sexes that could explain learning dimorphisms[97]. Recent
evidence highlights the important role of neuromodulators in recognizing and avoiding PA14[54,58].
Consistent with this, we demonstrate that the sensory modulation of NPR-5 expression is not only
crucial for male behavior but also for the maintenance of neuronal calcium response patterns.
Further studies are required to examine this effect in other neurons as well as its universality in
multiple behavioral contexts.

A key question arises regarding the potential influence of evolutionary pressures. Our data suggest

that while males do eventually learn to avoid PA14, they might exhibit short-term repression as a

behavioral adaptation. Drawing a parallel to models of foraging animals that exhibit

overharvesting or over-staying[98], it may be evolutionarily beneficial for *C. elegans* males to stay

in a patch of pathogenic bacterium, despite immunological activation, to maximize the chances of

encountering incoming hermaphrodites. Supporting this notion is the observed reduction in

attraction towards PA14 in mated males. One interpretation of this finding is that by removing the

need to search for mates, the mating pressure is reduced in males, thus allowing them to activate

the otherwise gated behavior. This observation is further supported by recent evidence of increased

mating by hermaphrodites previously exposed to pathogenic bacterium[99]. It is tempting to

speculate that these two pathways could have co-evolved to maximize organismal fitness under

dire conditions. In such a scenario, the overall evolutionary demand of the species is fulfilled,

albeit in a complex sex-specific manner. A fascinating example is the lower participation of the

male bumblebee in cognitive tasks[100], even though they share comparable learning abilities with

sterile worker females[101], suggesting that nuanced sex-specific roles alter the perceived learning

objective for each sex. Therefore, considering the sex differences in decision-making behavior

across organisms, it stands to reason that sexual selection impinges on learning behaviors to favor

sex-specific outcomes. By assessing the influence of sexual identity at key levels involved in

learning, such as molecular and neuronal circuits, we emphasize the importance of considering sex

differences in learning and memory disorders.

**Methods**

<u>Strains and experimental design</u>

Nematodes were cultivated on regular NGM plates on *Escherichia coli* strain OP50 at room

temperature unless otherwise mentioned. *him-5*(e1490) were used as a source for wild-type males.

All transgenic and bacterial strains are listed in Supplementary Data 3.

<u>PA14 aversive learning</u>

Animals were age-synchronized by bleaching, unless otherwise specified, and were grown at room

temperature on OP50 plates until they were late L4, sex-separated into males and females for

behavioral assays as young adults. The training and assay plates were slightly modified from [27].

Briefly, OP50 and PA14 were streaked from glycerol stocks onto LB plates and incubated at 37°C

overnight. They were then moved to 4°C and used for up to a week. To prepare OP50 or PA14

cultures used in the training and assay plates, a small amount of bacteria from streaked plates were

incubated in 15 ml tubes containing 5 ml LB at 37°C overnight at 250 rpm. After which, the

specified amount of overnight culture (200 μl bacteria for the training plates and 10 μl for the assay

plates) was plated onto dried NGM plates and incubated at 37°C overnight and then at 26°C for

24 hours. Before being used for behavioral assays, the plates were acclimatized to room

temperature (22-23°C). Care was taken to avoid overly dried or contaminated plates. Before each

experiment, a test batch of training and assay plates were verified for behavioral consistency using

young adult hermaphrodites, failure of which resulted in fresh streaks from a glycerol stock.

For the training, sex-separated animals were washed using distilled water or M9 and introduced

into training plates for 6 hours at room temperature unless otherwise mentioned. Excess liquid was

dried with a Kim-wipe. For the choice assay, the worms were washed off the training plates and

washed twice in distilled water or M9. Then, they were placed on the choice assay plates and dried

with a Kim-wipe. Once dried, they were left at RT for an hour before being scored manually.

Choice Index $= \frac{\#\ worms\ on\ PA14 - \#\ worms\ on\ OP50}{Total\ \#\ worms}$

Learning index = Choice index$_{naïve}$ – Choice index$_{trained}$

<u>PA14 larval aversive learning</u>

To prepare the worms, L1-arrested worms were grown on regular NGM plates until they were L3,

approximately 28 hours. For the training, L3 worms were washed and put on either OP50 (control)

or PA14 training plates and then incubated at RT for 6 hours. After which, they were washed off

the plates and washed twice in distilled water or M9. Then, they were placed on choice assay plates

and dried with a Kim-wipe. Once dried, they were left at RT for an hour and then, using a sharp

surgical blade, three 1.5cm plugs from the choice areas of the plate were separated as follows:

OP50 choice, PA14 choice and Indecision choice (point of insert) and plated onto respective food

plates. After 24 hours, the plates were sexed to determine the ratio of each sex in the choice made.

<u>Fast-Killing assay</u>

The fast killing plates were prepared as described[25]. Briefly, PSG media (Peptone-Glucose-

Sorbitol (1% Bacto-Peptone; 1% NaCl; 1% glucose; 0.15M sorbitol; 1.7% Bacto-Agar)) was

poured onto 3cm plates and then left to dry. The bacterial cultures were prepared as described

above. 10 μl of the overnight culture of PA14 was then used to inoculate the plates and spread

evenly using a spreader. The worms were also prepared as described above; sex-separated young

adults of each sex (50-80 animals) were introduced to the killing plates and incubated at RT for 21

hours; the number of live worms was counted every 6 hours, with dead worms removed from the

plates. Once tabulated, the survival percentage was calculated. The assay was done in three

technical replicates.

Slow-killing assay

The slow-killing assay was based on previous methods[25] with slight modifications. The bacterial

cultures and plates were prepared as described above, except 20 μl of the overnight culture of

PA14 was used to inoculate regular NGM plates and spread evenly using a spreader. Before

introducing the worms, 40 μl of 100 mM FUDR was dropped on to the sides of the plate. The

worms were prepared as described above, and sex-separated young adults of each sex (40-100

animals) were introduced to the killing plates and incubated at RT for 96 hours, during which, the

number of live worms was counted every 24 hours, with dead worms removed from the plates.

Once tabulated, the survival percentage was calculated. The assay was done in two biological

replicates.

Lawn-leaving assay

The bacterial cultures and plates were prepared as described above, except 30 μl of the overnight

culture of OP50 or PA14 inoculum used to inoculate normal 6 cm NGM plates. Next, sex-separated

age-synchronized animals were introduced and left at RT for 6-8 hours, and then the number of

worms inside or outside the lawn was counted.

Chemotaxis Assay

Olfactory preference assays for Diacetyl and Pyrazine was conducted as described in [24,40,102], with

slight modifications. Animals were sex-separated at late L4s, washed and placed on a 10 cm plate

equidistant from two spots of ethanol-diluted odors. After 60 mins, the animals within 1.5 cm of

each odor spot was manually counted to achieve the olfactory preference index, OPI = (b-a)/(a+b),

where a and b are the two odor spots.

Sex-exposed learning assay

The assay for the virgin groups was carried out as described above. However, for the mated groups,

hermaphrodites were introduced into the training plates at the start of the assay, and at the end of

6 hours, the males were separated, and the downstream processing was continued as usual.

IRG-1 imaging

Age-synchronized animals were incubated on OP50 or PA14 training plates for 6 hours and
imaged under a Zeiss LSM 880 confocal microscope at 25x magnification on an agar pad. The
PA14 groups were imaged first and the OP50 groups within 8 hours of incubation. Each image file
was subjected to the same post processing by converting it to 8-bit image z-projected using
maximum intensity and then subtracting background by rolling ball radius of 50 pixels and
thresholded between 4-5% before selecting the Region of Interest, which ran from start of the gut
until the end of the anterior region, using FIJI and the mean intensity was quantified.

AIY Rab-3 mCherry imaging

Age-synchronized animals were incubated on OP50 or PA14 training plates for 6 hours and
imaged under a Zeiss LSM 880 confocal at 63x magnification on an agar pad. Each image file was
subjected to the same post processing by converting it to 8-bit image, z-projected using maximum
intensity and then subtracting background using rolling ball radius of 50 and thresholded to
generate appropriate mask using analyze particles before selecting the Region of Interest in the red
channel for mean intensity quantification. Zone 2 and Zone 3 was identified as previously
described [103] and the mean intensity was quantified. To determine the delta between the training
states, the males were normalized to the hermaphrodites, with hermaphrodite mean intensity taking
a value of one.

Calcium imaging

Overnight cultures of OP50 and PA14 were centrifuged at 4000g for 10 minutes and the
supernatant was then filtered using a 0.22-micron syringe filter. Age-synchronized animals after 6
hours of training were used for imaging. Calcium imaging was performed according to Pechuk et
al. [8], with changes made to the stimulus type and duration of stimulus. Briefly, the imaging chips
were fabricated according to Chronis et al.[103], with the help of the Nanofabrication Unit at the
Weizmann Institute of Science. Two pumps control the flow of OP50 supernatant and PA14
supernatant into the microfluidic chip. The solutions were channeled through PVC tubes and using
a manual switch, the stimulus was presented to the worm, through stainless steel connectors. The
pumping rate during experiments was ~0.005 ml/min. The worms were loaded into worm inlet of
the chip by using a 1 ml syringe that had worms suspended in OP50 supernatant. 2 min habituation
was given prior to imaging with the lasers on. To ensure smooth imaging, 10 mM Levamisole was
added to all supernatants and 50 μM rhodamine B was added only to the PA14 supernatant to
visualize stimulus delivery. Confocal imaging was done using a 40x magnification water objective.

The total imaging duration was 40 sec, constituting 544 frames in total, and the stimulus duration

was 12 sec for AWC, 11 sec for AWB, and 15 sec for AIY, after an equal amount of initial time

to capture baseline for each neuron respectively. All imaging files were exported as tiff files and

the GCaMP fluorescence intensity was analyzed using FIJI. ROIs (regions of interest) of the somas

or the axon were drawn manually to best represent the signal, and their mean grey values were

quantified (Source Data). Downstream data processing was performed using MATLAB; for each

worm, the baseline fluorescent level (F0) was calculated by averaging the mean grey values of 100

frames before stimulus delivery, then for each frame, the ΔF was calculated by subtracting F0 from

the value of that time point, and the result was divided by F0, to normalize differences in the

fluorescence baseline levels between individuals (ΔF/F0). A similar analysis was carried out in the

stimulus channel of the images to align the responses to the start of the stimulus (AIY) or the end

of the stimulus (AWB and AWC). Once aligned, the ON response was defined as the difference

in mean of 10 seconds post stimulus start and the mean of 5 seconds pre-stimulus start. OFF

response was defined as the difference in mean of 10 seconds post stimulus stop and the mean of

5 seconds pre-stimulus start to keep the baseline for both ON and OFF responses consistent. The

latency of the response was defined as the time it took to reach 2 times the SD of the mean before

stimulus start. A similar threshold was used to calculate the time of neuronal activity from the start

of the stimulus to the end of the recording for each animal.

For calcium imaging with 2-nonaone as the stimulus, the same protocol was followed as described

above, except that the buffer used was NGM buffer and the stimulus was $10^5$ diluted 2-nonaone.

Transcriptomics analysis

Approximately 700–1000 animals of each sex were subjected to PA14 or OP50 training for 6 hours

and washed with M9 to remove any bacterial remnants. Each training group for each sex had 4

biological replicates. The worm pellet was flash-frozen in 500μl TRIzol using liquid nitrogen and

stored at -80°C. Total RNA was extracted using the TRIzol LS (Invitrogen) protocol. The RNA

from the post isopropanol precipitation step was resuspended in extraction buffer from the

PicoPure (Arcturus) RNA isolation kit, and further processing was performed according to the

manufacturer's recommendations. After determining the RNA concentration, using a Qubit

Fluorometer (model number 4, Invitrogen), and quality, using a Bioanalyzer (Agilent), the samples

with a RIN-Score >8 (Supplementary Data 1) were used to create RNA-seq libraries, as previously

described using the MARS-seq procedure[49]. The FASTQ files were then processed using the user-

friendly transcriptome analysis pipeline (UTAP) (https://utap.wexac.weizmann.ac.il/)[104]. The

reads were aligned against the *C. elegans* reference genome WBcel235. The data thus captured

was analyzed against the Wormbase database for Gene IDs. Using Python, PCA plots were plotted

using seaborn. The data was filtered by sex using the following parameters: Basemean >5 and p-

value >0.05 from each sex (naïve vs trained) for all analyses and genes with Basemean >5 was

taken ahead for only volcano plots. The p-values for neuronal genes were corrected using the

Benjamini-Hochberg method (Supplementary Data 2). The heatmaps and volcano plots were

plotted using the Heatmap and EnhancedVolcano functions of R studio (2022.12.0+353),

respectively. The resultant data was further taken for GO term analysis of genes from each sex

(naïve vs trained) using ShinyGo[105] and the circular barplot was made using ggplot of R studio

(2022.12.0+353).

RT-PCR

Total RNA was extracted for all groups using the protocol described above in 'Transcriptomics

analysis'. The isolated RNA was converted to cDNA using random hexamers from the

SuperScript[TM] IV First-Strand Synthesis system. After the concentration was assessed using

NanoDrop, qPCR reactions were set up in 10ul reaction mixtures as per manufacturer instructions

using Fast SYBR Green Master Mix and the StepOnePlus[TM] Real-Time PCR system. Ct values

were used to quantify the relative expression of the target gene using the $2-\Delta\Delta Ct$ method. *nhr-*

*23* was used as the housekeeping gene.

Dye staining of *npr-5::GFP* animals

Animals at the YA stage were washed with M9 buffer and incubated in 1ml M9 and 5μl DiD dye

(Vybrant DiD Cell-labelling solution, ThermoFisher), for 1 hour, on a rotor at ~40 rpm. The worms

were then transferred to food plates and the M9 allowed to dry. The animals were then sex-

separated and transferred to either OP50 or PA14 training plates and incubated for 6 hours. Next,

the worms were transferred to an agar pad and anesthetized with 5μl of 100mM $NaN_3$ and imaged

under a Zeiss LSM 880 using a 40x water objective. The Region of Interest was selected in FIJI,

using the DiD staining, for the positions of ASJ and AWB, while a co-injection marker was used

to assess the position of AIY. The mean intensity was quantified after background correction and

thresholding.

CRISPR/Cas-9 mediated *npr-5* deletion

The CRISPR strategy was based on Paix et al.[106]. Recombinant cas9 (IDT) was injected into N2

gonads with tracrRNA (IDT) and three crRNAs, one targeting the dpy-10 locus (Paix et al. 2015)

and two flanking the *npr-5* locus (GATAACGTTGCTGATGGTGG, located 52bp downstream of

*npr-5* start codon, and GAGACGGGCAAGATGAACAC, located 54bp downstream of stop

codon). The repair template was the ssODN

CCATTTCGACCATTTCAACCACAACGACTCCCTCCGTTCATCTTGCCCGTCTCTCGAA

AAATTCCATTAT, composed of 35nt homology arms from each side of the *npr-5* locus. Rol/dpy

F1 progeny were screened by PCR to identify the *npr-5(ety11)* deletion allele. Three primers were

included in the reaction: TTCGATGCATGGTTAGTTCG (*npr-5* CR_F1),

TTTCCATGTCACTCGTCAGC (*npr-5* CR_R1), GGAAATTCCCGGAAAATGAT (*npr-5*

CR_R2), resulting in a 347bp band in WT animals and a 577bp band in ety11 animals.

The *syb5786* allele was generated by SunyBiotech.

<u>Transgenic strains/molecular cloning</u>

The feminization of the gut was generated by expressing *tra-2[intracellular]* under the *elt-2*

promoter, cloned by Gibson assembly[90]. *elt-2* promoter was amplified from a plasmid provided by

Prof. Sivan Korenblit.

To express *npr-5* cDNA specifically in AWB, *npr-5* cDNA was obtained from Genescript and

fused to 3.5kb promoter of *str-1* and upstream of SL2::NLS::tagRFP::unc-54 3'UTR using PCR

fusion[107].

Primers used are in Supplementary Data 3.

<u>RNAi</u>

RNA interference of select genes was induced using the feeding method [108]. Briefly, L4

hermaphrodites were fed HT115 bacteria carrying the dsRNA for the relevant genes including a

control empty vector. Multiple plates for each gene were maintained to achieve sufficient young

adult males to perform the learning assay as described above. Fresh *pos-1* control was prepared

for each RNAi experiment to verify efficiency (Only batches > 95% were used for behavioral

assays).

<u>Quantification and statistical analysis</u>

Each figure legend contains details of the quantification, statistical tests performed and the sample

sizes of each individual experiment. All statistical analysis was performed on GraphPad Prism 10.

**Acknowledgments**

We thank the members of Oren-Suissa lab for their insights regarding the manuscript. We especially thank Dr. Yehuda Salzberg who helped us in generating the *npr-5* CRISPR strain used in this study. We would like to thank Mario De Bono for furnishing us with much-needed strains. Some strains were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). We would also like to thank Prof. Sivan Korenblit for plasmids and Prof. Ravid Straussman for providing us with *S. aureus*. We would like to extend our gratitude to the Nano-Fabrication unit and the Genomics Sandbox unit at the Life Science Core Facility of Weizmann Institute of Science for helping us with the PDMS chips and the transcriptomics, respectively. We thank WormBase, the online database of *C. elegans* genes, for the excellent curation of various information regarding each gene. MOS acknowledges financial support from the European Research Council ERC-2019-STG 850784, Israel Science Foundation grant 961/21, Dr. Barry Sherman Institute for Medicinal Chemistry, Sagol Weizmann-MIT Bridge Program and the Azrieli Foundation. MOS is the incumbent of the Jenna and Julia Birnbach Family Career Development Chair.

**Authors contribution**

SPK conducted and analyzed the experiments. RH performed the library preparation and sequencing for the transcriptomics experiment. AG performed and analyzed the *npr-5* dye staining experiment. MOS supervised and designed the experiments. SPK and MOS wrote the paper.

**Competing interests**

The authors declare no competing interests.

**Data availability**

All data generated and used for analysis are included in the manuscript along with the supplementary information files and Source Data files. The sequence data generated in the study has been deposited in the GEO database under the accession code GSE273091. The RNA-seq data is provided in Supplementary Data 1 and Supplementary Data 2. Quantifications for calcium imaging are added to the Source Data. Raw images for all the imaging experiments can be found at (10.34933/2c05a150-4238-4273-93ec-b419d2c5429f). Source Data are provided in this paper.

**Code availability**

Code generated in this study is available at (10.34933/2c05a150-4238-4273-93ec-b419d2c5429f).

**References**

1. Mery, F. Natural variation in learning and memory. *Curr. Opin. Neurobiol.* **23**, 52–56 (2013).

2. Crossley, M., Benjamin, P. R., Kemenes, G., Staras, K. & Kemenes, I. A circuit mechanism linking past and future learning through shifts in perception. *Sci. Adv.* **9**, eadd3403 (2023).

3. Asok, A., Leroy, F., Rayman, J. B. & Kandel, E. R. Molecular Mechanisms of the Memory Trace. *Trends Neurosci.* **42**, 14–22 (2019).

4. Janicke, T., Häderer, I. K., Lajeunesse, M. J. & Anthes, N. Darwinian sex roles confirmed across the animal kingdom. *Sci. Adv.* **2**, e1500983 (2016).

5. Connallon, T. The geography of sex-specific selection, local adaptation, and sexual dimorphism. *Evolution* **69**, (2015).

6. Paletta, P., Bass, N., Aspesi, D. & Choleris, E. Sex Differences in Brain Function and Dysfunction. *Curr Top Behav Neurosci* **62**, 207–234 (2022).

7. Gall, C. M., Le, A. A. & Lynch, G. Sex differences in synaptic plasticity underlying learning. *J. Neurosci. Res.* **101**, 764–782 (2023).

8. Pechuk, V. *et al.* Reprogramming the topology of the nociceptive circuit in C. elegans reshapes sexual behavior. *Curr Biol* (2022) doi:10.1016/j.cub.2022.08.038.

9. Beck, K. D. & Luine, V. N. Evidence for sex-specific shifting of neural processes underlying learning and memory following stress. *Physiol Behav* **99**, 204–211 (2010).

10. Ehmann, B. *et al.* Immature wild orangutans acquire relevant ecological knowledge through sex-specific attentional biases during social learning. *Plos Biol* **19**, e3001173 (2021).

11. Carazo, P., Noble, D. W. A., Chandrasoma, D. & Whiting, M. J. Sex and boldness explain individual differences in spatial learning in a lizard. *Proc Royal Soc B Biological Sci* **281**, 20133275 (2014).

12. Chiu, H. *et al.* A circuit logic for sexually shared and dimorphic aggressive behaviors in Drosophila. *Cell* **184**, 507-520.e16 (2021).

13. Verzijden, M. N. *et al.* The impact of learning on sexual selection and speciation. *Trends Ecol. Evol.* **27**, 511–519 (2012).

14. Møller, A. P. The Behavior of Animals, 2nd Edition. 315–341 (2021) doi:10.1002/9781119109556.ch12.

15. Molina-García, L. & Barrios, A. Sex differences in learning—shared principles across taxa.
*Curr. Opin. Physiol.* **6**, 65–74 (2018).

16. McCarthy, M. M. & Arnold, A. P. Reframing sexual differentiation of the brain. *Nature*
*Neuroscience* **14**, 677–683 (2011).

17. Rahmani, A. & Chew, Y. L. Investigating the molecular mechanisms of learning and
memory using Caenorhabditis elegans. *J. Neurochem.* **159**, 417–451 (2021).

18. Sakai, N. *et al.* A sexually conditioned switch of chemosensory behavior in C. elegans. *PLoS*
*ONE* **8**, e68676 (2013).

19. Ghosh, D. D., Nitabach, M. N., Zhang, Y. & Harris, G. Multisensory integration in C.
elegans. *Curr. Opin. Neurobiol.* **43**, 110–118 (2017).

20. Stein, G. M. & Murphy, C. T. C. elegans positive olfactory associative memory is a
molecularly conserved behavioral paradigm. *Neurobiol. Learn. Mem.* **115**, 86–94 (2014).

21. Kaletsky, R. *et al.* C. elegans interprets bacterial non-coding RNAs to learn pathogenic
avoidance. *Nature* **586**, 445–451 (2020).

22. Sasakura, H. & Mori, I. Behavioral plasticity, learning, and memory in C. elegans. *Curr.*
*Opin. Neurobiol.* **23**, 92–99 (2013).

23. Setty, H. *et al.* Sexually dimorphic architecture and function of a mechanosensory circuit in
C. elegans. *Biorxiv* 2022.02.18.481005 (2022) doi:10.1101/2022.02.18.481005.

24. Ryan, D. A. *et al.* Sex, age, and hunger regulate behavioral prioritization through dynamic
modulation of chemoreceptor expression. *Curr Biol* **24**, 2509–2517 (2014).

25. Tan, M.-W., Mahajan-Miklos, S. & Ausubel, F. M. Killing of Caenorhabditis elegans by
Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proc National Acad*
*Sci* **96**, 715–720 (1999).

26. Liu, H. *et al.* Forgetting generates a novel state that is reactivatable. *Sci Adv* **8**, eabi9071
(2022).

27. Zhang, Y., Lu, H. & Bargmann, C. I. Pathogenic bacteria induce aversive olfactory learning
in Caenorhabditis elegans. *Nature* **438**, 179–184 (2005).

28. Irazoqui, J. E. *et al.* Distinct Pathogenesis and Host Responses during Infection of C. elegans
by P. aeruginosa and S. aureus. *PLoS Pathog.* **6**, e1000982 (2010).

29. Otarigho, B., Butts, A. F. & Aballay, A. Neuronal NPR-15 modulates molecular and
behavioral immune responses via the amphid sensory neuron-intestinal axis in C. elegans. *eLife*
**12**, RP90051 (2024).

30. Estes, K. A., Dunbar, T. L., Powell, J. R., Ausubel, F. M. & Troemel, E. R. bZIP
transcription factor zip-2 mediates an early response to Pseudomonas aeruginosa infection in
Caenorhabditis elegans. *Proc National Acad Sci* **107**, 2153–2158 (2010).

31. JebaMercy, G. & Balamurugan, K. Effects of sequential infections of Caenorhabditis elegans
with Staphylococcus aureus and Proteus mirabilis. *Microbiol. Immunol.* **56**, 825–835 (2012).

32. Troemel, E. R. *et al.* p38 MAPK Regulates Expression of Immune Response Genes and
Contributes to Longevity in C. elegans. *PLoS Genet.* **2**, e183 (2006).

33. Powell, J. R., Kim, D. H. & Ausubel, F. M. The G protein-coupled receptor FSHR-1 is
required for the Caenorhabditis elegans innate immune response. *Proc. Natl. Acad. Sci.* **106**,
2782–2787 (2009).

34. Miller, E. V., Grandi, L. N., Giannini, J. A., Robinson, J. D. & Powell, J. R. The Conserved
G-Protein Coupled Receptor FSHR-1 Regulates Protective Host Responses to Infection and
Oxidative Stress. *PLoS ONE* **10**, e0137403 (2015).

35. Prakash, D. *et al.* 1-Undecene from Pseudomonas aeruginosa is an olfactory signal for flight-
or-fight response in Caenorhabditis elegans. *Embo J* **40**, e106938 (2021).

36. O'Donnell, M. P., Fox, B. W., Chao, P.-H., Schroeder, F. C. & Sengupta, P. A
neurotransmitter produced by gut bacteria modulates host sensory behaviour. *Nature* **583**, 415–
420 (2020).

37. Singh, J. & Aballay, A. Intestinal infection regulates behavior and learning via
neuroendocrine signaling. *Elife* **8**, e50033 (2019).

38. Ha, H. *et al.* Functional organization of a neural network for aversive olfactory learning in
Caenorhabditis elegans. *Neuron* **68**, 1173–1186 (2010).

39. Jin, X., Pokala, N. & Bargmann, C. I. Distinct Circuits for the Formation and Retrieval of an
Imprinted Olfactory Memory. *Cell* **164**, 632–643 (2016).

40. Bargmann, C. I., Hartwieg, E. & Horvitz, H. R. Odorant-selective genes and neurons mediate
olfaction in C. elegans. *Cell* **74**, 515–527 (1993).

41. Troemel, E. R., Kimmel, B. E. & Bargmann, C. I. Reprogramming Chemotaxis Responses:
Sensory Neurons Define Olfactory Preferences in C. elegans. *Cell* **91**, 161–169 (1997).

42. Zhang, Y. Neuronal mechanisms of Caenorhabditis elegans and pathogenic bacteria
interactions. *Curr. Opin. Microbiol.* **11**, 257–261 (2008).

43. Hawk, J. D. *et al.* Integration of Plasticity Mechanisms within a Single Sensory Neuron of
C. elegans Actuates a Memory. *Neuron* **97**, 356-367.e4 (2018).

44. Chandra, R. *et al.* Sleep is required to consolidate odor memory and remodel olfactory
synapses. *Cell* **186**, 2911-2928.e20 (2023).

45. Satoh, Y. *et al.* Regulation of Experience-Dependent Bidirectional Chemotaxis by a Neural
Circuit Switch in Caenorhabditis elegans. *J. Neurosci.* **34**, 15631–15637 (2014).

46. Hendi, A., Kurashina, M. & Mizumoto, K. Intrinsic and extrinsic mechanisms of synapse
formation and specificity in C. elegans. *Cell. Mol. Life Sci.* **76**, 2719–2738 (2019).

47. Richmond, J. E. & Broadie, K. S. The synaptic vesicle cycle: exocytosis and endocytosis in
Drosophila and C. elegans. *Curr. Opin. Neurobiol.* **12**, 499–507 (2002).

48. Colón-Ramos, D. A., Margeta, M. A. & Shen, K. Glia promote local synaptogenesis through
UNC-6 (netrin) signaling in C. elegans. *Science* **318**, 103–106 (2007).

49. Jaitin, D. A. *et al.* Massively parallel single-cell RNA-seq for marker-free decomposition of
tissues into cell types. *Science* **343**, 776–779 (2014).

50. Haque, R. *et al.* Sex-specific developmental gene expression atlas unveils dimorphic gene
networks in C. elegans. *Nat. Commun.* **15**, 4273 (2024).

51. Okkema, P. G. & Kimble, J. Molecular analysis of tra-2, a sex determining gene in
C.elegans. *EMBO J.* **10**, 171–176 (1991).

52. Barr, M. M. & Sternberg, P. W. A polycystic kidney-disease gene homologue required for
male mating behaviour in C. elegans. *Nature* **401**, 386–389 (1999).

53. Kawli, T. & Tan, M.-W. Neuroendocrine signals modulate the innate immunity of
Caenorhabditis elegans through insulin signaling. *Nat Immunol* **9**, 1415–1424 (2008).

54. Yu, Y., Zhi, L., Guan, X., Wang, D. & Wang, D. FLP-4 neuropeptide and its receptor in a
neuronal circuit regulate preference choice through functions of ASH-2 trithorax complex in
Caenorhabditis elegans. *Sci Rep-uk* **6**, 21485 (2016).

55. Deng, P. *et al.* Mitochondrial UPR repression during Pseudomonas aeruginosa infection
requires the bZIP protein ZIP-3. *Proc National Acad Sci* **116**, 6146–6151 (2019).

56. Wu, T. *et al.* Pheromones Modulate Learning by Regulating the Balanced Signals of Two
Insulin-like Peptides. *Neuron* **104**, 1095-1109.e5 (2019).

57. Zheng, Z. *et al.* GABAergic synapses suppress intestinal innate immunity via insulin
signaling in Caenorhabditis elegans. *Proc National Acad Sci* **118**, e2021063118 (2021).

58. Marquina-Solis, J. *et al.* Peptidergic signaling controls the dynamics of sickness behavior in
Caenorhabditis elegans. *Biorxiv* 2022.04.16.488560 (2022) doi:10.1101/2022.04.16.488560.

59. Chang, H. C., Paek, J. & Kim, D. H. Natural polymorphisms in C. elegans HECW-1 E3
ligase affect pathogen avoidance behaviour. *Nature* **480**, 525–529 (2011).

60. Chen, Z. *et al.* Two Insulin-like Peptides Antagonistically Regulate Aversive Olfactory
Learning in C. elegans. *Neuron* **77**, 572–585 (2013).

61. Lee, K. & Mylonakis, E. An Intestine-Derived Neuropeptide Controls Avoidance Behavior
in Caenorhabditis elegans. *Cell Reports* **20**, 2501–2512 (2017).

62. Cao, X., Kajino-Sakamoto, R., Doss, A. & Aballay, A. Distinct Roles of Sensory Neurons in
Mediating Pathogen Avoidance and Neuropeptide-Dependent Immune Regulation. *Cell Reports*
**21**, 1442–1451 (2017).

63. Zhang, X. & Zhang, Y. DBL-1, a TGF-β, is essential for Caenorhabditis elegans aversive
olfactory learning. *Proc National Acad Sci* **109**, 17081–17086 (2012).

64. Meisel, J. D., Panda, O., Mahanti, P., Schroeder, F. C. & Kim, D. H. Chemosensation of
Bacterial Secondary Metabolites Modulates Neuroendocrine Signaling and Behavior of
C. elegans. *Cell* **159**, 267–280 (2014).

65. Dasgupta, M. *et al.* NHR-49 Transcription Factor Regulates Immunometabolic Response and
Survival of Caenorhabditis elegans during Enterococcus faecalis Infection. *Infect. Immun.* **88**,
e00130-20 (2020).

66. Lanjuin, A., VanHoven, M. K., Bargmann, C. I., Thompson, J. K. & Sengupta, P. Otx/otd
Homeobox Genes Specify Distinct Sensory Neuron Identities in C. elegans. *Dev. Cell* **5**, 621–
633 (2003).

67. Liu, F. *et al.* The homeodomain transcription factor CEH-37 regulates PMK-1/p38 MAPK
pathway to protect against intestinal infection via the phosphatase VHP-1. *Cell. Mol. Life Sci.* **80**,
312 (2023).

68. O'Rourke, E. J. & Ruvkun, G. MXL-3 and HLH-30 transcriptionally link lipolysis and
autophagy to nutrient availability. *Nat. Cell Biol.* **15**, 668–676 (2013).

69. Hammarlund, M., Hobert, O., Miller, D. M. & Sestan, N. The CeNGEN Project: The
Complete Gene Expression Map of an Entire Nervous System. *Neuron* **99**, 430–433 (2018).

70. Mariani, L., Lussi, Y. C., Vandamme, J., Riveiro, A. & Salcini, A. E. The H3K4me3/2
histone demethylase RBR-2 controls axon guidance by repressing the actin-remodeling gene
wsp-1. *Development* **143**, 851–863 (2016).

71. Nono, M. *et al.* Intestine-to-Germline Transmission of Epigenetic Information
Intergenerationally Ensures Systemic Stress Resistance in C. elegans. *Cell Rep.* **30**, 3207-
3217.e4 (2020).

72. Wani, K. A., Goswamy, D. & Irazoqui, J. E. Nervous system control of intestinal host
defense in C. elegans. *Curr. Opin. Neurobiol.* **62**, 1–9 (2020).

73. Sellegounder, D., Yuan, C.-H., Wibisono, P., Liu, Y. & Sun, J. Octopaminergic Signaling
Mediates Neural Regulation of Innate Immunity in Caenorhabditis elegans. *mBio* **9**, e01645-18
(2018).

74. Reddy, K. C., Andersen, E. C., Kruglyak, L. & Kim, D. H. A Polymorphism in npr-1 Is a
Behavioral Determinant of Pathogen Susceptibility in C. elegans. *Science* **323**, 382–384 (2009).

75. Styer, K. L. *et al.* Innate Immunity in Caenorhabditis elegans Is Regulated by Neurons
Expressing NPR-1/GPCR. *Science* **322**, 460–464 (2008).

76. Fruyt, N. D., Yu, A. J., Rankin, C. H., Beets, I. & Chew, Y. L. The role of neuropeptides in
learning: Insights from C. elegans. *Int. J. Biochem. Cell Biol.* **125**, 105801 (2020).

77. Gøtzsche, C. R. & Woldbye, D. P. D. The role of NPY in learning and memory.
*Neuropeptides* **55**, 79–89 (2016).

78. Pleil, K. E. *et al.* NPY signaling inhibits extended amygdala CRF neurons to suppress binge
alcohol drinking. *Nat Neurosci* **18**, 545–552 (2015).

79. Zhou, Z. *et al.* Genetic variation in human NPY expression affects stress response and
emotion. *Nature* **452**, 997–1001 (2008).

80. Naveilhan, P. *et al.* Normal feeding behavior, body weight and leptin response require the
neuropeptide Y Y2 receptor. *Nat Med* **5**, 1188–1193 (1999).

81. Cohen, H. *et al.* The Neuropeptide Y (NPY)-ergic System is Associated with Behavioral
Resilience to Stress Exposure in an Animal Model of Post-Traumatic Stress Disorder.
*Neuropsychopharmacol* **37**, 350–363 (2012).

82. Cohen, M. *et al.* Coordinated Regulation of Foraging and Metabolism in C. elegans by
RFamide Neuropeptide Signaling. *Cell Metab* **9**, 375–385 (2009).

83. Bhardwaj, A., Pandey, P. & Babu, K. Control of Locomotory Behavior of Caenorhabditis
elegans by the Immunoglobulin Superfamily Protein RIG-3. *Genetics* **214**, 135–145 (2020).

84. Lemieux, G. A. *et al.* Kynurenic Acid Is a Nutritional Cue that Enables Behavioral Plasticity.
*Cell* **160**, 119–131 (2015).

85. Choi, M.-K., Liu, H., Wu, T., Yang, W. & Zhang, Y. NMDAR-mediated modulation of gap
junction circuit regulates olfactory learning in C. elegans. *Nat. Commun.* **11**, 3467 (2020).

86. Hilbert, Z. A. & Kim, D. H. Sexually dimorphic control of gene expression in sensory
neurons regulates decision-making behavior in C. elegans. *Elife* **6**, 386 (2017).

87. Flavell, S. W. *et al.* Serotonin and the neuropeptide PDF initiate and extend opposing
behavioral states in C. elegans. *Cell* **154**, 1023–1035 (2013).

88. Mukhopadhyay, S., Lu, Y., Shaham, S. & Sengupta, P. Sensory Signaling-Dependent
Remodeling of Olfactory Cilia Architecture in C. elegans. *Dev. Cell* **14**, 762–774 (2008).

89. Komatsu, H., Mori, I., Rhee, J.-S., Akaike, N. & Ohshima, Y. Mutations in a Cyclic
Nucleotide–Gated Channel Lead to Abnormal Thermosensation and Chemosensation in C.
elegans. *Neuron* **17**, 707–718 (1996).

90. Yang, W. *et al.* Redundant neural circuits regulate olfactory integration. *Plos Genet* **18**,
e1010029 (2022).

91. Sammut, M. *et al.* Glia-derived neurons are required for sex-specific learning in C. elegans.
*Nature* **526**, 385–390 (2015).

92. Gutiérrez-Menéndez, A., Méndez, M. & Arias, J. L. Learning and metabolic brain
differences between juvenile male and female rats in the execution of different training regimes
of a spatial memory task. *Physiol Behav* **267**, 114203 (2023).

93. Rama, S., Zbili, M. & Debanne, D. Signal propagation along the axon. *Curr. Opin.*
*Neurobiol.* **51**, 37–44 (2018).

94. Morais, L. H., Schreiber, H. L. & Mazmanian, S. K. The gut microbiota–brain axis in
behaviour and brain disorders. *Nat Rev Microbiol* **19**, 241–255 (2021).

95. Hamson, D. K., Roes, M. M. & Galea, L. A. M. Comprehensive Physiology. *Compr. Physiol.*
**6**, 1295–1337 (2017).

96. Frick, K. M., Kim, J., Tuscher, J. J. & Fortress, A. M. Sex steroid hormones matter for
learning and memory: estrogenic regulation of hippocampal function in male and female rodents.
*Learn. Mem.* **22**, 472–493 (2015).

97. Ernst, D. A., Agcaoili, G. A., Merrill, A. N. & Westerman, E. L. A learning experience elicits
sex-dependent neurogenomic responses in Bicyclus anynana butterflies. *Mol Ecol* (2023)
doi:10.1111/mec.16920.

98. Kilpatrick, Z. P., Davidson, J. D. & Hady, A. E. Uncertainty drives deviations in normative
foraging decision strategies. *J Roy Soc Interface* **18**, 20210337 (2021).

99. Wu, T. *et al.* Pathogenic bacteria modulate pheromone response to promote mating. *Nature*
**613**, 324–331 (2023).

100. Manning, T. H., Austin, M. W., MuseMorris, K. & Dunlap, A. S. Equivalent learning, but
unequal participation: Male bumble bees learn comparably to females, but participate in
cognitive assessments at lower rates. *Behav Process* **193**, 104528 (2021).

101. Wolf, S. & Chittka, L. Male bumblebees, Bombus terrestris, perform equally well as
workers in a serial colour-learning task. *Anim Behav* **111**, 147–155 (2016).

102. Lee, K. & Portman, D. S. Neural sex modifies the function of a C. elegans sensory circuit.
*Current Biology* **17**, 1858–1863 (2007).

103. Chronis, N., Zimmer, M. & Bargmann, C. I. Microfluidics for in vivo imaging of neuronal
and behavioral activity in Caenorhabditis elegans. *Nature Methods* **4**, 727–731 (2007).

104. Kohen, R. *et al.* UTAP: User-friendly Transcriptome Analysis Pipeline. *BMC Bioinform.*
**20**, 154 (2019).

105. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical enrichment tool for animals and plants.
*Bioinformatics* **36**, 2628–2629 (2019).

106. Paix, A., Folkmann, A., Rasoloson, D. & Seydoux, G. High Efficiency, Homology-Directed
Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein Complexes.
*Genetics* **201**, 47–54 (2015).

107. Hobert, O. PCR fusion-based approach to create reporter gene constructs for expression
analysis in transgenic C. elegans. *BioTechniques* **32**, 728–730 (2002).

108. Ahringer, J. Reverse genetics. *WormBook* 1–43 (2006) doi:10.1895/wormbook.1.47.1.

978