Acquisition of auditory discrimination mediated by different processes through two distinct circuits linked to the lateral striatum

Susumu Setogawa[1,2,3,4,5,*], Takashi Okauchi[4], Di Hu[4], Yasuhiro Wada[6], Keigo Hikishima[7], Hirotaka Onoe[8], Kayo Nishizawa[3], Nobuyuki Sakayori[9], Hiroyuki Miyawaki[1,2], Takuma Kitanishi[2], Kenji Mizuseki[1,2], Yilong Cui[4], Kazuto Kobayashi[3*]

[1]Department of Physiology, Graduate School of Medicine, Osaka Metropolitan University, Osaka 545-8585, Japan

[2]Department of Physiology, Osaka City University Graduate School of Medicine, Osaka 545-8585, Japan

[3]Department of Molecular Genetics, Institute of Biomedical Sciences, Fukushima Medical University, Fukushima 960-1295, Japan

[4]Laboratory for Biofunction Dynamics Imaging, RIKEN Center for Biosystems Dynamics Research, Kobe 650-0047, Japan

[5]Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

[6]Laboratory for Pathophysiological and Health Science, RIKEN Center for Biosystems Dynamics Research, Kobe 650-0047, Japan

[7]Medical Devices Research Group, Health and Medical Research Institute, National Institute of Advanced Industrial Science and Technology, Tsukuba 305-8564, Japan

[8]Human Brain Research Center, Kyoto University Graduate School of Medicine, Kyoto 606-8507, Japan

[9]Department of Physiology and Oral Physiology, Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima 734-8553, Japan

*Address for correspondence:

Kazuto Kobayashi, Ph.D.

E-mail: kazuto@fmu.ac.jp

Department of Molecular Genetics, Institute of Biomedical Sciences, Fukushima Medical

University, Fukushima 960-1295, Japan.

Tel. +81-24-547-1667

Susumu Setogawa, Ph.D.

E-mail: setogawa@omu.ac.jp

Department of Physiology, Graduate School of Medicine, Osaka Metropolitan University, Osaka

545-8585, Japan

Tel. +81-6-6645-3717

Abstract

The striatum, the central hub of cortico-basal ganglia loops, contains functionally heterogeneous subregions distinguished by the topographic patterns of structural connectivity. These subregions mediate various processes of procedural learning. However, it remains unclear when and how striatal subregions engage in the acquisition of sensory stimulus-based decision-making. A neuroimaging of regional brain activity shows that the anterior dorsolateral striatum (aDLS) and posterior ventrolateral striatum (pVLS) are activated in a different temporal pattern during the acquisition phase of auditory discrimination. Chronic and transient pharmacologic manipulations show that the aDLS promotes the behavioral strategy driven by the stimulus-response association while suppressing that by the response-outcome association, and that the pVLS contributes to forming and maintaining the stimulus-response strategy. Electrophysiological recording indicates that subpopulations of aDLS neurons predominantly represent the outcome of specific behaviors at the initial period of discrimination learning, and that pVLS subpopulations encode the beginning and ending of each behavior according to the progress of learning. In addition, other subpopulations of striatal neurons indicate sustained activation after obtaining reward with distinct patterns reflecting the stimulus-response associations. Our findings demonstrate that aDLS and pVLS neurons integrate the new learning of auditory discrimination in spatiotemporally and functionally different manners.

Introduction

The neural network connecting the cerebral cortex, basal ganglia, and thalamus forms functionally heterogeneous loops distinguished by the topographic patterns of connectivity[1,2], and is implicated in various learning processes by the integration of cognitive, sensorimotor, and reward information[3,4]. The striatum is a key node in the cortico-basal ganglia loops, and its segregated subregions are considered to mediate procedural learning, such as motor skills and instrumental behavior, through parallel processing via different loops[5-7]. Previous studies in humans and non-human primates have reported that the neural activity within the striatum shifts from the associative subregions (caudate and anterior putamen) to the sensorimotor subregions (posterior putamen) as motor learning progresses[8-11]. In rodents, an instrumental behavior initiates as goal-directed actions and then transits to stimulus-response habits[12,13], requiring the dorsomedial striatum (DMS; a homologue to the caudate in primates) and the dorsolateral striatum (DLS; a homologue to the putamen in primates), respectively[14,15]. Based on these results, a model of procedural learning has been proposed, in which the functional dominance changes from the associative to sensorimotor subregions in the striatum during the learning phases[16].

In contrast, deviations from the prior model are evident in other behavioral tasks on decision-making based on external sensory cues. For example, the neural activity in the caudate in humans is associated with learning of a visuomotor association task and the execution of learned behavior[17,18]. The tail of the caudate in non-human primates contributes to the automatic, habitual behavior in decision-making task using complex visual stimuli[19]. In rodents, the DLS is involved in the acquisiton and execution of the discrimination task with visual and auditory cues, wheras the DMS madiates the performance of learned behavior[20,21]. Moreover, other striatal subregions, such as the posterior striatum and ventrolateral striatum in rodents are also engaged in the auditory discrimination[22-24]. Based on these results, a hypothesis has arisen that the acquisition of sensory cue-based decision-making requires wide-range spatiotemporal processing within the striatum. However, the neural mechanism by which the functional circuits linked to striatal subregions

change throughout learning processes is not yet fully understood.

To investigate a large-scale striatal reorganization in accordance with the acquisition of

external cue-dependent decisions, we conducted a two-alternative auditory discrimination task[25].

Using a small-animal neuroimaging technique, we found that the anterior dorsolateral striatum

(aDLS) and posterior ventrolateral striatum (pVLS) are activated to the highest level at the middle

and late stages during the acquisition phase of auditory discrimination, respectively. Then, chronic

pharmacologic manipulations confirmed that the aDLS and pVLS are necessary for discrimination

learning, and the transient manipulations also ascertained the function of these subregions in the

learning at the corresponding stages. The resultant effects of the manipulations on the behavioral

strategy showed that the aDLS promotes the strategy based on the stimulus-response association,

while suppressing that based on response-outcome associations. Additionally, the pVLS engages in

the formation and maintenance of the stimulus-response association strategy. Electrophysiological

recording of striatal neurons indicated that subpopulations of aDLS neurons mainly represent the

outcome of specific behaviors at the initial period of discrimination learning, whereas pVLS

subpopulations encode the beginning and ending of each behavior in association with the progress

of the learning. In addition, other subpopulations of aDLS and pVLS neurons show sustained

activation after obtaining reward with distinct patterns of the combination between the stimulus and

response. These findings reveal that the aDLS and pVLS regulate the acquisition of auditory

discrimination through distinct spatiotemporal and functional manners, challenging the prior model

which proposed a transition from the associative to sensorimotor striatal subregions during learning.

Results

Leaning processes of auditory discrimination

We employed a two-alternative auditory discrimination task, in which animals are required to make

new associations between a tone instruction cue (2 or 10 kHz) and a response (pressing the left or

right lever)[25], setting the interval between the stimulus presentation and lever insertion to 3 s

(Figure 1A). The success rate gradually increased, reaching a plateau around Day 13, which

persisted through Day 24 (Figure 1B). When the success rates across Days 2, 6, 10, 13 and 24 were

compared, significant differences were observed between Days 2 and 6, Days 6 and 10, as well as

Days 10 and 13, showing no significant difference between Days 13 and 24 (Figure 1C). In contrast,

the response time and omission rate did not vary among the training days (Figures 1D and 1E), and

the variation of response bias decreased along with the progress of auditory discrimination (Figure

1F). Based on changes observed in the success rate, we categorized the learning processes into two

phases: the acquisition phase, which lasted until Day 13; and the learned phase, which began after

Day 13.

Distinct brain activity patterns in striatal subregions during learning processes

To investigate dynamic changes in regional brain activity in the entire striatum during auditory

discrimination in the same animals, we conducted the positron emission tomography (PET) for

small animals with 2-deoxy-2-[^{18}F]fluoro-$_D$-glucose (^{18}F-FDG), which measures cerebral glucose

metabolism correlated to brain activity[26-28] (Figure 2A). We performed ^{18}F-FDG-PET scanning on

Day 4 of the single lever press task without sound-based decision as a control condition and during

a series of the trainings including the acquisition (Days 2, 6, and 10) and learned (Day 24) phases

(Figure 2B). The average number of lever presses in the single lever press task exceeded 80 times

(81.6 ± 1.3, mean $\pm$ s.e.m.) on Day 2 and maintained that level until Day 4 (Figure 2C). In the

following auditory discrimination task, the success rate gradually increased along with the training

days (Figure 2D), showing a significant difference among sessions in which we performed the ^{18}F-

FDG-PET scanning (Figure 2E). The response time and omission rate were consistent among the

days, and the variation of response bias became smaller as the task progresses (Figure 2E). These

performances were similar to those in the intact rats (compared to Figures 1C–1F), suggesting that

the procedure for ^{18}F-FDG-PET scans does not affect the acquisition of discrimination.

The regional brain activity related to the behavioral task was analyzed with a voxel-based

statistical parametric analysis[26], in which the ^{18}F-FDG uptake on either day of the discrimination

task was compared with that on the single lever task. The task-related brain activity in the whole

brain is summarized in Table S1. In the present study, we focused on the task-related activity in

striatal subregions. We found significant increases of the activity in the unilateral aDLS and the

bilateral pVLS on Days 6 and 10 during the acquisition phase (Figures 2F and 2G). However, no

significant change was observed in the DMS during the acquisition phase except a unilateral

decrease on Day 24 (Figure 2F). Next, to evaluate the activity related to the progress of learning,

^{18}F-FDG uptake on Days 6, 10, and 24 was compared with that on Day 2. The learning-dependent

brain activity was significantly increased in the bilateral aDLS and the unilateral pVLS on Days 6

and 10, respectively (Figure 2H), whereas the activity decreased in the bilateral DMS on Day 24

(Figure 2H). This decrease was also detected when ^{18}F-FDG uptakes were compared between Days

10 and 24 (Figure 2H), suggesting the down-regulation of the DMS activity in the learned phase.

To quantitatively validate changes in the brain activity in striatal subregions, we analyzed

the amount of ^{18}F-FDG uptake in the voxel of interest of these subregions (Figure 2I). The ^{18}F-FDG

uptake in the bilateral aDLS reached a peak on Day 6, and it decreased during the subsequent days

(Figure 2J), whereas the uptake in the bilateral pVLS gradually increased along with the progress of

learning, showing the maximal value on Day 10 (Figure 2K). The uptake in the DMS did not alter

during the acquisition phase, and it decreased on Day 24 (Figure 2L). These results suggest that the

aDLS and pVLS contribute to learning processes in a different temporal pattern during the

acquisition phase, whereas the DMS may be engaged to the execution of the behavior in the learned

phase.

Excitotoxic lesion of the aDLS and pVLS, but not the DMS, disrupts the acquisition of

auditory discrimination

To study whether the changed activity of striatal subregions is involved in learning processes, rats

received a bilateral injection of solution containing ibotenic acid (IBO; 8 mg/mL, 0.3 μL/site) or

phosphate buffered saline (PBS) into the subregions, and we conducted for the behavioral study.

The range of IBO lesion in the subregions was analyzed by immunostaining with a neuronal marker

NeuN after the behavioral tests (Figures 3A–3C). The number of single lever presses increased

similarly along with the training in both the PBS- and IBO-injected groups into the aDLS, pVLS, or

DMS along with the training (Figures 3D–3F). In the discrimination learning, increases of the

success rate were impaired in both IBO-injected aDLS and pVLS groups, compared to the

corresponding PBS-injected groups (Figures 3G and 3H). The response time in the aDLS-lesioned

group did not alter during the discrimination training, whereas that in the pVLS-lesioned group was

continuously prolonged (Figure. S1A and S1B), and the omission rate and response bias were

similar between the lesioned and the corresponding control groups (Figures S1D/S1G for aDLS

lesion and Figures S1E/S1H for pVLS lesion). In contrast, the success rate in the discrimination

task was comparable between the PBS- and IBO-injected groups into the DMS (Figure 3I). The

DMS lesion lengthened the response time through the learning (Figure S1C), whereas it did not

alter the omission rate and response bias (Figures S1F and S1I). These results indicate that the

aDLS and pVLS are required for the acquisition of auditory discrimination, whereas the DMS

contributes to the execution but not learning of the discriminative behavior, being consistent with

previous reports[29,30].

Previous studies have reported that the dorsal striatum is involved in both behavioral

strategies based on the response-outcome association and the stimulus-response association during

stimulus-response learning[12,13]. To assess the impact of striatal lesions on the behavioral strategies,

we analyzed the proportion of response attributed to two representative strategies in all responses of

each session. One is the "win-shift-win (WSW)" strategy, in which, after a correct response in the

previous trial, rats press the opposite lever in the current trial in response to a shift of the instruction

cue, resulting in the correct response. The WSW is considered to reflect the behavioral strategy

based on the stimulus-response association[31,32]. Another strategy is the "lose-shift-lose (LSL)"

strategy, in which, after an error response in the previous trial rats press the opposite lever in the

current trial despite a shift of the instruction cue, leading to another error response. The LSL is

considered to appear as a consequence of the behavioral strategy based on the response-outcome

association[31]. In the PBS-injected group into either of three striatal subregions, the proportion of

WSW and LSL strategies increased and decreased, respectively, during the acquisition phase

(Figures 3J–3L). In the IBO-lesioned group into the aDLS, the increase in the WSW proportion was

markedly impaired as compared to the corresponding control group, and the decrease in the LSL

proportion was delayed compared to the same group (Figure 3J). In the pVLS-lesioned group, the

increasing WSW proportion was also impaired as compared to the corresponding controls, although

the change in the LSL proportion was comparable to the control response (Figure 3K). In contrast,

in the DMS lesion group both WSW and LSL proportions did not show significant differences from

the corresponding values in their control group (Figure 3L). These results indicate that the aDLS

promotes the WSW strategy through stimulus-response association and suppresses the LSL strategy

rooted in response-outcome association, and that the pVLS selectively facilitates the WSW strategy

acquisition. Thus, these two subregions not only play roles in the acquisition of auditory

discrimination but also control distinct behavioral strategies during the learning processes.

After the completion of discrimination training, we tested locomotion and feeding behavior

of rats with aDLS and pVLS lesions. There was no difference in locomotor activity in the open field

and the amount of food intake in both lesioned groups (Figures S1J-S1M), suggesting that learning

deficits following excitotoxic lesions of the striatal subregions cannot be attributable to

abnormalities in these general motor behaviors.

Inhibiting aDLS and pVLS activity impairs discriminative behavior at different stages of the

acquisition phase

Our [18]F-FDG-PET imaging study indicated that the brain activity in the aDLS and pVLS was

increased in a different temporal pattern during the acquisition phase of auditory discrimination. To

examine whether neuronal activity in striatal subregions is linked with the processes at different

stages, we temporarily inhibited the activity in each subregion by the treatment with a GABA$_A$

receptor agonist muscimol (MUS). In a pilot study, we tested the effect of the bilateral injection of

several doses of MUS (0.1, 0.2, and 0.5 µg/µL) into the aDLS or pVLS on the single lever press

behavior and confirmed that the administration of the lowest dose of MUS did not affect the

frequency of single lever presses and the quantity of food intake (Figures S2A–S2D). In addition,

we divided the acquisition phase into three stages based on the success rate that included the early

($< 60\%$), middle (60–80%), and late (80–100%) stages. To achieve the stage-specific inactivation of

the aDLS or pVLS neurons, rats were given bilateral injections of MUS or saline (SAL) at three

different timings during a series of the training. The first injection was conducted on Day 2 to target

the early stage. The second and third injections were conducted on the day after the success rate had

reached 60% and 80% for the first time through the training, respectively, to target the middle or

late stage. After the injections, the rats were used for the behavioral test. Placement locations of the

injection cannula into the striatal subregions were assessed by cresyl violet staining after the

behavioral tests (Figures 4A and 4B). The success rates on the day before the injection (day N-1) at

the early, middle, and late stages were similar between the two groups that received the bilateral

injections of MUS or SAL (Figures S2E and S2F). Rats were then administered injections into the

aDLS on the next day (Day N), and the success rate in the MUS-injected groups significantly

decreased at the middle stage compared to the corresponding SAL-injected groups, but not at the

early and late stages (Figure 4C, left). In contrast, the success rate in the MUS-injected groups into

the pVLS significantly decreased at the late stage, but not at the early and middle stages, compared

with the respective SAL-injected groups (Figure 4D, left). To normalize the variation of learning

speed in the individual animals, we calculated the value by subtracting the success rate on Day N-1

from that on Day N in the individual animals. There were significant reductions in the subtracted

rate at the middle stage in the MUS-injected group into the aDLS (Figure 4C, right) and at the late

stage in the MUS-injected group into the pVLS (Figure 4D, right). These results indicate that the

aDLS and pVLS mainly function at the middle and late stages, respectively, in the acquisition phase

241 of auditory discrimination, supporting the results obtained from our imaging study.

242   Next, we analyzed the influence of temporal striatal inactivation on the behavioral strategy

243 at the middle and late stages of the acquisition phase. We calculated the proportion of WSW and

244 LSL strategies, and subtracted the values of WSW or LSL proportion on Day N-1 from those on

245 Day N. The proportion of both strategies on Day N-1 were comparable between the MUS- and

246 SAL-injected groups (Figures S2G and S2H). In the aDLS, the subtracted proportion of LSL

247 strategy at the middle stage significantly increased in the MUS-injected group as compared to the

248 SAL-injected group, although the subtracted proportion of WSW strategy tended to decrease in the

249 MUS-injected group as compared to the control group (Figure 4E), and the subtracted proportion of

250 WSW or LSL strategy at the late stage was similar between both groups (Figure 4F). In the pVLS,

251 the subtracted proportions of the WSW and LSL strategies at the middle stage were similar between

252 the MUS- and SAL-injected groups (Figure 4G), and the subtracted WSW proportion at the late

253 stage significantly decreased in the MUS-injected group as compared to the SAL-injected group,

254 although the subtracted LSL proportion was comparable between the two groups (Figure 4H). Our

255 results show that transient inhibition of aDLS activity at the middle stage affects the LSL strategy

256 and has a moderate effect on the WSW strategy, supporting that the aDLS mediates the acquisition

257 process of auditory discrimination at the middle stage at least partly through the suppression of the

258 behavior based on the response-outcome association. Although the impact of inhibition on the

259 WSW strategy was modest, excitotoxic lesion of the aDLS influenced the WSW strategy in addition

260 to the LSL strategy (see Figure 3J), suggesting that the aDLS acts to enhance the auditory

261 discrimination by promoting the stimulus-response association, together with the suppression of the

262 response-outcome association. The discrepancy in the extent of changes in the WSW strategy

263 between the two treatments may be generated from the difference in the timing or sustainability of

264 drug inhibition of the striatal subregion. In addition, our results show that pVLS inactivation at the

265 late stage selectively impairs the WSW strategy, supporting contribution of the pVLS to the process

266 via progression of the stimulus-response association.

Neuronal activity in the aDLS and pVLS during the acquisition phase

To examine the firing activity in striatal subregions during the acquisition phase of auditory

discrimination, we simultaneously recorded neuronal activity in the aDLS and pVLS at the

unilateral (left) side by using a multi-unit recording procedure (Figure 5A). Using immunostaining

for tyrosine hydroxylase (TH) after the recordings, we verified the location of electrodes within the

striatal subregions (Figure 5B). After the single lever press task, we conducted the auditory

discrimination task for the recordings. In correct trials, the delay period (0.5 ± 0.2 s) was inserted

between a lever press and a sound indicating the reward to distinguish neuronal activity related to

the two events (Figure 5C). The behavioral data in individual rats were divided into three grouped

sessions corresponding to the early, middle, and late stages, based on a sigmoid function analysis[33]

of the success rate in the task. The success rates in the grouped sessions were 52.7 ± 0.8 %, $61.6 \pm$

0.7 %, and 85.3 ± 1.7 % (mean $\pm$ s.e.m.) for the early, middle, and late stages, respectively, and the

rate was significantly increased throughout the three stages (Figure S3A). However, the response

time, omission rate, and response bias were similar among the stages (Figures S3B-S3D). Thus,

these stages corresponded to the timing window defined in the acquisition phase of the transient

MUS inhibition experiment of neuronal activity.

We identified 1,062 and 423 well-isolated neurons from the aDLS and pVLS, respectively,

across the three stages (Figure S3E) and focused on the major population of striatal neurons,

putative medium spiny neurons (MSNs; Figures S3F and S3G). The total numbers of MSNs at the

early, middle, and late stages were 295, 364, and 360 neurons for the aDLS, and 87, 110, and 190

neurons for pVLS, respectively. Based on the combinatorial pattern of the tone instruction cue and

lever press in our discrimination task, we categorized the electrophysiological data into four trial

types: (1) high-frequency tone/right lever press (HR) and (2) low-frequency tone/left lever press

(LL) as correct responses; and (3) low-frequency tone/right lever press (LR) and (4) high-frequency

tone/left lever press (HL) as error responses (see Figure 1A). We identified HR or LL type neurons

showing significant changes in firing rate related to the instruction cue onset (CO), choice response

(CR), reward sound (RS), or first licking (FL) as compared to the baseline firing rate in each of two

trials (Mann-Whitney U test, $p < 0.01$). These neurons were further divided into two groups based

on increased or decreased activity (Figures S4A and S4B). In the following analyses, we focused on

the HR and LL type neurons with increased event-related activity to explore the firing patterns of

neurons underlying behavioral processes of auditory discrimination.

Subpopulations of aDLS neurons, but not of pVLS neurons, show firing activity related to the

behavioral outcome

Previous studies have reported that neurons in the dorsal striatum show the outcome-related activity

in the instrumental conditioning tasks[24,34]. We thus investigated whether aDLS and pVLS neurons

exhibit a response related to the behavioral outcome during the acquisition of auditory

discrimination. In the aDLS, the firing activity of RS-related HR type (RS-HR) neurons showed a

higher extent in the increase during a period, when the RS is presented in the correct responses, in

the HR trial as compared with the LR, LL, or HL trial throughout the three stages (Figure 5D, top

rows). The rate of these neurons also moderately increased during the same period in the LR trial

compared to the LL or HL trial across the three stages, although the extent of the increase was lower

than that in the HR trial (Figure 5D, top rows). The area under the receiver operating characteristic

(auROC) values revealed a significant difference of the HR trial from the LR trial during the RS

period, whereas the values of HL and LL trials indicated a significant difference in the opposite

direction from the LR trial during the same period (Figure 5D, bottom rows). The averaged firing

rate during the RS period in the HR or LR trial appeared to modestly increase at the middle stage,

but no significant difference existed among the stages (Figure 5E). To directly compare the number

of neurons showing event-related activity among different stages, we calculated the number of RS-

HR neurons by matching the trial number at each stage (Figure 5F). The proportion decreased

progressively from the early to middle stages, and to the late stage. In addition, the firing rate of

RS-related LL type (RS-LL) neurons exhibited a higher increase in the LL trial and a modest

increase in the HL trial while the averaged firing rate during the RS period was similar among the

three stages. Furthermore, the proportion of RS-LL neurons reduced progressively through the

stages (Figures S5A–S5C). These results indicate that the aDLS contained neurons related to the

outcome of choice, showing a greater increase in the firing activity for the correct responses (HR

and LL trials) than the error responses (LR and HL trials), and that the proportion of these

subpopulations was gradually reduced through the progress of learning. In contrast, during the RS

period at all stages, the firing rate and auROC values of RS-HR and RS-LL neurons in the pVLS

revealed no significant differences across the four kinds of trial (Figures S5D and S5E), indicating

that the firing activity of pVLS neurons did not appear to be associated with the behavioral outcome.

Subpopulations of aDLS and pVLS neurons show sustained firing activity after the reward in

different manners

The dorsal striatum is known to demonstrate a response of neuronal activity after the positive

outcome in the instrumental behaviors[35,36]. We thus examined whether aDLS and pVLS neurons

show the response after obtaining the reward during the acquisition phase of auditory discrimination.

In the aDLS, the firing rate of FL-related HR type (FL-HR) neurons indicated a sustained elevation

after the FL in the HR trial as compared with any of the other three trials across the early, middle,

and late stages (Figure 6A, top rows). The auROC values revealed a significant difference of the LR,

LL, or HL trial from the HR trial across the three stages (Figure 6A, middle rows). In addition, we

calculated the averaged firing rate and increased firing period, defined as the total number of time

bins above 3 s.d. of the baseline firing rate for 5 s after the FL in the HR trial. The averaged firing

rate was comparable between the early and middle stages, but the rate was significantly lower at the

late stage compared to either of the other two stages (Figure 6B). The increased firing period was

also comparable between the early and middle stages, showing a significant reduction at the late

stage (Figure 6C). The proportions of FL-HR neurons were similar between the early and late

stages, although they showed a slight decrease at the middle stage (Figure 6D). There were few

neurons with significant correlations between the firing rate and licking performance (Figure 6E).

In addition, the firing rate of the FL-related LL type (FL-LL) neurons exhibited the largest increase in the LL trial at the three stages, showing a tendency to reduce the average firing rate and increased firing period after the FL across the stages (Figures S6A-S6C). The proportion of FL-LL neurons during the learning processes displayed similar changes as observed in the FL-HR neurons (Figure S6D). These results indicate that the aDLS contained neurons showing long-lasting, increased activity after the reward in the specific correct combination of the stimulus and response (HR or LL trial), and that the extent of firing activity and length of firing period in the response appeared to be gradually reduced with no correlation between the proportion of neurons and the progress of the learning stages.

In the pVLS, FL-HR neurons showed a sustained elevation in the firing rate after receiving a reward in the HR trial compared to that in the LR or HL trial throughout the three stages (Figure 6F, top rows). A similar elevation in the firing rate was also observed in the LL trial throughout the stages (Figure 6F, top rows). The auROC values revealed a significant difference of the LR or HL trial, but not the LL trial, from the HR trial throughout the stages (Figure 6F, middle rows). The average firing rate did not show significant changes among the three stages (Figure 6G), and the increased firing period was also similar across the stages (Figure 6H). The proportion of FL-HR neurons increased at the late stage compared to the early or middle stage (Figure 6I). The firing rate showed almost no significant correlation to the licking performance (Figure 6J). In addition, FL-LL neurons exhibited similar firing patterns and changes in the proportion of neurons across the stages (Figures S6F–S6J). Therefore, the pVLS included neurons representing the long-lasting increased activity after the reward in the two kinds of correct responses (HR and LL trials) with the increased number of these neurons at the late stage of learning.

Neurons in the pVLS, but not in the aDLS, exhibit firing patterns related to the beginning and

ending of the behavior

The dorsal striatum plays a key role in the acquisition and execution of behaviors based on the stimulus-response association, and the association is accompanied by a transient activation of neuronal firing immediately after the stimulus presentation or action in trials, which is considered to represent the beginning and ending of a learned behavior[37-39]. We first examined whether the firing activity of CO-related HR type (CO-HR) or CO-related LL type (CO-LL) neurons corresponds to the beginning of the behavior in the auditory discrimination task. In the aDLS, the firing rate of CO-HR or CO-LL neurons showed a significant difference at some time points in each trial compared to those in any of the other three trials, with a significant difference of auROC values between the trials (Figures S7A and S7C). However, the transiently increased activity did not emerge at the time points corresponding to the beginning of the behavior. In contrast, CO-HR neurons in the pVLS showed a transient increase in the firing rate immediately after the CO, and the activity appeared at the middle and late stages (Figure 7A, top). The transient activation was observed not only in the HR trial but also in the other three trials, being consistent with the results of a previous report[40]. The increased firing rate showed a slight but notable difference between the HR and LL trials with a significant difference in the auROC values between the two trials (Figure 7A, low), suggesting that different combinations between the stimulus and response may affect the level of firing activity. Differences in the averaged firing rate between the two kinds of trial was observed at the middle and late stages (Figure 7B). The proportion of CO-HR neurons increased at the middle and late stages compared to the early stage (Figure 7C). In addition, the firing rate of CO-LL neurons exhibited patterns similar to those observed in the CO-HR neurons (Figure S7E). The data indicate that subpopulations of pVLS neurons show firing activity related to the beginning of the behavior, with a rising number of neurons throughout discrimination learning.

Next, we examined the firing activity of CR-related HR type (CR-HR) or CR-related LL type (CR-LL) neurons, corresponding to the ending of a behavior in auditory discrimination. In the aDLS, the firing rate of these neurons did not exhibit the transient increase related to the ending of

the behavior, instead, there was a modest increase before the lever press (Figures S7B and S7D). In

contrast, CR-HR neurons in the pVLS were transiently activated immediately after the lever press

in the four kinds of trial at the middle and late stages, and the rate appeared to gradually decline

toward the RS presentation (Figure 7D). The averaged firing rate was similar between the HR and

LL trials though the stages (Figure 7E). The proportion of the CR-HR neurons increased at the

middle and late stages compared to the early stage (Figure 7F). In addition, the rate of CR-LL

neurons displayed similar firing patterns to those observed in the CR-HR neurons (Figure S7F).

Thus, subpopulations of pVLS neurons also represent the activity related to the ending of the

behavior in auditory discrimination, increasing the neuron number throughout the learning stages in

a similar pattern to neurons relevant to the beginning of the behavior in the task.

Discussion

Our findings demonstrate that the aDLS and pVLS are required for the acquisition of auditory

discrimination at different stages of learning, and that the DMS is not necessary for the acquisition

of discrimination, although it appears to contribute to the execution of actions, in particular to the

regulation of response time. These results not only support some recent studies reporting no

requirement of the DMS in the procedural learning[20,41,42], but also reveal important roles of two

different subregions in the lateral striatum in the external cue-dependent decision-making. We

propose that aDLS and pVLS neurons act to integrate the new learning of auditory discrimination in

spatiotemporally and functionally distinct manners, showing a remarkable difference from the prior

learning model that proposes the functional dominance changes from the DMS to DLS subregions

during procedural learning[12-16].

Our [18]F-FDG-PET imaging study demonstrated that the brain activity in the aDLS reaches

a peak at the middle stage of the acquisition phase of auditory discrimination (Figure 2I). The

results of excitotoxic lesion experiment of striatal subregions suggest that the aDLS functions to

enhance the acquisition of discrimination through the promotion of behavior based on the stimulus-

response association and the suppression of behavior based on the response-outcome association

(Figure 3J). Transient inhibition experiment of neuronal activity confirmed this aDLS function at

the middle stage of the learning, although the effect of drug inhibition was moderate for the

promotion of the stimulus-response behavior (Figures 4E and 4F). In addition, we found

subpopulations of aDLS neurons (RS-related neurons) which represent the behavioral outcome and

differentiate the correct (HR or LL) and incorrect (LR or HL) choices (Figures 5D and S5A). The

distinct responses of these neurons may explain the mechanism by which the aDLS regulates the

stimulus-response and response-outcome associations through distinct neural circuits leading to a

differential control of the two associations. Alternatively, the responses mainly promote the

behavior based on the stimulus-response association, thereby secondarily inhibiting the response-

outcome association. Indeed, some previous studies have suggested that an action based on either

the stimulus-response association or response-outcome association is generated as a result of

competition between different striatal subregions[43]. We also found other subpopulations of aDLS

neurons (FL-related neurons) showing long-lasting increased activity after obtaining reward in

response to a specific combination between the stimulus and response (HR or LL trial) (Figures 6A

and S6A). This activity may function as the feedback signal to promote the stimulus and response

association, resulting in the enhancement of discrimination learning. The proportion of neurons

related to the behavioral outcome showed a gradual reduction through the progress of auditory

discrimination (Figures 5E and S5C), and the extent of the firing activity and the length of the firing

period of neurons showing sustained activation after reward were also decreased at the late stage of

learning (Figures 6B and 6C). The learning-dependent changes in the properties of these

subpopulations in the aDLS may explain the mechanism underlying the earlier acquisition process

of the auditory discrimination.

The [18]F-FDG-PET imaging indicated that the pVLS activity was gradually elevated

through the acquisition phase of auditory discrimination, showing the maximal value at the late

stage (Figure 2J). The lesion experiment suggested that the pVLS drives the formation of

discrimination learning via the selective progression of the stimulus-response association (Figure

3K), and the transient inhibition experiment ascertained the pVLS function at the late stage of

learning (Figures 4G and 4H). Previous studies have shown that the stimulus-response association

is encoded in the DLS or VLS as the transient neuronal activity representing the beginning and

ending of a learned behavior[37-39,44]. A recent study reported that the VLS is causally related to

learned performance based on the auditory stimulus, while the DLS shows a weaker relationship to

the behavior[22]. In the present study, we found subpopulations of pVLS neurons that represent the

beginning and ending of a behavior (CO- and CR-related neurons, respectively) (Figures 7A, 7D,

S7E, and S7F), showing an increased number of neurons along with the progress of discrimination

learning (Figures 7C and 7F). These data suggest that pVLS neurons contribute to the formation of

the stimulus-response association through the learning processes. In addition, we found another

subpopulation of pVLS neurons (FL-related neurons) that are continuously activated after the

reward outcome in response to two kinds of correct choices (HR and LL trials) (Figures 6F and

S6F), and the proportion of neurons was increased at the late stage (Figures 6I and S6I). This

activity may act as a feedback signal to consolidate the stimulus and response association, leading

to the maintenance of associative memory at the late stage during the acquisition phase of learning.

In the present study, we identified neurons with four kinds of event-related activity in the

aDLS and pVLS, and classified them into the HR and LL type neurons in which their firing activity

showed a significant increase in the HR and LL trials, respectively, as compared to the baseline

(Figures 5–7 and S5-S7). Parts of these two types of neurons with the respective event-related

activity overlapped, and the numbers of overlapping neurons varied between striatal subregions

(Figures S8A and S8B). In the pVLS, the overlapping neurons for the same event-related activities

were 35.1%, 45.2%, and 67.0% in the CO, CR, and FL, respectively. In contrast, the aDLS had

lower overlaps of 12.6% and 22.7% in the RS and FL, respectively. Thus, distinct neurons in the

aDLS tend to have properties of a single trial type while the same neurons in the pVLS seem to

possess properties of both types. One possible explanation for the diversity in the proportion of

overlapping neurons between the two subregions may be the difference of input patterns into striatal

neurons or plastic changes in functional connectivity during learning processes. The detailed

mechanism by which aDLS and pVLS neurons react to the events in a different manner should be

investigated in the future.

Anatomically, striatal subregions are known to form heterogeneous cortico-basal ganglia

loops in mice[2,45,46]. We examined the neural circuits linked to the aDLS and pVLS in rats by using

bidirectional tracers. The results showed that two striatal subregions receive innervations from

different subregions in the cerebral cortex, intralaminar thalamus, and substantia nigra pars

compacta, and send projections to different subregions in the basal ganglia nuclei, including the

globus pallidus, entopeduncular nucleus, and substantia nigra pars reticulata (Figure S9),

confirming that the aDLS and pVLS form the heterogenous cortico-basal ganglia loops in the rat

brain. These data suggest that the aDLS and pVLS mediate the acquisition of auditory

discrimination via the parallel loops, and cooperate to achieve neural processing related to the dual

regulation of the stimulus-response and response-outcome associations. Further studies are needed

to elucidate the mechanism by which the two kinds of loops are processed separately and integrate

the new learning of discrimination.

Materials and Methods

Animals

Animal care and handling procedures were carried out in accordance with the guidelines established

by RIKEN Center for Biosystems Dynamics Research, Fukushima Medical University, and Osaka

City University. All procedures were approved by their Institutional Animal Care and Use

Committees. All efforts were made to minimize the number of animals used and their suffering

throughout the course of the experiments. Male Long-Evans rats (8–13 weeks old, Institute for

Animal Reproduction (Ibaraki, Japan)) were used for the present study. The rats were maintained at

22 ± 2 °C and 60% humidity in a 12-h light/12-h dark cycle, and food and water were continuously

available unless stated otherwise.

Surgery

The rats underwent a stereotaxic surgery under isoflurane anesthesia (1–5%). For excitotoxic lesion

of striatal neurons, IBO was dissolved in PBS (0.8 mg/mL, Wako Pure Chemical), and IBO solution

(0.3 μL per site) was injected bilaterally with the coordinates (mm) from bregma and dura for the

aDLS (AP +1.6, ML ±3.3, DV -3.9), pVLS (AP -1.6, ML ±4.6, and DV -5.6), or DMS (AP -1.6, ML

±1.4, and DV -4.8) according to the atlas of rat brain[47]. Injection was performed through a glass

pipette (diameter: 60 μm) at a constant velocity of 0.1 mL/min with the microinfusion pump

(EICOM). For transient inactivation of striatal neurons, two guide cannulae (length: 10 mm, outer

diameter: 26 gauge) were bilaterally placed on the skull and fixed using four screws and dental

cement. The chip of the cannula was placed at 1 mm above the target coordinates. A solution

containing MUS (Sigma-Aldrich) in saline (0.1 μg/μL, 0.25 μL per site) was injected into the target

coordinates through the internal cannula (outer diameter: 30 gauge) at a constant velocity of 0.25

μL/min. For anterograde and retrograde tracing, a solution containing cholera toxin subunit B (CTb;

1.0 mg/mL) conjugated to Alexa 488 or Alexa 555 (Invitrogen) as a bidirectional tracer was injected

into the aDLS and pVLS (0.3 μL/site), respectively. The coordinates for the injection into striatal

subregions were the same as described above. For the electrophysiological experiments, surgery

was performed as described previously[48]. A 64-channel silicon probe (Buzsaki64spl; NeuroNexus)

attaching to a 3D-printed microdrive with a movable screw was chronically implanted with the

coordinates (mm) from the bregma and dura into the left aDLS (AP +1.6, ML -3.3, and DV -2.5)

and pVLS (AP -1.6, ML -4.6, and DV -3.6). A faraday cage composed of copper mesh was secured

on the skull with dental cement to reduce electrical noise and avoid damage to the implants. The

rats were allowed to recover for 1 week after the surgeries.

Behavioral analysis

Behavioral tasks were conducted in operant conditioning chambers ($30.5 \times 24.1 \times 29.2$ cm; ENV-

007CT; MED Associates) equipped with two retractable levers on both sides of a reward port in the

front panel (ENV-203M-45, ENV-652AM, and ENV-201A; MED Associates). A multi-tone

generator (ENV-223; MED Associates) and room light were mounted on the top of the center and

corner panels on the rear side, respectively. For the electrophysical experiment, a 3D printed reward

port was used to detect the licking and to prevent contact with the head attachment and front panel.

The behavioral experiment in intact rats was performed using grain pellets (F0021, Bioserv) as a

food reward. To prevent spillover of radioactivity into the temporal cortex as a consequence of

excessive ^{18}F-FDG accumulation into the muscles by their activity to eat the pellets[49,50], the PET

imaging experiments were carried out using 0.02% saccharine solution (B0131, Tokyo Chemical

Industry) as a liquid reward. The electrophysiological experiment was also carried out using the

saccharine solution. The pharmacological experiment was performed using the pellets.

The rats were subjected to dietary restrictions of 12 g/day (CE-2, CLEA Japan) and water

access was limited to 6 to 9 h in accordance with the reward used in the experiments for 3 days

before the beginning of the training schedule, and the body weights of the rats were maintained at

~85% of their normal weights. The training schedule consisted of four continuous steps. In the first

step (magazine training), the rats were habituated to the operant conditioning chambers, and the

reward was presented with the click every 20 s. Each daily session was started by illumination of

the room light, and continued for 30 min. The sessions were conducted for 3 consecutive days. In

the second step (shaping), only one lever on the left or right side was inserted. When the rats

pressed the lever, they earned a reward accompanied by a click. Each daily session was started by

illumination of the light, and lasted 30 min or until the criteria, in which the rats received 90

rewards. When the number of lever presses on either side reached the criteria, the lever was inserted

in the opposite side on the next day. When the rats achieved the criteria on both sides for 2 or 3 days,

they moved to the following step. In the third step (single lever press task), each trial was started by

illumination of the light and insertion of one lever on either the left or right side in a pseudorandom

manner with equal frequencies on both sides for every set of four trials. When the rats pressed the

lever, the reward was presented with the sound, and the lever was immediately retracted. The room

light turned off 4 s after the lever press. When they did not press the lever within 10 s after the start,

the trial was terminated when the light turned off. The trial was repeated every 20 s. Each daily

session continued for 30 min, and the sessions were conducted for 4 consecutive days. The number

of lever presses in each session was counted. In the final step (two-alternative auditory

discrimination task), each trial was started by the presentation of a high (10 kHz) or low (2 kHz)

frequency tone as an instruction cue in a pseudorandom manner. After three seconds, the room light

was illuminated, and two retractable levers were inserted at the same time. The rats were required to

press the left or right levers in response to the low or high tone cues, respectively. When the rats

pressed the correct lever, the reward was presented with the sound, and the levers were immediately

retracted. Correct response allowed the rats to access the reward: the retractable spout delivering

water was inserted for 1 s for the PET imaging experiment; a pellet was delivered by a food

dispenser for the pharmacological experiment; and three drops of water were delivered through a

solenoid valve for the electrophysiological experiment. The room light turned off 4 s after the

correct response. On the other hand, error response resulted in the retraction of levers and the

turning off of the room light. No response within 10 s after the lever insertion was recorded as an

omission response. The success rate in each session was calculated by dividing the number of

correct response trials with the number of lever press trials minus the number of omission trials.

The trial was repeated every 20 s. Each daily session continued for 60 min, and the sessions were

conducted for 14 or 24 days according to the experiments.

To assess the behavioral strategies in the auditory discrimination task, a trial-by-trial

analysis was conducted; the WSW strategy was defined as a correct response, in which, after a

correct response during the previous trial, the rats press the opposite lever during the current trial in

response to the shift of the tone instruction cue; the LSL strategy was defined as an error response,

in which, after an error response during the previous trial, the rats press the opposite lever during

the current trial despite the shift of the instruction cue. The proportion of both strategies was

calculated by dividing the number of each strategy with the number of lever press trials minus the

number of omission trials.

For the transient inactivation experiment, MUS injections were bilaterally performed 20

min before start of the session. For habituation, all rats were injected with saline bilaterally into the

corresponding striatal subregions on the last day (Day 4) of the single lever press task. During the

auditory discrimination task, the first injection was conducted on Day 2 of the task. The second and

third injections were conducted the next day, after the success rate had exceeded 60% and 80%,

respectively. Each session continued for 15 min.

For the electrophysiological experiments, a delay period (0.5 ± 0.2 s) was added in the

auditory discrimination task between the lever press and reward sound to distinguish the motor- and

reward-related responses of neurons. When the rats pressed the lever, the lever was immediately

retracted. After the delay period, the reward was presented with a sound. The room light turned off

4 s after the lever press. The trial was repeated every 20 s, and each daily session continued for 60

min.

PET imaging analysis

An [18]F-FDG-PET scan was performed with a microPET Focus220 (Siemens Medical Solutions,

Knoxville) designed for high-resolution imaging of small animals (spatial resolution of 1.4 mm in

full width at half maximum at the center of the field of view) as described previously[26]. Serial [18]F-

FDG-PET scans were performed on the last day of the single lever press task and on Days 2, 6, 10,

and 24 of the auditory discrimination task. Under the awake condition, a solution of [18]F-FDG (ca.

74 MBq/0.4 mL) was intravenously injected just before the beginning of each behavioral session

through an indwelling catheter attached to the tail. After the 30-min session, the rats were returned

to their home cage, and then a 30-min static PET scan was performed under anesthesia with 2.0–

2.5% isoflurane 55-min after the [18]F-FDG injection. During the PET scan, the body temperature of

the anesthetized rats was maintained at 37°C using a small animal warmer (BWT-100A, Bio

Research Center). The acquired emission data were sorted into a single sinogram and reconstructed

by a standard 2D filtered back projection (FBP) with a ramp filter and cutoff frequency at 0.5 cycles

per pixel or by a statistical maximum a *posteriori* probability algorithm (MAP), 12 iterations with

point spread function effect.

Voxel-based analysis of PET images was conducted according to the method described

previously[26]. Briefly, individual MAP-reconstructed FDG (MAP-FDG) images were aligned to a

FDG template image of Long Evans rats using PMOD software package (version 3.2, PMOD

Technologies), and then the aligned images were transformed into the space of a T2 weighted MRI

templates for the rats (https://www.nitrc.org/projects/tpm_rat/). The transformation parameters

obtained from individual MAP-FDG images were used to match the individual FBP-reconstructed

FDG (FBP-FDG) images with the MRI templates. The voxel size of the template was magnified by

10 to a size similar to that of a human brain. All FBP-FDG images were resampled with a voxel

size of $1.2 \times 1.2 \times 1.2$ mm and spatially smoothed through an isotropic Gaussian kernel (6-mm

FWHM). Voxel-based statistical analysis was assessed by using the Statistical Parametric Mapping

software (version 8; https://www.fil.ion.ucl.ac.uk/spm/). The voxel of interest (VOI) was

determined as voxels with more than 85% of the maximum T-value within activated areas obtained

by voxel-based statistical comparison of FBP-FDG images. The average regional [18]F-FDG uptake

was calculated on the basis of the values of the VOI in FBP-FDG images.

Electrophysiological recording

Extracellular multi-unit recording was conducted at a sampling rate of 20 kHz from the striatum of

freely moving rats by using a 512-ch acquisition board (Open Ephys) via 64-ch recording

headstages (C3325; Intan Technologies). The tips of the probe were lowered by turning the screw (>

140 µm/day) until it reached the target region (the aDLS and pVLS) during the single lever press

task. The tips were sometimes lowered by the same procedure (140 to 280 µm/day) after the daily

recording session to improve cell yield.

Spike sorting was performed automatically by the Kilosort software

(https://github.com/cortex-lab/KiloSort). Clear noise clusters were removed in the first manual

sorting using the phy GUI (https://github.com/cortex-lab/phy). The remaining clusters were then re-

clustered by using the Klusta Kwik (https://klusta.readthedocs.io/en/latest/), and the clusters were

adjusted in the second manual sorting by using the phy GUI again. The sorted clusters satisfying all

the following criteria were used for further analysis: L-ratio < 0.05, isolation distance > 15, and ISI

index < 0.2[51,52]. Well-isolated single neurons (n = 1,485; aDLS, n = 1,062; pVLS, n = 423) were

classified as putative medium spiny neurons (MSNs), fast spiking interneurons (FSIs), tonically

active neurons (TANs), and unclassified interneurons (UIs). Putative MSNs and TANs were

separated from FSIs and UIs by trough-to-peak duration of mean spike waveform (> 0.6 ms)

according to previous studies, with slight modifications[53,54]. Putative FSIs were then separated from

UIs by utilizing the proportion of time associated with long inter-spike intervals. This was achieved

by summing the inter-spike intervals longer than 2 s and subsequently dividing the resulting sum by

the total recording time ($Prop_{ISIs} > 2$ s)[55]. Neurons with values of $Prop_{ISIs} > 2$ s less than 0.4 were

classified as putative FSIs, while neurons with $Prop_{ISIs} > 2$ s greater than 0.4 were classified as

putative UIs. The putative MSNs and TANs were separated by measuring the post-spike

suppression[55]. We measured the length of suppressed time during which the firing rate of a neuron

was suppressed following a spike by counting the number of 1-ms bins in its autocorrelation

function until the rate of the neuron was equal to or greater than its averaged rate over the 600-ms to

900-ms autocorrelation bins. Putative TANs were separated by post-spike suppression of less than

40 ms, and the remaining neurons were classified as putative MSNs. Because the numbers of FSIs,

TANs, and UIs were low (79 out of 1,485, 5.3%), they were excluded from further analysis.

To divide the three stages during the acquisition phase, the sigmoidal function was fitted

using the following four-parameter logistic equation:

$$f_{(x)} = \frac{1 - \gamma - (1 - \lambda)}{1 + \exp\left(-\beta(x - \alpha)\right)} + \gamma$$

Where α is the inflection point, β is a variable for slope, and λ and γ are parameters related to the

lower asymptote (γ) and upper asymptote (λ) of the learning curve. The function was used to

evaluate the day when the success rate reached 20% ($t_{20\%}$) or 90% ($t_{90\%}$) of the maximum rate

change defined by the difference between the γ and λ. Three stages were defined according to the

following criteria: early stage (up to 3 days before the middle stage); middle stage (3 days including

before and after the day of $t_{20\%}$); and late stage (3 days including the day of $t_{90\%}$ and before). One rat

learned the discrimination faster, and the success rate reached $t_{90\%}$ 2 days after the day of $t_{20\%}$. One

day before the day of $t_{90\%}$ was assigned to the late stage. For the following analysis, spike data in

each striatal subregion were separated into the early, middle, and late stages ($n = 295$, 364 and 360,

respectively, for the aDLS; and $n = 87$, 110 and 190, respectively, for the pVLS).

666   To determine the neurons showing significant changes in firing rate related to the task

events, electrophysiological data were grouped into four trial types based on the instruction cue

(high or low tone) and the choice (right or left lever); HR, high tone and right lever press; HL, high

tone and left lever press; LR, low tone and right lever press; and LL, low tone and left lever press.

Neurons with three or more trial data were analyzed. We identified the HR or LL type neurons,

detecting significant increases or decreases of the firing rate by applying the Wilcoxon rank-sum

test ($p < 0.01$) to trial-by-trial spike number between the baseline period (BL, -1,500 to -500 ms or

-1,100 to -500 ms before presentation of the instruction cue) and any of four peri-event periods (CO,

0 to +1,000 ms after the onset of the instruction cue; CR, -700 to +300 ms around the time of lever

press; RS, -300 to 300 ms around the time of sound onset instructing the reward delivery; and FL, 0

to +1,000 ms after the time of first licking after the CR). Any spikes around the events were

averaged in 50-ms bins. The event-related activity was shown as averaged firing rates aligned based

on the timing of an event for each trial. The auROC values were calculated by using the spike

counts between the HR or LR trials and any of the other three trials. Bins showing significant

differences in the auROC data were determined by applying the Wilcoxon rank-sum test ($p < 0.05$)

to the distribution of 200 ms bin-by-bin auROC values and 0.5.

To compare the number of neurons showing event-related activity among different stages,

the number of HR or LL type neurons were calculated by matching the trial number at each stage.

Twenty trials were randomly selected for a neuron, and neurons with less than 20 trials were

removed from the analysis. The neurons with event-related activity were identified as described

above. The proportion (percentage) of neurons relative to the total number of neurons was

calculated at each individual stage. The same analysis was repeated 30 times and the distribution of

the proportion of neurons with event-related activity at the stage was obtained. The cumulative

probability against the proportion was calculated using cumulative distribution function (cdfplot,

MATLAB code from Mathworks).

Licking responses were detected with an infrared photobeam sensor placed in the 3D

printed reward port, and generated timestamps for the onset of each spout contact according to the

analog voltage signal. The licking responses corresponding to four kinds of trial in the FL-related

neurons were shown as averaged rates (50-ms bins) aligned based on the timing of the CR. To

calculate the correlation coefficient between neuronal activity and licking responses, the spikes and

licking for 5 s after the FL were counted every trial for each FL-related neuron. Trials in which

neurons never responded were removed from both the spike and licking data. The correlation

coefficients between the two remaining data were calculated for each neuron (corr, MATLAB code

from Mathworks), and neurons showing significant differences were determined by applying the

Pearson correlation coefficient ($p < 0.01$).

Histology

After the behavioral experiments, the rats were anesthetized with pentobarbital (50 mg/kg body

weight) and isoflurane (3%), then transcardially perfused 4% paraformaldehyde in phosphate-buffer

(PB). The brains were post-fixed overnight at 4 □ and cryoprotect with 10, 20, and 30% sucrose in

PB. Brain tissues were cut into 30-µm sections by using a cryostat (Leica). For the lesion

experiment, cresyl violet staining or immunostaining were performed to define the lesioned areas in

the striatum. The sections were stained with a mouse monoclonal anti-NeuN IgG (1:1000 dilution;

Millipore, Cat# MAB377, RRID:AB_2298772), and then with a goat anti-mouse IgG conjugated

with Alexa Fluor 488 (1:400 dilution; Thermo Fisher Scientific, Cat# A-11029,

RRID:AB_2534088). The sections were counterstained with 0.3 µM 4',6-diamidino-2-phenylindole

(DAPI; Sigma-Aldrich). For transient inactivation experiment, sections were stained by cresyl

violet to visualize the positions of guide cannulae. For the electrophysiological experiments, small

lesions were made by applying an electrolytic current (3 µA, 10 s) through top and bottom channels

of each shank by using the stimulus isolator (A365, World Precision Instrument). Sections through

the striatum were stained with a mouse monoclonal anti-TH IgG (1:1,000 dilution; Millipore, Cat#

MAB318, RRID:AB_2201528), and then with a goat anti-mouse IgG conjugated with Alexa Fluor

488 (1:400 dilution; Thermo Fisher Scientific, Cat# A-11029, RRID:AB_2534088). The recording

sites were verified by the staining with DAPI and red fluorescent Nissl stain solution (1:200

dilution; Thermo Fisher Scientific, Cat# N21482, RRID:AB_2620170). Fluorescence and

immunostained signals were visualized under fluorescence microscopes (BZ-X800; Keyence) and

confocal microscope (A1R; Nikon).

Statistical analysis

Statistical analysis was performed using MATLAB (MathWorks) and SPSS Statistics (Ver. 27,

IBM). Comparisons of data between two groups were performed using the Student's t-test (paired),

the Wilcoxon signed rank test (paired), and the Wilcoxon rank-sum test (unpaired). Comparisons of

data among three groups were constructed using a one-way repeated measures ANOVA, two-way

repeated measures ANOVA, or the Kruskal-Wallis test (without assuming normal distribution and

homogeneity of variances). All statistical tests were two-sided. The Bonferroni post-hoc test and

Mann–Whitney U-test with Bonferroni adjustment were carried out as needed to compare means.

For boxplots, the data points below the lower (25th percentile) -1.5 interquartile range (IQR) and

above the upper (75th percentile) quartiles +1.5 IQR were regarded as outliers, and they are not

included in the boxplots.

Acknowledgements

We thank Drs. Keiji Ota, Hikaru Yokoyama, and Sho Aoki for instructing the behavioral analysis

and Dr. Hideyuki Matsumoto for technical advice and discussion. We thank the Research Support

Platform, Osaka Metropolitan University Graduate School of Medicine for microscopic imaging.

This work was supported by grants-in-aid for JSPS Fellows (19J01997) and Scientific Research (C)

(21K11556) from the Japan Society for the Promotion of Science (S.S.); a grant-in-aid for Scientific

Research on Transformative Research Areas (A) Adaptive Circuit Census (21H05244) from the

Ministry of Education, Science, Sports, and Culture of Japan (K.K.); the Grant for Casio Science

Promotion Foundation, Takeda Science Foundation; and the Nakatomi Foundation (S.S.).

Author contributions

S.S., Y.C., and K.K. designed the study. S.S. performed the small-animal neuroimaging, multi-unit

recordings, tracing studies, and pharmacological experiments with technical support from all

authors; T.O., D.H., Y.W., H.O., K.H., and Y.C. assisted with the neuroimaging studies. K.N.

performed the tracing study, and N.S. assisted with the immunostaining. H.M., T.K., and K.M.

assisted with the multi-unit extracellular recordings. S.S. and K.K. wrote the manuscript with

assistance from all coauthors.

Declaration of interest

The authors declare no competing interest.

Data availability

All source data underlying the graphs and charts are available in Supplementary Data. All other data

are available upon request. The datasets generated and analyzed in this study are deposited on

Mendeley data (DOI: 10.17632/ghzr838bdk.1).

References

1. Alexander G.E., Crutcher M.D (1990) Functional architecture of basal ganglia circuits: neural substrates of parallel processing *Trends Neurosci* **13**:266-271. DOI: https://doi.org/10.1016/0166-2236(90)90107-l

2. Foster N.N., *et al.* (2021) The mouse cortico-basal ganglia-thalamic network *Nature* **598**:188-194. DOI: https://doi.org/10.1038/s41586-021-03993-3

3. Kim H.F., Hikosaka O (2015) Parallel basal ganglia circuits for voluntary and automatic behaviour to reach rewards *Brain* **138**:1776-1800. DOI: https://doi.org/10.1093/brain/awv134

4. Yin H.H., Knowlton B.J (2006) The role of the basal ganglia in habit formation *Nat Rev Neurosci* **7**:464-476. DOI: https://doi.org/10.1038/nrn1919

5. Yin H.H., *et al.* (2009) Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill *Nat Neurosci* **12**:333-341. DOI: https://doi.org/10.1038/nn.2261

6. Kupferschmidt D.A., Juczewski K., Cui G., Johnson K.A., Lovinger D.M (2017) Parallel, but dissociable, processing in discrete corticostriatal inputs encodes skill learning *Neuron* **96**:476-489 e5. DOI: https://doi.org/10.1016/j.neuron.2017.09.040

7. Thorn C.A., Atallah H., Howe M., Graybiel A.M (2010) Differential dynamics of activity changes in dorsolateral and dorsomedial striatal loops during learning *Neuron* **66**:781-795. DOI: https://doi.org/10.1016/j.neuron.2010.04.036

8. Doyon J., *et al.* (2009) Contributions of the basal ganglia and functionally related brain structures to motor learning *Behav Brain Res* **199**:61-75. DOI: https://doi.org/10.1016/j.bbr.2008.11.012

9. Lehéricy S., *et al.* (2005) Distinct basal ganglia territories are engaged in early and advanced motor sequence learning *Proc Natl Acad Sci U S A* **102**:12566-12571. DOI: https://doi.org/10.1073/pnas.0502762102

10. Miyachi S., Hikosaka O. Lu X (2002) Differential activation of monkey striatal neurons in the early and late stages of procedural learning *Exp Brain Res* **146**:122-126. DOI:

789 https://doi.org/10.1007/s00221-002-1213-7

790 11. Miyachi S., Hikosaka O., Miyashita K., Kárádi Z., Rand M.K (1997) Differential roles of

791   monkey striatum in learning of sequential hand movement *Exp Brain Res* **115**:1-5. DOI:

792   https://doi.org/10.1007/pl00005669

793 12. Dickinson A (1985) Actions and habits: the development of behavioural autonomy *Phil Trans R*

794   *Soc Lond B* **308**:67-78. DOI: https://doi.org/10.1098/rstb.1985.0010

795 13. O'Hare J.K.*, et al.* (2016) Pathway-specific striatal substrates for habitual behavior *Neuron*

796   **89**:472-479. DOI: https://doi.org/10.1016/j.neuron.2015.12.032

797 14. Yin H.H., Knowlton B.J., Balleine B.W (2004) Lesions of dorsolateral striatum preserve

798   outcome expectancy but disrupt habit formation in instrumental learning. *Eur J Neurosci*

799   **19**:181-189. DOI: https://doi.org/10.1111/j.1460-9568.2004.03095.x

800 15. Yin H.H., Ostlund S.B., Knowlton B.J., Balleine B.W (2005) The role of the dorsomedial

801   striatum in instrumental conditioning *Eur J Neurosci* **22**:513-523. DOI:

802   https://doi.org/10.1111/j.1460-9568.2005.04218.x

803 16. Redgrave P.*, et al.* (2010) Goal-directed and habitual control in the basal ganglia: implications

804   for Parkinson's disease *Nat Rev Neurosci* **11**:760-772. DOI: https://doi.org/10.1038/nrn2915

805 17. Toni I., Passingham R.E (1999) Prefrontal-basal ganglia pathways are involved in the learning

806   of arbitrary visuomotor associations: a PET study *Exp Brain Res* **127**:19-32. DOI:

807   https://doi.org/10.1007/s002210050770

808 18. Seger C.A., Cincotta C.M (2005) The roles of the caudate nucleus in human classification

809   learning *J Neurosci* **25**:2941-2951. DOI: https://doi.org/10.1523/JNEUROSCI.3401-04.2005

810 19. Kim H.F., Hikosaka O (2013) Distinct basal ganglia circuits controlling behaviors guided by

811   flexible and stable values *Neuron* **79**:1001-1010. DOI:

812   https://doi.org/10.1016/j.neuron.2013.06.044

813 20. Featherstone R.E., McDonald R.J (2004) Dorsal striatum and stimulus-response learning:

814   lesions of the dorsolateral, but not dorsomedial, striatum impair acquisition of a simple

discrimination task *Behav Brain Res* **150**:15-23. DOI: https://doi.org/10.1016/S0166-

4328(03)00218-3

21. Featherstone R.E., McDonald R.J (2005) Lesions of the dorsolateral striatum impair the

acquisition of a simplified stimulus-response dependent conditional discrimination task

Neuroscience **136**, 387-395. DOI: https://doi.org/10.1016/j.neuroscience.2005.08.021

22. Lee J., Wang W., Sabatini B.L (2020) Anatomically segregated basal ganglia pathways allow

parallel behavioral modulation *Nat Neurosci* **23**:1388-1398. DOI:

https://doi.org/10.1038/s41593-020-00712-5

23. Xiong Q., Znamenskiy P., Zador A.M (2015) Selective corticostriatal plasticity during

acquisition of an auditory discrimination task *Nature* **521**:348-351. DOI:

https://doi.org/10.1038/nature14225

24. Guo L., Weems J.T., Walker W.I., Levichev A., Jaramillo S (2019) Choice-selective neurons in

the auditory cortex and in its striatal target encode reward expectation *J Neurosci* **39**:3687-3697.

DOI: https://doi.org/10.1523/JNEUROSCI.2585-18.2019

25. Nishizawa K., *et al.*(2012) Striatal indirect pathway contributes to selection accuracy of learned

motor actions *J Neurosci* **32**:13421-13432. DOI: https://doi.org/10.1523/JNEUROSCI.1969-

12.2012

26. Cui Y., *et al.* (2015) A voxel-based analysis of brain activity in high-order trigeminal pathway

in the rat induced by cortical spreading depression *Neuroimage* **108**:17-22. DOI:

https://doi.org/10.1016/j.neuroimage.2014.12.047

27. Kornblum H.I., *et al.* (2000) In vivo imaging of neuronal activation and plasticity in the rat

brain by high resolution positron emission tomography (microPET) *Nat Biotechnol* **18**:655-660.

DOI: https://doi.org/10.1038/76509

28. Zimmer E.R.*, et al.* (2017) [18 F] FDG PET signal is driven by astroglial glutamate transport

Nat Neurosci **20**:393-395. DOI: https://doi.org/10.1038/nn.4492

29. Featherstone R.E., McDonald R.J (2004) Dorsal striatum and stimulus-response learning:

lesions of the dorsolateral, but not dorsomedial, striatum impair acquisition of a stimulus-

response-based instrumental discrimination task, while sparing conditioned place preference

learning *Neuroscience* **124**:23-31. DOI: https://doi.org/10.1016/j.neuroscience.2003.10.038

30. Featherstone R.E., McDonald R.J (2005) Lesions of the dorsolateral or dorsomedial striatum

impair performance of a previously acquired simple discrimination task *Neurobiol Learn Mem*

84:159-167. DOI: https://doi.org/10.1016/j.nlm.2005.08.003

31. Bergstrom H.C.*, et al.* (2018) Dorsolateral striatum engagement interferes with early

discrimination learning *Cell Rep* **23**:2264-2272. DOI:

https://doi.org/10.1016/j.celrep.2018.04.081

32. Packard M.G., Hirsh R., White N.M (1989) Differential effects of fornix and caudate nucleus

lesions on two radial maze tasks: evidence for multiple memory systems *J Neurosci* **9**:1465-

1472. DOI: https://doi.org/10.1523/JNEUROSCI.09-05-01465.1989

33. Bathellier B., Tee S.P., Hrovat C., Rumpel S (2013) A multiplicative reinforcement learning

model capturing learning dynamics and interindividual variability in mice *Proc Natl Acad Sci*

U S A **110**:19950-19955. DOI: https://doi.org/10.1073/pnas.1312125110

34. Nonomura S.*, et al.* (2018) Monitoring and updating of action selection for goal-directed

behavior through the striatal direct and indirect pathways *Neuron* **99**:1302-1314 e5. DOI:

https://doi.org/10.1016/j.neuron.2018.08.002

35. Williams Z.M., Eskandar E.N (2006) Selective enhancement of associative learning by

microstimulation of the anterior caudate *Nat Neurosci* **9**:562-568. DOI:

https://doi.org/10.1038/nn1662

36. Smith K.S., Graybiel A.M (2016) Habit formation coincides with shifts in reinforcement

representations in the sensorimotor striatum *J Neurophysiol* **115:**1487-1498. DOI:

https://doi.org/10.1152/jn.00925.2015

37. Jog M.S., Kubota Y., Connolly C.I., Hillegaart V., Graybiel A.M (1999) Building neural

representations of habits *Science* **286**:1745-1749. DOI:

https://doi.org/10.1126/science.286.5445.1745

38. Barnes T.D., Kubota Y., Hu D., Jin D.Z., Graybiel A.M (2005) Activity of striatal neurons

reflects dynamic encoding and recoding of procedural memories *Nature* **437**:1158-1161. DOI:

https://doi.org/10.1038/nature04053

39. Jin X., Costa R.M (2010) Start/stop signals emerge in nigrostriatal circuits during sequence

learning *Nature* **466**:457-462. DOI: https://doi.org/10.1038/nature09263

40. Barnes T.D.*, et al.* (2011) Advance cueing produces enhanced action-boundary patterns of spike

activity in the sensorimotor striatum *J Neurophysiol* **105**:1861-1878. DOI:

https://doi.org/10.1152/jn.00871.2010

41. Smith A.C.W.*, et al.* (2021) Opposing roles for striatonigral and striatopallidal neurons in

dorsolateral striatum in consolidating new instrumental actions *Nat Commun* **12**:5121. DOI:

https://doi.org/10.1038/s41467-021-25460-3

42. Wolff S.B.E., Ko R., Ölveczky B.P (2022) Distinct roles for motor cortical and thalamic inputs

to striatum during motor skill learning and execution *Sci Adv* **8:**eabk0231. DOI:

https://doi.org/10.1126/sciadv.abk0231

43. Gremel C.M.*, et al.* (2016) Endocannabinoid modulation of orbitostriatal circuits gates habit

formation *Neuron* **90**:1312-1324. DOI: https://doi.org/10.1016/j.neuron.2016.04.043

44. Chen Z.*, et al.* (2021) Direct and indirect pathway neurons in ventrolateral striatum

differentially regulate licking movement and nigral responses *Cell Rep* **37**:109847. DOI:

https://doi.org/10.1016/j.celrep.2021.109847

45. Hunnicutt B.J.*, et al.* (2016) A comprehensive excitatory input map of the striatum reveals

novel functional organization *eLife* **5**:e19103. DOI: https://doi.org/10.7554/eLife.19103

46. Hintiryan H.*, et al.* (2016) The mouse cortico-striatal projectome *Nat Neurosci* **19**:1100-1114.

DOI: https://doi.org/10.1038/nn.4332

47. Paxinos G., Watson C (1998) The rat brain in stereotaxic coordinates IV Ed (Academic Press).

48. Kitanishi T., Umaba R., Mizuseki K (2021) Robust information routing by dorsal subiculum

neurons *Sci Adv* **7**:eabf1913. DOI: https://doi.org/10.1126/sciadv.abf1913

49. Endepols H.*, et al.* (2010) Effort-based decision making in the rat: an [18F]fluorodeoxyglucose

micro positron emission tomography study *J Neurosci* **30**:9708-9714. DOI:

https://doi.org/10.1523/JNEUROSCI.1202-10.2010

50. Marx C.*, et al.* (2012) Conflict processing in the rat brain: behavioral analysis and functional

μPET imaging using [F]fluorodeoxyglucose *Front Behav Neurosci* **6**:4. DOI:

https://doi.org/10.3389/fnbeh.2012.00004

51. Fee M.S., Mitra P.P., Kleinfeld D (1996) Automatic sorting of multiple unit neuronal signals in

the presence of anisotropic and non-Gaussian variability *J Neurosci Methods* **69**:175-188. DOI:

https://doi.org/10.1016/S0165-0270(96)00050-7

52. Schmitzer-Torbert N., Jackson J., Henze D., Harris K., Redish A.D (2005) Quantitative

measures of cluster quality for use in extracellular recordings *Neuroscience* **131**:1-11. DOI:

https://doi.org/10.1016/j.neuroscience.2004.09.066

53. Bakhurin K.I., Mac V., Golshani P., Masmanidis S.C (2016) Temporal correlations among

functionally specialized striatal neural ensembles in reward-conditioned mice *J Neurophysiol*

115:1521-1532. DOI: https://doi.org/10.1152/jn.01037.2015

54. Kim N.*, et al.* (2019) A striatal interneuron circuit for continuous target pursuit *Nat Commun*

10: 2715. DOI: https://doi.org/10.1038/s41467-019-10716-w

55. Peters A.J., Fabre J.M.J., Steinmetz N.A., Harris K.D., Carandini M (2021) Striatal activity

topographically reflects cortical activity *Nature* **591**:420-425. DOI:

https://doi.org/10.1038/s41586-020-03166-8

Figure 1. Behavioral performance of two-alternative auditory discrimination learning.

(A) Schematic diagram of the auditory discrimination task. Each trial started by presentation of a

tone instruction cue with the high (10 kHz) or low (2 kHz) frequency. Three seconds later, the room

light was illuminated, and two retractable levers were inserted at the same time. The rats were

required to press the right and left levers in response to the high and low tones, respectively. (B)

Learning curve of the auditory discrimination in intact rats (n = 14). Inset indicates the cumulative

curve of number achieved the success rate of more than 90%. (C) Success rate at Days 2, 6, 10, 13,

and 24 (one-way repeated ANOVA, F[1.897,24.659] = 106.706, p = 1.1 × 10^{-12}, *post hoc*

Bonferroni test, Day 2 vs. Day 6, p = 0.001, Day 6 vs. Day 10, p = 3.6 × 10^{-4}, Day 10 vs. Day 13, p

= 0.040, and Day 13 vs. Day 24, p = 0.080). (D) Response time (one-way repeated ANOVA,

F[2.203,28.642] = 2.386, p = 0.105). (E) Omission rate (F[4,52] = 0.699, p = 0.596). (F) Response

bias (one-way repeated ANOVA, F[1.129,14.674] = 0.250, p = 0.653). Data are indicated as the

mean ± s.e.m., and individual data are overlaid. *p < 0.05, **p < 0.01, and ***p < 0.001.

Figure 2. Dissociable brain activity patterns during auditory discrimination learning among

three striatal subregions. (A) Schematic illustration of an awake rat received intravenously [18]F-

FDG injection through an indwelling catheter attached to the tail. The rat was conducted for the

behavioral experiment and then used for microPET imaging. (B) Schedule for the behavioral

training and [18]F-FDG-PET scan. After [18]F-FDG injection, rats (n = 14 rats) were subjected to the

behavioral experiment (30 min) and returned to the home cage (15 min). Ten min before the PET

scan started, the rats were anesthetized and then the scan was started (30 min). (C and D) Learning

curves of the single lever press task (C) and auditory discrimination task (D). Arrowheads indicate

the PET scan days. (E) Behavioral performance on the scan days during the auditory discrimination.

Success rate (one-way repeated ANOVA, $F[3,39] = 125.012$, $p = 4.8 \times 10^{-20}$, *post hoc* Bonferroni

test; Day 2 vs. Day 6, $p = 5.0 \times 10^{-7}$; Day 6 vs. Day 10, $p = 0.001$; Day 10 vs. Day 24, $p = 0.003$),

response time (one-way repeated ANOVA, $F[3,39] = 0.156$, $p = 0.926$), omission rate (one-way

repeated ANOVA, $F[3,39] = 1.559$, $p = 0.215$), and response bias (one-way repeated ANOVA,

$F[1.779,23.130] = 17.734$, $p = 3.6 \times 10^{-5}$) are shown. (F) Representative images of coronal sections

that compared the brain activity in the single lever press task with the activity on Day 2, 6, 10, or 24

in the discrimination task. Left panels show schematic illustrations of striatal subregions. (G)

Horizontal (top) and coronal (bottom) images of striatal activation areas shown on Day 10 vs. single

lever in (F). (H) Representative images of coronal section that compared the brain activity on Day 2

with that on Day 6, 10, or 24 in the discrimination task. Color bars indicate the T-values, and a

value of 3.8 was used as the threshold corresponding to the uncorrected threshold ($p < 0.001$). (I)

Schematic pictures showing voxel of interests for the aDLS, pVLS, and DMS (green, left

hemisphere; and purple, right hemisphere). (J-L) Regional [18]F-FDG uptakes in the aDLS (J, one-

way repeated ANOVA, left aDLS, $F[4,52] = 10.322$, $p = 3.0 \times 10^{-6}$, *post hoc* Bonferroni test; single

lever vs. Day 6, $p = 0.016$; Day 2 vs. Day 6, $p = 6.5 \times 10^{-5}$; Day 2 vs. Day 10, $p = 0.019$; Day 2 vs.

Day 24, $p = 0.017$; Day 6 vs. Day 24, $p = 0.041$; right aDLS, $F[4,52] = 7.462$, $p = 7.9 \times 10^{-5}$, *post*

hoc Bonferroni test, Day 2 vs. Day 6, $p = 2.9 \times 10^{-4}$; Day 2 vs. Day 10, $p = 0.036$; Day 6 vs. Day 24,

$p = 0.005$), pVLS (K, one-way repeated ANOVA, left pVLS, $F[2.368,30.784] = 4.152$, $p = 0.020$,

post hoc Bonferroni test; single lever vs. Day 10, $p = 1.8 \times 10^{-5}$; right pVLS, $F[4,52] = 5.995$, $p =$

4.8×10^{-4}, *post hoc* Bonferroni test, single lever vs. Day 10, $p = 0.001$; Day 2 vs. Day 10, $p = 0.011$;

Day 6 vs. Day 10, $p = 0.032$), and DMS (L, one-way repeated ANOVA, left DMS, $F[4,52] = 12.836$,

$p = 2.4 \times 10^{-7}$, *post hoc* Bonferroni test; Day 2 vs. Day 24, $p = 1.8 \times 10^{-5}$; Day 6 vs. Day 24, $p = 4.8$

$\times 10^{-4}$; Day 10 vs. Day 24, $p = 5.5 \times 10^{-5}$; right DMS, $F[4,52] = 10.717$, $p = 2.0 \times 10^{-6}$, *post hoc*

Bonferroni test; single lever vs. Day 24, $p = 0.036$; Day 2 vs. Day 24, $p = 0.011$; Day 6 vs. Day 24,

$p = 2.6 \times 10^{-4}$; Day 10 vs. Day 24, $p = 0.011$). Arrowheads indicate the day with the most

activations throughout the learning process. Data are indicated as the mean ± s.e.m., and individual

data are overlaid. The anteroposterior coordinates from bregma (mm) are shown (F–I). Scale bar; 2

mm (I). *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$.

Figure 3. **Impacts of excitotoxic lesion of striatal subregions on the acquisition of auditory**

discrimination. Rats were given intracranial injection of PBS or IBO solution into the aDLS ($n = 8$

for each injection), pVLS ($n = 8$ for each injection), or DMS ($n = 7$ for PBS injection, and $n = 6$ for

IBO injection) before start of the single lever press task. (A-C) Representative images of NeuN

immunostaining and individual schematic illustrations showing lesioned area in the aDLS (A),

pVLS (B), or DMS (C). Dotted lines in the images indicate the range of the lesioned area. ac,

anterior commissure; cc, corpus callosum; and LV, lateral ventricle. (D-F) Learning curves during

the single lever press task in the groups injected into the aDLS (D, two-way repeated ANOVA,

group, $F[1,14] = 1.275$, $p = 0.278$, day, $F[1.512,21.171] = 26.281$, $p = 7.0 \times 10^{-6}$, group × day,

$F[1.512,21.171] = 0.834$, $p = 0.418$), pVLS (E, two-way repeated ANOVA, group, $F[1,14] = 2.542$,

$p = 0.133$, day, $F[1.547,21.658] = 40.183$, $p = 2.2 \times 10^{-7}$, group × day, $F[1.547,21.658] = 0.953$, $p =$

0.380), or DMS (F, two-way repeated ANOVA, group, $F[1,11] = 0.025$, $p = 0.876$, day,

$F[1.205,13.252] = 14.818$, $p = 0.001$, group × day, $F[1.205,13.252] = 0.610$, $p = 0.478$). (G-I)

Learning curves during the auditory discrimination task in the groups injected into the aDLS (G,

two-way repeated ANOVA, group, $F[1,14] = 11.578$, $p = 0.004$, day, $F[2.805,39.274] = 60.733$, $p =$

1.8×10^{-14}, group × day, $F[2.805,39.274] = 3.863$, $p = 0.018$), pVLS (H, two-way repeated ANOVA,

group, $F[1,14] = 7.221$, $p = 0.018$, day, $F[3.646,51.041] = 51.794$, $p = 9.2 \times 10^{-17}$, group × day,

$F[3.646,51.041] = 1.612$, $p = 0.190$), or DMS (I, two-way repeated ANOVA, group, $F[1,11] = 0.469$,

$p = 0.507$, day, $F[2.677,29.452] = 48.860$, $p = 3.9 \times 10^{-11}$, group × day, $F[2.677,29.452] = 1.094$, p

$= 0.362$). (J-L) Changes in the proportion of the WSW or LSL strategy through the discrimination

learning in the groups treated into the aDLS (J, two-way repeated ANOVA, group, $F[1,14] = 13.451$,

$p = 0.003$, day, $F[3.048,42.673] = 54.777$, $p = 9.1 \times 10^{-15}$, group × day, $F[3.048,42.673] = 3.288$, p

$= 0.029$ for the WSW; group, $F[1,14] = 5.039$, $p = 0.041$, day, $F[3.866,54.119] = 14.578$, $p = 5.2 \times$

10^{-8}, group × day, $F[3.866,54.119] = 2.471$, $p = 0.057$ for the LSL), pVLS (K, two-way repeated

ANOVA, group, $F[1,14] = 9.251$, $p = 0.009$, day, $F[3.206,44.881] = 47.168$, $p = 2.7 \times 10^{-14}$, group ×

day, $F[3.206,44.881] = 1.862$, $p = 0.146$ for the WSW; group, $F[1,14] = 0.762$, $p = 0.397$, day,

$F[4.773,66.815] = 24.018$, $p = 1.9 \times 10^{-13}$, group × day, $F[4.773,66.815] = 1.338$, $p = 0.260$ for the

LSL), or DMS (L, two-way repeated ANOVA, group, $F[1,11] = 0.703$, $p = 0.420$, day,

$F[3.052,33.568] = 48.424$, $p = 2.3 \times 10^{-12}$, group × day, $F[3.052,33.568] = 0.846$, $p = 0.480$ for the

WSW; group, $F[1,11] = 0.002$, $p = 0.965$, day, $F[3.442,37.866] = 13.787$, $p = 1.0 \times 10^{-6}$, group ×

day, $F[3.442, 37.866] = 0.581$, $p = 0.654$ for the LSL). Data are indicated as the mean ± s.e.m., and

individual data are overlaid (except for panels J–L). The anteroposterior coordinates from bregma

(mm) are shown (A–C). Scale bars; 2 mm (A–C). *$p < 0.05$ and **$p < 0.01$.

Figure 4. Influence of transient inhibition of striatal subregions at different timings on the

performance of auditory discrimination. Rats received intracranial injection of SAL or MUS

solution into the aDLS (n = 8 for each injection) or pVLS (n = 6 for each injection). (A and B)

Representative images of cresyl violet staining and schematic illustrations showing placement sites

of the tip of guide cannula in the aDLS (A) or pVLS (B). Dotted lines in the images indicate the

position of the cannula placement. Ac, anterior commissure; cc, corpus callosum; and LV, lateral

ventricle. (C and B) Effects of transient striatal inhibition on the performance. For aDLS inhibition

(C), success rate (left, two-way repeated ANOVA, stage, $F[2,28]$ = 28.708, p = 1.7×10^{-7}, group,

$F[1,14]$ = 8.840, p = 0.010, stage × group, $F[2,28]$ = 0.757, p = 0.478; unpaired Student's t-test,

early, $t[14]$ = 0.558, p = 0.586; middle, $t[14]$ = 2.764, p = 0.015; Welch's t-test, late, $t[12.010]$ =

1.617, p = 0.132, p = 0.016 after a Bonferroni correction method) and subtracted success rate on

day N-1 from that on day N (right, two-way repeated ANOVA, stage, $F[2,28]$ = 1.231, p = 0.307,

group, $F[1,14]$ = 10.797, p = 0.005, stage × group, $F[2,28]$ = 2.360, p = 0.113; unpaired Student's t-

test, early, $t[14]$ = 0.285, p = 0.780, late, $t[14]$ = 1.271, p = 0.225, Welch's t-test, middle, $t[9.603]$ =

3.331, p = 0.008, p = 0.016 after a Bonferroni correction method). For pVLS inhibition (D), success

rate (left, two-way repeated ANOVA, stage, $F[2,20]$ = 17.642, p = 3.8×10^{-5}, group, $F[1,10]$ =

43.942, $p = 5.9 \times 10^{-5}$, stage × group, $F[2,20] = 4.729$, $p = 0.012$; simple main effect, early, $p =$

0.055, middle, $p = 0.109$, late, $p = 4.2 \times 10^{-5}$) and subtracted success rate on day N-1 from that on

day N (right, two-way repeated ANOVA, stage, $F[2,20] = 12.105$, $p = 3.6 \times 10^{-4}$, group, $F[1,10] =$

16.310, $p = 0.002$, stage × group, $F[2,20] = 8.500$, $p = 0.002$; simple main effect, early, $p = 0.595$,

middle, $p = 0.085$, late, $p = 3.0 \times 10^{-6}$). (E and F) Subtracted proportion of the WSW or LSL

strategy in the groups injected into the aDLS at the middle stage (E, unpaired Student's t-test, $t[14]$

$= 2.038$, $p = 0.061$ for the WSW, and $t[14] = 2.714$, $p = 0.017$ for the LSL) and the late stage (F,

unpaired Student's t-test, $t[14] = 0.898$, $p = 0.384$ for the WSW, and $t[14] = 0.226$, $p = 0.824$ for the

LSL). (G and H) Subtracted proportion of the WSW or LSL strategy in the groups injected into the

pVLS at the middle stage (G, unpaired Student's t-test, $t(10) = 1.924$, $p = 0.083$ for the WSW, and

Welch's t-test, $t[6.095] = 1.364$, $p = 0.221$ for the LSL) and the late stage (H, unpaired Student's t-

test, $t[10] = 3.629$, $p = 0.005$ for the WSW, and $t[10] = 1.577$, $p = 0.146$ for the LSL). Data are

indicated as the mean ± s.e.m., and individual data are overlaid. The anteroposterior coordinates

from bregma (mm) are shown (A and B). Scale bars; 2 mm (A and B). *$p < 0.05$, **$p < 0.01$, and

***$p < 0.001$.

1032

Figure 5. Multi-unit recording of neurons in striatal subregions and firing activity related to the behavioral outcome of RS-HR neurons in the aDLS. (A) Schematic illustration of a freely moving rat used for simultaneous multi-unit recordings in the aDLS and pVLS. Six rats were used

for the following analysis. (B) Representative images showing positions of electrode tips (arrows)

in the aDLS and pVLS (left rows). The recording sites estimated by electrode tracks and electrical

marks in individual rats are shown (center and right rows). ac, anterior commissure; LV, lateral

ventricle. (C) Sequence of some events in correct and error trials. The delay periods were

pseudorandomly added between a correct lever press and the reward sound (0.5 ± 0.2 s). The room

light turned off the extended time (4 s) after the correct response or immediately after the error

response. (D) Mean firing rate (top rows) and auROC values (bottom rows) of RS-HR neurons in

the aDLS during the period when the RS is presented (green shadows, 500 ± 200 ms after the lever

press) at the early ($n = 43$ neurons), middle ($n = 37$ neurons), and late ($n = 44$ neurons) stages. Time

bins with significant differences between the mean firing rate in the HR or LR trial and any of the

rates in other three trials (Wilcoxon signed rank test) or the distribution of auROC values and 0.5

(Wilcoxon rank-sum test) are represented by the circles at the top. (E) Averaged firing rate during

the RS period in the HR and LR trials at the early, middle, and late stages (Kruskal-Wallis test: HR

trial, $\chi^2 = 3.028$, $p = 0.220$; and LR trial, $\chi^2 = 1.249$, $p = 0.536$). (F) Cumulative probability of the

proportion of RS-HR neuron number at the three stages (Kruskal-Wallis test, $\chi^2 = 45.378$, $p = 1.4 \times$

10^{-10}; *post hoc* Tukey-Kramer test, $p = 7.1 \times 10^{-5}$ for early vs. middle, $p = 1.0 \times 10^{-9}$ for early vs.

late, and $p = 0.040$ for middle vs. late). Data are indicated as the mean $\pm$ s.e.m. (D) or the median

and quartiles with the maximal and minimal values (E). The anteroposterior coordinates from

bregma (mm) are shown (B). Scale bars; 2 mm (B). $*p < 0.05$ and $***p < 0.001$.

1055

Figure 6. Sustained activity after the reward of FL-HR neurons in the aDLS and pVLS. (A)

Firing rate (top rows), auROC (middle rows), and licking rate (bottom rows) after the reward of FL-

HR neurons in the aDLS at the early (n = 27 neurons), middle (n = 31 neurons), and late (n = 49

neurons) stages. Time bins with significant differences between the mean firing rate in the HR trial

and any of the rate in other three trials (Wilcoxon signed rank test) or the distribution of auROC

values and 0.5 (Wilcoxon rank-sum test) are represented by the circles at the top. Timings of the RS

and FL are shown. (B) Averaged firing rate for 5 s after the FL in the HR trial (Kruskal-Wallis test,

χ^2 = 16.396, p = 2.8 × 10^{-4}, *post hoc* Tukey-Kramer test, early vs. middle, p = 0.726, early vs. late, p

= 7.2 × 10^{-4}, middle vs. late, p = 0.009). (C) Averaged total number of time bins above 3 S.D. of the

baseline firing of aDLS neurons for 5 s after the FL in the HR trial (Kruskal-Wallis test, χ^2 =

12.803, p = 0.002, *post hoc* Tukey-Kramer test, early vs. middle, p = 0.173, early vs. late, p = 0.001,

middle vs. late, p = 0.219). (D) Cumulative probability of the proportion of FL-HR neuron number

(Kruskal-Wallis test, χ^2 = 39.280, p = 3.0 × 10^{-9}, *post hoc* Tukey-Kramer test, early vs. middle, p

$= 1.3 \times 10^{-7}$, early vs. late, $p = 0.995$, middle vs. late, $p = 2.3 \times 10^{-7}$). (E) Distribution of the

correlation coefficient between the numbers of spikes of FL-HR neurons and licking. Closed

column indicates the number of neurons showing significant correlations between the two

parameters (5 out of 107 neurons). (F) Firing rate (top rows), auROC (middle rows), and licking

rate (bottom rows) after the reward of FL-HR neurons in the pVLS at the early ($n = 17$ neurons),

middle ($n = 11$ neurons), and late ($n = 64$ neurons) stages. Time bins with significant differences

between the rate in the HR trial and any of the rate in other trials (Wilcoxon signed rank test) or the

distribution of auROC values and 0.5 (Wilcoxon rank-sum test) are represented by the circles at the

top. The timings of the RS and FL are shown. (G) Averaged firing rate of pVLS neurons after the

FL in the HR trial (Kruskal-Wallis test, $\chi^2 = 7.652$, $p = 0.022$; *post hoc* Tukey-Kramer test, early vs.

middle, $p = 0.975$, early vs. late, $p = 0.076$, and middle vs. late, $p = 0.096$). (H) Averaged total

number of time bins above 3 S.D. of the baseline firing after the FL in the HR trial (Kruskal-Wallis

test, $\chi^2 = 0.023$, $p = 0.989$). (I) Cumulative probability of the proportion of FL-HR neuron number

(Kruskal-Wallis test, $\chi^2 = 47.922$, $p = 3.9 \times 10^{-11}$; *post hoc* Tukey-Kramer test, early vs. middle, $p =$

0.011, early vs. late, $p = 1.7 \times 10^{-4}$, and middle vs. late, $p = 9.7 \times 10^{-10}$). (J) Distribution of the

correlation coefficient between the number of spikes in the pVLS and the licking. Closed column

indicates the number of neurons showing significant correlations between the two parameters (5 out

of 92 neurons). Data are indicated as the mean ± s.e.m. (A and F) or the median and quartiles with

the maximal and minimal values (B, C, G, and H). **$p < 0.01$ and ***$p < 0.001$.

1088

Figure 7. Transient activity related to the beginning and ending of a behavior of CO-HR and CR-HR neurons in the pVLS. (A) Firing rate (top rows) and auROC values (bottom rows) of CO-HR neurons in the pVLS at the early ($n = 4$ neurons), middle ($n = 21$ neurons), and late ($n = 24$ neurons) stages. Time bins with significant differences between the mean firing rate in the HR trial and either of the rate in other three trials (Wilcoxon signed rank test) or the distribution of auROC values and 0.5 (Wilcoxon rank-sum test) are represented by the circles at the top. (B) Averaged firing rate of CO-HR neurons in HR and LL trials (0-600 ms after the CO) (Wilcoxon signed rank test; early, $p = 0.875$, middle, $p = 0.006$, and late, $p = 0.004$). (C) Cumulative probability of the proportion of CO-HR neuron number (Kruskal-Wallis test, $\chi^2 = 51.114$, $p = 8.0 \times 10^{-12}$; *post hoc* Tukey-Kramer test, early vs. middle, $p = 9.6 \times 10^{-10}$, early vs. late, $p = 0.002$, and middle vs. late, $p = 6.4 \times 10^{-4}$). (D) Firing rate (top rows) and auROC values (bottom rows) of CR-HR neurons in the pVLS at the early ($n = 11$ neurons), middle ($n = 22$ neurons), and late ($n = 41$ neurons) stages. Time bins with significant differences between the mean rate in the HR trial and any of the rate in other trials (Wilcoxon signed rank test) or the distribution of auROC values and 0.5 (Wilcoxon rank-sum test) are represented by the circles at the top. The timing of the RS is presented. (E) Averaged firing rate of CR-HR neurons in HR and LL trials (0-600 ms after the CR) (Wilcoxon signed rank test; early, $p = 0.278$, middle, $p = 0.758$, and late, $p = 0.279$). (F) Cumulative probability of the

1106 proportion of CR-HR neuron number (Kruskal-Wallis test, $\chi^2 = 13.786$, $p = 0.001$; *post hoc* Tukey-

1107 Kramer test, early vs. middle, $p = 0.003$, early vs. late, $p = 0.005$, and middle vs. late, $p = 0.976$).

1108 Data are indicated as the mean ± s.e.m. (A and B) or the median and quartiles with the maximal and

1109 minimal values (B and E). **$p < 0.01$ and ***$p < 0.001$.

1110

Article and author information

Susumu Setogawa

Department of Physiology, Osaka Metropolitan University Graduate School of Medicine, Osaka 545-8585, Japan

ORCID: 0000-0002-9161-2656

Takashi Okauchi

Laboratory for Biofunction Dynamics Imaging, RIKEN Center for Biosystems Dynamics Research, Kobe 650-0047, Japan4Japan Society for the Promotion of Science, Tokyo 102-0083, Japa

Di Hu

Department of Neurophysiology, Graduate School of Biomedical & Health Sciences, Hiroshima University, Hiroshima 734-8553, Japan

Yasuhiro Wada

Laboratory for Pathophysiological and Health Science, RIKEN Center for Biosystems Dynamics Research, Kobe 650-0047, Japan

ORCID: 0000-0003-2096-1602

Keigo Hikishima

Medical Devices Research Group, Health and Medical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba 305-8564, Japan

ORCID: 0000-0003-1133-1297

Hirotaka Onoe

Human Brain Research Center, Kyoto University Graduate School of Medicine, Kyoto 606-8507,

1137   Japan

1138

1139   Kayo Nishizawa

1140   Department of Molecular Genetics, Institute of Biomedical Sciences, Fukushima Medical

1141   University School of Medicine, Fukushima 960-1295, Japan

1142   ORCID: 0009-0002-8398-588X

1143

1144   Nobuyuki Sakayori

1145   Department of Physiology and Oral Physiology, Graduate School of Biomedical and Health

1146   Sciences, Hiroshima University, Hiroshima 734-8553, Japan

1147   ORCID: 0000-0003-1611-4187

1148

1149   Hiroyuki Miyawaki

1150   Department of Physiology, Osaka Metropolitan University Graduate School of Medicine, Osaka

1151   545-8585, Japan

1152   ORCID: 0000-0003-1556-1348

1153

1154   Takuma Kitanishi

1155   Department of Physiology, Osaka City University Graduate School of Medicine, Osaka 545-8585,

1156   Japan

1157   ORCID: 0000-0001-8062-3854

1158

Kenji Mizuseki

Department of Physiology, Osaka Metropolitan University Graduate School of Medicine, Osaka

545-8585, Japan

ORCID: 0000-0002-1456-2149

Yilong Cui

Laboratory for Biofunction Dynamics Imaging, RIKEN Center for Biosystems Dynamics Research,

Kobe 650-0047, Japan[4]Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

ORCID: 0000-0002-8302-1899

1167

1168   Kazuto Kobayashi

1169   Department of Molecular Genetics, Institute of Biomedical Sciences, Fukushima Medical

1170   University School of Medicine, Fukushima 960-1295, Japan

1171   ORCID: 0000-0002-7617-2939