Title page

Manuscript title

Prdm1 Positively Regulates Liver Group 1 ILCs Cancer Immune Surveillance and Preserves Functional Heterogeneity

Authors and affiliations

Jitian He[1], Le Gao[1], Peiying Wang[1], Wing Keung Chan[4], Yiran Zheng[1], Yumo Zhang[1], Jiaman Sun[1], Jiming Wang[5], Xiaohong Li[1], Huaiyong Chen[3,6,7], Zhouxin Yang[2], Youwei Wang[1]

[1]Institute of Medical Engineering & Translational Medicine, Tianjin University, Tianjin 300072, China.

[2]Zhejiang Provincial Key Lab of Geriatrics and Geriatrics Institute of Zhejiang Province, Zhejiang Hospital, Hangzhou 310030, China.

[3]College of Pulmonary and Critical Care Medicine, 8th Medical Center, Chinese PLA General Hospital, Beijing, China

[4]Department of Internal Medicine, Division of Hematology, The Ohio State University, Columbus, OH 43210, USA.

[5]Tianjin Economic-Technological Development Area (TEDA) Hospital, Tianjin 300451, China

[6]Department of Basic Medicine, Haihe Hospital, Tianjin University, Tianjin 300350, China.

[7]Tianjin Key Laboratory of Lung Regenerative Medicine, Tianjin 300350, China.

Authorship notes: JH, LG, and PW contributed equally to this work.

Co-corresponding authors: Xiaohong Li (xhli18@tju.edu.cn), Huaiyong Chen

(huaiyong.chen@foxmail.com), Zhouxin Yang (yangzhouxin@hotmail.com), and Youwei Wang

(youwei.wang@tju.edu.cn)

Address correspondence to: Youwei Wang, Institute of Medical Engineering & Translational

Medicine, Tianjin University, Tianjin 300072, China. Email: youwei.wang@tju.edu.cn. Or to

Zhouxin Yang, Zhejiang Provincial Key Lab of Geriatrics and Geriatrics Institute of Zhejiang

Province, Zhejiang Hospital, Hangzhou 310030, China. Email: yangzhouxin@hotmail.com. Or to

Huaiyong Chen, Department of Basic Medicine, Haihe Hospital, Tianjin University, Tianjin

300350, China. Email: huaiyong.chen@foxmail.com.

Abstract

Group 1 innate lymphoid cells (ILCs) comprise conventional natural killer (cNK) cells and type 1 innate lymphoid cells (ILC1s). The main functions of liver cNK cells and ILC1s not only include directly killing target cells but also regulating local immune microenvironment of the liver through the secretion of cytokines. Uncovering the intricate mechanisms by which transcriptional factors regulate and influence the functions of liver cNK cells and ILC1s, particularly within the context of liver tumors, presents a significant opportunity to amplify the effectiveness of immunotherapies against liver malignancies. Using Ncr1-drived conditional knockout mouse model, our study reveals the regulatory role of *Prdm1* in shaping the composition and maturation of cNK cells. Although *Prdm1* did not affect the killing function of cNK cells in an *in vivo* cytotoxicity model, a significant increase in cancer metastasis was observed in *Prdm1* knockout mice. Interferon-gamma (IFN-γ), granzyme B, and perforin secretion decreased significantly in *Prdm1* deficient cNK cells and liver ILC1s. scRNA sequencing data also provided evidences that Prdm1 maintains functional subsets of cNK cells and liver ILC1s and facilitates communications between cNK cells, liver ILC1s and macrophages. The present study unveiled a novel regulatory mechanism of Prdm1 in cNK cells and liver ILC1s, showing promising potential for developing innovative immune therapy strategies against liver cancer.

Main text

Introduction

Group 1 ILCs consist of cNK cells and ILC1s (1, 2), with distinct developmental trajectories and effect molecules (3). Both cNK cells and ILC1s play indispensable roles in combatting viral infections (4), maintaining local immune homeostasis (5), eradicating malignant transformed cells (6), and fostering cross-talk with adaptive immunity (7). cNK cells demonstrate potent cellular cytotoxicity, facilitating direct elimination of target cells (8). On the contrary, a defining attribute of ILC1s is their predominant cytokine-mediated functions, with limited cellular killing capacity (9). In a state of homeostasis, liver group 1 ILCs ($CD45^+CD3^-NK1.1^+NKp46^+$) can be discriminated into cNK cells and ILC1s by the differential expression of CD49a and CD49b (2): cNK cells are marked by the expression of CD49b, while liver ILC1s exhibit a distinctive positivity for CD49a. Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) is also expressed on liver ILC1s, but not on cNK cells (10). Transcriptional factors (TFs) such as T-bet, Nfil3, PLZF, and ID2 are required for cNK cells and liver ILC1s development or generation of their progenitors (11-14). Eomes is considered to be necessary for NK cell maturation, but not for ILC1s (15). Liver environment facilitated T-bet expression in the early stage of NK cells development, which results in Eomes repression. The repression of T-bet is required for $Eomes^+$ NK cells (16). On the contrary, deficiency of Hobit results in the depletion of ILC1s, while only leaving little impact on liver NK cells (17). These studies suggest that some TFs may, or usually, play different roles in NK cells and ILC1s.

During the development of liver cancers, the functions of immune cells are often inhibited,

resulting in the formation of an immunosuppressive tumor microenvironment, and sometimes

even systemic immune suppression (18). Transforming a cold tumor into a hot tumor with a

stronger immune response is one of the goals of many cancer immunotherapies (19). Research

targeting the immune system is showing increasing clinical promise in liver cancer treatment (20).

One of our recent studies also found that Toll-like receptor agonists can significantly enhance the

anti-tumor effect of Sorafenib by reconstructing the tumor immune microenvironment and

reshaping the vascular system (21).

The anti-tumor activity of cNK cells in the liver is relatively well-established, whereas the

relationship between liver ILC1 and tumors is still a topic of controversy. Clinical data showed

that liver tumors with a higher infiltration of NK cells are associated with a better prognosis (22).

Decreased NK cell activity is closely correlated with the malignancy of liver tumors and

represents a significant risk factor for recurrence (23). The tumor microenvironment orchestrates

the transformation of cNK cells into ILC1s in a TGF-β-dependent manner, resulting in their

diminished capacity to control tumor growth and metastasis. This process ultimately promotes

tumor immunoevasion (24). However, researches also revealed that, distinct from directly

inhibiting tumor growth, the primary function of ILC1 is to suppress the seeding of metastatic

tumor cells in liver tissue (6). Comprehensive research in this field is essential to harness the

precision of group 1 ILCs targeted immunotherapy for liver cancers.

The transcription factor network governs the function of group 1 ILCs and the balance

between cNK and ILC1. Our research team, along with others, has observed that one of the

transcription factor in TGF-β pathway, Smad4, promotes the shift in balance from cNK cells

towards ILC1 in a TGF-beta-independent pathway, and simultaneously, it positively regulates the

expression of another transcription factor, PR domain 1 (Prdm1/Blimp1) (25, 26). Prdm1 plays a

crucial role in the differentiation of B cells into plasma cells and the homeostasis of T cells

(27-29). The function and regulatory network of Prdm1 in NK cells is distinct from B cells and T

cells, for its expression is independent of B-cell lymphoma 6 (Bcl-6) and Interferon Regulatory

actor 4 (IRF4) but relies on T-bet. In the study that identified Prdm1 as an essential transcriptional

factor for NK cell maturation (30), no significant differences were observed in IFN-γ production

or cytotoxicity between Prdm1-deficient and wild-type NK cells. Although Prdm1 expression is

dependent on IL-15 in immature NK cells and can be further upregulated by IL-12 or IL-21, it

plays a role in downregulating the expression of certain cytokine receptors, such as CD25,

consequently diminishing the responsiveness of NK cells to IL-2 (31). Notably, some studies even

suggested that Prdm1 suppressed the secretion of IFN-γ in NK cells (32). These findings imply

that, although Prdm1 promotes NK cell development, it may function as a negative regulator of

their activity. However, direct evidence supporting the impact of Prdm1 on NK cell anti-tumor

capabilities is still lacking. Furthermore, there has been no investigation into its influence on the

homeostasis of cNK cells and ILC1s in the liver, nor has there been an exploration of the

underlying mechanisms.

In the current study, we found that the deletion of *Prdm1* in Ncr1[+] cells resulted in an

imbalance in the homeostasis of liver group 1 ILCs, with a shift towards cNK cells. The data also

support the essential role of Prdm1 in the cancer surveillance mediated by cNK cells. While

Prdm1 positively regulates genes associated with cellular cytotoxicity, it concurrently exerts

inhibitory control over certain positive regulators of NK cell development and functionality, such

as JunB. Using single-cell RNA sequencing, we have identified a subset of cNK cells within the

liver characterized by elevated JunB expression. These cells exhibit decreased expression of genes

associated with cellular cytotoxicity, and their abundance significantly increases following Prdm1

knockout. Furthermore, our data also support that Prdm1 promotes the cross-talk between group 1

ILCs and macrophages.

Results

Prdm1 **promotes group 1 ILCs homeostasis and terminal maturation**

Examination of 363 liver hepatocellular carcinoma (LIHC) patient samples from The Cancer

Genome Atlas (TCGA) revealed a positive correlation between the expression of NK

cell-associated genes (33) (*NCR1*, *KLRB1*, *CD160*, *PRF1,* etc.) and *PRDM1* expression (Figure

1A). The patients are ordered from highest to lowest based on the expression of NK-Prdm1 for

survival analysis (Figure 1B). Notably, patients exhibiting higher levels of NK-PRDM1

expression (above the median) experienced better survival outcomes compared to those with lower

levels of NK-PRDM1 expression (below the median) (Figure 1C). Similar results were also found

in skin cutaneous melanoma (SKCM, n=454) and lung adenocarcinoma (LUAD, n=497) patients

(Supplemental Figure 1, A-F). Patients within the highest quartile of NK-PRDM1 signature

expression demonstrated enhanced overall survival, a result that achieved statistical significance in

LUAD and SKCM patients (Supplemental Figure 1, G-I). These data suggested that *PRDM1* in

NK cells might be essential for immune surveillance in solid tumors, including liver cancer, and

prompted us to investigate the function and mechanism of *PRDM1* in NK cells and ILC1 within

the context of liver cancer.

Ncr1-Cre mice were crossed with *Prdm1$^{fl/fl}$* mice to generate *Ncr1-cre Prdm1$^{fl/fl}$* mice, which

specifically knockout exons 6-8 of *Prdm1* (34) in NKp46 positive cells (Figure 1D). The mice

carrying $Ncr1^{Cre/+}Prdm1^{fl/fl}$ were referred to as $Prdm1^{\Delta Ncr1}$ mice, and the mice carrying

$Ncr1^{+/+}Prdm1^{fl/fl}$ were referred to as $Prdm1^{+/+}$ mice or wild-type (WT) mice. To further validate

the deletion of *Prdm1* in NKp46$^+$ cells of $Prdm1^{\Delta Ncr1}$ mice, CD3$^-$NK1.1$^+$NKp46$^+$ cells were sorted

and the expression of *Prdm1* was measured by real-time RT-PCR. The expression of *Prdm1* was

almost undetectable in CD3$^-$NK1.1$^+$NKp46$^+$ cells of $Prdm1^{\Delta Ncr1}$ mice, while was similar between

total splenocytes in $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice (Figure 1, E and F) indicating successful, and

specific knockout of *Prdm1* in NKp46$^+$ cells.

Proportion and absolute number of cNK cells in blood, bone marrow, lung, liver, spleen, and

lymph nodes were analyzed by flow cytometry. Compared with $Prdm1^{+/+}$ mice, the percentage

and absolute number of cNK cells (CD45$^+$CD3$^-$NK1.1$^+$NKp46$^+$) among lymphocytes was

decreased in all of these tissues, whereas increased number of NK cells were observed in bone

marrow (Figure 1G; Supplemental Figure 2A). NK cell terminal maturation can be divided into

four stages according to the expression of CD11b and CD27 (35). Stage IV (CD11b$^+$CD27$^-$) NK

cells show higher level of effector molecules than any other stages and are considered as the most

mature NK cells (36). The maturation of cNK cells (gated by CD45$^+$CD3$^-$NK1.1$^+$NKp46$^+$CD49b$^+$)

from blood, bone marrow, lung, liver, spleen, and lymph nodes were assessed, based on the

expression of CD11b and CD27. Compared with $Prdm1^{+/+}$ mice, the proportion of the most

mature CD11b$^+$CD27$^-$ NK cells were significantly decreased in all of the analyzed tissues in

$Prdm1^{\Delta Ncr1}$ mice (Figure 1H; Supplemental Figure 2B). Killer cell lectin-like receptor subfamily G

member 1 (KLRG1) is a lectin-like receptor which was considered as another marker of NK cell

maturation (37). In both $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice, cNK cells from the liver and lung had the

highest expression of KLRG1, following by blood and spleen (Supplemental Figure 2C). The

lowest KLRG1 expression was observed in cNK cells derived from lymph nodes and bone marrow,

indicating the presence of the most immature cNK cells in these tissues. Consistent with

CD11b/CD27 based maturation analysis, a significant loss of KLRG1$^+$ NK cells in cNK cells was

observed in *Prdm1$^{\Delta Ncr1}$* mice (Supplemental Figure 2C). Together, these data comfired that *Prdm1*

is required for the terminal maturation of cNK cells among various tissues.

Prdm1 is required for group 1 ILCs to control tumor metastasis

Two subpopulations of liver group 1 ILCs (gated by CD45$^+$CD3$^-$NK1.1$^+$NKp46$^+$) were

further analyzed based on the expression of CD49a and CD49b (Figure 2A). Compared with

Prdm1$^{+/+}$ mice, the *Prdm1$^{\Delta Ncr1}$* mice exhibited an increased percentage of cNK cells

(CD49a$^-$CD49b$^+$) and reduced proportion of ILC1s (CD49a$^+$CD49b$^-$) (Figure 2A). Of note, the

absolute number of both cNK cells and ILC1s were decreased in *Prdm1$^{\Delta Ncr1}$* mice, with a more

robustly reduction in ILC1s (Figure 2A), which underscored the crucial role of *Prdm1* in

maintaining the quantity of both liver cNK cells and ILC1s. Expression level of CD49b was

slightly upregulated in *Prdm1$^{\Delta Ncr1}$* cNK cells and NKp46$^+$ cells in the liver and other tissues

(Supplemental Figure 2, D and E). Increased CD49a expression was also observed in *Prdm1$^{\Delta Ncr1}$*

liver ILC1s, while it showed decreased expression in NKp46$^+$ cells in the liver, bone marrow, and

lymph nodes (Supplemental Figure 2, F and G). These results indicated the essential role of Prdm1

in maintaining the balance and hemostasis of cNK cells and ILC1s.

The decreased quantity of NK cell homeostasis and maturation of cNK cells due to *Prdm1*

loss motivated us to further explore whether deficiency of Prdm1 impaired NK cell cytotoxicity. A

B2M-deficient cell-based *in vivo* cytotoxicity assay was used to evaluate the effect of *Prdm1* on

the cytotoxicity of NK cells (38, 39). B2M-deficient cells do not have detectable Major

Histocompatibility Complex I (MHC-$\square$) on the cell surface, making them the target of NK cells

(40). Health NK cells will reject B2M-deficient donor cells efficiently and the elimination was

used to quantify the cytotoxicity of NK cells. Although significant impaired homeostasis and

maturation of NK cells were observed in *Prdm1$^{\Delta Ncr1}$* mice, no significant difference in the *in vivo*

cytotoxicity assay were observed between *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$* mice (Supplemental Figure 2,

H and I).

Besides their direct cytotoxic capabilities, NK cells' anti-tumor potential is also influenced by

additional factors. These include their ability to counteract tumor-induced immune suppression

and exhaustion, enhancing their effectiveness against cancer cells. Furthermore, NK cells can

secrete cytokines that activate other immune cells, thereby orchestrating a broader immune

response for the elimination of tumors. Moreover, the cytotoxicity assay, due to its relatively short

duration, might not fully represent the anti-tumor activity of NK cells when continuously exposed

to immune inhibitory signals in the tumor microenvironment. Therefore, we initiated an *in vivo*

tumor model to further investigate the impact of *Prdm1* on NK cell anti-tumor capability. B16F10

is a melanoma cell line with low expression of MHC-$\square$, which was susceptible to NK cell killing

and usually used to evaluate NK cell anti-tumor capacity (41-43). The B16F10 cells were

intravenously (for lung metastasis) or intrasplenic (for liver metastasis) administrated in the mice.

The melanoma nodes were quantified three (intravenous injection) or two (intrasplenic injection)

weeks after tumor inoculation. Compared with *Prdm1$^{+/+}$* mice, deficiency of *Prdm1* resulted in

more metastasis nodules in both lung (~ 2-fold) and liver (~ 4-fold) (Figure 2, B and D).

Histological analysis further confirmed the increased frequency of metastasis tumor foci in

Prdm1^{ΔNcr1} mice (Figure 2, C and E).

In agreement with our *in vivo* data, we also observed decreased IFN-γ secretion in *Prdm1^{ΔNcr1}*

mice-derived splenic cNK cells, liver cNK cells, and liver ILC1s when stimulated by IL-18 alone

or IL-12/IL-18 (Figure 2F; Supplemental Figure3; Supplemental Figure 12B), which indicated that

Prdm1 is required for full activation of cNK cells and ILC1s in the context of IFN-γ production.

These data implying that *Prdm1* is indispensable for NK cell mediated tumor surveillance.

Bulk RNA-seq depicts *Prdm1*-mediated functions in cNK cells

Bulk RNA sequencing of splenic cNK cells (CD3$^-$NK1.1$^+$NKp46$^+$) was conducted to uncover

the molecular mechanisms by which *Prdm1* regulates NK cell anti-tumor immunity (Figure 3A).

Differentially expressed genes (DEGs) between *Prdm1^{+/+}* and *Prdm1^{ΔNcr1}* mice were determined

using a criterion of $\log_2$ (fold change) > 0.5 and P < 0.05. 445 DEGs were identified out of 17434

protein-coding genes, which consisted of 223 upregulated genes and 222 downregulated genes

(Figure 3B).

Gene Ontology (GO) analysis revealed the enrichment of glucuronate metabolism and

lymphocyte differentiation in upregulated genes in *Prdm1^{ΔNcr1}* mice derived NK cells (Figure 3C),

both of which were associated with cellular growth and development. In contrast, leukocyte

mediate cytotoxicity, immune receptor activity, and integrin binding was enriched in the genes

which decreased their expression level in in *Prdm1^{ΔNcr1}* mice (Figure 3C).

Gene Set Enrichment Analysis (GSEA) showed that NF-kappa B signaling pathway enriched

in *Prdm1*-deficient cNK cells (Figure 3D), suggesting the potential targets by *Prdm1* to regulate

NK cell function. Increased expression of multiple TFs such as *Junb*, *Batf3*, *Nfkb1*, *Tcf7*, and

Nr4a2 was observed in *Prdm1* knockout cNK cells, suggesting they might be suppressed by

Prdm1 (Figure 3E). Downregulation of granzyme B (*Gzmb*), Perfroin (*Prf1*) were observed in

Prdm1 deficient NK cells (Figure 3E), implied decreased anti-tumor ability, which was consistent

with increased melanoma metastasis in *Prdm1$^{\Delta Ncr1}$* mice (Figure 2, B and D). CXCR6 and

CX3CR1 was considered to play an important role in promoting the egress of NK cells from bone

marrow. Decreased expression of *Cxcr6* and *Cx3cr1* in *Prdm1$^{\Delta Ncr1}$* NK (Figure 3E) might be the

reason for the increased quantity of NK cells in bone marrow and decreased number in peripheral

tissues (Figure 1G). As a result of the reduced expression levels of *Cxcr6* and *Cx3cr1*, NK cells

may not be able to egress from the bone marrow and accumulated therein. Consistent with

decreased production of IFN-γ after stimulated by IL-12/IL-18 (Figure 2F), decreased expression

of *Il18rap* and *Il12rb2* were observed in *Prdm1$^{\Delta Ncr1}$* cNK cells (Figure 3E), implying impaired

response to cytokine stimulation.

To confirm the result of RNA-sequencing, the expression of fractalkine receptor (CX3CR1),

granzyme B and perforin were analyzed by flow cytometry. The percentage of CX3CR1$^+$ cNK

cells was significantly decreased in multiple tissues of *Prdm1$^{\Delta Ncr1}$* mice, while the proportion of

CX3CR1$^+$ ILC1 was increased in the liver (Figure 3, F and G). Lower GZMB and PRF1

production was observed in *Prdm1*-deficient splenic cNK cells, liver cNK cells and ILC1s (Figure

3, H-K; Supplemental Figure 4, A-I). Notably, the proportion of GZMB$^+$ and PRF1$^+$ cNK cells

was decreased among almost all of the maturation stages of cNK cells (Figure 3, J and K). The

relative mean fluorescent intensities (MFIs) of GZMB and PRF1 consistently show a reduction

across all developmental stages in *Prdm$^{\Delta Ncr1}$* NK cells (Supplemental Figure 4, H and I). Yet, no

statistical difference of PRF1 was found within the CD11b⁻CD27+ and CD11b⁺CD27⁺ subsets,

likely due to the relatively lower perforin levels in these populations (Supplemental Figure 4I).

These findings suggest that Prdm1 may directly influence cytotoxic molecule in NK cells, rather

than impacting their anti-tumor abilities solely by affecting the maturation phenotype of

Prdm1-deficient NK cells.

scRNA-seq reveals distinct properties of two clusters from liver group I ILCs following *Prdm1*

knockout

To further investigate the effect of *Prdm1* in liver cNK cells and ILC1s and the changes in the

haptic immune microenvironment caused by the deficiency of *Prdm1* in group 1 ILCs, single-cell

RNA sequencing (scRNA-seq) was performed for liver CD45$^+$ cells (Figure 4A; Supplemental

Figure 5A). Initial quality control revealed high-quality of cell purity, library assembly, and

sequencing (Supplemental Figure 5B). 10978 cells passed the quality criteria and were selected for

further analysis (6,161 from *Prdm1$^{+/+}$* mice, 4,817 from *Prdm1^{ΔNcr1}* mice). Unsupervised

clustering of all sequenced cells based on transcript signatures identified twelve distinct clusters

(Supplemental Figure 5C), including B cells, epithelial cells (ECs), CD4$^+$ T cells, CD8$^+$ T cells,

NKT cells, ILC1s, cNK cells, dendritic cells (DCs), monocyte-derived macrophages (MDMs),

Monocytes, Kupffer cells (KCs), Neutrophils, and a small number of undefined cells (Figure 4A).

In *Prdm1^{ΔNcr1}* mice, an increased proportion of MDMs, CD4$^+$ T cells, CD8$^+$ T cells, Monocytes,

and DCs was observed, alongside a decreased proportion of ECs, cNK cells, ILC1s, and NKT

cells (Supplemental Figure 5D). Liver cNK cells and ILC1s were identified and discriminated

based on the expression of surface markers and distinctive TFs (Supplemental Figure 5, E and F).

Compared with other clusters, cNK cells and ILC1s highly expressed *Ncr1* and *Klrb1c* (NK1.1)

(Supplemental Figure 5E). cNK cells expressed high levels of *Itga2* (CD49b) and Eomes, while

ILC1s had high levels expression of *Itga1* (CD49a) and *Tnfsf10* (Supplemental Figure 5, F and G).

Consistent with our flow cytometry data, both cNK cells and ILC1s have significant reduced

proportion in *Prdm1*$^{\Delta Ncr1}$ mouse (Supplemental Figure 5H). In group 1 ILCs from *Prdm1*$^{\Delta Ncr1}$ mice,

there was an increase in the proportion of cNK cells accompanied by a decrease in ILC1s (Figure

4, B and C).

To better understand the specific function of *Prdm1* in liver cNK cells and ILC1s, the two

subpopulations of liver group 1 ILCs were further analyzed separately using unsupervised

clustering and visualized by Uniform Manifold Approximation and Projection (UMAP) (Figure 4,

D and E). Based on the cluster specific gene expression signature (Supplemental Figure 6A), the

subpopulation of liver cNK cells were referred as "*Prf1*^{hi}", "*Junb*^{hi}", and "*Cxcr3*^{hi}" cNK cells

(Figure 4D), with different distribution in *Prdm1*$^{\Delta Ncr1}$ and *Prdm1*$^{+/+}$ genotype (Figure 4F;

Supplemental Figure 6B).

The *Prf1*^{hi} cNK cell cluster was defined by high expression of cytolysis-related genes,

including *Ncr1*, *Gzma*, *Gzmb*, *Prf1*, and *Fgl2* (Figure 4G; Supplemental Figure 6, C and D),

indicating the strong target-killing ability of this cluster. Although this cluster is present in both

Prdm1$^{\Delta Ncr1}$ and *Prdm1*$^{+/+}$ mice, there is a significant reduction in *Prdm1*$^{\Delta Ncr1}$ mice (Figure 4F),

indicating the importance of Prdm1 in maintaining this group of cells. GO analysis further

revealed the enrichment signatures of cytolysis, response to virus, and lymphocyte mediated

immunity in the genes upregulated in *Prf1*^{hi} cNK cell cluster, further confirming the cytotoxic

effects of this cluster (Figure 4H). These data underscore the crucial role of *Prdm1* in maintaining

NK cells with immune effector functions.

The *Junb*^hi liver cNK cell cluster distinguished themselves by higher expression of *Junb*

compared to other clusters (Figure 4G; Supplemental Figure 6, C and D). The predominant

majority (92.98%) of *Junb*^hi liver cNK cells are derived from *Prdm1*^*ΔNcr1* mice, with less than ten

percent (7.02%) originating from *Prdm1*^*+/+* mice (Figure 4F). Many signal transduction elements,

gene expression regulator, and transcriptional factors, such as *Nfkbia*, *Tnfaip3*, *Nr4a1/2/3*, *Batf3*,

Fos, *Fosb*, *Tcf7*, and *Kit* were upregulated in the *Junb*^hi liver cNK cells (Figure 4G; Supplemental

Figure 6D). The expression of cytotoxicity related genes, such as *Gzmb* and *Prf1*, in *Junb*^hi cluster

was also lower than other cNK cell clusters (Figure 4G). GO analysis showed that the genes

upregulated in *Junb*^hi liver cNK cells enriched in cell differentiation, cell activation, and

transcriptional regulation (Figure 4H). GSEA indicates that the NF-kappa B, IL-17, MAPK, and

TNF signaling pathways were upregulated in this clusters (Figure 4, I-L). GSEA also showed that

mitochondrial related pathways, such as mitochondrial protein, oxidative phosphorylation, and

respiratory electron transport chain were suppressed in *Junb*^hi cNK cell cluster (Supplemental

Figure 6, E-G). Increased proportion of *Junb*^hi cluster in *Prdm1* deficient cNK cells suggested

impaired anti-tumor activity, which was consistent with more melanoma metastasis in *Prdm1*^*ΔNcr1*

mice and lower expression of cytotoxicity-related genes in splenic cNK cells based on bulk

RNA-sequencing.

The *Cxcr3*^hi cNK cell cluster was characterized with high expression of *Cxcr3*, *Ccr2*, and

some genes encoding ribosomal subunits such as *Rps7* (Supplemental Figure 6A). Expression of

tissue-resident markers Cd69 was also highly expressed in this clusters (Supplemental Figure 6D).

The enrichment of chemokine receptors in the genes upregulated in the *Cxcr3*^hi cluster implying a

greater likelihood of this cluster being tissue-resident compared with other cNK cell clusters (Figure 4H). To further confirmed tissue-resident properties of this clusters, we calculated the module score based on top30 DEGs in ILC1 versus cNK clusters, including *Cxcr6*, *Itga1*, *Cd160*, *Cd226*, etc. *Cxcr3*^{hi} cNK clusters have the highest score among all cNK clusters (Supplemental Figure 6H), indicating the similarity with liver ILC1s. In the tumor microenvironment, reports indicated that NK cells could transform into ILC1s (24). If this conversion of cNK cells into ILC1s also occurred under normal physiological conditions, then *Cxcr3*^{hi} cNK cell cluster might be the most susceptible to such transformation.

The significant enrichment of ribosomal subunits and cytoplasmic translation in *Cxcr3*^{hi} cluster (Figure 4H) implied their distinct and active metabolic profile and the capability to mount immune responses. The remarkably decreased proportion of *Cxcr3*^{hi} cNK cell cluster in *Prdm1*$^{\Delta Ncr1}$ mice (Figure 4F), which emphasized the critical role of *Prdm1* in maintaining this cluster of liver cNK cells, consistent with the flow cytometry result that showed an increase in the number of NK cells in the bone marrow and a decrease in NK cells in peripheral tissues (Figure 1G). To confirm the regulatory effect of Prdm1 on the migration of NK cells and ILC1 through chemokine receptors, and to validate the scRNA-seq data, we further evaluated the expression of CX3CR1 by flow cytometry. A decrease in the proportion of CX3CR1^{+} NK cells was observed across all tested organs following Prdm1 knockout. However, it is noteworthy that, in contrast to the trend in NK cells, CX3CR1 expression was increased in liver ILC1s after Prdm1 knockout (Figure 3, F and G). This not only substantiates the involvement of Prdm1 in managing NK cell migration but also underscores its distinctive regulatory impacts on the chemokine receptor expressions within NK cells and ILC1 populations. Bulk RNA sequencing data also found that the

expression of *Cx3cr1* and *Cxcr6* decreased in *Prdm1*$^{\Delta Ncr1}$ cNK cells (Figure 3E). These findings

supported the hypothesis that *Prdm1*, through regulating chemokine receptor expression levels,

influenced the distribution of NK cells in the bone marrow and peripheral tissues, particularly

within the liver tissue.

Three clusters of ILC1s were identified from liver ILC1s (Figure 5A), comprising "*Il7r*^{hi}",

"*Klra*^{hi}", and "*Gzmb*^{hi}" ILC1s. *Prdm1*$^{+/+}$ and *Prdm1*$^{\Delta Ncr1}$ ILC1s seemed to co-cluster largely and

have minor difference within the proportion of clusters (Figure 5, B and C; Supplemental Figure

7B). The first two clusters of ILC1s were characterized by higher expression of *Il7r* (CD127) and

Klra5 separately, while the *Gzmb*^{hi} ILC1 cluster was identified by elevated expression of both

Gzma and *Gzmb* (Figure 5D; Supplemental Figure 7, A, C and D). Additionally, both *Gzma* and

Gzmb expression were downregulated, and *Junb* was upregulated, in *Prdm1*$^{\Delta Ncr1}$ mice derived

cNK cells and ILC1s compared to those from *Prdm1*$^{+/+}$ mice (Figure 5E). The *Il7r*^{hi} ILC1s cluster

was distinguishable from other ILC1 clusters by its high expression of *Il7r*, IL18RA (*Il18r1*), and

IFN-γ (*Ifng*) (Figure 5B; Supplemental Figure 7C). The high expression of *Il18r1* and *Ifng* in *Il7r*^{hi}

ILC1s indicated this cluster of cells was highly responsive to IL-18 (Supplemental Figure 7D).

GSEA and GO analysis showed that IL-17, NF-kappa B, TNF, MAPK signaling pathway and T

cell differentiation were activated in the *Il7r*^{hi} ILC1 cluster (Figure 5, F-H; Supplemental Figure 7,

E-G). Module scores, calculated based on the expression of feature genes within the *Junb*^{hi} cNK

cell cluster, revealed a comparable *Junb*^{hi} signature expression pattern within the *Il7r*^{hi} ILC1

cluster (Figure 5I). Several ILC3 signature genes, such as *Rora, Tmem176a,* and *Tmem176b* (44),

highly expressed in this cluster (Supplemental Figure 7D). Considering the close relationship

between IL-17 mediated immunity response and ILC3 (1, 45), it is plausible that *Il7r*^{hi} ILC1

cluster may be attributed, at least in part, to potential plasticity between ILC1 and ILC3 subsets.

The second liver ILC1 cluster, characterized by high expression of Ly49E (*Klra5*) and Ly49G

(*Klra7*), was designated as the *Klra*hi ILC1 cluster (Figure 5D; Supplemental Figure 7D). Notably,

there was an elevated proportion of *Klra*hi ILC1s in *Prdm1*ΔNcr1 ILC1s (39.7%) compared to

Prdm1+/+ ILC1s (28.1%) (Figure 5C). Liver Ly49E+ ILC1s have been identified as possessing

greater cytotoxic potential and a more robust viral response compared to liver Ly49E- ILC1s (46).

The *Klra*hi cluster exhibited notably high expression of *Ccl5* (Supplemental Figure 7D). Previous

research has underscored the pivotal role of CCL5, produced by both cNK cells and ILC1s, in

facilitating the accumulation of DCs within the tumor microenvironment, thereby impeding tumor

immune evasion, as highlighted in studies (47, 48). The expression of *Ccl5* was reduced in the

*Klra*hi cluster of *Prdm1*ΔNcr1 ILC1s compared to *Prdm1*+/+ ILC1s (Supplemental Figure 7D), which

could potentially have a detrimental impact on the ability of *Klra*hi ILC1s to develop a connection

between innate and adaptive immune responses.

The *Gzmb*hi ILC1 cluster was identified according to the high expression of *Gzma*, *Gzmb*, and

Fgl2 (Supplemental Figure 7D) compared to other ILC1 clusters. GO analysis also revealed the

enrichment in cytolysis and stimulus-response capacity (Figure 5H) of the *Gzmb*hi ILC1 cluster.

Consistent with the *Prf1*hi cNK cell cluster, the proportion of the *Gzmb*hi cluster among liver ILC1s

exhibited a considerable reduction in *Prdm1*ΔNcr1 mice compared to *Prdm1*+/+ mice, and the

expression of *Gzma* also downregulated in *Prdm1*ΔNcr1 mice (Figure 5B). Previous reports showed

that GzmA+ ILC1 constituted the main population of liver ILC1s at birth, with the potential

target-killing ability (49, 50). Within cNK cells, *Il12rb2*, *Il18r1* and *Il18rap* was highly expressed

in *Prf1*hi and *Cxcr3*hi cNK clusters (Supplemental Figure 6I), indicating the IL-18 receptor

expression correlated with the NK cell maturation. While in ILC1, these receptors mostly

expressed on $Il7r^{hi}$ and $Gzmb^{hi}$ ILC1 clusters (Supplemental Figure 7D). Significant decreased of

$Il18r1$ expression in $Prdm1^{\Delta Ncr1}$ cNK cells and ILC1s may associated with the impaired ability to

produce IFN-γ.

To investigate the universal transcriptional program between group 1 ILCs across liver and

spleen, we have explored DEGs in $Prdm1^{\Delta Ncr1}$ liver cNK cells and ILC1s using our scRNA-seq

data. Compared with liver ILC1s, more valuable DEGs was observed in liver cNK cells, including

Junb, *Kit*, *Tcf7*, *Gzmb*, *Prf1*, etc. (Supplemental Figure 8, A and B). Through the integration of the

bulk RNA-seq and scRNA-seq data, we identified 17 DEGs that are regulated by Prdm1 among

liver cNK cells, splenic cNK cells, and liver ILC1s. *Batf3*, *Junb*, *Tcf7*, and *Nfkb1* was upregulated,

whereas *Gzmb*, *Prf1*, and *Fgl2* downregulated, in both $Prdm1^{\Delta Ncr1}$ liver and splenic cNK cells

(Figure 5J). Cxcr6 was downregulated in liver ILC1s and splenic cNK cells in $Prdm1^{\Delta Ncr1}$ mice

(Figure 5J). Previous research found that spleen NK cells could be divided into three distinct

groups based on their expression levels of CD27, CD62L, CD49a, and CD49b (51).

$CD27^{+}CD62L^{-}$ NK cells have remarkable high expression of Batf3, while it was only barely

expressed in $CD27^{+}CD62L^{+}$ and $CD27^{-}CD62L^{+}$ NK cells (51). Based the sequencing data

published by Flommersfeld et al., (GSE180978), a notable negative correlation was observed

between the expression levels of Prdm1 and Batf3 (Supplemental Figure 8I). On top of that, our

findings unveiled the negative regulatory influence of Prdm1 on Batf3 within both spleen and liver

NK cells. This discovery highlights a potential upstream mechanism that may influence the

hemostasis of the spleen NK cell subpopulations through Batf3.

We also compared the gene expression patterns between *Prdm1* and *Hobit* (homologue of

Blimp1) with two published scRNA-seq data (50, 52). Following the knockout of Hobit, the DEGs

were primarily identified within ILC1s. Conversely, after the knockout of Prdm1, a greater

number of DEGs were observed in cNK cells. This indicates that Prdm1 likely possesses a broader

range of target genes within cNK cells, whereas Hobit appears to have a more pronounced impact

on gene expression within ILC1s (Supplemental Figure 8, C-F). There are some overlaps between

the downstream transcriptional profile of *Prdm1* and *Hobit* in liver cNK cells and ILC1s

(Supplemental Figure 8, G and H), such as *Junb*, *Fosb*, *Tcf7*, *Kit*, *Gzmb*, *Prf1*, and *Cxcr6* was

simultaneously upregulated or downregulated in both *Prdm1*$^{\Delta Ncr1}$ and *Hobit*^{KO} liver cNK cells or

ILC1s, indicating the similar regulatory networks of Prdm1 and Hobit.

Prdm1 facilitates the intercellular communication between liver group 1 ILCs and

macrophages

The reciprocal crosstalk between group 1 ILCs and macrophages plays a critical role in

maintaining liver immune homeostasis and anti-cancer immune surveillance (53, 54). The scRNA

sequencing analysis identified two well-established subpopulations of liver macrophages: the

resident Kupffer Cells (KCs) and the Monocyte-Derived Macrophages (MDMs) (Figure 6, A-C;

Supplemental Figure 9A). When comparing the total proportion of macrophages within the

immune cell population of the liver between WT and *Prdm1*$^{\Delta Ncr1}$ mice, there is an increase in

Prdm1$^{\Delta Ncr1}$ mice (Figure 6C). To confirm these findings, we utilized flow cytometry to define

macrophages, including both KCs and MDMs, gating by CD45$^+$Ly6G$^-$F4/80$^+$CD11b$^+$ (Figure 6D).

Our analysis showed that, following the deletion of *Prdm1* in Group 1 ILCs, there is a significant

increase in both the proportion and number of macrophages in the liver (Figure 6D).

According to the transcriptional profile, liver macrophages further clustered and were labeled

as "$Ly6c2^{hi}$"; "$Cxcl2^{hi}$"; "$Ear2^{hi}$" MDMs, and "$Mrc1^{hi}$"; "$C1q^{hi}$" KCs (Figure 6A, Supplemental

Figure 9, A-E). Increased proportion of MDMs and KCs was observed in $Prdm1^{\Delta Ncr1}$ cells

(Supplemental Figure 9B). Within MDMs clusters, $Ly6c2^{hi}$ MDMs mainly compose of $Prdm1^{+/+}$

cells, while $Prdm1^{\Delta Ncr1}$ cells concentrated in $Cxcl2^{hi}$ cluster (Figure 6C). The scRNA-seq data

reveal that following Prdm1 knockout in NKp46$^+$ cells, there is a decrease in the proportion of

KCs within the macrophage population, while the proportion of MDMs increases (Figure 6D).

CX3CR1, a chemokine receptor, is extensively utilized to distinguish KCs and MDMs within

macrophages. Cells expressing CX3CR1 are identified as MDMs, whereas those without CX3CR1

expression are categorized as KCs (55). Employing flow cytometry and leveraging CX3CR1

expression, we assessed the ratios of KCs and MDMs. However, diverging from the scRNA-seq

findings, flow cytometry indicates that post-Prdm1 knockout in group 1 ILCs, there is a minor

increase in the proportion of KCs within the total liver macrophages, and a decrease in the

proportion of MDMs (Figure 6D; Supplemental Figure 9B). This discrepancy could stem from the

different bases of classification: scRNA-seq defines KCs based on gene expression profiles,

whereas flow cytometry differentiates between KCs and MDMs using the single surface marker,

CX3CR1. Analysis of the macrophage subsets identified by scRNA-seq reveals that, while MDM

clusters generally show high CX3CR1 expression, there exists a subset within MDMs, labeled

$Mrc1^{hi}$, that also exhibits high levels of CX3CR1 (Supplemental Figure 9C). Consequently, if flow

cytometry solely employs CX3CR1 for differentiating between KCs and MDMs, it could result in

disparities when compared to scRNA-seq outcomes. Both KCs and MDMs has significantly

increased in $Prdm1^{\Delta Ncr1}$ mice, which was consist with the scRNA-seq data (Supplemental Figure 9,

B and F). Despite the decrease in the proportion of *Ly6c2*^{hi} MDMs in *Prdm1*$^{\Delta Ncr1}$ mice, the

expression levels of *Ly6c2* exhibited minimal variation between WT and *Prdm1*$^{\Delta Ncr1}$ mice

(Supplemental Figure 9D). Intriguingly, within certain cellular subsets, notably the *Ear2*^{hi} cluster,

the *Ly6c2* expression levels in KO mice were found to be higher than those in WT mice.

Additionally, we employed flow cytometry to examine Ly6C expression within the macrophages.

Similar with the scRNA-seq findings, there were no notable differences in Ly6C expression levels

between WT and KO mice (Figure 6E; Supplemental Figure 9G).

High-resolution interactions among liver cNK cells, ILC1s, and macrophages were

established and compared between *Prdm1*$^{+/+}$ and *Prdm1*$^{\Delta Ncr1}$ mice using the CellChat program

(56). Interactions between ILC1s and total macrophages were higher than that between cNK cells

and macrophages (Supplemental Figure 10, A, C, E, and G). Cross-talk between liver group 1

ILCs with macrophages enriched in macrophage migration inhibitory factor (MIF), MHC-I, CXC

chemokine ligand (CXCL), Thy-1 cell surface antigen (THY1), and C-type lectin (CLEC)

pathways (Supplemental Figure 10, B, D, F, and H). Although the quantity of macrophages

significantly increases in *Prdm1*$^{\Delta Ncr1}$ mice, there is a significant decrease in the interaction number

and interaction strength between liver group 1 ILCs and macrophages (~1.5 fold) in *Prdm1*$^{\Delta Ncr1}$

mice (Figure 6, F and G). The reduction of interaction mostly occurred in the cross-talk of

ILC1-MDM and ILC1-KC, whereas no difference was observed in cNK-MDM and cNK-KC

interaction (Supplemental Figure 10, A-H). A reduction in the interaction of ligand-receptor, such

as Mif-CD74, Cxcl16-Cxcr6, and Cxcl10-Cxcr3 was observed in *Prdm1*$^{\Delta Ncr1}$ mice compared to

Prdm1$^{+/+}$ mice (Supplemental Figure 11). Compared to *Prdm1*$^{+/+}$ mice, the information flow of

CXCL and MIF pathways significantly decreased in *Prdm1*$^{\Delta Ncr1}$ mice (Figure 6, H and I;

 Supplemental Figure 10, B, D, F, and H). These pathways play a crucial role in facilitating

macrophage migration. The CXCL signaling was sent from $Ly6c2^{hi}$ $Cxcl2^{hi}$ MDMs and $C1q^{hi}$ KC,

targeting all ILC1 clusters and $Cxcr3^{hi}$ cNK cell clusters (Figure 6J). Of note, although the

population of $Cxcl2^{hi}$ macrophage primarily comprised cells from $Prdm1^{\Delta Ncr1}$ mice, the interaction

within the CXCL pathway between macrophages and group 1 ILCs was obviously less than

$Prdm1^{+/+}$ sample (Figure 6J). These changes could be linked to a decreased population of ILC1s

and $Cxcr3^{hi}$ cNK cell cluster in $Prdm1^{\Delta Ncr1}$ mice, implying that the homeostasis of $Cxcl2^{hi}$

macrophages required sufficient signals from cNK cells and ILC1s. The impaired CXCL-CXCR

interactions might subsequently lead to reduced recruitment and activation of group 1 ILCs and

macrophages within the tumor microenvironment.

Prdm1 safeguards group 1 ILCs from exhaustion-like phenotypes in the tumor

microenvironment

The suppression of mitochondrial related pathways in $Junb^{hi}$ cNK cell cluster, along with a

significant increase of this cNK cell cluster in $Prdm1^{\Delta Ncr1}$ mice, encouraged us to explore

mitochondrial   function   through   flow   cytometry.   MitoTracker,   MitoSOX,   and

Tetramethylrhodamine methyl ester (TMRM) were used to assess the mitochondrial mass,

superoxide production, and mitochondrial membrane potential. A substantial decrease in MFIs of

MitoTracker was observed in $Prdm1^{\Delta Ncr1}$ splenic cNK cells, liver cNK cells, and liver ILC1s when

compared to their $Prdm1^{+/+}$ counterparts (Figure 7A). This observation aligns with the enrichment

of downregulated genes from $Prdm1$ deficient sample in mitochondrial related pathway, as

revealed by RNA sequencing data (Supplemental Figure 6, D-F). There was no significant

difference in MitoSOX and TMRM between $Prdm1^{\Delta Ncr1}$ and $Prdm1^{+/+}$ mice (Figure 7, B and C),

which suggested that the ATP synthesize capacity was minimally affected by $Prdm1$.

IFN-γ is a critical cytokine for NK cells mediated cancer surveillance(57, 58) and impaired

production of IFN-γ was considered as a key hallmark of exhausted NK cells (59, 60). To evaluate

the IFN-γ secreting capacity of liver cNK cells and ILC1s in tumor microenvironment, B16F10

tumor cells were inoculated to the liver via splenic injection and the IFN-γ levels in response to

stimulation of IL-12 and/or IL-18 were assessed by flow cytometry (Supplemental Figure 12A).

The proportion changes of cNK cells and ILC1s in $Prdm1^{\Delta Ncr1}$ mice was similar with the no

tumor-burden condition, while the number of both cNK cells and ILC1s have significant

decreased in tumor-bearing liver (Figure 7D). Compared with $Prdm1^{+/+}$ mice, significant deceased

of IFN-γ were observed in $Prdm1^{\Delta Ncr1}$ mice liver cNK cells and ILC1s under the combinate

stimulation of IL-12/IL-18 (Figure 7E; Supplemental Figure 12B), which was more remarkable in

liver ILC1s. Similar trends were observed when IL-12 or IL-18 was used alone, although only

liver ILC1s showed a significant decrease in response to IL-18 stimulation (Figure 7E). These

findings were consistent with the heavy tumor burden observed in $Prdm1^{\Delta Ncr1}$ mice.

Discussion

Prdm1 is a pivotal transcription factor that has attracted substantial research interest due to its

role in lymphocytes. In a study involving systemic knockout combined with competitive

transplantation, it was found that Prdm1 promotes NK cell maturation and the expression of Gzmb.

On the contrary, the same study also found that NK cells with Prdm1 deficiency exhibit

heightened proliferation, increased survival, enhanced migratory abilities towards tumors, and

greater cytotoxicity against subcutaneously implanted RMAS tumors (30). Using Ncr1-driven

conditional knockout transgenic mice, which specifically delete *Prdm1* in group 1 ILCs, we not

only validated *Prdm1*'s positive regulation of NK cell maturation, but also demonstrated its

indispensable role in NK cell anti-tumor activity. The compromised mitochondrial function and

reduced IFN-γ, granzyme B, and perforin production appear to be potential contributing factors to

Prdm1-mediated cancer surveillance. Reduction of CX3CR1$^+$ NK cells in multiple tissues, and

decreased expression of *Cx3cr1* and *Cxcr6* was observed in *Prdm1*^{ΔNcr1} splenic cNK cells (Figure

3F), both of which are essential for NK cells egressing from bone marrow (61, 62). Our results not

only confirmed a decrease in cytotoxic molecules in *Prdm1*-deficient NK cells (30) but also

showed that the reduction in Gzmb and perforin is not solely attributable to the diminished

maturation of these cells. Mature NK cells in bone marrow obtained the expression of CX3CR1

and acquired ability to enter the circulation (61, 62). The quantity of cNK cells increased

exclusively in the bone marrow, with reductions observed in all other tissues (Figure 1G),

indicating *Prdm1* might regulate chemokine receptors to facilitate the egression of cNK cells from

the bone marrow to peripheral tissues. In addition, higher expression of *Cxcr6* compared to cNK

cells is also a key factor for liver tissue residence of ILC1s (63). Decreased expression of *Cxcr6* in

Prdm1 deficient group 1 ILCs may also contribute to the balance shift towards cNK cells.

Furthermore, although both liver NK cells and liver ILC1s require Prdm1 to maintain their

quantity, liver ILC1s demonstrate a more pronounced dependency on Prdm1. However, it is

currently widely believed that liver NK cells and liver ILC1s originate from different progenitors.

It is worth noting that while we observed changes in the NK and ILC1 proportions after *Prdm1*

knockout, our data does not support the hypothesis that Prdm1 affects progenitor differentiation

decisions, thereby influencing the fate selection of NK and ILC1. Further research may be needed

to elucidate how Prdm1 regulates the balance between NK cells and ILC1s.

scRNA sequencing analysis reveals that both liver cNK cells and ILC1s can be further

divided into three subgroups based on their gene expression patterns. Junb is a crucial

transcriptional factor for the cytotoxic function of CD8$^+$ T cells and NK cells. However, excessive

Junb expression has been found to promote T cell exhaustion (64). Our previous study showed

that as NK cells mature, the expression level of *Prdm1* increased while the expression level of

Junb gradually decreased (25). The current study demonstrated that in NK cells, the expression

level of *Junb* significantly increases upon the deletion of Prdm1, indicating that *Junb* expression is

suppressed by Prdm1. As *Junb* expression decreases with NK cell maturation, and it is inhibited

by the gradually increasing Prdm1 during maturation. This implies that constraining *Junb*

expression is likely a fundamental prerequisite for NK cell maturation. However, the precise

mechanism by which Prdm1 downregulates *Junb* in NK cells still needs further research.

Furthermore, *Junb*^{hi} NK cells exhibit lower expression levels of cytotoxic genes and reduced

mitochondrial-related signaling pathways. Mitochondria are pivotal organelles crucial for cellular

metabolism. Disruptions in mitochondrial function have been linked to T Cell exhaustion,

attributed to glycolytic reprogramming (65). Similarly, mitochondrial fragmentation has been

closely associated with NK cell exhaustion (66). However, the concept of NK cell exhaustion isn't

as firmly established as it is for T cells. Exhausted NK cells should primarily exhibit diminished

functions. This is characterized by a diminished ability to destroy tumor cells, a reduced capability

to activate other components of the immune system, and compromised proliferation and survival

rates. Additionally, this reduced functionality is associated with a decline in the expression of

molecules responsible for cytotoxic activity, lower production of IFN-γ, and metabolic

disturbances that may arise from mitochondrial dysfunction. While our current data is not

sufficient to definitively classify these cells as exhausted NK cells, it supports that a subpopulation,

referred to $Junb^{hi}$ cluster, demonstrates an exhaustion-like phenotype. The significant increase in

this cell population following $Prdm1$ knockout in NK cells may potentially be one of the reasons

why $Prdm1^{\Delta Ncr1}$ mice lose their tumor-killing capacity. Whether the excessive expression of JunB

in NK cells is also a contributing factor to their exhaustion, similar to T cells(64), requires further

investigation.

The scRNA-seq data revealed that $Prdm1$ plays distinct roles in regulating cNK cells and

ILC1s despite being required for the quantity of both lineages. Specifically, $Prdm1$ appears to be

more involved in promoting the resistance against exhaustion in cNK cells, whereas in ILC1s, it

may play a role in the plasticity between ILC1s and ILC3s. In both our previous study and a study

by Colonna et al. (25, 67), it was demonstrated that $Smad4$, a transcriptional factor involved in

TGF-β signal pathway, upregulated $Prdm1$ in NK cells and depletion of $Smad4$ resulted in a

decreased ratio of NK cells to ILC1s in the liver. However, knocking out $Prdm1$ in Ncr1$^+$ cells

increased the ratio of NK cells to ILC1s in liver group 1 ILCs (Figure 2A). These findings suggest

the possibility of a Smad4-independent pathway through which $Prdm1$ promotes the maintenance

of ILC1s, or that $Prdm1$ plays a more significant role in maintaining ILC1s compared to its role in

NK cells.

Previous studies have identified Hobit and Prdm1 as central regulators instructing

tissue-dependent programs and retention of diverse tissue-resident lymphocytes (17, 50, 52). Liver

ILC1s required Hobit, but not necessary for cNK cells (6). Expression of Prdm1 was remarkably

higher in cNK cells versus ILC1s (17). While in our study, cNK cells and liver ILC1s reduced

simultaneously in *Prdm1^{ΔNcr1}* mice, and even more significant in ILC1s. This indicates that while

Prdm1 is expressed at lower levels in ILC1s, its role in preserving the quantity of ILC1s may be

more crucial. Thus, Prdm1 and Hobit may have parallel program in instructing ILC1s functional

development and maturation. Prdm1 and Hobit directly bound and repressed *Tcf7* (17), which

encoded TCF-1, a TF binding and limiting the activity of Gzmb regulatory element (68). Gzmb

has been demonstrated directly bound and activated by Junb in NK cells, which suggested Gzmb

expression regulated by multiple Prdm1/Hobit downstream signals (25). In human T cells, binding

motif of JUNB was enriched in the binding sites of PRDM1 (69), indicating the essential role of

PRDM1-JUNB axis during NK cell and T cell development. In NK cells deficient in Prdm1

expression, we noted a decrease in Gzmb levels alongside with an elevation in Junb expression.

This indicates that Prdm1 not only facilitates the expression of Gzmb in NK cells but also

suppresses Junb expression. Given that Junb is recognized as a positive regulator of Gzmb (70),

this presents a complex interplay that seems contradictory. Therefore, it is imperative to develop a

theoretical framework to comprehensively understand and interpret this paradoxical relationship.

Our previous research has demonstrated that with the progression of NK cell maturation, there is a

gradual increase in Prdm1 expression, accompanied by a subtle reduction in JunB levels (25).

Overexpression of JunB can induce lymphocyte exhaustion (64). Based on these findings, we

hypothesize that during the initial stages of NK cell development, JunB might enhance the

expression of certain molecules associated with cytotoxicity, thereby aiding NK cells in acquiring

the capacity to eliminate target cells. During these initial stages, Prdm1 level are comparatively

low, thus exerting a weaker inhibitory effect on JunB. As NK cells proceed to maturity, there's a

progressive increase in Prdm1 level, which then exerts a stronger inhibitory influence on JunB,

contributing to the reduced JunB level in terminal matured NK cells. Consequently, this reduces

the potential for NK cell exhaustion caused by elevated levels of JunB.

Chronic inflammation is a crucial factor in promoting tumorigenesis, and macrophages play a

significant role in this process. Macrophages interact with both cNK cells and ILC1s. However,

the TFs that regulate these interactions are poorly understood. Fortunately, recent advances in

scRNA-seq technology and the CellChat software tool (56) have allowed us to gain a better

understanding of the *Prdm1* signaling pathway in group 1 ILCs and its impact on macrophages at

the transcriptional level. Increased metastasis in *Prdm1$^{\Delta Ncr1}$* mice may be due to a decrease in the

killing ability of NK cells, making them more prone to exhaustion, or it could be due to

abnormalities in group 1 ILCs, leading to a decrease in the anti-tumor abilities of macrophages

and an enhancement of their pro-tumor capabilities. It is worth noting that normal

ILC1-macrophage interactions are more prevalent than the interaction between NK cells and

macrophages (Supplemental Figure 10). We also found that CXCL signaling based interaction

remarkably diminished in *Prdm1$^{\Delta Ncr1}$* mice, which suggested CXCL-CXCR may contribute to

keep the sufficient interaction between group 1 ILC and macrophage. Specifically, *Prdm1* in group

1ILCs may be critical in preventing the overactivation of macrophages that can lead to cancer

development. Increased population of both MDMs and KCs in *Prdm1$^{\Delta Ncr1}$* mice, as well as

different distribution of macrophage clusters, indicating the homeostasis of macrophages require

environment with functional cNK cells and ILC1s. Although not all macrophage phenotypes have

been verified in this study, the present research serves primarily to offer initial insights and

preliminary data for investigating the interactions between group 1 ILCs and macrophages. It aims

to inspire further research into the role of transcription factors within the liver and the cancer

microenvironment.

While our findings underscore the importance of Prdm1 in liver cNK cells and ILC1s tumor

immune surveillance, it does not be validated in human NK cells, whereas previous studies have

found that PRDM1 might inhibit the proliferation and function of human NK cells (32, 71).

Furthermore, we not provided an in-depth evaluation in multiple tumor models. Further research

may provide deeper insight into the role of PRDM1 in the anti-tumor function of human NK cells,

enabling a more direct investigation of its application in cancer therapies. Given its important role

in preserving liver cNK cells and ILC1s functional heterogeneity, enhancing Prdm1 function in

human NK cells could potentially be a strategy to promote NK cell-based immunotherapy for

cancer.

Methods

Mice

Prdm1$^{fl/fl}$ mice were purchased from The Jackson Laboratory. *Ncr1-iCre* and *B2m$^{-/-}$* mice

were purchased from Shanghai Model Organisms Center, Inc.. Six- to twelve-week-old littermates

were used for the experiment.

Experimental metastasis model

For lung metastasis model, 0.3×10^6 B16F10 cells were intravenous injected into mice.

Three weeks later, mice were euthanized for analysis. For liver metastasis model, mice were

inoculation with 0.5×10^6 B16F10 via intrasplenic injection. Three weeks later, mice were

euthanized for analysis. Lung and liver from tumor-bearing mice were fixed in 10% formalin and

embedded in paraffin. Sections were stained with H&E.

***In vivo* cytotoxicity assay**

Donor splenocytes harvested from B2m deficient ($B2m^{-/-}$) mice were labeled with 5 μM

CFDA-SE. Donor splenocytes harvested from B2m-adequate ($B2m^{+/+}$) mice were labeled with 5

μM eF670. Labeled splenocytes from two mouse strains were mixed in a 1:1 ratio, and 1×10^{7}

cells in total were injected i.v. into $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice. One day after administration,

spleen and liver cells were isolated from recipient mice, and the population of labeled cells was

analyzed by flow cytometry. Rejection % was quantified according to the following formula:

$1-(\%B2m^{+/+}/\%B2m^{-/-})_{Prdm1^{+/+} \text{ or } Prdm1^{\Delta Ncr1}} \times 100\%$.

Cell isolation

Mice were perfused with 1◊ PBS by portal vein puncture before harvesting tissues. Liver and

lung was digested with 0.05% collagenase II for 30 minutes and filtered through 70 μm cell

strainers, and mononuclear cells were isolated after subjected to density gradient using 30% and

70% percoll. Spleen were also removed and pressed through 70 μm filterers to obtain splenocytes.

Peripheral blood mononuclear cells were obtained from peripheral blood after lysis of red blood

cells (Biolegend, 420301). Flushing femurs and mechanical disruption of inguinal lymph nodes

were performed to obtain cells from bone marrow and lymph nodes.

Real-time RT-PCR

RNA was extracted from FACS-sorted NK cells or splenocytes using RNASimple Total RNA

Kit (TIANGEN Biotech, 4992858) and subsequently reverse transcribed to cDNA with

SuperScript VILO Master Mix (Thermo Fisher Scientific, 11755050) according to

manufacturer's instructions. qPCR was performed with SYBR Green Mix (Thermo Fisher

Scientific, A25742) and CFX Opus 96 Real-Time PCR System (Bio-Rad). The relative mRNA

expression level was calculated using 2^{-ddCt} method. Primer sequences: *Prdm1*:

5'-CAGAAACACTACTTGGTACA-3'; 5'-GATTGCTTGTGCTGCTAA-3'.

Flow cytometry

Flow cytometry and cell sorting were performed with a Cytoflex S/SRT (Beckman Coulter).

The following antibodies were used (all purchased from BioLegend unless otherwise indicated):

CD45-PE-Cy7 (catalog 103114, clone 30-F11); CD3ε-PerCP-Cy5.5 (catalog 100327, clone

145-2C11); NK1.1-BV421 (catalog 108741, clone PK136); CD335-AF647 (catalog 560755, clone

29A1.4, BD Bioscience); CD49a-PE (catalog 562115, clone Ha31/8, BD Bioscience);

CD49b-FITC (catalog 108906, clone DX5); CD27-BV510 (catalog 124229, clone LG.3A10);

CD11b-AF700 (catalog 101222, clone M1/70); KLRG1-APC (catalog 561620, clone 2F1, BD

Bioscience); CD49a-BV421 (catalog 740046, clone Ha31/8, BD Bioscience); IFN-γ-PE (catalog

505807, clone XMG1.2); Granzyme B (catalog 372207, clone QA16A02); Perforin (catalog

154305, clone S16009A); CD49b-APC-Cy7 (catalog 108919, clone DX5) ; IgG-PE (catalog

402203, clone 27-35); CD335-BV510 (catalog 137623, clone 29A1.4);; CD3ε-APC-Cy7 (catalog

100330, clone 145-2C11); CD11b-BV421 (catalog 101235, clone M1/70); CD3ε-BV421 (catalog

100335, clone 145-2C11); CD3ε-FITC (catalog 553061, clone 145-2C11, BD Bioscience);

CX3CR1-PE (catalog 149005, clone SA011F11); Ly6G-PerCP-Cy5.5 (catalog 127615, clone

1A8); Ly6C-BV510 (catalog 108437, clone RB6-8C5); F4/80-APC-Cy7 (catalog 123118, clone

BM8). For mitochondrial metabolic assay, fresh cells were incubated in 37□ media for 30min

with 100 nM MitoTracker Green (catalog M7514, Invitrogen), 100 nM TMRM (catalog T668;

Invitrogen), and 10 uM MitoSOX Red (catalog M36008; Invitrogen), respectively. Surface-stained

after washing with PBS and then detected by flow cytometry. For intracellular IFN-γ staining,

Cells freshly obtained from liver and spleen were stimulated 12 hours with or without cytokine.

GolgiStop (BD Biosciences) was added 4 hours before intracellular staining.

Bulk RNA sequencing

Total RNA from FACS sorted splenic NK cells of $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice was

extracted by TRIzol reagent (Invitrogen), then reverse transcribed into cDNA. Library

construction was prepared using Illumina mRNA Library kit, and sequencing was performed by

the BGISEQ-500. Standard methods were used to analyze the RNA-seq data, including aligning

the reads to the genome by HISAT2 (v2.1.0) (72), and gene expression values (Counts) were

calculated using RSEM (v1.3.1) (73). DEGs were identified using DEseq2 (v1.4.5) (74) with a

cutoff of $\log_2$(fold change) > 0.5 and P < 0.05. The "clusterProfiler" package (v4.4.4) (75) and

gene sets from molecular signatures database (MSigDB) were used for GSEA and GO analysis.

The heatmap was plotted using the "Pheatmap" package (v1.0.12).

Single-cell RNA sequencing

FACS-sorted liver CD45$^+$ cells with more than 80% cell viability were used for library

preparation. Each sample contained cells from three *Prdm1$^{+/+}$* or *Prdm1$^{\Delta Ncr1}$* mice. Gel

Bead-in-Emulsions (GEMs) were generated using the 10X Genomics Chromium system, which

combinates Master Mix, Single Cell 3' v3.1 Gel Beads, and Partitioning Oil with single cells.

GEMs were mixed with cell lysate and reverse transcription reagent to produce full-length cDNA.

After incubation, the GEMs were broken, and recovered cDNA were amplified via PCR.

Fragmentation, End repair, A-tailing, and Adaptor Ligation were performed to obtain final

libraries, which contain P5 and P7 sequences. The 3' library was sequenced on Novaseq 6000 with

approximately 50k read pairs/cell sequencing depth. The "Seurat" R package (v4.2.0) (76) was

used for data analysis. Initial quality control was performed to filter out the low-quality cells and

cell doublets. Cells with 200-5500 expressed genes and no more than 10% mitochondrial genes

were considered for high-quality. Doublets were filtered with the "scDblFinder" Package (v1.10.0)

(77). After quality control, we totally recovered 6161 cells and 4817 cells from *Prdm1$^{+/+}$* and

Prdm1$^{\Delta Ncr1}$ mice, respectively. Principal component analysis (PCA) was used for cluster analysis.

The first 15 PCs were picked for clustering and further visualized by UMAP. Clusters-specific

marker was defined using the "FindAllMarkers" function, and clusters were manually annotated

based on the top 30 or 15 markers. The "clusterProfiler" package and gene set from MSigDB were

used for GSEA and GO analysis. "CellChat" package (v1.4.0) (56) was utilized to predict the

cell-to-cell communication from scRNA-seq data.

TCGA datasets assay

The normalized gene expression and survival datasets of cancer patients collected in The

Cancer Genome Atlas (TCGA) were downloaded from UCSC Xena (http://xena.ucsc.edu/) (78).

NK cell-associated genes including *CD160*, *CD244*, *CTSW*, *FASLG*, *GZMA*, *GZMB*, *GZMH*,

IL18RAP, *IL2RB*, *KIR2DL4*, *KLRB1*, *KLRC3*, *KLRD1*, *KLRF1*, *KLRK1*, *NCR1*, *NKG7*, *PRF1*,

XCL1, *XCL2,* according to the previous study (33). NK cell-associated genes, together with

PRDM1, constitute the *NK-PRDM1* signature in this study. The mean expression of per genes was

ordered from high-to-low and plotted by heatmap using the "Pheatmap" package. The overall

survival of patients in the high and low expression of *NK-PRDM1* signature was selected for

analysis. Kaplan-Meier curves were plotted by GraphPad Prism.

Statistics

For experiment results, two-tailed t tests were used to measure the continuous and normally

distributed between the two independent groups. Paired t-tests were used to determine the

statistical significance between two paired groups. Log-rank tests were used to compare the

overall survival distribution between the two groups of patients. A P value less than 0.05 was

considered significant and data were presented as mean ± SEM.

Study approval

All animal experiments were approved by The Tianjin University Animal Care and Use

Committee. No human subjects were performed in this study.

Author contributions

734   Jitian He, and Youwei Wang designed experiments, performed experiments, analyzed the data

735 and wrote the manuscript. Jitian He, Le Gao, Peiying Wang, Yiran Zheng, and Yumo Zhang

736 performed experiments. Wing Keung Chan, Jiming Wang, Huaiyong Chen, and Zhouxin Yang

737 analyzed the data, interpreted results and reviewed the manuscript.

Acknowledgments

This work was supported by National Key Research and Development Plan of China (2022YFF1202901); National Natural Science Foundation of China (82372801); and The Zhejiang Provincial Natural Science Foundation of China (LY21H150002).

Declaration of interests

745 The authors have declared that no conflict of interest exists.

References

1. Spits H, Artis D, Colonna M, Diefenbach A, Di Santo JP, Eberl G, et al. Innate lymphoid cells--a proposal for uniform nomenclature. *Nat Rev Immunol.* 2013;13(2):145-9.

2. Peng H, Jiang X, Chen Y, Sojka DK, Wei H, Gao X, et al. Liver-resident NK cells confer adaptive immunity in skin-contact inflammation. *J Clin Invest.* 2013;123(4):1444-56.

3. Bai L, Vienne M, Tang L, Kerdiles Y, Etiennot M, Escaliere B, et al. Liver type 1 innate lymphoid cells develop locally via an interferon-gamma-dependent loop. *Science.* 2021;371(6536).

4. Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C, et al. ILC1 Confer Early Host Protection at Initial Sites of Viral Infection. *Cell.* 2017;171(4):795-808 e12.

5. Schuster IS, Wikstrom ME, Brizard G, Coudert JD, Estcourt MJ, Manzur M, et al. TRAIL+ NK cells control CD4+ T cell responses during chronic viral infection to limit autoimmunity. *Immunity.* 2014;41(4):646-56.

6. Ducimetiere L, Lucchiari G, Litscher G, Nater M, Heeb L, Nunez NG, et al. Conventional NK cells and tissue-resident ILC1s join forces to control liver metastasis. *Proc Natl Acad Sci U S A.* 2021;118(27).

7. Fumagalli V, Venzin V, Di Lucia P, Moalli F, Ficht X, Ambrosi G, et al. Group 1 ILCs regulate T cell-mediated liver immunopathology by controlling local IL-2 availability. *Science immunology.* 2022;7(68):eabi6112.

8. Kiessling R, Klein E, and Wigzell H. "Natural" killer cells in the mouse. I. Cytotoxic cells with specificity for mouse Moloney leukemia cells. Specificity and distribution according to genotype. *European journal of immunology.* 1975;5(2):112-7.

9. Nabekura T, Riggan L, Hildreth AD, O'Sullivan TE, and Shibuya A. Type 1 Innate Lymphoid Cells Protect Mice from Acute Liver Injury via Interferon-gamma Secretion for Upregulating Bcl-xL Expression in Hepatocytes. *Immunity.* 2020;52(1):96-108 e9.

10. Takeda K, Hayakawa Y, Smyth MJ, Kayagaki N, Yamaguchi N, Kakuta S, et al. Involvement of tumor necrosis factor-related apoptosis-inducing ligand in surveillance of tumor metastasis by liver natural killer cells. *Nat Med.* 2001;7(1):94-100.

11. Townsend MJ, Weinmann AS, Matsuda JL, Salomon R, Farnham PJ, Biron CA, et al. T-bet regulates the terminal maturation and homeostasis of NK and Valpha14i NKT cells. *Immunity.* 2004;20(4):477-94.

12. Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, et al. Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature.* 1999;397(6721):702-6.

13. Gascoyne DM, Long E, Veiga-Fernandes H, de Boer J, Williams O, Seddon B, et al. The basic leucine zipper transcription factor E4BP4 is essential for natural killer cell development. *Nature immunology.* 2009;10(10):1118-24.

14. Constantinides MG, McDonald BD, Verhoef PA, and Bendelac A. A committed precursor to innate lymphoid cells. *Nature.* 2014;508(7496):397-401.

15. Gordon SM, Chaix J, Rupp LJ, Wu J, Madera S, Sun JC, et al. The transcription factors T-bet and Eomes control key checkpoints of natural killer cell maturation. *Immunity.* 2012;36(1):55-67.

16. Daussy C, Faure F, Mayol K, Viel S, Gasteiger G, Charrier E, et al. T-bet and Eomes instruct
the development of two distinct natural killer cell lineages in the liver and in the bone marrow.
J Exp Med. 2014;211(3):563-77.

17. Mackay LK, Minnich M, Kragten NA, Liao Y, Nota B, Seillet C, et al. Hobit and Blimp1
instruct a universal transcriptional program of tissue residency in lymphocytes. *Science.*
2016;352(6284):459-63.

18. Lee JC, Green MD, Huppert LA, Chow C, Pierce RH, and Daud AI. The Liver-Immunity
Nexus and Cancer Immunotherapy. *Clin Cancer Res.* 2022;28(1):5-12.

19. Galon J, and Bruni D. Approaches to treat immune hot, altered and cold tumours with
combination immunotherapies. *Nature reviews Drug discovery.* 2019;18(3):197-218.

20. Ruff SM, Shannon AH, Beane JD, and Pawlik TM. Highlighting novel targets in
immunotherapy for liver cancer. *Expert review of gastroenterology & hepatology.*
2022;16(11-12):1029-41.

21. He Y, Zhan L, Shi J, Xiao M, Zuo R, Wang C, et al. The Combination of R848 with Sorafenib
Enhances Antitumor Effects by Reprogramming the Tumor Immune Microenvironment and
Facilitating Vascular Normalization in Hepatocellular Carcinoma. *Adv Sci (Weinh).*
2023:e2207650.

22. Nersesian S, Schwartz SL, Grantham SR, MacLean LK, Lee SN, Pugh-Toole M, et al. NK cell
infiltration is associated with improved overall survival in solid cancers: A systematic review
and meta-analysis. *Transl Oncol.* 2021;14(1):100930.

23. Lee HA, Goh HG, Lee YS, Jung YK, Kim JH, Yim HJ, et al. Natural killer cell activity is a
risk factor for the recurrence risk after curative treatment of hepatocellular carcinoma. *BMC
Gastroenterol.* 2021;21(1):258.

24. Gao Y, Souza-Fonseca-Guimaraes F, Bald T, Ng SS, Young A, Ngiow SF, et al. Tumor
immunoevasion by the conversion of effector NK cells into type 1 innate lymphoid cells.
Nature immunology. 2017;18(9):1004-15.

25. Wang Y, Chu J, Yi P, Dong W, Saultz J, Wang Y, et al. SMAD4 promotes
TGF-beta-independent NK cell homeostasis and maturation and antitumor immunity. *J Clin
Invest.* 2018;128(11):5123-36.

26. Cortez VS, Ulland TK, Cervantes-Barragan L, Bando JK, Robinette ML, Wang Q, et al.
SMAD4 impedes the conversion of NK cells into ILC1-like cells by curtailing non-canonical
TGF-beta signaling. *Nat Immunol.* 2017;18(9):995-1003.

27. Keller AD, and Maniatis T. Identification and characterization of a novel repressor of
beta-interferon gene expression. *Genes Dev.* 1991;5(5):868-79.

28. Kallies A, Hawkins ED, Belz GT, Metcalf D, Hommel M, Corcoran LM, et al. Transcriptional
repressor Blimp-1 is essential for T cell homeostasis and self-tolerance. *Nat Immunol.*
2006;7(5):466-74.

29. Martins GA, Cimmino L, Shapiro-Shelef M, Szabolcs M, Herron A, Magnusdottir E, et al.
Transcriptional repressor Blimp-1 regulates T cell homeostasis and function. *Nat Immunol.*
2006;7(5):457-65.

30. Kallies A, Carotta S, Huntington ND, Bernard NJ, Tarlinton DM, Smyth MJ, et al. A role for
Blimp1 in the transcriptional network controlling natural killer cell maturation. *Blood.*
2011;117(6):1869-79.

31. Akman B, Hu X, Liu X, Hatipoglu T, You H, Chan WC, et al. PRDM1 decreases sensitivity of

human NK cells to IL2-induced cell expansion by directly repressing CD25 (IL2RA). *Journal*
of leukocyte biology. 2021;109(5):901-14.

32. Smith MA, Maurin M, Cho HI, Becknell B, Freud AG, Yu J, et al. PRDM1/Blimp-1 controls
effector cytokine production in human NK cells. *J Immunol*. 2010;185(10):6058-67.

33. Cursons J, Souza-Fonseca-Guimaraes F, Foroutan M, Anderson A, Hollande F,
Hediyeh-Zadeh S, et al. A Gene Signature Predicting Natural Killer Cell Infiltration and
Improved Survival in Melanoma Patients. *Cancer Immunol Res*. 2019;7(7):1162-74.

34. Shapiro-Shelef M, Lin K-I, McHeyzer-Williams LJ, Liao J, McHeyzer-Williams MG, and
Calame K. Blimp-1 Is Required for the Formation of Immunoglobulin Secreting Plasma Cells
and Pre-Plasma Memory B Cells. *Immunity*. 2003;19(4):607-20.

35. Hayakawa Y, and Smyth MJ. CD27 dissects mature NK cells into two subsets with distinct
responsiveness and migratory capacity. *J Immunol*. 2006;176(3):1517-24.

36. Chiossone L, Chaix J, Fuseri N, Roth C, Vivier E, and Walzer T. Maturation of mouse NK
cells is a 4-stage developmental program. *Blood*. 2009;113(22):5488-96.

37. Huntington ND, Tabarias H, Fairfax K, Brady J, Hayakawa Y, Degli-Esposti MA, et al. NK
cell maturation and peripheral homeostasis is associated with KLRG1 up-regulation. *J*
Immunol. 2007;178(8):4764-70.

38. Bix M, Liao NS, Zijlstra M, Loring J, Jaenisch R, and Raulet D. Rejection of class I
MHC-deficient haemopoietic cells by irradiated MHC-matched mice. *Nature*.
1991;349(6307):329-31.

39. Bern MD, Parikh BA, Yang L, Beckman DL, Poursine-Laurent J, and Yokoyama WM.
Inducible down-regulation of MHC class I results in natural killer cell tolerance. *The Journal*
of experimental medicine. 2019;216(1):99-116.

40. Karre K, Ljunggren HG, Piontek G, and Kiessling R. Selective rejection of H-2-deficient
lymphoma variants suggests alternative immune defence strategy. *Nature*.
1986;319(6055):675-8.

41. Viant C, Fenis A, Chicanne G, Payrastre B, Ugolini S, and Vivier E. SHP-1-mediated
inhibitory signals promote responsiveness and anti-tumour functions of natural killer cells.
Nat Commun. 2014;5:5108.

42. Cillo C, Dick JE, Ling V, and Hill RP. Generation of drug-resistant variants in metastatic B16
mouse melanoma cell lines. *Cancer Res*. 1987;47(10):2604-8.

43. Sathe P, Delconte RB, Souza-Fonseca-Guimaraes F, Seillet C, Chopin M, Vandenberg CJ, et al.
Innate immunodeficiency following genetic ablation of Mcl1 in natural killer cells. *Nat*
Commun. 2014;5:4539.

44. Robinette ML, Fuchs A, Cortez VS, Lee JS, Wang Y, Durum SK, et al. Transcriptional
programs define molecular characteristics of innate lymphoid cell classes and subsets. *Nat*
Immunol. 2015;16(3):306-17.

45. Klose CS, Kiss EA, Schwierzeck V, Ebert K, Hoyler T, d'Hargues Y, et al. A T-bet gradient
controls the fate and function of CCR6-RORgammat+ innate lymphoid cells. *Nature*.
2013;494(7436):261-5.

46. Chen Y, Wang X, Hao X, Li B, Tao W, Zhu S, et al. Ly49E separates liver ILC1s into
embryo-derived and postnatal subsets with different functions. *The Journal of experimental*
medicine. 2022;219(5).

47. Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrerizo M, Sammicheli S, et al.

NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting Cancer Immune Control. *Cell.* 2018;172(5):1022-37 e14.

48. Kirchhammer N, Trefny MP, Natoli M, Brucher D, Smith SN, Werner F, et al. NK cells with tissue-resident traits shape response to immunotherapy by inducing adaptive antitumor immunity. *Sci Transl Med.* 2022;14(653):eabm9043.

49. Di Censo C, Marotel M, Mattiola I, Muller L, Scarno G, Pietropaolo G, et al. Granzyme A and CD160 expression delineates ILC1 with graded functions in the mouse liver. *European journal of immunology.* 2021;51(11):2568-75.

50. Friedrich C, Taggenbrock R, Doucet-Ladeveze R, Golda G, Moenius R, Arampatzi P, et al. Effector differentiation downstream of lineage commitment in ILC1s is driven by Hobit across tissues. *Nat Immunol.* 2021;22(10):1256-67.

51. Flommersfeld S, Bottcher JP, Ersching J, Flossdorf M, Meiser P, Pachmayr LO, et al. Fate mapping of single NK cells identifies a type 1 innate lymphoid-like lineage that bridges innate and adaptive recognition of viral infection. *Immunity.* 2021;54(10):2288-304 e7.

52. Yomogida K, Bigley TM, Trsan T, Gilfillan S, Cella M, Yokoyama WM, et al. Hobit confers tissue-dependent programs to type 1 innate lymphoid cells. *Proc Natl Acad Sci U S A.* 2021;118(50).

53. Tu Z, Bozorgzadeh A, Pierce RH, Kurtis J, Crispe IN, and Orloff MS. TLR-dependent cross talk between human Kupffer cells and NK cells. *J Exp Med.* 2008;205(1):233-44.

54. Park MD, Reyes-Torres I, LeBerichel J, Hamon P, LaMarche NM, Hegde S, et al. TREM2 macrophages drive NK cell paucity and dysfunction in lung cancer. *Nat Immunol.* 2023;24(5):792-801.

55. Heymann F, Peusquens J, Ludwig-Portugall I, Kohlhepp M, Ergen C, Niemietz P, et al. Liver inflammation abrogates immunological tolerance induced by Kupffer cells. *Hepatology.* 2015;62(1):279-91.

56. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and analysis of cell-cell communication using CellChat. *Nat Commun.* 2021;12(1):1088.

57. Takeda K, Nakayama M, Sakaki M, Hayakawa Y, Imawari M, Ogasawara K, et al. IFN-gamma production by lung NK cells is critical for the natural resistance to pulmonary metastasis of B16 melanoma in mice. *Journal of leukocyte biology.* 2011;90(4):777-85.

58. Lin Q, Rong L, Jia X, Li R, Yu B, Hu J, et al. IFN-gamma-dependent NK cell activation is essential to metastasis suppression by engineered Salmonella. *Nature communications.* 2021;12(1):2537.

59. Roe K. NK-cell exhaustion, B-cell exhaustion and T-cell exhaustion-the differences and similarities. *Immunology.* 2022;166(2):155-68.

60. Zhang Q, Bi J, Zheng X, Chen Y, Wang H, Wu W, et al. Blockade of the checkpoint receptor TIGIT prevents NK cell exhaustion and elicits potent anti-tumor immunity. *Nature Immunology.* 2018;19(7):723-32.

61. Sciume G, De Angelis G, Benigni G, Ponzetta A, Morrone S, Santoni A, et al. CX3CR1 expression defines 2 KLRG1+ mouse NK-cell subsets with distinct functional properties and positioning in the bone marrow. *Blood.* 2011;117(17):4467-75.

62. Chea S, Possot C, Perchet T, Petit M, Cumano A, and Golub R. CXCR6 Expression Is Important for Retention and Circulation of ILC Precursors. *Mediators Inflamm.* 2015;2015:368427.

63. Hudspeth K, Donadon M, Cimino M, Pontarini E, Tentorio P, Preti M, et al. Human
liver-resident CD56(bright)/CD16(neg) NK cells are retained within hepatic sinusoids via the
engagement of CCR5 and CXCR6 pathways. *Journal of autoimmunity.* 2016;66:40-50.

64. Lynn RC, Weber EW, Sotillo E, Gennert D, Xu P, Good Z, et al. c-Jun overexpression in CAR
T cells induces exhaustion resistance. *Nature.* 2019;576(7786):293-300.

65. Wu H, Zhao X, Hochrein SM, Eckstein M, Gubert GF, Knopper K, et al. Mitochondrial
dysfunction promotes the transition of precursor to terminally exhausted T cells through
HIF-1alpha-mediated glycolytic reprogramming. *Nature communications.* 2023;14(1):6858.

66. Zheng X, Qian Y, Fu B, Jiao D, Jiang Y, Chen P, et al. Mitochondrial fragmentation limits NK
cell-based tumor immunosurveillance. *Nature immunology.* 2019;20(12):1656-67.

67. Cheng HP, Huang CJ, Tsai ML, Ong HT, Cheong SK, Choo KB, et al. MicroRNA-362
negatively and positively regulates SMAD4 expression in TGF-beta/SMAD signaling to
suppress cell migration and invasion. *Int J Med Sci.* 2021;18(8):1798-809.

68. Jeevan-Raj B, Gehrig J, Charmoy M, Chennupati V, Grandclement C, Angelino P, et al. The
Transcription Factor Tcf1 Contributes to Normal NK Cell Development and Function by
Limiting the Expression of Granzymes. *Cell reports.* 2017;20(3):613-26.

69. Guo H, Wang M, Wang B, Guo L, Cheng Y, Wang Z, et al. PRDM1 Drives Human Primary T
Cell Hyporesponsiveness by Altering the T Cell Transcriptome and Epigenome. *Front
Immunol.* 2022;13:879501.

70. Babichuk CK, and Bleackley RC. Mutational analysis of the murine granzyme B gene
promoter in primary T cells and a T cell clone. *J Biol Chem.* 1997;272(30):18564-71.

71. Kucuk C, Iqbal J, Hu X, Gaulard P, De Leval L, Srivastava G, et al. PRDM1 is a tumor
suppressor gene in natural killer cell malignancies. *Proc Natl Acad Sci U S A.*
2011;108(50):20119-24.

72. Kim D, Langmead B, and Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods.* 2015;12(4):357-60.

73. Li B, and Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics.* 2011;12:323.

74. Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

75. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment
tool for interpreting omics data. *Innovation (Camb).* 2021;2(3):100141.

76. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated
analysis of multimodal single-cell data. *Cell.* 2021;184(13):3573-87 e29.

77. Germain PL, Lun A, Garcia Meixide C, Macnair W, and Robinson MD. Doublet identification
in single-cell sequencing data using scDblFinder. *F1000Res.* 2021;10:979.

78. Caicedo HH, Hashimoto DA, Caicedo JC, Pentland A, and Pisano GP. Overcoming barriers to
early disease intervention. *Nat Biotechnol.* 2020;38(6):669-73.

Graphical abstract

Human *NK-PRDM1* signature

NK-PRDM1 high (n=181)
NK-PRDM1 low (n=181)

P = 0.0177

Survival in liver cancer patients

Figure 1. *Prdm1* promotes NK cell homeostasis and terminal maturation.
(**A**) Correlation between the average expression of NK cell-associated signature and *PRDM1* in LIHC (Liver Hepatocellular Carcinoma; n=363) patients sourced from TCGA datasets. (**B**) Heatmap of the ordered, z-score normalized expression values for *PRDM1* and NK cell-associated genes in liver cancer patients. High and low expression of *NK-PRDM1* signature are indicated. (**C**) Prognostic value of the *NK-PRDM1* signature for overall survival of liver cancer patients comparing high and low samples with a median cutoff. (**D**) Schematic representation of the Prdm1-conditional knockout mouse model. Targeted exon 6-8 of *Prdm1* (top) is flanked with *loxP* sites (middle). Ncr1-expressed Cre recombinase was used to generate the *Prdm1^ΔNcr1* allele (bottom). (**E** and **F**) Real-time RT-PCR quantification of *Prdm1* expression in NKp46+ cells (**E**) and splenocytes (**F**) to determine the presence of *Prdm1* (n=5). (**G**) Representative flow cytometric plots (left) and quantification (right) of the proportion and absolute number of CD45+CD3-NKp46+ cells among lymphocytes in liver, lung, and bone marrow (n=10). (**H**) Representative flow cytometric plots (left) of the CD11b and CD27 expression within CD3-NK1.1+NKp46+CD49b+ cells in liver, lung, and bone marrow (n=10). Right panel showed the percentage of distinct maturation stages of NK cells. Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; r, pearson correlation coefficient; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 2. *Prdm1* is required for group 1 ILCs to control tumor metastasis.

(**A**) Representative flow cytometric plots (left) of liver cNK cells (CD49a⁻CD49b⁺) and ILC1s (CD49a⁺CD49b⁻) from *Prdm1⁺/⁺* and *Prdm1^ΔNcr1* mice. The two bar graphs (right) quantitate the percentages and absolute numbers of cells respectively (n=7). (**B** and **D**) Representative image (left) and quantification (right) of tumor nodes on the livers (n=7) (**B**) and lungs (n=10) (**D**) of *Prdm1⁺/⁺* and *Prdm1^ΔNcr1* mice at day 14 or 21 after inoculation with B16F10 melanoma cells. (**C** and **E**) Representative histopathological images of liver (**C**) and lung (**E**) tissues stained by hematoxylin-eosin to detect tumor metastasis. Red bar indicates 500 μm distance under the microscope. (**F**) Liver cells and splenocytes were co-stimulated in the presence or absence of IL-12 and IL-18 for 12 hours. GolgiStop was added 4 hours before intracellular staining of IFN-γ. The graphs showed percentage of IFN-γ⁺ splenic cNK cells, liver cNK cells and ILC1s from *Prdm1⁺/⁺* and *Prdm1^ΔNcr1* mice (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 3. Bulk RNA-seq reveals *Prdm1*-mediated functions in splenic cNK cells.
(**A**) Splenic cNK cells and liver CD45+ cells were sorted from *Prdm1+/+* and *Prdm1ΔNcr1* mice using flow cytometry, and prepared for bulk RNA-seq and single-cell RNA-seq analysis. (**B**) Volcano plot of the bulk RNA-seq differentially expressed genes (DEGs) (log₂|fold change|>0.5; *P*<0.05) in splenic cNK cells between *Prdm1+/+* and *Prdm1ΔNcr1* mice. Upregulated and downregulated genes in *Prdm1ΔNcr1* cells were highlighted in red and blue. (**C**) Enriched Gene Ontology (GO) terms of DEGs in *Prdm1ΔNcr1* cells compared *Prdm1+/+* cells. The Enrichment gene set in upregulated (red) and downregulated (blue) genes were indicated in different colour. Bar length represents statistical significance. (**D**) Gene Set Enrichment Analysis (GSEA) showing the enrichment of NF-kappa B signaling pathway of DEGs in *Prdm1ΔNcr1* cells compared *Prdm1+/+* cells. NES, normalized enrichment score. (**E**) Heatmap of selected genes from DEGs. Shown is z-score transformed expression of DEGs. (**F**) Representative flow cytometric plots (left) and cumulative data (right) of the percentage and absolute numbers of CX3CR1+ cells in liver cNK cells and ILC1s (n=5). (**G**) Quantification of CX3CR1+ cells in NK cells in blood, bone marrow, lung, liver, spleen, and lymph nodes (n=5). (**H** and **I**) Representative flow cytometric plots (left) and cumulative data (right) showing the proportion of GZMB+ (**H**) and PRF1+ (**J**) liver cNK cells and ILC1s from *Prdm1+/+* and *Prdm1ΔNcr1* mice (n=5). (**J** and **K**) Proportion of GZMB+ (**J**) and PRF1+ (**K**) splenic NK cells at different maturation stages was analyzed by flow cytometry (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 4. Different properties of cNK clusters following *Prdm1* knockout.
(**A**) Uniform manifold approximation and projection (UMAP) visualization of liver CD45⁺ cells from *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* mice. Twelve clusters were defined and indicated by distinct colours. Each dot represents a single cell. (**B**) UMAP visualization of liver cNK and ILC1 clusters. Cells were colored by genotypes (*Prdm1⁺/⁺*-blue; *Prdm1ᐞNcr1*-red). (**C**) Percentages of cNK cells and ILC1s in total group 1 ILCs (left), and their distribution in each cluster (right). (**D** and **E**) UMAP visualization of three different liver cNK clusters from two mouse strains. (**F**) Proportions of cNK cells among total cNK cells (left; 211 cells in *Prdm1⁺/⁺*, and 141 cells in *Prdm1ᐞNcr1*) and within clusters (right). (**G**) Violin plots showing the normalized expression of select genes in different cNK clusters. (**H**) Enriched GO term of marker genes in three cNK clusters. Dot size represents enriched gene number, and color intensity represents significance. (**I-L**) GSEA plots (left) depicting the enrichment of NF-kappa B (**I**), TNF (**J**), IL-17 (**K**), and MAPK (**L**) signaling pathway in *Junb*ʰⁱ cNK cluster compared with clusters of *Prf1*ʰⁱ and *Cxcr3*ʰⁱ cNK cells. Right panel showed dynamic relative expression of the given gene sets from cluster1 to cluster3 between *Prdm1⁺/⁺* and *Prdm1ᐞNcr1*. Dots represent the average expression of given gene set in each cell, which was calculated through the sum of normalized expression of each individual gene within the designated gene set in every single cell. NES, normalized enrichment score.

Figure 5. Different properties of ILC1 clusters following *Prdm1* knockout..

(**A and B**) UMAP visualization of three different liver ILC1 clusters from two mouse strains. (**C**) Proportions of ILC1s among total ILC1s in different genotypes (left; 114 cells in *Prdm1*[+/+], and 63 cells in *Prdm1*[ΔNcr1]) and within each cluster (right). (**D**) Violin plots showing the normalized expression of select genes in different ILC1 clusters. (**E**) Ridge plots showing the normalized expression of *Gzmb, Prf1,* and *Junb* in cNK and ILC1 clusters between *Prdm1*[+/+] and *Prdm1*[ΔNcr1] cells. (**F** and **G**) GSEA plots (left) depicting the enrichment of IL-17 signaling pathway (**F**) and T cell differentiation (**G**) in *Il7r*[hi] ILC1 cluster compared with clusters of *Klra*[hi] and *Gzmb*[hi] ILC1s. Right panel showed dynamic relative expression of the given gene sets from cluster1 to cluster3 between *Prdm1*[+/+] and *Prdm1*[ΔNcr1]. Dots represent the average expression of given gene set in each cell, which was calculated through the sum of normalized expression of each individual gene within the designated gene set in every single cell. NES, normalized enrichment score. (**H**) Enriched GO term of marker genes in three ILC1 clusters. Dot size represents enriched gene number, and color intensity represents significance. (**I**) Violin plot showing the *Junb*[hi] signature score for cNK cell and ILC1 clusters, calculated using the signature genes of *Junb*[hi] cNK cluster. (**J**) Venn diagram showing overlapping and unique DEGs in comparisons within liver cNK cells, ILC1s and splenic cNK cells between *Prdm1*[+/+] and *Prdm1*[ΔNcr1](left), and 17 overlapped DEGs were shown at the right panel.

Figure 6. *Prdm1* facilitates the intercellular communication between liver group 1 ILCs and macrophages.
(**A** and **B**) UMAP visualization of monocyte-derived macrophages (MDMs) and Kupffer cells (KCs) cluster (**A**) between *Prdm1*+/+ and *Prdm1*ΔNcr1 (**B**). (**C**) Proportions of total macrophages in liver immune cells (left), and proportions of MDMs and KCs among total macrophages in different genotypes (510 cells in *Prdm1*+/+, and 624 cells in *Prdm1*ΔNcr1). (**D**) Representative flow cytometric plots (left) and cumulative data (right) of the percentage and absolute numbers of liver total macrophages (CD45+Ly6G-CD11b+F4/80+), MDMs (CX3CR1-), and KCs (CX3CR1+) between *Prdm1*+/+ and *Prdm1*ΔNcr1 (n=7). (**E**) Representative flow cytometric plots (left) and cumulative data (right) of the percentage and absolute numbers of Ly6C- and Ly6C+ cells in MDMs. (**F** and **G**) Circle plots (**F**) and summary data (**G**) illustrating the interaction numbers and strength of significant enriched ligand–receptor pairs among cluster of liver cNK cells, ILC1s, and macrophages from *Prdm1*+/+ (left) and *Prdm1*ΔNcr1 (right) cells. The thickness of the line indicates the number of enrich pairs, and the arrow reflects the direction of the interaction. (**H**) Heatmap of overall signaling pattern recognized from ligand-receptor pairs, which contained the sum of signaling from the sender and target cells. (**I**) Bar graphs showing the information flow in selected active signaling patterns between *Prdm1*+/+ and *Prdm1*ΔNcr1 cells. Relative information flow was calculated as the sum of the communication probability in given signaling patterns. (**J**) Chord plot of the CXCL signaling interaction network among cluster of liver cNK cells, ILC1s, and macrophages between *Prdm1*+/+ and *Prdm1*ΔNcr1 cells. Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 7. *Prdm1* **safeguards group 1 ILCs from exhaustion-like phenotypes in the tumor microenvironment.**
(**A-C**) The Mitochondrial mass (MitoTracker Green staining; n=8) (**A**), Mitochondrial ROS (MitoSOX staining; n=5) (**B**), and Mitochondrial membrane potential (TMRM staining; n=5) (**C**) of splenic cNK cells, liver cNK cells and ILC1s were analyzed by flow cytometry. Representative flow cytometric plots (left) and cumulative data (right) showing the relative mean fluorescence intensities (MFIs) of each group. (**D**) Representative flow cytometric plots (left) and cumulative data (right) showing the percentage and absolute number of liver cNK cells and ILC1s in *Prdm1*$^{+/+}$ and *Prdm1*$^{\Delta Ncr1}$ tumor-bearing mice at day 14 after inoculation with B16F10 melanoma cells via intrasplenic injection(n=5). (**E**) Percentages of IFN-γ$^+$ liver cNK cells and ILC1s from *Prdm1*$^{+/+}$ and *Prdm1*$^{\Delta Ncr1}$ tumor-bearing mice (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.