Title page

Manuscript title

Prdm1 Positively Regulates Liver Type 1 ILCs Cancer Immune Surveillance and Preserves Functional Heterogeneity

Authors and affiliations

Jitian He[1], Le Gao[1], Peiying Wang[1], Wing Keung Chan[4], Yiran Zheng[1], Yumo Zhang[1], Jiming Wang[5], Huaiyong Chen[3,6,7], Zhouxin Yang[2], Youwei Wang[1]

[1]Institute of Medical Engineering & Translational Medicine, Tianjin University, Tianjin 300072, China.

[2]Zhejiang Provincial Key Lab of Geriatrics and Geriatrics Institute of Zhejiang Province, Zhejiang Hospital, Hangzhou 310030, China.

[3]College of Pulmonary and Critical Care Medicine, 8th Medical Center, Chinese PLA General Hospital, Beijing, China

[4]Department of Internal Medicine, Division of Hematology, The Ohio State University, Columbus, OH 43210, USA.

[5]Tianjin Economic-Technological Development Area (TEDA) Hospital, Tianjin 300451, China

[6]Department of Basic Medicine, Haihe Hospital, Tianjin University, Tianjin 300350, China.

[7]Tianjin Key Laboratory of Lung Regenerative Medicine, Tianjin 300350, China.

Authorship notes: JH, LG, and PW contributed equally to this work.

Co-corresponding authors: Huaiyong Chen (huaiyong.chen@foxmail.com), Zhouxin Yang

(yangzhouxin@hotmail.com), Youwei Wang (youwei.wang@tju.edu.cn)

Address correspondence to: Youwei Wang, Institute of Medical Engineering & Translational

Medicine, Tianjin University, Tianjin 300072, China. Email: youwei.wang@tju.edu.cn. Or to

Zhouxin Yang, Zhejiang Provincial Key Lab of Geriatrics and Geriatrics Institute of Zhejiang

Province, Zhejiang Hospital, Hangzhou 310030, China. Email: yangzhouxin@hotmail.com. Or to

Huaiyong Chen, Department of Basic Medicine, Haihe Hospital, Tianjin University, Tianjin

300350, China. Email: huaiyong.chen@foxmail.com.

Abstract

Liver Type 1 innate lymphoid cells (ILCs) comprise conventional natural killer (cNK) cells and ILC1s. The main functions of Type 1 ILCs not only include directly killing target cells but also regulating the local immune microenvironment of the liver through the secretion of cytokines. Uncovering the intricate mechanisms by which transcriptional factors regulate and influence the functions of Type 1 ILCs, particularly within the context of liver tumors, presents a significant opportunity to amplify the effectiveness of immunotherapies against liver malignancies. Using Ncr1-drived conditional knockout mouse model, our study reveals the regulatory role of *Prdm1* in shaping the composition and maturation of liver Type 1 ILCs. Notably, Prdm1 regulates the ratio between NK cells and ILC1s, promoting a shift in the balance towards the direction of NK cells. Although *Prdm1* did not affect the killing function of cNK cells in an *in vivo* cytotoxicity model, a significant increase in cancer metastasis was observed in *Prdm1* knockout mice. IFN-γ, granzyme B, and perforin secretion decreased significantly in *Prdm1* deficient Type 1 ILCs. scRNA sequencing data also provided evidence that Prdm1 sustains functional subsets of liver type 1 ILCs and facilitates communications between Type 1 ILCs and macrophages. The present study unveiled a novel regulatory mechanism of Prdm1 in liver Type 1 ILCs, showing promising potential for developing innovative immune therapy strategies against liver cancer.

Main text

Introduction

Liver Type 1 ILCs consist of cNK cells and ILC1s (1), with distinct developmental trajectories and effect molecules (2). Both NK cells and ILC1s play indispensable roles in combatting viral infections (3), maintaining local immune homeostasis (4), eradicating malignant transformed cells (5), and fostering cross-talk with adaptive immunity (6). NK cells demonstrate potent cellular cytotoxicity, facilitating direct elimination of target cells (7). On the contrary, a defining attribute of ILC1s is their predominant cytokine-mediated functions, with limited cellular killing capacity (8). In a state of homeostasis, liver Type 1 ILCs ($CD45^+CD3^-NK1.1^+NKp46^+$) can be discriminated into cNK cells and ILC1s by the differential expression of CD49a and CD49b (1): liver cNK cells are marked by the expression of CD49b, while liver ILC1s exhibit a distinctive positivity for CD49a. Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) is also expressed on liver ILC1s, but not on liver NK cells (9, 10). Transcriptional factors (TFs) such as T-bet, Nfil3, PLZF, and ID2 are also required for Type 1 ILCs development or generation of their progenitors (11-14). Eomes is considered to be necessary for NK cell maturation, but not for ILC1s (15). On the contrary, deficiency of Hobit results in the depletion of ILC1s, while only leaving little impact on liver NK cells (16). These studies suggest that TFs may, or usually, play different roles in NK cells and ILC1s.

During the development of liver cancers, the functions of immune cells are often inhibited, resulting in the formation of an immunosuppressive tumor microenvironment, and sometimes even systemic immune suppression (17). Transforming a cold tumor into a hot tumor with a

stronger immune response is one of the goals of many cancer immunotherapies (18). Research

targeting the immune system is showing increasing clinical promise in liver cancer treatment (19).

Our recent study also found that Toll-like receptor agonists can significantly enhance the

anti-tumor effect of Sorafenib by reconstructing the tumor immune microenvironment and

reshaping the vascular system (20).

The anti-tumor activity of NK cells in the liver is relatively well-established, whereas the

relationship between ILC1 and tumors is still a topic of controversy. Clinical research findings

have shown that liver tumors with a higher infiltration of NK cells are associated with a better

prognosis (21). Decreased NK cell activity is closely correlated with the malignancy of liver

tumors and represents a significant risk factor for recurrence (22). The tumor microenvironment

orchestrates the transformation of NK cells into ILC1s in a TGF-β-dependent manner, resulting in

their diminished capacity to control tumor growth and metastasis. This process ultimately

promotes tumor immunoevasion (23). However, research has also revealed that, distinct from

directly inhibiting tumor growth, the primary function of ILC1 is to suppress the seeding of

metastatic tumor cells in liver tissue (5). Comprehensive research in this field is essential to

harness the precision of Type 1 ILCs targeted immunotherapy for liver cancers.

The transcription factor network governs the function of Type 1 ILCs and the balance

between NK and ILC1. Our research team, along with others, has observed that Smad4 promotes

the shift in balance from NK cells towards ILC1 in a TGF-beta-independent pathway. This study

also revealed that Smad4 positively regulates the expression of another transcription factor,

Prdm1(24, 25). PR domain 1 (Prdm1/Blimp1) plays a crucial role in the differentiation of B cells

into plasma cells and the homeostasis of T cells (26-28). The function and regulatory network of

Prdm1 in NK cells is distinct from B cells and T cells, for its expression is independent of B-cell

lymphoma 6 (Bcl-6) and Interferon Regulatory actor 4 (IRF4) but relies on T-bet. In the study that

identified Prdm1 as an essential transcriptional factor for NK cell maturation (29), no significant

differences were observed in IFN-γ production or cytotoxicity between Prdm1-deficient and

wild-type NK cells. Although Prdm1 expression is dependent on IL-15 in immature NK cells and

can be further upregulated by IL-12 or IL-21, it plays a role in downregulating the expression of

certain cytokine receptors, such as CD25, consequently diminishing the responsiveness of NK

cells to IL-2 (30). Notably, some studies have even suggested that Prdm1 suppresses the secretion

of IFN-γ by NK cells (31). These findings imply that, although Prdm1 promotes NK cell

development, it may function as a negative regulator of their activity. However, direct evidence

supporting the impact of Prdm1 on NK cell anti-tumor capabilities is still lacking. Furthermore,

there has been no investigation into its influence on the homeostasis of NK cells and ILC1s in the

liver, nor has there been an exploration of the underlying mechanisms.

In the current study, we found that the deletion of *Prdm1* in Ncr1[+] cells resulted in an

imbalance in the homeostasis of liver Type 1 ILCs, with a shift towards cNK cells. The data also

support the essential role of Prdm1 in the cancer surveillance mediated by NK cells. While Prdm1

positively regulates genes associated with cellular cytotoxicity, it concurrently exerts inhibitory

control over certain positive regulators of NK cell development and functionality, such as JunB.

Using single-cell RNA sequencing, we have identified a subset of NK cells within the liver

characterized by elevated JunB expression. These cells exhibit decreased expression of genes

associated with cellular cytotoxicity, and their abundance significantly increases following Prdm1

knockout.

Results

Prdm1 promotes Type 1 ILCs homeostasis and terminal maturation

Examination of 363 liver hepatocellular carcinoma (LIHC) patient samples from The Cancer

Genome Atlas (TCGA) revealed a positive correlation between the expression of NK

cell-associated genes (*NCR1*, *NCR3*, *KLRB1*, *CD160*, and *PRF1*) (32) and *PRDM1* expression

(Figure 1A). Patients with top and bottom quartiles of *NK-PRDM1* signature expression were

chosen for survival analysis (Figure 1B). Notably, patients with the *NK-PRDM1*hi signature had

better overall survival compared to the these with NK-*PRDM1*lo signature (Figure 1C). Similar

results were also found in skin cutaneous melanoma (SKCM, n=454) and lung adenocarcinoma

(LUAD, n=497) patients (Supplemental Figure 1, A-F). These data suggested that *PRDM1* in NK

cells might be essential for immune surveillance in some solid tumors, including liver cancer.

These findings prompted us to investigate the impact and mechanism of *PRDM1* in NK cells and

ILC1 within the context of liver cancer.

To achieve this, we developed a conditional knockout mouse model to specifically deplete

Prdm1 in Type 1 ILCs. *Ncr1-Cre* mice were crossed with *Prdm1*fl/fl mice to generate *Ncr1-cre*

*Prdm1*fl/fl mice, which specifically knockout exons 6-8 of *Prdm1* (33) in NKp46 positive cells

(Figure 1D). The mice carrying *Ncr1*Cre/+*Prdm1*fl/fl were referred to as *Prdm1*ΔNcr1 mice, and the

mice carrying *Ncr1*+/+*Prdm1*fl/fl were referred to as *Prdm1*+/+ mice or wild-type (WT) mice. To

further validate the deletion of *Prdm1* in NKp46+ cells of *Prdm1*ΔNcr1 mice, CD3-NK1.1+NKp46+

cells were sorted and the expression of *Prdm1* was quantified by real-time RT-PCR. The

expression of *Prdm1* was almost undetectable in *Prdm1$^{\Delta Ncr1}$* mice (Figure 1E), indicating

successful knockout of *Prdm1* in NKp46$^+$ cells.

Proportion and absolute number of cNK cells in blood, bone marrow, lung, liver, spleen, and

lymph nodes were analyzed by flow cytometry. Compared with *Prdm1$^{+/+}$* mice, the percentage of

cNK cells (CD3$^-$NK1.1$^+$NKp46$^+$) among lymphocytes was decreased in all of these tissues except

bone marrow and lymph nodes (Figure 1F; Supplemental Figure 2A). However, no significant

difference was observed in the percentage of cNK cells among bone marrow-derived lymphocytes

between *Prdm1$^{\Delta Ncr1}$* and *Prdm1$^{+/+}$* mice. The absolute number of cNK cells in blood, lung, liver,

and spleen also decreased in *Prdm1$^{\Delta Ncr1}$* mice (Figure 1F; Supplemental Figure 2A). Only a slight

decrease in the number of cNK cells was observed in the lymph nodes of *Prdm1$^{\Delta Ncr1}$* mice, which

did not reach statistical significance either (Supplemental Figure 2A). In contrast, the absolute

number of cNK cells in *Prdm1$^{fl/fl}$* mice bone marrow is moderately higher than *Prdm1$^{\Delta Ncr1}$* mice

(Figure 1F).

NK cell terminal maturation can be divided into four stages according to the expression of

CD11b and CD27 (34). Stage IV (CD11b$^+$CD27$^-$) NK cells show higher level of effector

molecules than any other stages and are considered as the most mature NK cells (35). Based on

the expression of CD11b and CD27, the maturation of cNK cells from blood, bone marrow, lung,

liver, spleen, and lymph nodes were assessed. Compared with *Prdm1$^{+/+}$* mice, the proportion of

most mature CD11b$^+$CD27$^-$ NK cells were significantly decreased in all of the analyzed tissues in

Prdm1$^{\Delta Ncr1}$ mice, without any exceptions (Figure 1G; Supplemental Figure 2B). Remarkably, an

increase in immature CD11b$^-$CD27$^+$ NK cells was also observed in these tissues. Killer cell

lectin-like receptor subfamily G member 1 (KLRG1) is a lectin-like receptor which was

considered as another marker of NK cell maturation (36). In both *Prdm1*[+/+] and *Prdm1*[ΔNcr1] mice,

cNK cells from the liver and lung had the highest expression of KLRG1, following by blood and

spleen (Supplemental Figure 2, C and D). The lowest KLRG1 expression was observed in cNK

cells derived from lymph nodes and bone marrow, indicating the presence of the most immature

cNK cells in these tissues. Consistent with CD11b/CD27 based maturation analysis, a significant

loss of KLRG1[+] NK cells in *Prdm1*[ΔNcr1] mice was observed compared to *Prdm1*[+/+] mice

(Supplemental Figure 2, C and D). Together, these data showed that *Prdm1* is required for the

terminal maturation of cNK cells among various tissues.

Prdm1 is required for Type 1 ILCs to control tumor metastasis

Two subpopulations of liver Type 1 ILCs (gated by CD45[+]CD3[-]NK1.1[+]NKp46[+]) were further

analyzed based on the expression of CD49a and CD49b (Figure 2A). Compared with *Prdm1*[+/+]

mice, the *Prdm1*[ΔNcr1] mice exhibited an increased percentage of cNK cells (CD49a[-]CD49b[+]) and

reduced proportion of ILC1s (CD49a[+]CD49b[-]) (Figure 2A). Of note, the absolute number of both

cNK cells and ILC1s were decreased in *Prdm1*[ΔNcr1] mice, with a more robustly reduction in ILC1s

(Figure 2A). This underscored the crucial role of *Prdm1* in maintaining the homeostasis of both

liver cNK cells and ILC1s.

The disruption of NK cell homeostasis and maturation due to *Prdm1* loss motivated us to

further explore whether deficiency of Prdm1 impaired NK cell cytotoxicity. A B2M-deficient

cell-based *in vivo* cytotoxicity assay was used to evaluate the effect of *Prdm1* on the cytotoxicity

of NK cells (37, 38). B2M-deficient cells do not have detectable Major Histocompatibility

Complex I (MHC-I) on the cell surface, making them the target of NK cells (39). Health NK cells

will reject B2M-deficient donor cells efficiently and the elimination was used to quantify the

cytotoxicity of NK cells. Although significant impaired homeostasis and maturation of NK cells

were observed in *Prdm1^{ΔNcr1}* mice, no significant difference in the *in vivo* cytotoxicity assay were

observed between *Prdm1^{+/+}* and *Prdm1^{ΔNcr1}* mice (Figure 2, B and C).

While our B2M-deficient cell-based assay did not reveal a significant impact of *Prdm1* on

NK cell cytotoxicity, both our results and those of other researchers have demonstrated *Prdm1*'s

role in positively regulating NK cell development and homeostasis. Besides direct cytotoxicity,

other factors influence the anti-tumor capability of NK cells, such as the ability to counteract

tumor-induced immune suppression and exhaustion, the secretion of cytokines to activate other

immune cells for tumor elimination. Moreover, the cytotoxicity assay, due to its relatively short

duration, might not fully represent the anti-tumor activity of NK cells when continuously exposed

to immune inhibitory signals in the tumor microenvironment. Therefore, we initiated an *in vivo*

tumor model to further investigate the impact of *Prdm1* on NK cell anti-tumor capabilities.

B16F10 is a melanoma cell line with low expression of MHC-I, which was susceptible to NK cell

killing and usually used to evaluate NK cell anti-tumor capacity (40, 41). The B16F10 cells were

intravenously (for lung metastasis) or intrasplenic (for liver metastasis) administrated in the mice.

The melanoma nodes were quantified three (intravenous injection) or two (intrasplenic injection)

weeks after tumor inoculation. Compared with *Prdm1^{+/+}* mice, deficiency of *Prdm1* resulted in

more metastasis nodules in both lung (~ 2-fold) and liver (~ 4-fold) (Figure 2, D and F).

Histological analysis further confirmed the increased frequency of metastasis tumor foci in

Prdm1^{ΔNcr1} mice (Figure 2, E and G).

In agreement with our *in vivo* data, we also observed decreased IFN-γ secretion in *Prdm1^{ΔNcr1}*

mice-derived splenic cNK cells, liver cNK cells, and liver ILC1s when stimulated by IL-18 alone

or IL-12/IL-18 (Figure 2H; Supplemental Figure3), which indicated that *Prdm1* is required for full

activation of cNK cells and ILC1s in the context of IFN-γ production. These data implying that

although similar cellular killing capacity was observed between *Prdm1*$^{+/+}$ and *Prdm1*^{ΔNcr1} mice,

Prdm1 is indispensable for NK cell mediated tumor surveillance.

Bulk RNA-seq depicts *Prdm1*-mediated functions in peripheral cNK cells

Bulk RNA sequencing of splenic cNK cells (CD3$^-$NK1.1$^+$NKp46$^+$) was conducted to uncover

the molecular mechanisms by which *Prdm1* regulates NK cell anti-tumor immunity (Figure 3A).

Differentially expressed genes (DEGs) between *Prdm1*$^{+/+}$ and *Prdm1*^{ΔNcr1} mice were determined

using a criterion of $\log_2$ (fold change) > 0.5 and *P* < 0.05. 445 DEGs were identified out of 17434

protein-coding genes, which consisted of 223 upregulated genes and 222 downregulated genes

(Figure 3B).

Gene Ontology (GO) analysis revealed the enrichment of glucuronate metabolism and

lymphocyte differentiation in upregulated genes in *Prdm1*^{ΔNcr1} mice derived NK cells (Figure 3C),

both of which were associated with cellular growth and development. In contrast, leukocyte

mediate cytotoxicity, immune receptor activity, and integrin binding was enriched in the genes

which decreased their expression level in in *Prdm1*^{ΔNcr1} mice (Figure 3C).

Gene Set Enrichment Analysis (GSEA) showed enriched NF-kappa B signaling pathway and

negative regulation of IL-6 production pathway in *Prdm1*-deficient cNK cells (Figure 3, D and E),

suggesting the potential targets by *Prdm1* to regulate NK cell function. Increased expression of

multiple TFs such as *Junb*, *Batf3*, *Nfkb1*, *Tcf7*, and *Nr4a2* was observed in *Prdm1* knockout cNK

cells, suggesting they might be suppressed by *Prdm1* (Figure 3F). Downregulation of granzyme B

(*Gzmb*), Perfroin (*Prf1*) were observed in *Prdm1* deficient NK cells (Figure 3F), implied

decreased anti-tumor ability, which was consistent with increased melanoma metastasis in

Prdm1^{ΔNcr1} mice (Figure 2D and F). *Cxcr6*, and *Cx3cr1* were also decreased in the NK cells

derived from *Prdm1^{ΔNcr1}* mice (Figure 3F), which was considered to play an important role in

promoting the egress of NK cells from bone marrow. This might be the reason for the quantity of

NK cells increased in bone marrow while decreasing in other peripheral tissues (Figure 1F). As a

result of the reduced expression levels of *Cxcr6* and *Cx3cr1*, NK cells were unable to egress from

the bone marrow and accumulated therein. Consistent with decreased production of IFN-γ after

stimulated by IL-12/IL-18 (Figure 2H), decreased expression of *Il18rap* and *Il12rb2* were

observed in *Prdm1^{ΔNcr1}* cNK cells (Figure 3F), implying impaired response to cytokine

stimulation.

To confirm the result of RNA-sequencing, the expression of granzyme B and perforin were

also analyzed by flow cytometry. Lower granzyme B and perforin production was observed in

Prdm1-deficient splenic cNK cells, liver cNK cells and ILC1s (Figure 3, G and I; Supplemental

Figure 4, A and B). Notably, granzyme B decreased among all of the maturation stages of cNK

cells. Probably because of relatively low level of perforin in CD11b$^-$CD27$^+$ and CD11b$^+$CD27$^+$

cNK cells, there is not statistical difference between *Prdm1^{+/+}* and *Prdm1^{ΔNcr1}* cNK cells. However,

significantly decrease in Perforin was observed in *Prdm1^{ΔNcr1}* mice derived CD11b$^+$CD27$^-$ NK

cells, which have the highest level of Perforin. (Figure 3, H and J; Supplemental Figure 4C).

These data implied that *Prdm1* directly regulates NK cells to exert cytotoxicity, not only due to the

less mature phenotype in *Prdm1*-deficient NK cells.

245   scRNA-seq reveals unique properties of two clusters from liver type I ILCs following *Prdm1*

246   knockout.

To further investigate the effect of *Prdm1* in liver cNK cells and ILC1s, as well as the

changes in the haptic immune microenvironment caused by the deficiency of *Prdm1* in Type 1

ILCs, single-cell RNA sequencing (scRNA-seq) was performed for CD45$^+$ cells sorted from the

liver (Figure 3A; Supplemental Figure 5A). Initial quality control revealed high-quality of cell

purity, library assembly, and sequencing (Supplemental Figure 5B). 10978 cells passed the quality

criteria and were selected for further analysis (6161 from *Prdm1$^{+/+}$* mice, 4,817 from *Prdm1^{ΔNcr1}*

mice). Unsupervised clustering of all sequenced cells based on transcript signatures identified

twelve distinct clusters, including B cells, Macrophages, CD4$^+$ T cells, CD8$^+$ T cells, dendritic

cells (DCs), NKT cells, cNK cells, ILC1s, Neutrophils, Monocytes, Basophils, and a small

number of undefined cells (Figure 4A). A higher percentage of Macrophages, CD8$^+$ T cells,

Monocytes, DCs, and a lower percentage of NK cells, ILC1s, CD4$^+$ T cells, and NKT cells was

observed in *Prdm1^{ΔNcr1}* mice (Supplemental Figure 5C). Liver cNK cells and ILC1s were

identified based on the expression of surface markers and TFs (Supplemental Figure 5D). For

example, liver cNK cells expressed high levels of *Itga2* (CD49b) and Eomes, while ILC1s had

high levels expression of *Itga1* (CD49a) and *Tnfsf10* (Supplemental Figure 5E). In liver Type 1

ILCs, increased percentage of cNK cells and decreased percentage of ILC1s were observed in

Prdm1^{ΔNcr1} mice (Figure 4, B and C), which was in line with the flow cytometry data (Figure 2A).

To better understand the specific function of *Prdm1* played in liver cNK cells and ILC1s, the

two subpopulations of liver Type 1 ILCs were further analyzed separately using unsupervised

clustering. Liver cNK cells and ILC1s were grouped into three clusters respectively and visualized

by Uniform Manifold Approximation and Projection (UMAP) (Figure 4, D-I). Based on the

cluster specific gene expression signature (Supplemental Figure 6, A and B), the subpopulation of

liver cNK cells were referred as "*Prf1*hi", "*Junb*hi ", and "*Cxcr3*hi" cNK cells (Figure 4, D-F).

The *Prf1*hi cNK cell cluster was defined by high expression of cytolysis-related genes,

including *Ncr1*, *Gzma*, *Gzmb*, *Prf1*, and *Fgl2* (Figure 4J; Supplemental Figure 6C), indicating the

strong target-killing ability of this cluster. GO analysis further revealed the enrichment signatures

of cytolysis, response to virus, and lymphocyte mediated immunity in the genes upregulated in

*Prf1*hi cNK cell cluster, further confirming the cytotoxic effects of this cluster (Figure 4K). While

this cluster is present in both *Prdm1*$^{\Delta Ncr1}$ and *Prdm1*$^{+/+}$ mice, there is a significant reduction in

Prdm1$^{\Delta Ncr1}$ mice (Figure 4F). Additionally, both *Gzma* and *Gzmb* expression were downregulated

in *Prdm1*$^{\Delta Ncr1}$ mice derived cNK cells and ILC1s compared to those from *Prdm1*$^{+/+}$ mice (Figure

5A). These data underscore the crucial role of *Prdm1* in maintaining NK cells with immune

effector functions.

The *Junb*hi liver cNK cell cluster distinguished themselves by higher expression of *Junb*

compared to other clusters (Figure 4J). The expression of *Junb* was also upregulated in *Prdm1*$^{\Delta Ncr1}$

derived cNK cells and ILC1s (Figure 5A). The predominant majority (92.98%) of *Junb*hi liver

cNK cells are derived from *Prdm1*$^{\Delta Ncr1}$ mice, with less than ten percent (7.02%) originating from

Prdm1$^{+/+}$ mice (Figure 4F). Many signal transduction elements, gene expression regulator, and

transcriptional factors, such as *Nfkbia*, *Tnfaip3*, *Nr4a1/2/3*, *Batf3*, *Fos*, *Fosb*, *Tcf7*, and *Kit* were

upregulated in the *Junb*hi liver cNK cells (Figure 4J; Supplemental Figure 6C). The expression of

cytotoxicity related genes, such as *Gzmb* and *Prf1*, in *Junb*hi cluster was also lower than other cNK

cells (Figure 4J). GO analysis showed that the genes upregulated in *Junb*hi liver cNK cells

enriched in cell differentiation, cell activation, and transcriptional regulation (Figure 4K). GSEA

indicates that the NF-kappa B, IL-17, MAPK, and TNF signaling pathways were upregulated in

this clusters (Figure 5, E-H). GSEA also showed that mitochondrial related pathways, such as

mitochondrial protein, oxidative phosphorylation, and respiratory electron transport chain were

suppressed in $Junb^{hi}$ cNK cell cluster (Supplemental Figure 6, D-F). Increased proportion of

$Junb^{hi}$ cluster in $Prdm1$ deficient cNK cells suggested impaired anti-tumor activity, which was

consistent with more melanoma metastasis in $Prdm1^{\Delta Ncr1}$ mice and lower expression of

cytotoxicity-related genes in splenic cNK cells based on bulk RNA-sequencing.

The $Cxcr3^{hi}$ cNK cell cluster was characterized with high expression of $Cxcr3$, $Ccr2$, and

some genes encoding ribosomal subunits such as $Rps7$ (Supplemental Figure 6C). The enrichment

of chemokine receptors in the genes upregulated in the $Cxcr3^{hi}$ cluster implying a greater

likelihood of them being tissue-resident compared with other cNK cell clusters (Figure 4K). The

significant enrichment of ribosomal subunits and cytoplasmic translation in $Cxcr3^{hi}$ cluster (Figure

4K) revealed their distinct and active metabolic profile and the capability to mount immune

responses. The $Cxcr3^{hi}$ cNK cell cluster had a remarkably decreased proportion in $Prdm1^{\Delta Ncr1}$

mice (Figure 4F), which emphasized the critical role of $Prdm1$ in maintaining this cluster of liver

cNK cells. Consisting with the flow cytometry result that showed an increase in the number of NK

cells in the bone marrow and a decrease in NK cells in peripheral tissues, bulk RNA sequencing

data also found that the expression of $Cx3cr1$ and $Cxcr6$ decreased in $Prdm1^{\Delta Ncr1}$ cNK cells. These

findings supported the hypothesis that $Prdm1$, through regulating chemokine receptor expression

levels, influenced the distribution of NK cells in the bone marrow and peripheral tissues,

particularly within the liver tissue.

Three clusters of ILC1s were identified from liver ILC1s (Figure 4, G-I), comprising "$Il7r^{hi}$",

"$Klra^{hi}$", and "$Gzma^{hi}$" ILC1s. The first two clusters of ILC1s were characterized by higher

expression of $Il7r$ (CD127) and $Klra5$ separately, while the $Gzma^{hi}$ ILC1 cluster was identified by

elevated expression of both $Gzma$ and $Gzmb$ (Supplemental Figure 7, A-C). The $Il7r^{hi}$ ILC1s

cluster was distinguishable from other ILC1 clusters by its high expression of $Il7r$, IL18RA

($Il18r1$), and IFN-γ ($Ifng$) (Figure 5B; Supplemental Figure 7C). The high expression of $Il18r1$

and $Ifng$ in $Il7r^{hi}$ ILC1s indicated this cluster of cells was highly responsive to IL-18. Module

scores, calculated based on the expression of feature genes within the $Junb^{hi}$ cNK cell cluster,

revealed a comparable $Junb^{hi}$ signature expression pattern within the $Il7r^{hi}$ ILC1 cluster (Figure

5C). GO analysis and GSEA showed that IL-17, NF-kappa B, TNF, MAPK signaling pathway and

T cell differentiation were activated in the $Il7r^{hi}$ ILC1 cluster (Figure 5D, I and J; Supplemental

Figure 7D-F). Considering the close relationship between IL-17 mediated immunity response and

ILC3 (42), it is plausible that the observed heterogeneity within this cluster may be attributed, at

least in part, to potential plasticity between ILC1 and ILC3 subsets. These two clusters appear

reminiscent of an intermediate stage in the conversion process from cNK cells to ILC1s, as

previously described (23, 43). There was no significant difference in the proportion of this cluster

between WT and in $Prdm1^{\Delta Ncr1}$ mice.

The second liver ILC1 cluster, characterized by high expression of Ly49E ($Klra5$) and Ly49G

($Klra7$), was designated as the $Klra^{hi}$ ILC1 cluster (Figure 5B). Notably, there was an elevated

proportion of $Klra^{hi}$ ILC1s in $Prdm1^{\Delta Ncr1}$ ILC1s (39.7%) compared to $Prdm1^{+/+}$ ILC1s (28.1%).

Liver Ly49E^{+} ILC1s have been identified as possessing greater cytotoxic potential and a more

robust viral response compared to liver Ly49E^{-} ILC1s (44). The $Klra^{hi}$ cluster exhibited notably

high expression of *Ccl5* (Supplemental Figure 7C). Previous research has underscored the pivotal

role of CCL5, produced by both cNK cells and ILC1s, in facilitating the accumulation of DCs

within the tumor microenvironment, thereby impeding tumor immune evasion, as highlighted in

studies (32, 45). The expression of *Ccl5* was reduced in the *Klra*^{hi} cluster of *Prdm1^{ΔNcr1}* ILC1s

compared to *Prdm1$^{+/+}$* ILC1s (Supplemental Figure 7C). This decrease in *Ccl5* expression could

potentially have a detrimental impact on the ability of *Klra*^{hi} ILC1s to develop a connection

between innate and adaptive immune responses.

The *Gzma*^{hi} ILC1 cluster was identified according to the high expression of *Gzma*, *Gzmb*, and

Fgl2 (Supplemental Figure 7C), compared to other ILC1 clusters. GO analysis also revealed the

enrichment in cytolysis and stimulus-response capacity (Figure 5D) of the *Gzma*^{hi} ILC1 cluster.

Consistent with the *Prf1*^{hi} cNK cell cluster, the proportion of the *Gzma*^{hi} cluster among liver ILC1s

exhibited a considerable reduction in *Prdm1^{ΔNcr1}* mice compared to *Prdm1$^{+/+}$* mice, and the

expression of *Gzma* also downregulated in *Prdm1^{ΔNcr1}* mice (Figure 5B). Previous reports showed

that GzmA^{+} ILC1 constituted the main population of liver ILC1s at birth, with the potential

target-killing ability (46, 47).

***Prdm1* facilitates the intercellular communication between liver Type 1 ILCs and**

macrophages

The reciprocal crosstalk between Type 1 ILCs and macrophages plays a critical role in

maintaining liver immune homeostasis and anti-cancer immune surveillance. Based on the

scRNA-seq data, four clusters were generated from liver macrophages according to the

transcriptional profile. They were labeled as "*Ly6c2*^{hi}"; "*Cxcl2*^{hi}"; "*Ear2*^{hi}"; and "*C1q*^{hi}"

macrophages (Mac) (Figure 6A, Supplemental Figure 8, A-D).

High-resolution interactions among liver cNK cells, ILC1s, and macrophages were

established and compared between $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice using the CellChat program

(48). Interactions between ILC1s and macrophages were higher than that between cNK cells and

macrophages (Supplemental Figure 9, A and B). There is a significant decrease in the interaction

number between liver Type 1 ILCs (including cNK cells and ILC1s) and macrophages (~2 fold) in

$Prdm1^{\Delta Ncr1}$ mice (Figure 6, B and C). A reduction in the interaction of ligand-receptor, such as

Mif-CD74, Cxcl16-Cxcr6, and Cxcl10-Cxcr3 was observed in $Prdm1^{\Delta Ncr1}$ mice compared to

$Prdm1^{+/+}$ mice (Supplemental Figure 10). Liver cNK cells and ILC1s contributed equally to

macrophages through most pathways, while CXC chemokine ligand (CXCL), Thy-1 cell surface

antigen (THY1), and C-type lectin (CLEC) pathways participated more in ILC1-Mac interaction

than cNK-Macrophage interaction (Supplemental Figure 9, C and D). Compared to $Prdm1^{+/+}$ mice,

the information flow of CXCL and macrophage migration inhibitory factor (MIF) pathways

significantly decreased in $Prdm1^{\Delta Ncr1}$ mice (Figure 6, D and E). These pathways play a crucial role

in facilitating macrophage migration. The pathway Thrombospondin (THBS) was upregulated in

$Prdm1^{\Delta Ncr1}$, which was consistent with the higher expression of $Thbs1$ in $Prdm1^{\Delta Ncr1}$ macrophages

(Supplemental Figure 8C). The CXCL signaling was sent from $Cxcl2^{hi}$ macrophage and $C1q^{hi}$

macrophage, targeting all ILC1 clusters and $Cxcr3^{hi}$ cNK cell clusters (Figure 6F). Of note,

although the population of $Cxcl2^{hi}$ macrophage primarily comprised cells from $Prdm1^{\Delta Ncr1}$ mice,

the interaction within the CXCL pathway between macrophages and Type 1 ILCs was only

observed in $Prdm1^{+/+}$ sample (Figure 6F). These changes could be linked to a decreased

population of ILC1s and $Cxcr3^{hi}$ cNK cell cluster in $Prdm1^{\Delta Ncr1}$ mice, implying that the

homeostasis of *Cxcl2*^{hi} macrophages required sufficient signals from cNK cells and ILC1s. The

impaired CXCL-CXCR interactions might subsequently lead to reduced recruitment and

activation of Type 1 ILCs and macrophages within the tumor microenvironment.

***Prdm1* safeguards Type 1 ILCs from exhaustion-like phenotypes in the tumor**

microenvironment

The suppression of mitochondrial related pathways in *Junb*^{hi} cNK cell cluster, along with a

significant increase of this cNK cell cluster in *Prdm1*^{ΔNcr1} mice, encouraged us to explore

mitochondrial   function   through   flow   cytometry.   MitoTracker,   MitoSOX,   and

Tetramethylrhodamine methyl ester (TMRM) were used to assess the mitochondrial mass,

superoxide production, and mitochondrial membrane potential. A substantial decrease in mean

fluorescent intensities (MFIs) of MitoTracker was observed in *Prdm1*^{ΔNcr1} splenic cNK cells, liver

cNK cells, and liver ILC1s when compared to their *Prdm1*^{+/+} counterparts (Figure 6A). This

observation aligns with the enrichment of downregulated genes from *Prdm1* deficient sample in

mitochondrial related pathway, as revealed by RNA sequencing data (Supplemental Figure 6, D-F).

There was no significant difference in MitoSOX and TMRM between *Prdm1*^{ΔNcr1} and *Prdm1*^{+/+}

mice (Figure 7, B and C), which suggested that the ATP synthesize capacity was minimally

affected by *Prdm1*.

IFN-γ is a critical cytokine for NK cells mediated cancer surveillance(49, 50) and impaired

production of IFN-γ was considered as a key hallmark of exhausted NK cells (51, 52). To evaluate

the IFN-γ secreting capacity of liver cNK cells and ILC1s in tumor microenvironment, B16F10

tumor cells were inoculated to the liver via splenic injection and the IFN-γ levels in response to

stimulation of IL-12 and/or IL-18 were assessed by flow cytometry (Supplemental Figure 11).

Compared with *Prdm1^(+/+)*, significant deceased of IFN-γ were observed in *Prdm1^(ΔNcr1)* liver cNK

cells and ILC1s under the combinate stimulation of IL-12/IL-18 (Figure 7D), which was more

remarkable in liver ILC1s. Similar trends were observed when IL-12 or IL-18 was used alone,

although only liver ILC1s showed a significant decrease in response to IL-18 stimulation (Figure

7D). These findings were consistent with the heavy tumor burden observed in *Prdm1^(ΔNcr1)* mice.

Discussion

Using Ncr1-driven conditional knockout transgenic mice, which specifically delete *Prdm1* in

Type 1 ILCs, we not only validated *Prdm1*'s positive regulation of NK cell maturation, but also

demonstrated its indispensable role in NK cell anti-tumor activity. The compromised

mitochondrial function and reduced IFN-γ, granzyme B, and perforin production appear to be

potential contributing factors to Prdm1-mediated cancer surveillance. Decreased expression of

Cx3cr1 and *Cxcr6* was observed in *Prdm1^(ΔNcr1)* splenic cNK cells (Figure 3F), both of which are

essential for NK cells egressing from bone marrow (53, 54). The quantity of cNK cells increased

exclusively in the bone marrow, with reductions observed in all other tissues (Figure 1F). This

indicated that *Prdm1* might regulate chemokine receptors to facilitate the egression of cNK cells

from the bone marrow to peripheral tissues. In addition, higher expression of *Cxcr6* compared to

cNK cells is also a key factor for liver tissue residence of ILC1s (55). Decreased expression of

Cxcr6 in *Prdm1* deficient Type 1 ILCs may also contribute to the balance shift towards cNK cells.

scRNA sequencing analysis reveals that both liver cNK cells and ILC1s can be further

divided into three subgroups based on their gene expression patterns. Within the cNK cells, the

abundance of *Junb*hi cluster substantially increased in *Prdm1^ΔNcr1* mice (75.8%). Notably, these

cells account for less than 5% of WT cNK cells. High *Junb* expression stood out as a prominent

feature of this subgroup. Junb is a crucial transcriptional factor for the cytotoxic function of CD8[+]

T cells and NK cells. However, excessive *Junb* expression has been found to promote T cell

exhaustion (56). Our previous study showed that as NK cells mature, the expression level of

Prdm1 increased while the expression level of *Junb* gradually decreased (57). Our data

demonstrated that in NK cells, the expression level of *Junb* significantly increases upon the

deletion of Prdm1, indicating that *Junb* expression is suppressed by Prdm1. As *Junb* expression

decreases with NK cell maturation, and it is inhibited by the gradually increasing Prdm1 during

maturation. This implies that constraining *Junb* expression is likely a fundamental prerequisite for

NK cell maturation. However, the precise mechanism by which Prdm1 downregulates *Junb* in NK

cells still needs further research. Furthermore, *Junb*hi NK cells exhibit lower expression levels of

cytotoxic genes and reduced mitochondrial-related signaling pathways. Mitochondrial

fragmentation has also been confirmed to be closely associated with NK cell exhaustion(58).

While our current data is not sufficient to definitively classify these cells as exhausted NK cells, it

supports that this particular group of NK cells demonstrates an exhausted phenotype. The

significant increase in this cell population following *Prdm1* knockout in NK cells may potentially

be one of the reasons why *Prdm1^ΔNcr1* mice lose their tumor-killing capacity.

Our scRNA-seq data revealed that *Prdm1* plays distinct roles in regulating cNK cells and

ILC1s despite being required for both lineages. Specifically, *Prdm1* appears to be more involved

in promoting the resistance against exhaustion in cNK cells, whereas in ILC1s, it may play a role

in the plasticity between ILC1s and ILC3s. In both our previous study and a study by Colonna et

al. (57, 59), it was demonstrated that *Smad4*, a transcriptional factor involved in TGF-β signal

pathway, upregulated *Prdm1* in NK cells and depletion of *Smad4* resulted in a decreased ratio of

NK cells to ILC1s in the liver. However, knocking out *Prdm1* in Ncr1[+] cells increased the ratio of

NK cells to ILC1s in liver type 1 ILCs (Figure 2A). These findings suggest the possibility of a

Smad4-independent pathway through which *Prdm1* promotes the maintenance of ILC1s, or that

Prdm1 plays a more significant role in maintaining ILC1s compared to its role in NK cells.

Chronic inflammation is a crucial factor in promoting tumorigenesis, and macrophages play a

significant role in this process. Macrophages interact with both cNK cells and ILC1s. However,

the TFs that regulate these interactions are poorly understood. Fortunately, recent advances in

scRNA-seq technology and the CellChat software tool (48) have allowed us to gain a better

understanding of the *Prdm1* signaling pathway in Type 1 ILCs and its impact on macrophage

function at the transcriptional level. It is worth noting that normal ILC1-macrophage interactions

are more prevalent than those with NK cells (Supplemental Figure 9, A and B). Moreover, the

finding of increased liver metastasis in *Prdm1[ΔNcr1]* mice indicated the crucial role of *Prdm1* in

maintaining homeostasis between Type 1 ILCs and macrophages. Specifically, *Prdm1* may be

critical in preventing the overactivation of macrophages that can lead to cancer development.

Methods

Mice

Prdm1[fl/fl] mice were purchased from The Jackson Laboratory. *Ncr1-iCre* and *B2m[-/-]* mice

were purchased from Shanghai Model Organisms Center, Inc.. Six- to twelve-week-old littermates

were used for the experiment.

Experimental metastasis model

For lung metastasis model, 0.3×10^6 B16F10 cells were intravenous injected into mice. Three weeks later, mice were euthanized for analysis. For liver metastasis model, mice were inoculation with 0.5×10^6 B16F10 via intrasplenic injection. Three weeks later, mice were euthanized for analysis. Lung and liver from tumor-bearing mice were fixed in 10% formalin and embedded in paraffin. Sections were stained with H&E.

In vivo **cytotoxicity assay**

Donor splenocytes harvested from B2m deficient ($B2m^{-/-}$) mice were labeled with 5 μM CFDA-SE. Donor splenocytes harvested from B2m-adequate ($B2m^{+/+}$) mice were labeled with 5 μM eF670. Labeled splenocytes from two mouse strains were mixed in a 1:1 ratio, and 1×10^7 cells in total were injected i.v. into $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice. One day after administration, spleen and liver cells were isolated from recipient mice, and the population of labeled cells was analyzed by flow cytometry. Rejection % was quantified according to the following formula:

$[1-(\%B2m^{+/+}/\%B2m^{-/-})_{Prdm1^{+/+} \text{ or } Prdm1^{\Delta Ncr1}}] \times 100\%$.

Flow cytometry

Flow cytometry and cell sorting were performed with a Cytoflex S/SRT (Beckman Coulter). Data were analyzed using FlowJo software. The following antibodies were used (all purchased from BioLegend unless otherwise indicated): CD45-PE-Cy7 (catalog 103114, clone 30-F11); CD3ε-PerCP-Cy5.5 (catalog 100327, clone 145-2C11); NK1.1-BV421 (catalog 108741, clone PK136); CD335-AF647 (catalog 560755, clone 29A1.4, BD Bioscience); CD49a-PE (catalog

562115, clone Ha31/8, BD Bioscience); CD49b-FITC (catalog 108906, clone DX5);

CD27-BV510 (catalog 124229, clone LG.3A10); CD11b-AF700 (catalog 101222, clone M1/70);

KLRG1-APC (catalog 561620, clone 2F1, BD Bioscience); CD49a-BV421 (catalog 740046, clone

Ha31/8, BD Bioscience); IFN-γ-PE (catalog 505807, clone XMG1.2); Granzyme B (catalog

372207, clone QA16A02); Perforin (catalog 154305, clone S16009A); CD49b-APC-Cy7 (catalog

108919, clone DX5) ; IgG-PE (catalog 402203, clone 27-35); CD335-BV510 (catalog 137623,

clone 29A1.4); TIGIT-PE (catalog 622205, clone A17200C); CD3ε-APC-Cy7 (catalog 100330,

clone 145-2C11). For mitochondrial metabolic assay, fresh cells were incubated in 37℃ media for

30min with 100 nM MitoTracker Green (catalog M7514, Invitrogen), 100 nM TMRM (catalog

T668; Invitrogen), and 10 uM MitoSOX Red (catalog M36008; Invitrogen), respectively.

Surface-stained after washing with PBS and then detected by flow cytometry. For intracellular

IFN-γ staining, Cells freshly obtained from liver and spleen were stimulated 12 hours with or

without cytokine. GolgiStop (BD Biosciences) was added 4 hours before intracellular staining.

Bulk RNA sequencing

Total RNA from FACS sorted splenic NK cells of $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice was

extracted by TRIzol reagent (Invitrogen), then reverse transcribed into cDNA. Library

construction was prepared using Illumina mRNA Library kit, and sequencing was performed by

the BGISEQ-500. Standard methods were used to analyze the RNA-seq data, including aligning

the reads to the genome by HISAT2 (v2.1.0) (60), and gene expression values (Counts) were

calculated using RSEM (v1.3.1) (61). DEGs were identified using DEseq2 (v1.4.5) (62) with a

cutoff of $\log_2$(fold change) > 0.5 and P < 0.05. The "clusterProfiler" package (v4.4.4) (63) and

gene sets from molecular signatures database (MSigDB) were used for GSEA and GO analysis.

The heatmap was plotted using the "Pheatmap" package (v1.0.12).

Single-cell RNA sequencing

FACS-sorted liver CD45$^+$ cells with more than 80% cell viability were used for library

preparation. Each sample contained cells from three *Prdm1$^{+/+}$* or *Prdm1$^{\Delta Ncr1}$* mice. Gel

Bead-in-Emulsions (GEMs) were generated using the 10X Genomics Chromium system, which

combinates Master Mix, Single Cell 3' v3.1 Gel Beads, and Partitioning Oil with single cells.

GEMs were mixed with cell lysate and reverse transcription reagent to produce full-length cDNA.

After incubation, the GEMs were broken, and recovered cDNA were amplified via PCR.

Fragmentation, End repair, A-tailing, and Adaptor Ligation were performed to obtain final

libraries, which contain P5 and P7 sequences. The 3' library was sequenced on Novaseq 6000 with

approximately 50k read pairs/cell sequencing depth. The "Seurat" R package (v4.2.0) (64) was

used for data analysis. Initial quality control was performed to filter out the low-quality cells and

cell doublets. Cells with 200-5500 expressed genes and no more than 10% mitochondrial genes

were considered for high-quality. Doublets were filtered with the "scDblFinder" Package (v1.10.0)

(65). After quality control, we totally recovered 6161 cells and 4817 cells from *Prdm1$^{+/+}$* and

Prdm1$^{\Delta Ncr1}$ mice, respectively. Principal component analysis (PCA) was used for cluster analysis.

The first 15 PCs were picked for clustering and further visualized by UMAP. Clusters-specific

marker was defined using the "FindAllMarkers" function, and clusters were manually annotated

based on the top 30 or 15 markers. The "clusterProfiler" package and gene set from MSigDB were

used for GSEA and GO analysis. "CellChat" package (v1.4.0) (48) was utilized to predict the

cell-to-cell communication from scRNA-seq data.

TCGA datasets assay

The normalized gene expression and survival datasets of cancer patients collected in The

Cancer Genome Atlas (TCGA) were downloaded from UCSC Xena (http://xena.ucsc.edu/) (66).

The mean expression of NK cell-associated genes (*NCR1*, *NCR3*, *PRF1*, *KLRB1*, *CD160*) was

defined as NK signature according to the previous study (32). NK cell-associated genes, together

with *PRDM1*, constitute the *NK-PRDM1* signature in this study. The mean expression of per genes

was ordered from high-to-low and plotted by heatmap using the "Pheatmap" package. The overall

survival of patients in the high and low quartiles expression of *NK-PRDM1* signature was selected

for analysis. Kaplan-Meier curves were plotted by GraphPad Prism.

Statistics

For experiment results, two-tailed t tests were used to measure the continuous and normally

distributed between the two independent groups. Paired t-tests were used to determine the

statistical significance between two paired groups. Log-rank tests were used to compare the

overall survival distribution between the two groups of patients. A P value less than 0.05 was

considered significant and data were presented as mean±SEM.

Study approval

All animal experiments were approved by The Tianjin University Animal Care and Use

Committee (protocol 00000000202010100023). No human subjects were performed in this study.

Author contributions

544   Jitian He, and Youwei Wang designed experiments, performed experiments, analyzed the data

545   and wrote the manuscript. Jitian He, Le Gao, Peiying Wang, Yiran Zheng, and Yumo Zhang

546   performed experiments. Wing Keung Chan, Jiming Wang, Huaiyong Chen, and Zhouxin Yang

547   analyzed the data, interpreted results and reviewed the manuscript.

Acknowledgments

This work was supported by The National Key Research and Development Program of China (2022YFF1202900), and The Zhejiang Provincial Natural Science Foundation of China (LY21H150002).

Declaration of interests

555   The authors have declared that no conflict of interest exists.

References

1.   Peng H, Jiang X, Chen Y, Sojka DK, Wei H, Gao X, et al. Liver-resident NK cells confer

adaptive immunity in skin-contact inflammation. *J Clin Invest.* 2013;123(4):1444-56.

2.   Bai L, Vienne M, Tang L, Kerdiles Y, Etiennot M, Escaliere B, et al. Liver type 1 innate

lymphoid cells develop locally via an interferon-gamma-dependent loop. *Science.*

2021;371(6536).

3.   Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C, et al. ILC1 Confer Early

Host Protection at Initial Sites of Viral Infection. *Cell.* 2017;171(4):795-808 e12.

4.   Schuster IS, Wikstrom ME, Brizard G, Coudert JD, Estcourt MJ, Manzur M, et al. TRAIL+

NK cells control CD4+ T cell responses during chronic viral infection to limit autoimmunity.

Immunity. 2014;41(4):646-56.

5.   Ducimetiere L, Lucchiari G, Litscher G, Nater M, Heeb L, Nunez NG, et al. Conventional NK

cells and tissue-resident ILC1s join forces to control liver metastasis. *Proc Natl Acad Sci U S*

A. 2021;118(27).

6.   Fumagalli V, Venzin V, Di Lucia P, Moalli F, Ficht X, Ambrosi G, et al. Group 1 ILCs regulate

T cell-mediated liver immunopathology by controlling local IL-2 availability. *Science*

immunology. 2022;7(68):eabi6112.

7.   Kiessling R, Klein E, and Wigzell H. "Natural" killer cells in the mouse. I. Cytotoxic cells

with specificity for mouse Moloney leukemia cells. Specificity and distribution according to

genotype. *European journal of immunology.* 1975;5(2):112-7.

8.   Nabekura T, Riggan L, Hildreth AD, O'Sullivan TE, and Shibuya A. Type 1 Innate Lymphoid

Cells Protect Mice from Acute Liver Injury via Interferon-gamma Secretion for Upregulating

Bcl-xL Expression in Hepatocytes. *Immunity.* 2020;52(1):96-108 e9.

9.   Takeda K, Cretney E, Hayakawa Y, Ota T, Akiba H, Ogasawara K, et al. TRAIL identifies

immature natural killer cells in newborn mice and adult mouse liver. *Blood.*

2005;105(5):2082-9.

10.   Takeda K, Hayakawa Y, Smyth MJ, Kayagaki N, Yamaguchi N, Kakuta S, et al. Involvement

of tumor necrosis factor-related apoptosis-inducing ligand in surveillance of tumor metastasis

by liver natural killer cells. *Nat Med.* 2001;7(1):94-100.

11.   Townsend MJ, Weinmann AS, Matsuda JL, Salomon R, Farnham PJ, Biron CA, et al. T-bet

regulates the terminal maturation and homeostasis of NK and Valpha14i NKT cells. *Immunity.*

2004;20(4):477-94.

12.   Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, et al. Development of

peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor

Id2. *Nature.* 1999;397(6721):702-6.

13.   Gascoyne DM, Long E, Veiga-Fernandes H, de Boer J, Williams O, Seddon B, et al. The basic

leucine zipper transcription factor E4BP4 is essential for natural killer cell development.

Nature immunology. 2009;10(10):1118-24.

14.   Constantinides MG, McDonald BD, Verhoef PA, and Bendelac A. A committed precursor to

innate lymphoid cells. *Nature.* 2014;508(7496):397-401.

15.   Gordon SM, Chaix J, Rupp LJ, Wu J, Madera S, Sun JC, et al. The transcription factors T-bet

and Eomes control key checkpoints of natural killer cell maturation. *Immunity.*

2012;36(1):55-67.

16.   Mackay LK, Minnich M, Kragten NA, Liao Y, Nota B, Seillet C, et al. Hobit and Blimp1

601   instruct a universal transcriptional program of tissue residency in lymphocytes. *Science.*

602   2016;352(6284):459-63.

603 17. Lee JC, Green MD, Huppert LA, Chow C, Pierce RH, and Daud AI. The Liver-Immunity

604   Nexus and Cancer Immunotherapy. *Clin Cancer Res.* 2022;28(1):5-12.

605 18. Galon J, and Bruni D. Approaches to treat immune hot, altered and cold tumours with

606   combination immunotherapies. *Nature reviews Drug discovery.* 2019;18(3):197-218.

607 19. Ruff SM, Shannon AH, Beane JD, and Pawlik TM. Highlighting novel targets in

608   immunotherapy for liver cancer. *Expert review of gastroenterology & hepatology.*

609   2022;16(11-12):1029-41.

610 20. He Y, Zhan L, Shi J, Xiao M, Zuo R, Wang C, et al. The Combination of R848 with Sorafenib

611   Enhances Antitumor Effects by Reprogramming the Tumor Immune Microenvironment and

612   Facilitating Vascular Normalization in Hepatocellular Carcinoma. *Adv Sci (Weinh).*

613   2023:e2207650.

614 21. Nersesian S, Schwartz SL, Grantham SR, MacLean LK, Lee SN, Pugh-Toole M, et al. NK cell

615   infiltration is associated with improved overall survival in solid cancers: A systematic review

616   and meta-analysis. *Transl Oncol.* 2021;14(1):100930.

617 22. Lee HA, Goh HG, Lee YS, Jung YK, Kim JH, Yim HJ, et al. Natural killer cell activity is a

618   risk factor for the recurrence risk after curative treatment of hepatocellular carcinoma. *BMC*

619   *Gastroenterol.* 2021;21(1):258.

620 23. Gao Y, Souza-Fonseca-Guimaraes F, Bald T, Ng SS, Young A, Ngiow SF, et al. Tumor

621   immunoevasion by the conversion of effector NK cells into type 1 innate lymphoid cells. *Nat*

622   *Immunol.* 2017;18(9):1004-15.

24.   Cortez VS, Ulland TK, Cervantes-Barragan L, Bando JK, Robinette ML, Wang Q, et al.

SMAD4 impedes the conversion of NK cells into ILC1-like cells by curtailing non-canonical

TGF-beta signaling. *Nature immunology.* 2017;18(9):995-1003.

25.   Wang Y, Chu J, Yi P, Dong W, Saultz JN, Wang Y, et al. SMAD4 promotes

TGF-beta-independent NK cell homeostasis and maturation and antitumor immunity. *The*

Journal of clinical investigation. 2018.

26.   Keller AD, and Maniatis T. Identification and characterization of a novel repressor of

beta-interferon gene expression. *Genes Dev.* 1991;5(5):868-79.

27.   Kallies A, Hawkins ED, Belz GT, Metcalf D, Hommel M, Corcoran LM, et al. Transcriptional

repressor Blimp-1 is essential for T cell homeostasis and self-tolerance. *Nat Immunol.*

2006;7(5):466-74.

28.   Martins GA, Cimmino L, Shapiro-Shelef M, Szabolcs M, Herron A, Magnusdottir E, et al.

Transcriptional repressor Blimp-1 regulates T cell homeostasis and function. *Nat Immunol.*

2006;7(5):457-65.

29.   Kallies A, Carotta S, Huntington ND, Bernard NJ, Tarlinton DM, Smyth MJ, et al. A role for

Blimp1 in the transcriptional network controlling natural killer cell maturation. *Blood.*

2011;117(6):1869-79.

30.   Akman B, Hu X, Liu X, Hatipoglu T, You H, Chan WC, et al. PRDM1 decreases sensitivity of

human NK cells to IL2-induced cell expansion by directly repressing CD25 (IL2RA). *Journal*

of leukocyte biology. 2021;109(5):901-14.

31.   Smith MA, Maurin M, Cho HI, Becknell B, Freud AG, Yu J, et al. PRDM1/Blimp-1 controls

effector cytokine production in human NK cells. *J Immunol.* 2010;185(10):6058-67.

32. Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrerizo M, Sammicheli S, et al.

NK Cells Stimulate Recruitment of cDC1 into the Tumor Microenvironment Promoting

Cancer Immune Control. *Cell.* 2018;172(5):1022-37 e14.

33. Shapiro-Shelef M, Lin K-I, McHeyzer-Williams LJ, Liao J, McHeyzer-Williams MG, and

Calame K. Blimp-1 Is Required for the Formation of Immunoglobulin Secreting Plasma Cells

and Pre-Plasma Memory B Cells. *Immunity.* 2003;19(4):607-20.

34. Hayakawa Y, and Smyth MJ. CD27 dissects mature NK cells into two subsets with distinct

responsiveness and migratory capacity. *J Immunol.* 2006;176(3):1517-24.

35. Chiossone L, Chaix J, Fuseri N, Roth C, Vivier E, and Walzer T. Maturation of mouse NK

cells is a 4-stage developmental program. *Blood.* 2009;113(22):5488-96.

36. Huntington ND, Tabarias H, Fairfax K, Brady J, Hayakawa Y, Degli-Esposti MA, et al. NK

cell maturation and peripheral homeostasis is associated with KLRG1 up-regulation. *J

Immunol.* 2007;178(8):4764-70.

37. Bix M, Liao NS, Zijlstra M, Loring J, Jaenisch R, and Raulet D. Rejection of class I

MHC-deficient haemopoietic cells by irradiated MHC-matched mice. *Nature.*

1991;349(6307):329-31.

38. Bern MD, Parikh BA, Yang L, Beckman DL, Poursine-Laurent J, and Yokoyama WM.

Inducible down-regulation of MHC class I results in natural killer cell tolerance. *The Journal

of experimental medicine.* 2019;216(1):99-116.

39. Karre K, Ljunggren HG, Piontek G, and Kiessling R. Selective rejection of H-2-deficient

lymphoma variants suggests alternative immune defence strategy. *Nature.*

1986;319(6055):675-8.

40.   Viant C, Fenis A, Chicanne G, Payrastre B, Ugolini S, and Vivier E. SHP-1-mediated

668   inhibitory signals promote responsiveness and anti-tumour functions of natural killer cells.

669   *Nat Commun.* 2014;5:5108.

670   41.   Cillo C, Dick JE, Ling V, and Hill RP. Generation of drug-resistant variants in metastatic B16

671   mouse melanoma cell lines. *Cancer Res.* 1987;47(10):2604-8.

672   42.   Spits H, Artis D, Colonna M, Diefenbach A, Di Santo JP, Eberl G, et al. Innate lymphoid

673   cells--a proposal for uniform nomenclature. *Nat Rev Immunol.* 2013;13(2):145-9.

674   43.   Park E, Patel S, Wang Q, Andhey P, Zaitsev K, Porter S, et al. Toxoplasma gondii infection

675   drives conversion of NK cells into ILC1-like cells. *Elife.* 2019;8.

676   44.   Chen Y, Wang X, Hao X, Li B, Tao W, Zhu S, et al. Ly49E separates liver ILC1s into

677   embryo-derived and postnatal subsets with different functions. *J Exp Med.* 2022;219(5).

678   45.   Kirchhammer N, Trefny MP, Natoli M, Brucher D, Smith SN, Werner F, et al. NK cells with

679   tissue-resident traits shape response to immunotherapy by inducing adaptive antitumor

680   immunity. *Sci Transl Med.* 2022;14(653):eabm9043.

681   46.   Di Censo C, Marotel M, Mattiola I, Muller L, Scarno G, Pietropaolo G, et al. Granzyme A and

682   CD160 expression delineates ILC1 with graded functions in the mouse liver. *European*

683   *journal of immunology.* 2021;51(11):2568-75.

684   47.   Friedrich C, Taggenbrock R, Doucet-Ladeveze R, Golda G, Moenius R, Arampatzi P, et al.

685   Effector differentiation downstream of lineage commitment in ILC1s is driven by Hobit across

686   tissues. *Nat Immunol.* 2021;22(10):1256-67.

687   48.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and

688   analysis of cell-cell communication using CellChat. *Nat Commun.* 2021;12(1):1088.

689   49.   Takeda K, Nakayama M, Sakaki M, Hayakawa Y, Imawari M, Ogasawara K, et al.

690   IFN-gamma production by lung NK cells is critical for the natural resistance to pulmonary

691   metastasis of B16 melanoma in mice. *Journal of leukocyte biology.* 2011;90(4):777-85.

692   50.   Lin Q, Rong L, Jia X, Li R, Yu B, Hu J, et al. IFN-gamma-dependent NK cell activation is

693   essential to metastasis suppression by engineered Salmonella. *Nature communications.*

694   2021;12(1):2537.

695   51.   Roe K. NK-cell exhaustion, B-cell exhaustion and T-cell exhaustion-the differences and

696   similarities. *Immunology.* 2022;166(2):155-68.

697   52.   Zhang Q, Bi J, Zheng X, Chen Y, Wang H, Wu W, et al. Blockade of the checkpoint receptor

698   TIGIT prevents NK cell exhaustion and elicits potent anti-tumor immunity. *Nature*

699   *Immunology.* 2018;19(7):723-32.

700   53.   Sciume G, De Angelis G, Benigni G, Ponzetta A, Morrone S, Santoni A, et al. CX3CR1

701   expression defines 2 KLRG1+ mouse NK-cell subsets with distinct functional properties and

702   positioning in the bone marrow. *Blood.* 2011;117(17):4467-75.

703   54.   Chea S, Possot C, Perchet T, Petit M, Cumano A, and Golub R. CXCR6 Expression Is

704   Important for Retention and Circulation of ILC Precursors. *Mediators Inflamm.*

705   2015;2015:368427.

706   55.   Hudspeth K, Donadon M, Cimino M, Pontarini E, Tentorio P, Preti M, et al. Human

707   liver-resident CD56(bright)/CD16(neg) NK cells are retained within hepatic sinusoids via the

708   engagement of CCR5 and CXCR6 pathways. *J Autoimmun.* 2016;66:40-50.

709   56.   Lynn RC, Weber EW, Sotillo E, Gennert D, Xu P, Good Z, et al. c-Jun overexpression in CAR

710   T cells induces exhaustion resistance. *Nature.* 2019;576(7786):293-300.

57.   Wang Y, Chu J, Yi P, Dong W, Saultz J, Wang Y, et al. SMAD4 promotes

TGF-beta-independent NK cell homeostasis and maturation and antitumor immunity. *J Clin*

Invest. 2018;128(11):5123-36.

58.   Zheng X, Qian Y, Fu B, Jiao D, Jiang Y, Chen P, et al. Mitochondrial fragmentation limits NK

cell-based tumor immunosurveillance. *Nature immunology.* 2019;20(12):1656-67.

59.   Cheng HP, Huang CJ, Tsai ML, Ong HT, Cheong SK, Choo KB, et al. MicroRNA-362

negatively and positively regulates SMAD4 expression in TGF-beta/SMAD signaling to

suppress cell migration and invasion. *Int J Med Sci.* 2021;18(8):1798-809.

60.   Kim D, Langmead B, and Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. *Nat Methods.* 2015;12(4):357-60.

61.   Li B, and Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics.* 2011;12:323.

62.   Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

63.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment

tool for interpreting omics data. *Innovation (Camb).* 2021;2(3):100141.

64.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated

analysis of multimodal single-cell data. *Cell.* 2021;184(13):3573-87 e29.

65.   Germain PL, Lun A, Garcia Meixide C, Macnair W, and Robinson MD. Doublet identification

in single-cell sequencing data using scDblFinder. *F1000Res.* 2021;10:979.

66.   Caicedo HH, Hashimoto DA, Caicedo JC, Pentland A, and Pisano GP. Overcoming barriers to

early disease intervention. *Nat Biotechnol.* 2020;38(6):669-73.

Figures legends

Figure 1. *Prdm1* promotes type 1 ILCs homeostasis and terminal maturation.

(**A**) Correlation between the average expression of NK cell-associated genes(*NCR1*, *NCR3*, *PRF1*, *CD160*, *KLRB1*) and *PRDM1* in LIHC (Liver Hepatocellular Carcinoma; n=363) patients sourced from TCGA datasets. (**B**) Heatmap of the ordered, z-score normalized expression values for *PRDM1* and NK cell-associated genes in liver cancer patients. The top and bottom quartiles of all patient samples are indicated. (**C**) Prognostic value of the *NK-PRDM1* signature for overall survival of liver cancer patients comparing high and low quartiles. (**D**) Schematic representation of the Prdm1-conditional knockout mouse model. Targeted exon 6-8 of *Prdm1* (top) is flanked with *loxP* sites (middle). Ncr1-expressed Cre recombinase was used to generate the $Prdm1^{\Delta Ncr1}$ allele (bottom). (**E**) Real-time RT-PCR quantification of *Prdm1* expression in NKp46$^+$ cells to determine the presence of *Prdm1* (n=5). (**F**) Representative flow cytometric plots (left) and quantification (right) of the proportion and absolute number of CD3$^-$NKp46$^+$ cells among lymphocytes in liver, lung, and bone marrow (n=8). (**G**) Representative flow cytometric plots (left) of the CD11b and CD27 expression within CD3$^-$NK1.1$^+$NKp46$^+$ cells in liver, lung, and bone marrow (n=7). Right panel showed the percentage of distinct stages of NK cells. Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; r, pearson correlation coefficient; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 2. *Prdm1* is required for type I ILCs to control tumor metastasis.

(**A**) Representative flow cytometric plots (left) of liver cNK cells (CD49a⁻CD49b⁺) and ILC1s

(CD49a⁺CD49b⁻) from *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* mice. The two bar graghs (right) quantitate the

percentages and absolute numbers of cells respectively (n=7). (**B and C**) Splenocytes from *B2m⁻/⁻*

and *B2m⁺/⁺* were labeled with CFDASE and eF670 respectively. Labeled cells were 1:1 mixed and

injected i.v. into *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* recipient mice to evaluate *in vivo* NK cell target-killing

ability. Representative flow cytometric plots (left) of transferred cells recovered from recipient

mice and percentage (right) of NK cell-specific rejection of donor cells in spleen (n=6) (**B**) and

liver (n=7) (**C**) between *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* mice. (**D and F**) Image (left) and quantification

(right) of tumor nodes on the livers (n=7) (**D**) and lungs (n=10) (**F**) of *Prdm1⁺/⁺* and *Prdm1ᐞNcr1*

mice at day 14 or 21 after inoculation with B16F10 melanoma cells. (**E and G**) Histopathological

images of liver (**E**) and lung (**G**) tissues stained by hematoxylin-eosin to detect tumor metastasis.

Red bar indicates 500 μm distance under the microscope. (**H**) Liver cells and splenocytes were

costimulated in the presence or absence of IL-12 and IL-18 for 12 hours. GolgiStop was added 4

hours before intracellular staining of IFN-γ. The graphs showed percentage of IFN-γ⁺ splenic cNK

cells, liver cNK cells and ILC1s from *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* mice (n=5). Data are presented as

the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between

littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *,

P<0.05; **, *P*<0.01, ns, not significant.

Figure 3. Bulk RNA-seq depicts *Prdm1*-mediated functions in peripheral cNK cells.

(**A**) Splenic cNK cells and liver CD45⁺ cells were sorted from *Prdm1⁺/⁺* and *Prdm1ᐞNcr1* mice

using flow cytometry, and prepared for bulk RNA-seq and single-cell RNA-seq analysis. (**B**)

Volcano plot of the bulk RNA-seq differentially expressed genes ($\log_2$(fold change)>0.5; *P*<0.05)

in splenic cNK cells between *Prdm1*[+/+] and *Prdm1*[ΔNcr1] mice. Upregulated and downregulated

genes in *Prdm1*[ΔNcr1] cells were highlighted in red and blue. (**C**) Enriched Gene Ontology (GO)

terms of DEGs in *Prdm1*[ΔNcr1] cells compared *Prdm1*[+/+] cells. The Enrichment gene set in

upregulated (red) and downregulated (blue) genes were indicated in different colours. Bar length

represents statistical significance. (**D and E**) Gene Set Enrichment Analysis (GSEA) showing the

enrichment of negative regulation of IL-6 production (**D**) and NF-kappa B signaling pathway (**E**)

of DEGs in *Prdm1*[ΔNcr1] cells compared *Prdm1*[+/+] cells. NES, normalized enrichment score. (**F**)

Heatmap of selected genes from DEGs. Shown is z-score transformed expression of DEGs. (**G**

and **I**) Representative flow cytometric plots (left) and cumulative data (right) showing the relative

mean fluorescence intensities (MFIs) of Granzyme B (**G**) and Perforin (**I**) in liver cNK cells and

ILC1s from *Prdm1*[+/+] and *Prdm1*[ΔNcr1] mice (n=5). (**H and J**) Relative MFIs of Granzyme B (**H**)

and Perforin (**J**) in splenic NK cells at different stages of maturation was analyzed by flow

cytometry (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test.

Differences were evaluated between littermates. Each circle and square on graphs represents an

individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 4. scRNA-seq reveals unique properties of two clusters from liver type I ILCs

following *Prdm1* knockout.

(**A**) Uniform manifold approximation and projection (UMAP) visualization of liver CD45[+] cells

from *Prdm1*[+/+] and *Prdm1*[ΔNcr1] mice. Twelve clusters were defined and indicated by distinct

colours. Each dot represents a single cell. (**B**) UMAP visualization of liver cNK and ILC1 clusters.

Cells were colored by origins (*Prdm1$^{+/+}$*-blue; *Prdm1$^{\Delta Ncr1}$*-red). (**C**) Percentages of cNK cells and

ILC1s in type I ILCs (left), and their distribution in each cluster (right). (**D-I**) UMAP visualization

of three different liver cNK (**D**) and ILC1 (**G**) clusters from two mouse strains (**E** and **H**).

Proportions of cNK cells (**F**) and ILC1s (**I**) among total cells (cNK cells or ILC1s) (left) and

within clusters (right) were calculated respectively. (**J**) Violin plots showing the normalized

expression of select genes in different cNK clusters. (**K**) Enriched GO term of marker genes in

three cNK clusters. Dot size represents enriched gene number, and color intensity represents

significance.

Figure 5. *Junb*^{hi} signature is associated with the activation of multiple signaling pathways.

(**A**) Ridge plots showing the normalized expression of *Gzmb, Prf1*, and *Junb* in cNK and ILC1

clusters between *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$* cells. (**B**) Violin plots showing the normalized

expression of select genes in different ILC1 clusters. (**C**) Violin plot showing the *Junb*^{hi} signature

score for cNK cell and ILC1 clusters, calculated using the signature genes of *Junb*^{hi} cNK cluster.

(**D**) Enriched GO term of marker genes in three ILC1 clusters. Dot size represents enriched gene

number, and color intensity represents significance. (**E-J**) GSEA plots (left) depicting the

enrichment of NF-kappa B (**E**) , TNF (**F**) , IL-17 (**G**) , and MAPK (**H**) signaling pathway in *Junb*^{hi}

cNK cluster compared with clusters of *Prf1*^{hi} and *Cxcr3*^{hi} cNK cells, and the enrichment of IL-17

signaling pathway (**I**) and T cell differentiation (**J**) in *Il7r*^{hi} ILC1 cluster compared with clusters of

Klra^{hi} and *Gzma*^{hi} ILC1s. Right panel showed dynamic relative expression of the given gene sets

from cluster1 to cluster3 between *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$*. Dots represent the average expression

of given gene set in each cell, which was calculated through the sum of normalized expression of

each individual gene within the designated gene set in every single cell. NES, normalized

enrichment score.

Figure 6. *Prdm1* facilitates the intercellular communication between liver type I ILCs and

macrophages.

(**A**) UMAP visualization of macrophages (Mac) cluster. Four cell subsets were defined. (**B** and **C**)

Circle plots (**B**) and summary data (**C**) illustrating the significant enriched ligand–receptor pairs

among cluster of liver cNK cells, ILC1s, and macrophages from *Prdm1*$^{+/+}$ (left) and *Prdm1*$^{\Delta Ncr1}$

(right) cells. The thickness of the line indicates the number of enrich pairs, and the arrow reflects

the direction of the interaction. (**D**) Heatmap of overall signaling pattern recognized from

ligand-receptor pairs, which contained the sum of signaling from the sender and target cells. (**E**)

Bar graphs showing the information flow in selected active signaling patterns between *Prdm1*$^{+/+}$

and *Prdm1*$^{\Delta Ncr1}$ cells. Relative information flow was calculated as the sum of the communication

probability in given signaling patterns. (**F**) Chord plot of the CXCL signaling interaction network

among cluster of liver cNK cells, ILC1s, and macrophages in *Prdm1*$^{+/+}$ cells.

Figure 7. *Prdm1* safeguards type I ILCs from exhaustion-like phenotypes in the tumor

microenvironment.

(**A-C**) The Mitochondrial mass (MitoTracker Green staining; n=8) (**A**), Mitochondrial ROS

(MitoSOX staining; n=5) (B), and Mitochondrial membrane potential (TMRM staining; n=5) (**C**)

of splenic cNK cells, liver cNK cells and ILC1s were analyzed by flow cytometry. Representative

flow cytometric plots (left) and cumulative data (right) showing the relative mean fluorescence

intensities (MFIs) of each group. (**D**) Percentages of IFN-γ$^+$ liver cNK cells and ILC1s from

Prdm1$^{+/+}$ and *Prdm1^{ΔNcr1}* tumor-bearing mice at day 14 after inoculation with B16F10 melanoma

cells via intrasplenic injection (n=5). Data are presented as the mean±SEM and were analyzed by

2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on

graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Supplemental Figure 1. High expression of *PRDM1-NK* signature predicts better overall

survival of cancer patients.

(**A and D**) Correlation between the average expression of NK cell-associated genes(*NCR1*, *NCR3*,

PRF1, *CD160*, *KLRB1*) and *PRDM1* in skin cutaneous melanoma (SKCM, n=454) (**A**) and lung

adenocarcinoma (**D**) (LUAD, n=497) patients from TCGA datasets. (**B and E**) Heatmap of the

ordered, z-score normalized expression values for *PRDM1* and NK cell-associated genes in skin

cutaneous melanom (**B**) and lung cancer (**E**) patients. The top and bottom quartiles of all patient

samples are indicated. (**C and F**) Prognostic value of the *NK-PRDM1* signature for overall

survival of skin cutaneous melanom (**C**) and lung cancer (**F**) patients comparing high and low

quartiles. *P*, *P*-value; r, pearson correlation coefficient.

Supplemental Figure 2. *Prdm1* plays an essential role in type I ILCs homeostasis and

maturation.

(**A**) Representative flow cytometric plots (left) and quantification (right) of the proportion and

absolute number of CD3$^-$NKp46$^+$ cells among lymphocytes in blood, spleen, and lymph nodes

(n=8). (**B**) Representative flow cytometric plots (left) of the CD11b and CD27 expression within

CD3$^-$NK1.1$^+$NKp46$^+$ cells in blood, spleen, and lymph nodes (n=7). Right panel showed the

percentage of distinct stages of NK cells. **(C and D)** Representative flow cytometry analyses **(C)**

and quantification **(D)** of KLRG1 expression in CD3⁻NK1.1⁺NKp46⁺ cells in blood, bone marrow,

lung, liver, spleen, and lymph nodes (n=5). Data are presented as the mean±SEM and were

analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and

square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not

significant.

Supplemental Figure 3. *Prdm1* affects the IFN-γ secretion ability of type I ILCs.

(A) Representative flow cytometric plot showing the frequency of IFN-γ⁺ cNK cells and ILC1s in

spleen and liver between *Prdm1*⁺/⁺ and *Prdm1*^{ΔNcr1} mice (n=5). Liver cells and splenocytes were

costimulated in the presence or absence of IL-12 and IL-18 for 12 hours. GolgiStop was added 4

hours before intracellular staining of IFN-γ. Data are presented as the mean±SEM and were

analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and

square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not

significant.

Supplemental Figure 4. *Prdm1* deficiency impairs production of Granzyme B and Perforin in

type I ILC.

(A and B) Representative flow cytometric plot (left) and summary data (right) showing the

relative mean fluorescence intensities (MFIs) of Granzyme B **(A)** and Perforin **(B)** in splenic NK

cells from *Prdm1*⁺/⁺ and *Prdm1*^{ΔNcr1} mice (n=5). **(C)** Splenic NK cells were divided into different

stages according to the expression of maturation markers CD11b and CD27. Representative flow

cytometric plot showing the MFIs of Granzyme B and Perforin in evaluated between littermates.

Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences

were evaluated between littermates. Each circle and square on graphs represents an individual

mouse; *P*, *P*-value; r, pearson correlation coefficient; *, $P<0.05$; **, $P<0.01$, ns, not significant.

Supplemental Figure 5. scRNA-seq identified subsets of liver CD45$^+$ cell from *Prdm1$^{+/+}$* and

***Prdm1$^{\Delta Ncr1}$* mice.**

(**A**) Liver CD45$^+$ cells from *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$* mice were FACS-sorted for scRNA-seq. (**B**)

Quality control of the scRNA-seq data. Violin plot showing the nFeature_RNA, nCount_RNA,

and percentage of mitochondrial genes of sequencing cells. nFeature_RNA represents the number

of genes detected in each cell, and nCount_RNA represents the total number of molecules

detected within a cell. (**C**) Population of twelve liver CD45$^+$ cell clusters between *Prdm1$^{+/+}$* and

Prdm1$^{\Delta Ncr1}$ mice. (**D**) Dot plot showing the normalized expression of marker genes in liver cNK

cells and ILC1s clusters between *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$*. The dot size represents the percentage

of cells expressing selected genes, and color intensity represents the average expression. (**E**)

Violin plots showing the expression of marker genes *Itga2, Itga1, Eomes, and Tnfsf10* between

liver cNK cells and ILC1s between *Prdm1$^{+/+}$* and *Prdm1$^{\Delta Ncr1}$* mice.

Supplemental Figure 6. Cluster-specific markers of liver cNK cell clusters.

(**A**) Heatmap showing the expression of top 30 upregulated DEGs by log fold change (computed

using Wilcox test in the "FindAllMarkers" function of Seurat, avg_log2FC>0.25; P<0.05), across

the three liver cNK cell clusters (*Prf1^{hi}* cNK, *Junb^{hi}* cNK, *Cxcr3^{hi}* cNK) within *Prdm1$^{+/+}$* and

Prdm1$^{\Delta Ncr1}$ mice. (**B**) Feature plots showing the normalized expression of selected markers for

liver cNK cell populations. (**C**) Violin plots showing the normalized expression of DEGs for each

cNK cluster within $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice. (**D-F**) GSEA of the enrichment of

Mitochondrial protein (**D**), Oxidative phosphorylation (**E**), and Respiratory electron transport

chain (**F**) in $Junb^{hi}$ cNK cluster compared with clusters of $Prf1^{hi}$ and $Cxcr3^{hi}$ cNK cells. NES,

normalized enrichment score.

Supplemental Figure 7. Cluster-specific markers of liver ILC1 clusters.

(**A**) Heatmap showing the expression of top 30 upregulated DEGs by log fold change (computed

using Wilcox test in the "FindAllMarkers" function of Seurat, avg_log2FC>0.25; P<0.05), across

the three ILC1 clusters ($Il7r^{hi}$ ILC1, $Klra^{hi}$ ILC1, $Gzma^{hi}$ ILC1) within $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$

mice. (**B**) Feature plots showing the normalized expression of selected markers for liver ILC1

populations. (**C**) Normalized expression of DEGs for each ILC1 cluster within $Prdm1^{+/+}$ and

$Prdm1^{\Delta Ncr1}$ mice. (D) GSEA plots (left) depicting the enrichment of NF-kappa B (**D**) , TNF (**E**),

and MAPK (**F**) signaling pathways) in $Il7r^{hi}$ ILC1 cluster compared with clusters of $Klra^{hi}$ and

$Gzma^{hi}$ ILC1s. Right panel showed dynamic relative expression of the given gene sets from

cluster1 to cluster3 between $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$. Dots represent the average expression of

given gene set in each cell, which was calculated through the sum of normalized expression of

each individual gene within the designated gene set in every single cell. NES, normalized

enrichment score.

Supplemental Figure 8. Identification of four distinct macrophage clusters.

(**A**) Heatmap showing the top 15 upregulated DEGs across four macrophage clusters (computed

using Wilcox test in the "FindAllMarkers" function of Seurat, avg_log2FC>0.25; P<0.05). (**B**)

Percentage of macrophages from $Prdm1^{+/+}$ and $Prdm1^{\Delta Ncr1}$ mice among total cells (left) and

within clusters (right). (**C**) Violin plots showing the expression of selected markers of each

macrophage cluster within *Prdm1$^{+/+}$* and *Prdm1^{ΔNcr1}* mice. (**D**) GO analysis of four distinct

macrophage clusters. Dot size represents enriched gene number, and color intensity represents

significance.

Supplemental Figure 9. Cell-Cell communication between cNK cells-macrophages and

ILC1s-macrophages.

(**A and B**) Circle plots (**A**) and cumulative data (**B**) showing the interaction numbers between

cNK cells-macrophages and ILC1s-macrophages. The thickness of the line indicates the number

of enrich pairs, and the arrow reflects the direction of the interaction. (**C**) Heatmap of overall

signaling pattern recognized from ligand-receptor pairs between cNK cells-macrophages and

ILC1s-macrophages, which contained the sum of signaling from the sender and target cells. (**D**)

Bar graphs showing the information flow in selected active signaling patterns between cNK

cells-macrophages and ILC1s-macrophages. Relative information flow was calculated as the sum

of the communication probability in given signaling patterns.

Supplemental Figure 10. Ligand-receptor interaction between type I ILCs and macrophages.

Bubble plots showing the significant ligand-receptor pairs between cNK cells, ILC1s, and

macrophages within *Prdm1$^{+/+}$* and *Prdm1^{ΔNcr1}* mice. The highlighted Mif-Cd74 and Cxcl-Cxcr

signaling was significantly decreased in *Prdm1^{ΔNcr1}* mice. Dot size represents the *P*-value, and

color intensity represents the communication probabilities. Empty space indicates a

communication probability of zero. *P*-value were calculated by the one-sided permutation test.

Supplemental Figure 11: *Prdm1* maintains the IFN-γ production of liver type 1 ILCs in

tumor microenvironment.

Representative flow cytometric plot of the frequency of IFN-γ$^+$ liver cNK cells and ILC1s from

Prdm1$^{+/+}$ and *Prdm1*^{ΔNcr1} tumor-bearing mice at day 14 after inoculation with B16F10 melanoma

cells via intrasplenic injection (n=5). Data are presented as the mean±SEM and were analyzed by

2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on

graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Graphical abstract

Figure 1. *Prdm1* promotes type I ILCs homeostasis and terminal maturation.
(**A**) Correlation between the average expression of NK cell-associated genes(*NCR1*, *NCR3*, *PRF1*, *CD160*, *KLRB1*) and *PRDM1* in LIHC (Liver Hepatocellular Carcinoma; n=363) patients sourced from TCGA datasets. (**B**) Heatmap of the ordered, z-score normalized expression values for *PRDM1* and NK cell-associated genes in liver cancer patients. The top and bottom quartiles of all patient samples are indicated. (**C**) Prognostic value of the *NK-PRDM1* signature for overall survival of liver cancer patients comparing high and low quartiles. (**D**) Schematic representation of the Prdm1-conditional knockout mouse model. Targeted exon 6-8 of *Prdm1* (top) is flanked with *loxP* sites (middle). Ncr1-expressed Cre recombinase was used to generate the *Prdm1^ΔNcr1* allele (bottom). (**E**) Real-time RT-PCR quantification of *Prdm1* expression in NKp46+ cells to determine the presence of *Prdm1* (n=5). (**F**) Representative flow cytometric plots (left) and quantification (right) of the proportion and absolute number of CD3-NKp46+ cells among lymphocytes in liver, lung, and bone marrow (n=8). (**G**) Representative flow cytometric plots (left) of the CD11b and CD27 expression within CD3-NK1.1+NKp46+ cells in liver, lung, and bone marrow (n=7). Right panel showed the percentage of distinct stages of NK cells. Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; r, pearson correlation coefficient; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 2. *Prdm1* is required for type I ILCs to control tumor metastasis.
(**A**) Representative flow cytometric plots (left) of liver cNK cells (CD49a⁻CD49b⁺) and ILC1s (CD49a⁺CD49b⁻) from *Prdm1⁺/⁺* and *Prdm1^ΔNcr1^* mice. The two bar graghs (right) quantitate the percentages and absolute numbers of cells respectively (n=7). (**B** and **C**) Splenocytes from *B2m⁻/⁻* and *B2m⁺/⁺* were labeled with CFDASE and eF670 respectively. Labeled cells were 1:1 mixed and injected i.v. into *Prdm1⁺/⁺* and *Prdm1^ΔNcr1^* recipient mice to evaluate *in vivo* NK cell target-killing ability. Representative flow cytometric plots (left) of transferred cells recovered from recipient mice and percentage (right) of NK cell-specific rejection of donor cells in spleen (n=6) (**B**) and liver (n=7) (**C**) between *Prdm1⁺/⁺* and *Prdm1^ΔNcr1^* mice. (**D** and **F**) Image (left) and quantification (right) of tumor nodes on the livers (n=7) (**D**) and lungs (n=10) (**F**) of *Prdm1⁺/⁺* and *Prdm1^ΔNcr1^* mice at day 14 or 21 after inoculation with B16F10 melanoma cells. (**E** and **G**) Histopathological images of liver (**E**) and lung (**G**) tissues stained by hematoxylin-eosin to detect tumor metastasis. Red bar indicates 500 μm distance under the microscope. (**H**) Liver cells and splenocytes were costimulated in the presence or absence of IL-12 and IL-18 for 12 hours. GolgiStop was added 4 hours before intracellular staining of IFN-γ. The graphs showed percentage of IFN-γ⁺ splenic cNK cells, liver cNK cells and ILC1s from *Prdm1⁺/⁺* and *Prdm1^ΔNcr1^* mice (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 3. Bulk RNA-seq depicts *Prdm1*-mediated functions in peripheral cNK cells.
(**A**) Splenic cNK cells and liver CD45+ cells were sorted from *Prdm1+/+* and *Prdm1ΔNcr1* mice using flow cytometry, and prepared for bulk RNA-seq and single-cell RNA-seq analysis. (**B**) Volcano plot of the bulk RNA-seq differentially expressed genes (log2(fold change)>0.5; *P*<0.05) in splenic cNK cells between *Prdm1+/+* and *Prdm1ΔNcr1* mice. Upregulated and downregulated genes in *Prdm1ΔNcr1* cells were highlighted in red and blue. (**C**) Enriched Gene Ontology (GO) terms of DEGs in *Prdm1ΔNcr1* cells compared *Prdm1+/+* cells. The Enrichment gene set in upregulated (red) and downregulated (blue) genes were indicated in different colours. Bar length represents statistical significance. (**D** and **E**) Gene Set Enrichment Analysis (GSEA) showing the enrichment of negative regulation of IL-6 production (**D**) and NF-kappa B signaling pathway (**E**) of DEGs in *Prdm1ΔNcr1* cells compared *Prdm1+/+* cells. NES, normalized enrichment score. (**F**) Heatmap of selected genes from DEGs. Shown is z-score transformed expression of DEGs. (**G** and **I**) Representative flow cytometric plots (left) and cumulative data (right) showing the relative mean fluorescence intensities (MFIs) of Granzyme B (**G**) and Perforin (**I**) in liver cNK cells and ILC1s from *Prdm1+/+* and *Prdm1ΔNcr1* mice (n=5). (**H** and **J**) Relative MFIs of Granzyme B (**G**) and Perforin (**I**) in splenic NK cells at different stages of maturation was analyzed by flow cytometry (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.

Figure 4. scRNA-seq reveals unique properties of two clusters from liver type I ILCs following *Prdm1* knockout.
(**A**) Uniform manifold approximation and projection (UMAP) visualization of liver CD45+ cells from *Prdm1+/+* and *Prdm1ΔNcr1* mice. Twelve clusters were defined and indicated by distinct colours. Each dot represents a single cell. (**B**) UMAP visualization of liver cNK and ILC1 clusters. Cells were colored by origins (*Prdm1+/+*-blue; *Prdm1ΔNcr1*-red). (**C**) Percentages of cNK cells and ILC1s in type I ILCs (left), and their distribution in each cluster (right). (**D-I**) UMAP visualization of three different liver cNK (**D**) and ILC1 (**G**) clusters from two mouse strains (**E** and **H**). Proportions of cNK cells (**F**) and ILC1s (**I**) among total cells (cNK cells or ILC1s) (left) and within clusters (right) were calculated respectively. (**J**) Violin plots showing the normalized expression of select genes in different cNK clusters. (**K**) Enriched GO term of marker genes in three cNK clusters. Dot size represents enriched gene number, and color intensity represents significance.

Figure 5. *Junb*hi signature is associated with the activation of multiple signaling pathways.
(**A**) Ridge plots showing the normalized expression of *Gzmb, Prf1*, and *Junb* in cNK and ILC1 clusters between *Prdm1+/+* and *Prdm1ΔNcr1* cells.
(**B**) Violin plots showing the normalized expression of select genes in different ILC1 clusters. (**C**) Violin plot showing the *Junb*hi signature score for cNK cell and ILC1 clusters, calculated using the signature genes of *Junb*hi cNK cluster. (**D**) Enriched GO term of marker genes in three ILC1 clusters. Dot size represents enriched gene number, and color intensity represents significance. (**E-J**) GSEA plots (left) depicting the enrichment of NF-kappa B (**E**) , TNF (**F**) , IL-17 (**G**) , and MAPK (**H**) signaling pathway in *Junb*hi cNK cluster compared with clusters of *Prf1*hi and *Cxcr3*hi cNK cells, and the enrichment of IL-17 signaling pathway (**I**) and T cell differentiation (**J**) in *Il7r*hi ILC1 cluster compared with clusters of *Klra*hi and *Gzma*hi ILC1s. Right panel showed dynamic relative expression of the given gene sets from cluster1 to cluster3 between *Prdm1+/+* and *Prdm1ΔNcr1*. Dots represent the average expression of given gene set in each cell, which was calculated through the sum of normalized expression of each individual gene within the designated gene set in every single cell. NES, normalized enrichment score.

Figure 6. *Prdm1* facilitates the intercellular communication between liver type I ILCs and macrophages.
(**A**) UMAP visualization of macrophages (Mac) cluster. Four cell subsets were defined. (**B** and **C**) Circle plots (**B**) and summary data (**C**) illustrating the significant enriched ligand–receptor pairs among cluster of liver cNK cells, ILC1s, and macrophages from *Prdm1^+/+^* (left) and *Prdm1^ΔNcr1^* (right) cells. The thickness of the line indicates the number of enrich pairs, and the arrow reflects the direction of the interaction. (**D**) Heatmap of overall signaling pattern recognized from ligand-receptor pairs, which contained the sum of signaling from the sender and target cells. (**E**) Bar graphs showing the information flow in selected active signaling patterns between *Prdm1^+/+^* and *Prdm1^ΔNcr1^* cells. Relative information flow was calculated as the sum of the communication probability in given signaling patterns. (**F**) Chord plot of the CXCL signaling interaction network among cluster of liver cNK cells, ILC1s, and macrophages in *Prdm1^+/+^* cells.

Figure 7. *Prdm1* safeguards type I ILCs from exhaustion-like phenotypes in the tumor microenvironment.
(A-C) The Mitochondrial mass (MitoTracker Green staining; n=8) (**A**), Mitochondrial ROS (MitoSOX staining; n=5) (**B**), and Mitochondrial membrane potential (TMRM staining; n=5) (**C**) of splenic cNK cells, liver cNK cells and ILC1s were analyzed by flow cytometry. Representative flow cytometric plots (left) and cumulative data (right) showing the relative mean fluorescence intensities (MFIs) of each group. (**D**) Percentages of IFN-γ^+ liver cNK cells and ILC1s from *Prdm1*[+/+] and *Prdm1*[ΔNcr1] tumor-bearing mice at day 14 after inoculation with B16F10 melanoma cells via intrasplenic injection (n=5). Data are presented as the mean±SEM and were analyzed by 2-tailed, paired t-test. Differences were evaluated between littermates. Each circle and square on graphs represents an individual mouse; *P*, *P*-value; *, *P*<0.05; **, *P*<0.01, ns, not significant.