**Sequencing, de novo assembly of *Ludwigia* plastomes, and comparative analysis within**

**the Onagraceae family**

Barloy-Hubler F.[3], Le Gac A.-L.[1], Boury C.[2], Guichoux E.[2], and Barloy D.[1*]

1-DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, IFREMER, INRAE,

Rennes, France.

2- Université de Bordeaux, INRAE, BIOGECO, Cestas, France

3- Université de Rennes 1, CNRS, UMR 6553 ECOBIO, Rennes, France.

* Corresponding author: dominique.barloy@agrocampus-ouest.fr

**Abstract**

The Onagraceae family, which belongs to the order Myrtales, consists of approximately 657 species and 17 genera. This family includes the genus *Ludwigia* L., which is comprised of 82 species. In this study, we focused on the two aquatic invasive species *Ludwigia grandiflora* subsp. *hexapetala* (*Lgh*) and *Ludwigia peploides* subsp *montevidensis* (*Lpm*) largely distributed in aquatic environments in North America and in Europe. Both species have been found to degrade major watersheds leading ecological and economical damages. Genomic resources for Onagraceae are limited, with only *Ludwigia octovalvis* (Lo) plastid genome available for the genus *Ludwigia* L. at the time of our study. This scarcity constrains phylogenetic, population genetics, and genomic studies. To brush up genomic ressources, new complete plastid genomes of *Ludwigia grandiflora* subps. *hexapetala* (*Lgh*) and *Ludwigia peploides* subsp. *montevidensis* (*Lpm*) were generated using a combination of MiSeq (Illumina) and GridION (Oxford Nanopore) sequencing technologies. These plastomes were then compared to the published *Ludwigia octovalvis* (*Lo*) plastid genome, which was re-annotated by the authors. We initially sequenced and assembled the chloroplast (cp) genomes of *Lpm* and *Lgh* using a hybrid strategy combining short and long reads sequences. We observed the existence of two *Lgh* haplotypes and two potential *Lpm* haplotypes. *Lgh*, *Lpm,* and *Lo* plastomes were similar in terms of genome size (around 159 Kb), gene number, structure, and inverted repeat (IR) boundaries, comparable to other species in the Myrtales order. A total of 45 to 65 SSRs (simple sequence repeats), were detected, depending on the species, with the majority consisting solely of A and T, which is common among angiosperms. Four chloroplast genes (*matK*, *accD*, *ycf2* and *ccsA)* were found under positive selection pressure, which is commonly associated with plant development, and especially in aquatic plants such as *Lgh,* and *Lpm*. Our hybrid sequencing approach revealed the presence of two *Lgh* plastome haplotypes which will help to advance phylogenetic and evolutionary studies, not only specifically for *Ludwigia*, but also the Onagraceae family and Myrtales order. To enhance the robustness of our findings, a larger dataset of chloroplast genomes would be beneficial.

**Keywords**

Water primrose, *Ludwigia* sp., Onagraceae, chloroplast genome, long and short reads, hybrid assembly, haplotype

**Introduction**

The Onagraceae family belongs to the order Myrtales which includes approximately 657 species of herbs, shrubs, and trees across 17 genera grouped into two subfamilies: subfam. Ludwigioideae W. L. Wagner and Hoch, which only has one genus (*Ludwigia* L.), and subfam. Onagroideae which contains six tribes and 21 genera [1]. *Ludwigia* L. is composed of 83 species[2][3] . The current classification for *Ludwigia* L., which are composed of several hybrid and/or polyploid species, lists 23 sections. A recent molecular analysis is clarified and supported several major relationships in the genus but has challenged the complex sectional classification of *Ludwigia* L.[4].

The diploid species *Ludwigia peploides* (Kunth) Raven subsp. *montevidensis* (Spreng.) [5] (named here *Lpm*) (2n=16), and the decaploid species, *Ludwigia grandiflora* (Michx) Greuter & Burdet subsp. *hexapetala* (Hook. & Arn) Nesom & Kartesz (named here *Lgh*) (2n=80), reproduce essentially by clonal propagation, which suggests that there is a low genetic diversity within the species [6]. *Lgh* and *Lpm* are native to South America and are considered as one of the most aggressive aquatic invasive plants [7]. Largely distributed in aquatic environments in North America and in Europe [8], both species have been found to degrade major watersheds as well as aquatic and riparian ecosystems  [9] leading ecological and economical damages. In France, both species occupied aquatic habitats, such as static or slow-flowing waters, riversides, and have recently been observed in wet meadows [10]. The transition from an aquatic to a terrestrial habitat has led to the emergence of two *Lgh* morphotypes [11]. The appearance of metabolic and morphological adaptations could explain the ability to acclimatize to terrestrial conditions, and this phenotypic plasticity involves various genomic and epigenetic modifications [12].

Adequate genomic resources are necessary in order to be identify the genes and metabolic pathways involved in the adaptation process leading to plant invasion [13] with genomic information making it possible to predict and control invasiveness [14]. However, even though the number of terrestrial plant genomes has increased considerably over the last 20 years, only a small fraction (~ 0.16%) have been sequenced, with some clades being significantly more represented than others [15]. Thus, for the Onagraceae family (which includes *Ludwigia* sp.), only a handful of chloroplast sequences (plastomes) are available, and the complete genome has not yet been sequenced. If *Lpm* is a diploid species (2n=2x=16) with a relatively small genome size (262 Mb), *Lgh* is a decaploid species (2n=10x=80) with a large size genome of 1419 Mb [16]. Obtaining a reference genome for these two non-model species without having a genome close to the *Ludwigia* species is challenging and development of plastome and/or

mitogenome will be a first step to generate genomic resource. As of April 2023, there are 10,712 reference plastomes listed on GenBank (Release 255: April 15 2023), with the vast majority (10,392 genomes) belonging to Viridiplantae (green plants). However, in release 255, the number of plastomes available for the Onagraceae family is limited, with only 36 plastomes currently listed. Among these, 15 plastomes are from the tribe Epilobieae, with 11 in the Epilobium genus and 4 in the Chamaenerion genus. Additionally, there are 23 plastomes from the tribe Onagreae, with 17 in the Oenothera genus, 5 in the Circaea genus, and only one in the *Ludwigia* genus. The *Ludwigia octovalvis* chloroplast genome was released in 2016 as a unique haplotype of approximately 159 kb [17]. *L. octovalvis* belongs to sect. *Macrocarpon* (Micheli) H.Hara while *Lpm* and *Lgh* belong to *Jussieae* section [18][19]. Generally, the inheritance of chloroplast genomes is considered to be maternal in angiosperms. However, biparentally inherited chloroplast genomes could potentially exist in approximately 20% of angiosperm species [20][21]. Both maternal and biparental inheritance are described in the Onagraceae family. In tribe Onagreae, *Oenothera* subsect. *Oenothera* are known to have biparental plastid inheritance [22][23]. In tribe Epilobieae, biparental plastid inheritance was also reported in *Epilobium* L. with mainly maternal transmission, and very low proportions of paternally transmitted chloroplasts [24].

The chloroplast is the symbolic organelle of plants and plays a fundamental role in photosynthesis. Chloroplasts evolved from cyanobacteria through endosymbiosis and thereby inherited components of photosynthesis reactions (photosystems, electron transfer and ATP synthase) and gene expression systems (transcription and translation)[25]. In general, chloroplast genomes (plastomes) are highly conserved in size, structure, and genetic content. They are rather small (120-170 kb,[26]), with a quadripartite structure comprising two long identical inverted repeats (IR, 10–30 kb) separated by large and a small single copy regions (LSC and SSC, respectively). They are also rich in genes, with around 100 unique genes encoding key proteins involved in photosynthesis, and a comprehensive set of ribosomal RNAs (rRNAs) and transfer RNAs (tRNAs)[27]. Plastomes are generally circular but linear shapes also exist [28]. Chloroplast DNA usually represents 5-20% of total DNA extracted from young leaves and therefore low-coverage whole genome sequencing can generate enough data to assemble an entire chloroplast genome [29].

If we refer to their GenBank records, more than 95% of these plastomes were sequenced by so-called short read techniques (mostly Illumina). However, in most seed plants, the plastid genome exhibits two large inverted repeat regions (60 to 335 kb,[29]), which are longer than the short read lengths (< 300 bp). This leads to incomplete or approximate assemblies [30].

Recent long-read sequencing (> 1000 bp) provides compelling evidence that terrestrial plant plastomes exhibit two structural haplotypes. These haplotypes are present in equal proportions and differ in their inverted repeat (IR) orientation [31]. This shows the importance of using the so-called third generation sequence (TGS, PacBio or Nanopore) to correctly assemble the IRs of chloroplasts and to identify any different structural haplotypes. The current problem with PacBio or Nanopore long read sequencing is the higher error rate compared to short read technology [32][33][34]. Thus, a hybrid strategy which combines long reads (to access the genomic structure) and short reads (to correct sequencing errors) could be effective [30][35].

Here, we report the newly sequenced complete plastid genomes of *Ludwigia grandiflora* subps. *hexapetala* (*Lgh*) and *Ludwigia peploides* subsp *montevidensis* (*Lpm*), using a combination of different sequencing technologies, as well as a re-annotated comparative genomic analysis of the published *Ludwigia octovalvis* (*Lo*) plastid. The main objectives of this study are (1) to assemble and annotate the plastomes of two new species of *Ludwigia* sp., (2) to reveal the divergent sequence hotspots of the plastomes in this genus and in the Onagraceae (3) to identify the genes under positive selection.

To achieve this, we utilized long read sequencing data from Oxford Nanopore and short read sequencing data from Illumina to assemble the *Lgh* plastomes and compared these assemblies with those obtained solely from long reads of *Lpm*. We also compared both plastomes to the published plastome of *Lo*. Our findings demonstrated the value of *de novo* assembly in reducing assembly errors and enabling accurate reconstruction of full heteroplasmy. We also evaluated the performance of a variety of software for sequence assembly and correction in order to define a workflow that will be used in the future to assemble *Ludwigia* sp. mitochlondrial and nuclear genomes. Finally, the three new *Ludwigia* plastomes generated by our study make it possible to extend the phylogenetic study of the Onagraceae family and to compare it with previously published analyses [4][36][37].

**Material and Methods**

**Plant sampling and experimental design**

The original plant materials were collected in June of 2018 near to Nantes (France) and formal identified by D. Barloy. *L. grandiflora* subsp. *hexapetala (Lgh)* plants were taken from the Mazerolles swamps (N47 23.260, W1 28.206), and *L. peploides* subsp. *montevidensis* (*Lpm*) plants from La Musse (N 47.240926, W -1.788688)*)*. Plants were cultivated in a growth chamber in a mixture of $^1/_3$ soil, $^1/_3$ sand, $^1/_3$ loam with flush water level, at 22°C and a 16 h/8

h (light/dark) cycle. A single stem of 10 cm for each species was used for vegetative propagation in order to avoid potential genetic diversity. *De novo* shoots, taken three centimeters from the apex, were sampled for each species. Samples for gDNA extraction were pooled and immediately snap-frozen in liquid nitrogen, then lyophilized over 48 h using a Cosmos 20K freeze-dryer (Cryotec, Saint-Gély-du-Fesc, France) and stored at room temperature. All the plants were destroyed after being used as required by French authorities for invasive plants (article 3, prefectorial decree n°2018/SEE/2423).

Due to high polysaccharide content and polyphenols in *Lpm* and *Lgh* tissues and as no standard kit provided good DNA quality for sequencing, genomic DNA extraction was carried out using a modified version of the protocol proposed by Panova et al in 2016, with three purification steps [38].

40 mg of lyophilized buds were ground at 30 Hz for 60 s (Retsch MM200 mixer mill, FISHER). The ground tissues were lysed with 1 ml CF lysis buffer (MACHEREY-NAGEL) supplemented with 20 μl RNase and incubated for 1 h at 65°C under agitation. 20 μl proteinase K was then added before another incubation for 1 h at 65°C under agitation. To avoid breaking the DNA during pipetting, the extracted DNA was recovered using a Phase-lock gel tube as described in Belser [39]. The extracts were transferred to 2 ml tubes containing phase-lock gel, and an equal volume of PCIA (Phenol, Chloroform, Isoamyl Alcohol; 25:24:1) was added. After shaking for 5 min, tubes were centrifuged at 11000 g for 20 min. The aqueous phase was transferred into a new tube containing phase-lock gel and extraction with PCIA was repeated. DNA was then precipitated after addition of an equal volume of binding buffer C4 (MACHEREY-NAGEL) and 99% ethanol overnight at 4°C or 1 h in ice then centrifuged at 800 rpm for 10 min. After removal of the supernatant, 1 ml of CQW buffer was added then the pellet of DNA was re-suspended. Next, DNA purification was carried out by adding a 2 ml mixture of wash buffer PW2 (MACHEREY-NAGEL), wash buffer B5 (MACHEREY-NAGEL), and ethanol at 99% in equal volumes, followed by centrifugation at 800 rpm for 10 min. This DNA purification step was carried out twice. Finally, the DNA pellet was dried in the oven at 70°C for 30 min then re-suspended in 100 μl elution buffer BE (MACHEREY-NAGEL) (5 mM Tris solution, pH 8.5) after 10 min incubation at 65°C under agitation.

A second purification step was performed using a PCR product extraction from gel agarose kit from Macherey-Nagel (MN) NucleoSpin® Gel and PCR Clean-up kit and restarting the above protocol from the step with the addition of CQW buffer then PW2 buffer.

The third purification step consisted of DNA purification using a Macherey-Nagel (MN)
NucleoMag kit for clean-up and size selection. Finally, the DNA was resuspended after a 5 min
incubation at 65°C in 5 mM TRIS at pH 8.5.

The quantity and quality of the gDNA was verified using a NanoDrop spectrometer,
electrophoresis on agarose gel and ethidium bromide staining under UV light and Fragment
Analyzer (Agilent Technologies) of the University of Rennes1.

**Library preparation and sequencing**

MiSeq (Illumina) and GridION (Oxford Nanopore Technologies, referred to here as
ONT) sequencing were performed at the PGTB (doi:10.15454/1.5572396583599417E12). *Lgh*
and *Lpm* genomic DNA were re-purified using homemade SPRI beads (1.8X ratio). *Lgh* has a
large genome size of 1419 Mb, 5-fold larger than *Lpm* genome 262 Mb [16]. SR (Illumina, one
run) and LR (Oxford Nanopore, three runs) sequencing were therefore carried out for *Lgh* and
only LR sequencing for *Lpm* (one run). For Illumina sequencing, 200 ng of *Lgh* DNA was used
according to the QIAseq FX DNA Library Kit protocol (Qiagen). The final library was checked
on TapeStation D5000 screentape (Agilent Technologies) and quantified using a QIAseq
Library Quant Assay Kit (Qiagen). The pool was sequenced on an Illumina MiSeq using V3
chemistry and 600 cycles (2x300bp). For ONT sequencing, around 8 μg of *Lgh* and *Lpm* DNA
were size selected using a Circulomics SRE kit (according to the manufacturer's instructions)
before library preparation using a SQK-LSK109 ligation sequencing kit following ONT
recommendations. Basecalling in High Accuracy - Guppy version: 4.0.11 (MinKNOW
GridION release 20.06.9) was performed for the 48 h of sequencing. Long reads (LR) and short
reads (SR) were available for *Lgh* and only LR for *Lpm*.

**Chloroplast assemblies**

Quality controls and preprocessing of sequences were conducted using Guppy v4.0.14 for
long reads (via Oxford Nanopore Technology Client access) and fastp v0.20.0 [40]  for short
reads, using Q15, since increasing the Phred quality to 20 or higher has no effect on the number
of sequences retained (66%). A preliminary draft assembly was performed using *Lgh* short-
reads (SR, 2*23,067,490 reads) with GetOrganelle v1.7.0 [41] and NOVOPlasty v4.2.1[42],
and chloroplastic short and long reads were extracted by mapping against this draft genome.
Chloroplastic short reads were then *de novo* assemble using Velvet (version 1.2.10) [43],
ABySS (version 2.1.5 [44][45]), MEGAHIT (1.1.2,[46]), and SPAdes (version 3.15.4,[47]),
without and with prior error correction. The best k-mer parameters were tested using kmergenie
[48] and k=99 was found to be optimal. For ONT reads, *Lgh* (550,516 reads) and *Lpm* (68,907
reads) reads were self-corrected using CANU 1.8 [49] or SR-corrected using Ratatosk [50] and

*de novo* assembly using CANU [49] and FLYE 2.8.2 [51]  run with the option --meta and –

plasmids. For all these assemblers, unless otherwise specified, we used the default parameters.

**Post plastome assembly validation**

As we used many assemblers and different strategies, we produced multiple contigs that

needed to be analyzed and filtered in order to retain only the most robust plastomes. For that,

all assemblies were evaluated using the QUality ASsessment Tool (QUAST) for quality

assessment [52] and visualized using BANDAGE [53], both using default parameters.

BANDAGE compatible graphs (.gfa format) were created with the megahit_toolkit for

MEGAHIT [46] and with gfatools for ABySS [45]. Overlaps between fragments were manually

checked and ambiguous "IUPAC or N" nucleotides were also biocured with Illumina reads

when available.

**Chloroplast genome annotation**

Plastomes were annotated via the GeSeq [54] using ARAGORN and tRNAscan_SE to

predict tRNAs and rRNAs and tRNAscan_SE to predict tRNAs and rRNAs and via Chloe

prediction site [55]. The previously reported *Lo* chloroplast genome was also similarly re-

annotated to facilitate genomic comparisons. Gene boundaries, alternative splice isoforms,

pseudogenes and gene names and functions were manually checked and biocurated using

Geneious (v.10). Finally, plastomes were represented using OrganellarGenomeDRAW

(OGDRAW)[56]. These genomes were submitted to GenBank at the National Center of

Biotechnology Information (NCBI) with specific accession numbers (for *Lgh* haplotype 1,

(LGH1) OR166254 and *Lgh* haplotype 2, (LGH2) OR166255; for *Lpm* haplotype, (LPM)

OR166256) using annotation tables generated through GB2sequin [57].

**SSRs and Repeat Sequences Analysis**

Simple Sequence Repeats (SSRs) were analyzed through the MISA web server [58], with

parameters set to 10, 5, 4, 3, 3, and 3 for mono-, di-, tri-, tetra-, penta-, and hexa-nucleotides,

respectively. Direct, reverse and palindromic repeats were identified using RepEx [59].

Parameters used were: for inverted repeats (min size 15 nt, spacer = local, class = exact); for

palindromes (min size 20 nt); for direct repeats (minimum size 30 nt, minimum repeat similarity

97%). Tandem repeats were identified using Tandem Repeats Finder[60], with parameters set

to two for the alignment parameter match and seven for mismatches and indels. The IRa region

was removed for all these analyses to avoid over representation of the repeats.

**Comparative chloroplast genomic analyses**

*Lgh* and *Lpm* plastomes were compared with the reannotated and biocurated *Lo* plastome

using mVISTA program [61], with the LAGAN alignment algorithm [62] and a cut-off of 70%

identity.

Nucleotide diversity (Pi) was analyzed using the software DnaSP v.6.12.01 [63] [64]with

step size set to 200 bp and window length to 300 bp. IRscope [65] was used for the analyses of

inverted repeat (IR) region contraction and expansion at the junctions of chloroplast genomes.

To assess the impact of environmental pressures on the evolution of these three *Ludwigia*

species, we calculated the nonsynonymous (Ka) and synonymous (Ks) substitutions and their

ratios ($\omega$ = Ks/Ks) using TBtools [66] to measure the selective pressure. Genes with $\omega < 1$, $\omega =$

1, and $1 < \omega$ were considered to be under purifying selection (negative selection), neutral

selection, and positive selection, respectively.

**Phylogenetic analysis of *Ludwigia* based on MatK sequences**

We performed a phylogenetic analysis on the *Ludwigia* genus using the MatK, only protein

coding barcode available for a large number of *Ludwigia* species. All MatK amino acid

sequences were aligned with the FFT-NS-2 (Fast Fourier Transform-based Narrow Search)

algorithm and BLOSUM62 scoring matrix using MAFFT 7 [67]. The phylogenetic tree analysis

was conducted using the rapid hill-climbing algorithm (command line: -f d) in RAxML 8.2.11

[68], with GAMMA JTT (Jones-Taylor-Thornton) protein model. Node support was assessed

through fast bootstrapping (-f a) with 1,000 non-parametric bootstrap pseudo-replicates.

*Circaea* MatK were selected as outgroup, and all accession numbers are indicated on the

phylogenetic tree labels.

**Graphic representation**

Statistical analyses were performed using R software in RStudio integrated development

environment (R Core Team, 2015, RStudio: Integrated Development for R. RStudio, Inc.,

Boston, MA, http://www.rstudio.com/). Figures were realized using ggplot2, ggpubr, tidyverse,

dplyr, gridExtra, reshape2, and viridis packages. SNPs were represented using trackViewer [69]

and genes represented using gggenes packages.

**Results**

**Plastome short read assembly**

The chloroplastic fraction of *Lgh* short reads (SR) was extracted by mapping against the two

draft haplotypes generated by GetOrganelle, which differ only by a "flip-flop" of the SSC region

(Figure 1). Since the assembly by NOVOplasty did not provide any additional information

compared to GetOrganelle, it was not retained. This subset (1,360,507 reads) was assembled

using ABySS, Velvet, MEGAHIT and SPAdes in order to identify the best assembler for this
plant model. As shown in Figure 2, both the number and size of contigs depend greatly on the
algorithms used and the correction step. The effect of prior read correction is notable for
MEGAHIT and Velvet, especially concerning the increase in the size of the large alignment
(Add. Figure 1A), loss of misassemblies, and reduction of the number of mismatches (Add.
Figure 1B). Investigating results via BANDAGE (Add. Figure 2), we observed that ABySS and
SPAdes suggest the tripartite structure with the long single-copy (LSC) region as the larger
circle in the graph (blue), joined to the small single-copy region (green) by one copy of the
inverted repeats (IRs, red), both IRs being collapsed in a segment of approximately twice the
coverage. For Velvet and MEGAHIT, graphs confirm the significant fragmentation of the
assemblies, which is improved by prior correction of the reads.

In conclusion, none of the short-read assemblers tested in our study produced a complete
plastome. The best result was achieved by SPAdes using corrected short reads (mean coverage
1900 X) to assemble a plastome consisting of three contigs: 90,272 bp (corresponding to LSC),
19,788 bp (corresponding to SSC), and 24,762 bp (corresponding to one of the two copies of
the IR).

**Plastome long read assembly**

Chloroplast fractions of *Lgh* long reads (28,882 reads) were assembled using CANU or
FLYE. With raw data, CANU generates a unique contig (NGA50 112648) corresponding to
haplotype 2, whereas FLYE makes two contigs (NGA50 133687) that reconstruct haplotype 1.
Self-corrected LR leads to fragmentation into two (CANU) or three (FLYE) contigs which both
reconstruct haplotype 1, with an large gap corresponding to one of the IR copies for CANU.
Finally, SR-correction by RATATOSK allows CANU to assemble two redundant contigs
reproducing haplotype 2 while FLYE makes two contigs corresponding to haplotype 1 (Add.
Figure 3A). In conclusion, the two *Lgh* haplotypes were reconstructed (average coverage 700X)
and the most complete and accurate hybrid assemblies (99.94% accuracy, Additional Figure
3B) were submitted to GenBank.

Unfortunately, due to the absence of short read data, we could only perform self-corrected
long read assembly for *Lpm* using CANU. We also compared CANU and FLYE assembler
efficiency, and found that assembly using CANU produces 13 contigs whereas FLYE produces
12 contigs. In both cases, only three contigs are required to reconstitute a complete cpDNA
assembly (no gap, no N), with an SSC region oriented like those of the *Lgh* haplotype 2 and the
*Lo* plastome. Although it is more than likely that these two SSC region orientations also exist
for *Lpm*, the low number of nanopore sequences generated (68907 reads) and absence of

Illumina short reads prevented us from demonstrating the existence of both haplotypes. As a

result, only the "haplotype 2" generated sequence was deposited to Genbank.

**Annotation and comparison of *Ludwigia* plastomes**

**1. General Variations**

Plastomes of the three species of *Ludwigia* sp., *Lgh*, *Lpm* and *Lo,* are circular double-

stranded DNA molecules (Figure 3) which are all (as shown in Table 1) approximately the same

size: *Lo* is 159,396 bp long, making it the smallest, while *Lgh* is the largest with 159,584 bp,

and *Lpm* is intermediate at 159,537 bp. The overall GC content is almost the same for the three

species (37.4% for *Lo*, 37.3 % for *Lgh* and *Lpm*) and the GC contents of the IR regions are

higher than those of the LSC and SSC regions (approximately 43.5 % compared to 35% and

*ca.*32% respectively). Between the three species, the lengths of the total chloroplasts, LSC,

SSC, and IR are broadly similar (approximately 90.2 kb for LSC, 19.8 kb for SSC and 24.8 kb

for IB, see details Table 1) and the three plastomes are perfectly syntenic if we orient the SSC

fragments the same way.

All three *Ludwigia* sp. plastomes contain the same number of functional genes (134 in total)

encoding 85 proteins (embracing 7 duplicated in the IR region: *ndh*B, *rpl2*, *rpl23*, *rps7*, *rps12*,

*ycf2*, *ycf15*), 37 tRNAs (including *trn*K-UUU which contains *mat*K), and 8 rRNAs (16S, 23S,

5S, and 4.5S as duplicated sets in the IR). Among these genes, 18 contain introns, of which six

are tRNAs (Table 2). Only the *rps12* gene is a trans-spliced gene. A total of 46 genes are

involved in photosynthesis, and 71 genes related to transcription and translation, including a

bacterial-like RNA polymerase and 70S ribosome, as well as a full set of transfer RNAs

(tRNAs) and ribosomal RNAs (rRNAs). Six other protein-coding genes are involved in

essential functions, such as *acc*D*,* which encodes the β-carboxyl transferase subunit of acetyl-

CoA carboxylase, an important enzyme for fatty acid synthesis; *mat*K encodes for maturase K,

which is involved in the splicing of group II introns; *cem*A, a protein located in the membrane

envelope of the chloroplast is involved in the extrusion of protons and thereby indirectly allows

the absorption of inorganic CO2 in the plastids; *clp*P1 which is involved in proteolysis, and;

*ycf1*, *ycf2*, two ATPases members of the TIC translocon. Finally, a highly pseudogenized *ycf15*

locus was annotated in the IR even though premature stop codons indicate loss of functionality.

**2. Segments Contractions/Expansion**

The junctions between the different chloroplast segments were compared between three

*Ludwigia* sp. (*Lpm*, *Lgh* and *Lo),* and we found that the overall resemblance of *Ludwigia* sp.

plastomes was confirmed at all junctions (Figure 4A). In all three genomes, *rpl22, rps19, and*

*rpl2* were located around the LSC/IRb border, and *rpl2*, *trn*H, and *psb*A were located at the

IRa/LSC edge. The JSB (junction between IRb and SSC) is either located in the *ndh*F gene or the *ycf1* gene depending on the orientation of the SSC region (Figure 4B). The *ycf1* gene was initially annotated as a 1139 nt pseudogene that we biocurate as a larger gene (5302 nt) with a frameshift due to a base deletion, compared to *Lg* and *Lo* which both carry a complete *ycf1* gene.

If we compare *Ludwigia* sp. chloroplastic LSC/SCC/IR junctions (via IRscope) with representative Onagraceae plastomes of *Chamaenerion sp. conspersum* (MZ353638) and *sp. angustifolium* (NC_052848)*, Circaea sp. cordata* (NC_060876) and *sp. alpina* (NC_061010), *Epilobium amurense* (NC_061015) and *Oenothera villosa* subsp. *strigosa* (NC_061365) and *Oenothera lindheimeri* (MW538951) (Figure 5), we can observe that the gene positions at the JLB (junction of LSC/IRb) and JLA (junction of IRa/LSC) boundary regions are well-preserved throughout the entire family, whereas those at the JSB and JSA regions differ. Concerning JSB (junction of IRb/SSC), in the five Onagraceae genera studied, *ndh*F is duplicated, with the exception of *Circaea* sp. and *Ludwigia* sp. For *Oenothera villosa*, the first copy of *ndh*F, which is located in the IRb, overlaps the JSB border, whereas for *Oenothera lindheimeri*, *Epibolium amurense* and Chamaenerion sp., *ndh*F is only located in inverted repeats. Only *Circaea* sp. and *Ludwigia* sp. have a unique copy of this locus, and it is found in the SSC segment (Figure 5). At the JSA border (junction of SSC/Ira), in *Circaea* sp., the *ycf1* gene crosses the IRa/SSC boundary and extends into the IRa region.

When comparing the respective sizes of chloroplast fragments (IR/SSC/LSC) in Onagraceae, it can be observed that *Ludwigia* species exhibit expansions in the SSC and LSC regions which are not compensated by significant contractions in the IR regions. This is likely due to the relocation of the *ndh*F in the SSC region and *rps19* in the LSC region. Additionally, there may be significant size variations in the intergenic region between *trnI* and *ycf2*, as well as the intergenic segment containing the *ycf15* pseudogene (Add. Figure 4).

### 3. Repeats and SSRs analysis

In this study, we analyzed the nature and distribution of single sequence repeats (SSR), as their polymorphism is an interesting indicator in phylogenetic analyses. A total of 65 (*Lgh*), 48 (*Lpm*) and 45 (*Lo*) SSRs were detected, the majority being single nucleotide repeats (38–21), followed by tetranucleotides (12–10) and then di-, tri- and penta-nucleotides (Add. Figure 5A). Mononucleotide SSRs are exclusively composed of A and T, indicating a bias towards the use of the A/T bases, which is confirmed for all SSRs (Add. Figure 5B). In addition, the SSRs are mainly distributed in the LSC region for the three species, which is probably biased by the fact that LSC is the longest segment of the plastome (Add. Figure 5C). The analysis of SRR

locations revealed that most were distributed in non-coding regions (intergenic regions and

introns, Add. Figure 5D).

The chloroplast genomes of the three *Ludwigia* species were also screened for long repeat

sequences. They were counted in a non-redundant way (if smaller repetitions were included in

large repeats, only the large ones were considered). Four types of repeats (tandem, palindromic

inverted and direct) were surveyed in the three *Ludwigia* sp. plastomes. No inverted repeats

were detected with the criteria used.

For the three other types of repeats, here are their distributions:

***Tandem repeats* (Table 3A)**: Perfect tandem repeats (TRs) with more than 15 bp were

examined. Twenty-two *loci* were identified in the three *Ludwigia* sp. plastomes (*Lgh*, *Lpm*, *Lo*),

heterogeneously distributed as shown in Table 3A: 13 loci (plus one imperfect) in *Lo*, nine loci

(plus one imperfect) in *Lgh* and seven loci (plus two imperfect) in *Lpm*. It can therefore be seen

that the TR distributions (occurrence and location) are specific to each plastome, since only

four pairs are common to the three species. Thus, nine TRs are unique to *Lo*, three to *Lpm* and

three to *Lgh.* Two pairs are common to *Lgh* and *Lpm* and one is common to *Lo* and *Lgh.* TRs

are mainly intergenic or intronic but are detected in two genes (*acc*D and *ycf1)*. These genes

have accelerated substitution rates, although this does not generate a large difference in their

lengths. This point will be developed later in this article.

***Direct repeats* (Table 3B)**: There are few direct (non-tandem) repeats (DRs) in the

chloroplast genomes of *Ludwigia* sp. A single direct repeat of 41 nt is common to the three

species, at 2 kb intervals, in *psaB* and *psaA* genes. This DR corresponds to an amino acid repeat

[WLTDIAHHHLAIA] which corresponds to a region predicted as transmembrane. We then

observe three direct repeats conserved in *Lpm* and *Lgh* in *ycf1*, *acc*D and *clp*P1 respectively,

two unique DRs in *Lo* (in the *acc*D gene and *rps12-clp*P1 intergene) and one in *Lgh* (in the

*clp*P1 intron 1 and *clp*P1 intron 2).

***Palindromes* (Table 3C)**: Palindromic repeats make up the majority of long repetitions,

with the numbers of perfect repeats varying from 19, 24 and 26 in *Lo*, *Lgh* and *Lpm*,

respectively, and the number of quasi-palindromes (1 mutation) varying between 8, 3 and 6.

They are mainly found in the intronic and intergenic regions, with the exception of six genic

locations in *psb*D, *ndh*K, *ccs*A and *rpl22,* and two palindromic sequences in *ycf2.* These gene

palindromic repeats do not seem to cause genetic polymorphism in *Ludwigia* and can be

considered as silent.

Thirteen palindromes are common to the three species (including 2 with co-variations in

*Lo*). 13 others present in *Lpm* and *Lgh* correspond to quasi-palindromes (QPs) in *Lo* due to

mutated bases, and conversely, three *Lo* perfect palidromes are mutated in *Lpm* and *Lgh*.

Finally, only five palindromes are species specific. Two in particular are located in the

hypervariable intergenic spacer *ndh*F-*rpl32*, and are absent in *Lo* due to a large deletion of 160

nt.

**4.  Repeat distribution in LSC, SSC and IR segments**

In the IRa/IRb regions, repeats are only identified in the first 9 kb region between *rpl2* and

*ycf2*: a tandem repeat in the *Lpm rpl2* intron, and a tetranucleotide repeat, [TATC]*3, located

in the *ycf2* gene in the 3 species. In *ycf2* we also found 1 common palindrome (16 nt), a single

palindrome in *Lo* (20 nt, absent following an A:G mutation in the 2 other species), as well as a

shared tandem repeat (24 nt), and an additional 15 nt tandem repeat in *Lo* which adds 4 amino

acids to protein sequence.

In the SSC region, the repeats are almost all located in the intergenic and/or intronic

regions, with a hotspot between *ndh*F and *ccs*A. There is also a shared microsatellite in *ndh*F,

and a palidrome (16 nt) in *ccs*A which is absent in *Lo* (due to an A:C mutation), resulting in a

synonymous mutation (from isoleucine to leucine). We also observed multiple and various

repeats in the *ycf1* gene: 3 common poly-A repeats (from 10 to 13 nt), 3 species-specific

microsatellites (ATAG)*3 and (ACCA)*4 in *Lgh* and (CAAC)*3 in *Lo,* as well as two direct

repeats of 32 nt (37 nt spacing), which were absent from *Lo* due to a G:T SNP. Two tandem

repeats were also observed in *Lo* and *Lgh*. Neither of these repeats are at the origin of the

frameshift causing the pseudogenization of *ycf1* in *Lo*, this latter being due to a single deletion

of an A at position 3444 of the gene.

Finally, in the LSC region, the longest segment, which consequently contains the maximum

number of repeats, we still observed a preferential localization in the intergenic and intronic

regions since only genes *atp*A, *rpo*C2, *rpo*B, *psb*D, *psb*A, *psb*B, *ndh*K and *clp*P1 contain either

mononucleotic repeats (poly A and T), palindromes, or microsatellites (most often common to

the three species and without affecting the sequences of the proteins produced). As mentioned

earlier, the only exception is the *acc*D gene, which contains several direct and tandem repeats

in *Lgh* and *Lpm,* corresponding to a region of 174 nt (58 amino acids) missing in *Lo* and,

conversely, a direct repeat of 40 nucleotides, in a region of 147 nt (49 aa), which is present in

*Lo* and missing in the other two species. These tandem repeats lead to the presence of four

copies of 9 amino acids [DESENSNEE] in *Lgh* and *Lpm*, two of which form a larger duplication

of 17 aa [FLSDSDIDDESENSNEE]. Similarly, the TRs present only in *Lo* generate two perfect

9 amino acid repeats [EELSEDGEE], included in two longer degenerate repeats of 27 nt (Add.

Figure 6). It should be noted that though these TRs do not disturb the open reading phases, it is

still possible for them to form an intron which is not translated. Different functional studies will

be necessary to clarify this point. The presence of polymorphisms of the *acc*D gene between

*Lo* and the two species (*Lpm*, *Lgh*) is interesting because *acc*D, that encodes a subunit of acetyl-

CoA carboxylase (EC 6.4.1.2). This enzyme is essential in fatty acid synthesis and also

catalyzes the synthesis of malonyl-CoA, which is necessary for the growth of dicots, plant

fitness and leaf longevity, and is involved in the adaptation to specific ecological niches [70].

Large *acc*D expansions due to TRs have also been described in other plants such as *Medicago*

[71] and *Cupressophytes* [72]. Some authors have suggested that these inserted repeats are not

important for acetyl-CoA carboxylase activity as the reading frame is always preserved, and

they assume that these repeats must have a regulatory role [73].

**5.   Sequence Divergence Analysis and Polymorphic Loci Identification**

Determination of divergent regions by MVista, using *Lo* as a reference, confirmed that the

three *Ludwigia* sp. plastomes are well preserved if the SSC segment is oriented in the same way

(Add. Figure 7). Sliding window analysis (Figure 6) indicated variations in definite coding

regions, notably *clp*P, *acc*D, *ndh*5, *ycf1* with high Pi values, and to a lesser extent, *rps16*, *mat*K,

*ndh*K, *pet*A, *ccs*A and four tRNAs (*trn*H, *trn*D, *trn*T and *trn*N). These polymorphic *loci* could

be suitable for inferring genetic diversities in *Ludwigia* sp.

A comparative analysis of the sizes of protein coding genes sizes also shows that the *rps11*

gene initially annotated in *Lo* is shorter than those which have been newly annotated in *Lgh* and

*Lpm* (345 bp instead of 417 bp). Comparative analysis by BLAST shows that it is the long form

which is annotated in other Myrtales, and the observation of the locus in *Lo* shows a frameshift

mutation (deletion of a nucleotide in position 311). Functional analysis would be necessary to

check whether the *rps11* frameshift mutation produces shorter proteins that have lost their

function. And only obtaining the complete genome will verify whether copies of some of these

genes have been transferred to mitochondrial or nuclear genomes. Such *rps11* horizontal

transfers have been reported for this gene in the mitochondrial genomes of various plant

families[74]. This also applies to *ycf1*, found as a pseudogene in *Lo* (as specified previously),

although it is not known if this reflects a gene transfer or a complete loss of function [75][76].

Moreover, there is a deletion of nine nucleotides in the 3' region of the *rpl32* gene in *Lgh* and

*Lpm*, leading to a premature end of the translation and the deletion of the last 4 amino acids

[QRLD], which are replaced by a K. However, if we look carefully at the preserved region as

defined by the RPL32 domain (CHL00152, member of the superfamily CL09115), we see that

the later amino acids are not important for *rpl32* function since they are not found in the

orthologs.

Our results show that the Ka/Ks ratio is less than 1 for most genes (Figure 7). This indicates adaptive pressures to maintain the protein sequence except for *mat*K (1.17 between *Lgh* and *Lpm*), *acc*D (2.48 between *Lgh* and *Lo* and 2.16 between *Lpm* and *Lo*), *ycf2* (4.3 between both *Lgh-Lp* and *Lo*) and *ccs*A (1.4 between both *Lgh-Lpm* and *Lo*), showing a positive selection for these genes, and a possible key role in the processes of the species' ecological adaptations. As we have already described the variability in the *acc*D sequence, we will focus on *ycf2*, *mat*K, and *ccs*A variations.

Concerning *ccs*A, the variations observed, although significant, concern only five amino acids, and modifications do not seem to affect the C-type cytochrome synthase gene function.

Concerning *ycf2*, our analysis shows that this gene is highly polymorphic with 256 SNPs that provoke 10 deletions, 7 insertions, 21 conservative and 49 non-conservative substitutions in *Lo* (Add. Figure 8), compared to *Lgh* and *Lpm* (100 % identical). This gene has been shown as "variant" in other plant species such as *Helianthus tuberosus* [77].

The *mat*K gene has been used as a universal barcoding locus to enable species discrimination of terrestrial plants [78], and is often, together with the *rbc*L gene, the only known genetic resource for many plants. Thus, we propose a phylogenetic tree from *Ludwigia mat*K sequences (Figure 8). It should however be noted that this tree contains only 149 amino acids common to all the sequences (out of the 499 in the complete protein). As only three complete *Ludwigia* plastomes are available at the time of our study, we cannot specify whether these barcodes are faithful to the phylogenomic history of *Ludwigia* in the same way as the complete plastome. In any case, for this tree, we can see that *Lo* stands apart from the other *Ludwigia* sp., *Lpm* and *Lgh*, and that the *L. grandiflora* subsp. *hexapetala* belongs to the same branch as the species *L. ovalis* (aquatic taxon used in aquariums [79]), *L. stolonifera* (native to the Nile, found in a variety of habitats, from freshwater wetlands to brackish and marine waters) [80] and *L. adscendens* (common weed of rice fields in Asia) [81]. *Lpm* is in a sister branch, close to the *L. grandiflora* subsp. *hexapetala*, forming a phylogenetic group corresponding to subsect Jussiaea (in green, Figure 8).

**Discussion**

In the present study, we first sequenced and *de novo* assembled the chloroplast (cp) genomes of *Ludwigia peploides* (*Lpm*) and *Ludwigia grandiflora* (*Lgh*), two species belonging to the Onagraceae family. We employed a hybrid strategy and demonstrated the presence of two cp haplotypes in *Lgh* and one haplotype in *Lpm*, although the presence of both haplotypes in *Lpm* is likely. Furthermore, we compared these genomes with those of other species in the

Onagraceae family to expand our knowledge of genome organization and molecular evolution in these species.

Our findings demonstrate that the utilization of solely short reads has failed to produce complete *Ludwigia* plastomes, likely due to challenges posed by long repeats and rearrangements. On the other hand, relying solely on long reads resulted in a lower quality sequence due to insufficient coverage and sequencing errors. After conducting our research, we discovered that, for *Lgh* plastomes, hybrid assembly, which incorporates both long and short read sequences, resulted in the most superior complete assemblies. This innovative approach capitalizes on the advantages of both sequencing technologies, harnessing the accuracy of short read sequences and the length of long read sequences. In the case of our study on *Lgh* plastome reconstruction, hybrid assembly was the most complete and effective, similarly to studies on other chloroplasts, such as those in *Eucalyptus* [82], *Falcataria* [83], *Carex* [84] or *Cypripedium* [85].

In our study, we were able to identify the presence of two haplotypes in *Lgh*, which is a first for *Ludwigia* (and more broadly within Onagraceae)*,* as the plastome of *L. octovalvis* was only delivered in one haplotype [86].

Due to the unavailability of sequence data for *Ludwigia octovalvis* and the fact that we only have long reads for *Ludwigia peploides*, none of which large enough to cover the SSC/IR junctions, we are unable to conclusively identify the presence of these two forms in the *Ludwigia* genus. However, we believe that they are likely to be present. Unfortunately, the current representation of plastomes in GenBank primarily consists of short-read data, which may result in an underrepresentation of this polymorphism. It is unfortunate that structural heteroplasmy, which is expected to be widespread in angiosperms, has been overlooked. Existence of two plastome haplotypes has been identified in the related order of Myrtales (Eucalyptus sp.), in 58 species of Angiosperms, [87], Asparagales (*Ophrys apifera* orchid [88]), Brassicales (*Carica papaya*, *Vasconcellea pubescens* [89]), Solanales (*Solanum tuberosum* [90]), Laurales (Avocado *Persea americana* [91]) and Rhamnaceae (*Rhamnus crenata* [92]). However, the majority of reference plastomes in the current GenBank database (Release 260: April 15, 2024) are described as a single haplotype, indicating an underrepresentation of structural heteroplasmy in angiosperm chloroplasts. This underscores the importance of sequencing techniques, as the database is predominantly composed of short-read data (98%), which are less effective than long reads or hybrid assemblies at detecting flip-flop phenomena in the LSC region.

The chloroplast genome sizes for the three genera of Onagraceae subfam. Onagroideae varied as follows: *Circaea* sp. ranged from 155,817 bp to 156,024 bp, *Chamaenerion* sp. ranged from

159,496 bp to 160,416 bp, and Epilobium sp. ranged from 160,748 bp to 161,144 bp [93]. Our

study revealed that the size of the complete chloroplast of *Ludwigia* (Onagraceae subfamily

Ludwigioideae) ranged from 159,369 bp to 159,584 bp, which is remarkably similar to other

Onagraceae plants (average length of 162,030 bp). Furthermore, *Ludwigia* plastome sizes are

consistent with the range observed in Myrtales (between 152,214 to 171,315 bp [94]). In the

same way, similar overall GC content was found in *Ludwigia sp.* (from 37.3 to 37.4%), *Circaea*

*sp.* (37.7 to 37.8%), *Chamaenerion sp.* and *Epilobium sp.* (38.1 to 38.2%,[93]) and more

generally for the order Myrtales (36.9–38.9%, with the average GC content being 37%,[94]).

Higher GC content of the IR regions (43.5%) found in *Ludwigia* sp. has already been shown in

the Myrtales order (39.7–43.5%) and in other families/orders such as Amaranthaceae (order

Caryophyllales *[95]*) or Lamiaceae (order Lamiales [96]), and is mainly due to the presence of

the four GC rich rRNA genes.

The complete chloroplast genomes of the three *Ludwigia* species encoded an identical set of

134 genes including 85 protein-coding genes, 37 tRNA genes and eight ribosomal RNAs,

consistent with gene content found in the Myrtales order, with a gene number varying from 123

to 133 genes with 77–81 protein-coding genes, 29–31 tRNA gene and four rRNA genes [94].

Chloroplast genes have been selected during evolution due to their functional importance[97].

In our current study, we made the noteworthy discovery that *mat*K, *acc*D, *ycf2*, and *ccs*A genes

were subjected to positive selection pressure. These genes have frequently been reported in

literature as being associated with positive selection, and are known to play crucial roles in

plant development conditions. *Lgh* and *Lpm* are known to thrive in aquatic environments, where

they grow alongside rooted emergent aquatic plants, with their leaves and stems partially

submerged during growth, as reported by Wagner et al. in 2007 [1]. Both species possess the

unique ability of vegetative reproduction, enabling them to establish themselves rapidly in

diverse habitats, including terrestrial habitats, as noted by Haury et al [98]. Additionally, *Lo* is

a wetland plant that typically grows in gullies and at the edges of ponds, as documented by

Wagner *et al.* in 2007 [1]. Given their ability to adapt to different habitats, these species may

have evolved specialized mechanisms to cope with various abiotic stresses, such as reduced

carbon and oxygen availability or limited access to light in submerged or emergent conditions.

Concerning *matK*, Barthet et al [99] demonstrated the relationship between light and

developmental stages, and MatK maturase activity, suggesting important functions in plant

physiology. This gene has recently been largely reported to be under positive selection in an

aquatic plant (*Anubias* sp.,[100]), and more generally in terrestrial plants (*Pinus sp* [101]or

*Chrysosplenium sp.* [102]). The *acc*D gene has been described as an essential gene required for

leaf development [103] and longevity in tobacco (*Nicotiana tabacum*)[104]. Under drought stress, plant resistance can be increased by inhibiting *acc*D [105], and conversely, enhanced in response to flooding stress by upregulating *acc*D accumulation [106]. Hence, we can hypothesize that the positive selection observed on the *acc*D gene can be explained by the submerged and emerged constraints undergone by *Ludwigia* species. The *ycf2* gene seems to be subject to adaptive evolution in *Ludwigia species*. Its function, although still vague, would be to contribute to a protein complex generating ATP for the TIC machinery (proteins importing into the chloroplasts [107][108]), as well as plant cell survival [109][110]. The *ccs*A gene positive selection is found in some aquatic plants such as *Anubia sp.*[100], marine flowering plants as *Zostera* species [111], and some species of Lythraceae [105]. The *ccs*A gene is required for cytochrome c biogenesis [112] and this hemoprotein plays a key role in aerobic and anaerobic respiration, as well as photosynthesis [113]. Furthermore, we showed that *Lgh* colonization is supported by metabolic adjustments mobilizing glycolysis and fermentation pathways in terrestrial habitats, and the aminoacyl-tRNA biosynthesis pathway, which are key components of protein synthesis in aquatic habitats [114]. It can be assumed that the ability of *Ludwigia* to invade aquatic and wet environments, where the amount of oxygen and light can be variable, leads to a high selective pressure on genes involved in respiration and photosynthesis.

Molecular markers are often used to establish population genetic relationships through phylogenetic studies. Five chloroplasts (*rps16*, *rpl16*, *trn*L-*trn*F, *trn*L-CD, *trnG*) and two nuclear markers (ITS, *waxy*) were used in previous phylogeny studies of *Ludwigia sp.*[115]. However, no SSR markers had previously been made available for the *Ludwigia* genus, or more broadly, the Onagraceae. In this study, we identified 45 to 65 SSR markers depending on the *Ludwigia* species. Most of them were AT mononucleotides, as already recorded for other angiosperms [116][117]. In addition, we identified various genes with highly mutated regions that can also be used as SNP markers. Chloroplast SSRs (cpSSRs) represent potentially useful markers showing high levels of intraspecific variability due to the non-recombinant and uniparental inheritance of the plastomes [118][119]. Chloroplast SSR characteristics for *Ludwigia sp.* (location, type of SSR) were similar to those described in most plants. While the usual molecular markers used for phylogenetic analysis are nuclear DNA markers, cpSSRs have also been used to explore cytoplasmic diversity in many studies [120][121][122]. To conclude, the 13 highly variable loci and cpSSRs identified in this study are potential markers for population genetics or phylogenetic studies of *Ludwigia* species, and more generally, Onagraceae.

Concerning the MatK-based phylogenetic tree, its topology is generally congruent with the first molecular classification of Liu *et al.* [115] as all *Ludwigia* from sect *Jussiaea* (clade B1) and sect. *Ludwigia* (clade A1) and sect. *Isnardia* (clade A2) branched together. In this MatK-based tree, *Ludwigia prostrata*, a species absent from previously published phylogenetic studies, positions itself alone at the root of the *Ludwigia* tree. This species, sole member of section *Nematopyxis*, is related as having no close relatives [123], finding supported by our work. We also observed that *Ludwigia ovalis* branches within sect. *Jussiaea*, as its 258 amino acids partial MatK sequence (ca. half of the complete sequence) is identical to the MatK proteins of *L. grandiflora*, *L. stolonifera* and *L. adscendens*. Its phylogenetic placement remains unresolved: classified alone by Raven (1963) [5] and Wagner (2017) [22] in sect. *Miquelia*, later positioned by Liu et al. (2017)[4] within the *Isnardia-Microcarpium* section (using nuclear DNA) or as sister to it (using plastid DNA). For this reason, conducting a whole plastome analysis would be valuable to provide insights into *L. ovalis* phylogenetic positioning. Another species positioned on the margins of sect. *Isnardia* (clade A2) is *Ludwigia suffruticosa* (previously classified in sect. *Microcarpium*), which branches within sect. *Ludwigia* (clade A1). This positioning raises questions about the current grouping of sections *Isnardia*, *Michelia*, and *Microcarpium* into a single section *Isnardia* as proposed by Liu et al. (2023) [124] and highlights that plastid protein coding markers can provide differing phylogenetic insights. Finally, the last species positioned differently of this clade (clade B4) is *Ludwigia decurrens* (sect. *Pterocaulon*) which clusters with *L. leptocarpa* (clade B3) and *L. bonariensis* (clade B4a). However, it is important to note that in their study, Liu et al. (2017) indicate that clade B4 is moderately supported and that the two members of sect. *Pterocaulon*, *L. decurrens* and *L. nervosa*, diverge in all trees [4]. In summary, acquiring complete plastomes for *Ludwigia* sp. could significantly enhance our understanding of the phylogeny of this complex genus. Furthermore, comparing nuclear and plastid phylogenies would help determine if they reflect the same evolutionary history and whether plastid phylogeny alone can accurately reconstruct the phylogeny of *Ludwigia* genus.

**Conclusion**

In this study, we conducted the first-time sequencing and assembly of the complete plastomes of *Lpm* and *Lgh*, which are the only available genomic resources for functional analysis in both species. We were able to identify the existence of two haplotypes in *Lgh*, but further investigations will be necessary to confirm their presence in *Lo* and *Lpm*, and more broadly, within the *Ludwigia* genus. Comparison of all 10 Onagraceae plastomes revealed a high degree

of conservation in genome size, gene number, structure, and IR boundaries. However, to further elucidate the phylogenetic analysis and evolution in *Ludwigia* and Onagraceae, additional chloroplast genomes will be necessary, as highlighted in recent studies of Iris and Aristidoideae species [125].

**Declarations**

- Availability of data and materials

The datasets generated and/or analysed during the current study were available in GenBank (for *Lgh* haplotype 1, (LGH1) OR166254 and *Lgh* haplotype 2, (LGH2) OR166255; for *Lpm* haplotype, (LPM) OR166256). Chloroplastic short and long reads are available upon request.

- Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article

- Funding

The post-doctoral research grant of Anne-Laure Le Gac was supported by the Conseil regional Bretagne (SAD18001).

- Acknowledgements

We are grateful to Luis Portillo-Lemus for developing the high molecular weight genomic DNA extraction protocol. All sequencing experiments were performed at the PGTB (doi:10.15454/1.5572396583599417E12).

**Figure 1:** Two structural haplotypes of *L. grandiflora* plastomes representing the flip-flop organization of SSC segment.

**Figure 2:** Comparative results of *L. grandiflora* short read (SR) assemblies. **A:** Total number

of contigs obtained with the uncorrected (dark green) and corrected (light green) chloroplast

SRs for the 4 assemblers (ABySS, MEGAHIT, Velvet and SPAdes). **B:** Comparison of the size

of contigs assembled by the 4 tools using corrected or uncorrected SRs. **C:** Boxplot showing

the distribution of these contigs by size and the improvement brought by the prior correction of

the SRs with the long reads for each tool.

**692**

**Figure 3:** Circular representation of annotations plastomes in *Ludwigia octovalis*, *Ludwigia grandiflora* and *Ludwigia peploides* using ogdraw. Each card contains four circles. From the center outwards, the first circle shows forward and reverse repeats (red and green arcs, respectively). The next circle shows tandem repeats as bars. The third circle shows the microsatellite sequences. Finally, the fourth and fifth circles show the genes colored according to their functional categories (see colored legend). Only the haplotype 1 of *L. grandiflora* is represented as haplotype 2 only diverge by the orientation of the SSC segment. Accession numbers are indicated for each plastome.

**Figure 4:** Comparison of the borders of LSC, SSC, and IR regions in Onograceae plastomes. **A:** Comparison of the junction between large single-copy (LSC, light blue), inverted repeat (IR, orange) and short single-copy (SSC, light green) regions among the chloroplast genomes of *L. octovalvis*, *L. peploides* and *L. grandiflora* (both haplotypes). Genes are denoted by colored boxes and the gaps between genes and boundaries are indicated by base lengths (bp). JLB: junction line between LSC and IRb; JSB: junction line between IRb and SSC; JSA: junction line between SSC and IRa; JLA: junction line between IRa and LSC. **B:** Comparison of SSC boundaries in haplotype 1 (*L. peploides* and *L. grandiflora* haplotype 1) and haplotype 2 (*L. octovalvis* and *L. grandiflora* haplotype 2) plastomes.

**Figure 5:** Comparison of LSC, SSC and IR regions boundaries in Onograceae chloroplast genomes. Representative sequences from each genus have been chosen (noted R on the diagram) except for *Oenothera lindheimeri* (only 89.35 % identity with others *Oenothera*), *Circaea alpina* (99.5 % identity but all others *Circaea* are 99.9% identical) and *Chamaenerion conspersum* (99% but all others *Chamaenerion* are ca. 99.7 identical). As shown in Figure 7, the 3 *Ludwigia* plastomas had the same structure, *L. octovalvis* was chosen as a representative of this genus. JLB: junction of LSC/IRb; JSB: junction of IRb/SSC; JSA: junction of SSC/IRa; JLA: junction of IRa/LSC. Accession numbers : *Chamaenerion sp. conspersum* (MZ353638), *Chamaenerion sp. angustifolium* (NC_052848)**,** *Circaea sp. cordata* (NC_060876), *Circaea sp. alpina* (NC_061010), *Epilobium amurense* (NC_061015), *Oenothera villosa* subsp. *strigosa* (NC_061365) and *Oenothera lindheimeri* (MW538951).

**Figure 6:** Illustration of nucleotide diversity of the three *Ludwigia* chloroplast genome sequences. The graph was generated using DnaSP software version 6.0 (windows length: 800 bp, step size: 200 bp) [64][63]. The x-axis corresponds to the base sequence of the alignment, and the y-axis represents the nucleotide diversity ($\pi$ value). LSC, SSC and IR segments were indicated under the line representing the genes coding the proteins (in light blue) the tRNAs (in pink) and the rRNAs (in red). The genes marking diversity hotspots are noted at the top of the peaks.

736  **Figure 7:** The Ka/Ks ratios of the 80 protein-coding genes of *Ludwigia* plastomes. The blue

737  curve represents *L. grandiflora* versus *L. peploides*, purple curve denotes *L. grandiflora* versus

738  *L. octovalvis* and green curve *L. peploides* versus *L. octovalvis*. Four genes (*matK*, *accD*, *ycf2*

739  and *ccsA*) have Ka/Ks ratios greater than 1.0, whereas the Ka/Ks ratios of the other genes were

740  less than 1.0.

**Figure 8:** Phylogenetic tree based on *Ludwigia* MatK protein sequences. Only six *Ludwigia* sequences are complete (yellow star), the others correspond to amino acids ranging from 128 to 289 aa, with an average of 244 aa. Clades are named and colored regarding the *Ludwigia* phylogeny proposed by Liu et al. (2017) [4]. The sections are based on the works of Raven (1963) [5], Wagner et al (2017) [22] and Liu et al. (2023) [124]. The scale bar indicates the branch length.

749    Table 1. The general characteristics of the 3 *Ludwigia* plastomes

| | *L.octovalvis\** | *L.grandiflora* | *L.peploides* |
|---|---|---|---|
| **Size (bp)** | | | |
| | 159,396 | 159,584 | 159,537 |
| LSC | 90,183 | 90,272 | 90,156 |
| SSC | 19,703 | 19,788 | 19,799 |
| IR | 24,755 | 24,762 | 24,791 |
| **GC%** | | | |
| | 37.4 | 37.3 | 37.3 |
| LSC | 35.2 | 35.1 | 35.1 |
| SSC | 32 | 31.7 | 31.7 |
| IR | 43.5 | 43.5 | 43.4 |

\* KX827312 (ref)

753    Table 2. Genes present in the plastomes of *Ludwigia*

| FUNCTION | NAME |
|---|---|
| Photosynthesis | |
| **Rubisco** | *rbc*L |
| **Photosystem I (PSI)** | *psa*A, *psa*B, *psa*C, *psa*I, *psa*J |
| **PSI assembly factors** | *ycf*3# (*paf*I), *ycf*4 (*paf*II) |
| **Photosystem II** | *psb*A, *psb*B, *psb*C, *psb*D, *psb*E, *psb*F, *psb*H, *psb*I, *psb*J, *psb*K, *psb*L, *psb*M, *pbf*1 (*psb*N) *psb*T, *psb*Z |
| **ATP synthase** | *atp*A, *atp*B, *atp*E, *atp*F#, *atp*H, *atp*I |
| **Cytochrome *b6f*** | *pet*A, *pet*B#, *pet*D#, *pet*G, *pet*L, *pet*N |
| **Cytochrome biogenesis** | *ccs*A |
| **NADPH dehydrogenase** | *ndh*A#, *ndh*B**#, *ndh*C, *ndh*D, *ndh*E, *ndh*F, *ndh*G, *ndh*H, *ndh*I, *ndh*J |
| Transcription and translation | |
| **Transcription** | *rpo*A, *rpo*B, *rpo*C1#, *rpo*C2 |
| **Small ribosomal proteins** | *rps*2, *rps*3, *rps*4, *rps*7**, *rps*8, *rps*11, *rps*12**#, *rps*14, *rps*15, *rps*16#, *rps*18, *rps*19 |
| **Large ribosomal proteins** | *rpl*2**#, *rpl*14, *rpl*16#, *rpl*20, *rpl*22, *rpl*23**, *rpl*32, *rpl*33, *rpl*36 |
| **Translation initiation** | *inf*A |
| **Ribosomal RNA** | *rrn*5**, *rrn*4,5**, *rrn*16**, *rrn*23** |
| **Transfer RNA** | *trn*A-UGC**#,*trn*C-GCA,*trn*D-GUC,*trn*E-UUC,*trn*F-GAA,*trn*fM-CAU,*trn*G-GCC,*trn*G-UCC#,*trn*H-GUG,,*trn*I-CAU**,*trn*I-GAU**#,*trn*K-UUU#,*trn*L-CAA**,*trn*L-UAA#,*trn*L-UAG,*trn*M-CAU,*trn*N-GUU**,*trn*P-UGG,*trn*Q-UUG,*trn*R-ACG**,*trn*R-UCU,*trn*S-GCU,*trn*S-GGA,*trn*S-UGA,*trn*T-GGU,*trn*T-UGU,*trn*V-GAC**,*trn*V-UAC#,*trn*W-CCA,*trn*Y-GUA |
| Other functions | |
| **Group II intron splicing** | *mat*K |
| **Inorganic carbon uptake** | *cem*A |
| **Protease** | *clp*P1# |
| **Fatty acid synthesis/Heat tolerance** | *acc*D |
| **TIC machinery (protein import)** | *ycf*1 (Tic214), *ycf*2** |
| ***Unknown function pseudogene*** | *ycf*15** |
| | ** duplicated in IR region, # spliced genes |

## Table 3A : Tandem repeats

| Sequence | Lo | Lgh | Lpm | Length | Region | Locus | Comments |
|---|---|---|---|---|---|---|---|
| TTGTAGTCAGGGGTGTAGTACTAT | ▓ | | | 24 | IRs | ycf2 | |
| TAGAAGAGAGTGCAG | | X | X | 15 | IRs | ycf2 | 15 nt deletion in Lgh and Lpm |
| ATGAAATATCGTATAATGAAGTACCACACGAGTGGATAT | X | X | ▓ | 39 | IRs | rpl2 intron | 39 nt deletion in Lgh and Lo |
| AAAAATAGGATAGGAT | | X | X | 16 | LSC | ycf1-trnH-GUG | 56 nt deletion in Lgh and Lpm |
| TAAATTAATATCTATATA | | X | X | 18 | LSC | psbZ-trnG-GCC | 18 nt deletion in Lgh and Lpm |
| TTTTCTATCTATCTTATATCAA | | X | X | 22 | LSC | trnK-UUU-rps16 | 22 nt deletion in Lgh and Lpm |
| AGATCCATAACATCATCAAA | | X | X | 20 | LSC | rps16 intron | 22 nt deletion in Lgh and Lpm |
| TATTAGTTATTAATATTATTAGA | | X | X | 23 | LSC | trnP-UGG-psaJ | 23 nt deletion in Lgh and Lpm |
| AATAATATATAATAACTTAAATA | | X | X | 23 | LSC | rpl33-rps18 | 33 et 44 nt nt deletion in in Lgh et Lpm, respectively |
| TTTTTATTTAACATGCTATCAAATCAACAATGCCATACCGTAGGGCATCTGTT | | X | X | 53 | LSC | rpl20-clpP1 | 107 nt deletion in Lgh and Lpm |
| ATATATTTCGATTCAATTC | X | | X | 19 | LSC | trnH-GUG-psbA | 3 copies in a 57 nt deletion in Lo and Lpm |
| ATAGAAATATCAGTATTTGAGTG | X | | X | 23 | LSC | atpH-atpI | 23 nt deletion in Lo and Lpm |
| TTAATTTTAATTGAAGAA | X | | X | 18 | LSC | psbJ-psbL | 17 and 24 nt deletion in Lo and Lpm, respectively |
| TTAAAGAATATTAATATTC | imperfect TR | | | 19 | LSC | trnR-UCU-atpA | A -> C mutation in second copy in Lo |
| TATTATTATTATTAAT | X | X | | 16 | LSC | atpH-atpI | 16 nt deletion in Lgh and Lo |
| TCTAAGGCTGAAATAAGG | X | X | | 18 | LSC | pafI intron | 18 nt deletion in Lgh and Lo |
| TGTGAATCTATCTAT | | | X | 15 | LSC | trnS-UGA-psbZ | 8 nt deletion in Lpm |
| TTTTTTCTAGTA | | | | 12 | LSC | pafI intron | |
| CTAGTTATTGACATGG | | imperfect TR | imperfect TR | 16 | LSC | psaJ-rpl33 | G -> A mutation in second in Lpm et Lgh |
| ATTTTTATTAACTCT | X | | imperfect TR | 15 | SSC | ycf1 | T->A mutation in first copy in Lpm, other sequence in first copy in Lo |
| AATCAAATAGTTGAT | | X | X | 15 | SSC | ycf1 | other sequence in first copy of Lpm and Lgh |
| ATAATAATATATTTATTATTAATTAATA | X | | | 28 | SSC | ndhF-rpl32 | 160 nt deletion in Lo |

## Table 3B : Direct repeats

| Sequence | Lo | Lgh | Lpm | Size (nt) | Spacers (nt) | Region | Locus | Comments |
|---|---|---|---|---|---|---|---|---|
| TTCAATTGGAACGGACGATTCGTCAATCATCT | ▓ | | | 32 | 37 | SSC | ycf1 | 2 copies. In Lo, one mutation (G->A) in the second copie |
| CATCGATGATGAAAGTGAAAACAGTAATGAAGAGG | X | | | 35 | 28 - 22 - 11 | LSC | accD | 3 perfects copies and 1 mutated (G->A) copie in Lgh and Lpm. Region of 174 nt deleted in Lo |
| AGATGGTGAAGAACCTTATGAAGATGGTGAAGAACCTTATG | | X | X | 41 | 22 | LSC | accD | Region of 147 nt deleted in Lgh and Lpm |
| TATCAAATCAACAATGCCATACCGTAGGGCAT | | X | X | 32 | | LSC | rps12-clpP1 | 3 copies |
| TTAAGAGCCGTACAGGCACCTTTTGATGCATACGG | X | | | | 408 in Lpm , 406 in | LSC | clpP1 | 2 copies. In Lgh, one mutation (C->T) in the second copie |
| TTAAGAGCCGTACAGGCACTTTTTGATGCATACGG | X | | X | 35 | 811 | LSC | clpP1 intron 1- intron 2 | |
| TGCAATAGCCAAATGATGATGAGCAATATCAGTCAGCCATA | | | | 41 | 2178 | LSC | psaB-psaA | |

## Table 3C : Palindromic repeats

| Common perfect palindromic repeats | | Locus | Comment |
|---|---|---|---|
| AGACTCTCATGAGAGTCT ATTAAATAGAATATTCTATTTAAT | | trnC-GCA - cemN trnE-UUC-trnT-GGU | |
| TTGGTAAATTTACCAA | | psbD | |
| TTCATTTCAATTTCAATTGAAATTGAAATGAA | | trnC-CAU-vcf2 | 2 copies in IR |
| GAAAAAGGGCTTTTTC | | vcf2 | 2 copies in IR |
| TCTCAAATGATTAATCATTTGAGA | | trnL-UAA intron | |
| GGATTACTAGTAATCC | | trnD-GUC-trnY-GUA | |
| TTTGAATGCATTCAAA | | trnG-UCC intron | |
| ATATATTCGAATATAT | | trnG-UCC -trnR-UCU | |
| TAGTAATTAATTACTA | | trnG-GCC-trnfM-CAU | |
| CCAGTATGCATACTGG | | ndhK | |

| Common palindromic repeats with covariation | | Locus | |
|---|---|---|---|
| in L. octovalvis | in L. grandiflora et L. peploides | | |
| ATAGAATCTATATTCTATTAGAATATAGATTCTAT | ATCGAATCTATATTCTATTAGAATATAGATTCGAT | ndhC-trnV-UAC | |
| ATGTATATATATCGAT | ATCTATATATATAGAT | trnE-UUC-trnT-GGU | |

| Common palindromic and quasi-palindromic repeats | | Locus | Comment |
|---|---|---|---|
| in L. octovalvis | in L. grandiflora and L. peploides | | |
| TTTAACGAATATTAATATT t GTTAAA      TTAA c GAATATTAATATTCTTTAA | TTTAACGAATATTAATATTCGTTAAA TTAAAGAATATTAATATTCTTTAA | trnR-UCU-atpA | |
| AATTGTA c TTACAATT | AATTGTAATTACAATT | ccsA | |
| AGGAAGATTGATCAATCTT t CT | AGGAAGATTGATCAATCTTCCT | | |
| TTTA c TAATATTACTAA | TTAGTAATATTACTAA | rpl22 | |
| ATATAGAATAT c CTATAT | ATATAGAATATTCTATAT | | |
| ACATATCGATGATA g GT | ACATATCGATGATATGT | | |
| AATTACTAATTTCTATTACTATGTTCAATTGAACATAGTAATAGAAATTAGTAATT | AATTACTAATTTCTATTACT t TGTTCAATTGAACATAGTAATAGAAATTAGTAATT | atpH-atpI | |
| TAGTTAGAATTCTAACTA | TAGTT c GAATTCTAACTA | trnT-UGU-trnL-UAA | |
| TATTTTTTCTAGAAAAAATA | TATTTTTTCTAGAA g AAATA | ycf1 | 2 copies in IR |

| in L. octovalvis and L. peploides | in L. grandiflora | | |
|---|---|---|---|
| CCCATCAATCATGATTG t TGGG | CCCATCAATCATGATTGATGGG | psbN-trnD-GUC | |

| in L. octovalvis and L. grandiflora | in L. peploides | | |
|---|---|---|---|
| ATGAAAAAATCGATTTTTTTCAT | ATGATAAAAATAGATTTTT a TCAT | trnK-UUU-rps16 | |
| ATGAAAAAAATCGATTTTTTTCAT- ATGATAAAAATCGATTTTTATCAT | ATGATAAAAATA n ATTTTTATCAT | trnK-UUU-rps16 | |

| Unique palidromic repeats | | | |
|---|---|---|---|
| L. peploides | | | |
| TTATATATATATATATAA | | rpl32-ndhF | Full deletion in L. octovalvis , 6 bases deletion in L. grandiflora |
| L. octovalvis | | | |
| ATTGAAATTCGAATTTCAAT | | psbZ-trnG-GCC | Full deletion in L. grandiflora and L. peploides |
| L. peploides and L. grandiflora | | | |
| AAAAAATGGATCCATTTTTT | | trnL-UAG -rpl32 | 3 bases deleted and 3 bases mutated in L. octovalvis |
| AATATATTATTATAATAATATATT | | rpl32-ndhF | Full deletion in L. octovalvis |
| TATATTTATTATTAATTAATAATAAATATA | | rpl32-ndhF | Full deletion in L. octovalvis |
| L. octovalvis | | | |
| ATTGAAATTCGAATTTCAAT | | psbZ-trnG-GCC | Full deletion in L. grandiflora and L. peploides |
| L. peploides and L. grandiflora | | | |
| AAAAAATGGATCCATTTTTT | | trnL-UAG -rpl32 | 3 bases deleted and 3 bases mutated in L. octovalvis |
| AATATATTATTATAATAATATT | | rpl32-ndhF | Full deletion in L. octovalvis |
| TATATTTATTATTAATTAATAATAAATATA | | rpl32-ndhF | Full deletion in L. octovalvis |

 *Lo = Ludwigia octovalvis; Lgh = L. grandiflora* subsp. *hexapetala; Lpm = L. peploides* subsp. *montevidensis.*

**REFERENCES**

[1]     W. L. Wagner, P. C. Hoch, and P. H. Raven, "Revised classification of the Onagraceae," *Systematic Botany Monographs*, 2007.

[2]     R. A. Levin *et al.*, "Family-level relationships of Onagraceae based on chloroplast rbcL and ndhF data," *Am J Bot*, vol. 90, no. 1, 2003, doi: 10.3732/ajb.90.1.107.

[3]     R. A. Levin *et al.*, "Paraphyly in Tribe Onagreae: Insights into Phylogenetic Relationships of Onagraceae Based on Nuclear and Chloroplast Sequence Data," *Syst Bot*, vol. 29, no. 1, 2004, doi: 10.1600/036364404772974293.

[4]     S. H. Liu, P. C. Hoch, M. Diazgranados, P. H. Raven, and J. C. Barber, "Multi-locus phylogeny of ludwigia (Onagraceae): Insights on infra- generic relationships and the current classification of the genus," *Taxon*, vol. 66, no. 5, 2017, doi: 10.12705/665.7.

[5]     Raven P.H., "The_Old_World_species_of_Ludwigia_including_Jussia," *Reinwardtia*, vol. 6, pp. 327–427, 1963.

[6]     S. Dandelot, R. Verlaque, A. Dutartre, and A. Cazaubon, "Ecological, dynamic and taxonomic problems due to Ludwigia (Onagraceae) in France," in *Hydrobiologia*, 2005. doi: 10.1007/s10750-005-4455-0.

[7]     A. M. Reddy *et al.*, "Biological control of invasive water primroses, Ludwigia spp., in the United States: A feasibility assessment." J. Aquat. Plant Manage. 59s: 2021

[8]     A. Hussner, M. Windhaus, and U. Starfinger, "From weed biology to successful control: an example of successful management of Ludwigia grandiflora in Germany," *Weed Res*, vol. 56, no. 6, 2016, doi: 10.1111/wre.12224.

[9]     B. J. Grewell, M. D. Netherland, and M. J. Skaer Thomason, "Establishing research and management priorities for invasive water primroses (Ludwigia spp.)," *Aquatic Plant Control Research Program, US Army Corps of Engineers, Engineer Research and Development Center, Environmental Laboratory Technical Report ERDC/ELTR-15-X*, no. February, 2016.

[10]    E. Lambert, A. Dutartre, J. Coudreuse, and J. Haury, "Relationships between the biomass production of invasive Ludwigia species and physical properties of habitats in France," *Hydrobiologia*, vol. 656, no. 1, 2010, doi: 10.1007/s10750-010-0440-3.

[11]    J. Haury, A. Druel, T. Cabral, Y. Paulet, M. Bozec, and J. Coudreuse, "Which adaptations of some invasive Ludwigia spp. (Rosidae, Onagraceae) populations occur in contrasting hydrological conditions in Western France?," *Hydrobiologia*, vol. 737, no. 1, 2014, doi: 10.1007/s10750-014-1815-7.

[12]    K. Billet, J. Genitoni, M. Bozec, D. Renault, and D. Barloy, "Aquatic and terrestrial morphotypes of the aquatic invasive plant, Ludwigia grandiflora, show distinct morphological and metabolomic responses," *Ecol Evol*, vol. 8, no. 5, 2018, doi: 10.1002/ece3.3848.

[13]    M. Gioria, P. E. Hulme, D. M. Richardson, and P. Pyšek, "Annual Review of Plant Biology Why Are Invasive Plants Successful?," *Annu. Rev. Plant Biol. 2023*, vol. 74, p. 2023, 2023, doi: 10.1146/annurev-arplant-070522.

[14]    L. Moravcová, P. Pyšek, V. Jarošík, and J. Pergl, "Getting the right traits: Reproductive and dispersal characteristics predict the invasiveness of herbaceous plant species," *PLoS One*, vol. 10, no. 4, Apr. 2015, doi: 10.1371/journal.pone.0123634.

[15]    R. A. Marks, S. Hotaling, P. B. Frandsen, and R. VanBuren, "Representation and participation across 20 years of plant genome sequencing," *Nat Plants*, vol. 7, no. 12, 2021, doi: 10.1038/s41477-021-01031-8.

[16]    D. Barloy, L. Portillo-Lemus, S. Krueger-Hadfield, V. Huteau, and O. Coriton, "Genomic relationships among diploid and polyploid species of the genus Ludwigia L. section Jussiaea using a combination of molecular cytogenetic, morphological, and

crossing investigations," *Peer Community Journal*, vol. 4, 2024, doi:
10.24072/pcjournal.364.

[17] S. H. Liu, C. Edwards, P. C. Hoch, P. H. Raven, and J. C. Barber, "Complete plastome
sequence of ludwigia octovalvis (Onagraceae), a globally distributed wetland plant,"
*Genome Announc*, vol. 4, no. 6, 2016, doi: 10.1128/genomeA.01274-16.

[18] E. Zardini and P. H. Raven, "A New Section of Ludwigia (Onagraceae) with a Key to
the Sections of the Genus," *Syst Bot*, vol. 17, no. 3, 1992, doi: 10.2307/2419486.

[19] P. C. Hoch, W. L. Wagner, and P. H. Raven, "The correct name for a section of Ludwigia
L. (Onagraceae)," *PhytoKeys*, vol. 50, no. 1, 2015, doi: 10.3897/phytokeys.50.4887.

[20] Y. Hu, Q. Zhang, G. Rao, and Sodmergen, "Occurrence of plastids in the sperm cells of
caprifoliaceae: Biparental plastid inheritance in angiosperms is unilaterally derived from
maternal inheritance," *Plant Cell Physiol*, vol. 49, no. 6, 2008, doi: 10.1093/pcp/pcn069.

[21] Q. Zhang and Sodmergen, "Why does biparental plastid inheritance revive in
angiosperms?," *J Plant Res*, vol. 123, no. 2, 2010, doi: 10.1007/s10265-009-0291-z.

[22] W. L. Wagner, P. C. Hoch, and P. H. Raven, *Systematic botany monographs: Revised
classification of the Onagraceae*. 2007.

[23] K. Jones and R. E. Cleland, "Oenothera, Cytogenetics and Evolution," *Kew Bull*, vol.
29, no. 1, 1974, doi: 10.2307/4108389.

[24] U. K. Schmitz and K. V. Kowallik, "Plastid inheritance in Epilobium," *Curr Genet*, vol.
11, no. 1, 1986, doi: 10.1007/BF00389419.

[25] N. Sato, "Are cyanobacteria an ancestor of chloroplasts or just one of the gene donors
for plants and algae?," *Genes (Basel)*, vol. 12, no. 6, 2021, doi: 10.3390/genes12060823.

[26] J. M. Gualberto, D. Mileshina, C. Wallet, A. K. Niazi, F. Weber-Lotfi, and A. Dietrich,
"The plant mitochondrial genome: Dynamics and maintenance," *Biochimie*, vol. 100, no.
1. 2014. doi: 10.1016/j.biochi.2013.09.016.

[27] J. Tonti-Filippini, P. G. Nevill, K. Dixon, and I. Small, "What can we do with 1000
plastid genomes?," *Plant Journal*, vol. 90, no. 4, 2017, doi: 10.1111/tpj.13491.

[28] D. J. Oldenburg and A. J. Bendich, "The linear plastid chromosomes of maize: terminal
sequences, structures, and implications for DNA replication," *Curr Genet*, vol. 62, no.
2, 2016, doi: 10.1007/s00294-015-0548-0.

[29] A. D. Twyford and R. W. Ness, "Strategies for complete plastid genome sequencing,"
*Mol Ecol Resour*, vol. 17, no. 5, 2017, doi: 10.1111/1755-0998.12626.

[30] W. Wang, M. Schalamun, A. Morales-Suarez, D. Kainer, B. Schwessinger, and R.
Lanfear, "Assembly of chloroplast genomes with long- and short-read data: A
comparison of approaches using Eucalyptus pauciflora as a test case," *BMC Genomics*,
vol. 19, no. 1, 2018, doi: 10.1186/s12864-018-5348-8.

[31] W. Wang, R. Lanfear, and B. Gaut, "Long-Reads Reveal That the Chloroplast Genome
Exists in Two Distinct Versions in Most Plants," *Genome Biol Evol*, vol. 11, no. 12,
2019, doi: 10.1093/gbe/evz256.

[32] M. Ferrarini *et al.*, "An evaluation of the PacBio RS platform for sequencing and de novo
assembly of a chloroplast genome," *BMC Genomics*, vol. 14, no. 1, 2013, doi:
10.1186/1471-2164-14-670.

[33] M. Jain *et al.*, "Nanopore sequencing and assembly of a human genome with ultra-long
reads," *Nat Biotechnol*, vol. 36, no. 4, 2018, doi: 10.1038/nbt.4060.

[34] F. J. Rang, W. P. Kloosterman, and J. de Ridder, "From squiggle to basepair:
Computational approaches for improving nanopore sequencing read accuracy," *Genome
Biology*, vol. 19, no. 1. 2018. doi: 10.1186/s13059-018-1462-9.

[35] A. Scheunert, M. Dorfner, T. Lingl, and C. Oberprieler, "Can we use it? On the utility of
de novo and reference-based assembly of Nanopore data for plant plastome sequencing,"
*PLoS One*, vol. 15, no. 3, 2020, doi: 10.1371/journal.pone.0226234.

[36] A. M. Bedoya and S. Madriñán, "Evolution of the aquatic habit in Ludwigia (Onagraceae): Morpho-anatomical adaptive strategies in the Neotropics," *Aquat Bot*, vol. 120, no. PB, 2015, doi: 10.1016/j.aquabot.2014.10.005.

[37] S. H. Liu *et al.*, "Disentangling Reticulate Evolution of North Temperate Haplostemonous Ludwigia (Onagraceae)," *Annals of the Missouri Botanical Garden*, vol. 105, no. 2, 2020, doi: 10.3417/2020479.

[38] M. Panova *et al.*, "DNA extraction protocols for whole-genome sequencing in marine organisms," in *Methods in Molecular Biology*, vol. 1452, 2016. doi: 10.1007/978-1-4939-3774-5_2.

[39] C. Belser *et al.*, "Chromosome-scale assemblies of plant genomes using nanopore long reads and optical maps," *Nature Plants*, vol. 4, no. 11. 2018. doi: 10.1038/s41477-018-0289-4.

[40] S. Chen, Y. Zhou, Y. Chen, and J. Gu, "Fastp: An ultra-fast all-in-one FASTQ preprocessor," in *Bioinformatics*, 2018. doi: 10.1093/bioinformatics/bty560.

[41] J. J. Jin *et al.*, "GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of organelle genomes," *Genome Biol*, vol. 21, no. 1, 2020, doi: 10.1186/s13059-020-02154-5.

[42] N. Dierckxsens, P. Mardulyn, and G. Smits, "NOVOPlasty: De novo assembly of organelle genomes from whole genome data," *Nucleic Acids Res*, vol. 45, no. 4, 2017, doi: 10.1093/nar/gkw955.

[43] D. R. Zerbino and E. Birney, "Velvet: Algorithms for de novo short read assembly using de Bruijn graphs," *Genome Res*, vol. 18, no. 5, pp. 821–829, May 2008, doi: 10.1101/gr.074492.107.

[44] J. T. Simpson, K. Wong, S. D. Jackman, J. E. Schein, S. J. M. Jones, and I. Birol, "ABySS: A parallel assembler for short read sequence data," *Genome Res*, vol. 19, no. 6, 2009, doi: 10.1101/gr.089532.108.

[45] S. D. Jackman *et al.*, "ABySS 2.0: Resource-efficient assembly of large genomes using a Bloom filter," *Genome Res*, vol. 27, no. 5, 2017, doi: 10.1101/gr.214346.116.

[46] D. Li *et al.*, "MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies and community practices," *Methods*, vol. 102. 2016. doi: 10.1016/j.ymeth.2016.02.020.

[47] A. Bankevich *et al.*, "SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing," *Journal of Computational Biology*, vol. 19, no. 5, 2012, doi: 10.1089/cmb.2012.0021.

[48] R. Chikhi and P. Medvedev, "Informed and automated k-mer size selection for genome assembly," *Bioinformatics*, vol. 30, no. 1, 2014, doi: 10.1093/bioinformatics/btt310.

[49] S. Koren, B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman, and A. M. Phillippy, "Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting and repeat separation," *Genome Res*, vol. 27, no. 5, 2017, doi: 10.1101/gr.215087.116.

[50] G. Holley *et al.*, "Ratatosk: hybrid error correction of long reads enables accurate variant calling and assembly," *Genome Biol*, vol. 22, no. 1, 2021, doi: 10.1186/s13059-020-02244-4.

[51] M. Kolmogorov, J. Yuan, Y. Lin, and P. A. Pevzner, "Assembly of long, error-prone reads using repeat graphs," *Nat Biotechnol*, vol. 37, no. 5, 2019, doi: 10.1038/s41587-019-0072-8.

[52] A. Gurevich, V. Saveliev, N. Vyahhi, and G. Tesler, "QUAST: Quality assessment tool for genome assemblies," *Bioinformatics*, vol. 29, no. 8, pp. 1072–1075, Apr. 2013, doi: 10.1093/bioinformatics/btt086.

919 [53] R. R. Wick, M. B. Schultz, J. Zobel, and K. E. Holt, "Bandage: Interactive visualization
920    of de novo genome assemblies," *Bioinformatics*, vol. 31, no. 20, 2015, doi:
921    10.1093/bioinformatics/btv383.

922 [54] M. Tillich *et al.*, "GeSeq - Versatile and accurate annotation of organelle genomes,"
923    *Nucleic Acids Res*, vol. 45, no. W1, 2017, doi: 10.1093/nar/gkx391.

924 [55] X. Zhong, "Assembly, annotation and analysis of chloroplast genomes," 2020. [Doctoral
925    Thesis, The University of Western Australia].

926 [56] S. Greiner, P. Lehwark, and R. Bock, "OrganellarGenomeDRAW (OGDRAW) version
927    1.3.1: Expanded toolkit for the graphical visualization of organellar genomes," *Nucleic
928    Acids Res*, vol. 47, no. W1, 2019, doi: 10.1093/nar/gkz238.

929 [57] P. Lehwark and S. Greiner, "GB2sequin - A file converter preparing custom GenBank
930    files for database submission," *Genomics*, vol. 111, no. 4, 2019, doi:
931    10.1016/j.ygeno.2018.05.003.

932 [58] S. Beier, T. Thiel, T. Münch, U. Scholz, and M. Mascher, "MISA-web: A web server for
933    microsatellite prediction," *Bioinformatics*, vol. 33, no. 16, 2017, doi:
934    10.1093/bioinformatics/btx198.

935 [59] M. Gurusaran, D. Ravella, and K. Sekar, "RepEx: Repeat extractor for biological
936    sequences," *Genomics*, vol. 102, no. 4, pp. 403–408, Oct. 2013, doi:
937    10.1016/j.ygeno.2013.07.005.

938 [60] G. Benson, "Tandem repeats finder: A program to analyze DNA sequences," *Nucleic
939    Acids Res*, vol. 27, no. 2, 1999, doi: 10.1093/nar/27.2.573.

940 [61] K. A. Frazer, L. Pachter, A. Poliakov, E. M. Rubin, and I. Dubchak, "VISTA:
941    Computational tools for comparative genomics," *Nucleic Acids Res*, vol. 32, no. WEB
942    SERVER ISS., 2004, doi: 10.1093/nar/gkh458.

943 [62] M. Brudno *et al.*, "LAGAN and Multi-LAGAN: Efficient tools for large-scale multiple
944    alignment of genomic DNA," *Genome Research*, vol. 13, no. 4. 2003. doi:
945    10.1101/gr.926603.

946 [63] J. Rozas and R. Rozas, "DnaSP version 3: An integrated program for molecular
947    population genetics and molecular evolution analysis," *Bioinformatics*, vol. 15, no. 2.
948    1999. doi: 10.1093/bioinformatics/15.2.174.

949 [64] J. Rozas *et al.*, "DnaSP 6: DNA sequence polymorphism analysis of large data sets," *Mol
950    Biol Evol*, vol. 34, no. 12, 2017, doi: 10.1093/molbev/msx248.

951 [65] A. Amiryousefi, J. Hyvönen, and P. Poczai, "IRscope: an online program to visualize the
952    junction sites of chloroplast genomes," *Bioinformatics*, vol. 34, no. 17, 2018, doi:
953    10.1093/bioinformatics/bty220.

954 [66] C. Chen *et al.*, "TBtools: An Integrative Toolkit Developed for Interactive Analyses of
955    Big Biological Data," *Mol Plant*, vol. 13, no. 8, 2020, doi: 10.1016/j.molp.2020.06.009.

956 [67] K. Katoh, K. Misawa, K. I. Kuma, and T. Miyata, "MAFFT: A novel method for rapid
957    multiple sequence alignment based on fast Fourier transform," *Nucleic Acids Res*, vol.
958    30, no. 14, 2002, doi: 10.1093/nar/gkf436.

959 [68] A. Stamatakis, "RAxML version 8: A tool for phylogenetic analysis and post-analysis of
960    large phylogenies," *Bioinformatics*, vol. 30, no. 9, pp. 1312–1313, May 2014, doi:
961    10.1093/bioinformatics/btu033.

962 [69] J. Ou and L. J. Zhu, "trackViewer: a Bioconductor package for interactive and integrative
963    visualization of multi-omics data," *Nature Methods*, vol. 16, no. 6. Nature Publishing
964    Group, pp. 453–454, Jun. 01, 2019. doi: 10.1038/s41592-019-0430-y.

965 [70] T. Konishi and Y. Sasaki, "Compartmentalization of two forms of acetyl-CoA
966    carboxylase in plants and the origin of their tolerance toward herbicides," *Proc Natl Acad
967    Sci U S A*, vol. 91, no. 9, pp. 3598–3601, 1994, doi: 10.1073/pnas.91.9.3598.

[71] S. Wu *et al.*, "Extensive genomic rearrangements mediated by repetitive sequences in
plastomes of Medicago and its relatives," *BMC Plant Biol*, vol. 21, no. 1, p. 421, 2021,
doi: 10.1186/s12870-021-03202-3.
[72] J. Li, Y. Su, and T. Wang, "The Repeat Sequences and Elevated Substitution Rates of
the Chloroplast accD Gene in Cupressophytes," *Front Plant Sci*, vol. 9, p. 533, 2018,
doi: 10.3389/fpls.2018.00533.
[73] C. Gurdon and P. Maliga, "Two distinct plastid genome configurations and
unprecedented intraspecies length variation in the accD coding region in Medicago
truncatula," *DNA Res*, vol. 21, no. 4, pp. 417–427, 2014, doi: 10.1093/dnares/dsu007.
[74] A. O. Richardson and J. D. Palmer, "Horizontal gene transfer in plants," *J Exp Bot*, vol.
58, no. 1, pp. 1–9, 2007, doi: 10.1093/jxb/erl148.
[75] J. de Vries, F. L. Sousa, B. Bolter, J. Soll, and S. B. Gould, "YCF1: A Green TIC?,"
*Plant Cell*, vol. 27, no. 7, pp. 1827–1833, 2015, doi: 10.1105/tpc.114.135541.
[76] E. Filip and L. Skuza, "Horizontal Gene Transfer Involving Chloroplasts," *Int J Mol Sci*,
vol. 22, no. 9, 2021, doi: 10.3390/ijms22094484.
[77] Q. Zhong, S. Yang, X. Sun, L. Wang, and Y. Li, "The complete chloroplast genome of
the Jerusalem artichoke (Helianthus tuberosus L.) and an adaptive evolutionary analysis
of the ycf2 gene," *PeerJ*, vol. 7, p. e7596, 2019, doi: 10.7717/peerj.7596.
[78] S. Antil *et al.*, "DNA barcoding, an effective tool for species identification: a review,"
*Mol Biol Rep*, vol. 50, no. 1, pp. 761–775, 2023, doi: 10.1007/s11033-022-08015-7.
[79] J. Li *et al.*, "Removal effects of aquatic plants on high-concentration phosphorus in
wastewater during summer," *J Environ Manage*, vol. 324, p. 116434, 2022, doi:
10.1016/j.jenvman.2022.116434.
[80] A. T. Soliman, R. S. Hamdy, and A. B. Hamed, "Ludwigia stolonifera (Guill. & Perr.)
PH Raven, insight into its phenotypic plasticity, habitat diversity and associated species,"
*Egyptian Journal of Botany*, vol. 58, no. 3, pp. 605–626, 2018.
[81] A. Kamoshita, H. Ikeda, J. Yamagishi, B. Lor, and M. Ouk, "Residual effects of
cultivation methods on weed seed banks and weeds in Cambodia," *Weed Biol Manag*,
vol. 16, no. 3, pp. 93–107, 2016.
[82] W. Wang, M. Schalamun, A. Morales-Suarez, D. Kainer, B. Schwessinger, and R.
Lanfear, "Assembly of chloroplast genomes with long-and short-read data: a comparison
of approaches using Eucalyptus pauciflora as a test case," *BMC Genomics*, vol. 19, pp.
1–15, 2018.
[83] V. P. D. Anita, D. D. Matra, and U. J. Siregar, "Chloroplast genome draft assembly of
Falcataria moluccana using hybrid sequencing technology," *BMC Res Notes*, vol. 16, no.
1, p. 31, 2023, doi: 10.1186/s13104-023-06290-6.
[84] S. Xu *et al.*, "Chloroplast genomes of four Carex species: Long repetitive sequences
trigger dramatic changes in chloroplast genome structure," *Front Plant Sci*, vol. 14, p.
1100876, 2023, doi: 10.3389/fpls.2023.1100876.
[85] Y. Y. Guo, J. X. Yang, H. K. Li, and H. S. Zhao, "Chloroplast Genomes of Two Species
of Cypripedium: Expanded Genome Size and Proliferation of AT-Biased Repeat
Sequences," *Front Plant Sci*, vol. 12, p. 609729, 2021, doi: 10.3389/fpls.2021.609729.
[86] S.-H. Liu, C. Edwards, P. C. Hoch, P. H. Raven, and J. C. Barber, "Complete plastome
sequence of Ludwigia octovalis (Onagraceae), a globally distributed wetland plant,"
*Genome Announc*, vol. 4, no. 6, pp. e01274-16, 2016.
[87] W. Wang and R. Lanfear, "Long-reads reveal that the chloroplast genome exists in two
distinct versions in most plants," *Genome Biol Evol*, vol. 11, no. 12, pp. 3372–3381,
2019.
[88] R. M. Bateman, P. J. Rudall, A. R. M. Murphy, R. S. Cowan, D. S. Devey, and O. A.
Perez-Escobar, "Whole plastomes are not enough: phylogenomic and morphometric

exploration at multiple demographic levels of the bee orchid clade Ophrys sect. Sphegodes," *J Exp Bot*, vol. 72, no. 2, pp. 654–681, 2021, doi: 10.1093/jxb/eraa467.

[89] Z. Lin *et al.*, "Comparative analysis of chloroplast genomes in Vasconcellea pubescens A.DC. and Carica papaya L," *Sci Rep*, vol. 10, no. 1, p. 15799, 2020, doi: 10.1038/s41598-020-72769-y.

[90] G. A. Lihodeevskiy and E. P. Shanina, "The Use of Long-Read Sequencing to Study the Phylogenetic Diversity of the Potato Varieties Plastome of the Ural Selection," *Agronomy*, vol. 12, no. 4, p. 846, 2022.

[91] O. Nath *et al.*, "A haplotype resolved chromosomal level avocado genome allows analysis of novel avocado genes," *Hortic Res*, vol. 9, p. uhac157, 2022, doi: 10.1093/hr/uhac157.

[92] K. Wanichthanarak *et al.*, "Revisiting chloroplast genomic landscape and annotation towards comparative chloroplast genomes of Rhamnaceae," *BMC Plant Biol*, vol. 23, no. 1, p. 59, 2023.

[93] Y. Luo *et al.*, "Comparative Analysis of Complete Chloroplast Genomes of 13 Species in Epilobium, Circaea, and Chamaenerion and Insights Into Phylogenetic Relationships of Onagraceae," *Front Genet*, vol. 12, p. 730495, 2021, doi: 10.3389/fgene.2021.730495.

[94] X. F. Zhang, J. B. Landis, H. X. Wang, Z. X. Zhu, and H. F. Wang, "Comparative analysis of chloroplast genome structure and molecular dating in Myrtales," *BMC Plant Biol*, vol. 21, no. 1, p. 219, 2021, doi: 10.1186/s12870-021-02985-9.

[95] J. Xu, X. Shen, B. Liao, J. Xu, and D. Hou, "Comparing and phylogenetic analysis chloroplast genome of three Achyranthes species," *Sci Rep*, vol. 10, no. 1, p. 10818, 2020, doi: 10.1038/s41598-020-67679-y.

[96] C. Lian *et al.*, "Comparative analysis of chloroplast genomes reveals phylogenetic relationships and intraspecific variation in the medicinal plant Isodon rubescens," *PLoS One*, vol. 17, no. 4, p. e0266546, 2022, doi: 10.1371/journal.pone.0266546.

[97] T. K. Mohanta, A. K. Mishra, A. Khan, A. Hashem, E. F. Abd Allah, and A. Al-Harrasi, "Gene Loss and Evolution of the Plastome," *Genes (Basel)*, vol. 11, no. 10, 2020, doi: 10.3390/genes11101133.

[98] J. Haury, A. Druel, T. Cabral, Y. Paulet, M. Bozec, and J. Coudreuse, "Which adaptations of some invasive Ludwigia spp.(Rosidae, Onagraceae) populations occur in contrasting hydrological conditions in Western France?," *Hydrobiologia*, vol. 737, pp. 45–56, 2014.

[99] M. M. Barthet and K. W. Hilu, "Expression of matK: functional and evolutionary implications," *Am J Bot*, vol. 94, no. 8, pp. 1402–1412, 2007.

[100] L. Li, C. Liu, K. Hou, and W. Liu, "Comparative Analyses of Plastomes of Four Anubias (Araceae) Taxa, Tropical Aquatic Plants Endemic to Africa," *Genes (Basel)*, vol. 13, no. 11, 2022, doi: 10.3390/genes13112043.

[101] U. Zeb *et al.*, "Comparative genome sequence and phylogenetic analysis of chloroplast for evolutionary relationship among Pinus species," *Saudi J Biol Sci*, vol. 29, no. 3, pp. 1618–1627, 2022, doi: 10.1016/j.sjbs.2021.10.070.

[102] Z. Wu *et al.*, "Analysis of six chloroplast genomes provides insight into the evolution of Chrysosplenium (Saxifragaceae)," *BMC Genomics*, vol. 21, no. 1, p. 621, 2020, doi: 10.1186/s12864-020-07045-4.

[103] V. Kode, E. A. Mudd, S. Iamtham, and A. Day, "The tobacco plastid accD gene is essential and is required for leaf development," *Plant J*, vol. 44, no. 2, pp. 237–244, 2005, doi: 10.1111/j.1365-313X.2005.02533.x.

[104] Y. Madoka, K. Tomizawa, J. Mizoi, I. Nishida, Y. Nagano, and Y. Sasaki, "Chloroplast transformation with modified accD operon increases acetyl-CoA carboxylase and causes

extension of leaf longevity and increase in seed yield in tobacco," *Plant Cell Physiol*,
vol. 43, no. 12, pp. 1518–1525, 2002, doi: 10.1093/pcp/pcf172.

[105] H. Gu *et al.*, "Drought stress triggers proteomic changes involving lignin, flavonoids and
fatty acids in tea plants," *Sci Rep*, vol. 10, no. 1, p. 15504, 2020, doi: 10.1038/s41598-
020-72596-1.

[106] B. Bharadwaj *et al.*, "Physiological and genetic responses of lentil (Lens culinaris) under
flood stress," *Plant Stress*, p. 100130, 2023.

[107] S. Kikuchi *et al.*, "A Ycf2-FtsHi Heteromeric AAA-ATPase Complex Is Required for
Chloroplast Protein Import," *Plant Cell*, vol. 30, no. 11, pp. 2677–2703, 2018, doi:
10.1105/tpc.18.00357.

[108] T. B. Schreier *et al.*, "Plastidial NAD-Dependent Malate Dehydrogenase: A
Moonlighting Protein Involved in Early Chloroplast Development through Its Interaction
with an FtsH12-FtsHi Protease Complex," *Plant Cell*, vol. 30, no. 8, pp. 1745–1769,
2018, doi: 10.1105/tpc.18.00121.

[109] A. Drescher, S. Ruf, T. Calsa Jr., H. Carrer, and R. Bock, "The two largest chloroplast
genome-encoded open reading frames of higher plants are essential genes," *Plant J*, vol.
22, no. 2, pp. 97–104, 2000, doi: 10.1046/j.1365-313x.2000.00722.x.

[110] J. Xing *et al.*, "The plastid-encoded protein Orf2971 is required for protein translocation
and chloroplast quality control," *Plant Cell*, vol. 34, no. 9, pp. 3383–3399, 2022, doi:
10.1093/plcell/koac180.

[111] J. Chen *et al.*, "Chloroplast genomic comparison provides insights into the evolution of
seagrasses," *BMC Plant Biol*, vol. 23, no. 1, p. 104, 2023, doi: 10.1186/s12870-023-
04119-9.

[112] Z. Xie and S. Merchant, "The plastid-encoded ccsA gene is required for heme attachment
to chloroplast c-type cytochromes," *J Biol Chem*, vol. 271, no. 9, pp. 4632–4639, 1996,
doi: 10.1074/jbc.271.9.4632.

[113] R. Kranz, R. Lill, B. Goldman, G. Bonnard, and S. Merchant, "Molecular mechanisms
of cytochrome c biogenesis: three distinct systems," *Mol Microbiol*, vol. 29, no. 2, pp.
383–396, 1998, doi: 10.1046/j.1365-2958.1998.00869.x.

[114] K. Billet, J. Genitoni, M. Bozec, D. Renault, and D. Barloy, "Aquatic and terrestrial
morphotypes of the aquatic invasive plant, Ludwigia grandiflora, show distinct
morphological and metabolomic responses," *Ecol Evol*, vol. 8, no. 5, pp. 2568–2579,
2018, doi: 10.1002/ece3.3848.

[115] S.-H. Liu, P. C. Hoch, M. Diazgranados, P. H. Raven, and J. C. Barber, "Multi-locus
phylogeny of Ludwigia (Onagraceae): insights on infra-generic relationships and the
current classification of the genus," *Taxon*, vol. 66, no. 5, pp. 1112–1127, 2017.

[116] P. Maheswari, C. Kunhikannan, and R. Yasodha, "Chloroplast genome analysis of
Angiosperms and phylogenetic relationships among Lamiaceae members with particular
reference to teak (Tectona grandis L.f)," *J Biosci*, vol. 46, 2021, [Online]. Available:
https://www.ncbi.nlm.nih.gov/pubmed/34047286

[117] Y. Zhang *et al.*, "The Complete Chloroplast Genome Sequences of Five Epimedium
Species: Lights into Phylogenetic and Taxonomic Analyses," *Front Plant Sci*, vol. 7, p.
306, 2016, doi: 10.3389/fpls.2016.00306.

[118] L. S. Huang *et al.*, "Development of high transferability cpSSR markers for individual
identification and genetic investigation in Cupressaceae species," *Ecol Evol*, vol. 8, no.
10, pp. 4967–4977, 2018, doi: 10.1002/ece3.4053.

[119] P. Leontaritou, F. N. Lamari, V. Papasotiropoulos, and G. Iatrou, "Exploration of
genetic, morphological and essential oil variation reveals tools for the authentication and
breeding of Salvia pomifera subsp. calycina (Sm.) Hayek," *Phytochemistry*, vol. 191, p.
112900, 2021, doi: 10.1016/j.phytochem.2021.112900.

[120] M. Snoussi, L. Riahi, M. Ben Romdhane, A. Mliki, and N. Zoghlami, "Chloroplast DNA Diversity of Tunisian Barley Landraces as Revealed by cpSSRs Molecular Markers and Implication for Conservation Strategies," *Genet Res (Camb)*, vol. 2022, p. 3905957, 2022, doi: 10.1155/2022/3905957.

[121] S. L. Song, P. E. Lim, S. M. Phang, W. W. Lee, D. D. Hong, and A. Prathep, "Development of chloroplast simple sequence repeats (cpSSRs) for the intraspecific study of Graciliaria tenuistipitata (Gracilariales, Rhodophyta) from different populations," *BMC Res Notes*, vol. 7, p. 77, 2014, doi: 10.1186/1756-0500-7-77.

[122] G. L. Wheeler, H. E. Dorman, A. Buchanan, L. Challagundla, and L. E. Wallace, "A review of the prevalence, utility, and caveats of using chloroplast simple sequence repeats for studies of plant biology," *Appl Plant Sci*, vol. 2, no. 12, 2014, doi: 10.3732/apps.1400059.

[123] P. H. Raven and W. Tai, "Observations of Chromosomes in Ludwigia (Onagraceae)," 1979. [Online]. Available: https://about.jstor.org/terms

[124] S. H. Liu, K. H. Hung, T. W. Hsu, P. C. Hoch, C. I. Peng, and T. Y. Chiang, "New insights into polyploid evolution and dynamic nature of Ludwigia section Isnardia (Onagraceae)," *Bot Stud*, vol. 64, no. 1, Dec. 2023, doi: 10.1186/s40529-023-00387-8.

[125] J. L. Feng *et al.*, "Comparison Analysis Based on Complete Chloroplast Genomes and Insights into Plastid Phylogenomic of Four Iris Species," *Biomed Res Int*, vol. 2022, 2022, doi: 10.1155/2022/2194021.