Sequencing, de novo assembly of *Ludwigia* plastomes, and comparative analysis within

the Onagraceae family

Barloy-Hubler F.[3], Le Gac A.-L.[1], Boury C.[2], Guichoux E.[2], and Barloy D.[1*]

1-DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, IFREMER, INRAE,

Rennes, France.

2- Université de Bordeaux, INRAE, BIOGECO, Cestas, France

3- Université de Rennes 1, CNRS, UMR 6553 ECOBIO, Rennes, France.

* Corresponding author: dominique.barloy@agrocampus-ouest.fr

Abstract

The Onagraceae family, which belongs to the order Myrtales, consists of approximately 657 species and 17 genera, including the genus *Ludwigia* L., which is comprised of 82 species. There are few genomic resources for Onagraceae, which limits phylogenetic and population genetics, as well as genomic studies. In this study, new complete plastid genomes of *Ludwigia grandiflora subps. hexapetala* (*Lgh*) and *Ludwigia peploides subsp montevidensis* (*Lpm*) were generated using a combination of different sequencing technologies. These plastomes were then compared to the published *Ludwigia octovalvis* (*Lo*) plastid genome, which was re-annotated by the authors. We initially sequenced and assembled the chloroplast (cp) genomes of *Lpm* and *Lgh* using a hybrid strategy. We observed the existence of two *Lgh* haplotypes and two potential *Lpm* haplotypes. *Lgh*, *Lpm,* and *Lo* plastomes were similar in terms of genome size, gene number, structure, and inverted repeat (IR) boundaries, comparable to other species in the Myrtales order. A total of 45 to 65 SSRs (simple sequence repeats), were detected, depending on the species, with the majority consisting solely of A and T, which is common among angiosperms. Four chloroplast genes (*matK*, *accD*, *ycf2* and *ccsA)* were found under positive selection pressure, which is commonly associated with plant development, and especially in aquatic plants such as *Lgh,* and *Lpm.* Our hybrid sequencing approach revealed the presence of two *Lgh* plastome haplotypes which will help to advance phylogenetic and evolutionary studies, not only specifically for Ludwigia, but also the Onagraceae family and Myrtales order. To enhance the robustness of our findings, a larger dataset of chloroplast genomes would be beneficial.

Keywords

Water primrose, Ludwigia, Onagraceae, chloroplast genome, long and short reads, hybrid assembly, haplotype

Introduction

The Onagraceae family belongs to the order Myrtales which includes approximately 657 species of herbs, shrubs, and trees across 17 genera grouped into two subfamilies: subfam. Ludwigioideae W. L. Wagner and Hoch, which only has one genus (*Ludwigia* L.), and subfam. Onagroideae which contains six tribes and 21 genera [1]. *Ludwigia* L. is composed of 83 species [2, 3]. The current classification for Ludwigia L., which are composed of several hybrid and/or polyploid species, lists 23 sections. A recent molecular analysis is clarified and supported several major relationships in the genus but has challenged the complex sectional classification of Ludwigia L. [4].

The diploid species *Ludwigia peploides* (Kunth) Raven subsp. *montevidensis* (Spreng.) Raven (1963) (named here *Lpm*) (2n=16), and the decaploid species, *Ludwigia grandiflora* (Michx.) Greuter & Burdet (1987) subsp. *hexapetala* (Hook. & Arn.) (Nesom & Kartesz 2000) (named here *Lgh*) (2n=80), reproduce essentially by clonal propagation, which suggests that there is a low genetic diversity within the species (Dandelot et al, 2005). *Lgh* and *Lpm* are native to South America and are described as aquatic invasive plants in Europe [5]. In France, both species occupied aquatic habitats, such as static or slow-flowing waters, riversides, and have recently been observed in wet meadows [6]. The transition from an aquatic to a terrestrial habitat has led to the emergence of two *Lgh* morphotypes [7]. The appearance of metabolic and morphological adaptations could explain the ability to acclimatize to terrestrial conditions, and this phenotypic plasticity involves various genomic and epigenetic modifications [8].

Adequate genomic resources are necessary in order to be able to analyze the different (epi)genomic mechanisms of this acclimation. However, even though the number of terrestrial plant genomes has increased considerably over the last 20 years, only a small fraction (~ 0.16%) have been sequenced, with some clades being significantly more represented than others [9]. Thus, for the Onagraceae family (which includes *Ludwigia* sp.), only a handful of chloroplast

sequences (plastomes) are available, and the complete genome has not yet been sequenced. As

of April 2023, there are 10,712 reference plastomes (RefSeq) listed on GenBank, with the vast

majority (10,392 genomes) belonging to Viridiplantae (green plants). However, the number of

plastomes available for the Onagraceae family is limited, with only 36 plastomes currently

listed. Among these, 15 plastomes are from the tribe Epilobieae, with 11 in the Epilobium genus

and 4 in the Chamaenerion genus. Additionally, there are 23 plastomes from the tribe Onagreae,

with 17 in the Oenothera genus, 5 in the Circaea genus, and only one in the Ludwigia genus.

The *Ludwigia octovalvis* chloroplast genome was released in 2016 as a unique haplotype of

approximately 159 kb (Liu et al., 2016). *L. octovalvis* belongs to sect. *Macrocarpon* (Micheli)

H.Hara of the genus Ludwigia while *Lpm* and *Lgh* belong to jussieae section [10],[11].

Generally, the inheritance of chloroplast genomes is considered to be maternal in angiosperms.

However, biparentally inherited chloroplast genomes could potentially exist in approximately

20% of angiosperm species [12, 13] . Both maternal and biparental inheritance are described in

the Onagraceae family. In tribe Onagreae, *Oenothera* subsect. *Oenothera* are known to have

biparental plastid inheritance [1, 14]. In tribe Epilobieae, biparental plastid inheritance was also

reported in *Epilobium* L. with mainly maternal transmission, and very low proportions of

paternally transmitted chloroplasts [15].

The chloroplast is the symbolic organelle of plants and plays a fundamental role in

photosynthesis. Chloroplasts evolved from cyanobacteria through endosymbiosis and thereby

inherited components of photosynthesis reactions (photosystems, electron transfer and ATP

synthase) and gene expression systems (transcription and translation) [16]. In general,

chloroplast genomes (plastomes) are highly conserved in size, structure, and genetic content.

They are rather small (120-170 kb, [17]), with a quadripartite structure comprising two long

identical inverted repeats (IR, 10–30 kb) separated by large and a small single copy regions

(LSC and SSC, respectively). They are also rich in genes, with around 100 unique genes

encoding key proteins involved in photosynthesis, and a comprehensive set of ribosomal RNAs

(rRNAs) and transfer RNAs (tRNAs) [18]. Plastomes are generally circular but linear shapes

also exist [19]. Chloroplast DNA usually represents 5-20% of total DNA extracted from young

leaves and therefore low-coverage whole genome sequencing can generate enough data to

assemble an entire chloroplast genome [20].

If we refer to their GenBank records, more than 95% of these plastomes were sequenced by

so-called short read techniques (mostly Illumina). However, in most seed plants, the plastid

genome exhibits two large inverted repeat regions (60 to 335 kb, [20]), which are longer than

the short read lengths (< 300 bp). This leads to incomplete or approximate assemblies [21].

Recent long-read sequencing (> 1000 bp) provides strong evidence that terrestrial plant

plastomes have two structural haplotypes, present in equal proportions and differing in IR

orientation of the [22]. This shows the importance of using the so-called third generation

sequence (TGS, PacBio or Nanopore) to correctly assemble the IRs of chloroplasts and to

identify any different structural haplotypes. The current problem with PacBio or Nanopore long

read sequencing is the higher error rate compared to short read technology [23-25]. Thus, a

hybrid strategy which combines long reads (to access the genomic structure) and short reads (to

correct sequencing errors) could be effective [21, 26].

Here, we report the newly sequenced complete plastid genomes of *Ludwigia grandiflora*

subps. *hexapetala* (*Lgh*) and *Ludwigia peploides* subsp *montevidensis* (*Lpm*), using a

combination of different sequencing technologies, as well as a re-annotated comparative

genomic analysis of the published *Ludwigia octovalvis* (*Lo*) plastid. The main objectives of this

study are (1) to assemble and annotate the plastomes of two new species of Ludwigia sp., (2)

to reveal the divergent sequence hotspots of the plastomes in this genus and in the Onagraceae

(3) to identify the genes under positive selection.

To achieve this, we utilized long read sequencing data from Oxford Nanopore and short read

sequencing data from Illumina to assemble the *Lgh* plastomes and compared these assemblies

with those obtained solely from long reads of *Lpm*. We also compared both plastomes to the

published plastome of *Lo*. Our findings demonstrated the value of *de novo* assembly in reducing

assembly errors and enabling accurate reconstruction of full heteroplasmy. We also evaluated

the performance of a variety of software for sequence assembly and correction in order to define

a workflow that will be used in the future to assemble *Ludwigia* sp. mitochlondrial and nuclear

genomes. Finally, the three new Ludwigia plastomes generated by our study make it possible

to extend the phylogenetic study of the Onagraceae family and to compare it with previously

published analyses [4, 27, 28].

Material and Methods

Plant sampling and experimental design

The original plant materials were collected in June of 2018 near to Nantes (France) and

formal identified by D. Barloy. *L. grandiflora* subsp. *hexapetala (Lgh)* plants were taken from

the Mazerolles swamps (N47 23.260, W1 28.206), and *L. peploides* subsp. *montevidensis* (*Lpm*)

plants from La Musse (N 47.240926, W -1.788688)*)*. Plants were cultivated in a growth

chamber in a mixture of $^1/_3$ soil, $^1/_3$ sand, $^1/_3$ loam with flush water level, at 22°C and a 16 h/8

h (light/dark) cycle. A single stem of 10 cm for each species was used for vegetative

propagation in order to avoid potential genetic diversity. *De novo* shoots, taken three

centimeters from the apex, were sampled for each species. Samples for gDNA extraction were

pooled and immediately snap-frozen in liquid nitrogen, then lyophilized over 48 h using a

Cosmos 20K freeze-dryer (Cryotec, Saint-Gély-du-Fesc, France) and stored at room

temperature. All the plants were destroyed after being used as required by French authorities

for invasive plants (article 3, prefectorial decree n°2018/SEE/2423).

Due to high polysaccharide content and polyphenols in *Lpm* and *Lgh* tissues, genomic DNA

extraction was carried out using a modified version of the protocol proposed by Panova et al in

2016, with three purification steps [29].

40 mg of lyophilized buds were ground at 30 Hz for 60 s (Retsch MM200 mixer mill,

FISHER). The ground tissues were lysed with 1 ml CF lysis buffer (MACHEREY-NAGEL)

supplemented with 20 µl RNase and incubated for 1 h at 65°C under agitation. 20 µl proteinase

K was then added before another incubation for 1 h at 65°C under agitation. To avoid breaking

the DNA during pipetting, the extracted DNA was recovered using a Phase-lock gel tube as

described in Belser [30]. The extracts were transferred to 2 ml tubes containing phase-lock gel,

and an equal volume of PCIA (Phenol, Chloroform, Isoamyl Alcohol; 25:24:1) was added.

After shaking for 5 min, tubes were centrifuged at 11000 g for 20 min. The aqueous phase was

transferred into a new tube containing phase-lock gel and extraction with PCIA was repeated.

DNA was then precipitated after addition of an equal volume of binding buffer C4

(MACHEREY-NAGEL) and 99% ethanol overnight at 4°C or 1 h in ice then centrifuged at 800

rpm for 10 min. After removal of the supernatant, 1 ml of CQW buffer was added then the

pellet of DNA was re-suspended. Next, DNA purification was carried out by adding a 2 ml

mixture of wash buffer PW2 (MACHEREY-NAGEL), wash buffer B5 (MACHEREY-

NAGEL), and ethanol at 99% in equal volumes, followed by centrifugation at 800 rpm for 10

min. This DNA purification step was carried out twice. Finally, the DNA pellet was dried in

the oven at 70°C for 30 min then re-suspended in 100 µl elution buffer BE (MACHEREY-

NAGEL) (5 mM Tris solution, pH 8.5) after 10 min incubation at 65°C under agitation.

A second purification step was performed using a PCR product extraction from gel agarose

kit from Macherey-Nagel (MN) NucleoSpin® Gel and PCR Clean-up kit and restarting the

above protocol from the step with the addition of CQW buffer then PW2 buffer.

The third purification step consisted of DNA purification using a Macherey-Nagel (MN)

NucleoMag kit for clean-up and size selection. Finally, the DNA was resuspended after a 5 min

incubation at 65°C in 5 mM TRIS at pH 8.5.

The quantity and quality of the gDNA was verified using a NanoDrop spectrometer,

electrophoresis on agarose gel and ethidium bromide staining under UV light and Fragment

Analyzer (Agilent Technologies) of the University of Rennes1.

Library preparation and sequencing

MiSeq (Illumina) and GridION (Oxford Nanopore Technologies, referred to here as

ONT) sequencing were performed at the PGTB (doi:10.15454/1.5572396583599417E12). *Lgh*

and *Lpm* genomic DNA were re-purified using homemade SPRI beads (1.8X ratio). For

Illumina sequencing, 200 ng of *Lgh* DNA was used according to the QIAseq FX DNA Library

Kit protocol (Qiagen). The final library was checked on TapeStation D5000 screentape (Agilent

Technologies) and quantified using a QIAseq Library Quant Assay Kit (Qiagen). The pool was

sequenced on an Illumina MiSeq using V3 chemistry and 600 cycles (2x300bp). For ONT

sequencing, around 8 μg of *Lgh* and *Lpm* DNA were size selected using a Circulomics SRE kit

(according to the manufacturer's instructions) before library preparation using a SQK-LSK109

ligation sequencing kit following ONT recommendations. Basecalling in High Accuracy -

Guppy version: 4.0.11 (MinKNOW GridION release 20.06.9) was performed for the 48 h of

sequencing. Long reads (LR) and short reads (SR) were available for *Lgh* and only LR for *Lpm*.

Chloroplast assemblies

Quality controls and preprocessing of sequences were performed using Guppy v4.0.14 base

call software for ONT and fastp v0.20.0 [31] using the default quality filter (phred quality >=

Q15) for Illumina reads. After filtering and trimming, *Lgh* Illumina-reads (SR, 2*23,067,490

reads) were used for reference-guided assembly using GetOrganelle v1.7.0 [32] and

NOVOPlasty v4.2.1 [33] using the parameters recommended for green plants. As we sequenced

total DNA, chloroplast reads are mixed with nuclear and mitochondrial reads. Chloroplastic

reads were then extracted by mapping against draft *Lgh* plastomes generated by GetOrganelle

in order to reduce computational costs. Chloroplastic R1 and R2 reads were used with and

without prior error correction using ONT reads with BayesHammer [34] for *de novo* assemblies

using ABySS (version 2.1.5 [35, 36]), MEGAHIT (1.1.2, [37]), Velvet (version 1.2.10) using

VelvetOptimiser and SPAdes (version 3.15.4,[38]). The best k-mer parameters were tested

using kmergenie [39] and Velvetoptimizer. The best results were obtained with k=99, and was

therefore the parameter used for all MiSeq assemblers. For ONT reads, *Lgh* (550,516 reads)

and *Lpm* (68,907 reads) LR reads were self-corrected using CANU 1.8 [40] and *de novo*

assembly using CANU and FLYE 2.8.2 run with the option --meta and –plasmids [41]. Hybrid

correction and assemblies of LR reads using SR data were realized for *Lgh* using Ratatosk [42].

Post plastome assembly validation

As we used many assemblers and different strategies, we produced multiple contigs that

needed to be analyzed and filtered in order to retain only the most robust plastomes. For that,

all assemblies were evaluated using the QUality ASsessment Tool (QUAST) for quality

assessment (http://cab.cc.spbu.ru/quast/) and visualized using BANDAGE [43]. BANDAGE

compatible graphs (.gfa format) were created with the megahit_toolkit for MEGAHIT and with

gfatools for ABySS. Overlaps between fragments were manually checked and ambiguous

"IUPAC or N" nucleotides were also biocured with Illumina reads when available.

Chloroplast genome annotation

Plastomes were annotated via the GeSeq [44] using ARAGORN and tRNAscan_SE to

predict tRNAs and rRNAs and tRNAscan_SE to predict tRNAs and rRNAs and via Chloe

prediction site (https://chloe.plantenergy.edu.au). The previously reported *Lo* chloroplast

genome was also similarly re-annotated to facilitate genomic comparisons. Gene boundaries,

alternative splice isoforms, pseudogenes and gene names and functions were manually checked

and biocurated using Geneious (v.10). Finally, plastomes were represented using

OrganellarGenomeDRAW (OGDRAW) [45]. These genomes were submitted to GenBank at

the National Center of Biotechnology Information (NCBI) with specific accession numbers (for

Lgh haplotype 1, (LGH1) OR166254 and *Lgh* haplotype 2, (LGH2) OR166255; for *Lpm*

haplotype, (LPM) OR166256) using annotation tables generated through GB2sequin [46].

SSRs and Repeat Sequences Analysis

Simple Sequence Repeats (SSRs) were analyzed through the MISA web server [47], with

parameters set to 10, 5, 4, 3, 3, and 3 for mono-, di-, tri-, tetra-, penta-, and hexa-nucleotides,

respectively. Direct, reverse and palindromic repeats were identified using RepEx. Parameters

used were: for inverted repeats (min size 15 nt, spacer = local, class = exact); for palindromes

(min size 20 nt); for direct repeats (minimum size 30 nt, minimum repeat similarity 97%).

Tandem repeats were identified using Tandem Repeats Finder [48], with parameters set to two

for the alignment parameter match and seven for mismatches and indels. The IRa region was

removed for all these analyses to avoid over representation of the repeats.

Comparative chloroplast genomic analyses

Lgh and *Lpm* plastomes were compared with the reannotated and biocurated *Lo* plastome

using mVISTA program [49], with the LAGAN alignment algorithm [50] and a cut-off of 70%

identity.

Nucleotide diversity (Pi) was analyzed using the software DnaSP v.6.12.01 [51, 52] with

step size set to 200 bp and window length to 300 bp. IRscope [53] was used for the analyses of

inverted repeat (IR) region contraction and expansion at the junctions of chloroplast genomes.

To assess the impact of environmental pressures on the evolution of these three Ludwigia

species, we calculated the nonsynonymous (Ka) and synonymous (Ks) substitutions and their

ratios (ω = Ks/Ks) using TBtools [54] to measure the selective pressure. Genes with $\omega < 1$, ω =

1, and $1 < \omega$ were considered to be under purifying selection (negative selection), neutral

selection, and positive selection, respectively.

Plastome phylogenomic analyses

We reconstructed phylogenetic relationships among plastomes of Onagraceae. The FFT-

NS-2 method in MAFFT 7 [55] was used to align all plastomes with one of the IRs removed to

avoid data duplication. Phylogenetic tree analysis, based on the maximum likelihood (ML)

method, was conducted in RAxML 8.2.9 [56] using the GTR+G model with node support

assessed by fast-bootstrap (-f a) using 1,000 non-parametric bootstrap pseudo-replicates.

Graphic representation

Statistical analyses were performed using R software in RStudio integrated development

environment (R Core Team, 2015, RStudio: Integrated Development for R. RStudio, Inc.,

Boston, MA, http://www.rstudio.com/). Figures were realized using ggplot2, ggpubr, tidyverse,

dplyr, gridExtra, reshape2, and viridis packages. SNPs were represented using trackViewer [53]

and genes represented using gggenes packages.

Results

Plastome short read assembly

We first used GetOrganelle and Novoplasty with the short reads of *Lgh*. GetOrganelle

reconstructed two identical haplotypes that differ only by a "flip-flop" of the SSC region, as

shown in Figure 1, while NOVOplasty generated three contigs that can be reassembled to form

either one of the two haplotypes, as the SSC region can be oriented in both directions.

In order to evaluate the best strategies for assembly of *Lgh* plastomes, we used these

haplotypes to map and extract short (Illumina) and long (Nanopore) chloroplast reads.

Chloroplastic short reads, with or without prior correction using long reads, were assembled

using ABySS, Velvet, MEGAHIT and SPAdes. As shown in Figure 2A, the number of contigs

obtained is highly variable depending on the tools and strategies. Without correction, Velvet

and MEGAHIT generate highly fragmented assemblies with 1103 and 722 contigs respectively,

while ABySS and SPAdes produce a more reasonable number of contigs (65 and 12

respectively). In addition, all contigs produced by all assemblers except SPAdes, are

predominantly less than 500 nt long (Figure 2B from 68 to 95% of the contigs). By correcting

the reads before assembly, we saw a noticeable improvement in the number of contigs, which

decreased for all assemblers while increasing their sizes. However, except for SPAdes, the

number of contigs below 500 nt remained preponderant (from 54 to 70%, Figure 2C).

An assessment of the assembly quality using QUAST, and based on an alignment on *Lgh*

haplotype 1 generated by GetOrganelle as a reference, shows that none of the assemblers

produces a complete plastome. By comparing the contribution of small contigs (less than 200,

500 and 1000 nt), we noticed an invariant coverage rate for SPAdes (85%), while the fraction

of covered genome decreases drastically for Velvet (from 92% to 22%), correlated with a

duplication rate going from 2 to 1 (Add. Figure 1A). We noted that compared to an assembly

with all reads, the use of reads superior to 1 kb slightly decreases the coverage rate to reach an

optimal duplication ratio equal to 1. The effect of prior read correction is notable for MEGAHIT

and Velvet, especially concerning the increase in the size of the large alignment (Add. Figure

1A), loss of misassemblies, and reduction of the number of mismatches (Add. Figure 1B).

Investigating all possible scaffolds generated by BANDAGE visualization (Add. Figure 2)

shows that ABySS and SPAdes assembly graphs suggest a tripartite structure with the long

single-copy (LSC) region as the larger circle in the graph (blue), joined to the small single-copy

region (green) by one copy of the inverted repeats (IRs, red). In all graphs, the two IRs are a

collapsed repeat of approximately twice the coverage. For Velvet and MEGAHIT, graphs

confirm the significant fragmentation of the assemblies, which is improved by prior correction

of the reads.

In conclusion, SPAdes is the most efficient tool, and gives the best result using corrected short

reads to assemble a resolved plastome, with three contigs of 90,272 bp (corresponding to LSC),

19,788 bp (corresponding to SSC) and 24,762 bp (corresponding to a copy of the IR).

Plastome long read assembly

Chloroplast fractions of *Lgh* long reads were selected through mapping. Raw, self-

corrected reads were assembled using CANU, and short-read-corrected comparative assembly

was carried out using CANU or FLYE. Using raw data, CANU generates a unique contig

corresponding to haplotype 2, whereas FLYE makes two contigs that reconstruct haplotype 1.

Assemblies using long reads corrected by CANU leads to fragmentation into two (CANU

assembler) or three (FLYE assembler) contigs which both reconstruct haplotype 1, with an large

gap corresponding to one of the IR copies for the CANU assembler. Finally, correction by

RATATOSK allows CANU to assemble two redundant contigs, with contig 1 reproducing the

entire haplotype 2, while FLYE makes two contigs corresponding to haplotype 1 (Add. Figure

3A). In conclusion, the two haplotypes proposed by GetOrganelle using the short reads were

likewise reconstructed randomly using the long reads, with or without correction.

Consequently, both haplotypes were retained as *Lgh* plastomes and submitted to Genbank.

Hybrid assembly is the most complete and the most faithful, with an average of 1.6 contigs (all

assemblers combined), and optimal correction using RATATOSK, which permits plastomes

with 99.94% accuracy compared to biocurated molecules (Add. Figure 3B).

Unfortunately, due to the absence of short read data, we could only perform self-corrected

long read assembly for *Lpm* using CANU. We also compared CANU and FLYE assembler

efficiency, and found that assembly using CANU produces 13 contigs whereas FLYE produces

12 contigs. In both cases, only three contigs are required to reconstitute a complete cpDNA

assembly (no gap, no N), with an SSC region oriented like those of the *Lgh* haplotype 2 and the

Lo plastome. Although it is more than likely that these two SSC region orientations also exist for *Lpm*, the low number of nanopore sequences generated (68907 reads) and absence of Illumina short reads prevented us from demonstrating the existence of both haplotypes. As a result, only the "haplotype 2" generated sequence was deposited to Genbank.

Annotation and comparison of *Ludwigia* plastomes

1. General Variations

Plastomes of the three species of Ludwigia sp*., Lgh*, *Lpm* and *Lo,* are circular double-stranded DNA molecules (Figure 3) which are all (as shown in Table 1) approximately the same size: *Lo* being slightly smallest and *Lgh* the biggest. The overall GC content is almost the same for the three species and the GC contents of the IR regions are higher than those of the LSC and SSC regions. Between the three species, the lengths of the total chloroplasts, LSC, SSC, and IR are broadly similar (Table 2) and the three plastomes are perfectly syntenic if we orient the SSC fragments the same way.

All three Ludwigia sp. plastomes contain the same number of functional genes (134 in total) encoding 85 proteins (embracing 7 duplicated in the IR region: *ndhB*, *rpl2*, *rpl23*, *rps7*, *rps12*, *ycf2*, *ycf15*), 37 tRNAs (including trnK-UUU which contains *matK*), and 8 rRNAs (16S, 23S, 5S, and 4.5S as duplicated sets in the IR). Among these genes, 18 contain introns, of which six are tRNAs (Table 2). Only the *rps12* gene is a trans-spliced gene. A total of 46 genes are involved in photosynthesis, and 71 genes related to transcription and translation, including a bacterial-like RNA polymerase and 70S ribosome, as well as a full set of transfer RNAs (tRNAs) and ribosomal RNAs (rRNAs). Six other protein-coding genes are involved in essential functions, such as *accD,* which encodes the β-carboxyl transferase subunit of acetyl-CoA carboxylase, an important enzyme for fatty acid synthesis; *matK* encodes for maturase K, which is involved in the splicing of group II introns; *cemA*, a protein located in the membrane envelope of the chloroplast is involved in the extrusion of protons and therby indirectly allows

the absorption of inorganic CO2 in the plastids; *clpP1* which is involved in proteolysis, and;

ycf1, ycf2, two ATPases members of the TIC translocon. Finally, a highly pseudogenized *ycf15*

locus was annotated in the IR even though premature stop codons indicate loss of functionality.

2.   Segments Contractions/Expansion

The junctions between the different chloroplast segments were compared between three

Ludwigia sp. (*Lpm*, *Lgh* and *Lo),* and we found that the overall resemblance of Ludwigia sp.

plastomes was confirmed at all junctions (Figure 4A). In all three genomes, *rpl22, rps19, and*

rpl2 were located around the LSC/IRb border, and *rpl2*, *trnH*, and *psbA* were located at the

IRa/LSC edge. The JSB (junction between IRb and SSC) is either located in the *ndhF* gene or

the *ycf1* gene depending on the orientation of the SSC region (Figure 4B). The *ycf1* gene was

initially annotated as a 1139 nt pseudogene that we biocurate as a larger gene (5302 nt) with a

frameshift due to a base deletion, compared to *Lg* and *Lo* which both carry a complete *ycf1*

gene.

If we compare Ludwigia sp. junctions with those of other Onagraceae plastomes (Figure

5), we can observe that the JLB and JLA connections are well-preserved in the whole family,

whereas JSB and JLA differ. Concerning JSB, in the five Onagraceae family genera with

available plastomes, *ndhF* is duplicated, with the exception of Circaea sp. and Ludwigia sp. For

Oenothera villosa, the first copy of *ndhF*, which is located in the IRb, overlaps the JSB border,

whereas for *Oenothera lindheimeri*, *Epibolium amurense* and Chamaenerion sp*.*, *ndhF* is only

located in inverted repeats. Only Circaea sp. and Ludwigia sp. have a unique copy of this locus,

and it is found in the SSC segment (Figure 5). At the JSA border, in Circaea sp., the *ycf1* gene

crosses the IRa/SSC boundary and extends into the IRa region.

When comparing the respective sizes of chloroplast fragments (IR/SSC/LSC) in

Onagraceae, it can be observed that *Ludwigia* species exhibit expansions in the SSC and LSC

regions which are not compensated by significant contractions in the IR regions. This is likely

due to the relocation of the *ndhF* in the SSC region and *rps19* in the LSC region. Additionally,

there may be significant size variations in the intergenic region between *trnI* and *ycf2*, as well

as the intergenic segment containing the *ycf15* pseudogene (Add. Figure 4).

3. Repeats and SSRs analysis

In this study, we analyzed the nature and distribution of single sequence repeats (SSR), as

their polymorphism is an interesting indicator in phylogenetic analyses. A total of 65 (*Lgh*), 48

(*Lpm*) and 45 (*Lo*) SSRs were detected, the majority being single nucleotide repeats (38–21),

followed by tetranucleotides (12–10) and then di-, tri- and penta-nucleotides (Add. Figure 5A).

Mononucleotide SSRs are exclusively composed of A and T, indicating a bias towards the use

of the A/T bases, which is confirmed for all SSRs (Add. Figure 5B). In addition, the SSRs are

mainly distributed in the LSC region for the three species, which is probably biased by the fact

that LSC is the longest segment of the plastome (Add. Figure 5C). The analysis of SRR

locations revealed that most were distributed in non-coding regions (intergenic regions and

introns, Add. Figure 5D).

The chloroplast genomes of the three *Ludwigia* species were also screened for long repeat

sequences. They were counted in a non-redundant way (if smaller repetitions were included in

large repeats, only the large ones were considered). Four types of repeats (tandem, palindromic

inverted and direct) were surveyed in the three *Ludwigia* sp. plastomes. No inverted repeats

were detected with the criteria used.

For the three other types of repeats, here are their distributions:

Tandem repeats **(Table 3A)**: Perfect tandem repeats (TRs) with more than 15 bp were

examined. Twenty-two *loci* were identified in the three *Ludwigia* sp. plastomes (*Lgh*, *Lpm*, *Lo*),

heterogeneously distributed as shown in Table 3A: 13 loci (plus one imperfect) in *Lo*, nine loci

(plus one imperfect) in *Lgh* and seven loci (plus two imperfect) in *Lpm*. It can therefore be seen

that the TR distributions (occurrence and location) are specific to each plastome, since only

four pairs are common to the three species. Thus, nine TRs are unique to *Lo*, three to *Lpm* and

three to *Lgh.* Two pairs are common to *Lgh* and *Lpm* and one is common to *Lo* and *Lgh.* TRs

are mainly intergenic or intronic but are detected in two genes (*accD* and *ycf1)*. These genes

have accelerated substitution rates, although this does not generate a large difference in their

lengths. This point will be developed later in this article.

Direct repeats **(Table 3B)**: There are few direct (non-tandem) repeats (DRs) in the

chloroplast genomes of *Ludwigia* sp. A single direct repeat of 41 nt is common to the three

species, at 2 kb intervals, in *psaB* and *psaA* genes. This DR corresponds to an amino acid repeat

[WLTDIAHHHLAIA] which corresponds to a region predicted as transmembrane. We then

observe three direct repeats conserved in *Lpm* and *Lgh* in *ycf1*, *accD* and *clpP1* respectively,

two unique DRs in *Lo* (in the *accD* gene and *rps12-clpP1* intergene) and one in *Lgh* (in the

clpP1 intron 1 and *clpP1* intron 2).

Palindromes **(Table 3C)**: Palindromic repeats make up the majority of long repetitions,

with the numbers of perfect repeats varying from 19, 24 and 26 in *Lo*, *Lgh* and *Lpm*,

respectively, and the number of quasi-palindromes (1 mutation) varying between 8, 3 and 6.

They are mainly found in the intronic and intergenic regions, with the exception of six genic

locations in *psbD*, *ndhK*, *ccsA* and *rpl22,* and two palindromic sequences in *ycf2.* These gene

palindromic repeats do not seem to cause genetic polymorphism in *Ludwigia* and can be

considered as silent.

Thirteen palindromes are common to the three species (including 2 with co-variations in

Lo). 13 others present in *Lpm* and *Lgh* correspond to quasi-palindromes (QPs) in *Lo* due to

mutated bases, and conversely, three *Lo* perfect palidromes are mutated in *Lpm* and *Lgh*.

Finally, only five palindromes are species specific. Two in particular are located in the

hypervariable intergenic spacer *ndhF-rpl32*, and are absent in *Lo* due to a large deletion of 160

nt.

4. Repeat distribution in LSC, SSC and IR segments

In the IRa/IRb regions, repeats are only identified in the first 9 kb region between *rpl2* and *ycf2*: a tandem repeat in the *Lpm rpl2* intron, and a tetranucleotide repeat, [TATC]*3, located in the *ycf2* gene in the 3 species. In *ycf2* we also found 1 common palindrome (16 nt), a single palindrome in *Lo* (20 nt, absent following an A:G mutation in the 2 other species), as well as a shared tandem repeat (24 nt), and an additional 15 nt tandem repeat in *Lo* which adds 4 amino acids to protein sequence.

In the SSC region, the repeats are almost all located in the intergenic and/or intronic regions, with a hotspot between *ndhF* and *ccsA*. There is also a shared microsatellite in *ndhF,* and a palidrome (16 nt) in *ccsA* which is absent in *Lo* (due to an A:C mutation), resulting in a synonymous mutation (from isoleucine to leucine). We also observed multiple and various repeats in the *ycf1* gene: 3 common poly-A repeats (from 10 to 13 nt), 3 species-specific microsatellites (ATAG)*3 and (ACCA)*4 in *Lgh* and (CAAC)*3 in *Lo,* as well as two direct repeats of 32 nt (37 nt spacing), which were absent from *Lo* due to a G:T SNP. Two tandem repeats were also observed in *Lo* and *Lgh*. Neither of these repeats are at the origin of the frameshift causing the pseudogenization of *ycf1* in *Lo*, this latter being due to a single deletion of an A at position 3444 of the gene.

Finally, in the LSC region, the longest segment, which consequently contains the maximum number of repeats, we still observed a preferential localization in the intergenic and intronic regions since only genes *atpA*, *rpoC2*, *rpoB*, *psbD*, *psbA*, *psbB*, *ndhK* and *clpP1* contain either mononucleotic repeats (poly A and T), palindromes, or microsatellites (most often common to the three species and without affecting the sequences of the proteins produced). As mentioned earlier, the only exception is the *accD* gene, which contains several direct and tandem repeats in *Lgh* and *Lpm,* corresponding to a region of 174 nt (58 amino acids) missing in *Lo* and, conversely, a direct repeat of 40 nucleotides, in a region of 147 nt (49 aa), which is present in

Lo and missing in the other two species. These tandem repeats lead to the presence of four

copies of 9 amino acids [DESENSNEE] in *Lgh* and *Lpm*, two of which form a larger duplication

of 17 aa [FLSDSDIDDESENSNEE]. Similarly, the TRs present only in *Lo* generate two perfect

9 amino acid repeats [EELSEDGEE], included in two longer degenerate repeats of 27 nt (Add.

Figure 6). It should be noted that though these TRs do not disturb the open reading phases, it is

still possible for them to form an intron which is not translated. Different functional studies will

be necessary to clarify this point. The presence of polymorphisms of the *accD* gene between

Lo and the two species (*Lpm*, *Lgh*) is interesting because *accD*, that encodes a subunit of acetyl-

CoA carboxylase (EC 6.4.1.2). This enzyme is essential in fatty acid synthesis and also

catalyzes the synthesis of malonyl-CoA, which is necessary for the growth of dicots, plant

fitness and leaf longevity, and is involved in the adaptation to specific ecological niches [57].

Large *accD* expansions due to TRs have also been described in other plants such as *Medicago*

[58] and *Cupressophytes* [59]. Some authors have suggested that these inserted repeats are not

important for acetyl-CoA carboxylase activity as the reading frame is always preserved, and

they assume that these repeats must have a regulatory role [60].

5. Sequence Divergence Analysis and Polymorphic Loci Identification

Determination of divergent regions by MVista, using *Lo* as a reference, confirmed that the

three Ludwigia sp. plastomes are well preserved if the SSC segment is oriented in the same way

(Add. Figure 7). Sliding window analysis (Figure 6) indicated variations in definite coding

regions, notably *clpP*, *accD*, *ndh5*, *ycf1* with high Pi values, and to a lesser extent, *rps16*, *matK*,

ndhK, *petA*, *ccsA* and four tRNAs (*trnH*,*trnD*, *trnT* and *trnN*). These polymorphic *loci* could be

suitable for inferring genetic diversities in Ludwigia sp.

A comparative analysis of the sizes of protein coding genes sizes also shows that the *rps11*

gene initially annotated in *Lo* is shorter than those which have been newly annotated in *Lgh* and

Lpm (345 bp instead of 417 bp). Comparative analysis by BLAST shows that it is the long form

which is annotated in other Myrtales, and the observation of the locus in *Lo* shows a frameshift

mutation (deletion of a nucleotide in position 311). Functional analysis would be necessary to

check whether the *rps11* frameshift mutation produces shorter proteins that have lost their

function. And only obtaining the complete genome will verify whether copies of some of these

genes have been transferred to mitochondrial or nuclear genomes. Such *rps11* horizontal

transfers have been reported for this gene in the mitochondrial genomes of various plant

families [61]. This also applies to *ycf1*, found as a pseudogene in *Lo* (as specified previously),

although it is not known if this reflects a gene transfer or a complete loss of function [62,

63].Moreover, there is a deletion of nine nucleotides in the 3' region of the *rpl32* gene in *Lgh*

and *Lpm*, leading to a premature end of the translation and the deletion of the last 4 amino acids

[QRLD], which are replaced by a K. However, if we look carefully at the preserved region as

defined by the RPL32 domain (CHL00152, member of the superfamily CL09115), we see that

the later amino acids are not important for *rpl32* function since they are not found in the

orthologs.

Our results show that the Ka/Ks ratio is less than 1 for most genes (Figure 7). This indicates

adaptive pressures to maintain the protein sequence except for *matK* (1.17 between *Lgh* and

Lpm), *accD* (2.48 between *Lgh* and *Lo* and 2.16 between *Lpm* and *Lo*), *ycf2* (4.3 between both

Lgh-Lp and *Lo*) and *ccsA* (1.4 between both *Lgh-Lpm* and *Lo*), showing a positive selection for

these genes, and a possible key role in the processes of the species' ecological adaptations. As

we have already described the variability in the *accD* sequence, we will focus on *ycf2*, *matK,*

and *ccsA* variations.

Concerning *ccsA*, the variations observed, although significant, concern only five amino

acids, and modifications do not seem to affect the C-type cytochrome synthase gene function.

Concerning *ycf2*, our analysis shows that this gene is highly polymorphic with 256 SNPs

that provoke 10 deletions, 7 insertions, 21 conservative and 49 non-conservative substitutions

in *Lo* (Add. Figure 8), compared to *Lgh* and *Lpm* (100 % identical). This gene has been shown

as "variant" in other plant species such as *Helianthus tuberosus* [64].

The *matK* gene has been used as a universal barcoding locus to enable species

discrimination of terrestrial plants [65], and is often, together with the *rbcL* gene, the only

known genetic resource for many plants. Thus, we propose a phylogenetic tree from *Ludwigia*

matK sequences (Figure 8). It should however be noted that this tree contains only 149 amino

acids common to all the sequences (out of the 499 in the complete protein). As only three

complete *Ludwigia* plastomes are available, we cannot specify whether these barcodes are

faithful to the phylogenomic history of *Ludwigia* in the same way as the complete plastome. In

any case, for this tree, we can see that *Lo* stands apart from the other Ludwigia sp., *Lpm* and

Lgh, and that the *L. grandiflora* subsp. *hexapetala* belongs to the same branch as the species *L.*

ovalis (aquatic taxon used in aquariums [66]), *L. stolonifera* (native to the Nile, found in a

variety of habitats, from freshwater wetlands to brackish and marine waters) [67] and *L.*

adscendens (common weed of rice fields in Asia) [68]. *Lpm* is in a branch unique to its species

but close to the *L. grandiflora* subsp. *hexapetala* branch.

Discussion

In the present study, we first sequenced and *de novo* assembled the chloroplast (cp) genomes

of *Ludwigia peploides* (*Lpm*) and *Ludwigia grandiflora* (*Lgh*), two species belonging to the

Onagraceae family. We employed a hybrid strategy and demonstrated the presence of two cp

haplotypes in *Lgh* and one haplotype in *Lpm*, although the presence of both haplotypes in *Lpm*

is likely. Furthermore, we compared these genomes with those of other species in the

Onagraceae family to expand our knowledge of genome organization and molecular evolution

in these species.

Our findings demonstrate that the utilization of solely short reads has failed to produce complete

plastomes, likely due to challenges posed by long repeats and rearrangements. On the other

hand, relying solely on long reads resulted in a lower quality sequence due to insufficient

coverage and sequencing errors. After conducting our research, we discovered that hybrid

assembly, which incorporates both long and short read sequences, resulted in the most superior

complete assemblies. This innovative approach capitalizes on the advantages of both

sequencing technologies, harnessing the accuracy of short read sequences and the length of long

read sequences. Our findings corroborate with similar results obtained in studies on other

chloroplasts, such as those in *Eucalyptus* [21], *Falcataria* [69], *Carex* [70] or *Cypripedium* [71].

As a result of this strategy, we were able to successfully identify the presence of two haplotypes

in *Lgh*, which is a first for *Ludwigia,* as the plastome of *L. octovalvis* was only delivered in the

form of one haplotype [72]. Due to the unavailability of sequence data for *Ludwigia octovalvis*

and our exclusive use of long reads for *Ludwigia peploides*, we are unable to conclusively

identify the presence of these two forms in the *Ludwigia* genus. However, we believe that they

are likely to be present. Unfortunately, the current representation of plastomes in GenBank

primarily consists of short-read data, which may result in an underrepresentation of this

polymorphism. It is unfortunate that structural heteroplasmy, which is expected to be

widespread in angiosperms, has been overlooked. Notably, the existence of two plastome

haplotypes has been widely identified in the closely related order of Myrtales (Eucalyptus sp.),

and more commonly in Angiosperms, with 58 species across 16 other orders showing similar

patterns [22]. Recent studies have also revealed the presence of two cp haplotypes in other

orders, such as *Asparagales* (*Ophrys apifera* orchid, [73]), Brassicales (*Carica papaya*,

Vasconcellea pubescens, [74]), Solanales (*Solanum tuberosum*, [75]), and even in a new order,

Laurales (Avocado *Persea americana*, [76]). Furthermore, Wanichthanarak *et al*. (2023)

revisited 24 Ramanaceae published chloroplast genomes and identified a likely form of

heteroplasmy in *Rhamnus crenata* [77]. These emerging findings highlight the potential underestimation of the occurrence of two chloroplast haplotypes in angiosperms and emphasize the importance of identifying and considering such heteroplasmy during *de novo* or revisited chloroplast genome assemblies.

The chloroplast genome sizes for the three genera of Onagraceae subfam. Onagroideae varied as follows: Circaea sp. ranged from 155,817 bp to 156,024 bp, Chamaenerion sp. ranged from 159,496 bp to 160,416 bp, and Epilobium sp. ranged from 160,748 bp to 161,144 bp [78]. Our study revealed that the size of the complete chloroplast of *Ludwigia* (Onagraceae subfamily Ludwigioideae) ranged from 159,369 bp to 159,584 bp, which is remarkably similar to other Onagraceae plants (average length of 162,030 bp). Furthermore, *Ludwigia* plastome sizes are consistent with the range observed in Myrtales (between 152,214 to 171,315 bp, [79]). In the same way, similar overall GC content was found in *Ludwigia sp.* (from 37.3 to 37.4%), *Circaea sp.* (37.7 to 37.8%), *Chamaenerion sp.* and *Epilobium sp.* (38.1 to 38.2%, [78]) and more generally for the order Myrtales (36.9–38.9%, with the average GC content being 37%, [79]). Higher GC content of the IR regions (43.5%) found in *Ludwigia* sp. has already been shown in the Myrtales order (39.7–43.5%) and in other families/orders such as Amaranthaceae (order Caryophyllales, [80]) or Lamiaceae (order Lamiales, [81]), and is mainly due to the presence of the four GC rich rRNA genes.

The complete chloroplast genomes of the three Ludwigia species encoded an identical set of 134 genes including 85 protein-coding genes, 37 tRNA genes and eight ribosomal RNAs, consistent with gene content found in the Myrtales order, with a gene number varying from 123 to 133 genes with 77–81 protein-coding genes, 29–31 tRNA gene and four rRNA genes [79]. Chloroplast genes have been selected during evolution due to their functional importance [82]. In our current study, we made the noteworthy discovery that *matK*, *accD*, *ycf2*, and *ccsA* genes were subjected to positive selection pressure. These genes have frequently been reported in

literature as being associated with positive selection, and are known to play crucial roles in

plant development conditions. *Lgh* and *Lpm* are known to thrive in aquatic environments, where

they grow alongside rooted emergent aquatic plants, with their leaves and stems partially

submerged during growth, as reported by Wagner et al. in 2007 [1]. Both species possess the

unique ability of vegetative reproduction, enabling them to establish themselves rapidly in

diverse habitats, including terrestrial habitats, as noted by Haury et al [7]. Additionally, *Lo* is a

wetland plant that typically grows in gullies and at the edges of ponds, as documented by

Wagner *et al.* in 2007 [1]. Given their ability to adapt to different habitats, these species may

have evolved specialized mechanisms to cope with various abiotic stresses, such as reduced

carbon and oxygen availability or limited access to light in submerged or emergent conditions.

Concerning *matK*, Barthet et al [83] demonstrated the relationship between light and

developmental stages, and MatK maturase activity, suggesting important functions in plant

physiology. This gene has recently been largely reported to be under positive selection in an

aquatic plant (*Anubias* sp., [84]), and more generally in terrestrial plants (*Pinus sp*, [85] or

Chrysosplenium sp. [86]). The *accD* gene has been described as an essential gene required for

leaf development [87] and longevity [88] in tobacco (*Nicotiana tabacum*). Under drought stress,

plant resistance can be increased by inhibiting *accD* [89], and conversely, enhanced in response

to flooding stress by upregulating *accD* accumulation [90]. Hence, we can hypothesize that the

positive selection observed on the *accD* gene can be explained by the submerged and emerged

constraints undergone by *Ludwigia* species. The *ycf2* gene seems to be subject to adaptive

evolution in *Ludwigia species*. Its function, although still vague, would be to contribute to a

protein complex generating ATP for the TIC machinery (proteins importing into the

chloroplasts [91, 92]), as well as plant cell survival [93, 94]. The *ccsA* gene positive selection

is found in some aquatic plants such as *Anubia sp.* [84], marine flowering plants as

Zostera species [95], and some species of Lythraceae [89]. The *ccsA* gene is required for

cytochrome c biogenesis [96] and this hemoprotein plays a key role in aerobic and anaerobic

respiration, as well as photosynthesis [97]. Furthermore, we showed that *Lgh* colonization is

supported by metabolic adjustments mobilizing glycolysis and fermentation pathways in

terrestrial habitats, and the aminoacyl-tRNA biosynthesis pathway, which are key components

of protein synthesis in aquatic habitats [8] It can be assumed that the ability of *Ludwigia* to

invade aquatic and wet environments, where the amount of oxygen and light can be variable,

leads to a high selective pressure on genes involved in respiration and photosynthesis.

Molecular markers are often used to establish population genetic relationships through

phylogenetic studies. Five chloroplasts (*rps16*, *rpl16*, trnL-trnF, trnL-CD, *trnG*) and two

nuclear markers (ITS, *waxy*) were used in previous phylogeny studies of *Ludwigia sp.* [4].

However, no SSR markers had previously been made available for the *Ludwigia* genus, or more

broadly, the Onagraceae. In this study, we identified 45 to 65 SSR markers depending on the

Ludwigia species. Most of them were AT mononucleotides, as already recorded for other

angiosperms [98, 99]. In addition, we identified various genes with highly mutated regions that

can also be used as SNP markers. Chloroplast SSRs (cpSSRs) represent potentially useful

markers showing high levels of intraspecific variability due to the non-recombinant and

uniparental inheritance of the plastomes [100, 101]. Chloroplast SSR characteristics for

Ludwigia sp. (location, type of SSR) were similar to those described in most plants. While the

usual molecular markers used for phylogenetic analysis are nuclear DNA markers, cpSSRs have

also been used to explore cytoplasmic diversity in many studies [102-104]. To conclude, the 13

highly variable loci and cpSSRs identified in this study are potential markers for population

genetics or phylogenetic studies of *Ludwigia* species, and more generally, Onagraceae.

Conclusion

In this study, we conducted the first-time sequencing and assembly of the complete plastomes of *Lpm* and *Lgh*, which are the only available genomic resources for functional analysis in both species. We were able to identify the existence of two haplotypes in both *Lpm* and *Lgh*, while the absence of the *Lo* genome precluded further investigation for this species. Comparison of all 10 Onagraceae plastomes revealed a high degree of conservation in genome size, gene number, structure, and IR boundaries. However, to further elucidate the phylogenetic analysis and evolution in Ludwigia and Onagraceae, additional chloroplast genomes will be necessary, as highlighted in recent studies of Iris and Aristidoideae species [105].

Declarations

- Availability of data and materials

The datasets generated and/or analysed during the current study were available in GenBank (for *Lgh* haplotype 1, (LGH1) OR166254 and *Lgh* haplotype 2, (LGH2) OR166255; for *Lpm* haplotype, (LPM) OR166256).

- Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article

- Funding

The post-doctoral research grant of Anne-Laure Le Gac was supported by the Conseil regional Bretagne (SAD18001).

- Acknowledgements

We are grateful to Luis Portillo-Lemus for developing the high molecular weight genomic DNA extraction protocol. All sequencing experiments were performed at the PGTB (doi:10.15454/1.5572396583599417E12).

Figure 1: Two structural haplotypes of *L. grandiflora* plastomes representing the flip-flop organization of SSC segment.

Figure 2: Comparative results of *L. grandiflora* short read (SR) assemblies. **A:** Total number of contigs obtained with the uncorrected (dark green) and corrected (light green) chloroplast SRs for the 4 assemblers (ABySS, MEGAHIT, Velvet and SPAdes). **B:** Comparison of the size of contigs assembled by the 4 tools using corrected or uncorrected SRs. **C:** Boxplot showing the distribution of these contigs by size and the improvement brought by the prior correction of the SRs with the long reads for each tool.

Figure 3: Circular representation of annotations plastomes in *Ludwigia octovalis*, *Ludwigia grandiflora* and *Ludwigia peploides* using ogdraw. Each card contains four circles. From the center outwards, the first circle shows forward and reverse repeats (red and green arcs,

653   respectively). The next circle shows tandem repeats as bars. The third circle shows the

654   microsatellite sequences. Finally, the fourth and fifth circles show the genes colored according

655   to their functional categories (see colored legend). Only the haplotype 1 of *L. grandiflora* is

656   represented as haplotype 2 only diverge by the orientation of the SSC segment.

Figure 4: Comparison of the borders of LSC, SSC, and IR regions in Onograceae plastomes.
A: Comparison of the junction between large single-copy (LSC, light blue), inverted repeat (IR, orange) and short single-copy (SSC, light green) regions among the chloroplast genomes of *L. octovalvis*, *L. peploides* and *L. grandiflora* (both haplotypes). Genes are denoted by colored boxes and the gaps between genes and boundaries are indicated by base lengths (bp). JLB: junction line between LSC and IRb; JSB: junction line between IRb and SSC; JSA: junction line between SSC and IRa; JLA: junction line between IRa and LSC. **B:** Comparison of SSC boundaries in haplotype 1 (*L. peploides* and *L. grandiflora* haplotype 1) and haplotype 2 (*L. octovalvis* and *L. grandiflora* haplotype 2) plastomes.

Figure 5: Comparison of LSC, SSC and IR regions boundaries in Onograceae chloroplast genomes. Representative sequences from each genus have been chosen (noted R on the diagram) except for *Oenothera lindheimeri* (only 89.35 % identity with others *Oenothera*), *Circaea alpina* (99.5 % identity but all others *Circaea* are 99.9% identical) and *Chamaenerion conspersum* (99% but all others *Chamaenerion* are ca. 99.7 identical). As shown in Figure 7, the 3 *Ludwigia* plastomas had the same structure, *L. octovalvis* was chosen as a representative of this genus.

Figure 6: Illustration of nucleotide diversity of the three *Ludwigia* chloroplast genome sequences. The graph was generated using DnaSP software version 6.0 (windows length: 800 bp, step size: 200 bp). The x-axis corresponds to the base sequence of the alignment, and the y-axis represents the nucleotide diversity (π value). LSC, SSC and IR segments were indicated under the line representing the genes coding the proteins (in light blue) the tRNAs (in pink) and the rRNAs (in red). The genes marking diversity hotspots are noted at the top of the peaks.

687
688 **Figure 7:** The Ka/Ks ratios of the 80 protein-coding genes of *Ludwigia* plastomes. The blue

689 curve represents *L. grandiflora* versus *L. peploides*, purple curve denotes *L. grandiflora* versus

690 *L. octovalvis* and green curve *L. peploides* versus *L. octovalvis*. Four genes (*matK*, *accD*, *ycf2*

691 and *ccsA*) have Ka/Ks ratios greater than 1.0, whereas the Ka/Ks ratios of the other genes were

692 less than 1.0.

Figure 8: Phylogenetic tree based on the *Ludwigia matK* marker. Only four sequences are complete (499 aa, yellow star), the others correspond to barcode amplification ranging from 128 to 386 aa, with an average of 244 aa.

Table 1. The general characteristics of the 3 Ludwigia plastomes

	L.octovalvis*	L.grandiflora	L.peploides
Size (bp)			
	159;396	159;584	159;537
LSC	90;183	90;272	90;156
SSC	19;703	19;788	19;799
IR	24;755	24;762	24;791
GC%			
	37;4	37;3	37;3
LSC	35;2	35;1	35;1
SSC	32	31;7	31;7
IR	43;5	43;5	43;4

* KX827312 (ref)

699
700

Table 2 : Genes present in the plastomes of *Ludwigia*

FUNCTION	NAME
Photosynthesis	
Rubisco	rbcL
Photosystem I (PSI)	psaA; psaB; psaC; psaI; psaJ
PSI assembly factors	ycf3# (pafI); ycf4 (pafII)
Photosystem II	psbA; psbB; psbC; psbD; psbE; psbF; psbH; psbI; psbJ; psbK; psbL; psbM; pbf1 (psbN) psbT; psbZ
ATP synthase	atpA; atpB; atpE; atpF#; atpH; atpI
Cytochrome *b6f*	petA; petB#; petD#; petG; petL; petN
Cytochrome biogenesis	ccsA
NADPH dehydrogenase	ndhA#; ndhB**#; ndhC; ndhD; ndhE; ndhF; ndhG; ndhH; ndhI; ndhJ
Transcription and translation	
Transcription	rpoA; rpoB; rpoC1#; rpoC2
Small ribosomal proteins	rps2; rps3; rps4; rps7**; rps8; rps11; rps12**#; rps14; rps15; rps16#; rps18; rps19
Large ribosomal proteins	rpl2**#; rpl14; rpl16#; rpl20; rpl22; rpl23**; rpl32; rpl33; rpl36
Translation initiation	infA
Ribosomal RNA	rrn5**; rrn4;5**; rrn16**; rrn23**
Transfer RNA	trnA-UGC**#;trnC-GCA;trnD-GUC;trnE-UUC;trnF-GAA;trnfM-CAU;trnG-GCC;trnG-UCC#;trnH-GUG;;trnI-CAU**;trnI-GAU**#;trnK-UUU#;trnL-CAA**;trnL-UAA#;trnL-UAG;trnM-CAU;trnN-GUU**;trnP-UGG;trnQ-UUG;trnR-ACG**;trnR-UCU;trnS-GCU;trnS-GGA;trnS-UGA;trnT-GGU;trnT-UGU;trnV-GAC**;trnV-UAC#;trnW-CCA;trnY-GUA
Other functions	
Group II intron splicing	matK
Inorganic carbon uptake	cemA
Protease	clpP1#
Fatty acid synthesis/Heat tolerance	accD
TIC machinery (protein import)	ycf1 (Tic214); ycf2**
Unknown function	ycf15**
	** duplicated in IR region; # spliced genes

Table 3:

Table 3A : Tandem repeats

Sequence	L octovalis (Lo)	L grandiflora (Lg)	L peploides (Lp)	Length	Region	Locus	Comments
TTGTAGTCAGGGGTGTAGTACTAT				24	IRs	ycf2	
TAGAAGAGAGTGCAG		X	X	15	IRs	ycf2	15 nt deletion in L.g and L.p
ATGAAATATCGTATAATGAAGTACCACACGAGTGGATAT	X	X	X	39	IRs	rpl2 intron	39 nt deletion in L.g and L.o
AAAAATAGGATAGGAT		X	X	16	LSC	ycf1-trnH-GUG	56 nt deletion in L.g and L.p
TAAATTAATATCTATATA		X	X	18	LSC	psbZ-trnG-GCC	18 nt deletion in L.g and L.p
TTTTCTATCTATCTTATATCAA		X	X	22	LSC	trnK-UUU-rps16	22 nt deletion in L.g and L.p
AGATCCATAACATCATCAAA		X	X	20	LSC	rps16 intron	22 nt deletion in L.g and L.p
TATTAGTTATTAATATTATTAGA		X	X	23	LSC	trnP-UGG-psaJ	23 nt deletion in L.g and L.p
AATAATATATAATAACTTAAATA		X	X	23	LSC	rpl33-rps18	33 et 44 nt deletion in in L.g et L.p, respectively
TTTTTATTTAACATGCTATCAAATCAACAATGCCATACCGTAGGGCATCTGTT		X	X	53	LSC	rpl20-clpP1	107 nt deletion in L.g and L.p
ATATATTTCGATTCAATTC	X		X	19	LSC	trnH-GUG-psbA	3 copies in a 57 nt deletion in L.o and L.p
ATAGAAATATCAGTATTTGAGTG	X		X	23	LSC	atpH-atpI	23 nt deletion in L.o and L.p
TTAATTTTAATTGAAGAA	X		X	18	LSC	psbJ-psbL	17 and 24 nt deletion in L.o and L.p, respectively
TTAAAGAATATTAATATTC	imperfect TR			19	LSC	trnR-UCU-atpA	A -> C mutation in second copy in L.o
TATTTATTATTTAAT	X	X		16	LSC	atpH-atpI	16 nt deletion in L.g and L.o
TCTAAGGGCTGAAATAAGG	X	X		18	LSC	pafI intron	18 nt deletion in L.g and L.o
TGTGAATCTATCTAT			X	15	LSC	trnS-UGA-psbZ	8 nt deletion in L.p
TTTTTTCTAGTA				12	LSC	pafI intron	
CTAGTTATTGACATGG		imperfect TR	imperfect TR	16	LSC	psaJ-rpl33	G -> A mutation in second in L.p et L.g
ATTTTTATTTAACTCT	X	X	imperfect TR	15	SSC	ycf1	T->A mutation in first copy in L.p, other sequence in first copy in L.o
AATCAAATAGTTGAT		X	X	15	SSC	ycf1	other sequence in first copy of L.p and L.g
ATAATAATATATTTATTATTAATTAATA	X			28	SSC	ndhF-rpl32	160 nt deletion in L.o

Table 3B : Direct repeats

Sequence	L. octovalis (Lo)	L. grandiflora (Lg)	L. peploides (Lp)	Size (nt)	Spacers (nt)	Region	Locus	Comments
TTCAATTGGAACGGACGATTCGTCAATCATCT				32	37	SSC	ycf1	2 copies. In L.o, one mutation (G->A) in the second copie
CATCGATGATGAAAGTGAAAACAGTAATGAAGAGG	X			35	28 - 22 - 11	LSC	accD	3 perfects copies and 1 mutated (G->A) copie in L.g and L.p. Region of 174 nt deleted in L.o
AGATGGTGAAGAACCTTATGAAGATGGTGAAGAACCTTATG		X	X	41	22	LSC	accD	Region of 147 nt deleted in L.g and L.p
TATCAAATCAACAATGCCATACCGTAGGGCAT		X	X	32	22 - 21	LSC	rps12-clpP1	3 copies
TTAAGAGCCGTACAGGCACCTTTTGATGCATACGG	X				408 in L.p, 406 in L.g	LSC	clpP1	2 copies. In L.g, one mutation (C->T) in the second copie
TTAAGAGCCGTACAGGCACTTTTTGATGCATACGG	X		X	35	811	LSC	clpP1 intron 1-intron 2	
TGCAATAGCCAAATGATGATGAGCAATATCAGTCAGCCATA				41	2178		psaB & psaA	

Table 3C : Palindromic repeats

			Locus	
Common perfect palindromic repeats				
AGACTCTCATGAGAGTCT			trnC-GCA - petN	
ATTAAATAGAATATTCTATTTAAT			trnE-UUC-trnT-GGU	
TTGGTAAATTTACCAA			psbD	
TTCATTTCAATTTCAATTGAAATTGAAATGAA			trnI-CAU-ycf2	
GAAAAAGGCCTTTTTC			ycf2	2 copies in IR
TCTCAAATGATTAATCATTTGAGA			trnL-UAA intron	
GGATTACTAGTAATCC			trnD-GUC-trnY-GUA	
TTTGAATGCATTCAAA			trnG-UCC intron	
ATATATTCGAATATAT			trnG-UCC -trnR-UCU	
TAGTAATTAATTACTA			trnG-GCC-trnfM-CAU	
CCAGTATGCATACTGG			ndhK	
Common palindromic repeats with covariation				
in L octovalvis	in L grandiflora et L pepoides			
ATAGAATCTATATTCTATTAGAATATAGATTCTAT	ATCGAATCTATATTCTATTAGAATATAGATTCGAT		ndhC-trnV-UAC	
ATGTATATATATCGAT	ATCTATATATATAGAT		trnE-UUC-trnT-GGU	
Common palindromic and quasi-palindromic repeats				
in L octovalvis	in L grandiflora and L pepoides			
TTTAACGAATATTAATATT t GTTAAA	TTTAACGAATATTAATATTCGTTAAA		trnR-UCU-atpA	
TTAA c GAATATTAATATTCTTTAA	TTAAAGAATATTAATATTCTTTAA		trnR-UCU-atpA	
AATTGTA c TTACAATT	AATTGTAATTACAATT		ccsA	
AGGAAGATTGATCAATCTT t CT	AGGAAGATTGATCAATCTTCCT		trnL-UAG-rpl32	
TTA c TAATATTACTAA	TTAGTAATATTACTAA		trnK-UUU intron	
ATATAGAATAT c CTATAT	ATATAGAATATTCTATAT		psbZ-trnG-GCC	
ACATATCATGATA g GT	ACATATCATGATATGT		rpl22	
AATTACTAATTTCTATTACTATGTTCAATTGAACATAGTAATAGAAATTAGTAATT	AATTACTAATTTCTATTACT t TGTTCAATTGAACATAGTAATAGAAATTAGTAATT		atpH-atpI	
TAGTTAGAATTCTAACTA	TAGTT c GAATTCTAACTA		trnT-UGU-trnL-UAA	
TATTTTTTCTGAAAAAATA	TATTTTTTCTAGAA g AAATA		ycf2	2 copies in IR
in L octovalvis and L pepoides	in L grandiflora			
CCCATCAATCATCGATTG t TGGG	CCCATCAATCATCGATTGATGGG		psbN-trnD-GUC	
in L octovalvis and L grandiflora	in L pepoides			
ATGAAAAAAATCGATTTTTTTCAT	ATGATAAAAATAGATTTTT a TCAT		trnK-UUU-rps16	
ATGA A AAAAATCGATTTTTT t TCAT- ATGATAAAAATCGATTTTTATCAT	ATGATAAAAATA g ATTTTTATCAT		trnK-UUU-rps16	
Unique palindromic repeats				
L pepoides				
TTATATATATATATATAA			rpl32-ndhF	Full deletion in L. octovalvis , 6 bases deletion in L. grandiflora
L octovalvis				
ATTGAAATTCGAATTTCAAT			psbZ-trnG-GCC	Full deletion in L. grandiflora and L. peploides
L peploides and L. grandiflora				
AAAAAATGGATCCATTTTTT			trnL-UAG-rpl32	3 bases deleted and 3 bases mutated in L. octovalvis
AATATATATTATTATAATAATATATATT			rpl32-ndhF	Full deletion in L. octovalvis
TATATTTATTTATTAATTAATAATAAATAATA			rpl32-ndhF	Full deletion in L. octovalvis

Lo = Ludwigia octovalvis; Lgh = L. grandiflora subsp. *hexapetala; Lpm = L. peploides* subsp. *montevidensis.*

A

	ABySS		MEGAHIT		VELVET		SPAdes	
	not corrected	corrected	not corrected	corrected	not corrected	corrected	not corrected	corrected
Using all contigs								
Genome fraction (%)	86.868	85.279	86.428	85.158	91.927	86.796	84.682	84.483
Duplication ratio	1.047	1.042	1.796	1.041	2.002	1.128	1.042	1
Largest alignment	56 588	41 262	30 904	90 352	3531	17 235	90 399	90 272
Using contigs > 200 nt								
Genome fraction (%)	86.419	85.279	86.377	85.057	76.589	86.181	84.682	84.483
Duplication ratio	1.028	1.042	1.681	1.029	1.177	1.11	1.042	1
Largest alignment	56 588	41 262	30 904	90 352	3531	17 235	90 399	90 272
Using contigs > 500 nt								
Genome fraction (%)	85.564	84.517	83.503	84.774	45.468	79.279	84.682	84.483
Duplication ratio	1.009	1.012	1.041	1.004	1.015	1.054	1.042	1
Largest alignment	56 588	41 262	30 904	90 352	3531	17 235	90 399	90 272
Using contigs > 1000 nt								
Genome fraction (%)	83.701	84.199	81.256	84.545	22.194	66.438	84.563	84.483
Duplication ratio	1	1.002	1.007	1.001	1	1.011	1.026	1
Largest alignment	56 588	41 262	30 904	90 352	3531	17 235	90 399	90 272

B

	ABySS		MEGAHIT		VELVET		SPAdes	
	not corrected	corrected	not corrected	corrected	not corrected	corrected	not corrected	corrected
Using all contigs								
NGA50	15 215	26 577	19 986	90 352	469	2796	90 399	90 272
LGA50	3	3	3	1	93	9	1	1
Misassemblies								
# misassemblies	0	0	4	0	0	0	0	0
Misassembled contigs length	0	0	1595	0	0	0	0	0
Mismatches								
# mismatches per 100 kbp	109.53	107.19	1036.93	45.24	499.16	229.11	96.57	0
# indels per 100 kbp	12.4	10.58	62.99	16.26	27.92	74.88	19.17	0
# N's per 100 kbp	0	0	0	0	0	6.1	0	0
Using contigs > 500 nt								
NGA50	15 215	26 577	19 986	90 352	–	2796	90 399	90 272
LGA50	3	3	3	1	–	9	1	1
Misassemblies								
# misassemblies	0	0	1	0	0	0	0	0
Misassembled contigs length	0	0	665	0	0	0	0	0
Mismatches								
# mismatches per 100 kbp	62.39	46.17	123.32	8.1	221.33	148.48	96.57	0
# indels per 100 kbp	2.9	2.2	4.33	1.47	28.51	63.74	19.17	0
# N's per 100 kbp	0	0	0	0	0	7.22	0	0
Using contigs > 1000 nt								
NGA50	15 215	26 577	19 986	90 352	–	2796	90 399	90 272
LGA50	3	3	3	1	–	9	1	1
Misassemblies								
# misassemblies	0	0	0	0	0	0	0	0
Misassembled contigs length	0	0	0	0	0	0	0	0
Mismatches								
# mismatches per 100 kbp	0	0	37.51	0.74	64.94	61.6	67.87	0
# indels per 100 kbp	1.5	0.74	0	0.74	25.41	56.93	19.49	0
# N's per 100 kbp	0	0	0	0	0	9.25	0	0

Supp. Figure 1: QUAST evaluation of performance of the four assembly tools (using corrected or uncorrected SRs). **A:** Comparison of plastome fraction, duplication rate and size of the largest alignment obtained. **B:** Comparison of classic metrics (NGA50 and LGA50), number of errors (misassemblies and mismatches) produced.

719
720 **Supp. Figure 2:** BANDAGE visualization of the *L. grandiflora* plastome assembly graphs on corrected or
721 uncorrected SRs. Contigs are colored according to their BLAST match to the LSC (blue), SSC (green), and
722 IR (red) segments
723
724

B

Assembler : CANU	LSC	IR	SSC
No correction	99.793 (187 nt)	99.960 (30 nt)	99.778 (27 nt)
Canu-correction	99.811 (171 nt)	99.899 (25 nt)	99.828 (34 nt)
Ratatosk-correction	99.952 (43 nt)	99.879 (10 nt)	99.864 (27 nt)

Assembler : FLYE	LSC	IR	SSC
No correction	99.798 (182 nt)	99.903 (24 nt)	99.783 (50 nt)
Canu-correction	99.831 (153 nt)	99.907 (23 nt)	99.843 (31 nt)
Ratatosk-correction	99.975(23 nt)	99.988 (3 nt)	99.995 (3 nt)

725
726 **Supp. Figure 3:** Graphs representing the assemblies of *L. grandiflora* long reads. **A:** Contigs are represented
727 in light blue and the three segments (LSC, SSC and IR) in dark blue, green and yellow, respectively. B:
728 Comparative effectiveness of CANU and RATATOSK correctors.
729
730

731
Supp. Figure 4: Comparison of LSC, SSC and IR sizes in the Onagraceae. **A:** Comparison of the sizes of
LSC, SSC and IR segments in the Onogracae family (*Chamaenerion* in blue, *Circaea* in yellow, *Epibolium*
in dark purple, *Ludwigia* in light green and *Oenothera* in dark green). **B:** Maximum likelihood tree made
using RAxML (model GTR-GAMMA, algorithm Rapid Hill-climbing) on multiple sequences alignment of
Onogracae plastomes made using MAFFT. **C:** Average size of the different chloroplast segments (LSC,
SSC and IR) for the 5 genres of Onogracae. IR size corresponds to the sum of the two copies.
738

739
740
741 **Supp. Figure 5:** Comparative analysis of Simple-Sequence Repeats (SSRs) in *Ludwigia* chloroplast
742 genomes. A: SSR numbers detected in the three species, by repeat class types (mono, di-, tri-, tetra and
743 pentanucleotides). **B:** Frequency of SSR motifs by repeat class types. **C:** Frequency of SSRs in LSC, SSC
744 and IR regions. **D:** Repartition of SSRs in intergenic, protein-coding and intronic regions.

745
746

Supp. Figure 6: Diagram showing the position of tandem repeats in the *accD* gene. *L. octovalis* (in red) and *L. peploides* and *L. grandiflora* (in green). We also observe the consequences of these repetitions on the insertion of amino acids, also repeated.

754
Supp. Figure 7: Comparison of the three *Ludwigia* plastomes using mVISTA, with the *L. octovalvis* as a
reference. **A:** The y-axis represents the identity percentage (between 50 and 100%). The arrows show the
genes (in green: proteins genes, in purple: rRNAs and in fuchsia: tRNAs). Blue blocks indicate exonic
regions. LCS, IR and SSC regions are also distinguished (in dark blue, red and green, respectively). The
second line corresponds to *L. grandiflora* haplotype 2 (For this haplotype, SSC segment is oriented like *L.*
octovalvis) and the third line corresponds to *L. peploides* for which the SSC region has been artificially
oriented in the same way as the two other plastomes to allow comparison. **B:** Small box showing a part of

the alignment and presenting the consequences if we do not artificially orient the SSC segments in the same
direction for the analysis.
764

Supp. Figure 8: Lollipop diagram allowing the visualization of SNPs and their translational effects on the *ycf2*. **A:** localization of the 256 single nucleotide polymorphisms (SNP) observed by comparing *L. grandiflora-L. peploides* with *L. octovalvis.* Two regions particularly dense in SNPs (between 3420 and 3460 and between 6100 and 6600) have been zoomed into to allow better reading. **B:** Effect of these SNPs on the translated sequence of *L. octovalvis*, compared to Ycf2 of the other two species: non conservative mutation: red square; conservative mutation: circle green; deletion: triangle_point_up blue and insertion: triangle_point_down, orange. As for A, two regions were zoomed into in order to distinguish each mutation.

References

1. Wagner WL, Hoch PC, Raven PH: **Revised classification of the Onagraceae**. *Systematic Botany Monographs* 2007.

2. Levin RA, Wagner WL, Hoch PC, Nepokroeff M, Pires JC, Zimmer EA, Sytsma KJ: **Family-level relationships of Onagraceae based on chloroplast rbcL and ndhF data**. *American Journal of Botany* 2003, **90**(1):107-115.

3. Levin RA, Wagner WL, Hoch PC, Hahn WJ, Rodriguez A, Baum DA, Katinas L, Zimmer EA, Sytsma KJ: **Paraphyly in tribe Onagreae: insights into phylogenetic relationships of Onagraceae based on nuclear and chloroplast sequence data**. *Systematic Botany* 2004, **29**(1):147-164.

4. Liu S-H, Hoch PC, Diazgranados M, Raven PH, Barber JC: **Multi-locus phylogeny of Ludwigia (Onagraceae): insights on infra-generic relationships and the current classification of the genus**. *Taxon* 2017, **66**(5):1112-1127.

5. Hussner A, Windhaus M, Starfinger U: **From weed biology to successful control: an example of successful management of Ludwigia grandiflora in Germany**. *Weed Research* 2016, **56**(6):434-441.

6. Lambert E, Dutartre A, Coudreuse J, Haury J: **Relationships between the biomass production of invasive Ludwigia species and physical properties of habitats in France**. *Hydrobiologia* 2010, **656**:173-186.

7. Haury J, Druel A, Cabral T, Paulet Y, Bozec M, Coudreuse J: **Which adaptations of some invasive Ludwigia spp.(Rosidae, Onagraceae) populations occur in contrasting hydrological conditions in Western France?** *Hydrobiologia* 2014, **737**:45-56.

8.   Billet K, Genitoni J, Bozec M, Renault D, Barloy D: **Aquatic and terrestrial**

morphotypes of the aquatic invasive plant, Ludwigia grandiflora, show distinct

morphological and metabolomic responses. *Ecol Evol* 2018, **8**(5):2568-2579.

9.   Marks RA, Hotaling S, Frandsen PB, VanBuren R: **Representation and participation**

across 20 years of plant genome sequencing. *Nat Plants* 2021, **7**(12):1571-1578.

10.   Zardini E, Raven PH: **A new section of Ludwigia (Onagraceae) with a key to the**

sections of the genus. *Systematic botany* 1992:481-485.

11.   Hoch PC, Wagner WL, Raven PH: **The correct name for a section of Ludwigia**

L.(Onagraceae). *PhytoKeys* 2015(50):31.

12.   Hu Y, Zhang Q, Rao G: **Occurrence of plastids in the sperm cells of Caprifoliaceae:**

biparental plastid inheritance in angiosperms is unilaterally derived from

maternal inheritance. *Plant and cell physiology* 2008, **49**(6):958-968.

13.   Zhang Q: **Why does biparental plastid inheritance revive in angiosperms?** *Journal*

of plant research 2010, **123**(2):201-206.

14.   Cleland RE: **Oenothera. Cytogenetics and evolution**. 1972.

15.   Schmitz UK, Kowallik K-V: **Plastid inheritance in Epilobium**. *Current genetics* 1986,

11(1):1-5.

16.   Sato N: **Are Cyanobacteria an Ancestor of Chloroplasts or Just One of the Gene**

Donors for Plants and Algae? *Genes (Basel)* 2021, **12**(6).

17.   Gualberto JM, Mileshina D, Wallet C, Niazi AK, Weber-Lotfi F, Dietrich A: **The plant**

mitochondrial genome: dynamics and maintenance. *Biochimie* 2014, **100**:107-120.

18.   Tonti-Filippini J, Nevill PG, Dixon K, Small I: **What can we do with 1000 plastid**

genomes? In*.*, vol. 90: Wiley Online Library; 2017: 808-818.

19.   Oldenburg DJ, Bendich AJ: **The linear plastid chromosomes of maize: terminal**

sequences, structures, and implications for DNA replication. *Current genetics* 2016,

62:431-442.

20.   Twyford AD, Ness RW: **Strategies for complete plastid genome sequencing**.

Molecular ecology resources 2017, **17**(5):858-868.

21.   Wang W, Schalamun M, Morales-Suarez A, Kainer D, Schwessinger B, Lanfear R:

Assembly of chloroplast genomes with long-and short-read data: a comparison of

approaches using Eucalyptus pauciflora as a test case. *BMC genomics* 2018, **19**:1-

15.

22.   Wang W, Lanfear R: **Long-reads reveal that the chloroplast genome exists in two**

distinct versions in most plants. *Genome Biology and Evolution* 2019, **11**(12):3372-

3381.

23.   Ferrarini M, Moretto M, Ward JA, Šurbanovski N, Stevanović V, Giongo L, Viola R,

Cavalieri D, Velasco R, Cestaro A: **An evaluation of the PacBio RS platform for**

sequencing and de novo assembly of a chloroplast genome. *BMC genomics* 2013,

14(1):1-12.

24.   Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, Tyson JR, Beggs AD,

Dilthey AT, Fiddes IT: **Nanopore sequencing and assembly of a human genome with**

ultra-long reads. *Nature biotechnology* 2018, **36**(4):338-345.

25.   Rang FJ, Kloosterman WP, de Ridder J: **From squiggle to basepair: computational**

approaches for improving nanopore sequencing read accuracy. *Genome biology*

2018, **19**(1):90.

26.   Scheunert A, Dorfner M, Lingl T, Oberprieler C: **Can we use it? On the utility of de**

novo and reference-based assembly of Nanopore data for plant plastome

sequencing. *PLoS One* 2020, **15**(3):e0226234.

27. Bedoya AM, Madriñán S: **Evolution of the aquatic habit in Ludwigia (Onagraceae):**

Morpho-anatomical adaptive strategies in the Neotropics. *Aquatic Botany* 2015,

120:352-362.

28. Liu S-H, Yang H-A, Kono Y, Hoch PC, Barber JC, Peng C-I, Chung K-F:

Disentangling Reticulate Evolution of North Temperate Haplostemonous

Ludwigia (Onagraceae) 1, 2. *Annals of the Missouri Botanical Garden* 2020,

105(2):163-182.

29. Panova M, Aronsson H, Cameron RA, Dahl P, Godhe A, Lind U, Ortega-Martinez O,

Pereyra R, Tesson SV, Wrange A-L: **DNA extraction protocols for whole-genome**

sequencing in marine organisms. *Marine genomics: Methods and protocols* 2016:13-

44.

30. Belser C, Istace B, Denis E, Dubarry M, Baurens F-C, Falentin C, Genete M, Berrabah

W, Chèvre A-M, Delourme R: **Chromosome-scale assemblies of plant genomes using**

nanopore long reads and optical maps. *Nature plants* 2018, **4**(11):879-887.

31. Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ preprocessor**.

Bioinformatics 2018, **34**(17):i884-i890.

32. Jin J-J, Yu W-B, Yang J-B, Song Y, DePamphilis CW, Yi T-S, Li D-Z: **GetOrganelle:**

a fast and versatile toolkit for accurate de novo assembly of organelle genomes.

Genome biology 2020, **21**:1-31.

33. Dierckxsens N, Mardulyn P, Smits G: **NOVOPlasty: de novo assembly of organelle**

genomes from whole genome data. *Nucleic acids research* 2017, **45**(4):e18-e18.

34. Nikolenko SI, Korobeynikov AI, Alekseyev MA: **BayesHammer: Bayesian clustering**

for error correction in single-cell sequencing. In: *BMC genomics: 2013*. Springer: 1-

11.

35.   Jackman SD, Vandervalk BP, Mohamadi H, Chu J, Yeo S, Hammond SA, Jahesh G,

870   Khan H, Coombe L, Warren RL: **ABySS 2.0: resource-efficient assembly of large**

871   **genomes using a Bloom filter**. *Genome research* 2017, **27**(5):768-777.

872   36.   Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJ, Birol I: **ABySS: a parallel**

873   **assembler for short read sequence data**. *Genome research* 2009, **19**(6):1117-1123.

874   37.   Li D, Luo R, Liu C-M, Leung C-M, Ting H-F, Sadakane K, Yamashita H, Lam T-W:

875   **MEGAHIT v1. 0: a fast and scalable metagenome assembler driven by advanced**

876   **methodologies and community practices**. *Methods* 2016, **102**:3-11.

877   38.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

878   Nikolenko SI, Pham S, Prjibelski AD: **SPAdes: a new genome assembly algorithm**

879   **and its applications to single-cell sequencing**. *Journal of computational biology* 2012,

880   **19**(5):455-477.

881   39.   Chikhi R, Medvedev P: **Informed and automated k-mer size selection for genome**

882   **assembly**. *Bioinformatics* 2014, **30**(1):31-37.

883   40.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable**

884   **and accurate long-read assembly via adaptive k-mer weighting and repeat**

885   **separation**. *Genome research* 2017, **27**(5):722-736.

886   41.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA: **Assembly of long, error-prone reads**

887   **using repeat graphs**. *Nature biotechnology* 2019, **37**(5):540-546.

888   42.   Holley G, Beyter D, Ingimundardottir H, Møller PL, Kristmundsdottir S, Eggertsson

889   HP, Halldorsson BV: **Ratatosk: hybrid error correction of long reads enables**

890   **accurate variant calling and assembly**. *Genome Biology* 2021, **22**(1):1-22.

891   43.   Wick RR, Schultz MB, Zobel J, Holt KE: **Bandage: interactive visualization of de**

892   **novo genome assemblies**. *Bioinformatics* 2015, **31**(20):3350-3352.

893   44.   Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S:

894   **GeSeq–versatile and accurate annotation of organelle genomes**. *Nucleic acids*

895   *research* 2017, **45**(W1):W6-W11.

896   45.   Greiner S, Lehwark P, Bock R: **OrganellarGenomeDRAW (OGDRAW) version 1.3.**

897   **1: expanded toolkit for the graphical visualization of organellar genomes**. *Nucleic*

898   *acids research* 2019, **47**(W1):W59-W64.

899   46.   Lehwark P, Greiner S: **GB2sequin-A file converter preparing custom GenBank files**

900   **for database submission**. *Genomics* 2019, **111**(4):759-761.

901   47.   Beier S, Thiel T, Münch T, Scholz U, Mascher M: **MISA-web: a web server for**

902   **microsatellite prediction**. *Bioinformatics* 2017, **33**(16):2583-2585.

903   48.   Benson G: **Tandem repeats finder: a program to analyze DNA sequences**. *Nucleic*

904   *acids research* 1999, **27**(2):573-580.

905   49.   Frazer KA, Pachter L, Poliakov A, Rubin EM, Dubchak I: **VISTA: computational**

906   **tools for comparative genomics**. *Nucleic acids research* 2004, **32**(suppl_2):W273-

907   W279.

908   50.   Brudno M, Do CB, Cooper GM, Kim MF, Davydov E, Green ED, Sidow A, Batzoglou

909   S, Program NCS: **LAGAN and Multi-LAGAN: efficient tools for large-scale**

910   **multiple alignment of genomic DNA**. *Genome research* 2003, **13**(4):721-731.

911   51.   Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-

912   Onsins SE, Sánchez-Gracia A: **DnaSP 6: DNA sequence polymorphism analysis of**

913   **large data sets**. *Molecular biology and evolution* 2017, **34**(12):3299-3302.

914   52.   Rozas J, Rozas R: **DnaSP version 3: an integrated program for molecular**

915   **population genetics and molecular evolution analysis**. *Bioinformatics (Oxford,*

916   *England)* 1999, **15**(2):174-175.

53.   Amiryousefi A, Hyvönen J, Poczai P: **IRscope: an online program to visualize the**
junction sites of chloroplast genomes. *Bioinformatics* 2018, **34**(17):3030-3031.

54.   Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, Xia R: **TBtools: an**
integrative toolkit developed for interactive analyses of big biological data.
Molecular plant 2020, **13**(8):1194-1202.

55.   Katoh K, Misawa K, Kuma Ki, Miyata T: **MAFFT: a novel method for rapid multiple**
sequence alignment based on fast Fourier transform. *Nucleic acids research* 2002,
30(14):3059-3066.

56.   Stamatakis A: **The RAxML v8. 2. X Manual**. *Heidleberg Institute for Theoretical*
Studies.

57.   Konishi T, Sasaki Y: **Compartmentalization of two forms of acetyl-CoA carboxylase**
in plants and the origin of their tolerance toward herbicides. *Proc Natl Acad Sci U*
S A 1994, **91**(9):3598-3601.

58.   Wu S, Chen J, Li Y, Liu A, Li A, Yin M, Shrestha N, Liu J, Ren G: **Extensive genomic**
rearrangements mediated by repetitive sequences in plastomes of Medicago and
its relatives. *BMC Plant Biol* 2021, **21**(1):421.

59.   Li J, Su Y, Wang T: **The Repeat Sequences and Elevated Substitution Rates of the**
Chloroplast accD Gene in Cupressophytes. *Front Plant Sci* 2018, **9**:533.

60.   Gurdon C, Maliga P: **Two distinct plastid genome configurations and**
unprecedented intraspecies length variation in the accD coding region in Medicago
truncatula. *DNA Res* 2014, **21**(4):417-427.

61.   Richardson AO, Palmer JD: **Horizontal gene transfer in plants**. *J Exp Bot* 2007,
58(1):1-9.

62.   de Vries J, Sousa FL, Bolter B, Soll J, Gould SB: **YCF1: A Green TIC?** *Plant Cell*
2015, **27**(7):1827-1833.

63.   Filip E, Skuza L: **Horizontal Gene Transfer Involving Chloroplasts**. *Int J Mol Sci*
2021, **22**(9).

64.   Zhong Q, Yang S, Sun X, Wang L, Li Y: **The complete chloroplast genome of the**
Jerusalem artichoke (Helianthus tuberosus L.) and an adaptive evolutionary
analysis of the ycf2 gene. *PeerJ* 2019, **7**:e7596.

65.   Antil S, Abraham JS, Sripoorna S, Maurya S, Dagar J, Makhija S, Bhagat P, Gupta R,
Sood U, Lal R *et al*: **DNA barcoding, an effective tool for species identification: a**
review. *Mol Biol Rep* 2023, **50**(1):761-775.

66.   Li J, Wang Y, Cui J, Wang W, Liu X, Chang Y, Yao D, Cui J: **Removal effects of**
aquatic plants on high-concentration phosphorus in wastewater during summer. *J*
Environ Manage 2022, **324**:116434.

67.   Soliman AT, Hamdy RS, Hamed AB: **Ludwigia stolonifera (Guill. & Perr.) PH**
Raven, insight into its phenotypic plasticity, habitat diversity and associated
species. *Egyptian Journal of Botany* 2018, **58**(3):605-626.

68.   Kamoshita A, Ikeda H, Yamagishi J, Lor B, Ouk M: **Residual effects of cultivation**
methods on weed seed banks and weeds in Cambodia. *Weed Biology and*
Management 2016, **16**(3):93-107.

69.   Anita VPD, Matra DD, Siregar UJ: **Chloroplast genome draft assembly of Falcataria**
moluccana using hybrid sequencing technology. *BMC Res Notes* 2023, **16**(1):31.

70.   Xu S, Teng K, Zhang H, Gao K, Wu J, Duan L, Yue Y, Fan X: **Chloroplast genomes**
of four Carex species: Long repetitive sequences trigger dramatic changes in
chloroplast genome structure. *Front Plant Sci* 2023, **14**:1100876.

71.   Guo YY, Yang JX, Li HK, Zhao HS: **Chloroplast Genomes of Two Species of**
Cypripedium: Expanded Genome Size and Proliferation of AT-Biased Repeat
Sequences. *Front Plant Sci* 2021, **12**:609729.

72. Liu S-H, Edwards C, Hoch PC, Raven PH, Barber JC: **Complete plastome sequence**

of Ludwigia octovalvis (Onagraceae), a globally distributed wetland plant. *Genome*

announcements 2016, **4**(6):e01274-01216.

73. Bateman RM, Rudall PJ, Murphy ARM, Cowan RS, Devey DS, Perez-Escobar OA:

Whole plastomes are not enough: phylogenomic and morphometric exploration at

multiple demographic levels of the bee orchid clade Ophrys sect. Sphegodes. *J Exp*

Bot 2021, **72**(2):654-681.

74. Lin Z, Zhou P, Ma X, Deng Y, Liao Z, Li R, Ming R: **Comparative analysis of**

chloroplast genomes in Vasconcellea pubescens A.DC. and Carica papaya L. *Sci*

Rep 2020, **10**(1):15799.

75. Lihodeevskiy GA, Shanina EP: **The Use of Long-Read Sequencing to Study the**

Phylogenetic Diversity of the Potato Varieties Plastome of the Ural Selection.

Agronomy 2022, **12**(4):846.

76. Nath O, Fletcher SJ, Hayward A, Shaw LM, Masouleh AK, Furtado A, Henry RJ, Mitter

N: **A haplotype resolved chromosomal level avocado genome allows analysis of**

novel avocado genes. *Hortic Res* 2022, **9**:uhac157.

77. Wanichthanarak K, Nookaew I, Pasookhush P, Wongsurawat T, Jenjaroenpun P,

Leeratsuwan N, Wattanachaisaereekul S, Visessanguan W, Sirivatanauksorn Y,

Nuntasaen N: **Revisiting chloroplast genomic landscape and annotation towards**

comparative chloroplast genomes of Rhamnaceae. *BMC Plant Biology* 2023,

23(1):59.

78. Luo Y, He J, Lyu R, Xiao J, Li W, Yao M, Pei L, Cheng J, Li J, Xie L: **Comparative**

Analysis of Complete Chloroplast Genomes of 13 Species in Epilobium, Circaea,

and Chamaenerion and Insights Into Phylogenetic Relationships of Onagraceae.

Front Genet 2021, **12**:730495.

79.   Zhang XF, Landis JB, Wang HX, Zhu ZX, Wang HF: **Comparative analysis of**
chloroplast genome structure and molecular dating in Myrtales. *BMC Plant Biol*
2021, **21**(1):219.

80.   Xu J, Shen X, Liao B, Xu J, Hou D: **Comparing and phylogenetic analysis**
chloroplast genome of three Achyranthes species. *Sci Rep* 2020, **10**(1):10818.

81.   Lian C, Yang H, Lan J, Zhang X, Zhang F, Yang J, Chen S: **Comparative analysis of**
chloroplast genomes reveals phylogenetic relationships and intraspecific variation
in the medicinal plant Isodon rubescens. *PLoS One* 2022, **17**(4):e0266546.

82.   Mohanta TK, Mishra AK, Khan A, Hashem A, Abd Allah EF, Al-Harrasi A: **Gene Loss**
and Evolution of the Plastome. *Genes (Basel)* 2020, **11**(10).

83.   Barthet MM, Hilu KW: **Expression of matK: functional and evolutionary**
implications. *American Journal of Botany* 2007, **94**(8):1402-1412.

84.   Li L, Liu C, Hou K, Liu W: **Comparative Analyses of Plastomes of Four Anubias**
(Araceae) Taxa, Tropical Aquatic Plants Endemic to Africa. *Genes (Basel)* 2022,
13(11).

85.   Zeb U, Wang X, AzizUllah A, Fiaz S, Khan H, Ullah S, Ali H, Shahzad K:
Comparative genome sequence and phylogenetic analysis of chloroplast for
evolutionary relationship among Pinus species. *Saudi J Biol Sci* 2022, **29**(3):1618-
1627.

86.   Wu Z, Liao R, Yang T, Dong X, Lan D, Qin R, Liu H: **Analysis of six chloroplast**
genomes provides insight into the evolution of Chrysosplenium (Saxifragaceae).
BMC Genomics 2020, **21**(1):621.

87.   Kode V, Mudd EA, Iamtham S, Day A: **The tobacco plastid accD gene is essential**
and is required for leaf development. *Plant J* 2005, **44**(2):237-244.

88.   Madoka Y, Tomizawa K, Mizoi J, Nishida I, Nagano Y, Sasaki Y: **Chloroplast**

1017   **transformation with modified accD operon increases acetyl-CoA carboxylase and**

1018   **causes extension of leaf longevity and increase in seed yield in tobacco**. *Plant Cell*

1019   *Physiol* 2002, **43**(12):1518-1525.

1020   89.   Gu H, Wang Y, Xie H, Qiu C, Zhang S, Xiao J, Li H, Chen L, Li X, Ding Z: **Drought**

1021   **stress triggers proteomic changes involving lignin, flavonoids and fatty acids in tea**

1022   **plants**. *Sci Rep* 2020, **10**(1):15504.

1023   90.   Bharadwaj B, Mishegyan A, Nagalingam S, Guenther A, Joshee N, Sherman SH, Basu

1024   C: **Physiological and genetic responses of lentil (Lens culinaris) under flood stress**.

1025   *Plant Stress* 2023:100130.

1026   91.   Kikuchi S, Asakura Y, Imai M, Nakahira Y, Kotani Y, Hashiguchi Y, Nakai Y, Takafuji

1027   K, Bedard J, Hirabayashi-Ishioka Y *et al*: **A Ycf2-FtsHi Heteromeric AAA-ATPase**

1028   **Complex Is Required for Chloroplast Protein Import**. *Plant Cell* 2018, **30**(11):2677-

1029   2703.

1030   92.   Schreier TB, Clery A, Schlafli M, Galbier F, Stadler M, Demarsy E, Albertini D, Maier

1031   BA, Kessler F, Hortensteiner S *et al*: **Plastidial NAD-Dependent Malate**

1032   **Dehydrogenase: A Moonlighting Protein Involved in Early Chloroplast**

1033   **Development through Its Interaction with an FtsH12-FtsHi Protease Complex**.

1034   *Plant Cell* 2018, **30**(8):1745-1769.

1035   93.   Drescher A, Ruf S, Calsa T, Jr., Carrer H, Bock R: **The two largest chloroplast**

1036   **genome-encoded open reading frames of higher plants are essential genes**. *Plant J*

1037   2000, **22**(2):97-104.

1038   94.   Xing J, Pan J, Yi H, Lv K, Gan Q, Wang M, Ge H, Huang X, Huang F, Wang Y *et al*:

1039   **The plastid-encoded protein Orf2971 is required for protein translocation and**

1040   **chloroplast quality control**. *Plant Cell* 2022, **34**(9):3383-3399.

1041   95.   Chen J, Zang Y, Shang S, Yang Z, Liang S, Xue S, Wang Y, Tang X: **Chloroplast**

1042   **genomic comparison provides insights into the evolution of seagrasses**. *BMC Plant*

1043   *Biol* 2023, **23**(1):104.

1044   96.   Xie Z, Merchant S: **The plastid-encoded ccsA gene is required for heme attachment**

1045   **to chloroplast c-type cytochromes**. *J Biol Chem* 1996, **271**(9):4632-4639.

1046   97.   Kranz R, Lill R, Goldman B, Bonnard G, Merchant S: **Molecular mechanisms of**

1047   **cytochrome c biogenesis: three distinct systems**. *Mol Microbiol* 1998, **29**(2):383-396.

1048   98.   Maheswari P, Kunhikannan C, Yasodha R: **Chloroplast genome analysis of**

1049   **Angiosperms and phylogenetic relationships among Lamiaceae members with**

1050   **particular reference to teak (Tectona grandis L.f)**. *J Biosci* 2021, **46**.

1051   99.   Zhang Y, Du L, Liu A, Chen J, Wu L, Hu W, Zhang W, Kim K, Lee SC, Yang TJ *et al*:

1052   **The Complete Chloroplast Genome Sequences of Five Epimedium Species: Lights**

1053   **into Phylogenetic and Taxonomic Analyses**. *Front Plant Sci* 2016, **7**:306.

1054   100.   Huang LS, Sun YQ, Jin Y, Gao Q, Hu XG, Gao FL, Yang XL, Zhu JJ, El-Kassaby YA,

1055   Mao JF: **Development of high transferability cpSSR markers for individual**

1056   **identification and genetic investigation in Cupressaceae species**. *Ecol Evol* 2018,

1057   **8**(10):4967-4977.

1058   101.   Leontaritou P, Lamari FN, Papasotiropoulos V, Iatrou G: **Exploration of genetic,**

1059   **morphological and essential oil variation reveals tools for the authentication and**

1060   **breeding of Salvia pomifera subsp. calycina (Sm.) Hayek**. *Phytochemistry* 2021,

1061   **191**:112900.

1062   102.   Snoussi M, Riahi L, Ben Romdhane M, Mliki A, Zoghlami N: **Chloroplast DNA**

1063   **Diversity of Tunisian Barley Landraces as Revealed by cpSSRs Molecular**

1064   **Markers and Implication for Conservation Strategies**. *Genet Res (Camb)* 2022,

1065   **2022**:3905957.

103. Song SL, Lim PE, Phang SM, Lee WW, Hong DD, Prathep A: **Development of**
chloroplast simple sequence repeats (cpSSRs) for the intraspecific study of
Gracilaria tenuistipitata (Gracilariales, Rhodophyta) from different populations.
BMC Res Notes 2014, **7**:77.

104. Wheeler GL, Dorman HE, Buchanan A, Challagundla L, Wallace LE: **A review of the**
prevalence, utility, and caveats of using chloroplast simple sequence repeats for
studies of plant biology. *Appl Plant Sci* 2014, **2**(12).

105. Feng JL, Wu LW, Wang Q, Pan YJ, Li BL, Lin YL, Yao H: **Comparison Analysis**
Based on Complete Chloroplast Genomes and Insights into Plastid Phylogenomic
of Four Iris Species. *Biomed Res Int* 2022, **2022**:2194021.

1076