1   A candidate sex determination locus in amphibians,
2   evolved by structural variation between X- and Y-chromosomes
3

Heiner Kuhl[1,*], Wen Hui Tan [2,*], Christophe Klopp[3], Wibke Kleiner[1],

Baturalp Koyun[1,4], Mitica Ciorpac[5,6], Romain Feron[7,8], Martin Knytl[9,10],

Werner Kloas[1], Manfred Schartl[11,12], Christoph Winkler[2,§], Matthias Stöck[1,§]

[1] Leibniz-Institute of Freshwater Ecology and Inland Fisheries, IGB,

Müggelseedamm 301 & 310, 12587 Berlin, Germany,

[2] Department of Biological Sciences and Centre for Bioimaging Sciences, National

University of Singapore, 14 Science Drive 4, Block S1A, Level 6, Singapore 117543,

[3] SIGENAE, Plate-forme Bio-informatique Genotoul, Mathématiques et Informatique

Appliquées de Toulouse, INRAe, 31326 Castanet-Tolosan, France

[4] Bilkent University, Department of Molecular Biology and Genetics, Faculty of

Science, SB Building, Ankara, Turkey 06800,

[5] Danube Delta National Institute for Research and Development, Tulcea 820112,

Romania,

[6] Advanced Research and Development Center for Experimental Medicine –

CEMEX, "Grigore T. Popa", University of Medicine and Pharmacy, Mihail

Kogălniceanu Street 9-13, Iasi 700259, Romania,

[7] Department of Ecology and Evolution, University of Lausanne, Lausanne,

Switzerland,

[8] Swiss Institute of Bioinformatics, Lausanne, Switzerland,

[9] Department of Cell Biology, Faculty of Science, Charles University, Viničná 7,

Prague, 12843, Czech Republic

[10] Department of Biology, McMaster University, 1280 Main Street West, Hamilton L8S

4K1, Ontario, Canada

[11] Developmental Biochemistry, Biocenter, University of Wuerzburg, Am Hubland,

97074 Wuerzburg, Germany,

[12] The Xiphophorus Genetic Stock Center, Department of Chemistry and

Biochemistry, Texas State University, San Marcos, Texas, TX, 78666, USA,

[*] These authors contributed equally to the paper.

[§] Co-corresponding authors.

Abstract

Most vertebrates develop distinct females and males, where sex is determined by repeatedly-evolved environmental or genetic triggers. Undifferentiated sex chromosomes and large genomes have caused major knowledge gaps in amphibians. Only a single master sex-determining gene is known in >8650 species, the *dmrt1*-paralogue (*dm-w*) of female-heterogametic clawed frogs (*Xenopus*; ZW♀/ZZ♂). Combining chromosome-scale female and male genomes of a non-model amphibian, the European green toad, *Bufo*(*tes*) *viridis*, with ddRAD- and whole genome pool-sequencing revealed a candidate master locus, governing a male-heterogametic system (XX♀/XY♂). Targeted sequencing across multiple taxa uncovered structural X/Y-variation in the 5'-regulatory region of the gene *bod1l*, where a Y-specific non-coding RNA, only expressed in males, suggests that this locus may initiate sex-specific differentiation. Developmental transcriptomes and RNA *in-situ* hybridization show timely and spatially relevant, sex-specific ncRNA-Y and *bod1l*-gene expression in primordial gonads with coinciding differential *H3K4me*-methylation in pre-granulosa/pre-Sertoli cells, pointing to a specific mechanism of amphibian sex determination.

Introduction

Most vertebrates occur as females and males. Vertebrate sex determination comprises environmental triggers, or simple to complex genetic (genotypic) sex determination – and perhaps combinations thereof – that evolved several times[1,2]. Likewise, the genetic and cellular developmental processes directing undifferentiated gonads towards testes or ovaries are diverse[3].

Genetic sex determination is governed by sex chromosomes that are typically differentiated and often microscopically distinguishable in mammals, birds, and several reptile groups[4]. With several exceptions, some reptilians and most other cold-blooded (poikilothermic) vertebrates, like fish and amphibians, exhibit poorly or undifferentiated (homomorphic) sex chromosomes, which are prevailing in the tree of life[5]. In vertebrates, three main evolutionary concepts explain this prevalence: (i) frequent turnovers, overthrowing sex-determining genes by novel autosomal mutations, or even the sex determination system (e.g. XY to WZ)[6], (ii) recurrent X-Y recombination, e.g. by occasional sex reversal to XY-females[7], and (iii) "jumping master genes" with conserved genetic triggers translocating to other chromosomes[8].

Conserved poorly or non-differentiated sex chromosomes are known in several
clades of poikilothermic vertebrates, e.g., in sturgeons, pikes, and skinks[9-11].

The majority of amphibians possess undifferentiated sex chromosomes with either
female (ZZ/ZW) or male (XY/XX) heterogamety[12,13]. Several examples of
cytogenetically differentiated sex chromosomes[13,14], a female W0/00 male sex
determination system[15], and several amphibians with multiple sex chromosomes
have been identified in both laboratory strains[16] and natural populations[17]. Notably,
to date, in >8,650 species of amphibians[18], the widespread combination of poorly or
undifferentiated sex chromosomes with large genome-sizes (see below) have
caused a basic knowledge gap with only a single though very well-characterized
amphibian master sex-determination gene known in a model species, the African
clawed frog (*Xenopus laevis*[12,19]) and some related species[20]. The *Dm-w* gene arose
after (and perhaps in response to) tetraploidization, is essential as recently shown in
knock-out females[21], and has been lost secondarily in some tetraploid species[22].

True toads, Bufonidae, share large genomes, conserved karyotypes[23] and
undifferentiated sex chromosomes[24]. Typical of most bufonid toads is also Bidder's
organ that develops in males and - in many species - also in females during early
gonadogenesis from the anterior parts of the primordial gonads, considered to
possibly present a rudimentary ovary[25]. Using sex-reversal experiments, castrating
adult males and regrowing functional ovaries from Bidder's organs by Harms[26], early
on, Ponse[27] inferred female heterogamety (ZW) in common toads (*B. bufo*). This was
also cytogenetically deduced in related *B. gargarizans*[28], and cane toads (*Rhinella*
marina[29]). With few exceptions, a meta-analysis of sex-differential F1-hybrid fitness
of interspecific crosses in bufonids likewise supported widespread female
heterogamety[30].

This situation contrasts with that in our target species, Palearctic green toads of the
Bufo(*tes*) *viridis* sub-group*. While their sex chromosomes are also
undifferentiated[24,31], sex-reversal experiments in '*B. viridis*' from Asia Minor[32]
suggested an XY-system. Molecular evidence for male heterogamety resulted from
sibship analyses using microsatellites and nuclear sequence markers[33-35], identifying
the largest linkage group (LG1), homologous to autosomal LG1 of *X. tropicalis* and
harboring *dmrt1*, as sex-linked in several diploid green toad species. All feature
strongly-reduced X/Y-recombination but X/Y-exchange, in the pseudo-autosomal
region, is suggested at evolutionary time scales[33]. However, phylogenetic analysis

revealed X- and Y-alleles of *dmrt1* to cluster by species and not by gametologue,

suggesting X/Y-recombination involving *dmrt1*, which rejected its role as master sex

determiner[35]. Details of sex determination have not been examined in allopolyploids

green toads[36].

In mammals (6,649 species[37]), genome assemblies are available for about one third

(2740[38]). Although there are more species described in amphibians (>8650[18]), owing

to large genome sizes and excessive repetitive regions, to date only >80 amphibian

genomes have been assembled, including 56 out of 7627 Anura, 19 out of 805

Urodela (Caudata) and 9 out of 221 Gymnophiona[18,38]. However, profiting from long-

read- and 3D-sequencing technologies, amphibian genomics is on the rise.

Here, we provide a chromosome-scale genome of a non-model amphibian, the

European green toad, *Bufo*(*tes*) *viridis*, and identify a specific candidate master sex

determining locus - present in multiple related green toad taxa - that evolved by

structural variation, involving a Y-specific ncRNA, in the regulatory region between X

and Y chromosome.

Results

Chromosome-scale genome assembly of *B. viridis*

We obtained a 3.8 Gbp chromosomal-level genome assembly using Oxford

nanopore (ONT) long-read (N50 = 11.6 kbp; sequencing coverage 21x) and Illumina

short-read data (2x150 bp; sequence coverage 47x) for the European green toad

(Fig.1a,b). N50 contig size of the final assembly was 1.9 Mbp. Eleven chromosomes

(scaffold N50 = 470 Mbp; consisting of 94% of the contigs total length), consistent

with the karyotype of *B. viridis* (Fig.1c), were constructed using chromosome

conformation capture (Hi-C) short reads (318 Gbp). Many unplaced contigs contain a

known long satellite DNA family[39]. Comparing the chromosomes to the *B. bufo*

assembly revealed genome-wide synteny and collinearity with few inversions,

despite significant differences in assembly size (VGP's aBufBuf1.1 = 5.0 Gbp;

Fig.1b). *De novo* analysis revealed 68.5% repeat content in *B. viridis*. Using

published data from related species and RNAseq data, we annotated 21,605 protein

coding genes in the genome. BUSCO (Geneset vertebrata_ODB9)-scoring of protein

predictions revealed 96.4% complete coding sequences and a low missing gene

fraction of 1.7% (C:96.4% [S:94.2%, D:2.2%], F:1.9%, M:1.7%, n:2586).

ddRADseq in *B. viridis* genetic families

We first attempted to identify the sex determination region by double digest restriction-site associated DNA sequencing (ddRADseq). In three genetic families of European green toads, we analyzed a total of 443,378 markers, present in at least one individual with a depth higher than 4. Of these, 96 were significantly associated with male sex ($p < 0.05$; Bonferroni correction), but none was found in all males (Supplementary Fig. 1), probably due to high stochasticity of random sequencing of large genome fractions. When mapped to the *B. viridis* genome, 64 of these 96 markers were located along the entire chromosome 1 (scf1) (Supplementary Fig. 2), one on chromosome 7 (scf7), five on unplaced scaffolds, and 26 were not aligned uniquely or with low quality (mapping quality, MQ < 20). Overall, this strongly supports a male (XY) genetic sex determination system on chromosome 1, however, with a very small sex-specific region that is not fully covered in each individual, and sex-linkage of most ddRAD markers is not evolutionarily conserved due to X-Y recombination, in the huge pseudo-autosomal region (PAR) of this chromosome 1. While this approach powerfully corroborated previous microsatellite-based results[33], it did not provide a sex-diagnostic marker.

Pool-seq yields a cross-amplifying marker in multiple taxa

To narrow down the sex determining region on scaffold 1, we used whole genome pool sequencing (from here: "pool-sex"), involving 22 female and 22 male phenotypically-sexed adult green toads from multiple European green toad populations (Supplementary Table 1), chosen to be closely related according to phylogeographic information[40,41] and presumably sharing the same sex chromosome, based on microsatellites[33-35]. Mapping pool-sex short reads to the reference, genome-wide revealed a single peak in a region on scaffold 1 (scf1), where the number of clustered heterozygous SNPs (N = 103) in males surpassed about nine times the maximum number of clustered heterozygous SNPs (N = 12) in females (Fig.1c, Supplementary Table 2; Supplementary Fig. 3). This result was later confirmed quantifying male-heterozygous SNPs on scaffold 1 of the male genome assembly as well as male-specific pool-seq coverage on the male genome (see below and Fig. 1c).

This supports a male-heterogametic (XY) system, involving a short 66 kb region (scf1:566,783,058-566,848,882 bp). This locus encompasses one third of the 5'-

upstream-region and two thirds of the gene *bod1l* (bufVir1.4895.2, biorientation of
chromosomes in cell division 1 like 1). Transcriptomes and cDNA-based PCR-
products of males of *B. viridis* exhibited consistent heterozygosity of exons 7 to 10
(scf1:566,831,904-566,833,300), while females were always homozygous.

The heterozygous 5'-region exhibited several INDELs between X and Y haplotypes,
enabling primer design. In females, BvXY1 amplified a single PCR-product - visible
as one band on the gel, and two to three bands in males, widely consistent with an
XX♀/XY♂-system, but suggesting multiple Y-specific signals in males, which might
be sex-chromosomal duplications, nonspecific PCR-amplicons or even PCR-
recombinants. Of note, BvXY1-primers amplified not only sex-specific products in
individual *B. viridis* (sensu lato) males and females of the pool-sex-approach but also
in multiple phenotypically-sexed males and females of this and four additional taxa
(*B. balearicus*, *B. siculus*, *B. turanensis, B. shaartusiensis*), comprising a
monophyletic radiation, probably >3 Mio. years old[36] (Fig. 1e-h). The sex-diagnostic
PCR-marker BvXY1 amplified a single specific 100-200 bp band in females of six
green toad taxa, but 2-3 bands in their males, one in females' size, presenting the X-
copy. The additional male bands with the initial BvXY1-primers yielded identical or
similar sequences indicating duplicates on the Y-homologue.

Synteny and FISH-TSA of *bod1l* dismiss duplication with translocation

New sex determining genes evolve by allelic diversification or from gene
duplications[42]. To elucidate the situation in *B. viridis*, we examined synteny and
localization of the candidate locus. *Bod1l* lies between the genes *Cpeb2* and *nkx3-2*
(NK3 homeobox 2), on the long arm of scaffold 1, an evolutionarily conserved gene
order likewise found in *B. bufo*, *Rana temporaria* and *Xenopus tropicalis*, suggesting
synteny throughout anurans.

Fluorescence *in-situ* hybridization with tyramide signal amplification (FISH-TSA) on
metaphase chromosomes of a male *B. viridis* revealed two signals, one on each
homologue, for a *bod1l*-specific probe at a position on the distal part of the short (p)
arm of chromosome 1 (Fig. 1f). Two (and not three) signals suggest that this gene
exists in two copies in XY-males of *B. viridis*, implying that both heterozygous
sequences evolved by structural diversification[42] rather than translocation of a male-
specific duplication, as shown for *dm-w* of female *Xenopus laevis*[19]. This result is
further supported by the pool-seq and long-read male coverage (see below).

Targeted sequencing in multiple related species sharing the sex marker

If the hypothetically sex-determining SNPs in coding *bod1l*-regions arose in a
common ancestor, we expected the five diploid taxa, sharing BvXY1 in 5'-*bod1l*-
regions (Fig. 1d), to share at least some exonic, possibly functional SNPs. To test
this, we applied targeted enrichment to multiple females and males of five taxa,
using an AmpliSeq custom DNA panel on 80 kb (93%) of the *bod1l*-X-copy, including
its 5'-region with BvXY1, and analyzed 8183 SNPs. Of note, none of the *bod1l*
coding SNPs*,* present in *B. viridis* (Supplementary File 3), was conserved across all
five toad taxa (Supplementary Files 4, 5, Supplementary Fig. 4a), suggesting that
this coding region (cds) does not contain the mechanistic key to green toad sex
determination. Instead, the targeted enrichment approach narrowed down the sex-
specific region in the candidate locus to the 5'-region of *bod1l*, containing BvXY1
(Supplementary Fig. 4b).

Y-chromosomal structure of *B. viridis* from long-read WGS

After first working with the female reference genome, X-haplotype and *bod1l*-coding
region, we then moved the focus back to the 5'-end of *bod1l* and the BvXY1-vicinity
on the Y-haplotype. To elucidate this Y-chromosomal structure, we reconstructed
this region, which showed about 15x coverage (haploid), from ca. 30x whole genome
long-reads of a single male *B. viridis* and aligned it to the X-copy of the *bod1l*-region.
Female-to-male pool-seq-coverage of X- and Y-haplotypes yielded the expected
diploid female and haploid male coverage (Supplementary Fig. 5 and 6).
Additionally, improved base-calling and haplotype-aware correction algorithms
became available and enabled us to also assemble a whole male genome from that
data, which supported the results of the manual approach (Supplementary File 2).

In the *bod1l*-region, the confidently assembled Y-haplotype-specific sequence
comprises about 55 kb with some greater-than-zero female pool-seq-coverage,
explicable by repeat-enriched fragments, not assignable to X or Y (Supplementary
Fig. 7). The male long-read whole genome assembly revealed several structural
changes between X- and Y-haplotypes in the 5'-*bod1l*-region. The *B. viridis*-Y
contains large insertions, mainly stemming from repetitive elements, and a non-
coding RNA (ncRNA, see below). None of the so-called 'usual suspects', genes that
often acquire sex-determination function in vertebrates[1], was detected in this region.

Larval gene expression in sibling *B. viridis* tadpoles from multiple localities

Pool-sequencing of genomic DNA in *B. viridis* had revealed consistent heterozygosity in males for *bod1l*-exons 7 to 10. Accordingly, RNAseq revealed sex-differential expression (scf1:566,831,904-566,833,300), where heterozygous *B. viridis* males showed 1 SNP in exon 7, and 14 SNPs in exon 10, while females were always homozygous (confirmed by cDNA-based PCR-products). To test whether *bod1l*-products are differentially composed after transcription of X- and Y-copy, we analyzed splice sites in developmental transcriptomes, including the sex determination period. The absence of significant sex-specific differences excluded alternative splicing. However, this analysis allowed manual improvement of the *bod1l*-gene model.

To study the expression of the candidate sex determination gene (*bod1l*),the adjacent ncRNA-Y and ca. 80 other genes (Supplementary File 6) playing roles in sexual development and/or sex determination in other vertebrates ('usual suspects'[1]), we then examined larval transcriptomes of six sibling groups, in total 54 individual toads from six locations, ranging from Gosner[43] stage 23 (day 10 after fertilization) until Gosner stage 46 (the last about 6 weeks after metamorphosis) as well as from two adult gonads.

Since gonadal differentiation is histologically detectable only at or after metamorphosis completion (>Gosner 46[25,35,43]), the critical period of genetic sex determination lies earlier, within our sampling period. In Gosner stage 46, 6 weeks after metamorphosis, we could anatomically distinguish between Bidder's organ and early testis, proofing sexual differentiation. Until Gosner stages 34-38 (Fig. 2), overall *bod1l*-expression varies in *B. viridis* between males and females (. From Gosner-stages 38 to 43, other genes, specific for male sexual development, like *sox9*, *sox10*, *sox17*, and *sf1*, show higher expression in males than in females. Other such genes change their expression distinctly after Gosner stage 38, like *ck18* (bufVir1.18618.1) or *11ßHD* (bufVir1.16559.1) or - in our series - were detected only in Gosner stage 46 and subadult testes (*dmrt1, amh*). Female-specific genes also changed their expression around Gosner stages 34-38; *wnt4*, *ihh*, *foxL2*, *ptch1*, *gli3*, *fshr* from Gosner stage 43 or 45; however, their expression was stronger in males than females. Expression of *gdf9, cyp19a1a*, and *cyp17a1* was detectable only in Gosner 46 and adult ovaries, with *cyp19a1a* and *cyp17a1* presenting good markers for the

timing of gonadal sexual differentiation in anurans. *Bod1l*-expression differences

between females and males (Fig. 2) lack statistical significance but occur at the right

developmental period (Gosner stages 27-44), as required for a sex-determining

gene. Early male transcriptomes (Gosner stages 23-44) contain also the bufonid-

specific Bidder's organ, considered similar to a rudimentary ovary, that develops

adjacent to the anterior primordial gonads[25]. Published qPCR-data suggested that

gene expression in adult Bidder's organs differs from that of testes and ovaries but

resembles the latter, consistent with classical, mid-20th-century male-to-female sex-

reversal experiments, when Bidder's organs grew and resumed oogenesis after the

removal of testes[27] (reviewed[25]). In green toads, Bidder's organ develops in males

but barely in females[25]. Given that the "ovary-like" primordial Bidder's organ is

contributing tissue to our RNAseq-analysis of male tadpoles until Gosner stage 44,

we assume a "female-bias" in males' transcriptomes (Gosner stages 23 to 44). In

conclusion and in line with the AmpliSeq-data from multiple species, only sharing the

BvXY1-marker across taxa, we consider the male-only expression of ncRNA-Y and

the quantitative differences in additive *bod1l*-expression between males and females

(Fig. 2) as sufficient to trigger green toad sex determination. Of note, there is a trend

of greater Y-haplotypic expression of *bod1l* between Gosner-stages 38 and 43

(coinciding with an increase of the ncRNA-Y expression, next paragraph) and a

distinctly greater additive *bod1l*-XY-expression in males than XX-expression in

females (Fig. 2, Supplementary Fig. 8). *Bod1l*-Y also shows the highest expression

in testis compared to other male organs (heart, brain, liver).

A non-coding RNA (ncRNA) is Y-specifically expressed upstream of *bod1l*

When further comparing RNAseq (mapped to Y- or X-haplotypes) from ovaries *vs.*

testes (in tadpoles plus adjacent Bidder's organ), we discovered a long non-coding

RNA ("ncRNA-Y", ca. 1.2 kb, 4 exons, Fig. 3) with its exons 1 and 2 being Y-specific,

and expressed exclusively in males, as confirmed by PCR from cDNAs (Fig. 3c,

Supplementary Fig. 9). The ncRNA-Y harbors the initially detected gDNA-marker

(BvXY1) in intron 3. We hypothesize that this male-specifically expressed ncRNA-Y

hints at an enhancer element, missing on the X, which regulates male *bod1l*-

expression and simultaneously transcribes the ncRNA-Y in a one-dimensional (1D)

manner from the opposite strand[44]. With ca. 80 kb, the putative enhancer sits in a

reasonable distance to the *bod1l*-promoter (Fig. 3a,b)[45]. We also tested 200 bp-

"sliding-windows" of the ncRNA-region for their *in-silico* potential as enhancers of
bod1l (Supplementary Table 3; Supplementary Fig. 10). The most stringent threshold
of 0.9 still predicted larger parts of the Y-specific region to function as strong
enhancers, intensified by numerous G-quadruplexes (>25 alone in intron 1 of
ncRNA-Y), known to accumulate at promoters and enhancers[46]. As shown by PCR
using cDNA, this long ncRNA-Y is also expressed in skin, liver, brain and heart of
males after metamorphosis as shown using cDNA, and exhibits simultaneous
expression of the Y-haplotype of *bod1l* with the strongest signals in testis (Fig. 2).
Due to its low expression level, it is only detectable in high-coverage transcriptomes
of the primordial male gonads and adult testes but not in heart, brain and liver high-
coverage transcriptomes. It comprises 5.8 kb on the gDNA level and can be male-
specifically PCR-amplified in all those six taxa containing the BvXY1-marker and
suggests a common ancestral feature (Fig. 3d).
320
Spatially different gene expression in the developing gonads
For a sex-determination gene, not only the accurate timing of its expression but also
its organ- and cell-specific spatial expression is essential. We provide evidence by
RNA *in-situ* hybridization in histological sections of gonads in Gosner stages 36-37,
where *bod1l* shows sex-specific differential expression. In female gonads, *bod1l*-
expression was evident in *vasa*-positive primordial germ cells (developing oogonia)
but absent from surrounding somatic pre-granulosa cells (Fig. 4a,c, Supplementary
Fig. 11). In contrast, *vasa*-positive male primordial germ cells (developing
spermatogonia) as well as a majority of adjacent somatic pre-Sertoli cells exhibited
bod1l-expression (Fig. 4b,d, green arrows). In Gosner stages 43-44, at the beginning
of metamorphosis, *bod1l*-expression in females remained similar to Gosner 36-37. In
males, however, *bod1l*-expression was no longer detectable in somatic pre-Sertoli
cells (Fig. 4e,f; Supplementary Fig. 11). This suggests a dynamic regulation of *bod1l*
in early male and female gonads, with expression in somatic cells exclusively in male
but not female gonads. This precedes the differences observed by RNAseq from
trunk sections in Gosner stage 43-44, when quantitative differences of sex-specific
bod1l-expression occur (Fig. 2).
It has been shown that *Bod1l* interacts with *Setd1a* to facilitate tri-methylation of
histone H3 at lysine 4 (H3K4me3[47]), with similarities in eukaryotes from yeast to
mammals[48]. To examine its status in male and female gonads, we performed

H3K4me3-immunostaining on Gosner 43-44 gonads (Fig. 4g,h). In females, gonads

exhibited distinct H3K4me3-signals in nuclei of both germ and somatic pre-granulosa

cells (Fig. 4g, white asterisks and arrowheads, respectively; Supplementary Fig. 11).

In contrast, H3K4me3 was strongly detected in male germ cell nuclei (Fig. 4h, white

asterisks) but hardly detectable or absent in nuclei of somatic pre-Sertoli cells (Fig.

4h, yellow arrowheads). In Gosner stage 46, six weeks after metamorphosis, *bod1l*

was expressed (RNAseq, PCR of cDNA) in both sexes in several somatic tissues

(brain, heart, liver, intestine) as well as in gonads.

XY-phylogeny in six Eurasian diploid taxa from targeted sequencing

For a sex-determining locus, evolved from a common ancestor, non-recombining loci

are expected to cluster by gametologue (i.e., X- or Y-specific haplotypes) and not by

species. Analyzing X- and Y-specific sequences obtained from the AmpliSeq custom

DNA panel and amplicon sequencing of ncRNA-Y of six diploid green toad taxa, we

show that in the 2,671 bp-region (scf1:566,788,797-566,791,468), containing the

sex-specific BvXY1-marker, sequences cluster by gametologue (Xs *vs.* Ys,

Supplementary Fig. 4c), while in the similarly-sized adjacent upstream region

(scf1:566,786,116-566,788,796), sequences clustered by taxon/species, suggesting

occasional recombination (Supplementary Fig. 4d). This provided evidence of both

the stop of recombination since the sequence arose in a common ancestor and the

apparent functional relevance of this region, since these species also do not share

coding sequences.

Discussion

Our data in *B. viridis* and five related diploid green toad taxa are in accordance with

a locus that comprises a ncRNA and structural changes in the 5'-regulatory region of

a gene (*bod1l*), as well as sex-specific ncRNA-expression in heterozygous (XY)

males *vs.* homozygous (XX) females, accompanied by strong phylogenetic and

poolseq-supported signal of local recombination arrest. Sex-specific expression

occurs at the right time, with male-restricted expression of the ncRNA from the Y,

much earlier than histologically detectable differentiation, and *bod1l*-expression in

the target organs (primordial gonads), and therein in the germ and surrounding

somatic cell populations (in male pre-Sertoli but not in female pre-granulosa cells).

These properties suggest that the region containing ncRNA-Y and *bod1l* presents a

candidate sex determination locus. Future evidence (e.g., using CRISPR/Cas9) is
desirable for its better mechanistic understanding. A single genome-wide peak of
heterozygosity between female and male revealed by pool-seq and coverage
differences on whole genome assemblies of both sexes (Fig. 1c), along with
evolutionary conservation of the structural XY-differentiation in several Eurasian taxa
sharing the Y-specific ncRNA, support its candidacy as the Palearctic green toad sex
determining locus. The expression of the ncRNA-Y and *bod1l* in Bidder's organ of
the differentiating male gonad (Gosner stage 46) may contradict the view that
Bidder's organ in this XX/XY system can be simply interpreted as a 'proto-ovary'.

The only known amphibian master sex determining gene, *dm-w* of clawed frogs
(*Xenopus*), governs female-heterogamety (ZW♀/ZZ♂) and evolved by gene
duplication as a *dmrt1*-paralog, involving neofunctionalization of a non-coding part of
a DNA-transposon[12,19-22,49]. Here, we provide evidence for a sex-locus in a male
heterogametic bufonid toad system (XX♀/XY♂), evolved by structural diversification
in a common ancestor, as shown by the associated ncRNA-Y-phylogeny (Fig. 3c).
To our knowledge, this reveals only the 2[nd] master sex determining locus in
amphibians.

Our data suggest either a sex-determining function of the ncRNA-Y itself or its
functional interaction with *bod1l*, e.g., as an enhancer[50,51].

Long non-coding RNAs (lncRNAs) play important roles in gonadal development and
sex differentiation[50,52], and thus the ncRNA-Y might suffice as a master sex
determinator, although, so far, we have not detected homology with genes known to
play roles in sexual development. We rather consider a regulatory effect on the
neighboring *bod1l*. The ncRNA-Y, harboring the partially duplicated BvXY1, contains
a neighboring signal for a G-quadruples-rich enhancer towards its 3'-end. This is a
mechanistic hint that the enhancer modifies *bod1l*-expression in XY-males (Fig. 3),
as shown by timely and spatially differential expression in germ and pre-Sertoli but
not in pre-granulosa cells (Fig. 4).

There are also several, partly interlinked pathways by which spatially and/or timely
differential *bod1l*-expression may determine sex in green toads.

(i) Sex determination of vertebrates differs regarding the timing of primordial germ
cells to enter meiosis[53], with females entering meiosis earlier than males, exhibiting
an arrest. *Bod1l* is well-known to maintain genomic integrity at sites of DNA-
replication damage[54]. *Bod1l* interacts towards its N-terminus with the histone

methyltransferase *Setd1a*, which triggers the histone chaperone function of

Fancd2[55], part of the Fanconi anemia group (*Fanc*), that plays roles in zebrafish sex

reversal[56]. Thus, differential *bod1l*-expression might cause a compromised DNA

repair system by shifting sex determination towards males.

(ii) Interaction of *Bod1l* with *Setd1a* and H3K4me3 refers to tri-methylation on lysine

residue 4 in the amino-terminal tail of histone H3, is associated with potentially active

promoter regions[57] and linked to double-strand breaks, essential for meiosis 1

crossing-over, ensuring chromosome segregation[58,59]. Thus, the differential *bod1l*-

expression might enable double-strand breaks as a precondition for the earlier start

of recombination in females but not in males. (iii) H3K4-methylation in mammals[57] is

associated with *de novo* DNA-methylation in germ cells, and essential for

maintaining transcription in the growing oocyte and for triggering zygotic genome

activation[60], with XY-mice deficient for the H3K9-demethylating enzyme *Jmjd1a*

showing increased H3K9-dimethylation and decreased activating marks of H3K4-

trimethylation across the *Sry*-locus, leading to a high frequency of sex reversal[2].

According to Dupont and Capel[61] "prior to sex determination" in placental mammals,

bipotential supporting gonadal cells of both genetic sexes maintain identical

transcriptional programs, allowing commitment to either Sertoli (testes) or granulosa

(ovary) cell fate. In pre-Sertoli and pre-granulosa cells, bivalent promoters of key sex

determining genes (e.g., *Sox9*, *Fgf9*, *Wnt4*, *Foxl2*) have both, H3K27me3 and

H3K4me3 marks[61]. After sex determination by the master sex determination gene

Sry, promoters to be activated in that sex lost the repressive H3K27me3 mark, while

genes to be silenced maintained their bivalency[62]. This suggested that (a) sex-

determining genes are poised for activation (or repression) at the bipotential stage,

(b) gene activation involves removal of H3K27me3 marks, while (c) repression of the

alternative supporting cell fate requires maintenance of existing H3K27me3

repressive marks rather than *de novo* deposition of these marks after commitment to

the Sertoli or granulosa program[61].

One hypothesis is that differential *bod1l*-expression (reminiscent of *Sry*) might

determine sex by changing H3K4-methylation in green toads during early gonadal

development, prior to metamorphosis.

To test the hypothesis that *bod1l-expression* coincides in time and space with

changes of H3K4-methylation in developing green toad gonads, we used H3K4me-

antibodies together with RNA *in-situ* hybridization (Fig. 4). In Gosner stage 43-44,

slightly prior to morphological differentiation, we found a strong difference between
male and female gonadal somatic cell H3K4-methylation. This suggests that spatially
different *bod1l*-expression results in sex-specific methylation patterns that initiate
sex-specific gonadal differentiation. However, opposite to the situation in mammals,
H3K4me3 is weaker in somatic cells of green toad males than those of females (Fig.
4), while increasing H3K4-methylation allows *Sry*-expression activation (male
determination). Another major difference is that the primordial gonad of bufonid
toads, specifically in green toad males, includes the "ovary-like" Bidder's organ.
Given the multiple potential links between *bod1l*-expression and sex-specific meiotic
stages shown above, we hypothesize that this gene may function as a suppressor of
the female sex determination pathway – a scenario found in many vertebrates[63].
Following differential *bod1l*-expression, in the bipotential gonad, ovarian pre-
granulosa cells transdifferentiate to testicular pre-Sertoli cells.
In contrast to the only known amphibian master sex determining gene in clawed
frogs (*Xenopus*), the green toad sex-locus with ncRNA-Y and *bod1l* did not evolve by
gene duplication but structural diversification in the 5'-region of the gene, namely by
Y-specific sequence insertion. This evolution of sex determination seems broadly
similar to allelic diversification, so far unknown in amphibians, but known from many
examples of master genes in fish[42].
462
Methods
Ethics and animal welfare statement
Our research complies with all relevant ethical regulations; the relevant animal
protection committee of the IGB and the city of Berlin (LAGESO) approved the study
protocol where relevant. Most adult toads were sampled more than 20 years ago by
either minimal invasive sampling methods using tail tips of larvae or saliva of adults,
obtained by buccal cotton swabs. We also used tissues of road kill toads and/or
collected a few adult animals in Greece. For sampling and export of toads and
tadpoles, permits were kindly provided by the Ministry of the Environment of Greece
(115790/229) 2014-2017. Developmental stages of tadpoles were obtained from
control groups of an animal experiment approved by the German State Office of
Health and Social Affairs (LaGeSo, Berlin, Germany; G0359/12) and a commercial
breeder. Developmental stages of larvae (tadpoles) and early metamorphosed toads
were observed under a binocular microscope and determined using the drawings of

developmental stages (table) by Gosner[43]. Tadpoles were reared in a 12/12 h
light/dark cycle at constantly 22 ± 1 °C in sufficiently aerated and regularly cleaned
tanks. Weekly monitored water parameters comprised: dissolved oxygen, nitrate,
ammonium, pH, conductivity, and hardness and kept as described[64]. Tadpoles were
fed SeraMicron (Sera, Germany), later supplied with TetraMin (Tetra, Germany). To
imitate natural conditions at metamorphosis, animals were transferred to glass
terraria at Gosner stage 46. Thus, the tadpoles were kept according to approved
practice (permit ZH 114 to IGB), including the provision of appropriate tank size,
sufficient rate of waterflow, natural photoperiod, *ad libitum* food supply, as well as
temperatures within the species' thermal tolerance range. This ensured that no pain,
suffering, distress or lasting harm was inflicted on the animals. According to the
permit G0359/12, tadpoles and juvenile (and two adult) toads were euthanized by an
overdose of Tricaine PHARMAQ 1000 MG/G (MS222) (concentration: 500 mg/L))
since this does not incur pain, suffering or distress.
491
Chromosome-scale female genome assembly
The female *B. viridis* reference genome was sequenced in-house at IGB using ONT
long reads (MinIon device, SQK-LSK109 kits and R9.4.1 flow cells) and Illumina
short reads (Novogene, Europe). The long reads and pre-assembled short reads
IDBA-UD assembly[65] were assembled using the WTDBG2.2 assembler[66].
Sequences were error-corrected using two rounds of Flye polish with long reads[67]
and four rounds of VCFconsensus using short reads[68]. The chromosome-scale
assembly was achieved using Arima HiC-data and the JUICER/3DDNA package and
manual curation by juicebox[69-71]. *De novo* repeat analysis of the assembly was
performed by the RepeatModeler / RepeatMasker package. Protein coding genes
were annotated using RNAseq data of *Bufonidae* as well as Hyloidea transcriptomes
and more distantly related amphibian proteins, available at NCBI. A reference
genome browser was created http://genomes.igb-berlin.de/Greentoad/.[72]
505
Assembly of Y-haplotype
Several iterative approaches were used to extract Y-haplotypic sequencing reads of
B. viridis GR57 male[73] long read data (Oxford Nanopore Promethion platform;
Novogene). First, we mapped the reads to the BvXY1-marker sequences to separate
Y reads of the region. Second, we performed a FLYE[67] assembly of the male data,

identified several contigs with variation compared to the X haplotype (*bod1l*-region),

and extracted the corresponding reads. The Y-specific reads were assembled, then

added to the female assembly and all male reads were mapped to the combined

sequences, which allowed finding more reads matching better to Y than to X contigs.

This procedure was iteratively repeated several times until the Y-haplotype and its

structure could be assembled. Pool-seq data was again mapped to the female

assembly with added Y-haplotype sequences by MINIMAP2[74] showing the expected

coverage differences between female and male pools. A dot plot comparison of X

versus Y haplotype involved the corresponding function of the NCBI Blast website.

Assembly of a male genome

Data that was used for the manual assembly of the Y-haplotype (see above), served

also to perform a complete de novo assembly of a male B. viridis individual. To

improve assembly results RAW data was first base-called by the most up-to-date

basecaller for ONT data (dorado-0.5.1-linux-x64 with dna_r9.4.1_e8_sup@v3.6

model). Then the reads were corrected using the haplotype-aware read correction

method, implemented in PECAT[75]. The corrected reads and the prior flye assembly

contigs (see above) were assembled using wtdbg2. Contigs were corrected by one

round of Flye polish (with corrected long-reads) and two rounds of VCFconsensus

(with male poolseq data). Male Hi-C data was mapped to the contigs using

CHROMAP[76] and chromosomal scaffolds were computed by YAHS[77]. Some

unplaced contigs could be placed into the YAHS scaffolds using RAGOUT2[78] and

synteny analysis with the female genome. The assembly was scored using BUSCO

analysis and had highly similar scores as the female assembly.

Double-digest RAD sequencing (ddRADseq)

DNA of three genetic families consisting for one data set of both parents (buccal

swabs) and 40 sexed mostly phenotypically offspring and for two other sample sets

of 25 or 15 siblings, was extracted using the DNeasy Tissue Kit or the BioSprint

robotic workstation (QIAGEN), following the manufacturer's protocols. Double-digest

RADseq libraries were prepared according to this study[79]. In brief, genomic DNA

digestion by two restriction enzymes (*Sbf*I, *Mse*I) and subsequent ligation of

fragments to adapters containing unique barcodes (4–8 bp) was followed by library-

PCR for 20 cycles for only different cut-sites containing fragments, and a gel-based

size selection, isolating the 400–500 bp fraction. RADseq libraries were single-end

sequenced on an Illumina HiSeq 2500, at a 20x coverage.

Raw ddRAD reads were demultiplexed with STACKS *process_radtags* version

2.55[80], using the enzymes sbfI and mseI (*--renz_1 sbfI --renz_2 mseI*) and the

parameters *-c* (clean data), -r (rescue barcodes and RAD-Tag cut sites), and -q

(discard reads with low quality scores).

Demultiplexed reads were used as input for the RADSEX workflow[81]. First, a table

containing the number of occurrences of each unique sequence in the reads file in

each individual was generated with *radsex process*, using default parameter values.

Then, the distribution of markers between males and females was computed with

radsex distrib using a minimum depth to consider a marker present in an individual of

5 (--min-depth 5). Markers were aligned to the genome of *B. viridis* (bufVir1) using

radsex map using a minimum depth of 5 (--min-depth 5) and other parameters set to

default. Results were plotted with the *SGTR* package, with the entire process

provided as a Snakemake workflow[82].

Poolseq approach using phenotypically sexed toads: 'poolsex'

Of 22 safely sexed adult toads of each sex from multiple populations in the range of

the European green toad taxa (*B. viridis viridis* and *B. viridis variabilis*), using buccal

swabs or tissues genomic DNA was extracted using the DNeasy Tissue Kit or the

BioSprint robotic workstation (QIAGEN), following the manufacturer's protocols.

Individual DNAs were quantified using the QuantiFluor® dsDNA System kit

(Promega), adjusted and equimolarly mixed to yield a female and a male pool.

Whole genome library preparation for 300 bp fragments and whole genome

sequencing was performed at NOVOGENE (UK), on a NovaSeq PE150 to yield 131

and 140 Gbp of data for male and female pool, respectively. Reads were mapped to

the reference genome using the short-read mode of MINIMAP2[74] (parameters -K

2000M -I 100G -a -x sr -t 80, and converted to sorted bam-files by SAMTOOLS[83].

Variants (SNPs, MNPs, INDELs) were called from both bam files using the

PLATYPUS variant caller[84] with integrated duplicate fragment reads removal (–

filterDuplicates = 1) and reassembly of reads (–assembleAll = 1). The resulting vcf-

file was screened for XY- or XX-specific variants using AWK-scripting and different

variant read coverage cutoffs (i.e. SNP read coverage in females and males >=15;

variant read count in females <=7% or >=93% of total SNP read coverage as a

condition for homozygosity; variant frequency in males between 45 - 55% of total

SNP read cov. as a condition for heterozygosity; for screening ZW SNPs conditions

for males and females were interchanged). Variants were clustered according to

their distance in the genome (maximal distance of variants in a cluster: 9000 bp),

counted using BEDTOOLS (merge, annotate[85]) and sorted by variant count. Regions

with highest variant counts or densities and neighboring regions were inspected for

genes related to sex determination. The same heterozygous SNP analysis was

repeated using a male genome assembly (see above). Additionally, an analysis of

male-specific poolsex coverage of the male genome was performed. Here, we used

the bam files (MQ60 reads only) to analyze bp-level sequencing coverage using

BEDTOOLS genomecov. The resulting bedgraph files for female poolseq and male

poolsex were combined by BEDTOOLS unionbedg. Coverage of both poolsex

datasets was normalized by their modal coverage peak (setting diploid coverage =>

1.0). Then bp positions were extracted, if they had values of 0.25 - 0.75 for male-

and < 0.05 for female-poolsex coverage. These genomic positions were clustered by

distance and the number of male-specific covered bases in each cluster was

counted BEDTOOLS(merge, annotate).

Sex-diagnostic PCR marker design

The region 566,783,058-566,848,882 bp on scaffold 1 includes several INDELs

apparent between the X and the Y haplotype, allowing design of a number of primers

that were primarily tested in gradient-PCRs on phenotypically sexed *B. viridis*. While

several primers showed sex-specific amplification in this species, one (BvXY1) could

be optimized to work across multiple species.

A collection of DNAs (extraction: see above) comprising samples of phenotypically

sexed green toads of the entire radiation[36] served to test sex-specificity of all PCR

markers. PCR-amplifications were carried out in 25 µL reactions containing 1.25 µL

of each primer (BvXY1_F: 5'_CTGAATAGCAATTGGACACAGC_3; BvXY1_R:

5'_ATGTGAGAGCTGTTACAATGGA_3'), 14.5 µL of water, 5 µL of AllTaq buffer

(5x), 0.5 µL of AllTaq polymerase (Qiagen), and 2 µL of DNA (concentration 4-20

ng/µL). Temperature regime comprised 2 min at 95°C (initial denaturation), and 38

cycles of (1 min at 95°C denaturation, 45 s at 59°C annealing, 45 s at 72°C

extension), and 5 min at 72°C final extension. PCR products (4 µL) were visualized

by electrophoresis on 4% high-resolution Sieve-agarose (Biozym) gels, run for 60
min at 70 V.

To PCR-amplify either a 2.5 kb fragment or nearly the entire ncRNA-Y, comprising
5.8 kb based on gDNA on the Y-chromosome in all diploid green toad taxa with sex-
specific BvXY1-markers, we used the same ingredients and volumes as above either
with primers ncRNA_Y_ex1F: 5'_GGGCAGTGAAGAGAAGTCCT_3',
ncRNA_Y_ex2R 5'_TTCACCATTCTCCAGCCTGT_3' for the 2.5 kb-fragment or
5.8kb_ncRNA_Y_F: 5'_GTGGTAGTTGCTGTGCCTTT_3', 5.8kb_ncRNA_Y_R:
5'_CCGGTGTGGTGTAAAATATCCA_3' for the 5.8 kb-piece. Temperature regimes
involved 2 min at 95°C, and 37 cycles of (1 min at 95°C denaturation; 2.5 min at
63°C (for the 2.5 kb) or 5 min at 59.4°C (for 5.8 kb) annealing; 1 min at 72°C
extension and 5 min at 72°C final extension (note extended annealing times). PCR
products (4 µL) were visualized by electrophoresis on 1.5% agarose (Biozym) gels,
run for 60 min at 80 V. PCR-products (21 µL) were barcoded using the NBD196 and
LSK110 ONT-kits, and sequenced using the ONT-MinIon device, as described.
Basecalling was performed using ONTs GUPPY version 6 and the super accuracy
base-calling model (SUP-model).

Phylogenetic tree of a 2.7 kb-region at the Y-specific region boundary
Consensus sequences of 100 ONT reads of Y-specific amplicons were calculated
using NGSPECIESID[86]. Homologous X-sequences were reconstructed from targeted
enrichment sequencing data of females using BCFTOOLS consensus[87] for the
region scf1:566,788,797-566,791,468. We also excluded regions of the sequence
not covered by enrichment data by supplying a masking file through parameter "-m"
of bcftools consensus. All resulting sequences were multiple aligned using MAFFT
(automatic alignment parameters). The alignment was filtered by GBLOCKS[88] to
remove columns with excessive missing data. IQTREE2[89] was used with parameter
"-alrt 1000 -bb 1000" to calculate a phylogenetic tree. To calculate a phylogenetic
tree of the upstream neighboring region (scf1:566,786,116-566,788,796) data from
target enrichment for males and females was used as described above.

Phenotypic sexing in multiple species
Phenotypic sex of adult toads of multiple diploid Eurasian green toad taxa had been
established during the breeding period when males exhibited secondary sexual

characters like nuptial excrescences and thorny glandular tips on the back, or were
caught producing male advertisement calls and/or male release calls. Females were
identified as filled with eggs, caught in amplexus, and/or while spawning, producing
female release calls and/or lacking the male secondary sexual characters. During
the spawning migrations, several road kill toads also allowed for anatomical
identification (possession of ovaries or testes) of phenotypic sex.
652

Testing enhancer activity of the Y-haplotype

654 We used the AI-based predictor "iEnhancer-Deep" in a 200 bp "sliding-window"
655 approach in the ncRNA region to test its *in-silico* potential as enhancers of *bod1l-*
656 *expression*[90].
657

Development gene expression in sibling tadpoles from multiple locations

In total 54 transcriptomes of sibling tadpoles comprising brothers and sisters (plus
three unsexed larvae) of four genetic families from four different geographic locations
as well as gonadal tissue of a subadult male and adult female from two additional
localities were examined by RNAseq (transcriptomics). Tadpoles presented five
developmental stages from untreated control groups of a previous experiment[91].
They were euthanized in MS222 and flash-frozen in liquid nitrogen. In tadpoles older
than Gosner stage 37, the intestine was removed prior to flash-freezing. In tadpoles
at Gosner stage 43-44, only the body segment that contained the gonads was used
for RNA extraction. In toadlets of Gosner stage 46 (specifically 6 weeks after
metamorphosis), ovaries, testes (separated from Bidder's organ), and several other
organs (brain, heart, liver, intestine) were dissected and stored in RNAlater prior to
flash-freezing. DNA from tailtips, or skin samples of each tadpole or buccal swabs
(juvenile toads) served for genetic sexing using BvXY1-primers. RNAs were
extracted using the TRIzol Reagent (Thermo Fisher Scientific, Waltham, USA)
according to the supplier's recommendation and cleaned using the RNeasy Mini Kit
(Qiagen). All RNAs included in this study exhibited RIN values >9.0. Library
processing and RNAseq were carried out by NOVOGENE (Cambridge, UK) on a
NovaSeq 6000 PE 150, generating 9 Gb of raw data per sample.
Transcriptome sequences of 54 male and female juvenile developmental stages and
two adult gonads were normalized and mapped to the genome using the nf-core
RNA-Seq pipeline in STAR aligner (https://github.com/alexdobin/STAR/releases,—

quantMode GeneCounts). A list comprising genes that play roles in sexual
development and/or sex determination in other vertebrates (also known as 'usual
suspects'[1]) is shown as Supplementary File 6). To plot *bod1l* and *ncRNA-Y*
expression trends over the developmental stages and separate expression analysis
of *bod1l-X* and *bod1l-Y,* we mapped all reads, including the male haplotype, to the
female assembly. As the *bod1l-Y* gene model is 3'-partial, we used the
corresponding parts of the *bod1l-X* gene model for quantification. Counts of reads
matching the different haplotypes' gene models were extracted using
FEATURECOUNTS (from SUBREAD-2.0.3 package[92] with mapping quality (MQ)
higher than 19, to increase haplotype specificity. Read counts were normalized by
total mapped reads of each sample (TPM normalization). Average expression and
standard deviation were calculated and plotted combining samples of the different
stages using Microsoft Excel software.
693

cDNA of developmental stages

Using the iScript RT Supermix for RT-qPCR kit (Biorad), 10 µL of RNA
(concentration of 80 ng/µL) from all developmental stages (as used for RNAseq)
were randomly reverse-transcribed into cDNA according to the manufacturer's
recommendations. To test allele-specific *bod1l*-gene expression, we designed
primers (bod1l_RNA1F: 5'_AACTGGAAGAAAAGCGCAAA_3'; bod1l_RNA1R:
5'_GTGCTCTTGGCATTCTGTGA_3') and amplified 577 bp from cDNA under the
same conditions as for BvXY1 that were Sanger-sequenced in both directions. To
test expression of the ncRNA-Y, we used primers designed to match its inter-exonic
regions NcRNAY_Ex1to2_F: 5'_CATGAGGAGAGCAGTAATGAACC_3',
NcRNAY_Ex3to2_R: 5'_GCTGTCATCAACCTGCTCAA_3' and amplified and
visualize 204 bp male-specific products from cDNA under the same conditions as for
BvXY1 but at 55°C annealing temperature.
707

Probe preparation for RNA in-situ hybridization

The sequence marker obtained by cDNA amplification using bod1l_RNA1F/R
primers was also used as a probe for RNA *in-situ* hybridization on histological gonad
sections of various developmental stages. As a probe of a control gene, expressed
in gonads of both sexes, we used a 501 bp cDNA fragment of *Vasa* (DDX4-DEAD-
Box-Helicase-4, VASA-homolog), amplified with the primers Vasa1F:

5'_AACCAAAAAGAAGGCGGATT_3';   Vasa1R:

5'_ATGCCAGGATTGCTGTAACC_3' under the same conditions as for BvXY1 but

with annealing temperature of 57°C.

***RNA in-situ* hybridization and H3K4me antibody staining**

We opened the body cavity of freshly euthanized tadpoles of three developmental

stages and fixed them in 4% paraformaldehyde (PFA) buffered in 1x PBS

(Thermofisher), pH = 7.0, at 4°C overnight. Samples were washed in filtered 1×

phosphate-buffered saline with tween 20 (PBST, ROTI®Stock) twice for 5 min, then

dehydrated in a series of methanol (MetOH) with shaking at room temperature (25%

MetOH in PBST for 10 min to 1 h, 50% MetOH in PBST for 10 min to 1 h, 75%

MetOH in PBST for 10 min to 1 h, 100% MetOH for 10 min to 1h, timing dependent

on tissue sample size). Samples were stored in 100% MetOH at -20°C until

histological sectioning.

Cryosectioning, RNA *in-situ* hybridization and immunohistochemistry were performed

as described[93] with slight modifications. Transverse cryosections (20 µm) were

collected on SuperFrost Plus glass slides (Menzel Gläser J1800AMNZ) and stored at

-20°C for up to 2 months. To generate *bod1l* and *vasa* riboprobes, primers

Bod1_RNA_Bv_1F (5'_AACTGGAAGAAAAGCGCAAA_3') and Bod1_RNA_Bv_1R

(5'_GTGCTCTTGGCATTCTGTGA_3') were used to amplify a 577 bp *bod1l*

sequence and primers vasa_1F (5'_AACCAAAAAGAAGGCGGATT_3') and

vasa_1R (5'_ATGCCAGGATTGCTGTAACC_3') were used to amplify a 501 bp *Vasa*

sequence from Gosner 36-37 *B. viridis* cDNA. These sequences were then ligated to

a pCR-bluntII-TOPO vector (Invitrogen) and plasmids were purified using a plasmid

DNA miniprep kit according to manufacturer's instructions (Promega A1460). For

synthesis of sense and antisense riboprobes, plasmids were linearized using *BamHI*

or *NotI* (New England Biolabs) and *in vitro*-transcribed using T7 or SP6 polymerase

(DIG RNA Labeling Mix, Roche), respectively. Rabbit polyclonal anti-histone H3 (tri

methyl K4) primary antibody (Abcam ab8580) and secondary goat anti-rabbit Alexa

Fluor 568 (1:500, Invitrogen A-11011) were used. Sections were counterstained with

0.25 µg/mL DAPI and mounted in Mowiol 4-88 (Calbiochem). Imaging was

performed using the Nikon SMZ18 stereomicroscope and Olympus FluoView

FV3000 confocal microscope. To process images, Bitplane Imaris and Adobe

Photoshop were used.

FISH-TSA

As a probe for fluorescence *in-situ* hybridization with tyramide signal amplification

(FISH-TSA), we designed novel primers to amplify a longer portion of *bod1l* from

cDNA (*bod1l*_Exon7To10_F: 5′_AACTGGAAGAAAAGCGCAAA_3′,

*bod1l*_Exon7To10_R: 5′_TGTGTTGTCTGGAATCACTGG_3′; same conditions as

for BvXY1 but with annealing Temperature of 60°C.), exhibiting the highest

heterozygosity in the coding segment of the gene in *B. viridis* males. A PCR-product

was produced using the PPP Master Mix (Top-Bio, Czech Republic) according to

manufacturer's instructions and purified[94]. The cDNA amplicons were cloned using

the TOPO-TA cloning kit and DH5α-T1R Competent Cells (Thermo Fisher Scientific).

Plasmid DNA was isolated from bacterial colonies using the EZNA Plasmid DNA

Mini Kit I (Omega Bio-Tek Inc.), sequenced, and tested to contain the correct insert

by BLAT search on the green toad genome browser (see Data availability). The

sequence of the *bod1l* clone was compared to the reference genome. Subsequently,

the *bod1l* locus was re-amplified and purified[94]. A 20-year-old tissue suspension

(bone marrow), obtained from a male European green toad from Germany, treated

with colchicine and fixed in methanol-acetic acid[31] was dropped onto a microscope

slide[95], followed by FISH. The probe was labelled with digoxigenin-11-dUTP (Jena

Bioscience) using DecaLabel DNA Labelling Kit (Thermo Fisher Scientific). The

FISH-TSA followed[96] with minor modifications[97]. Mitotic metaphase spreads were

counterstained with ProLongTM Diamond Antifade Mountant with the fluorescent

4',6-diamidino-2-phenylindole, DAPI stain (ThermoFisher Scientific). Microscopy

using Leica Microsystem (Wetzlar, Germany) and processing of metaphase images

followed[98].

Analysis of potential alternative splicing of bod1l

775 For the entire *bod1l* region, including the 3'- and 5-' UTR-regions (scf1:566495375-

776 567417162), aligned RNAseq-reads were extracted from each RNA-seq alignment

777 file, the splice site locations were retrieved and counted. Using all the splice sites, a

778 reference file of them was built. The splice event counts were merged for all samples

779 using the reference and stored as csv text file (Supplementary file 7).

Structural evaluation of the bod1l-*X and Y protein copies*

We used Alphafold2[99] via its cloud-based application and its local installation

ColabFold[99] and analyzed the structure predictions by RaptorX[100] for structural and

potential functional amino-acid differences between the X and the Y-copy of *B.*

viridis. The molecular data viewer iCn3D[101] was used to inspect the structures and

highlight specific amino acid residues (Supplementary text 1).

AmpliSeq custom DNA panel

After providing pre-publication access to the *B. viridis* genome-browser, the Illumina

concierge design service team developed two primer pools covering ca. 80 kb of the

bod1l-region (scf1:566,785,000-566,865,000) and 10 kb for a control gene (*amh*),

known to play roles in sexual development and sex determination in other

vertebrates and lying on the same scaffold (scf1:324181812-324192250). Overall,

the theoretical coverage was 92.3% for *bod1l*, and 98.2% for *Amh*.

Wet lab approaches followed the "AmpliSeq for Illumina On-Demand, Custom, and

Community Panels" (Document #1000000036408 v09; pp. 5-23[102]) for two primer

pools with 193-384 primer pairs in the standard workflow. Briefly, this included ten

steps, (i) quantification of DNA samples of multiple sexed green toads (QuantiFluor®

dsDNA System kit, Promega), (ii) target amplification in two separate wells for the

two primer pools, (iii) partial digestion of amplicons, (iv) ligation of individual

indices/sample, (v) library clean up and (vi) amplification, (vii) 2nd clean up, (viii)

checking libraries on the Bioanalyser for expected insert size and concentration

(QuantiFluor® dsDNA System kit, Promega), (ix) dilution and mixing, and (x)

sequencing, performed by Novogene on a NovaSeq PE250 for individual libraries

obtained for 54 diploid green toads (ca. 100 Mio bp raw data per library).

We included Palearctic green toads by the following numbers, taxa and sexes: 8 *B.*

siculus (4 ♂♂, 3 ♀♀), 9 *B. viridis* (5 ♂♂, 4 ♀♀), 8 *B. balearicus* (5 ♂♂, 3 ♀♀), 9 *B.*

shaartusiensis (4 ♂♂, 5 ♀♀), 10 *B. turanensis* (4 ♂♂, 5 ♀♀). Species' nomenclature

follows[32].

The sequence reads from target enrichment were mapped against the reference

genome using MINIMAP2[74] (-I 100G -a -x sr -t 4). SAMTOOLS[83] was used to add

read group and sample names, to fix mates and sort the bam file output. All BAM

files were merged and variant calling for the target regions was performed by

SAMTOOLS and BCFTOOLS[87] with parameters to cope for extreme sequencing

depth (samtools view -bu input.bam scf1:324181812-324192250 scf1:566783149-

566866298 | bcftools mpileup --annotate DP4 --max-depth 1000000000 -f
bufVir1d.fa - | bcftools call -mv -Ob -o calls.bcf). The output was converted to vcf
format and summarized by an awk script that filtered SNP position and calculated
allele frequency from DP4 annotation for each sample. The resulting table was
further inspected in Excel and SNPs that were heterozygous in males while
homozygous in females were filtered. To fine map the region of the sex marker male
specific SNPs of the different species were intersected (resulting in male specific
SNPs present in 2, 3, 4, 5 species) and plotted along the target regions in our
genome browser.
825
826

Data availability
All data generated in this study have been deposited in publicly accessible
databases. Genome assembly, whole genome sequencing (WGS) data and RNA-
seq reads have been submitted to GenBank on 25 February 2022 (submission ID:
SUB11111520, Bioproject: PRJNA292872, Biosample: SAMN26135295).
Alternatively, the IGB-Green Toad genome browser http://genomes.igb-
berlin.de/Greentoad/ provides easy access to the assembly and its annotation.
834

835 **Code availability**
836 Scripts and parameters are deposited (upon manuscript acceptance) at
837 https://github.com/HMPNK/BUFVIR.
838

839 **References:**
840

1. Herpin, A. & Schartl, M. Plasticity of gene-regulatory networks controlling sex
determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO*
Rep **16**, 1260-74 (2015).
2. Capel, B. Vertebrate sex determination: evolutionary plasticity of a fundamental
switch. *Nat Rev Genet* **18**, 675-689 (2017).
3. Bachtrog, D. *et al.* Sex determination: why so many ways of doing it? *PLoS Biol* **12**,
e1001899 (2014).
4. Rovatsos, M. *et al.* Conservation of sex chromosomes in lacertid lizards. *Mol Ecol* **25**,
3120-6 (2016).
5. Beukeboom LW, Perrin N. The evolution of sex determination. (Oxford, UK: Oxford
University Press, 2014).
6. Volff, J.N., Nanda, I., Schmid, M. & Schartl, M. Governing sex determination in fish:
regulatory putsches and ephemeral dictators. *Sex Dev* **1**, 85-99 (2007).

7.   Perrin, N. Sex reversal: a fountain of youth for sex chromosomes? *Evolution* **63**,
3043-9 (2009).
8.   Bertho, S., Herpin, A., Schartl, M. & Guiguen, Y. Lessons from an unusual vertebrate
sex-determining gene. *Philos Trans R Soc Lond B Biol Sci* **376**, 20200092 (2021).
9.   Kuhl, H. *et al.* A 180 Myr-old female-specific genome region in sturgeon reveals the
oldest known vertebrate sex determining system with undifferentiated sex
chromosomes. *Philos Trans R Soc Lond B Biol Sci* **376**, 20200089 (2021).
10.   Kostmann, A., Kratochvíl, L. & Rovatsos, M. Poorly differentiated XX/XY sex
chromosomes are widely shared across skink radiation. *Proc Biol Sci* **288**, 20202139
(2021).
11.   Pan, Q. *et al.* The rise and fall of the ancient northern pike master sex-determining
gene. *Elife* **10** (2021).
12.   Ito, M. Sex Determination and Differentiation in Frogs. in *Reproductive and*
Developmental Strategies: The Continuity of Life (eds. Kobayashi, K., Kitano, T.,
Iwao, Y. & Kondo, M.) 349-366 (Springer Japan, Tokyo, 2018).
13.   Schmid, M. & Steinlein, C. Sex chromosomes, sex-linked genes, and sex
determination in the vertebrate class Amphibia. in *Genes and Mechanisms in*
Vertebrate Sex Determination (eds. Scherer, G. & Schmid, M.) 143-176 (Birkhäuser
Basel, Basel, 2001).
14.   Schartl, M., Schmid, M. & Nanda, I. Dynamics of vertebrate sex chromosome
evolution: from equal size to giants and dwarfs. *Chromosoma* **125**, 553–571 (2016).
15.   Green, D.M. Cytogenetics of the endemic New Zealand frog, *Leiopelma hochstetteri*:
extraordinary supernumerary chromosome variation and a unique sex-chromosome
system. *Chromosoma* **97**, 55-70 (1988).
16.   Roco Á, S. *et al.* Coexistence of Y, W, and Z sex chromosomes in *Xenopus tropicalis*.
Proc Natl Acad Sci U S A **112**, E4752-61 (2015).
17.   Furman, B.L.S. *et al.* A frog with three sex chromosomes that co-mingle together in
nature: *Xenopus tropicalis* has a degenerate W and a Y that evolved from a Z
chromosome. *PLoS Genet* **16**, e1009121 (2020).
18.   AmphibiaWeb. See www. amphibiaweb.org (accessed June 2023).
19.   Yoshimoto, S. *et al.* A W-linked DM-domain gene, DM-W, participates in primary
ovary development in *Xenopus laevis. Proc Natl Acad Sci U S A* **105**, 2469-74 (2008).
20.   Cauret, C.M.S. *et al.* Developmental Systems Drift and the Drivers of Sex
Chromosome Evolution. *Mol Biol Evol* **37**, 799-810 (2020).
21   Cauret CMS, et al. (2023) Functional dissection and assembly of a small, newly
evolved, W chromosome-specific genomic region of the African clawed frog *Xenopus*
laevis. Plos Genetics **19**(10), e1010990 (2023).
22   Evans, B.J. et al. New insights into *Xenopus* sex chromosome genomics from the
Marsabit clawed frog *X. borealis. J Evol Biol* **35**, 1777-1790 (2022).
23.   Bogart, J.P. Karyotypes. in *Evolution of the genus Bufo* (ed. Blair, W.F.) 171-232
(Austin, Texas: University of Texas Press, 1972).
24.   Schmid, M. Chromosome banding in Amphibia I. Constitutive heterochromatin and
nucleolus organizer regions in *Bufo* and *Hyla. Chromosoma (Berl.)* **66**, 361-388
(1978).
25.   Piprek, R.P., Kloc, M. & Kubiak, J.Z. Bidder's organ--structure, development and
function. *Int J Dev Biol* **58**, 819-27 (2014).
26   Harms, W. Untersuchungen über das Biddersche Organ der männlichen und
weiblichen Kröten. *Z Anat Entwickl Gesch* **69**, 598–629 (1923).
27.   Ponse, K. La proportion sexuelle dans la descendance issue des oeufs produits par
l'organe de Bidder des crapauds femelles. *Rev. Suisse Zool.* **48**, 541–544 (1941).

28. Changxiang, W., Qun, L. & Wei, X. Studies of chromosome banding and sister
chromatid exchange in *Bufo bufo gargarizans*. *Acta Gen Sinica* **10**, 291–297 (1983).

29. Abramyan, J., Ezaz, T., Graves, J.A. & Koopman, P. Z and W sex chromosomes in
the cane toad (*Bufo marinus*). *Chromosome Res* **17**, 1015-24 (2009).

30. Malone, J.H. & Fontenot, B.E. Patterns of reproductive isolation in toads. *PLoS One*
3, e3900 (2008).

31. Stöck, M., Steinlein, C., Lamatsch, D.K., Schartl, M. & Schmid, M. Multiple origins
of tetraploid taxa in the Eurasian *Bufo viridis* subgroup. *Genetica* **124**, 255-72 (2005).

32. Ueda, H. Offspring of sex-reversed males in *Bufo viridis*. *Sci. Rep. Lab. Amphib. Biol.*
Hiroshima. Univ. **10**(1990).

33. Stöck, M. *et al.* Low rates of X-Y recombination, not turnovers, account for
homomorphic sex chromosomes in several diploid species of Palearctic green toads
(*Bufo viridis* subgroup). *J Evol Biol* **26**, 674-82 (2013).

34. Brelsford, A. *et al.* Homologous sex chromosomes in three deeply divergent anuran
species. *Evolution* **67**, 2434-2440 (2013).

35. Tamschick, S. *et al.* Sex chromosome conservation, *DMRT1* phylogeny and gonad
morphology in diploid Palearctic green toads (*Bufo viridis* subgroup). *Cytogenet*
Genome Res **144**, 315-324 (2014).

36. Betto-Colliard, C., Hofmann, S., Sermier, R., Perrin, N. & Stöck, M. Profound genetic
divergence and asymmetric parental genome contributions as hallmarks of hybrid
speciation in polyploid toads. *Proc Biol Sci* **285,** 20172667 (2018).

37. Burgin, C. *et al.* Mammal Diversity Database (Version 1.11) dataset. Zenodo
http://dx.doi.org/10.5281/zenodo.7830771 (2023).

38. NCBI. See www.ncbi.nlm.nih.gov (accessed June 2023).

39. Guzmán, K. *et al.* Identification and characterization of a new family of long satellite
DNA, specific of true toads (Anura, Amphibia, Bufonidae). *Scientific Reports* **12**,
13960 (2022).

40. Stöck M., et al. (2006): Evolution of mitochondrial relationships and biogeography of
Palearctic green toads (*Bufo viridis* subgroup) with insights in their genomic
plasticity. *Mol Phylogen Evol* **41**, 663-689 (2006).

41. Dufresnes C., *et al.* Phylogeography of Aegean green toads (*Bufo viridis* subgroup):
continental hybrid swarm vs. insular diversification with discovery of a new island
endemic. *BMC Evol Biol* **18**, 67. (2018).

42. Schartl, M. & Herpin, A. Sex Determination in Vertebrates. in *Encyclopedia of*
Reproduction (Second Edition) (ed. Skinner, M.K.) 159-167 (Academic Press,
Oxford, 2018).

43. Gosner, K.L. A simplified table for staging anuran embryos and larvae with notes on
identification. *Herpetologica*, **16**, 183–190 (1960).

44. Burren, O.S. *et al.* Chromosome contacts in activated T cells identify autoimmune
disease candidate genes. *Genome Biol* **18**, 165 (2017).

45. Jackson, D.A., Dickinson, P. & Cook, P.R. The size of chromatin loops in HeLa cells.
EMBO J **9**, 567-571 (1990).

46. Lyu, J., Shao, R., Kwong Yung, P.Y. & Elsässer, S.J. Genome-wide mapping of G-
quadruplex structures with CUT&Tag. *Nucleic Acids Res* **50**, e13 (2022).

47. Bayley, R. *et al.* H3K4 methylation by SETD1A/BOD1L facilitates RIF1-dependent
NHEJ. *Mol Cell* **82**, 1924-1939.e10 (2022).

48. Giovanni, C. *et al.* The BOD1L subunit of the mammalian SETD1A complex sustains
the expression of DNA damage repair genes despite restraining H3K4 trimethylation.
Preprint at https://www.biorxiv.org/content/10.1101/2023.04.06.535882v2 (2023).

49. Hayashi, S. *et al.* Neofunctionalization of a Noncoding Portion of a DNA Transposon
in the Coding Region of the Chimerical Sex-Determining Gene *dm-W* in *Xenopus*
Frogs. *Mol Biol Evol* **39**, msac138 (2022).

50. Burgos, M., *et al.* Non-coding RNAs: lncRNAs, miRNAs, and piRNAs in sexual
development. *Sex Dev*, **15**, 335–350 (2021).

51. Joshi M, & Rajender S. Long non-coding RNA (lncRNAs) in spermatogenesis, male
infertility. *Reprod Biol Endocrinol* **18**, 103 (2020).

52. Feng B., et al. lncRNA DMRT2-as acts as a transcriptional regulator of *dmrt2*
involving in sex differentiation in the Chinese tongue sol (*Cynoglossus semilaevis*).
Comp Biochem Physiol B Biochem Mol Biol **253**, 110542 (2021).

53. Bowles, J. & Koopman, P. Retinoic acid, meiosis and germ cell fate in mammals.
Development **134**, 3401-11 (2007).

54. Higgs, M.R. *et al.* *BOD1L* Is Required to Suppress Deleterious Resection of Stressed
Replication Forks. *Mol Cell* **59**, 462-77 (2015).

55. Higgs, M.R. *et al.* Histone Methylation by *SETD1A* Protects Nascent DNA through
the Nucleosome Chaperone Activity of *FANCD2*. *Mol Cell* **71**, e6, 25-41 (2018).

56. Rodríguez-Marí, A. & Postlethwait, J.H. The role of Fanconi anemia/BRCA genes in
zebrafish sex determination. *Methods Cell Biol* **105**, 461-90 (2011).

57. Smallwood, S.A. & Kelsey, G. *De novo* DNA methylation: a germ cell perspective.
Trends Genet **28**, 33-42 (2012).

58. Youds, J.L. & Boulton, S.J. The choice in meiosis – defining the factors that influence
crossover or non-crossover formation. *Journal of Cell Science* **124**, 501-513 (2011).

59. Ki, B.S. *et al.* Epigenetic regulator Cfp1 safeguards male meiotic progression by
regulating meiotic gene expression. *Experimental & Molecular Medicine* **54**, 1098-
1108 (2022).

60. Fan, H.-Y. & Sun, Q.-Y. Chapter 12 - Oocyte Meiotic Maturation. in *The Ovary*
(Third Edition) (eds. Leung, P.C.K. & Adashi, E.Y.) 181-203 (Academic Press,
2019).

61. Dupont, S. & Capel, B. The Chromatin State during Gonadal Sex Determination. *Sex*
Dev **15**, 308-316 (2021).

62. Garcia-Moreno, S.A. *et al.* CBX2 is required to stabilize the testis pathway by
repressing Wnt signaling. *PLoS Genet* **15**, e1007895 (2019).

63. Nakamura, M. Sex determination in amphibians. *Semin Cell Dev Biol* **20**, 271-82
(2009).

64. Wolf, J.C. *et al.* Effects of 17β-estradiol exposure on *Xenopus laevis* gonadal
histopathology. *Environ Toxicol Chem* **29**, 1091–1105 (2010).

65. Peng, Y., Leung, H.C., Yiu, S.M. & Chin, F.Y. IDBA-UD: a de novo assembler for
single-cell and metagenomic sequencing data with highly uneven depth.
Bioinformatics **28**, 1420-8 (2012).

66. Ruan, J. & Li, H. Fast and accurate long-read assembly with *wtdbg2*. *Nat Methods* **17**,
155-158 (2020).

67. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P.A. Assembly of long, error-prone
reads using repeat graphs. *Nat Biotechnol* **37**, 540-546 (2019).

68. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-8
(2011).

69. Durand, N.C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-
Resolution Hi-C Experiments. *Cell Syst* **3**, 95-8 (2016).

70. Dudchenko, O. *et al.* *De novo* assembly of the *Aedes aegypti* genome using Hi-C
yields chromosome-length scaffolds. *Science* **356**, 92-95 (2017).

1002	71.	Durand, N.C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps
1003		with Unlimited Zoom. *Cell Syst* **3**, 99-101 (2016).
1004	72.	Kent, W.J. *et al.* The human genome browser at UCSC. *Genome Res* **12**, 996-1006
1005		(2002).
1006	73.	Gerchen, J.F. *et al.* A single transcriptome of a green toad (*Bufo viridis*) yields
1007		candidate genes for sex determination and -differentiation and non-anonymous
1008		population genetic markers. PLoS One **11**, e0156419 (2016).
1009	74.	Li, H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics **34**,
1010		3094-3100 (2018).
1011	75.	Fan Nie, *et al. De novo* diploid genome assembly using long noisy reads via
1012		haplotype-aware error correction and inconsistent overlap identification. bioRxiv
1013		2022.09.25.509436; https://doi.org/10.1101/2022.09.25.509436 (2022).
1014	76.	Zhang, H. et al. Fast alignment and preprocessing of chromatin profiles with
1015		Chromap. Nat Comm **12,** 6566 (2021).
1016	77.	Zhou C., McCarthy S.A. & R. Durbin YaHS: yet another Hi-C scaffolding tool,
1017		Bioinformatics **39**, btac808 (2023).
1018	78.	Kolmogorov M, et al. Chromosome assembly of large and complex genomes using
1019		multiple references. *Genome Research* **28**, 1720-1732 (2018).
1020	79.	Brelsford, A., Dufresnes, C., & Perrin, N. High-density sex-specific linkage maps of a
1021		European tree frog (*Hyla arborea*) identify the sex chromosome without information
1022		on offspring sex. Heredity, **116**, 177–181 (2016).
1023	80.	Rochette N.C., Rivera-Colón A.G. & J.M. Catchen. Stacks 2: Analytical methods for
1024		paired-end sequencing improve RADseq-based population genomics. *Mol Ecol Res*
1025		**28**, 4737–4754. (2019).
1026	81.	Feron, R. *et al.* RADSex: A computational workflow to study sex determination using
1027		restriction site-associated DNA sequencing data. *Mol Ecol Resour* **21**, 1715-1731
1028		(2021).
1029	82.	https://github.com/RomainFeron/paper-sexdetermination-greentoad-2021.
1030	83.	Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
1031		2078-2079 (2009).
1032	84.	Rimmer, A. *et al.* Integrating mapping-, assembly- and haplotype-based approaches
1033		for calling variants in clinical sequencing applications. *Nat Genet* **46**, 912-918 (2014).
1034	85.	Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing
1035		genomic features. *Bioinformatics* **26**, 841-2 (2010).
1036	86.	Sahlin, K., Lim, M.C.W. & Prost, S. NGSpeciesID: DNA barcode and amplicon
1037		consensus generation from long-read sequencing data. *Ecol Evol* **11**, 1392-1398
1038		(2021).
1039	87.	Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10,** giab008
1040		(2021).
1041	88.	Castresana, J. Selection of conserved blocks from multiple alignments for their use in
1042		phylogenetic analysis. *Mol Biol Evol* **17**, 540-52 (2000).
1043	89.	Minh, B.Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
1044		Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020).
1045	90.	Kamran, H., Tahir, M., Tayara, H. & Chong, K.T. iEnhancer-Deep: A Computational
1046		Predictor for Enhancer Sites and Their Strength Using Deep Learning. *Applied*
1047		*Sciences* **12**, 2120 (2022).
1048	91.	Tamschick S., *et al.* Sex reversal assessments reveal different vulnerability to
1049		endocrine disruption between deeply diverged anuran lineages. *Scientific Reports* **6**,
1050		23825 (2016).

92. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-30 (2014).
93. Dasyani, M. *et al.* Lineage tracing of col10a1 cells identifies distinct progenitor
populations for osteoblasts and joint cells in the regenerating fin of medaka (*Oryzias*
latipes). *Dev Biol* **455**, 85-99 (2019).
94. Knytl, M. & Fornaini, N.R. Measurement of chromosomal arms and FISH reveal
complex genome architecture and standardized karyotype of model fish, genus
Carassius. *Cells* **10**(2021).
95. Courtet, M., Flajnik, M. & Du Pasquier, L. Major histocompatibility complex and
immunoglobulin loci visualized by in situ hybridization on *Xenopus* chromosomes.
Dev Comp Immunol **25**, 149-57 (2001).
96. Krylov, V., Tlapakova, T. & Macha, J. Localization of the single copy gene Mdh2 on
Xenopus tropicalis chromosomes by FISH-TSA. *Cytogenet Genome Res* **116**, 110-2
(2007).
97. Knytl, M., Tlapakova, T., Vankova, T. & Krylov, V. *Silurana* chromosomal
evolution: a new piece to the puzzle. *Cytogenet Genome Res* **156**, 223-228 (2018).
98. Seroussi, E. *et al.* Avian expression patterns and genomic mapping implicate leptin in
digestion and TNF in immunity, suggesting that their interacting adipokine role has
been acquired only in mammals. *Int J Mol Sci* **20**, 4489 (2019).
99. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods*
19, 679-682 (2022).
100. Peng, J. & Xu, J. RaptorX: exploiting structure information for protein alignment by
statistical inference. *Proteins* **79 Suppl 10**, 161-71 (2011).
101. Wang, J. *et al.* iCn3D, a web-based 3D viewer for sharing 1D/2D/3D representations
of biomolecular structures. *Bioinformatics* **36**, 131-135 (2020).
102. https://support.illumina.com/downloads/ampliseq-for-illumina-custom-and-
community-panels-reference-guide-1000000036408.html.
1078

Acknowledgements

Funding has been in part provided by the German Research Foundation (DFG) to MStöck and HK (STO493/8), and the Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB). WKloas and MStöck also received funding from the European Union's Horizon 2020 research and innovation programme, grant agreement No. 825753 (ERGO). FISH-TSA experiments were partly covered by the PJAC-project CZ.02.01.01/00/22_010/0002902, MSCA Fellowships CZ–UK (MK). We thank Petros Lymberakis, Natural History Museum of Crete, University of Crete, for help with the sampling and export permits of green toads by the Ministry of the Environment of Greece (115790/229) 2014-2017. The tissue and DNA collection of MStöck profited from donations of multiple international colleagues over 25 years; he thanks everyone. We are grateful to Sandra Bittmann (IGB) for HMW-DNA extraction, Eva Kreuz (IGB) for initial efforts to clone the SD-region, Julia Bindl and Nathalie Granath (IGB) for help with the DNA and RNA work, Hughes Parinello (Plateforme MGX) for preparing the HiC-libraries and sequencing, the BeGenDiv (Berlin, Germany) for

1094 providing lab space and Illumina-sequencing support for the ddRAD. We thank
1095 Sebastian Petri (Biozyme) for discussions on high-resolution agarose gel-
1096 electrophoresis.

1097

Authors' contributions:
HK: Contributed to study design, sequenced long reads, assembled and annotated
the genome, set up the genome browser, mapped and analyzed pool-seq sequences
and SNPs of the AmpliSeq custom DNA panel, performed phylogenetic analyses,
contributed to 1st manuscript draft
TWH: Made histological sections and performed all RNA *in-situ* hybridizations and
H3K4me immunostainings, contributed to 1st manuscript draft
CK: Analyzed all transcriptomes, and gene expression data, analyzed splice sites of
bod1l
WKleiner: Performed RNA extraction, bioanalyzer quantification and evaluation, did
most marker testing by PCRs, co-worked in the wet lab for the AmpliSeq custom
DNA panel
BK: Performed molecular work to characterize the ncRNA of multiple green toad
taxa, prepared RNA for transcriptomes and cDNA-work at Gosner stage 46
MC: Contributed to ddRAD wet lab
RF: Analyzed the ddRADseq data
MK: Performed FISH-TSA
WKloas: Contributed to study design, discussed developmental histology and data.
MSchartl: Advised in many respects, initiated some collaborations, discussed data
and study progress and results.
CW: Contributed to study design, supervised and analyzed RNA *in-situ* hybridization
and H3K4me immunostainings.
MStöck: Conceived the study, did all field work/sampling, raised tadpoles, prepared
and fixed tissues for chromosome and RNA *in situ* work, collected buccal swabs and
tissues over 25 years, sexed toads, extracted all DNAs, performed ddRAD wet lab,
developed all primers, did some marker testing by PCRs, co-worked in the wet lab
for the AmpliSeq custom DNA panel, wrote the 1st manuscript draft.
All authors contributed text to the manuscript, and read and improved the final
version.

1127

1128   **Competing interests**

1129   The authors declare to have no competing interests.

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142   **Figures**

1143
Figure 1. The chromosome-scale reference-genome of *Bufo*(*tes*) *viridis* and the sex-diagnostic
marker. a, Breeding pair of *B. viridis* in amplexus (© M. Stöck). **b,** Giemsa-stained karyotype of *B.*
viridis, changed from[31]. **c,** above: Mapping the pool-seq short reads to the *B. viridis* reference:
genome-wide a single region on scaffold 1 (scf1:566,783,058-566,848,882), where the number of
heterozygous SNPs (y-axis) on scaffold 1 (N = 103) in males surpassed about nine times the
maximum number of genomic positions (N = 12) where females exhibit heterozygous SNPs, provides
strong evidence for male heterogamety (XX/XY); for comparison with a ZZ/ZW system see
Supplementary Fig. 3; **c,** middle: male heterozygous SNPs on scaffold 1 of the male genome
assembly identifies the same region (note a slight shift of the peak showing the same locus due to
different reference genomes); **c,** below: male-specific pool-seq coverage on the male genome (for
these analyses on the entire female and male genomes: Supplementary file 2); y-axis for
heterozygous SNPs are counts of SNPs (above and middle subfigure) or Y-specifically pool-sex
covered base pairs (lower subfigure) in a cluster, where the maximal distance of variants is 9000 bp.

d, *B. viridis*-assembly (y-axis) compared to *B. bufo (x-axis)* assembly by the Vertebrate Genome

Project (VGP, GenBank accession number GCA_905171765.1) showing high synteny of both bufonid

genomes across all 11 chromosomes. **e,** Overview of sex-specific PCR-products of primers BvXY1 in

females (♀♀, left) and males (♂♂, right) of six (i-vi) nominal diploid green toad taxa, electrophoresis

result using 4 µL of PCR-products on a 4% high-resolution Sieve-agarose gel run for 60 min on 70 V;

Palearctic green toad taxa shown are: (i) *siculus*, (ii) *shaartusiensis (note, the highest band in males*

is weak), (iii) *viridis*, (iv) *variabiilis*, (v) *balearicus*, (vi) *turanensis*. **f,** FISH-TSA on a metaphase spread

from a *B. viridis* male with red *bod1l*-signals on chromosomes 1 on their short (p) arm (arrows),

chromosomes were counterstained with DAPI (blue-green), scale bar in lower left corner represents

10 µm. **g,** Range map of Palearctic green toad taxa, numbers (i-vi) and color-coding as in (e) and (h),

diploid (2n) taxa are indicated by circles, triploids (3n) by triangles and tetraploids (4n) by squares. **h,**

Dated mitochondrial phylogeny of Palearctic green toads (adapted from[36], with the phylogenetic

positions of the diploid taxa, in which BvXY1 is sex-diagnostic, indicated by color-coded arrows as in

(**e**); the three ploidy levels are provided to the right of the leaves, note that the sex marker is

seemingly conserved only in the diploid green toads; all polyploids are of hybrid origin[36].

1172

1173

Figure 2. Additive sex-specific XX- or XY-expression of *bod1l* and expression of the ncRNA-Y

in females and males in seven developmental stages, with numbers (N) of females and males

(N) per stage shown on bars, genetically sexed using PCR-marker BvXY1. Gosner[43] stages 23-

24, 10 days after fertilization, Gosner stages 27-29: 15 days after fertilization, Gosner stages 30-33:

18 days after fertilization, Gosner stages 34-38: 34 days after fertilization, Gosner stages 43-44:

beginning of metamorphosis, Gosner stages 45-46, up to six weeks after metamorphosis, the latter

with anatomically visible differentiation of testes and Bidder's organs in males, and ovaries in females,

subadult male and adult female. Numbers of available tadpole transcriptomes varied by stage; TPM:

Transcript reads Per Million RNAseq reads; note that TPM for the ncRNA-Y were multiplied 20x. See

Supplementary Fig. 8 for comparison of additive sex-specific expression of *bod1l* and ncRNA-Y and

haplotype-specific *bod1l*-expression in males.

1185

1186

1187

Figure 3. Structure of the X- and Y-haplotype of the candidate sex locus of *Bufo*(*tes*) *viridis*

with focus on the 5'-end of *bod1l* and the Y-specific ncRNA. a, X-chromosomal (above, pink) and

Y-chromosomal haplotype (below, blue) with color-coded homologous or X- and Y-specific INDELs,

and the Y-specific long (ca. 5.8 kb) non-coding RNA (ncRNA-Y), flanking a putative enhancer at the

5'-end of the gene, ca. 80 kilobases before its promoter. **b,** Graph of the signal of the AI-based

software *i-enhancer*, reaching from 0 to 2, predicting strong enhancer properties of the Y-specific 5'-

end of *bod1l*, including a G-quadruplex-rich region on the non-template strand, overlapping with the

ncRNA-Y. **c,** Y- and thus male-specific expression of the ncRNA-Y (204 bp PCR-product from cDNA)

in larval green toads, from 10 days after fertilization (Gosner stages 23-24) until early metamorphosis

(Gosner 46); in females, only non-specific products occurred, since X-copies of *bod1l* do not contain

this ncRNA (for details and positive controls of females: Supplementary Fig. 9). **d,** Unrooted

Maximum-Likelihood-based phylogenetic tree, showing that the ncRNA-Y evolved in a common

ancestor of diploid Palearctic green toads sharing this XY-system, with *siculus* and *viridis* spanning

the most divergent lineages; two PCR-products (2.5 and 5.8 kb), directly amplified by independent

PCRs from six related diploid green toad taxa, clustered by Y-gametologue and thus species (*

symbolized 100% bootstrap support; only support values >50% are shown).

		Expression in somatic cells		Expression in germ cells	
		Male	Female	Male	Female
Gosner 36-37	*Bod1L*	+	−	+	+
Gosner 43-44	*Bod1L*	−	−	+	+
	H3K4me3	low	high	high	high

Figure 4. ***Bod1l*-expression and H3K4me3-methylation in primordial gonads of *B. viridis*, i.e. in the female early ovary, and in male Bidder's organ, contributing to the testes anlage (Gosner stages 36-37) as well as in early metamorphosis (Gosner stages 43-44). a-f**, *Bod1l* expression shown by RNA *in-situ* hybridization (ISH) on 20 μm transverse cryosections of female (a, d) and male (b, c, e, f, h) gonads at developmental stages Gosner 36-37 (a-c) and 43-44 (d-f), the latter presenting early metamorphic stages. DAPI was used to stain nuclei. Magenta asterisks indicate nuclei of germ cells; magenta arrows: cytoplasm of germ cells; yellow arrows: nuclei of somatic cells not stained by *bod1l* RNA ISH; green arrows: nuclei of somatic cells stained by *bod1l* RNA ISH. Yellow boxes in b and e indicate zoomed images shown in c and f, respectively. **g,h,** H3K4me3 and DAPI staining on 20 μm transverse cryosections of female (g) and male (h) green toad Bidder's organ and gonads, Gosner stage 43-44. White asterisks indicate H3K4me3-positive germ cell nuclei; yellow boxes indicate somatic cells. Scale bars (a, b, d, e, g, h): 20 μm; (c, f) 5 μm. **i,** Summary of *bod1l*-expression and H3K4me3-methylation in somatic cells and germ cells in Gosner 36-37 and Gosner 43-44 male and female gonads.

Additional information

Supplementary File 1: Description of Supplementary Data.

Supplementary File 2: Comparison of female and male *B. viridis* genome assemblies.

Supplementary File 3: Analyses of SNPs causing 6 non-synonymous changes in the coding region of *bod1l* in 49 male and female transcriptomes as compared to the female reference genome. None of these changes was 100% associated with genetic sex as in the 5'-region of *bod1l*.

Supplementary File 4: Percentages of covered parts of ca. 80 kb of the *bod1l*-region by targeted sequencing in six diploid green toad taxa, using a panel based on the female *B. viridis* reference genome, compared to ca. 20 kb of the gene for *anti-Muellerian hormone* (*amh*). Taxa abbreviations: Bbal - *balearicus*, Bsha - *shaartusiensis*, Bsic - *siculus*; Btur - *turanensis*, Bvir - *viridis* and *variabilis*.

Supplementary File 5: SNP-comparisons in ca. 80 kb of the *bod1l*- (scf1: 566,783,513 - 566,865,063) and ca. 10 kb of the *amh*-regions (scf1: 324,181,734 – 324,192,327) by targeted sequencing in six diploid green toad taxa in a panel based on the female *B. viridis* reference genome. The coordinates of the cluster of male specific SNPs present in all five species are colored in red (scf1: 566,790,085 - 566,790,204). Values for color-coded SNP-comparisons with the reference genome: >=0.9: homozygous (green/yellow) = identical to reference; between 0.9 and 0.1: heterozygous (orange) = one allele identical to reference; <=0.1: homozygous(red) = completely different from reference; NA: SNP not covered by sequencing data in the relevant individual / taxon; M - male, F - female; position: SNP position on the female *B. viridis* reference genome; taxa abbreviations as in Supplementary File S3.

Supplementary File 6: Boxplots showing expression of 80 genes that play roles in sexual development and/or sex determination in other vertebrates (also known as 'usual suspects'[1], in developmental stages of *B. viridis*. X-axis showing the normalized intensity of expression; Y-axis marks six developmental stages (details as in Fig. 2).

Supplementary File 7: File of splice site read counts in *bod1l*-region.

Supplementary Figures

Supplementary Figure 1: **Analyses of ddRAD markers in three genetic families of *B. viridis* using RADsex[63].** Of the 443,378 markers present in at least one individual with a coverage >5; 96 markers were significantly associated with male sex (p < 0.05 after Bonferroni correction), but none of these markers were found in all males.

Supplementary Figure 2: **Alignment of ddRAD markers shown in Supplementary Figure 1 to the *B. viridis* reference genome.** 64 of the 96 markers significantly associated with sex were located on chromosome 1 (scaffold 1, scf1), one marker was aligned to chromosome 7 (scf7), five markers were aligned to unplaced scaffolds, and 26 markers were not aligned uniquely or with a mapping quality lower than 20.

Supplementary Figure 3: **Mapping pool-sex short reads to the *B. viridis* reference genome with SNP-heterozygosity in males (a) *vs*. females (b) on scaffold (scf) 1.**

Supplementary Figure 4: Male-specific SNPs of 6 species over *bod1l-* region. a SNPs show different blocks of male-specific heterozygosity between species, which exclude coding changes of *bod1l* as sex-determination mechanism. **b** Highest congruence of male-specific SNPs among all species is in the PCR marker/ncRNA-region (red ellipsis). **c** Phylogenetic tree of haplotypes from the marked region shows clustering by gametologue. **d** sequences from the upstream region were clustered by species.

Supplementary Figure 5: Coverage distributions of X/Y haplotypes depending on female/male pool-seq in *B. viridis*. a Y-haplotype, **b** X-haplotype.

Supplementary Figure 6: Female/male pool-seq coverage over haplotypes in *B. viridis*. a female pool-sequencing, **b** male pool-sequencing.

Supplementary Figure 7: X- and Y-haplotype comparisons by dot plots in the *bod1l*-region. a X-*versus*-Y dot plot, **b** Y-*versus*-Y, **c** X-*versus*-X.

Supplementary Figure 8: Comparison of additive sex-specific expression of bod1l and ncRNA-Y and haplotype-specific expression in males.

Supplementary Figure 9: a, Y- and thus male-specific expression of the ncRNA-Y (204 bp PCR-product from cDNA) in larval green toads. PCR-products from cDNA (204 bp), obtained with primers NcRNAY_Ex1to2_F/_R; to the left males,

to the right females; each sex is shown from Gosner stage 23-34 (10 days after
fertilization) until early metamorphosis (Gosner 43-44) and one adult gonad for each
sex; in females, only non-specific products occurred, since X-copies of *bod1l* do not
contain and thus do not express this ncRNA. **b,** Gel image of a control-PCR using
the females from (a).
Supplementary Figure 10: Graph of the signal of the AI-based software i-
enhancer. Signal bars reaching from 0 to 2, predicting strong enhancer properties of
the Y-specific 5'-end of *bod1l*, including a G-quadruplex-rich region on the non-
template strand, overlapping with the ncRNA-Y.
Supplementary Figure 11: *Vasa, bod1l* expression and H3K4me3-methylation
in *B. viridis* gonads, overview images and additional representative examples.
a-m, v*asa* and *bod1L* expression shown by RNA *in-situ* hybridization (ISH) on 20 µm
transverse cryosections of female (**a, b, g, h**) and male (**c, d, f, i-m**) gonads, Gosner
36-37. DAPI was used to stain nuclei. Sense controls for RNA ISH are shown in **a, c,**
e, **g**, right panels. Red boxes in **a, c, g, i** (left panel) indicate zoomed images shown
in **b, d, h, k,** respectively. Yellow boxes in **b, d, h, k** (right panel) indicate zoomed
images shown in **Fig. 4a-b,** respectively. Red boxes in l**, m** (left panel) indicate
zoomed images shown in right panels of l**, m**. *Bod1L* staining on three different
cryosections of male gonads are shown in k**, l, m**. Magenta asterisks indicate nuclei
of germ cells; magenta arrows: cytoplasm of germ cells; yellow arrows: somatic cells
not stained by RNA ISH; green arrows: somatic cells stained by RNA ISH. **n, o,**
H3K4me3 and DAPI staining on 20 µm transverse cryosections of female (**l**) and
male (**m**) *B. viridis* Bidder's organ and gonads, Gosner stage 43-44. Yellow boxes in
l, m indicate zoomed images shown in **Fig. 4g**, **h,** respectively. White asterisks
indicate H3K4me3-positive germ cell nuclei; white arrowheads, H3K4me3-positive
somatic cell nuclei; yellow arrowheads, H3K4me3-negative somatic cell nuclei. Scale
bars: (a, c, g, i) 50 µm, (b, d-f, h, k-o) 20 µm.
1317
Supplementary Tables
Supplementary Table 1: Localities/populations of European green toads that
contributed to the pool-seq experiment.

Supplementary Table 2: Summary of the pool sequencing approach for scaffold 1

(scf1) for SNPs homozygous in males/heterozygous females (Z/W) as well as SNPs

heterozygous in males/homozygous females (X/Y).

Supplementary Table 3: Signal of the AI-based software i-enhancer, reaching from

0 to 2, predicting strong enhancer properties of the Y-specific 5'-end of *bod1l*,

including a G-quadruplex-rich region on the non-template strand, overlapping with

the ncRNA-Y.

1328

1329 **Supplementary Texts**

1330 **Supplementary Text 1:** Potential impact of X- *vs*. Y-*bod1l* coding-sequences in *B.*

1331 *viridis.*