A novel amphibian sex determination candidate gene, evolved by structural variation in the regulatory region between X and Y

Heiner Kuhl[1,*], Wen Hui Tan [2,*], Christophe Klopp[3], Wibke Kleiner[1],

Baturalp Koyun[1,4], Mitica Ciorpac[5,6], Romain Feron[7,8], Martin Knytl[9,10],

Werner Kloas[1], Manfred Schartl[11,12], Christoph Winkler[2,§], Matthias Stöck[1,§]

[1] Leibniz-Institute of Freshwater Ecology and Inland Fisheries, IGB, Müggelseedamm 301 & 310, 12587 Berlin, Germany,

[2] Department of Biological Sciences and Centre for Bioimaging Sciences, National University of Singapore, 14 Science Drive 4, Block S1A, Level 6, Singapore 117543,

[3] SIGENAE, Plate-forme Bio-informatique Genotoul, Mathématiques et Informatique Appliquées de Toulouse, INRAe, 31326 Castanet-Tolosan, France

[4] Bilkent University, Department of Molecular Biology and Genetics, Faculty of Science, SB Building, Ankara, Turkey 06800,

[5] Danube Delta National Institute for Research and Development, Tulcea 820112, Romania,

[6] Advanced Research and Development Center for Experimental Medicine – CEMEX, "Grigore T. Popa", University of Medicine and Pharmacy, Mihail Kogălniceanu Street 9-13, Iasi 700259, Romania,

[7] Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland,

[8] Swiss Institute of Bioinformatics, Lausanne, Switzerland,

[9] Department of Cell Biology, Faculty of Science, Charles University, Viničná 7, Prague, 12843, Czech Republic

[10] Department of Biology, McMaster University, 1280 Main Street West, Hamilton L8S 4K1, Ontario, Canada

[11] Developmental Biochemistry, Biocenter, University of Wuerzburg, Am Hubland, 97074 Wuerzburg, Germany,

[12] The Xiphophorus Genetic Stock Center, Department of Chemistry and Biochemistry, Texas State University, San Marcos, Texas, TX, 78666, USA,

[*] These authors contributed equally to the paper.

[§] Co-corresponding authors.

Abstract

Most vertebrates develop distinct females and males, where sex is determined by repeatedly-evolved environmental or genetic triggers. Undifferentiated sex chromosomes and large genomes have caused major knowledge gaps in amphibians. Only a single master gene is known in >8650 species, the *dmrt1*-paralogue (*dm-w*) of female-heterogametic clawed frogs (*Xenopus*; ZW♀/ZZ♂). Combining a chromosome-scale genome of a non-model amphibian, the European green toad, *Bufo*(*tes*) *viridis*, with ddRAD- and whole genome pool-sequencing revealed a novel candidate master gene (*bod1l*), governing a male-heterogametic system (XX♀/XY♂). Targeted sequencing across multiple taxa and a male long-read assembly uncovered structural X/Y-variation in the 5'-regulatory region, where a Y-specific non-coding RNA, only expressed in males, suggests regional enhancer-properties. Developmental transcriptomes and RNA *in-situ* hybridization provide evidence for timely and spatially relevant, sex-specific *bod1l*-gene expression in primordial gonads with coinciding differential *H3K4me*-methylation in pre-granulosa/pre-Sertoli cells, pointing to a novel mechanism of amphibian sex determination.

Introduction

Most vertebrates occur as females and males. Vertebrate sex determination comprises environmental triggers, or simple to complex genetic (genotypic) sex determination – and perhaps combinations thereof – that evolved several times[1,2]. Likewise, the genetic and cellular developmental processes directing undifferentiated gonads towards testes or ovaries are diverse[3].

Genetic sex determination is governed by sex chromosomes that are typically differentiated and often microscopically distinguishable in mammals, birds, and several reptile groups[4]. With several exceptions, some reptilians and most other cold-blooded (poikilothermic) vertebrates, like fish and amphibians, exhibit poorly or undifferentiated (homomorphic) sex chromosomes. Three main evolutionary concepts explain this prevalence: (i) frequent turnovers, overthrowing sex-determining genes by novel autosomal mutations, or even the sex determination system (e.g. XY to WZ)[5], (ii) recurrent X-Y recombination, e.g. by occasional sex reversal to XY-females[6], and (iii) "jumping master genes" with conserved genetic triggers translocating to other chromosomes[7] (incl. refs.). Conserved poorly or non-differentiated sex chromosomes

are known in several clades of poikilothermic vertebrates, e.g. in sturgeons, pikes, and
skinks[8-10].

More than 95% of amphibians possess undifferentiated sex chromosomes with either
female (ZZ/ZW) or male (XY/XX) heterogamety[11,12]. Few examples of cytogenetically
differentiated sex chromosomes[12], a female W0/00 male sex determination system[13],
and several amphibians with multiple sex chromosomes have been identified in both
laboratory strains[14] and natural populations[15]. Notably, to date, in >8,650 species of
amphibians[16], the widespread combination of poorly or undifferentiated sex
chromosomes with large genome-sizes (see below) have caused a basic knowledge
gap with only a single though very well-characterized amphibian master sex-
determination gene known in a model species, the African clawed frog (*Xenopus*
laevis[11,17]) and some related species[18]. *Dm-w* arose after (and perhaps in response
to) tetraploidization, and has been lost secondarily in some tetraploid species[19].

True toads, Bufonidae, share large genomes, conserved karyotypes[20] and
undifferentiated sex chromosomes[21]. Typical of most bufonid toads is also Bidder's
organ that develops in males and - in many species - also in females during early
gonadogenesis from the anterior parts of the primordial gonads, considered to possibly
present a rudimentary ovary[22]. Using sex-reversal experiments, castrating males and
regrowing functional ovaries from Bidder's organs, early on, Ponse[23] inferred female
heterogamety (ZW) in common toads (*B. bufo*). This was also cytogenetically deduced
in related *B. gargarizans*[24], and cane toads (*Rhinella marina*[25]). With few exceptions,
a meta-analysis of sex-differential F1-hybrid fitness of interspecific crosses in bufonids
likewise supported widespread female heterogamety[26].

This situation contrasts with that in our target species, Palearctic green toads of the
Bufo(*tes*) *viridis* sub-group. While their sex chromosomes are also undifferentiated[21,27]
(refs. therein), sex-reversal experiments in '*B. viridis*' from Asia Minor[28] suggested an
XY-system. Molecular evidence for male heterogamety resulted from sibship analyses
using microsatellites and nuclear sequence markers[29-31], identifying the largest linkage
group (LG1), homologous to autosomal LG1 of *X. tropicalis* and harboring *dmrt1*, as
sex-linked in several diploid green toad species. All feature strongly-reduced X/Y-
recombination but occasional X/Y-exchange is suggested at evolutionary time
scales[29]. However, phylogenetic analysis revealed X- and Y-alleles of *dmrt1* to cluster
by species and not by gametologue, suggesting X/Y-recombination involving *dmrt1*,

which rejected its role as master sex determiner[31]. Details of sex determination have

not been examined in allopolyploids green toads[32].

In mammals (6,649 species[33]), genome assemblies are available for about one third

(2740[34]). Although there are more species described in amphibians (>8650[16]), owing

to large genome sizes and excessive repetitive regions, to date only >80 amphibian

genomes have been assembled, including 56 out of 7627 Anura, 19 out of 805 Urodela

(Caudata) and 9 out of 221 Gymnophiona[16,34]. However, profiting from long-read- and

3D-sequencing technologies, amphibian genomics is on the rise.

Here, we provide a chromosome-scale genome of a non-model amphibian, the

European green toad, *Bufo*(*tes*) *viridis*, and identify a novel candidate master sex

determining gene - present in multiple related green toad taxa - that evolved by

structural variation, involving a Y-specific ncRNA, in the regulatory region between X

and Y chromosome.

Results

Chromosome-scale genome assembly of *B. viridis*

We obtained a 3.8 Gbp chromosomal-level genome assembly using Oxford nanopore

(ONT) long-read (N50 = 11.6 kbp; sequencing coverage 21x) and Illumina short-read

data (2x150 bp; sequence coverage 47x) for the European green toad (Fig.1a,b). N50

contig size of the final assembly was 1.9 Mbp. Eleven chromosomes (scaffold N50 =

470 Mbp; consisting of 94% of the contigs total length), consistent with the karyotype

of *B. viridis* (Fig.1c), were constructed using chromosome conformation capture (Hi-

C) short reads (318 Gbp). Many unplaced contigs contain a known long satellite DNA

family[35]. Comparing the chromosomes to the *B. bufo* assembly revealed genome-wide

synteny and collinearity with few inversions, despite significant differences in assembly

size (VGP's aBufBuf1.1 = 5.0 Gbp; Fig.1b). *De novo* analysis revealed 68.5% repeat

content in *B. viridis.* Using published data from related species and RNAseq data, we

annotated 21,605 protein coding genes in the genome. BUSCO (Geneset

vertebrata_ODB9)-scoring of protein predictions revealed 96.4% complete coding

sequences and a low missing gene fraction of 1.7% (C:96.4% [S:94.2%, D:2.2%],

F:1.9%, M:1.7%, n:2586).

ddRADseq in *B. viridis* genetic families

We first attempted to identify the sex determination region by double digest restriction-site associated DNA sequencing (ddRADseq). In three genetic families of European green toads, we analyzed a total of 443,378 markers, present in at least one individual with a depth higher than 4. Of these, 96 were significantly associated with male sex ($p < 0.05$; Bonferroni correction), but none was found in all males (Supplementary Figure 1), probably due to high stochasticity of random sequencing of large genome fractions. When mapped to the *B. viridis* genome, 64 of these 96 markers were located along the entire chromosome 1 (scf1) (Supplementary Figure 2), one on chromosome 7 (scf7), five on unplaced scaffolds, and 26 were not aligned uniquely or with low quality (< 20). Overall, this strongly supports a male (XY) genetic sex determination system on chromosome 1, however, with a very small sex-specific region that is not fully covered in each individual. While this approach powerfully corroborated previous microsatellite-based results[29], it did not provide a sex-diagnostic marker.

Pool-seq whole genome sequencing yields a cross-amplifying marker in multiple taxa

To narrow down the sex determining region on scaffold 1, we used whole genome pool sequencing (pool-sex), involving 22 female and 22 male phenotypically-sexed adult green toads from multiple European green toad populations (Supplementary Table 1). Mapping pool-sex short reads to the reference, genome-wide revealed a single peak in a region on scaffold 1 (scf1), where the number of clustered heterozygous SNPs (N = 105) in males surpassed about nine times the maximum number of clustered heterozygous SNPs (N = 12) in females (Fig.1d, Supplementary Table 2; Supplementary Figure 3).

157
158
Fig. 1| The chromosome-scale reference-genome of *Bufo*(*tes*) *viridis* and the new sex-diagnostic
marker. a, Breeding pair of *B. viridis* in amplexus (© M. Stöck). **b,** *B. viridis*-assembly compared to *B.*
bufo assembly by the Vertebrate Genome Project (VGP, Genbank accession number
GCA_905171765.1) showing high synteny of both bufonid genomes across all 11 chromosomes. **c,**
Giemsa-stained karyotype of *B. viridis*, changed from[27]. **d,** Mapping the pool-seq short reads to the *B.*
viridis reference: genome-wide a single region (scf:566,783,058-566,848,882), where the number of
heterozygous SNPs (y-axis) on scaffold 1 (N = 105) in males surpassed about nine times the maximum
number of genomic positions (N = 12) where females exhibit heterozygous SNPs, provides strong
evidence for male heterogamety (XX/XY); for comparisons with a ZZ/ZW system see Supplementary
Fig. 2. **e,** Overview of sex-specific PCR-products of primers BvXY1 in females (♀♀, left) and males
(♂♂, right) of six (i-vi) nominal diploid green toad taxa, electrophoresis result using 4 µL of PCR-
products on a 4% high-resolution Sieve-agarose gel run for 60 min on 70 V; Palearctic green toad taxa
shown are: (i) *siculus*, (ii) *shaartusiensis*, (iii) *viridis*, (iv) *variabiilis*, (v) *balearicus*, (vi) *turanensis*. **f,**
FISH-TSA of a metaphase spread of a *B. viridis* male with red *bod1l*-signals on chromosomes 1 on their
short (p) arm (arrows), chromosomes were counterstained with DAPI (blue-green), scale bar in lower
left corner represents 10 µm. **g,** Range map of Palearctic green toad taxa, numbers (i-vi) and color-
coding as in (e) and (h), diploid (2n) taxa are indicated by circles, triploids (3n) by triangles and
tetraploids (4n) by squares. **h,** Dated mitochondrial phylogeny of Palearctic green toads (adapted
from[32], with the phylogenetic positions of the diploid taxa, in which BvXY1 is sex-diagnostic, indicated

by color-coded arrows as in (**e**); the three ploidy levels are provided to the right of the leaves, note that

the sex marker is seemingly conserved only in the diploid green toads; all polyploids are of hybrid

origin[32].

This strongly supports a male-heterogametic (XY) system, involving a short 66 kb

region (scf1:566,783,058-566,848,882 bp). It encompasses one third of the 5'-

upstream-region and two thirds of the gene *bod1l* (bufVir1.4895.2, biorientation of

chromosomes in cell division 1 like 1). Transcriptomes and cDNA-based PCR-

products of males of *B. viridis* exhibited consistent heterozygosity of exons 7 to 10

(scf1:566,831,904-566,833,300), while females were always homozygous.

The heterozygous 5'-region exhibited several INDELs between X and Y haplotypes,

enabling primer design. In females, BvXY1 amplified a single PCR-product - visible as

one band on the gel, and two to three bands in males, widely consistent with a an

XX♀/XY♂-system, but suggesting multiple Y-specific signals in males. Of note,

BvXY1-primers amplified not only sex-specific products in individual *B. viridis* (sensu

lato) males and females of the pool-sex-approach but also in multiple phenotypically-

sexed males and females of this and four additional taxa (*B. balearicus*, *B. siculus, B.*

turanensis, B. shaartusiensis), comprising a monophyletic radiation, probably >3 Mio.

years old[32] (Fig. 1e-h). The sex-diagnostic PCR-marker BvXY1 amplified a single

specific 100-200 bp band in females of six green toad taxa, but 2-3 bands in their

males, one in females' size, presenting the X-copy. The additional male bands with

the initial BvXY1-primers yielded identical or similar sequences indicating duplicates

on the Y-homologue.

Position of the sex-specific marker and FISH on the karyotype dismiss

duplication with translocation

New sex determining genes evolve by allelic diversification or from gene

duplications[36]. To elucidate the situation in *B. viridis*, we examined synteny and

localization of the candidate gene. *Bod1l* lies between the genes *Cpeb2* and *nkx3-2*

(NK3 homeobox 2), on the long arm of scaffold 1, an evolutionarily conserved gene

order likewise found in *B. bufo*, *Rana temporaria* and *Xenopus tropicalis*, suggesting

synteny throughout anurans.

Fluorescence *in-situ* hybridization with tyramide signal amplification (FISH-TSA) on

metaphase chromosomes of a male *B. viridis* revealed two signals, one on each

homologue, for a *bod1l*-specific probe at a position on the distal part of the short (p)

arm of chromosome 1 (Fig. 1f). Two (and not three) signals suggest that this gene

exists in two copies in XY-males, implying that both heterozygous sequences evolved

by structural diversification[36] rather than translocation of a male-specific duplication,

as shown for *dm-w* of female *Xenopus laevis*[17]. This result is further supported by the

pool-seq and long-read male coverage (see below).

Targeted sequencing analyses in multiple related species sharing the genetic

sex marker

If the hypothetically sex-determining SNPs in coding *bod1l*-regions arose in a common

ancestor, we expected the five diploid taxa, sharing BvXY1 in 5'-*bod1l*-regions (Fig.

1d), to share at least some exonic, possibly functional SNPs. To test this, we applied

targeted enrichment to multiple females and males of five taxa, using an AmpliSeq

custom DNA panel on 80 kb (93%) of the *bod1l*-X-copy, including its 5'-region with

BvXY1, and analyzed 8183 SNPs. Of note, none of the *bod1l* coding SNPs, present

in *B. viridis* (Supplementary File 2), was conserved across all five toad taxa

(Supplementary Files 3, 4, Supplementary Figure 4a), suggesting that this coding

region (cds) does not contain the mechanistic key to green toad sex determination.

Instead, the targeted enrichment approach narrowed down the sex-specific region to

the 5'-region of *bod1l*, containing BvXY1 (Supplementary Figure 4b).

Y-chromosomal structure of *B. viridis* from long-read ONT whole genome

sequencing

After first working with the female reference genome, X-haplotype and *bod1l*-coding

region, we then moved the focus back to the 5'-end of *bod1l* and the BvXY1-vicinity

on the Y-haplotype. To elucidate this Y-chromosomal structure, we reconstructed this

region from ca. 30x whole genome long-reads of a single male *B. viridis* and aligned

it to the X-copy of the *bod1l*-region. Female-to-male pool-seq-coverage of X- and Y-

haplotypes yielded the expected diploid female and haploid male coverage

(Supplementary Figure 5 and 6).

In the *bod1l*-region, the confidently assembled Y-haplotype specific sequence

comprises about 55 kb with some greater-than-zero female pool-seq-coverage,

explicable by repeat-enriched fragments, not assignable to X or Y (Supplementary

Figure 7). The male long-read assembly revealed several structural changes between

X- and Y-haplotypes in the 5'-*bod1l*-region. The *B. viridis*-Y contains large insertions,

mainly stemming from repetitive elements, and a non-coding RNA (ncRNA, see

below). None of the so-called 'usual suspects', genes that often acquire sex-

determination function in vertebrates[1], was detected in this region.

Gene expression during larval development in sibling *B. viridis* tadpoles from

multiple European localities

Pool-sequencing of genomic DNA in *B. viridis* had revealed consistent heterozygosity

in males for *bod1l*-exons 7 to 10. Accordingly, RNAseq revealed sex-differential

expression (scf1:566,831,904-566,833,300), where heterozygous *B. viridis* males

showed 1 SNP in exon 7, and 14 SNPs in exon 10, while females were always

homozygous (confirmed by cDNA-based PCR-products). To test whether *bod1l*-

products are differentially composed after transcription of X- and Y-copy, we analyzed

splice sites in developmental transcriptomes, including the sex determination period.

The absence of significant sex-specific differences excluded alternative splicing.

However, this analysis allowed manual improvement of the *bod1l*-gene model.

To study the expression of the candidate master sex determination gene (*bod1l*) and

ca. 80 other genes (Supplementary File 5) playing roles in sexual development and/or

sex determination in other vertebrates ('usual suspects'[1]), we then examined larval

transcriptomes of six sibling groups, in total 54 individual toads from six locations,

ranging from day 10 after fertilization until Gosner[37] stage 46 (the last about 6 weeks

after metamorphosis) as well as from two adult gonads.

Since gonadal differentiation is histologically detectable only at or after metamorphosis

completion (>Gosner 46[31,37]), the critical period of genetic sex determination lies

earlier, within our sampling period. Until stage 5a, overall *bod1l*-expression varies in

B. viridis between males and females (Gosner 34-38; Fig. 2). From stages 4 to 5b,

other genes, specific for male sexual development, like *sox9*, *sox10*, *sox17*, and *sf1*,

show higher expression in males than in females. Other such genes change their

expression distinctly after stage 4, like *ck18* (bufVir1.18618.1) or *11ßHD*

(bufVir1.16559.1) or - in our series - were detected only in Gosner 46 and subadult

testes (*dmrt1, amh*). Female-specific genes also changed their expression around

stage 4; *wnt4*, *ihh*, *foxL2*, *ptch1*, *gli3*, *fshr* from stage 5a or 5b; however, their

expression was stronger in males than females. Expression of *cyp19a1a*, *cyp17a1*

and *gdf9* was detectable only in stage 5b (Gosner 46) and adult ovaries. *Bod1l*-

expression differences between females and males (Fig. 2) lack statistical significance

but occur at the right developmental period (2 to 5a), as required for a sex-determining

gene. Early male transcriptomes (stages 1 to 5a) contain also the bufonid-specific

Bidder's organ, considered similar to a rudimentary ovary, that develops adjacent to

the anterior primordial gonads[22]. Published qPCR-data suggested that gene

expression in adult Bidder's organs differs from that of testes and ovaries but

resembles the latter, consistent with classical, mid-20th-century male-to-female sex-

reversal experiments, when Bidder's organs grew and resumed oogenesis after the

removal of testes[23] (reviewed[22]). In green toads, Bidder's organ develops in males but

barely in females[22]. Given that the "ovary-like" primordial Bidder's organ is contributing

tissue to our RNAseq-analysis of male tadpoles until stage 5a, we assume a "female-

bias" in males' transcriptomes (stages 1 to 5a). In conclusion and in line with the

AmpliSeq-data from multiple species, only sharing the BvXY1-marker across taxa, we

consider these slight quantitative differences between males and females as sufficient

to trigger green toad sex determination. Of note, there is a trend of greater Y-haplotypic

expression of *bod1l* between stages 4 to 5a (coinciding with an increase of the ncRNA-

Y expression, next paragraph) and a distinctly greater additive XY-expression in males

than XX-expression in females. *Bod1l*-Y also shows the highest expression in testis

compared to other male organs (heart, brain, liver).

Fig. 2 | Haplotype-specific and trend of XX- or XY- additive expression of *bod1l* and the ncRNA-Y in females and males in seven developmental stages. Expression in females *vs*. males. Stages: 1: 10 days after fertilization, 2: 15 days after fertilization, 3: 18 days after fertilization; 4: 34 days after fertilization/Gosner[37] stages 34-38, 5a: Gosner stages 43-44, beginning of metamorphosis, 5b: Gosner stage 46, six weeks after metamorphosis, 6: subadult male and adult female. Numbers of available tadpole transcriptomes varied by stage; TPM: Transcript reads Per Million RNAseq reads.

A non-coding RNA (ncRNA) is Y-specifically expressed upstream of *bod1l*

When further comparing RNAseq (mapped to Y- or X-haplotypes) from ovaries *vs.* testes (in tadpoles plus adjacent Bidder's organ), we discovered a long non-coding RNA ("ncRNA-Y", ca. 1.2 kb, 4 exons, Fig. 3) with its exons 1 and 2 being Y-specific, and expressed exclusively in males, as confirmed by PCR from cDNAs (Fig. 3c, Supplementary Fig. 8). The ncRNA-Y harbors the initially detected gDNA-marker (BvXY1) in intron 3. We hypothesize that this male-specifically expressed ncRNA-Y hints at an enhancer element, missing on the X, which regulates male *bod1l*-expression and simultaneously transcribes the ncRNA-Y in a one-dimensional (1D) manner from the opposite strand[38]. With ca. 80 kb, the putative enhancer sits in a reasonable distance to the *bod1l*-promoter (Fig. 3a,b)[39]. We also tested 200 bp-"sliding-windows" of the ncRNA-region for their *in-silico* potential as enhancers of

bod1l (Supplementary Table 3; Supplementary Fig. 9). The most stringent threshold

of 0.9 still predicted larger parts of the Y-specific region to function as strong

enhancers, intensified by numerous G-quadruplexes (>25 alone in intron 1 of ncRNA-

Y), known to accumulate at promoters and enhancers[40]. As shown by PCR using

cDNA, this long ncRNA-Y is also expressed in skin, liver, brain and heart of males

after metamorphosis as shown using cDNA, and exhibits simultaneous expression of

the Y-haplotype of *bod1l* with the strongest signals in testis (Fig. 2). Due to its low

expression level, it is only detectable in high-coverage transcriptomes of the primordial

male gonads and adult testes but not in heart, brain and liver high-coverage

transcriptomes. It comprises 5.8 kb on the gDNA level and can be male-specifically

PCR-amplified in all those six taxa containing the BvXY1-marker and suggests a

common ancestral feature (Fig. 3d).

Fig. 3 | Structure of the X and Y-haplotype of the *bod1l*-region of *Bufo*(*tes*) *viridis* with focus on

the 5'-end of the candidate master sex-determining gene and the Y-specific ncRNA. a, X-

chromosomal (above, pink) and Y-chromosomal haplotype (below, blue) with color-coded homologous

or X- and Y-specific INDELs, and the Y-specific long (ca. 5.8 kb) non-coding RNA (ncRNA-Y), flanking

a putative enhancer at the 5'-end of the gene, ca. 80 kilobases before its promoter. **b,** Graph of the

signal of the AI-based software *i-enhancer*, reaching from 0 to 2, predicting strong enhancer properties

of the Y-specific 5'-end of *bod1l*, including a G-quadruplex-rich region on the non-template strand,

overlapping with the ncRNA-Y. **c,** Y- and thus male-specific expression of the ncRNA-Y (204 bp PCR-

product from cDNA) in larval green toads, from 10 days after hatching until early metamorphosis; in

females, only non-specific products occurred, since X-copies of *bod1l* do not contain this ncRNA (for

details and positive controls of females: Supplementary Figure 8). **d,** Unrooted Maximum-Likelihood-

based phylogenetic tree, showing that the ncRNA-Y evolved in a common ancestor of diploid Palearctic
green toads sharing this XY-system, with *siculus* and *viridis* spanning the most divergent lineages; two
PCR-products (2.5 and 5.8 kb), directly amplified by independent PCRs from six related diploid green
toad taxa, clustered by Y-gametologue and thus species (* symbolized 100% bootstrap support; only
support values >50% are shown).

Spatially different gene expression in the developing gonads

For a master sex-determination gene, not only the accurate timing of its expression

but also its organ- and cell-specific spatial expression is essential. We provide

evidence by RNA *in-situ* hybridization in histological sections of gonads in Gosner

stage 36-37, where *bod1l* shows sex-specific differential expression. In female

gonads, *bod1l*-expression was evident in *vasa*-positive primordial germ cells

(developing oogonia) but absent from surrounding somatic pre-granulosa cells (Fig.

4a,c, Supplementary Figure 10). In contrast, *vasa*-positive male primordial germ cells

(developing spermatogonia) as well as a majority of adjacent somatic pre-Sertoli cells

exhibited *bod1l*-expression (Fig. 4b,d, green arrows). In Gosner stages 43-44, at the

beginning of metamorphosis, *bod1l*-expression in females remained similar to Gosner

36-37. In males, however, *bod1l*-expression was no longer detectable in somatic pre-

Sertoli cells (Fig. 4e,f; Supplementary Figure 10). This suggests a dynamic regulation

of *bod1l* in early male and female gonads, with expression in somatic cells exclusively

in male but not female gonads at the relevant stages (prior stage 4), when sex

determination occurs.

It has been shown that *Bod1l* interacts with *Setd1a* to facilitate tri-methylation of

histone H3 at lysine 4 (H3K4me3[41]), with similarities in eukaryotes from yeast to

mammals[42]. To examine its status in male and female gonads, we performed

H3K4me3-immunostaining on Gosner 43-44 gonads (Fig. 4g,h). In females, gonads

exhibited distinct H3K4me3-signals in nuclei of both germ and somatic pre-granulosa

cells (Fig. 4g, white asterisks and arrowheads, respectively; Supplementary Figure

10). In contrast, H3K4me3 was strongly detected in male germ cell nuclei (Fig. 4h,

white asterisks) but hardly detectable or absent in nuclei of somatic pre-Sertoli cells

(Fig. 4h, yellow arrowheads). In Gosner stage 46, six weeks after metamorphosis,

bod1l was expressed (RNAseq, PCR of cDNA) in both sexes in several somatic

tissues (brain, heart, liver, intestine) as well as in gonads.

		Expression in somatic cells		Expression in germ cells	
		Male	Female	Male	Female
Gosner 36-37	*Bod1L*	+	−	+	+
Gosner 43-44	*Bod1L*	−	−	+	+
	H3K4me3	low	high	high	high

Fig. 4 | *Vasa*- and *bod1l*-expression and H3K4me3-methylation in primordial gonads of *B. viridis*, i.e. in the female early ovary, and in male Bidder's organ with testes (Gosner stages 36-37) as well as in early metamorphosis (Gosner stages 43-44). a-f, *Vasa* and *bod1l* expression shown by RNA *in-situ* hybridization (ISH) on 20 μm transverse cryosections of female (a, c, e, g) and male (b, d, f, h) gonads at developmental stages Gosner 36-37 (a-d) and 43-44 (e, f), the latter presenting early metamorphic stages. DAPI was used to stain nuclei. Magenta asterisks indicate nuclei of germ cells; magenta arrows: cytoplasm of germ cells; yellow arrows: nuclei of somatic cells not stained by RNA ISH; green arrows: nuclei of somatic cells stained by RNA ISH. **g,h,** H3K4me3 and DAPI staining on 20 μm transverse cryosections of female (g) and male (h) green toad Bidder's organ and gonads, Gosner stage 43-44. White asterisks indicate H3K4me3-positive germ cell nuclei; white arrowheads, H3K4me3-positive somatic cell nuclei; yellow arrowheads, H3K4me3-negative somatic cell nuclei.

Scale bars: 20 μm. **i,** Summary of *bod1l*-expression and H3K4me3-methylation in somatic cells and
germ cells in Gosner 36-37 and Gosner 43-44 male and female gonads.

XY-phylogeny in six Eurasian diploid taxa from targeted sequencing

For a master sex-determining locus, evolved from a common ancestor, non-
recombining loci are expected to cluster by gametologue (i.e. X- or Y-specific
haplotypes) and not by species. Analyzing X- and Y-specific sequences obtained from
the AmpliSeq custom DNA panel and amplicon sequencing of ncRNA-Y of six diploid
green toad taxa, we show that in the 2,671 bp-region (scf1:566,788,797-566,791,468),
containing the sex-specific BvXY1-marker, sequences cluster by gametologue (Xs *vs.*
Ys, Supplementary Figure 4c), while in the similar-sized adjacent upstream region
(scf1:566,786,116-566,788,796), sequences clustered by taxon/species, suggesting
occasional recombination (Supplementary Figure 4d). This provided evidence of both
the stop of recombination since the sequence arose in a common ancestor and the
apparent functional relevance of this region, since these species also do not share
coding sequences.

Discussion

Our data in *B. viridis* and five related diploid green toad taxa are in accordance with a
gene that shows structural changes in its 5'-regulatory region on the genomic level,
and sex-specific gene expression in heterozygous (XY) males *vs.* homozygous (XX)
females, including a strong phylogenetic and poolseq-supported signal of local
recombination arrest. Its sex-specific expression occurs at the right time, coinciding
with that of an ncRNA on the Y, much earlier than histologically detectable
differentiation, in the target organs (primordial gonads), and therein in the germ and
surrounding somatic cell populations (in male pre-Sertoli but not in female pre-
granulosa cells). These properties qualify *bod1l* as a candidate master sex
determination gene. While future evidence (e.g. using CRISPR/Cas9) is desirable for
its better mechanistic understanding, a single genome-wide peak of heterozygosity
revealed by pool-seq and evolutionary conservation of the structural XY-differentiation
in several Eurasian taxa along with the shared Y-specific ncRNA, support its
candidacy as the Palearctic green toad master gene.

The only known amphibian master sex determining gene, *dm-w* of clawed frogs
(*Xenopus*), governs female-heterogamety (ZW♀/ZZ♂) and evolved by gene

duplication as a *dmrt1*-paralog, involving neofunctionalization of a non-coding part of

a DNA-transposon[11,17-19,43]. Here, we provide evidence for a 5'-upstream-region of a

gene in a male heterogametic bufonid toad system (XX♀/XY♂), evolved by structural

diversification in a common ancestor, as shown by the associated ncRNA-Y-

phylogeny (Fig. 3c). To our knowledge, this reveals only the 2nd master sex

determining gene in amphibians. *Bod1l* on the Y involves (on the opposite strand) an

ancestral insertion of a ncRNA-Y, harboring the partially duplicated BvXY1, and a

neighboring signal of a G-quadruples-rich enhancer towards its 3'-end. This is a

mechanistic hint that the enhancer modifies *bod1l*-expression in XY-males (Fig. 3), as

shown by timely and spatially differential expression in germ and pre-Sertoli but not in

pre-granulosa cells (Fig. 4).

There are several, partly interlinked pathways by which spatially and/or timely

differential *bod1l*-expression may determine sex in green toads.

(i) Sex determination of vertebrates differs regarding the timing of primordial germ cells

to enter meiosis[44], with females entering meiosis earlier than males, exhibiting an

arrest. *Bod1l* is well-known to maintain genomic integrity at sites of DNA-replication

damage[45]. *Bod1l* interacts towards its N-terminus with the histone methyltransferase

Setd1a, which triggers the histone chaperone function of *Fancd2*[46], part of the Fanconi

anemia group (*Fanc*), that plays roles in zebrafish sex reversal[47]. Thus, differential

bod1l-expression might cause a compromised DNA repair system by shifting sex

determination towards males.

(ii) Interaction of *Bod1l* with *Setd1a* and H3K4me3 refers to tri-methylation on lysine

residue 4 in the amino-terminal tail of histone H3, is associated with potentially active

promoter regions[48] and linked to double-strand breaks, essential for meiosis 1

crossing-over, ensuring chromosome segregation[49,50]. Thus, the differential *bod1l*-

expression might enable double-strand breaks as a precondition for the earlier start of

recombination in females but not in males. This suggests that *bod1l*-expression in

males at the SD stage is, though not significantly, lower than in females (Fig. 2: stage

4), presumably by decreased expression of both functional copies, since we have no

evidence for a compromised Y-allele.

(iii) H3K4-methylation in mammals[48] is associated with *de novo* DNA-methylation in

germ cells, and essential for maintaining transcription in the growing oocyte and for

triggering zygotic genome activation[51], with XY-mice deficient for the H3K9-

demethylating enzyme *Jmjd1a* showing increased H3K9-dimethylation and decreased

activating marks of H3K4-trimethylation across the *Sry*-locus, leading to a high

frequency of sex reversal[2].

According to Dupont and Capel[52] "prior to sex determination" in placental mammals,

bipotential supporting gonadal cells of both genetic sexes maintain identical

transcriptional programs, allowing commitment to either Sertoli (testes) or granulosa

(ovary) cell fate. In pre-Sertoli and pre-granulosa cells, bivalent promoters of key sex

determining genes (e.g. *Sox9*, *Fgf9*, *Wnt4*, *Foxl2*) have both, H3K27me3 and

H3K4me3 marks[52]. After sex determination by the master sex determination gene *Sry*,

promoters to be activated in that sex lost the repressive H3K27me3 mark, while genes

to be silenced maintained their bivalency[53]. This suggested that (a) sex-determining

genes are poised for activation (or repression) at the bipotential stage, (b) gene

activation involves removal of H3K27me3 marks, while (c) repression of the alternative

supporting cell fate requires maintenance of existing H3K27me3 repressive marks

rather than *de novo* deposition of these marks after commitment to the Sertoli or

granulosa program[52].

Our preferred hypothesis is that differential *bod1l*-expression (reminiscent of *Sry*)

might determine sex by changing H3K4-methylation in green toads during early

gonadal development, prior to metamorphosis.

To test the hypothesis that *bod1l-expression* coincides in time and space with changes

of H3K4-methylation in developing green toad gonads, we used H3K4me-antibodies

together with RNA *in-situ* hybridization (Fig. 4). In Gosner stage 43-44, slightly prior to

morphological differentiation, we found a strong difference between male and female

gonadal somatic cell H3K4-methylation. This suggests that spatially different *bod1l*-

expression results in sex-specific methylation patterns that initiate sex-specific

gonadal differentiation.

Given the multiple potential links between *bod1l*-expression and sex-specific meiotic

stages shown above, we hypothesize that this gene functions as a suppressor of the

female sex determination pathway – a scenario found in many vertebrates[54]. Following

differential *bod1l*-expression, in the bipotential gonad, ovarian pre-granulosa cells

transdifferentiate to testicular pre-Sertoli cells. In contrast to the only known amphibian

master gene in clawed frogs (*Xenopus*), green toad *bod1l* did not evolve by gene

duplication but structural diversification in the 5'-region of the gene, namely by Y-

specific sequence insertion with emergence of an enhancer, which is transcribing

ncRNA-Y. This evolution of sex determination seems broadly similar to allelic

diversification, so far unknown in amphibians, but known from many examples of
master genes in fish[36].
493
494 **References:**
495

1. Herpin, A. & Schartl, M. Plasticity of gene-regulatory networks controlling sex
determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO*
Rep **16**, 1260-74 (2015).
2. Capel, B. Vertebrate sex determination: evolutionary plasticity of a fundamental
switch. *Nat Rev Genet* **18**, 675-689 (2017).
3. Bachtrog, D. *et al.* Sex determination: why so many ways of doing it? *PLoS Biol* **12**,
e1001899 (2014).
4. Rovatsos, M. *et al.* Conservation of sex chromosomes in lacertid lizards. *Mol Ecol* **25**,
3120-6 (2016).
5. Volff, J.N., Nanda, I., Schmid, M. & Schartl, M. Governing sex determination in fish:
regulatory putsches and ephemeral dictators. *Sex Dev* **1**, 85-99 (2007).
6. Perrin, N. Sex reversal: a fountain of youth for sex chromosomes? *Evolution* **63**, 3043-
9 (2009).
7. Bertho, S., Herpin, A., Schartl, M. & Guiguen, Y. Lessons from an unusual vertebrate
sex-determining gene. *Philos Trans R Soc Lond B Biol Sci* **376**, 20200092 (2021).
8. Kuhl, H. *et al.* A 180 Myr-old female-specific genome region in sturgeon reveals the
oldest known vertebrate sex determining system with undifferentiated sex
chromosomes. *Philos Trans R Soc Lond B Biol Sci* **376**, 20200089 (2021).
9. Kostmann, A., Kratochvíl, L. & Rovatsos, M. Poorly differentiated XX/XY sex
chromosomes are widely shared across skink radiation. *Proc Biol Sci* **288**, 20202139
(2021).
10. Pan, Q. *et al.* The rise and fall of the ancient northern pike master sex-determining gene.
Elife **10** (2021).
11. Ito, M. Sex Determination and Differentiation in Frogs. in *Reproductive and*
Developmental Strategies: The Continuity of Life (eds. Kobayashi, K., Kitano, T., Iwao,
Y. & Kondo, M.) 349-366 (Springer Japan, Tokyo, 2018).
12. Schmid, M. & Steinlein, C. Sex chromosomes, sex-linked genes, and sex determination
in the vertebrate class Amphibia. in *Genes and Mechanisms in Vertebrate Sex*
Determination (eds. Scherer, G. & Schmid, M.) 143-176 (Birkhäuser Basel, Basel,
2001).
13. Green, D.M. Cytogenetics of the endemic New Zealand frog, Leiopelma hochstetteri:
extraordinary supernumerary chromosome variation and a unique sex-chromosome
system. *Chromosoma* **97**, 55-70 (1988).
14. Roco Á, S. *et al.* Coexistence of Y, W, and Z sex chromosomes in Xenopus tropicalis.
Proc Natl Acad Sci U S A **112**, E4752-61 (2015).
15. Furman, B.L.S. *et al.* A frog with three sex chromosomes that co-mingle together in
nature: Xenopus tropicalis has a degenerate W and a Y that evolved from a Z
chromosome. *PLoS Genet* **16**, e1009121 (2020).
16. AmphibiaWeb. See www. amphibiaweb.org (accessed June 2023).
17. Yoshimoto, S. *et al.* A W-linked DM-domain gene, DM-W, participates in primary
ovary development in Xenopus laevis. *Proc Natl Acad Sci U S A* **105**, 2469-74 (2008).
18. Cauret, C.M.S. *et al.* Developmental Systems Drift and the Drivers of Sex Chromosome
Evolution. *Mol Biol Evol* **37**, 799-810 (2020).

19. Evans, B.J. *et al.* New insights into Xenopus sex chromosome genomics from the
Marsabit clawed frog X. borealis. *J Evol Biol* **35**, 1777-1790 (2022).
20. Bogart, J.P. Karyotypes. in *Evolution of the genus Bufo* (ed. Blair, W.F.) 171-232
(Austin, Texas: University of Texas Press 1972).
21. Schmid, M. Chromosome banding in Amphibia I. Constitutive heterochromatin and
nucleolus organizer regions in Bufo and Hyla. *Chromosoma (Berl.)* **66**, 361-388 (1978).
22. Piprek, R.P., Kloc, M. & Kubiak, J.Z. Bidder's organ--structure, development and
function. *Int J Dev Biol* **58**, 819-27 (2014).
23. Ponse, K. La proportion sexuelle dans la descendance issue des oeufs produits par
l'organe de Bidder des crapauds femelles. *Rev. Suisse Zool.* **48**, 541–544 (1941).
24. Changxiang, W., Qun, L. & Wei, X. Studies of chromosome banding and sister
chromatid exchange in Bufo bufo gargarizans. *Acta Gen Sinica* **10**, 291–297 (1983).
25. Abramyan, J., Ezaz, T., Graves, J.A. & Koopman, P. Z and W sex chromosomes in the
cane toad (Bufo marinus). *Chromosome Res* **17**, 1015-24 (2009).
26. Malone, J.H. & Fontenot, B.E. Patterns of reproductive isolation in toads. *PLoS One* **3**,
e3900 (2008).
27. Stöck, M., Steinlein, C., Lamatsch, D.K., Schartl, M. & Schmid, M. Multiple origins of
tetraploid taxa in the Eurasian Bufo viridis subgroup. *Genetica* **124**, 255-72 (2005).
28. Ueda, H. Offspring of sex-reversed males in Bufo viridis. *Sci. Rep. Lab. Amphib. Biol.*
Hiroshima. Univ. **10**(1990).
29. Stöck, M. *et al.* Low rates of X-Y recombination, not turnovers, account for
homomorphic sex chromosomes in several diploid species of Palearctic green toads
(Bufo viridis subgroup). *J Evol Biol* **26**, 674-82 (2013).
30. Brelsford, A. *et al.* Homologous sex chromosomes in three deeply divergent anuran
species. *Evolution* **67**, 2434-40 (2013).
31. Tamschick, S. *et al.* Sex chromosome conservation, DMRT1 phylogeny and gonad
morphology in diploid Palearctic green toads (Bufo viridis subgroup). *Cytogenet*
Genome Res **144**, 315-24 (2014).
32. Betto-Colliard, C., Hofmann, S., Sermier, R., Perrin, N. & Stöck, M. Profound genetic
divergence and asymmetric parental genome contributions as hallmarks of hybrid
speciation in polyploid toads. *Proc Biol Sci* **285**(2018).
33. Burgin, C. *et al.* Mammal Diversity Database (Version 1.11) dataset. Zenodo
http://dx.doi.org/10.5281/zenodo.7830771 (2023).
34. NCBI. See www.ncbi.nlm.nih.gov (accessed June 2023).
35. Guzmán, K. *et al.* Identification and characterization of a new family of long satellite
DNA, specific of true toads (Anura, Amphibia, Bufonidae). *Scientific Reports* **12**,
13960 (2022).
36. Schartl, M. & Herpin, A. Sex Determination in Vertebrates. in *Encyclopedia of*
Reproduction (Second Edition) (ed. Skinner, M.K.) 159-167 (Academic Press, Oxford,
2018).
37. Gosner, K.L. A simplified table for staging anuran embryos and larvae with notes on
identification. *Herpetologica*, 183–190 (1960).
38. Burren, O.S. *et al.* Chromosome contacts in activated T cells identify autoimmune
disease candidate genes. *Genome Biol* **18**, 165 (2017).
39. Jackson, D.A., Dickinson, P. & Cook, P.R. The size of chromatin loops in HeLa cells.
Embo j **9**, 567-71 (1990).
40. Lyu, J., Shao, R., Kwong Yung, P.Y. & Elsässer, S.J. Genome-wide mapping of G-
quadruplex structures with CUT&Tag. *Nucleic Acids Res* **50**, e13 (2022).
41. Bayley, R. *et al.* H3K4 methylation by SETD1A/BOD1L facilitates RIF1-dependent
NHEJ. *Mol Cell* **82**, 1924-1939.e10 (2022).

42. Giovanni, C. *et al.* The BOD1L subunit of the mammalian SETD1A complex sustains
the expression of DNA damage repair genes despite restraining H3K4 trimethylation.
Preprint at https://www.biorxiv.org/content/10.1101/2023.04.06.535882v2 (2023).
43. Hayashi, S. *et al.* Neofunctionalization of a Noncoding Portion of a DNA Transposon
in the Coding Region of the Chimerical Sex-Determining Gene dm-W in Xenopus
Frogs. *Mol Biol Evol* **39**(2022).
44. Bowles, J. & Koopman, P. Retinoic acid, meiosis and germ cell fate in mammals.
Development **134**, 3401-11 (2007).
45. Higgs, M.R. *et al.* BOD1L Is Required to Suppress Deleterious Resection of Stressed
Replication Forks. *Mol Cell* **59**, 462-77 (2015).
46. Higgs, M.R. *et al.* Histone Methylation by SETD1A Protects Nascent DNA through the
Nucleosome Chaperone Activity of FANCD2. *Mol Cell* **71**, 25-41.e6 (2018).
47. Rodríguez-Marí, A. & Postlethwait, J.H. The role of Fanconi anemia/BRCA genes in
zebrafish sex determination. *Methods Cell Biol* **105**, 461-90 (2011).
48. Smallwood, S.A. & Kelsey, G. De novo DNA methylation: a germ cell perspective.
Trends Genet **28**, 33-42 (2012).
49. Youds, J.L. & Boulton, S.J. The choice in meiosis – defining the factors that influence
crossover or non-crossover formation. *Journal of Cell Science* **124**, 501-513 (2011).
50. Ki, B.S. *et al.* Epigenetic regulator Cfp1 safeguards male meiotic progression by
regulating meiotic gene expression. *Experimental & Molecular Medicine* **54**, 1098-
1108 (2022).
51. Fan, H.-Y. & Sun, Q.-Y. Chapter 12 - Oocyte Meiotic Maturation. in *The Ovary (Third*
Edition) (eds. Leung, P.C.K. & Adashi, E.Y.) 181-203 (Academic Press, 2019).
52. Dupont, S. & Capel, B. The Chromatin State during Gonadal Sex Determination. *Sex*
Dev **15**, 308-316 (2021).
53. Garcia-Moreno, S.A. *et al.* CBX2 is required to stabilize the testis pathway by
repressing Wnt signaling. *PLoS Genet* **15**, e1007895 (2019).
54. Nakamura, M. Sex determination in amphibians. *Semin Cell Dev Biol* **20**, 271-82
(2009).
618

Online content

Methods

Ethics and animal welfare statement

Our research complies with all relevant ethical regulations; the relevant animal

protection committee of the IGB and the city of Berlin (LAGESO) approved the study

protocol where relevant. Most adult toads were sampled more than 20 years ago by

either minimal invasive sampling methods using tail tips of larvae or saliva of adults,

obtained by buccal cotton swabs. We also used tissues of road kill toads and/or

collected a few adult animals in Greece. For sampling and export of toads and

tadpoles, permits were kindly provided by the Ministry of the Environment of Greece

(115790/229) 2014-2017. Developmental stages of tadpoles were obtained from

control groups of an animal experiment approved by the German State Office of Health

and Social Affairs (LaGeSo, Berlin, Germany; G0359/12) and a commercial breeder.

The tadpoles were kept according to approved practice (permit ZH 114 to IGB),
including the provision of appropriate tank size, sufficient rate of waterflow, natural
photoperiod, *ad libitum* food supply, as well as temperatures within the species'
thermal tolerance range. This ensured that no pain, suffering, distress or lasting harm
was inflicted on the animals. According to the permit G0359/12, tadpoles and juvenile
toads were euthanized by an overdose of Tricaine PHARMAQ 1000mg/g (MS222)
since this does not incur pain, suffering or distress.

Chromosome-scale genome assembly
The female *B. viridis* reference genome was sequenced in-house at IGB using ONT
long reads (MinIon device, SQK-LSK109 kits and R9.4.1 flow cells) and Illumina short
reads (Novogene, Europe). The long reads and pre-assembled short reads IDBA-UD
assembly[55] were assembled using the WTDBG2.2 assembler[56]. Sequences were
error-corrected using two rounds of Flye polish with long reads[57] and four rounds of
VCFconsensus using short reads[58]. The chromosome-scale assembly was achieved
using Arima HiC-data and the JUICER/3DDNA package and manual curation by
juicebox[59-61]. Protein coding genes were annotated using RNAseq data of *Bufonidae*
as well as Hyloidea transcriptomes and more distantly related amphibian proteins,
available at NCBI. A reference genome browser was created http://genomes.igb-
berlin.de/Greentoad/.[62]

Double-digest RAD sequencing (ddRADseq)
DNA of three genetic families consisting for one data set of both parents (buccal
swabs) and 40 sexed mostly phenotypically offspring and for two other sample sets of
25 or 15 siblings, was extracted using the DNeasy Tissue Kit or the BioSprint robotic
workstation (QIAGEN), following the manufacturer's protocols. Double-digest RADseq
libraries were prepared according to this study[30]. In brief, genomic DNA digestion by
two restriction enzymes (*Sbf*I, *Mse*I) and subsequent ligation of fragments to adapters
containing unique barcodes (4–8 bp) was followed by library-PCR for 20 cycles for
only different cut-sites containing fragments, and a gel-based size selection, isolating
the 400–500 bp fraction. RADseq libraries were single-end sequenced on an Illumina
HiSeq 2500, at a 20x coverage.
Raw ddRAD reads were demultiplexed with STACKS *process_radtags* version 2.55
(cite https://onlinelibrary.wiley.com/doi/full/10.1111/mec.15253), using the enzymes

sbfI and mseI (*--renz_1 sbfI --renz_2 mseI*) and the parameters *-c* (clean data), -r

(rescue barcodes and RAD-Tag cut sites), and -q (discard reads with low quality

scores).

Demultiplexed reads were used as input for the RADSEX workflow[63]. First, a table

containing the number of occurrences of each unique sequence in the reads file in

each individual was generated with *radsex process*, using default parameter values.

Then, the distribution of markers between males and females was computed with

radsex distrib using a minimum depth to consider a marker present in an individual of

5 (--min-depth 5). Markers were aligned to the genome of *B. viridis* (bufVir1) using

radsex map using a minimum depth of 5 (--min-depth 5) and other parameters set to

default. Results were plotted with the *SGTR* package, with the entire process provided

as a Snakemake workflow[64].

Pool-sex approach

Of 22 safely sexed adult toads of each sex from multiple populations in the range of

the European green toad taxa (*B. viridis viridis* and *B. viridis variabilis*), using buccal

swabs or tissues genomic DNA was extracted using the DNeasy Tissue Kit or the

BioSprint robotic workstation (QIAGEN), following the manufacturer's protocols.

Individual DNAs were quantified using the QuantiFluor® dsDNA System kit

(Promega), adjusted and equimolarly mixed to yield a female and a male pool. Whole

genome library preparation for 300 bp fragments and whole genome sequencing was

performed at NOVOGENE (UK), on a NovaSeq PE150 to yield 131 and 140 Gbp of

data for male and female pool, respectively. Reads were mapped to the reference

genome using the short read mode of MINIMAP2 (parameters -K 2000M -I 100G -a -

x sr -t 80[65], and converted to sorted bam-files by SAMTOOLS[66]. Variants (SNPs,

MNPs, INDELs) were called from both bam files using the PLATYPUS variant caller[67]

with integrated duplicate fragment reads removal (–filterDuplicates = 1) and

reassembly of reads (–assembleAll = 1). The resulting vcf-file was screened for XY-

or XX-specific variants using AWK-scripting and different variant read coverage cutoffs

(i.e. SNP read coverage in females and males >=15; variant read count in females

<=7% or >=93% of total SNP read coverage as a condition for homozygosity; variant

frequency in males between 45 - 55% of total SNP read cov. as a condition for

heterozygosity; for screening ZW SNPs conditions for males and females were

interchanged). Variants were clustered according to their distance in the genome

(maximal distance of variants in a cluster: 9000 bp), counted using BEDTOOLS

(merge, annotate[68]) and sorted by variant count. Regions with highest variant counts

or densities and neighboring regions were inspected for genes related to sex

determination.

Sex-diagnostic PCR marker design

The region 566,783,058-566,848,882 bp on scaffold 1 includes several INDELs

apparent between the X and the Y haplotype, allowing design of a number of primers

that were primarily tested in gradient-PCRs on phenotypically sexed *B. viridis*. While

several primers showed sex-specific amplification in this species, one (BvXY1) could

be optimized to work across multiple species.

A collection of DNAs (extraction: see above) comprising samples of phenotypically

sexed green toads of the entire radiation[32] served to test sex-specificity of all PCR

markers. PCR-amplifications were carried out in 25 µL reactions containing 1.25 µL of

each primer (BvXY1_F: 5'_CTGAATAGCAATTGGACACAGC_3; BvXY1_R:

5'_ATGTGAGAGCTGTTACAATGGA_3'), 14.5 µL of water, 5 µL of AllTaq buffer (5x),

0.5 µL of AllTaq polymerase (Qiagen), and 2 µL of DNA (concentration 4-20 ng/µL).

Temperature regime comprised 2 min at 95°C (initial denaturation), and 38 cycles of

(1 min at 95°C denaturation, 45 s at 59°C annealing, 45 s at 72°C extension), and 5

min at 72°C final extension. PCR products (4 µL) were visualized by electrophoresis

on 4% high-resolution Sieve-agarose (Biozym) gels, run for 60 min at 70 V.

To PCR-amplify either a 2.5 kb fragment or nearly the entire ncRNA-Y, comprising 5.8

kb based on gDNA on the Y-chromosome in all diploid green toad taxa with sex-

specific BvXY1-markers, we used the same ingredients and volumes as above either

with primers nRNA_Y_ex1F: 5'_GGGCAGTGAAGAGAAGTCCT_3', nRNA_Y_ex2R

5'_TTCACCATTCTCCAGCCTGT_3' for the 2.5 kb-fragment or

5.8kb_ncRNA_Y_F: 5'_GTGGTAGTTGCTGTGCCTTT_3', 5.8kb_ncRNA_Y_R:

5'_CCGGTGTGGTGTAAAATATCCA_3' for the 5.8 kb-piece. Temperature regimes

involved 2 min at 95°C, and 37 cycles of (1 min at 95°C denaturation; 2.5 min at 63°C

(for the 2.5 kb) or 5 min at 59.4°C (for 5.8 kb) annealing; 1 min at 72°C extension and

5 min at 72°C final extension (note extended annealing times). PCR products (4 µL)

were visualized by electrophoresis on 1.5% agarose (Biozym) gels, run for 60 min at

80 V. PCR-products (21 µL) were barcoded using the NBD196 and LSK110 ONT-kits,

and sequenced using the ONT-MinIon device, as described. Basecalling was

performed using ONTs GUPPY version 6 and the super accuracy base-calling model (SUP-model).

Phylogenetic tree of a 2.7 kb-region at the Y-specific region boundary

Consensus sequences of 100 ONT reads of Y-specific amplicons were calculated using NGSPECIESID[69]. Homologous X-sequences were reconstructed from targeted enrichment sequencing data of females using BCFTOOLS consensus[70] for the region scf1:566,788,797-566,791,468. We also excluded regions of the sequence not covered by enrichment data by supplying a masking file through parameter "-m" of bcftools consensus. All resulting sequences were multiple aligned using MAFFT (automatic alignment parameters). The alignment was filtered by GBLOCKS[71] to remove columns with excessive missing data. IQTREE2[72] was used with parameter "-alrt 1000 -bb 1000" to calculate a phylogenetic tree. To calculate a phylogenetic tree of the upstream neighboring region (scf1:566,786,116-566,788,796) data from target enrichment for males and females was used as described above.

Phenotypic sexing in multiple species

Phenotypic sex of adult toads of multiple diploid Eurasian green toad taxa had been established during the breeding period when males exhibited secondary sexual characters like nuptial excrescences and thorny glandular tips on the back, or were caught producing male advertisement calls and/or male release calls. Females were identified as filled with eggs, caught in amplexus, and/or while spawning, producing female release calls and/or lacking the male secondary sexual characters. During the spawning migrations, several road kill toads also allowed for anatomical identification (possession of ovaries or testes) of phenotypic sex.

Assembly of Y-haplotype

Several iterative approaches were used to extract Y-haplotypic sequencing reads of *B. viridis* GR57 male[73] long read data (Oxford Nanopore Promethion platform; Novogene). First, we mapped the reads to the BvXY1-marker sequences to separate Y reads of the region. Second, we performed a FLYE[57] assembly of the male data, identified several contigs with variation compared to the X haplotype (*bod1l*-region), and extracted the corresponding reads. The Y-specific reads were assembled, then added to the female assembly and all male reads were mapped to the combined

sequences, which allowed finding more reads matching better to Y than to X contigs.
This procedure was iteratively repeated several times until the Y-haplotype and its
structure could be assembled. Pool-seq data was again mapped to the female
assembly with added Y-haplotype sequences by MINIMAP2[65] showing the expected
coverage differences between female and male pools. A dot plot comparison of X
versus Y haplotype involved the corresponding function of the NCBI Blast website.
774
775 **Testing enhancer activity of the Y-haplotype**
776 We used the AI-based predictor "iEnhancer-Deep" in a 200 bp-"sliding-windows"
777 approach in the ncRNA region to test its *in-silico* potential as enhancers of *bod1l-*
778 *expression*[74].
779

Gene expression using RNAseq during development in sibling tadpoles from
multiple European populations
In total 54 transcriptomes of sibling tadpoles comprising brothers and sisters (plus
three unsexed larvae) of four genetic families from four different geographic locations
as well as gonadal tissue of a subadult male and adult female from two additional
localities were examined by RNAseq (transcriptomics). Tadpoles presented five
developmental stages from control groups of an ecotoxicological experiment[31]. They
were euthanized in MS222 and flash-frozen in liquid nitrogen. In tadpoles older than
Gosner stage 37, the intestine was removed prior to flash-freezing. In tadpoles at
Gosner stage 43-44, only the body segment that contained the gonads was used for
RNA extraction. In toadlets of Gosner stage 46 (specifically 6 weeks after
metamorphosis), ovaries, testes (separated from Bidder's organ), and several other
organs (brain, heart, liver, intestine) were dissected and stored in RNAlater prior to
flash-freezing. DNA from tailtips, or skin samples of each tadpole or buccal swabs
(juvenile toads) served for genetic sexing using BvXY1-primers. RNAs were extracted
using the TRIzol Reagent (Thermo Fisher Scientific, Waltham, USA) according to the
supplier's recommendation and cleaned using the RNeasy Mini Kit (Qiagen). All RNAs
included in this study exhibited RIN values >9.0. Library processing and RNAseq were
carried out by NOVOGENE (Cambridge, UK) on a NovaSeq 6000 PE 150, generating
9 Gb of raw data per sample.
Transcriptome sequences of 54 male and female juvenile developmental stages and
two adult gonads were normalized and mapped to the genome using the nf-core RNA-

Seq pipeline in STAR aligner (https://github.com/alexdobin/STAR/releases,—
quantMode GeneCounts). A list comprising genes that play roles in sexual
development and/or sex determination in other vertebrates (also known as 'usual
suspects'[1]) is shown as Supplementary File 5). To plot *bod1l* and *ncRNA-Y* expression
trends over the developmental stages and separate expression analysis of *bod1l-X*
and *bod1l-Y,* we mapped all reads, including the male haplotype, to the female
assembly. As the *bod1l-Y* gene model is 3'-partial, we used the corresponding parts
of the *bod1l-X* gene model for quantification. Counts of reads matching the different
haplotypes' gene models were extracted using FEATURECOUNTS (from SUBREAD-
2.0.3 package[75] with mapping quality (MQ) higher than 19, to increase haplotype
specificity. Read counts were normalized by total mapped reads of each sample (TPM
normalization). Average expression and standard deviation was calculated and plotted
combining samples of the different stages using spreadsheet software.
815

cDNA of developmental stages
Using the iScript RT Supermix for RT-qPCR kit (Biorad), 10 µL of RNA (concentration
of 80 ng/µL) from all developmental stages (as used for RNAseq) were randomly
reverse-transcribed into cDNA according to the manufacturer's recommendations. To
test allele-specific *bod1l*-gene expression, we designed primers (bod1l_RNA1F:
5'_AACTGGAAGAAAAGCGCAAA_3'; bod1l_RNA1R:
5'_GTGCTCTTGGCATTCTGTGA_3') and amplified 577 bp from cDNA under the
same conditions as for BvXY1 that were Sanger-sequenced in both directions. To test
expression of the ncRNA-Y, we used primers designed to match its inter-exonic
regions NcRNAY_Ex1to2_F: 5'_CATGAGGAGAGCAGTAATGAACC_3',
NcRNAY_Ex3to2_R: 5'_GCTGTCATCAACCTGCTCAA_3' and amplified and
visualize 204 bp male-specific products from cDNA under the same conditions as for
BvXY1 but at 55°C annealing temperature.
829

Probe preparation for RNA** in-situ **hybridization
The sequence marker obtained by cDNA amplification using bod1l_RNA1F/R primers
was also used as a probe for RNA *in-situ* hybridization on histological gonad sections
of various developmental stages. As a probe of a control gene, expressed in gonads
of both sexes, we used a 501 bp cDNA fragment of *Vasa* (DDX4-DEAD-Box-Helicase-
4, VASA-homolog), amplified with the primers Vasa1F:

5'_AACCAAAAAGAAGGCGGATT_3';   Vasa1R:

5'_ATGCCAGGATTGCTGTAACC_3' under the same conditions as for BvXY1 but

with annealing temperature of 57°C.

RNA** in-situ *hybridization and H3K4me antibody staining

We opened the body cavity of freshly euthanized tadpoles of three developmental

stages and fixed them in 4% paraformaldehyde (PFA) buffered in 1x PBS

(Thermofisher), pH = 7.0, at 4°C overnight. Samples were washed in filtered 1×

phosphate-buffered saline with tween 20 (PBST, ROTI®Stock) twice for 5 min, then

dehydrated in a series of methanol (MetOH) with shaking at room temperature (25%

MetOH in PBST for 10 min to 1 h, 50% MetOH in PBST for 10 min to 1 h, 75% MetOH

in PBST for 10 min to 1 h, 100% MetOH for 10 min to 1h, timing dependent on tissue

sample size). Samples were stored in 100% MetOH at -20°C until histological

sectioning.

Cryosectioning, RNA *in-situ* hybridization and immunohistochemistry were performed

as described[76] with slight modifications. Transverse cryosections (20 μm) were

collected on SuperFrost Plus glass slides (Menzel Gläser J1800AMNZ) and stored at

-20°C for up to 2 months. To generate *bod1l* and *vasa* riboprobes, primers

Bod1_RNA_Bv_1F (5'_AACTGGAAGAAAAGCGCAAA_3') and Bod1_RNA_Bv_1R

(5'_GTGCTCTTGGCATTCTGTGA_3') were used to amplify a 577 bp *bod1l* sequence

and primers vasa_1F (5'_AACCAAAAAGAAGGCGGATT_3') and vasa_1R

(5'_ATGCCAGGATTGCTGTAACC_3') were used to amplify a 501 bp *Vasa* sequence

from Gosner 36-37 *B. viridis* cDNA. These sequences were then ligated to a pCR-

bluntII-TOPO vector (Invitrogen) and plasmids were purified using a plasmid DNA

miniprep kit according to manufacturer's instructions (Promega A1460). For synthesis

of sense and antisense riboprobes, plasmids were linearized using *BamHI* or *NotI*

(New England Biolabs) and *in vitro*-transcribed using T7 or SP6 polymerase (DIG RNA

Labeling Mix, Roche), respectively. Rabbit polyclonal anti-histone H3 (tri methyl K4)

primary antibody (Abcam ab8580) and secondary goat anti-rabbit Alexa Fluor 568

(1:500, Invitrogen A-11011) were used. Sections were counterstained with 0.25 μg/mL

DAPI and mounted in Mowiol 4-88 (Calbiochem). Imaging was performed using the

Nikon SMZ18 stereomicroscope and Olympus FluoView FV3000 confocal

microscope. To process images, Bitplane Imaris and Adobe Photoshop were used.

FISH-TSA

As a probe for fluorescence *in-situ* hybridization with tyramide signal amplification (FISH-TSA), we designed novel primers to amplify a longer portion of *bod1l* from cDNA (*bod1l*_Exon7To10_F: 5′_AACTGGAAGAAAAGCGCAAA_3′, *bod1l*_Exon7To10_R: 5′_TGTGTTGTCTGGAATCACTGG_3′; same conditions as for BvXY1 but with annealing Temperature of 60°C.), exhibiting the highest heterozygosity in the coding segment of the gene in *B. viridis* males. A PCR-product was produced using the PPP Master Mix (Top-Bio, Czech Republic) according to manufacturer's instructions (annealing temperature 60°C) and purified[77]. The cDNA amplicons were cloned using the TOPO-TA cloning kit and DH5α-T1R Competent Cells (Thermo Fisher Scientific). Plasmid DNA was isolated from bacterial colonies using the EZNA Plasmid DNA Mini Kit I (Omega Bio-Tek Inc.), sequenced, and tested to contain the correct insert by BLAT search on the green toad genome browser (see Data availability). The sequence of the *bod1l* clone was compared to the reference genome. Subsequently, the *bod1l* locus was re-amplified and purified[77]. A 20 years old tissue suspension (bone marrow), obtained from a male European green toad from Germany, treated with colchicine and fixed in methanol-acetic acid[27] was dropped onto a microscope slide[78], followed by FISH. The probe was labelled with digoxigenin-11-dUTP (Jena Bioscience) using DecaLabel DNA Labelling Kit (Thermo Fisher Scientific). The FISH-TSA followed[79] with minor modifications[80]. Mitotic metaphase spreads were counterstained with ProLongTM Diamond Antifade Mountant with the fluorescent 4',6-diamidino-2-phenylindole, DAPI stain (ThermoFisher Scientific). Microscopy using Leica Microsystem (Wetzlar, Germany) and processing of metaphase images followed[81].

Analysis of potential alternative splicing of bod1l

For the entire *bod1l* region, including the 3'- and 5-' UTR-regions (scf1:566495375-567417162), aligned RNAseq-reads were extracted from each RNA-seq alignment file, the splice sites locations were retrieved and counted. Using all the splice sites, a reference file of them was built. The splice event counts were merged for all samples using the reference and stored as csv text file (Supplementary file 6).

Structural evaluation of the bod1l-*X and Y protein copies*

We used Alphafold2[82] via its cloud-based application and its local installation ColabFold[82] and analyzed the structure predictions by RaptorX[83] for structural and

904 potential functional amino-acid differences between the X and the Y-copy of *B. viridis*.
905 The molecular data viewer iCn3D[84] was used to inspect the structures and highlight
906 specific amino acid residues (Supplementary text 1).
907

AmpliSeq custom DNA panel

After providing pre-publication access to the *B. viridis* genome-browser, the Illumina
concierge design service team developed two primer pools covering ca. 80 kb of the
bod1l-region (scf1:566,785,000-566,865,000) and 10 kb for a control gene (*amh*),
known to play roles in sexual development and sex determination in other vertebrates
and lying on the same scaffold (scf1:324181812-324192250). Overall, the theoretical
coverage was 92.3% for *bod1l*, and 98.2% for *Amh*.
Wet lab approaches followed the "AmpliSeq for Illumina On-Demand, Custom, and
Community Panels" (Document #1000000036408 v09; pp. 5-23[85]) for two primer pools
with 193-384 primer pairs in the standard workflow. Briefly, this included ten steps, (i)
quantification of DNA samples of multiple sexed green toads (QuantiFluor® dsDNA
System kit, Promega), (ii) target amplification in two separate wells for the two primer
pools, (iii) partial digestion of amplicons, (iv) ligation of individual indices/sample, (v)
library clean up and (vi) amplification, (vii) 2nd clean up, (viii) checking libraries on the
Bioanalyser for expected insert size and concentration (QuantiFluor® dsDNA System
kit, Promega), (ix) dilution and mixing, and (x) sequencing, performed by Novogene
on a NovaSeq PE250 for individual libraries obtained for 54 diploid green toads (ca.
100 Mio bp raw data per library).
We included Palearctic green toads by the following numbers, taxa and sexes: 8 *B.*
siculus (4 ♂♂, 3 ♀♀), 9 *B. viridis* (5 ♂♂, 4 ♀♀), 8 *B. balearicus* (5 ♂♂, 3 ♀♀), 9 *B.*
shaartusiensis (4 ♂♂, 5 ♀♀), 10 *B. turanensis* (4 ♂♂, 5 ♀♀). Species' nomenclature
follows[32].
The sequence reads from target enrichment were mapped against the reference
genome using MINIMAP22 (-I 100G -a -x sr -t 4)[65]. SAMTOOLS[66] was used to add
read group and sample names, to fix mates and sort the bam file output. All BAM files
were merged and variant calling for the target regions was performed by SAMTOOLS
and BCFTOOLS[66] with parameters to cope for extreme sequencing depth (samtools
view -bu input.bam scf1:324181812-324192250 scf1:566783149-566866298 |
bcftools mpileup --annotate DP4 --max-depth 1000000000 -f bufVir1d.fa - | bcftools
call -mv -Ob -o $1.calls.bcf). The output was converted to vcf format and summarized

by an awk script that filtered SNP position and calculated allele frequency from DP4

annotation for each sample. The resulting table was further inspected in Excel and

SNPs that were heterozygous in males while homozygous in females were filtered. To

fine map the region of the sex marker male specific SNPs of the the different species

were intersected (resulting in male specific SNPs present in 2, 3, 4, 5 species) and

plotted along the target regions in our genome browser.

References:
946

55. Peng, Y., Leung, H.C., Yiu, S.M. & Chin, F.Y. IDBA-UD: a de novo assembler for
single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*
28, 1420-8 (2012).
56. Ruan, J. & Li, H. Fast and accurate long-read assembly with wtdbg2. *Nat Methods* **17**,
155-158 (2020).
57. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P.A. Assembly of long, error-prone
reads using repeat graphs. *Nat Biotechnol* **37**, 540-546 (2019).
58. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-8
(2011).
59. Durand, N.C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution
Hi-C Experiments. *Cell Syst* **3**, 95-8 (2016).
60. Dudchenko, O. *et al.* De novo assembly of the Aedes aegypti genome using Hi-C yields
chromosome-length scaffolds. *Science* **356**, 92-95 (2017).
61. Durand, N.C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps
with Unlimited Zoom. *Cell Syst* **3**, 99-101 (2016).
62. Kent, W.J. *et al.* The human genome browser at UCSC. *Genome Res* **12**, 996-1006
(2002).
63. Feron, R. *et al.* RADSex: A computational workflow to study sex determination using
restriction site-associated DNA sequencing data. *Mol Ecol Resour* **21**, 1715-1731
(2021).
64. https://github.com/RomainFeron/paper-sexdetermination-greentoad-2021.
65. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094-3100 (2018).
66. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-9 (2009).
67. Rimmer, A. *et al.* Integrating mapping-, assembly- and haplotype-based approaches for
calling variants in clinical sequencing applications. *Nat Genet* **46**, 912-918 (2014).
68. Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-2 (2010).
69. Sahlin, K., Lim, M.C.W. & Prost, S. NGSpeciesID: DNA barcode and amplicon
consensus generation from long-read sequencing data. *Ecol Evol* **11**, 1392-1398 (2021).
70. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**(2021).
71. Castresana, J. Selection of conserved blocks from multiple alignments for their use in
phylogenetic analysis. *Mol Biol Evol* **17**, 540-52 (2000).
72. Minh, B.Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020).

73. Gerchen, J.F. *et al.* A Single Transcriptome of a Green Toad (Bufo viridis) Yields
Candidate Genes for Sex Determination and -Differentiation and Non-Anonymous
Population Genetic Markers. *PLoS One* **11**, e0156419 (2016).
74. Kamran, H., Tahir, M., Tayara, H. & Chong, K.T. iEnhancer-Deep: A Computational
Predictor for Enhancer Sites and Their Strength Using Deep Learning. *Applied Sciences*
12, 2120 (2022).
75. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-30 (2014).
76. Dasyani, M. *et al.* Lineage tracing of col10a1 cells identifies distinct progenitor
populations for osteoblasts and joint cells in the regenerating fin of medaka (Oryzias
latipes). *Dev Biol* **455**, 85-99 (2019).
77. Knytl, M. & Fornaini, N.R. Measurement of Chromosomal Arms and FISH Reveal
Complex Genome Architecture and Standardized Karyotype of Model Fish, Genus
Carassius. *Cells* **10**(2021).
78. Courtet, M., Flajnik, M. & Du Pasquier, L. Major histocompatibility complex and
immunoglobulin loci visualized by in situ hybridization on Xenopus chromosomes.
Dev Comp Immunol **25**, 149-57 (2001).
79. Krylov, V., Tlapakova, T. & Macha, J. Localization of the single copy gene Mdh2 on
Xenopus tropicalis chromosomes by FISH-TSA. *Cytogenet Genome Res* **116**, 110-2
(2007).
80. Knytl, M., Tlapakova, T., Vankova, T. & Krylov, V. Silurana Chromosomal Evolution:
A New Piece to the Puzzle. *Cytogenet Genome Res* **156**, 223-228 (2018).
81. Seroussi, E. *et al.* Avian Expression Patterns and Genomic Mapping Implicate Leptin
in Digestion and TNF in Immunity, Suggesting That Their Interacting Adipokine Role
Has Been Acquired Only in Mammals. *Int J Mol Sci* **20**(2019).
82. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679-682 (2022).
83. Peng, J. & Xu, J. RaptorX: exploiting structure information for protein alignment by
statistical inference. *Proteins* **79 Suppl 10**, 161-71 (2011).
84. Wang, J. *et al.* iCn3D, a web-based 3D viewer for sharing 1D/2D/3D representations
of biomolecular structures. *Bioinformatics* **36**, 131-135 (2020).
85. https://support.illumina.com/downloads/ampliseq-for-illumina-custom-and-
community-panels-reference-guide-1000000036408.html.

Acknowledgements

Funding has been in part provided by the German Research Foundation (DFG) to
MStöck and HK (STO493/8), and the Leibniz-Institute of Freshwater Ecology and
Inland Fisheries (IGB). WKloas and MStöck also received funding from the European
Union's Horizon 2020 research and innovation programme, grant agreement No.
825753 (ERGO). FISH-TSA experiments were partly covered by the PJAC-project
CZ.02.01.01/00/22_010/0002902, MSCA Fellowships CZ–UK (MK). We thank Petros
Lymberakis, Natural History Museum of Crete, University of Crete, for help with the
sampling and export permits of green toads by the Ministry of the Environment of
Greece (115790/229) 2014-2017. The tissue and DNA collection of MStöck profited

from donations of multiple international colleagues over 25 years; he thanks everyone. We are grateful to Sandra Bittmann (IGB) for HMW-DNA extraction, Eva Kreuz (IGB) for initial efforts to clone the SD-region, Julia Bindl (IGB) for help with the DNA and RNA work, Hughes Parinello (Plateforme MGX) for preparing the HiC-libraries and sequencing, the BeGenDiv (Berlin, Germany) for providing lab space and Illumina-sequencing support for the ddRAD. We thank Sebastian Petri (Biozyme) for discussions on high-resolution agarose gel-electrophoresis.

Authors' contributions:

HK: Contributed to study design, sequenced long reads, assembled and annotated the genome, set up the genome browser, mapped and analyzed pool-seq sequences and SNPs of the AmpliSeq custom DNA panel, performed phylogenetic analyses, contributed to 1st manuscript draft

TWH: Made histological sections and performed all RNA *in-situ* hybridizations and H3K4me immunostainings, contributed to 1st manuscript draft

CK: Analyzed all transcriptomes, and gene expression data, analyzed splice sites of *bod1l*

WKleiner: Performed RNA extraction, bioanalyzer quantification and evaluation, did most marker testing by PCRs, co-worked in the wet lab for the AmpliSeq custom DNA panel

BK: Performed molecular work to characterize the ncRNA of multiple green toad taxa, prepared RNA for transcriptomes and cDNA-work at Gosner stage 46

MC: Contributed to ddRAD wet lab

RF: Analyzed the ddRADseq data

MK: Performed FISH-TSA

WKloas: Contributed to study design, discussed developmental histology and data.

MSchartl: Advised in many respects, initiated some collaborations, discussed data and study progress and results.

CW: Contributed to study design, supervised and analyzed RNA *in-situ* hybridization and H3K4me immunostainings.

MStöck: Conceived the study, did all field work/sampling, raised tadpoles, prepared and fixed tissues for chromosome and RNA *in situ* work, collected buccal swabs and tissues over 25 years, sexed toads, extracted all DNAs, performed ddRAD wet lab,

1061 developed all primers, did some marker testing by PCRs, co-worked in the wet lab for

1062 the AmpliSeq custom DNA panel, wrote the 1st manuscript draft.

1063 All authors contributed text to the manuscript, and read and improved the final version.

1064

1065 **Competing interests**

1066 The authors declare to have no competing interests.

1067

Data availability

All data generated in this study have been deposited in publicly accessible databases.

Genome assembly, whole genome sequencing (WGS) data and RNA-seq reads have

been submitted to GenBank on 25 February 2022 (submission ID: SUB11111520,

Bioproject: PRJNA292872, Biosample: SAMN26135295). Alternatively, the IGB-

Green Toad genome browser http://genomes.igb-berlin.de/Greentoad/ provides easy

access to the assembly and its annotation.

1075

1076 **Additional information**

1077

Supplementary File 1: Description of Supplementary Data.

Supplementary File 2: Analyses of SNPs causing 6 non-synonymous changes

in the coding region of *bod1l* in 49 male and female transcriptomes as compared

to the female reference genome. None of these changes was 100% associated with

genetic sex as in the 5'-region of *bod1l*.

Supplementary File 3: Percentages of covered parts of ca. 80 kb of the *bod1l*-

region by targeted sequencing in six diploid green toad taxa, using a panel

based on the female *B. viridis* reference genome, compared to ca. 20 kb of the

gene for *anti-Muellerian hormone* (*amh*). Taxa abbreviations: Bbal - *balearicus*,

Bsha - *shaartusiensis*, Bsic - *siculus*; Btur - *turanensis*, Bvir - *viridis* and *variabilis*.

Supplementary File 4: SNP-comparisons in ca. 80 kb of the *bod1l*- (scf1:

566,783,513 - 566,865,063) and ca. 10 kb of the *amh*-regions (scf1: 324,181,734 –

324,192,327) by targeted sequencing in six diploid green toad taxa in a panel

based on the female *B. viridis* reference genome. The coordinates of the cluster of

male specific SNPs present in all five species are colored in red (scf1: 566,790,085 -

566,790,204). Values for color-coded SNP-comparisons with the reference genome:

>=0.9: homozygous (green/yellow) = identical to reference; between 0.9 and 0.1:

heterozygous (orange) = one allele identical to reference; <=0.1: homozygous(red) = completely different from reference; NA: SNP not covered by sequencing data in the relevant individual / taxon; M - male, F - female; position: SNP position on the female *B. viridis* reference genome; taxa abbreviations as in Supplementary File S3.

Supplementary File 5: Boxplots showing expression of 80 genes that play roles in sexual development and/or sex determination in other vertebrates (also known as 'usual suspects'[1], in developmental stages of *B. viridis*. X-axis showing the normalized intensity of expression; Y-axis marks six developmental stages (details as in Fig. 2).

Supplementary File 6: File of splice site read counts in *bod1l*-region.

Supplementary Figures

Supplementary Figure 1: **Analyses of ddRAD markers in three genetic families of *B. viridis* using RADsex[63].** Of the 443,378 markers present in at least one individual with a coverage >5; 96 markers were significantly associated with male sex ($p < 0.05$ after Bonferroni correction), but none of these markers were found in all males.

Supplementary Figure 2: **Alignment of ddRAD markers shown in Supplementary Figure 1 to the *B. viridis* reference genome.** 64 of the 96 markers significantly associated with sex were located on chromosome 1 (scaffold 1, scf1), one marker was aligned to chromosome 7 (scf7), five markers were aligned to unplaced scaffolds, and 26 markers were not aligned uniquely or with a mapping quality lower than 20.

Supplementary Figure 3: **Mapping pool-sex short reads to the *B. viridis* reference genome with SNP-heterozygosity in males (a) *vs*. females (b) on scaffold (scf) 1.**

Supplementary Figure 4: Male-specific SNPs of 6 species over *bod1l*- region. a SNPs show different blocks of male-specific heterozygosity between species, which exclude coding changes of *bod1l* as sex-determination mechanism. **b** Highest congruence of male-specific SNPs among all species is in the PCR marker/ncRNA-region (red ellipsis). **c** Phylogenetic tree of haplotypes from the marked region shows clustering by gametologue. **d** sequences from the upstream region were clustered by species.

Supplementary Figure 5: Coverage distributions of X/Y haplotypes depending on female/male pool-seq in *B. viridis*. a Y-haplotype, **b** X-haplotype.

Supplementary Figure 6: Female/male pool-seq coverage over haplotypes in *B. viridis*. a female pool-sequencing, **b** male pool-sequencing.

Supplementary Figure 7: X- and Y-haplotype comparisons by dot plots in the *bod1l*-region. a X-*versus*-Y dot plot, **b** Y-*versus*-Y, **c** X-*versus*-X.

Supplementary Figure 8: a, Y- and thus male-specific expression of the ncRNA-Y (204 bp PCR-product from cDNA) in larval green toads. PCR-products from cDNA (204 bp), obtained with primers NcRNAY_Ex1to2_F/_R; to the left males, to the right females; each sex is shown from 10 days after hatching until early metamorphosis (Gonser 43-44) and one adult gonad for each sex; in females, only non-specific products occurred, since X-copies of *bod1l* do not contain and thus do not express this ncRNA. **b,** Gel image of a control-PCR using the females from (a)

Supplementary Figure 9: Graph of the signal of the AI-based software i-enhancer. Sinal bars reaching from 0 to 2, predicting strong enhancer properties of the Y-specific 5'-end of *bod1l*, including a G-quadruplex-rich region on the non-template strand, overlapping with the ncRNA-Y.

Supplementary Figure 10: *Vasa*, *bod1l* expression and H3K4me3-methylation in *B. viridis* gonads, overview images and additional representative examples.
a-k, *Vasa* and *bod1L* expression shown by RNA *in-situ* hybridization (ISH) on 20 µm transverse cryosections of female (**a, b, e, f**) and male (**c, d, g-k**) gonads, Gosner 36-37. DAPI was used to stain nuclei. Sense controls for RNA ISH are shown in **a, c, e, g**, right panels. Red boxes in **a, c, e, g** (left panel) indicate zoomed images shown in **b, d, f, h**, respectively. Yellow boxes in **b, d, f, h** (right panel) indicate zoomed images shown in **Fig. 4a**-**d,** respectively. Red boxes in **i, k** (left panel) indicate zoomed images shown in right panels of **i, k**. *Bod1L* staining on three different cryosections of male gonads are shown in **g, i, k**. Magenta asterisks indicate nuclei of germ cells; magenta arrows: cytoplasm of germ cells; yellow arrows: somatic cells not stained by RNA ISH; green arrows: somatic cells stained by RNA ISH. **l,m,** H3K4me3 and DAPI staining on 20 µm transverse cryosections of female (**l**) and male (**m**) *B. viridis* Bidder's organ and gonads, Gosner stage 43-44. Yellow boxes in **l, m** indicate zoomed images shown in **Fig. 4g** and **Fig. 4h,** respectively. White asterisks indicate H3K4me3-positive germ cell nuclei; white arrowheads, H3K4me3-positive somatic cell nuclei; yellow arrowheads, H3K4me3-negative somatic cell nuclei. Scale bars: 50 µm.

1161

Supplementary Tables

Supplementary Table 1: Localities/populations of European green toads that contributed to the pool-seq experiment.

Supplementary Table 2: Summary of the pool sequencing approach for scaffold 1 (scf1) for SNPs homozygous in males/heterozygous females (Z/W) as well as SNPs heterozygous in males/homozygous females (X/Y).

Supplementary Table 3: Signal of the AI-based software i-enhancer, reaching from 0 to 2, predicting strong enhancer properties of the Y-specific 5'-end of *bod1l*, including a G-quadruplex-rich region on the non-template strand, overlapping with the ncRNA-Y.

Supplementary Texts

Supplementary Text 1: Potential impact of X- *vs*. Y-*bod1l* coding-sequences in *B. viridis*.

a

X

Sex-specific PCR-marker (BvXY1) in 6 green toad taxa

Bod1L
19 exons, 52.544 kb

Y

ncRNA-Y

Bod1L

Putative enhancer for *Bod1L* on Y

Putative interaction in chromatin loop leads to sex-specific regulation (promotor to enhancer distance ~80 kb on the Y)

Putative promotor of *Bod1L* on X and Y

b

ncRNA-Y

Signal stregth enhancer prediction

Coordinates Y-haplotype

G-quadruplex-rich non-template strand

Homologous sequences: X and Y
X-specific insertions
Y-specific insertions

Duplicated motifs on X and/or Y…
…some with partial ncRNA-Y
Y-duplicated repeat-rich region

c

♂♂   100 bp ladder   ♀♀

204 bp

ncRNA-Y

d

82 | *viridis*, male reference genome
    viridis, 5.8 kb
54 *variabilis*, 2.5 kb
    kermanensis, 5.8 kb
    turanensis, 2.5 kb
    turanensis, 5.8 kb
    balearicus, 5.8 kb
    balearicus, 2.5 kb
    shaartusiensis, 2.5 kb
    shaartusiensis, 5.8 kb
    siculus, 2.5 kb
    siculus, 5.8 kb

0.002 changes/site

ncRNA-Y

		Expression in somatic cells		Expression in germ cells	
		Male	Female	Male	Female
Gosner 36-37	*Bod1L*	+	-	+	+
Gosner 43-44	*Bod1L*	-	-	+	+
	H3K4me3	low	high	high	high