Effect of Soybean and Seaweed-based Diets on Growth Performance, Feed Utilization, and Gut Microbiota of Tilapia: A Systematic Review and Meta-analysis

Leonildo dos Anjo Viagem[1,2,3*]; Jean Nepomuscene Hakizimana[1]; Cyrus Rumisha[2]; Brunno da Silva Cerozi[4] and Gerald Misinzo[1,5*]

[1] SACIDS Africa Centre of Excellence for Infectious Diseases, SACIDS Foundation for One Health, Sokoine University of Agriculture, PO Box 3297, Morogoro, Tanzania

[2] Department of Animal, Aquaculture and Range Sciences, College of Agriculture, Sokoine University of Agriculture, PO Box 3004, Morogoro, Tanzania

[3] Department of Food and Agricultural Sciences, Rovuma University, Cabo Delgado, Mozambique

[4] Department of Animal Science, College of Agriculture, University of São Paulo, Avenida Padua Dias, 11, PO Box 9, Piracicaba, São Paulo, Brazil

[5] Department of Veterinary Microbiology, Parasitology and Biotechnology, College of Veterinary Medicine and Biomedical Sciences, Sokoine University of Agriculture, PO Box 3019, Morogoro, Tanzania

*Corresponding author

E-mail: leonildo.viagem@sacids.org (LAV) and gerald.misinzo@sacids.org (GM)

Abstract

Tilapia, a significant aquaculture species globally, relies heavily on feed for its production. While numerous studies have investigated the impact of soybean and seaweed-based diets on tilapia, a comprehensive understanding remains elusive. This review aimed at evaluating and synthesizing the existing literature on these diets' effects, focusing on growth performance, feed utilization, and gut microbiota. A systematic search of databases was conducted using Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and a total of 57 studies were included in the qualitative analysis and 24 in the meta-analysis. The results indicated that soybean-based diets, at a 59.4% inclusion level improved the Specific Growth Rate (SGR) of tilapia with an effect size of -2.14 (95% CI: -2.92, -1.37; $p < 0.00001$; $I^2 = 99\%$) and did not improve the feed conversion rate (FCR), as the effect size was 1.80 (95% CI: 0.72, 2.89; $p = 0.001$; $I^2 = 100\%$). For seaweed-based diets, at a 15,9% inclusion level did not improve SGR, with an effect size of -0.74 (95% CI: -1.70, 0.22; $p = 0.13$; $I^2 = 99\%$), and the FCR with an effect size of -0.70 (95% CI: -1.94, 0.54; $p = 0.27$; $I^2 = 100\%$). Regarding the gut microbiota, was noted a lack of studies meeting the inclusion criteria for tilapia. However, findings from studies on other farmed fishes suggested that soybean and seaweed-based diets could have diverse effects on gut microbiota composition and promote the growth of beneficial microbiota. This study suggests that incorporating soybean-based diets at 59.4% inclusion can improve the SGR of tilapia. Seaweed-based diets, while not demonstrating improvement in the analyzed parameters with an inclusion level of 15.9%, have the potential to contribute to the sustainability of the aquaculture industry when incorporated at lower levels.

Keywords: soybean, seaweed, diet, gut microbiota, tilapia, fish

Introduction

Tilapia are among the most important freshwater fish species worldwide and play a crucial role in global aquaculture production. As adaptable species to various environmental conditions, tilapia can be reared in a wide range of farming systems, including ponds, cages, and recirculation aquaculture systems [1,2,3]. The success of tilapia farming largely depends on the formulation of cost-effective and nutritionally balanced feed and an appropriate feeding management strategy that promotes optimal growth performance and feed utilization [4].

Traditional feed ingredients for tilapia include fishmeal and fish oil due to their high protein content balanced essential amino acids profile and omega-3 [5-8], but they are expensive and can contribute to the overfishing of wild fish stocks [9-11]. Therefore, there is a growing interest in exploring alternative protein sources for fish feed, such as soybean and seaweed. Soybean is a cost-effective, and widely available plant protein source [12,13], while seaweeds offer unique nutritional properties and are natural forage for fish [14]. Both have been investigated as partial or total replacements to improve the growth performance and feed efficiency of tilapia. Their inclusion in this study can provide a comprehensive understanding of their individual nutritional properties and highlight their potential as feed options for tilapia. However, there are challenges in including seaweed and soybeans in tilapia diets, such as mismatches in fatty acids and amino acids profile, leading to sub-optimal growth and health [15,16]. Additionally, soybean and seaweed contain antinutritional factors such as saponins, tannins, phytic acid, lectins, protease inhibitors, and amylase inhibitors [16-18], which may negatively affect nutrient digestibility, palatability, and overall feed performance.

Apart from the nutritional aspects, the gut microbiota plays a crucial role in the overall health and growth of fish [19-21] and can participate in modulating feeding behavior and efficiency [22,23]. Dietary interventions can influence the composition and activity of the gut microbiota, thereby influencing the host's metabolism, immune system, and overall performance [24-27].

While individual studies have investigated the effects of soybean and seaweed-based diets on tilapia, a comprehensive understanding of their impact on growth performance, feed utilization, and gut microbiota is lacking. A systematic review and meta-analysis are essential to synthesize the available evidence and provide a comprehensive overview of the effects of these alternative feed ingredients on tilapia. Therefore, the present study evaluated and synthesized the existing literature on the effects of soybean and seaweed-based diets on growth performance, feed utilization, and gut microbiota of tilapia to achieve sustainability in aquaculture practices.

Methods

Study design

This study was conducted according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (S1 Table). The protocol related to this systematic review was registered on Open Science Framework (osf.io/zg72m) with registration DOI: https://doi.org/10.17605/OSF.IO/ZG72M.

Data source and search strategy

A systematic literature search was conducted using online databases including Google Scholar, PubMed, Wiley Online Library, and ScienceDirect, to identify relevant studies published between 2000 and 2022. The search terms were combined as follows: "seaweed" OR "macroalgae" NOT "microalgae" AND "growth performance" AND "feed utilization" AND "tilapia"; "soybean" NOT "soybean fermented" AND "growth performance" AND "feed utilization" AND "tilapia"; "seaweed" OR "macroalgae" NOT "microalgae" AND "gut microbiota" OR "intestinal microbiota" AND "tilapia" OR "other fish" NOT "human"; "soybean" NOT "soybean fermented" AND "gut microbiota" OR "intestinal microbiota" AND "tilapia" OR "other fish" NOT "human". Additionally, a manual search of references cited in relevant articles was performed to identify

additional studies that may have been missed in the database search. The search strategies and the respective number of studies from the four databases are detailed in the S2 Table.

Inclusion and exclusion criteria

In this systematic review, studies published from January 2000 to December 2022 were included based on the following criteria: (1) articles written in English, (2) articles focusing on tilapia production; (3) studies utilizing various soybean-based diets such as soybean defatted, soybean meal, full-fat soybean, soybean milk, and soybean protein; (4) studies utilizing seaweed-based diets, including seaweed extract and various species of green, brown, red seaweed (S3 Table) to partially or totally replace fish meal/oil; (5) full-text available; (6) studies published in peer-reviewed journals; (7) studies reporting SGR and FCR with their respective standard error or standard deviation, and (8) articles examining the effect of soybean and seaweed-based diets on intestinal microbiota of other fish than tilapia. Excluded from consideration were: (1) studies not written in English; (2) studies not related to the use of soybean and seaweed as tilapia feed; (3) articles not published in peer-reviewed scientific journals; (4) articles not available in full-text format; (5) studies that used microorganisms to ferment soybean/seaweed; and (6) studies utilizing prebiotics in tilapia/fish production that were not derived from soybean or seaweed; (7) article that did not describe the composition of the gut microbiota; (8) studies lacking fish meal or fish oil as control group.

Selection process

The screening of all papers was done in the Rayyan tool for systematic literature review (https://www.rayyan.ai/). After downloading the article in the databases (in RIS format), the files were imported to Rayyan to check for duplicates, screening the title, abstract, and full text by two independent reviewers (Leonildo dos Anjo Viagem-LAV and Jean Nepomuscen Hakizimana-JNH).

Data extraction

Data were extracted from each included study using a standardized data extraction form created in Microsoft Excel 2013 (Microsoft Corporation, Washington, USA). This extraction process was carried out independently by two reviewers (LAV and JNH). The extracted data encompassed basic information such as author and publication year, experimental details including sample size and experiment duration, the inclusion level of the diets (both applied and recommended), and information about the different experimental groups, namely the soybean-based diet group, seaweed-based diet group, and control group (fish meal or fish oil). Parameters extracted for these groups included specific growth rate (SGR), and feed conversion ratio (FCR), along with their corresponding standard mean or standard error. Additionally, details regarding the types of soybean and seaweed species utilized in the experiments were recorded (S3 Table).

Risk of bias in included studies

The study quality and risk of bias were evaluated using the Cochrane risk-of-bias tool in Cochrane Collaboration's software Review Manager Version 5.4.1 (RevMan 5.4.1). This tool contains seven domains: (1) random sequence bias (selection bias), (2) allocation concealment (selection bias), (3) blinding of participants and personnel (performance bias), (4) blinding of outcome assessment (detection bias), (5) incomplete outcome data (attrition bias), (6) selective reporting (reporting bias), and (7) other biases [28]. This systematic review focused on five domains, as the blinding of participants and personnel (performance bias) and the blinding of outcome assessment (detection bias) are not applicable to aquaculture studies. In the domain of other bias was analyzed the information related to sample size and experiment duration.

Meta-analysis

A meta-analysis was conducted to analyze the effect of soybean and seaweed on the SGR and FCR of tilapia. Effect sizes were calculated, and a random-effects model was employed. Publication

bias was assessed using funnel plots of the effect of soybean on SGR and FCR (S1 Fig) and

seaweed (S2 Fig). The heterogeneity of the included studies was assessed by the I-squared statistic

test and was classified as low (<25%); moderate (25-50%) and high (>50%) [29]. The subgroup

analysis and sensitivity were performed to explore possible causes of heterogeneity in the study.

All statistical analyses were conducted in RevMan 5.4.1 and the variables were analyzed as

continuous via standard mean difference with a 95% confidence interval (CI).

Results

Summary of included studies

Initially, a total of 17,991 studies were collected from the databases and 8 from other sources (Fig

1). After removing duplicates, 12,548 remained for screening. Based on eligibility criteria, 11,369

studies were excluded after titles and abstracts screening. The majority of the excluded studies

focused on other animals than fish (n = 5,805) followed by those not addressing the topic of tilapia

(n = 2,853) and seaweed or soybean (n = 2,351). Then, 1,179 full texts were assessed and 1,122

were excluded because most studies used probiotics to ferment soybean/seaweed (n = 910) and due

to a lack of data on Specific Growth Rate or Feed Conversion Rate (n = 122). After full-text

screening, 57 studies were included in the qualitative synthesis and 24 were included in the meta-

analysis.

The included studies described that the fish were in good health and were purchased from

companies that regularly conduct health inspections. These studies exhibited variations in

experimental design, duration of the experiment, manner of processing soybeans, species of

seaweed used in the experiment, and dietary inclusion level. In all the included papers, soybeans

and seaweed were studied individually and in this review were analyzed separately.

Fig 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)

flowchart of the study selection process. *n* represents the number of studies.

Quality and risk of bias assessment

The quality and risk of bias of included studies were assessed through the Cochrane criteria

showing positive results for allocation concealment (selection bias), incomplete outcome data

(attrition bias), selective reporting (reporting bias), and other bias (sample size and experiment

duration) (Fig 2A and 2B). The evaluation of the studies revealed unclear risk bias for domain

random sequence bias (selection bias) because most of the studies reported the random method but

the sequence generation was not described. The studies Abdel-Warith et al., (2013), El-Saidy et al.,

(2010) and Sharda et al., (2017) (Fig 3A) and Stadtlander et al., (2012), Radwan et al., (2022) and

Eissa et al., (2021) (Fig 3B) revealed high-risk bias for random sequence bias (selection bias). The

study by Stadtlander et al., (2012) showed unclear bias for other bias because the sample size was

not enough for this systematic review but the experiment duration was reported.

Fig 2. **The risk of bias graph shows the review author's judgment about each risk of bias**

item presented as percentages across all included studies on soybean (A) and seaweed (B).

The quality and risk of bias was assessed using the Cochrane risk-of-bias tool in Cochrane

Collaboration's software RevMan 5.4.1

Fig 3. **Risk of bias summary showing review author's judgment about each risk of bias item**

for each included study on soybean (A) and seaweed (B). The quality and risk of bias was

assessed using the Cochrane risk-of-bias tool in Cochrane Collaboration's software RevMan 5.4.1

Effect of soybean on growth performance and feed utilization

In this study, Specific Growth Rate (SGR) and Feed Conversion Rate (FCR) were used as

parameters of growth performance and feed utilization, respectively. Among the studies included

in the meta-analysis, twelve [11,16,30-39] described the effects of soybean on growth performance

and feed utilization of tilapia (*Oreochromis niloticus* and *Sarotherodon melanotheron*) and

involved 8,055 fish and 53 treatments. One article (Koumi et al., 2011) studied 2 species of tilapia,

and the analysis was performed separately. The results showed that the random pooled effect size

of SGR was -2.14 (95% CI: -2.92, -1.37; $p < 0.00001$; $I^2 = 99\%$) (Fig 4A) and for FCR was 1.80

(95% CI: 0.72, 2.89; $p = 0.001$; $I^2 = 100\%$) (Fig 4B). The average inclusion level of soybeans in

the diets, calculated from the recommendation in each analyzed study (S3 Table), was found to be

59,4%. These findings suggest that including soybeans at this level can enhance the SGR but does

not lead to an improvement in FCR.

Fig 4. Forest plot showing the effect of soybean on specific growth rate (A) and feed

conversion rate (B) of tilapia. The analysis was performed on RevMan 5.4.1, based on twelve

studies and the effect size index is the standardized mean difference. The random-effects model

was employed for this analysis and the studies in the analysis are assumed to be a random sample

from a universe of potential studies. The variables were analyzed as continuous via standard mean

difference with a 95% CI. ***O.n***: *Oreochromis niloticus*; ***S.m***: *Sarotherodon melanotheron*; **CI**:

Confidence Interval; **Tau²**: tau-squared; **df**: difference; **Chi²**: chi-squared; ***P:*** *p*-value; **I²:** squared

statistic test; **Z**: z-value; **SD** and **Std**: Standard Deviation; *same study but carried out on two

species of tilapia (*O. niloticus and S. melanotheron*).

Effect of seaweed on growth performance and feed utilization

Among the studies included in the meta-analysis, twelve [8,40-50] described the effects of

seaweed on growth performance and feed utilization of *O. niloticus* and hybrid red tilapia (*O.*

niloticus and *Oreochromis mossambicus*), involving 5,657 fish and 55 treatments of different

species including rede seaweed (*Laurencia obtusa*, *Pterocladia capillacea*; *Gracillaria arcuata*,

Porphyra yezoensis, *Gracilaria vermiculophylla,* and *Porphyra dioica*); green seaweed (*Ulva*

lactuca, *Enteromorpha Flaxusa,* and *Ulva Fasciata*) and brown seaweed (*Taonia atomaria* and

Cystoseira myrica) (S3 Table). Two papers (Silva et al., 2014 and Eissa et al., 2020) each studied three species of seaweed while one paper (Khalafalla and El-Hais, 2015) studied 2 species of seaweed, and the analysis was performed separately for each species. The results showed that the random pooled effect size of SGR was -0.74 (95% CI: -1.70, 0.22; $p = 0.13$; $I^2 = 99\%$) (Fig 5A), and for FCR was -0.70 (95% CI: -1.94, 0.54; $p = 0.27$; $I^2 = 100\%$) (Fig 5B). The inclusion level of seaweed was 15,9%. These results indicate that the inclusion level of seaweed at 15,9% did not improve the SGR and FCR of tilapia.

Fig 5. Forest plot showing the effect of seaweed on specific growth rate (A) and feed conversion rate (B) of tilapia. The analysis was performed on RevMan 5.4.1, based on seventeen studies and the effect size index is the standardized mean difference. The random-effects model was employed for this analysis and the studies in the analysis are assumed to be a random sample from a universe of potential studies. The variables were analyzed as continuous via standard mean difference with a 95% CI. **GRA**: *Gracillaria*; **UV**: *Ulva*; LA: *Laurencia*; POR: *Porphyra*; CY: *Cystoseira*; **CI**: Confidence Interval; **Tau^2**: tau-squared; **df**: difference; **Chi^2**: chi-squared; ***P:*** *p*-value; **I^2:** squared statistic test; **Z**: z-value; **SD** and **Std**: Standard Deviation; **same study but carried out on different species of seaweed.

Effect of soybean and seaweed on the gut microbiota of tilapia

The search conducted in the databases using the search strategies defined in this systematic review on tilapia was not able to identify studies that met the inclusion criteria on this specific topic. Most of the studies were done on other farmed fish (Table 1), showing that soybean and seaweed could have different effects on gut microbiota composition and diversity.

Table 1. The most abundant gut microbiota in other fish than tilapia fed with soybean and seaweed-based diet

Type of diet	Inclusion level	Fish species	Composition of gut microbiota	Reference
Soybean meal SBM)	25%	Atlantic salmon (*Salmo salar*)	*Vibrio spp., Arthrobacter agillis, Brachybacterium spp., Kocuria carniphila* and *Rhodococcus spp*	[51]
Green Seaweed (*Ulva ohnoi*)	5%	Flatfish (*Solea senegalensis*)	*Vibrionaceae* and *Shewanella*	[52]

253 **Table 1**. Continued

Type of diet	Inclusion level	Fish species	Composition of gut microbiota	Reference
Red Seaweed (*Gracilaria gracilis*)	8%	European seabass (*Dicentrarchus labrax*)	*Acinetobacter, Corynebacterium, Ilumatobacter, Kordia, Methylobacterium, Polaribacter, Pseudomonas* and *Sulfitobacter*	[53]
Red seaweed (*Asparagopsis taxiformis*)	3%	Atlantic salmon (*Salmo salar*)	*Aeromodaceae*	[54]
Soybean meal	10%	Atlantic salmon (*Salmo salar*)	*Psychrobacter okhotskensis, Arthrobacter rhombi, Arthrobacter psychrolactophilus, Psychrobacter cibarius* and *Psychrobacter fozii*	[55]
Red Seaweed (*Gracilaria* cornea) and green seaweed (*Ulva rigida*)	5 %, 15% and 25%	Gilthead seabream (*Sparus aurata*) juveniles	**Gracillaria** (15% and 25%): *Pseudomonas* and *Lactobacillus*; **Ulva** (5% and 15%): *Lactobacillus and Sphingomonas*; **Ulva** (25%): *Comamonas aquatica*	[56]
Soybean meal (SBM)	20%	Atlantic salmon (*Salmo salar* L.)	*Lactococcus lactis subsp. lactis, Weissella confusa,* and *Photobacterium phosphoreum*	[57]
Soybean meal (SBM)	50%	Atlantic salmon (*Salmo salar*)	*Aeromonas* and *Sporosarcina aquimarina*	[58]
Soybean meal (SBM)	50%	Rainbow trout (*Oncorhynchus mykiss*)	*Psychrobacter spp* and *Saccharomyces spp*	[59]
Soybean Meal (SBM)	20% – 100%	*Carassius auratus gibelio* ♀ + *Cyprinus carpio* ♂	*Cetobacterium* (10.53–17.46 %) and *Aeromonas* (4.68–13.88 %)	[60]
Green seaweed (*Ulva rigida*)	10%	Atlantic cod (*Gadus morhua*)	*Ruminococcaceae spp., Shewanella, Tyzerella, Rikenellaceae spp., Peptococcaceae, Bacteroides* and *Fusobacterium.*	[61]
Green seaweed (*Ulva rigida*)	25%	*Sparus aurata* juveniles	*Lactobacillus delbrueckii*	[62]
Soybean meal (SBM)	45%	Juvenile rainbow trout (*Oncorhynchus mykiss*)	*Aeromonas, Sphingomonas* and *Chryseomonas*	[63]
Red seaweed (*Gracilaria gracilis*)	2.5% and 5%	European seabass (*Dicentrarchus labrax*)	*Sphingomonas, Photobacterium, Staphylococcus* and *Vibrio*	[64]

Soybean Protein Isolate (SPI)	25% and 50%	Spotted seabass (*Lateolabrax maculatus*)	*Prevotella, Succiniclasticum,* and *Desulfovibrio*	[65]
Green seaweed (*Ulva ohnoi*)	5%	*Solea senegalensis*	*Pseudomonas* and *Mycoplasmataceae*	[66]

Discussion

Effect of soybean and seaweed-based diet on SGR and FCR

The findings of this review suggest an enhancement in SGR among tilapia fed with soybean-based

diets. This improvement could be attributed to soybean's high protein content and relatively

balanced amino acid profile [33,34]. The findings also reveal that soybean did not improve FCR,

likely due to the presence of antinutritional factors such as protease inhibitors, lectins, phytic acid,

saponins, phytoestrogens, antivitamins, and allergens [16,17]. These factors may compromise the

palatability of soybeans and thus affect their effectiveness in improving FCR. The results from

studies examining the impact of soybeans suggest that growth performance and feed utilization of

tilapia tend to decline as the inclusion levels of soybeans increase, the ideal level of inclusion

being up to 50% [31,35]. Similar studies conducted on other fish species such as *Liza*

haematocheila [67]; *Heteropneustes fossilis* [68]; *Takifugu obscurus* [69]; *Clarias gariepinus* [70],

Solea aegyptiaca [71]; *Sparus aurata* [72]; also indicate a decrease in growth parameters with

higher levels of soybean inclusion level. However, determining the optimal inclusion level and

type of soybean (e.g., defatted, fermented, full fat) requires further investigation since the studies

included in the meta-analysis utilized different inclusion levels and types of soybeans. On the other

hand, the results do not support the use of seaweed as an alternative protein source for tilapia, as it

did not improve the SGR and FCR. However, it should be noted that the studies included in the

meta-analysis used different species of seaweeds and different levels of inclusion.

Several studies have demonstrated that seaweed inclusion levels of up to 20% can effectively

replace fishmeal in tilapia diets without compromising the growth performance and feed utilization

[40,41,43,48,50,73]. Similar outcomes have been observed in other fish species as well [74-76].

The bioactive compounds present in soybeans and seaweed, such as Non-Starch Polysaccharides (NSPs) and Antinutritional Factors (ANFs) like saponins, lectins, and phytates, play significant roles in

influencing fish digestibility and metabolic functions [77,78]. Excessive NSPs may cause gut inflammation and reduce energy availability, while ANFs can interfere with digestive enzymes and nutrient absorption [79]. However, proper processing methods such as heat and enzymatic treatment, bioprocessing, and fermentation of these diets can mitigate these effects [18,79,80]. Furthermore, phytochemicals like isoflavones in soybeans possess antioxidant properties, although their impact may vary among fish species [81]. Additionally, certain seaweed compounds, like polysaccharides may promote gut health, enhancing overall digestion and nutrient absorption [82,83]. However, the specific impact can vary depending on the type and amount of seaweed, as well as the fish species involved [15,84]. For instance, several studies have demonstrated that the green seaweed *Ulva spp*, when included at levels of up to 5%, can enhance the growth performance of certain carnivorous fish [85,86], and omnivorous fish like tilapia of up to 20% of inclusion [87]. In addition to soybeans and seaweed, plant-based diets such as sunflower, canola, cottonseed, Moringa, and *Azolla* meals have shown promise in feeding tilapia and other fish [88-92]. *Azolla*, noted for its exceptional amino acid profile, exhibits the highest nutritional index in tilapia feed [93]. In the data analysis, heterogeneity was observed among the included studies, potentially due to variations in experimental design, number of fish used in each study, type of soybean, and species of seaweed used across different studies. Subgroup analyses were performed to explore potential sources of heterogeneity, but the findings were inconclusive.

Effect of soybean and seaweed-based diet on gut microbiota

In this review, it was not possible to find studies that describe the effect of soybean or seaweed on the gut microbiota of tilapia. However, studies in other fish species have shown diverse effects on the gut microbiota composition. For instance, beneficial bacteria like *Shewanella* were observed in

Flatfish and Atlantic cod fed with *Ulva spp* [52,61], while *Lactobacillus spp* were found in Gilthead seabream fed with *Gracilaria cornea* and *Ulva rigida* [56,62] along with *Lactococcus spp* [57]. Additionally, *Psychrobacter spp, and Arthrobacter spp* were noted in fish fed with soybean meal [51,55,59]. These microorganisms are known to possess probiotic properties, which could potentially enhance nutrient absorption and disease resistance [94-97]. The seaweed and soybean dietary components contain bioactive compounds such as isoflavones, saponins, and oligosaccharides in soybean [98-100], and polysaccharides, polyphenols, phycobiliproteins, and phlorotannins in seaweed [101-103]. These compounds can exert prebiotic effects, influencing the composition of the gut microbiota [104]. Prebiotics are essentially non-digestible substances that, when consumed, foster beneficial changes in the gut microbiome, promoting the development of a more favorable microbial community [105]. This shift can have positive implications for various aspects of health, including digestive function, immune system modulation, and antimicrobial properties, inhibiting the growth of potentially harmful bacteria [106-108]. It is important to note that the specific effects of soybean and seaweed on fish gut microbiota can vary depending on various factors, such as the fish species, the inclusion levels of these ingredients, and the species of seaweed. Studies investigating the utilization of various woody forages such as *Moringa oleifera*, fermented *M. oleifera*, *Folium mori*, and fermented *F. mori* meals in tilapia farming have indicated significant variations in gut microbiota composition [109,110]. These findings suggest that diets incorporating soybean, seaweed, and other plant-based diets may have diverse effects on the diversity and composition of fish intestinal microbiota.

Implications for tilapia production

The inclusion of soybean and seaweed-based diets into tilapia farming can contribute to the sustainability of the aquaculture industry by reducing reliance on fishmeal and fish oil. The improved SGR observed in tilapia fed with a soybean diet indicates the potential for cost-effectiveness and viability in tilapia production. Regarding seaweed-based diets, although they

may not directly improve growth performance and feed utilization parameters in tilapia, they offer
alternative feed options due to their rich content of bioactive compounds. These compounds can
have beneficial effects on fish growth and health. Additionally, the modulation of gut microbiota
by soybean and seaweed-based diets can improve tilapia health and nutrient utilization leading to
more sustainable and cost-effective tilapia production.

Limitations and future research directions

Despite the valuable insights gained from this study, limitations should be considered: (1) the
included studies may have exhibited variations in experimental design, type of soybean and
seaweed used, leading to heterogeneity in the data, (2) limitations for some outcomes, particularly
effect of soybean and seaweed on gut microbiota of tilapia. For future research, it is suggested: (1)
investigations into the optimal inclusion levels of soybean and seaweed-based ingredients and
exploring the potential seaweed species for tilapia diets should be conducted to maximize growth
performance and feed utilization; (2) long-term studies are needed to evaluate the potential impact
of these diets on the gut microbiota and its implications for growth performance, nutrient
utilization, and the health of tilapia and (3) investigate the effect of those diets in other species of
tilapia.

Conclusion

This systematic review and meta-analysis provide valuable insights into the effects of soybean and
seaweed-based diets on tilapia production. The results of the meta-analysis indicated that the
soybean inclusion level of 59.4% improved SGR but did not improve the FCR of tilapia. On the
other hand, the seaweed-based diet did improve both SGR and FCR at an inclusion level of 15.9%.
Based on the search strategies defined in this review, it was not possible to find studies that met the
inclusion criteria for the gut microbiota of tilapia. However, studies on soybean and seaweed in
other farmed fish have shown diverse effects on the gut microbiota composition and have the
potential to enhance the growth of beneficial gut microbiota and improve fish health. This

information is valuable for tilapia production as it offers an alternative dietary strategy for optimizing their growth.

References

1.   Arumugam M, Jayaraman S, Sridhar A, Venkatasamy V, Brown PB, Abdul Kari Z, et al. Recent Advances in Tilapia Production for Sustainable Developments in Indian Aquaculture and Its Economic Benefits. Fishes. 2023;8(4):1-31.

2.   FAO. Social and economic performance of tilapia farming in Brazil. Junning Cai KQ and NH, editor. Vol. 1130, Social and economic performance of tilapia farming in Brazil. Rome, Italy: FAO; 2017. 144 p.

3.   Prabu E, Rajagopalsamy CBT, Ahilan B, Jeevagan IJMA, Renuhadevi M. Tilapia – An Excellent Candidate Species for World Aquaculture: A Review. Annu Res Rev Biol. 2019;31(3):1–14.

4.   Ng WK, Romano N. A review of the nutrition and feeding management of farmed tilapia throughout the culture cycle. Rev Aquac. 2013;5(4):220–254.

5.   El-Sayed AFM. Nutrition and feeding. Tilapia Culture. 2020. 135–172 p.

6.   Norambuena F, Hermon K, Skrzypczyk V, Emery JA, Sharon Y, Beard A, et al. Algae in fish feed: Performances and fatty acid metabolism in juvenile Atlantic Salmon. PLoS One. 2015;10(4):1–17.

7.   Yildirim-Aksoy M, Lim C, Li MH, Klesius PH. Interaction between dietary levels of vitamins C and E on growth and immune responses in channel catfish, *Ictalurus punctatus* (Rafinesque). Aquac Res. 2008 Aug 1;39(11):1198–1209.

8.   Younis ESM, Al-Quffail AS, Al-Asgah NA, Abdel-Warith AWA, Al-Hafedh YS. Effect of dietary fish meal replacement by red algae, *Gracilaria arcuata*, on growth performance and body composition of Nile tilapia Oreochromis niloticus. Saudi J Biol Sci. 2018;25(2):198–203.

9.   Naylor RL, Hardy RW, Bureau DP, Chiu A, Elliott M, Farrell AP, et al. Feeding

aquaculture in an era of finite resources. Proc Natl Acad Sci U S A. 2009;106(36):15103–15110.

10.   Olsen RL, Hasan MR. A limited supply of fishmeal: Impact on future increases in global

aquaculture production. Trends Food Sci Technol. 2012;27(2):120–128.

11.   Zhao H, Jiang R, Xue M, Xie S, Wu X, Guo L. Fishmeal can be completely replaced by

soy protein concentrate by increasing feeding frequency in Nile tilapia (*Oreochromis niloticus*

GIFT strain) less than 2 g. Aquac Nutr. 2010;16(6):648–653.

12.   Kook MC, Cho SC, Hong YH, Park H. *Bacillus subtilis* fermentation for enhancement of

feed nutritive value of soybean meal. J Appl Biol Chem. 2014;57(2):183–188.

13.   Yun A, Kim J, Jeong HS, Lee KW, Kim HS, Kim PY, et al. Inclusion effect of soybean

meal, fermented soybean meal, and *Saccharina japonica* in extruded pellet for juvenile abalone

(*Haliotis discus*, Reeve 1846). Fish Aquat Sci. 2018;21(1):1–8.

14.   Rajauria G. Seaweeds: a sustainable feed source for livestock and aquaculture. In: Tiwari

B, TroyD(eds) Seaweed sustainability: food and non-food applications. Seaweed Sustainability.

Elsevier Inc.; 2015. 389–420 p.

15.   Pereira R, Valente LMP, Sousa-Pinto I, Rema P. Apparent nutrient digestibility of

seaweeds by rainbow trout (*Oncorhynchus mykiss*) and Nile tilapia (*Oreochromis niloticus*). Algal

Res. 2012;1(1):77–82.

16.   Abdel-Warith A-WA-m., Younis EM, Al-Asgah NA. Influence of Dietary Inclusion of

Full-Fat Soybean Meal and Amino Acids Supplementation on Growth and Digestive Enzymes

Activity of Nile Tilapia, *Oreochromis niloticus*. Turkish J Fish Aquat Sci. 2013;13:69–77.

17.   Hussain SM, Afzal M, Rana SA, Javid A, Iqbal M. Effect of phytase supplementation on

growth performance and nutrient digestibility of *Labeo rohita* fingerlings fed on corn gluten meal-

based diets. Int J Agric Biol. 2011;13(6):916–922.

18.   Samtiya M, Aluko RE, Dhewa T. Plant food anti-nutritional factors and their reduction

strategies: an overview. Food Prod Process Nutr. 2020;2(1):1–14.

19.   Huang Q, Sham RC, Deng Y, Mao Y, Wang C, Zhang T, et al. Diversity of gut

microbiomes in marine fishes is shaped by host-related factors. Mol Ecol. 2020;29(24):5019–5034.

20.   Nie L, Zhou QJ, Qiao Y, Chen J. Interplay between the gut microbiota and immune

responses of ayu (*Plecoglossus altivelis*) during Vibrio anguillarum infection. Fish Shellfish

Immunol. 2017;68:479–487.

21.   Xiong JB, Nie L, Chen J. Current understanding of the roles of gut microbiota in fish

disease and immunity. Zool Res. 2019;40(2):70–76.

22.   Leitão-Gonçalves R, Carvalho-Santos Z, Francisco AP, Fioreze GT, Anjos M, Baltazar

C, et al. Commensal bacteria and essential amino acids control food choice behavior and

reproduction. PLoS Biol. 2017;15(4):1–29.

23.   Yamazaki Y, Meirelles PM, Mino S., Suda KW., Hattori M, Thompson FL., Sakai Y, TS.

Individual *Apostichopus japonicus* fecal microbiome reveals a link with polyhydroxybutyrate

producers in host growth gaps. Sci Rep. 2016; 6(February):1–10

24.   Huang C, Liao W. Seasonal Variation in Gut Microbiota Related to Diet in *Fejervarya*

limnocharis. Animals. 2021;1–15.

25.   Piazzon MC, Calduch-Giner JA, Fouz B, Estensoro I, Simó-Mirabet P, Puyalto M, et al.

Under control: how a dietary additive can restore the gut microbiome and proteomic profile, and

improve disease resilience in a marine teleostean fish fed vegetable diets. Microbiome.

2017;5(1):1-23.

26.   Ringø E, Zhou Z, Vecino JLG, Wadsworth S, Romero J, Krogdahl Å, et al. Effect of

dietary components on the gut microbiota of aquatic animals. A never-ending story? Aquac Nutr.

2016 Apr 1;22(2):219–282.

27.   Wang T, Wu HX, Li WJ, Xu R, Qiao F, Du ZY, et al. Effects of dietary mannan

oligosaccharides (MOS) supplementation on metabolism, inflammatory response, and gut

microbiota of juvenile Nile tilapia (*Oreochromis niloticus*) fed with high carbohydrate diet. Fish

Shellfish Immunol. 2022 Nov;130:550–559.

28.   Higgins J, Green S (Sally E, Cochrane Collaboration. Assessing risk of bias in included

studies Cochrane handbook for systematic reviews of interventions. Cochrane, editor. 2017. 2017.

1p.

29.   Julian P T Higgins, Simon G Thompson, Jonathan J Deeks DGA. Measuring

inconsistency in meta-analyses. BMJ 2003;327. 2003;327(2):557–560.

30.   Agbo NW, Madalla N, Jauncey K. Mixtures of oilseed meals as dietary protein sources in

diets of juvenile Nile tilapia (*Oreochromis niloticus* L.). J Sci Technol. 2016;35(3):11-24.

31.   Ahmad N, Siddiqui PJA, Khan KM, Ali A, Tahir M, Akbar NU, et al. Effects of partial

substitution of fishmeal by soybean meal in Nile tilapia (*Oreochromis niloticus*) diet. J Anim Plant

Sci. 2020;30(2):364–370.

32.   Azaza MS, Kammoun W, Abdelmouleh A, Kraïem MM. Growth performance, feed

utilization, and body composition of Nile tilapia (*Oreochromis niloticus* L.) fed with differently

heated soybean-meal-based diets. Aquac Int. 2009;17(6):507–521.

33.   El-Saidy DMSD. Effect of using okara meal, a by-product from soymilk production as a

dietary protein source for Nile tilapia (*Oreochromis niloticus* L.) mono-sex males. Aquac Nutr.

2011;17(4):380–386.

34.   Lin S, Luo L. Effects of different levels of soybean meal inclusion in replacement for fish

meal on growth, digestive enzymes and transaminase activities in practical diets for juvenile

tilapia, *Oreochromis niloticus×O. aureus*. Anim Feed Sci Technol. 2011;168(1):80–87.

35.   Koumi AR, Koffi KM, Atsé BC, Kouame LP. Growth, feed efficiency and carcass

mineral composition of *Heterobranchus longifilis*, *Oreochromis niloticus* and *Sarotherodon*

melanotheron juveniles fed different dietary levels of soybean meal-based diets. African J

Biotechnol. 2011;10(66):14990–14998.

36.   Koumi AR, Atse BC, Kouame LP. Utilization of soya protein as an alternative protein

source in *Oreochromis niloticus* diet: Growth performance, feed utilization, proximate

composition and organoleptic characteristics. African J Biotechnol. 2009;8(1):91–97.

37.   Obirikorang KA, Gyamfi S, Goode ME, Amisah S, Edziyie RE, Quagrainie K, et al.

Effect of soybean meal diets on the growth performance, ammonia excretion rates, gut histology

and feed cost of Nile tilapia (*Oreochromis niloticus*) fry. Aquac Res. 2020 Sep 1;51(9):3520–3532.

38.   Sharda S, Saini V. Replacement of fishmeal with soybean meal in Nile tilapia

(*Oreochromis niloticus*) Diet. J Entomol Zool Stud. 2017;5(4):845–849.

39.   Yigit N. O. and Demir T. Use of Fermented Soybean Meal with Whey as a Protein

Source for Feeding Juvenile Tilapia (*Oreochromis niloticus*). Isr J Aquac - Bamidgeh.

2010;(January):1–7

40.   Ashour M, Mabrouk MM, Ayoub HF, El-Feky MMMM, Zaki SZ, Hoseinifar SH, et al.

Effect of dietary seaweed extract supplementation on growth, feed utilization, hematological

indices, and non-specific immunity of Nile Tilapia, *Oreochromis niloticus* challenged with

Aeromonas hydrophila. J Appl Phycol. 2020;32(5):3467–3479.

41.   Eissa HA, Hegazi MM, Elmor ME, Sharawy ZZ. Effects of partial substitution of fish

meal with different levels of marine macroalgae species on growth indices and RNA/DNA ratio of

hybrid red tilapia. Egypt J Aquat Biol Fish. 2021;25(3):395-409.

42.   Saleh NE, Shalaby SM, Sakr EM, Elmonem AIA, Michael FR. Effect of dietary inclusion

of *Ulva fasciata* on red hybrid tilapia growth and carcass composition. J Appl Aquac.

2014;26(3):197–207.

43.   Saleh HHE sayed. Can Artificial Feed be Replace by Fresh Macro Algae (*Enteromorpha*

flaxusa) in Hybrid Red Tilapia (*Oreochromis mossambicus* × *Oreochromis niloticus*) Juvenile

Nutrition☐? J Oceanogr Mar Res. 2020;8(2):1–7.

44.   EL-Sayed Hussein EM, Ebtehal EL-Sayed Hussein CM. Effect of seaweed supplemented

diets on Nile tilapia, *Oreochromis niloticus* performance. Int J Fish Aquat Stud. 2017;5(2):205–

210.

45.   Marinho G, Nunes C, Sousa-Pinto I, Pereira R, Rema P, Valente LMP. The IMTA-

cultivated Chlorophyta *Ulva spp*. as a sustainable ingredient in Nile tilapia (*Oreochromis niloticus*)

diets. J Appl Phycol. 2013;25(5):1359–1367.

46.   Radwan M, El-Sharkawy MA, Negm MA, Mohammadein A, Malki JSA, Al-Thomali

AW, et al. Dual effect of dietary seaweed of extract nanoparticles (GNS) with bionanocomposite

cellulose acetate membranes (CA/bio-AgNps) on growth performance and health status of the Nile

tilapia (*Oreochromis niloticus*): Specification on feed utilization, immune s. Front Mar Sci.

2022;9(October):1–18.

47.   Salem MES, Abdel-Ghany HM, Almisherfi HM. Role of dietary *Laurencia obtusa* in

enhancing growth, blood indices, and hypoxia resistance of red tilapia (*Oreochromis niloticus* x *O.*

mossambicus). J Appl Phycol. 2021;33(4):2617–2628.

48.   Silva DM, Valente LMP, Sousa-Pinto I, Pereira R, Pires MA, Seixas F, et al. Evaluation

of IMTA-produced seaweeds (Gracilaria, Porphyra, and Ulva) as dietary ingredients in Nile tilapia,

Oreochromis niloticus L., juveniles. Effects on growth performance and gut histology. J Appl

Phycol. 2015;27(4):1671–1680.

49.   Stadtlander T, Khalil WKB, Focken U, Becker K. Effects of low and medium levels of

red alga Nori (*Porphyra yezoensis* Ueda) in the diets on growth, feed utilization and metabolism in

intensively fed Nile tilapia, *Oreochromis niloticus* (L.). Aquac Nutr. 2013;19(1):64–73.

50.   Khalafalla MM and El-Hais A-e.M.A. Evaluation of Seaweeds *Ulva rigida* and

P*terocladia capillaceaas* Dietary Supplements in Nile Tilapia Fingerlings. J Aquac Res Dev.

2015;06(03):1-5.

51.   Bakke-McKellep AM, Penn MH, Salas PM, Refstie S, Sperstad S, Landsverk T, et al.

Effects of dietary soyabean meal, inulin and oxytetracycline on intestinal microbiota and epithelial

cell stress, apoptosis and proliferation in the teleost Atlantic salmon (*Salmo salar* L.). Br J Nutr.

2007;97(4):699–713.

52.   Cerezo IM, Fumanal M, Tapia-Paniagua ST, Bautista R, Anguís V, Fernández-Díaz C, et

al. Solea senegalensis Bacterial Intestinal Microbiota Is Affected by Low Dietary Inclusion of

Ulva ohnoi. Front Microbiol. 2022;12(February):1–13.

53.   Ferreira M, Abdelhafiz YAA, Abreu H, Silva J, Valente LMP, Viswanath K. *Gracilaria*

gracilis and *Nannochloropsis oceanica*, singly or in combination, in diets alter the intestinal

microbiota of European seabass (Dicentrarchus labrax). Front. Mar. Sci. 2022;9:1-16

54.   Thépot V, Campbell AH, Rimmer MA, Jelocnik M, Johnston C, Evans B, et al. Dietary

inclusion of the red seaweed Asparagopsis taxiformis boosts production, stimulates immune

response and modulates gut microbiota in Atlantic salmon, *Salmo salar*. Aquaculture. 2022;546:1-

13

55.   Ringø E, Sperstad S, Kraugerud OF, Krogdahl Å. Use of 16S rRNA gene sequencing

analysis to characterize culturable intestinal bacteria in Atlantic salmon (*Salmo salar*) fed diets

with cellulose or non-starch polysaccharides from soy. Aquac Res. 2008;39(10):1087–1100.

56.   Rico RM, Tejedor-Junco MT, Tapia-Paniagua ST, Alarcón FJ, Mancera JM, López-

Figueroa F, et al. Influence of the dietary inclusion of *Gracilaria cornea* and *Ulva rigida* on the

biodiversity of the intestinal microbiota of Sparus aurata juveniles. Aquac Int. 2016;24:965–984.

57.   Reveco FE, Øverland M, Romarheim OH, Mydland LT. Intestinal bacterial community

structure differs between healthy and inflamed intestines in Atlantic salmon (*Salmo salar* L.).

Aquaculture. 2014;420–421:262–269.

58.   Navarrete P, Fuentes P, De la Fuente L, Barros L, Magne F, Opazo R, et al. Short-term

effects of dietary soybean meal and lactic acid bacteria on the intestinal morphology and

microbiota of Atlantic salmon (*Salmo salar*). Aquac Nutr. 2013;19(5):827–836.

59.   Merrifield DL, Dimitroglou A, Bradley G, Baker RTM, Davies SJ. Soybean meal alters

autochthonous microbial populations, microvilli morphology and compromises intestinal

enterocyte integrity of rainbow trout, *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2009 Sep

1;32(9):755–766.

60.   Liu X, Han B, Xu J, Zhu J, Hu J, Wan W, et al. Replacement of fishmeal with soybean

meal affects the growth performance, digestive enzymes, intestinal microbiota and immunity of

Carassius auratus gibelio♀ × *Cyprinus carpio*♂. Aquac Reports. 2020;18:1-9.

61.   Keating C, Bolton-Warberg M, Hinchcliffe J, Davies R, Whelan S, Wan AHL, et al.

Temporal changes in the gut microbiota in farmed Atlantic cod (*Gadus morhua*) outweigh the

response to diet supplementation with macroalgae. Anim Microbiome. 2021;3:1–21.

62.   Abdala-Díaz RT, García-Márquez J, Rico RM, Gómez-Pinchetti JL, Mancera JM,

Figueroa FL, et al. Effects of a short pulse administration of *Ulva rigida* on innate immune

response and intestinal microbiota in Sparus aurata juveniles. Aquac Res. 202;52(7):3038–3051.

63.   Heikkinen J, Vielma J, Kemiläinen O, Tiirola M, Eskelinen P, Kiuru T, et al. Effects of

soybean meal based diet on growth performance, gut histopathology and intestinal microbiota of juvenile rainbow trout (*Oncorhynchus mykiss*). Aquaculture. 2006;261(1):259–268.

64. Gonçalves AT, Simões M, Costa C, Passos R, Baptista T. Modulatory effect of *Gracilaria gracilis* on European seabass gut microbiota community and its functionality. Sci Rep. 2022;12(1):1-16.

65. Wang Z, Yang L, Zhang C, Li X, Wang L, Lu K, et al. Substituting dietary fishmeal with soybean meal isolate influences hepatic lipid metabolism and gut microbiota in spotted seabass (*Lateolabrax maculatus*). Isr J Aquac - Bamidgeh. 2021;73:1–13.

66. Tapia-Paniagua ST, Fumanal M, Anguís V, Fernández-Díaz C, Alarcón FJ, Moriñigo MA, et al. Modulation of intestinal microbiota in *Solea senegalensis* fed low dietary level of *Ulva ohnoi*. Front Microbiol. 2019;10:1-16.

67. Liu T, Han T, Wang J, Liu T, Bian P, Wang Y, et al. Effects of replacing fish meal with soybean meal on growth performance, feed utilization and physiological status of juvenile redlip mullet *Liza haematocheila*. Aquac Reports. 2021;20:1-10.

68. Howlader S, Sumi KR, Sarkar S, Billah SM, Ali ML, Howlader J, et al. Effects of dietary replacement of fish meal by soybean meal on growth, feed utilization, and health condition of stinging catfish, *Heteropneustes fossilis*. Saudi J Biol Sci. 2023;30(3):1-7.

69. Ye H, Xu M, Liu Q, Sun Z, Zou C, Chen L, et al. Effects of replacing fish meal with soybean meal on growth performance, feed utilization and physiological status of juvenile obscure puffer, *Takifugu obscurus*. Comp Biochem Physiol Part - C Toxicol Pharmacol. 2019;216(November):75–81.

70. Abdel-Warith AA, Younis EM, Al-Asgah NA, Mahboob S. Effect of replacing fish meal by full fat soybean meal on growth performance, feed utilization and gastrointestinal enzymes in diets for african catfish *Clarias gariepinus*. Brazilian J Biol. 2020;80(3):535–543.

71. Bonaldo A, Roem AJ, Pecchini A, Grilli E, Gatta PP. Influence of dietary soybean meal levels on growth, feed utilization and gut histology of Egyptian sole (*Solea aegyptiaca*) juveniles. Aquaculture. 2006;261(2):580–586.

72.   Kokou F, Rigos G, Henry M, Kentouri M, Alexis M. Growth performance, feed

utilization and non-specific immune response of gilthead sea bream (*Sparus aurata* L.) fed graded

levels of a bioprocessed soybean meal. Aquaculture . 2012;364–365:74–81.

73.   Natify W, Droussi M, Berday N, Araba A, Benabid M. Effect of the seaweed Ulva

lactuca as a feed additive on growth performance, feed utilization and body composition of Nile

tilapia (*Oreochromis niloticus* L.). Int J Agron Agric Res. 2015;7(3):85–92.

74.   Sulaeman HA, Zainuddin, Laining A. Evaluation of seaweed, *Ulva lactuca* as a fresh diet

for nursery stage of golden rabbitfish, *Siganus guttatus*. IOP Conf Ser Earth Environ Sci.

2022;1119(1):1-11.

75.   Madibana MJ, Mlambo V, Lewis B, Fouché C. Effect of graded levels of dietary seaweed

(*Ulva sp.*) on growth, hematological and serum biochemical parameters in dusky kob,

Argyrosomus japonicus, sciaenidae. Egypt J Aquat Res. 2017;43(3):249–254.

76.   Seyedalhosseini SH, Salati AP, Mozanzadeh MT, Parrish CC, Shahriari A. Effects of

dietary seaweeds (*Gracilaria spp.* and *Sargassum spp.*) on growth, feed utilization, and resistance

to acute hypoxia stress in juvenile Asian seabass (*Lates calcarifer*). Aquac Reports.

2023;31(June):1-6.

77.   Sinha AK, Kumar V, Makkar HPS, De Boeck G, Becker K. Non-starch polysaccharides

and their role in fish nutrition - A review. Food Chem. 2011;127(4):1409–1426.

78.   Francis G, Makkar HPS, Becker K. Antinutritional factors present in plant-derived

alternate fish feed ingredients and their effects in fish. Vol. 199, Aquaculture. 2001. 197–227 p.

79.   Kumar V, Sakhawat Hossain M, A. Ragaza J, Rubio Benito M. The Potential Impacts of

Soy Protein on Fish Gut Health. Soybean Hum Consum Anim Feed. 2020;(September):91-115.

80.   Raja K, Kadirvel V, Subramaniyan T. Seaweeds, an aquatic plant-based protein for

sustainable nutrition - A review. Futur Foods. 2022;5(March):1-12.

81.   Yang B, Jiang WD, Wu P, Liu Y, Zeng YY, Jiang J, et al. Soybean isoflavones improve

the health benefits, flavour quality indicators and physical properties of grass carp

(*Ctenopharygodon idella*). PLoS One. 2019;14(1):1–21.

82.   Hoseinifar SH, Jahazi MA, Nikdehghan N, Van Doan H, Volpe MG, Paolucci M. Effects

of dietary polyphenols from agricultural by-products on mucosal and humoral immune and

antioxidant responses of convict cichlid (*Amatitlania nigrofasciata*). Aquaculture. 2020;517:1-31.

83.   Jahazi MA, Hoseinifar SH, Jafari V, Hajimoradloo A, Van Doan H, Paolucci M. Dietary

supplementation of polyphenols positively affects the innate immune response, oxidative status,

and growth performance of common carp, *Cyprinus carpio* L. Aquaculture. 2020;517:1-33.

84.   Siddik MAB, Francis P, Rohani MF, Azam MS, Mock TS, Francis DS. Seaweed and

Seaweed-Based Functional Metabolites as Potential Modulators of Growth, Immune and

Antioxidant Responses, and Gut Microbiota in Fish. Antioxidants. 2023;12(12):1-34.

85.   Wassef EA, El-Sayed AFM, Sakr EM. Pterocladia (Rhodophyta) and Ulva (Chlorophyta)

as feed supplements for European seabass, *Dicentrarchus labrax* L., fry. J Appl Phycol.

2013;25(5):1369–1376.

86.   Martínez-Antequera FP, Martos-Sitcha JA, Reyna JM, Moyano FJ. Evaluation of the

inclusion of the green seaweed ulva ohnoi as an ingredient in feeds for gilthead sea bream (*Sparus

aurata*) and european sea bass (*Dicentrarchus labrax*). Animals. 2021;11(6):1–19.

87.   Azaza MS, Mensi F, Ksouri J, Dhraief MN, Brini B, Abdelmouleh A, et al. Growth of

Nile tilapia (*Oreochromis niloticus* L.) fed with diets containing graded levels of green algae ulva

meal (*Ulva rigida*) reared in geothermal waters of southern Tunisia. J Appl Ichthyol.

2008;24(2):202–207.

88.   Rahmdel KJ, Noveirian HA, Falahatkar B, Lashkan AB. Effects of replacing fish meal

with sunflower meal on growth performance, body composition, hematological and biochemical

indices of common carp (*Cyprinus carpio*) fingerlings. Fish Aquat Life. 2018;26(2):121–129.

89.   Güroy D, Güroy B, Merrifield DL, Tekinay AA, Davies SJ, Şahin I. Effects of fish oil

and partial fish meal substitution with oilseed oils and meals on growth performance, nutrient

utilization and health of the rainbow trout *Oncorhynchus mykiss*. Aquac Int. 2012;20(3):481–497.

90.   Elabd H, Soror E, El-Asely A, El-Gawad EA, Abbass A. Dietary supplementation of

Moringa leaf meal for Nile tilapia *Oreochromis niloticus*: Effect on growth and stress indices.

Egypt J Aquat Res. 2019;45(3):265–271.

91.   Yue YR, Zhou QC. Effect of replacing soybean meal with cottonseed meal on growth,

feed utilization, and hematological indexes for juvenile hybrid tilapia, *Oreochromis niloticus×O.*

aureus. Aquaculture. 2008;284(1):185–189.

92.   Ibrahim A, Abouelella S, Adam H, Hassan N, Abdallah R. Effect of Azolla pinnata on

growth performance and survival rate of fingerlings of grass carp fish *Ctenopharyngodon idellus*

(Valenciennes, 1844). Aswan Univ J Environ Stud. 2021;2(4):249–258.

93.   Moyo NAG, Rapatsa-Malatji MM. A review and meta-analysis of selected plant protein

sources as a replacement of fishmeal in the diet of tilapias. Ann Anim Sci. 2023;23(3):681–690.

94.   Xia Z, Zhu M, Zhang Y. Effects of the probiotic *Arthrobacter sp.* CW9 on the survival

and immune status of white shrimp (*Penaeus vannamei*). Lett Appl Microbiol. 2014;58(1):60–64.

95.   Wuertz S, Beça F, Kreuz E, Wanka KM, Azeredo R, Machado M, et al. Two Probiotic

Candidates of the Genus *Psychrobacter* Modulate the Immune Response and Disease Resistance

after Experimental Infection in Turbot (*Scophthalmus maximus*, Linnaeus 1758). Fishes.

2023;8(3):12–14.

96.   Moroni F, Naya-Català F, Piazzon MC, Rimoldi S, Calduch-Giner J, Giardini A, et al.

The Effects of Nisin-Producing *Lactococcus lactis* Strain Used as Probiotic on Gilthead Sea Bream

(*Sparus aurata*) Growth, Gut Microbiota, and Transcriptional Response. Front Mar Sci.

2021;8(April):1-18.

97.   Vasyliuk OM, Skrotskyi SO, Khomenko LA, Babich T V. Probiotics Based on Lactic

Acid Bacteria for Aquaculture. Mikrobiol Zh. 2023;85(2):75–92.

98.   Hu Y, Zhang J, Xue J, Chu W, Hu Y. Effects of dietary soy isoflavone and soy saponin

on growth performance, intestinal structure, intestinal immunity and gut microbiota community on

rice field eel (*Monopterus albus*). Aquaculture. 2021;537:1-11..

99.   Zhou C, Lin H, Ge X, Niu J, Wang J, Wang Y, et al. The Effects of dietary soybean

isoflavones on growth, innate immune responses, hepatic antioxidant abilities and disease

resistance of juvenile golden pompano *Trachinotus ovatus*. Fish Shellfish Immunol [Internet].

2015;43(1):158–166.

100.   Chojnacka K. Biologically Active Compounds in Seaweed Extracts - the Prospects for

the Application. Open Conf Proc J. 2012;3(1):20–28.

101.   Hentati F, Tounsi L, Djomdi D, Pierre G, Delattre C, Ursu AV, et al. Bioactive

polysaccharides from seaweeds. Molecules. 2020;25(14):1–29.

102.   Kumar LRG, Paul PT, Anas KK, Tejpal CS, Chatterjee NS, Anupama TK, et al.

Phlorotannins–bioactivity and extraction perspectives. J Appl Phycol. 2022;34(4):2173–2185.

103.   Ramnani P, Chitarrari R, Tuohy K, Grant J, Hotchkiss S, Philp K, et al. Invitro

fermentation and prebiotic potential of novel low molecular weight polysaccharides derived from

agar and alginate seaweeds. Anaerobe. 2012;18(1):1–6.

104.   Alvanou M V, Feidantsis K, Staikou A, Apostolidis AP, Michaelidis B, Giantsis IA.

Probiotics , Prebiotics , and Synbiotics Utilization in Crayfish Aquaculture and Factors Affecting

Gut Microbiota. Microorganisms. 2023;11(5):1–30.

105.   Eom SH, Kim YM, Kim SK. Antimicrobial effect of phlorotannins from marine brown

algae. Food Chem Toxicol. 2012;50(9):3251–3255.

106.   Ford L, Stratakos AC, Theodoridou K, Dick JTA, Sheldrake GN, Linton M, et al.

Polyphenols from Brown Seaweeds as a Potential Antimicrobial Agent in Animal Feeds. ACS

Omega. 2020;5(16):9093-9103.

107.   Manso T, Lores M, de Miguel T. Antimicrobial Activity of Polyphenols and Natural

Polyphenolic Extracts on Clinical Isolates. Antibiotics. 2022;11(1):1–18.

108.   Pradhan B, Nayak R, Bhuyan PP, Patra S, Behera C, Sahoo S, et al. Algal Phlorotannins

as Novel Antibacterial Agents with Reference to the Antioxidant Modulation: Current Advances

and Future Directions. Mar Drugs. 2022;20(6):1-15

109.   Zhang XY, Zeng XN, Liu S, Wu F, Li YY, Pan Q. Effects of dietary four different

woody forages on gut microbiota of Nile tilapia (*Oreochromis niloticus*). Aquac Res.

2021;52(4):1733–1742.

110. Wu F, Chen B, Liu S, Xia X, Gao L, Zhang X, et al. Effects of woody forages on biodiversity and bioactivity of aerobic culturable gut bacteria of tilapia (*Oreochromis niloticus*). PLoS One. 2020;15(7):1–17.

Support information

S1 Table. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 Checklist

S2 Table. Complete database search strategies and the number of articles accessed in each database

S3 Table. Data extraction from eligible studies on the effect of soybean and seaweed-based diets on tilapia production

S1 Fig. Publication bias represented by funnel plot of individual studies about the effect of soybean on SGR (A) and FCR (B)

S2 Fig. Publication bias represented by funnel plot of individual studies about the effect of seaweed on SGR (A) and FCR (B)

Author Contributions

Conceptualization: Leonildo dos Anjo Viagem

Data curation: Leonildo dos Anjo Viagem and Jean Nepomuscene Hakizimana

Formal analysis: Leonildo dos Anjo Viagem

Methodology: Leonildo dos Anjo Viagem

Supervision: Cyrus Rumisha, Brunno da Silva Cerozi and Gerald Misinzo

Writing - original draft: Leonildo dos Anjo Viagem

Writing - review & editing: Jean Nepomuscene Hakizimana, Cyrus Rumisha, Brunno da Silva Cerozi and Gerald Misinzo

Competing Interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Data Availability

All data supporting the findings of this study are available within the paper and its supplementary information.

A

Study or Subgroup	Treated (Soybean)			Control (Fish meal)			Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Abdel-Warith et al., (2013)	2.31	0.04	120	2.62	0.08	40	7.4%	-5.85 [-6.59, -5.11]
Agbo et al., (2015)	2.83	0.12	120	3.72	0.12	60	7.3%	-7.39 [-8.22, -6.55]
Ahmad et al., (2020)	2.57	0.48	270	2.21	0.12	90	7.9%	0.85 [0.61, 1.10]
Azaza et al., (2008)	5.41	0.13	270	5.22	0.09	90	7.9%	1.56 [1.30, 1.83]
El-Saidy et al., (2010)	1.42	0.02	180	1.4	0.01	45	7.8%	1.08 [0.74, 1.42]
Koumi et al., ((2011)* (O.n)	1.45	0.7	240	1.6	0.1	120	7.9%	-0.26 [-0.48, -0.04]
Koumi et al., ((2011)* (S.m)	1.3	0.4	240	1.3	0.2	120	7.9%	0.00 [-0.22, 0.22]
Koumi et al., (2009)	1.4	0.1	150	1.6	0.1	150	7.9%	-1.99 [-2.27, -1.72]
Lin and Luo 2011	2.77	0.05	360	2.96	0.05	90	7.8%	-3.79 [-4.13, -3.45]
Obirikorang et al., (2019)	2.35	0.27	2025	2.7	0.53	675	7.9%	-0.99 [-1.08, -0.90]
Sharda et al., (2017)	0.75	0.22	120	3.22	0.22	30	6.4%	-11.17 [-12.51, -9.83]
Yigit et al., (2016)	14.61	0.5	375	14.92	0.5	75	7.9%	-0.62 [-0.87, -0.37]
Zhao et al., (2010)	6.14	0.06	1500	6.27	0.12	500	7.9%	-1.64 [-1.75, -1.52]
Total (95% CI)			**5970**			**2085**	**100.0%**	**-2.14 [-2.92, -1.37]**

Heterogeneity: Tau² = 1.95; Chi² = 1890.82, df = 12 (P < 0.00001); I² = 99%
Test for overall effect: Z = 5.44 (P < 0.00001)

B

Study or Subgroup	Treated (Soybean)			Control (Fish meal)			Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total		
Abdel-Warith et al., (2013)	1.23	0.02	120	1.09	0.02	40	7.4%	6.97 [6.12, 7.82]
Agbo et al., (2015)	3.22	0.48	120	2.07	0.26	60	7.7%	2.73 [2.31, 3.15]
Ahmad et al., (2020)	1.86	0.3	270	2.04	0.04	90	7.7%	-0.69 [-0.93, -0.44]
Azaza et al., (2008)	1.51	0.08	270	1.61	0.15	90	7.7%	-0.98 [-1.23, -0.73]
El-Saidy et al., (2010)	1.78	0.06	180	1.83	0.08	45	7.7%	-0.77 [-1.11, -0.44]
Koumi et al., ((2011)* (O.n)	1.2	0.2	240	1.1	0.2	120	7.7%	0.17 [-0.05, 0.39]
Koumi et al., ((2011)* (S.m)	2.65	0.3	240	2.4	0.2	120	7.7%	0.53 [0.31, 0.75]
Koumi et al., (2009)	1.2	0.1	150	1.1	0.2	150	7.7%	0.63 [0.40, 0.86]
Lin and Luo (2011)	1.9	0.05	360	1.69	0.05	90	7.7%	4.19 [3.83, 4.55]
Obirikorang et al., (2019)	1.68	0.11	2025	1.65	0.32	675	7.8%	0.16 [0.07, 0.25]
Sharda et al., (2017)	6.18	1.06	120	0.93	0.27	30	7.5%	5.45 [4.71, 6.19]
Yigit et al., (2016)	1.83	0.08	375	1.8	0.04	75	7.7%	0.37 [0.13, 0.62]
Zhao et al., (2010)	1	0.01	1500	0.95	0.01	500	7.8%	5.00 [4.81, 5.18]
Total (95% CI)			**5970**			**2085**	**100.0%**	**1.80 [0.72, 2.89]**

Heterogeneity: Tau² = 3.94; Chi² = 3430.64, df = 12 (P < 0.00001); I² = 100%
Test for overall effect: Z = 3.26 (P = 0.001)

A

Study or Subgroup	Treated (seaweed) Mean	SD	Total	Control (Fish meal) Mean	SD	Total	Weight	Std. Mean Difference IV, Random, 95% CI	Std. Mean Difference IV, Random, 95% CI
Ashour et al., (2020)	2.33	0.15	1440	2.2	0.001	360	6.0%	0.97 [0.85, 1.09]	
Eissa et al., (2021) (CY)**	2.58	0.09	180	2.63	0.09	60	6.0%	-0.55 [-0.85, -0.26]	
Eissa et al., (2021) (LA)**	2.6	0.07	180	2.63	0.09	60	6.0%	-0.40 [-0.69, -0.10]	
Eissa et al., (2021) (UV)**	2.47	0.09	180	2.63	0.09	60	6.0%	-1.77 [-2.11, -1.44]	
El-Sayed Saleh (2020)	2.09	0.18	100	2.97	0.06	50	5.8%	-5.79 [-6.54, -5.04]	
Hussein (2017)	2.72	0.3	135	2.76	0.1	45	6.0%	-0.19 [-0.53, 0.14]	
Khalafalla and El-Hais** (PT) (2015)	1.29	0.3	60	1.12	0.3	30	5.9%	0.56 [0.12, 1.01]	
Khalafalla and El-Hais** (UV) (2015)	1.36	0.3	60	1.12	0.3	30	5.9%	0.79 [0.34, 1.25]	
Marinho et al., (2013)	2.56	0.03	270	2.87	0.05	90	5.8%	-8.59 [-9.26, -7.91]	
Radwan et al., (2022)	1.87	0.8	900	1.39	0.03	180	6.0%	1.38 [1.21, 1.55]	
Saleh et al., (2014)	3.21	0.1	360	3.18	0.14	90	6.0%	2.83 [2.53, 3.12]	
Salem et al., (2021)	2.14	0.03	90	1.88	0.01	30	5.4%	9.74 [8.42, 11.06]	
Silva et al., (2014) (GRA)**	0.91	0.04	75	1.27	0.06	75	6.0%	-7.02 [-7.89, -6.16]	
Silva et al., (2014) (POR)**	1.23	0.06	75	1.27	0.06	75	6.0%	-0.66 [-0.99, -0.33]	
Silva et al., (2014) (UV)**	1.22	0.06	75	1.27	0.06	75	6.0%	-0.83 [-1.16, -0.50]	
Stadtlander et al., (2012)	2.18	0.09	10	2.09	0.14	5	5.6%	0.79 [-0.34, 1.91]	
Younis et al., (2017)	1.07	0.07	144	1.41	0.16	48	6.0%	-3.38 [-3.86, -2.91]	
Total (95% CI)			**4334**			**1363**	**100.0%**	**-0.74 [-1.70, 0.22]**	

Heterogeneity: Tau² = 3.99; Chi² = 2474.40, df = 16 (P < 0.00001); I² = 99%
Test for overall effect: Z = 1.52 (P = 0.13)

B

Study or Subgroup	Treated (seaweed) Mean	SD	Total	Fish meal Mean	SD	Total	Weight	Std. Mean Difference IV, Random, 95% CI	Std. Mean Difference IV, Random, 95% CI
Ashour et al., (2020)	1.42	0.03	1440	1.51	0.01	360	6.0%	-3.31 [-3.46, -3.15]	
Eissa et al., (2021) (CY)**	1.7	0.03	180	1.69	0.03	60	6.0%	0.33 [0.04, 0.63]	
Eissa et al., (2021) (LA)**	1.71	0.03	180	1.69	0.03	60	6.0%	0.66 [0.37, 0.96]	
Eissa et al., (2021) (UV)**	1.76	0.04	180	1.69	0.03	60	6.0%	1.85 [1.51, 2.18]	
El-Sayed Saleh (2020)	2.2	0.12	100	1.29	0.04	50	5.7%	8.98 [7.90, 10.07]	
Hussein (2017)	1.21	0.17	135	1.3	0.1	45	5.9%	-0.58 [-0.92, -0.23]	
Khalafalla and El-Hais** (PT) (2015)	1.46	0.31	60	1.74	0.31	30	5.9%	-0.90 [-1.35, -0.44]	
Khalafalla and El-Hais** (UV) (2015)	1.31	0.31	60	1.74	0.31	30	5.9%	-1.38 [-1.86, -0.89]	
Marinho et al., (2013)	2.56	0.03	270	2.87	0.05	90	5.9%	-8.59 [-9.26, -7.91]	
Radwan et al., (2022)	1.31	0.32	900	1.76	0.12	180	6.0%	-1.52 [-1.69, -1.35]	
Saleh et al., (2014)	1.1	0.05	360	1.4	0.02	90	5.9%	-6.56 [-7.05, -6.07]	
Salem et al., (2021)	1.53	0.04	90	1.93	0.01	30	5.5%	-11.33 [-12.84, -9.81]	
Silva et al., (2014) (GRA)**	2.03	0.09	75	1.41	0.04	75	5.7%	8.86 [7.79, 9.92]	
Silva et al., (2014) (POR)**	1.44	0.08	75	1.41	0.04	75	5.9%	0.47 [0.15, 0.80]	
Silva et al., (2014) (UV)**	1.37	0.15	75	1.41	0.04	75	5.9%	-0.36 [-0.69, -0.04]	
Stadtlander et al., (2012)	1.55	0.08	10	1.67	0.11	5	5.7%	-1.25 [-2.44, -0.06]	
Younis et al., (2017)	2.95	0.51	144	1.78	0.25	48	5.9%	2.54 [2.12, 2.95]	
Total (95% CI)			**4334**			**1363**	**100.0%**	**-0.70 [-1.94, 0.54]**	

Heterogeneity: Tau² = 6.70; Chi² = 3494.14, df = 16 (P < 0.00001); I² = 100%
Test for overall effect: Z = 1.10 (P = 0.27)