CellTracksColab—A platform for compiling, analyzing, and exploring tracking data

Guillaume Jacquemet[1,2,3,4] ✉

[1]Turku Bioscience Centre, University of Turku and Åbo Akademi University, FI-20520 Turku, FI
[2]Faculty of Science and Engineering, Cell Biology, Åbo Akademi University, FI-20520 Turku, FI
[3]Turku Bioimaging, University of Turku and Åbo Akademi University, FI- 20520 Turku, FI
[4]InFLAMES Research Flagship Center, Åbo Akademi University, FI- 20520 Turku, FI

In life sciences, tracking objects from movies enables researchers to quantify the behavior of single particles, organelles, bacteria, cells, and even whole animals. While numerous tools now allow automated tracking from video, a significant challenge persists in compiling, analyzing, and exploring the large datasets generated by these approaches. Here, we introduce CellTracksColab, an online platform tailored to simplify the exploration and analysis of tracking data by leveraging the free, cloud-based computational resources of Google Colab. CellTracksColab facilitates the amalgamation and analysis of results across multiple fields of view, conditions, and repeats, ensuring a holistic dataset overview. CellTracksColab also harnesses the power of high-dimensional data reduction and clustering, enabling researchers to identify distinct behavioral patterns and trends without bias. Finally, CellTracksColab also includes specialized analysis modules enabling spatial analyses (clustering, proximity to specific regions of interest). CellTracksColab is available for the broader scientific community here.

Cell Migration | Data analysis | Tracking |
Correspondence: *guillaume.jacquemet@abo.fi*

Introduction

With the evolution of imaging techniques, biological phenomena are acquired at greater speed and resolution, facilitating the recording of dynamic events. In life science, tracking has emerged as an indispensable tool for unparalleled insights into dynamic molecular and cellular behaviors. Parallel to this, segmentation methods relying on machine learning and deep learning are now greatly facilitating the implementation of complex tracking pipelines[1–4], enabling the quantitative analysis of dynamic behaviors. Yet, as the capabilities of tracking tools have expanded, so too have the challenges associated with analyzing the resulting data.

Multiple tools have been developed to help researchers compile tracking data; for instance, these include the Ibidi Chemotaxis tool (a Fiji plugin), the MotilityLab website (an online platform for CelltrackR[5], or TrackMateR[6]. These traditional analytical approaches, implemented by us and many others, typically reduce tracking datasets to population-level analyses where track metrics are averaged across different conditions. Yet, while practical, such analyses overlook the heterogeneity within biological data. Over the past two years, multiple tools, including CellPhe (an R toolbox[7]), Traject3D (a collection of Matlab scripts,[8]), and CellPlato (a

Python toolbox, [9,10]), have been designed to harness the high-dimensionality of tracking datasets to assist in the unbiased discovery of rare phenotypes. Yet, these tools often remain challenging to implement for users with no or little coding expertise.

Here, prompted by a need to streamline the analysis of our own tracking datasets, we release CellTracksColab, a novice-oriented, cloud-based platform designed to compile and explore tracking data. Drawing inspiration from ColabFold[11] and ZeroCostDL4Mic[12], CellTracksColab operates within the Google Colaboratory framework (abbreviated as Colab) and provides a simple graphical user interface (GUI) that necessitates only a web browser and a Google account for user access. All computational tasks are conducted in the cloud via Google Colab, eliminating the requirement for software installation or the establishment of complex Python or R environments. Through a simplified workflow, researchers can install essential software dependencies with a few mouse clicks, upload their tracking data, and start with their analyses. CellTracksColab extends beyond visualization and population analyses, empowering researchers to delve into the nuanced dynamics and behaviors encapsulated within their tracking experiments.

Results

The CellTracksColab framework. The CellTracksColab platform comprises a collection of Jupyter notebooks designed to streamline tracking data analysis (Fig. 1A). CellTracksColab utilizes Google Colab, providing users free access to computing resources such as RAM and disk space. This approach facilitates advanced data analysis and simplifies the user experience by eliminating the need for local installations.

CellTracksColab processes tracking data stored in CSV files, where each file corresponds to the tracking results from an individual video. These files must be organized and named according to the notebook's specifications for optimal functionality (see the documentation for details). Such organization is critical to accurately identify and categorize various aspects of the dataset, including experimental conditions, biological replicates, and fields of view. This structured approach to data organization not only streamlines the analysis process but also enables a thorough exploration and understanding of data variability and heterogeneity across the

Fig. 1. The CellTracksColab platform. (**A**) Schematic representation of the CellTracksColab workflow. (**B**) Visualization of tracks in a CellTracksColab notebook. (**C**) Statistical analysis of track metrics using CellTracksColab. This figure shows the analysis of breast cancer cell migration (expressing CTRL shRNA or MYO10-targeting shRNA) in different environments beneath a collagen gel and standard media. The directionality metric is presented in a Tukey boxplot format. Vertical whiskers extend to data points within 1.5× the interquartile range. Each biological replicate is uniquely color-coded for clarity. Accompanying the plot are mirrored heatmaps that illustrate the effect size (Cohen's d value) and statistical significance (p-values from randomization tests with Bonferroni Correction) across various conditions. (**D**) Dimensionality reduction and clustering visualization in CellTracksColab. This panel displays a 2D t-SNE projection of the entire dataset, utilizing comprehensive track metrics for the analysis. Data points are color-coded to reflect cluster groups identified through HDBSCAN analysis on the t-SNE projection, providing insights into track characteristics and similarities. (**E**) Spatial clustering analysis using Ripley's L function and Monte Carlo simulations in CellTracksColab. This graph illustrates the spatial distribution of tracks, where a blue curve above the zero line indicates clustering at a specific radius in the field of view. The Monte Carlo simulation results are included to assess the statistical significance of the observed patterns. (**F**) Measurement and analysis of object-to-region proximity using CellTracksColab. This example demonstrates the platform's utility in quantifying the distance of objects (marked as yellow dots) relative to a defined region of interest (denoted by the white edge). The tool allows tracking these distances over time and computing related metrics, facilitating in-depth spatial analysis.

dataset.

Under the constraints of the free version of Google Colab, the performance of CellTracksColab is bound by the resources available. However, optimization of the underlying code has been executed to ensure maximal efficiency in resource utilization. For instance, we analyzed more than 50,000 tracks (> 3 million objects from 117 videos) using CellTracksColab and the free version of Google Colab (all results presented in the manuscript can be replicated with the free version of Google Colab). CellTracksColab could accommodate one of our larger datasets encompassing over 536,000 tracks (> 56 million objects from 300 videos), but this required the additional RAM that Google Colab Plus provides.

Analyzing data using CellTracksColab. Once tracking data are stored in CellTracksColab dataframes, users can perform various operations such as viewing (Fig. 1B) and filtering and smoothing tracks (Fig. S1). Track smoothing can prove to be particularly beneficial before the computation of directionality metrics, especially when the tracked object exhibits jitteriness (e.g., nuclei) and the user's interest lies in discerning the overall movement of the cell.

Upon aggregating data, CellTracksColab can compute various track metrics or import them directly from prior analyses conducted in the tracking software (Fig. 1A). Users can then generate boxplots illustrating the distribution of different track metrics of interest (Fig. 1C). Additionally, several pertinent statistical metrics are calculated, such as Cohen's d value, which quantifies the effect size between groups (independently of sample size). CellTracksColab also calculates p-values to compare conditions employing a randomization test to examine the null hypothesis (assessing whether the conditions share the same distribution), both with and without Bonferroni Correction (which adjusts significance thresholds for multiple comparisons) (Fig. 1C). Importantly, PDF files of all plots and CSV files encapsulating all plot data are exported, enabling users to replot the results using their preferred software platforms conveniently.

CellTracksColab also enables users to perform quality control on their dataset, such as checking that their data is balanced between repeats and conditions. The user can resample their data when deemed too unbalanced before plotting track metrics of interest. In addition, CellTracksColab can compute similarities across different experimental conditions and replicates using various track metrics to ensure data reliability and meaningful analysis. The results are visualized using hierarchical clustering in the form of dendrograms, which aids in comparing similarities within and across different conditions and identifying outliers.

Inspired by CellPlato[10], CellTracksColab integrates Uniform Manifold Approximation and Projection (UMAP) or t-distributed Stochastic Neighbor Embedding (t-SNE) combined with Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN) to explore the inherent heterogeneity within tracking datasets unbiasedly (Fig. 1D). This combination allows for dimensionality reduction and effective clustering of tracks, revealing distinct sub-

populations or behaviors within the data. The platform provides capabilities to plot track metrics for each cluster, create heatmaps for an overview of data variation, and identify exemplar tracks for detailed analysis. Additionally, it generates a 'fingerprint' of cluster distributions across conditions, aiding in comparative studies and benchmarking.

CellTracksColab also includes specialized analysis modules enabling the spatial analysis of track data, enabling, for instance, the assessment of track clustering (Fig. 1E) or calculating the proximity of tracks to specific regions of interest (Fig. 1F). Due to the platform's flexible design, we anticipate the addition of additional analysis modules, both by our team and the user community.

Exploring T-cell migration using CellTracksColab. To showcase the capabilities of CellTracksColab, we first chose to reanalyze a small dataset of T-cells migrating on either Vascular cell adhesion protein 1 (VCAM) or Intercellular Adhesion Molecule 1 (ICAM), captured through brightfield microscopy (Fig. 2A)[1,13,14]. Automated cell tracking was achieved using StarDist and TrackMate algorithms[1]. The dataset encompasses ten fields of view (videos) spread across two conditions and three biological repeats. CellTracksColab compiled this dataset in 4 seconds, incorporating 2,297 tracks and 38,852 tracked objects.

After the computation of additional metrics, we first evaluated the dataset's balance and variability across different fields of view. Although the dataset exhibited some imbalance, we opted against resampling due to its relatively small size (Fig. S2A). Intriguingly, the FOV-based clustering analysis unveiled an unexpected alignment between two FOVs from the ICAM condition with those from the VCAM condition, hinting at potential similarities in cell tracking patterns (Fig. S2B). The condition and repeat-based clustering analysis further corroborated this observation. Specifically, the analysis revealed that the ICAM second biological repeat displayed a clustering pattern remarkably similar to those observed in VCAM repeats (Fig. S2C). This analysis indicates that one ICAM biological repeat does not behave as the other two, providing valuable information to the experimental scientist.

We further utilized CellTracksColab to plot key track metrics, including mean speed and directionality of T-cell migration. Our analysis confirmed that T-cells exhibit slower and less directional movement on VCAM than on ICAM surfaces (Fig. 2B). To delve deeper, we employed UMAP for dimensionality reduction, followed by HDBSCAN clustering. This approach revealed the presence of at least four distinct behavioral clusters within the cell population, suggesting varied migration patterns among T-cells (Fig. 2C-D). A fingerprinting plot then provided insights into the distribution of ICAM and VCAM tracks across these clusters, highlighting differing proportions (Fig. 2E). A notable observation was the higher percentage of ICAM tracks in clusters 2 and 3 compared to VCAM. CellTracksColab generates a heatmap representing the Z score of available track metrics for each cluster to facilitate rapid metric comparison across clusters (Fig. 2F). Clusters 2 and 3 are characterized

Fig. 2. Exploring T-cell migration using CellTracksColab. (**A**) T-cells plated on ICAM were recorded using a brightfield microscope and automatically tracked using StarDist and TrackMate. Detected cells (in magenta) and their tracks (colors indicate track ID) are displayed. Scale bar: 100 μm. (**B**) The 'track mean speed' and track 'directionality' metrics for each condition are summarised in Tukey boxplots. The effect size (d, Cohen's d value) and the statistical significance (p, p-values from randomization tests with Bonferroni correction) between the conditions are displayed. (**C**) 2D UMAP projection of the entire dataset, using all available track metrics for dimensionality reduction. Data points are color-coded based on VCAM and ICAM conditions. (**D**) Resultant clusters from the HDBSCAN analysis on the 2D UMAP projection. Euclidean distance served as the metric for clustering. Each identified cluster is color-coded. (**E**) Fingerprint plot showcasing the distribution percentage of track in each cluster across different conditions. (**F**) Heatmap representation, normalized using Z scores, displaying variations in selected track metrics among the clusters. Full heatmaps are available in the Zenodo archive of this dataset. (**G**) The 'track mean speed,' track 'directionality,' and 'mean (cell) circularity' metrics for each cluster are summarised in a Tukey boxplot format as in (**B**). (**H**) The 'track mean speed,' track 'directionality,' and 'mean (cell) circularity' metrics for each condition for Cluster 2 are summarised in a Tukey boxplot format as in (**B**). For all box plots, the vertical whiskers extend to data points within 1.5× the interquartile range, and the values for each track are shown as dots where each biological replicate is displayed next to each other from R1 to R3 (left to right). Plot axes are limited to 10x the interquartile range.

by faster, more directional tracks, with cells having a lower circularity (Fig. 2G). Finally, we compared track metrics between the ICAM and VCAM conditions within specific clusters. Focusing solely on tracks in cluster 2, we observed that amongst motile cells, cells plated on ICAM migrated faster and more directionally and tended to be more circular than those on VCAM (Fig. 2H). While we provide only brief examples here, we can delve deeper into the analysis, identify tracks belonging to each cluster, and match them back to the original video. Further analyses will depend on the user's interest in the biological phenomenon studied. This multifaceted analysis underscores CellTracksColab's utility in offering nuanced insights into cell migration dynamics under different conditions.

Studying cancer cell migration using CellTracksColab. Next, we analyzed a new dataset of collectively migrating cancer cells (Fig. 3A). In this dataset, breast cancer cells expressing a CTRL shRNA or a shRNA targeting the filopodial protein MYO10 were allowed to migrate either beneath a collagen gel or within standard media (data describing the migration behavior of these cells in standard media was previously reported here[15]). Automated cell tracking was achieved using StarDist and TrackMate algorithms[1]. This larger dataset encompasses 117 fields of view (videos) spread across four conditions and three biological repeats. CellTracksColab compiled this dataset in around 6 minutes, storing 50,010 tracks and 3,291,496 tracked objects.

As with the T-cell dataset, we first performed quality control steps after computing additional track metrics. In the case of the breast cancer cell dataset, our quality control revealed some challenges: First, the third biological repeat (R3) did not cluster with the others (Fig. S3A). Additionally, the dataset was unbalanced, with the third repeat contributing disproportionately more tracks (Fig. 3B). Together, this analysis could indicate an issue with this third biological repeat and signal to the experimenter that the experiments might need to be repeated a fourth time. In addition, given this imbalance, we deemed it imperative to resample the dataset to ensure that R3's data does not unduly influence the overall conclusions. To ensure the robustness of the resampling, CelltrackColab allows for performing a statistical comparison between the original and resampled data. The outcomes of this comparison are succinctly visualized in a heatmap, providing a clear and accessible way to assess the effects of resampling on the dataset's overall distribution (Fig. S3B). Post-resampling, the dataset contains 1,376 tracks for each condition and repeat (Fig. S3C).

In our analysis of the resampled dataset using CellTracksColab, we focused on key track metrics such as mean speed and directionality to elucidate patterns of collective migration. We find that shMYO10 cells migrate slower than control cells, a pattern that persists even under the influence of a collagen gel (Fig. 3C and Fig. S4A). Intriguingly, a closer examination of individual tracks revealed that cells in the third biological repeat (R3) exhibited a faster movement, yet this did not alter the overall migration pattern observed in the dataset (Fig. 3C). Without a collagen gel, we observed

no significant differences in the directionality of migration between the conditions (Fig. 3C and Fig. S4B). However, MYO10-silenced cells displayed increased directionality under the collagen gel compared to control cells, which was an unexpected finding (Fig. 3C).

Further analysis utilizing 2D UMAP projections of the whole dataset revealed the inherent challenges in cluster generation likely due to the similarity in track characteristics, a common occurrence in collective migration (Fig. 3D). Nevertheless, by employing the canberra distance, distinct clusters were successfully delineated (Fig. S4C). These clusters provided a clear 'fingerprint' for each condition, with Cluster 2 highlighting key differences between conditions (Fig. 3E). Cluster 2 is characterized by faster, more directional tracks (Fig. S4D). Within Cluster 2, MYO10-silenced cells showed increased directionality compared to CTRL cells in the presence of a collagen gel (Fig. S4E). Additionally, CTRL cells in this cluster moved faster than their MYO10-silenced counterparts (Fig. S4F).

In the study of collective migration, a critical aspect to consider is the distinct behavior of leading cells compared to those positioned further from the leading edge[16]. To address this, we used the CellTracksColab spatial analysis module to measure each track's distance to the leading edge over time (Fig. 3F). Interestingly, overall, we observed no significant differences among the conditions in the cells' capability to target the leading edge (Fig. 3G and Fig. S4E). Yet, the 'Direction Movement' metric revealed a complex scenario: while, on average, cell distance to the leading edge does not change between the beginning and the end of the track, the data distribution was broad. Many tracks were found closer to the leading edge by the end of the tracking period, while others were found to be further away (Fig. 3G).

Delving deeper, CellTracksColab allowed us to segregate the tracks based on their maximum distance to the leading edge, distinguishing between tracks proximal to and distant from the monolayer edge. This stratified analysis unveiled that, across all conditions, cells closer to the leading edge exhibited more directional movement compared to those further away (Fig. 3H). Notably, in the presence of a collagen gel, silencing MYO10 resulted in more directional movement, irrespective of the cells' proximity to the leading edge (Fig. 3H). This example highlights how CellTracksColab can help extract spatial insights from tracking data. Furthermore, the spatial metrics derived from these analyses can enrich dimensionality reduction analyses, potentially helping unveil additional nuanced behaviors in tracking data.

Studying filopodia dynamics using CellTracksColab. In our final example, we aimed to showcase the versatility of CellTracksColab by exploring a filopodia dynamics dataset[17], diverging from our previous focus on cell migration. This study involved U2OS cells expressing different MYO10 constructs, a protein-inducing filopodia formation that also accumulates at their tips[18] (Fig. 4A). We tracked MYO10 puncta in live cells to investigate the dynamics of filopodia induced by three MYO10 variants: the wild type (MYO10$^{\text{WT}}$), a mutant lacking the MyTH4/FERM

Fig. 3. Exploring cancer cell migration using CellTracksColab. (**A**) MCF10DCIS.com lifeact-RFP cells, labeled with SiR-DNA, were recorded live using a spinning disk confocal microscope and tracked using StarDist and TrackMate. Detected nuclei and tracks (colors indicate track ID) are displayed. Scale bar: 100 μm. (**B**) This panel presents a stacked histogram showcasing the number of tracks for each biological repeat under different conditions. Each biological repeat is color-coded, and each histogram segment's specific number of tracks is annotated. (**C**) The 'track mean speed' and 'directionality' metrics for each condition (resampled dataset) are summarised in a Tukey boxplot format. The p-value and Cohen's d-value heatmaps are shown in Fig. S4A and S4B. (**D**) 2D UMAP projection of the entire dataset, using all available track metrics for dimensionality reduction. Data points are color-coded based on the conditions. (**E**) A fingerprint plot showcasing the distribution percentage of each cluster across different conditions (clustering is shown in Fig. S4C). (**F**) Distance measurement of 10 selected tracks to the monolayer leading edge. The image on the left is the raw microscopy image. The image on the right was generated by CellTracksColab to visually validate that the distance measurements are correct. The yellow dots indicate the randomly selected tracks, and the red circles indicate the measured distance. The leading edge is displayed in white. Scale bar: 100 μm. (**G**) The 'Direction Movement' metric for each condition (whole dataset) is summarized in a Tukey boxplot format. This metric is calculated as EndDistance - StartDistance (distances of the track from the leading edge at the end and start of the tracking period, respectively). A positive value indicates moving away from the leading edge over time, and a negative value suggests moving closer. The p-value and Cohen's d-value heatmaps are shown in Fig. S4G. (**H**) After separating the tracks based on their maximal distance to the leading edge (close, distance < 75 μm; far, distance > 75 μm), the track 'directionality' metric for each condition (whole dataset) was summarised in a Tukey boxplot. The mirrored heatmap displaying the Cohen's d value between each condition is shown on the right. For all box plots, the vertical whiskers extend to data points within 1.5× the interquartile range, and the values for each track are shown as dots where each biological replicate is displayed next to each other from R1 to R3 (left to right). Plot axes are limited to 10x the interquartile range.

domain (MYO10$^{\Delta FERM}$), and a chimera (MYO10TH), where MYO10's FERM domain is replaced by that from TLN1[17]. The dataset encompassed three experimental conditions, each with three biological replicates, totaling 112 videos. Utilizing CellTracksColab, we efficiently compiled this dataset, which included 91,825 tracks and nearly 1.5 million tracked objects, in around 4 minutes.

We started the analysis by filtering out tracks with fewer than five spots, resulting in a refined dataset comprising 59,391 tracks and 1,392,540 objects. Utilizing UMAP coupled with clustering analysis (Fig. 4B and Fig. S5A), we identified several distinct clusters, providing a window into the intricate behaviors of filopodia (Fig. S5B). However, the fingerprint plot revealed a similar distribution of tracks across clusters within each experimental condition (Fig. 4C). Given our focus on differences between the MYO10 constructs, we did not extensively investigate individual clusters.

Quality control assessments highlighted that the biological repeats were not clustering cohesively and revealed an imbalance in the dataset across both repeats and conditions (Fig. S5C and S5D). Consequently, we resampled the data to ensure a more balanced representation before proceeding with the analysis of track metrics.

Post-resampling, we observed notable distinctions: MYO10^{WT} filopodia exhibited greater stability, evidenced by longer lifetimes and slower speeds, compared to MYO10$^{\Delta FERM}$ and MYO10TH filopodia (Fig. 4D). Interestingly, MYO10$^{\Delta FERM}$ filopodia had a larger area than MYO10^{WT}, and while MYO10TH filopodia also showed a statistically significant difference in area from MYO10^{WT}, the low Cohen's d value suggests a negligible practical difference between these conditions (Fig. 4D). This observation highlights the importance of using both Cohen's d and p-values when comparing conditions, as it provides a more nuanced understanding of the data. Importantly, we also found that MYO10TH filopodia have substantially higher spatial coverage than MYO10^{WT} and MYO10$^{\Delta FERM}$ filopodia (Fig. 4D). The spatial coverage metric quantifies the area covered by the track, suggesting that MYO10TH filopodia may possess an enhanced capability to scan their environment. This result is particularly useful as we could not identify a gain-of-function phenotype for the MYO10TH mutant until now.

Discussion

Here, we introduced the CellTracksColab platform, a comprehensive tool designed for the life sciences community, offering a user-friendly, cloud-based solution for tracking data analysis. CellTracksColab integrates several functionalities for tracking data analysis, including track visualization, population analysis, statistical assessments, and dimensionality reduction.

CellTracksColab, utilizing the Google Colaboratory framework, significantly simplifies access, overcoming common barriers such as complex software installations. Its intuitive graphical user interface effectively bridges the gap between sophisticated computational methods and researchers with limited programming skills, making it more inclusive. However, it's important to acknowledge the limitations of the Google Colaboratory environment. One of the primary constraints is the limited runtime, as sessions in Colab are typically capped, which can interrupt longer analytical processes. Additionally, the computing power (especially the runtime RAM) and speed offered by the free version of Colab may not suffice for massive datasets. There are also concerns regarding data privacy. To address this challenge, we plan to enable CellTracksColab notebooks to function seamlessly across different platforms via automatically generated Docker images[19]. This expansion will allow users to leverage their local computational resources, potentially offering increased runtime, greater computing power, and enhanced data privacy. Such improvements will ensure that CellTracksColab can be efficiently used in a broader range of research scenarios, broadening its applicability.

Existing image repositories, such as the BioImage Archive[20] and the Image Data Resource[21], demonstrate the feasibility and value of sharing microscopy data. Analyzed tracking datasets are also very valuable; they hold significant potential for reanalysis and meta-analysis and offer a more manageable alternative to storing raw video footage[22]. Yet, their widespread sharing remains limited, and publicly available datasets are relatively scarce. This scarcity is partly due to the need for a standardized format for sharing tracking results. CellTracksColab partially addresses this challenge by adopting a unified format for storing tracking data and simplifying the sharing of tracking datasets. Additionally, the platform includes a streamlined notebook designed for easy viewing and replotting previously analyzed datasets. This feature enhances data analysis's transparency and promotes reproducibility, aligning with the FAIR (Findable, Accessible, Interoperable, and Reusable) principles in scientific research.

Despite its robust capabilities, CellTracksColab has certain limitations we aim to address in future updates. Currently, the platform has limited support for analyzing tracks in 3D space. While the platform can handle 3D + time datasets for metric computation, quality control, and dimensionality reduction tasks, it falls short in certain areas. Specifically, some of its analysis modules, including track visualization and spatial analysis, are optimized only for 2D + time datasets. In addition, CellTracksColab is not currently adapted for lineage tracing studies. Researchers focusing on lineage tracing must explore alternative tools specifically developed for such analyses[1,3,22].

We also plan to enhance CellTracksColab by introducing additional analytical features. This includes the capability to examine time-series data, such as analyzing variations in fluorescent reporters within tracked objects over time, which could provide deeper insights into dynamic biological processes[24,25]. Furthermore, we intend to develop a data loader for other popular tracking tools, broadening the platform's compatibility and ease of integration with existing workflows. In conclusion, we believe that CellTracksColab represents a significant step forward in tracking data analysis

Fig. 4. Exploring filopodia dynamics using CellTracksColab. (**A**) A U2OS cell expressing MYO10[WT]-GFP was imaged live using an Airyscan confocal microscope. MYO10 puncta were then tracked using StarDist and TrackMate. Detected MYO10 puncta and tracks (colors indicate track ID) are displayed. Scale bar: 25 μm. (**B**) 2D UMAP projection of the entire dataset, using all available track metrics for dimensionality reduction. Data points are color-coded based on the conditions. (**C**) A fingerprint plot showcasing the distribution percentage of each cluster across different conditions (clustering is shown in Fig. S5A). (**D**) The 'track mean speed', the 'track duration, the spot 'mean area', and the track 'spatial coverage' for each condition are summarised in Tukey boxplots. The effect size (d, Cohen's d value) and the statistical significance (p, p-values from randomization tests with Bonferroni correction) between the MYO10WT and the indicated conditions are displayed. For all box plots, the vertical whiskers extend to data points within 1.5× the interquartile range, and the values for each track are shown as dots where each biological replicate is displayed next to each other from R1 to R3 (left to right). Plot axes are limited to 10x the interquartile range.

in life sciences. Its user-friendly design and robust analytical capabilities allow researchers to explore and understand the complexities of biological motion and behavior. As we continue to develop and enhance CellTracksColab, we anticipate it becoming a useful tool in the life sciences toolkit, aiding in discovering and understanding new biological insights.

Materials and Methods

Implementation. CellTracksColab is implemented as a series of interactive Jupyter notebooks hosted on Google Colaboratory (Colab). The platform utilizes Python as its primary programming language, leveraging various libraries such as Pandas for data manipulation, Matplotlib, Seaborn for data visualization, and UMAP and HDBSCAN for dimensionality reduction and clustering analyses. The notebook's architecture facilitates ease of use while providing robust data analysis capabilities. The notebooks are structured to guide users through each step of the analysis process, from data import and preprocessing to advanced statistical and spatial analyses. Each notebook contains detailed instructions and documentation to assist users in customizing the analysis to their specific datasets. CellTracksColab requires a Google account to access Google Colab's cloud-based computational

resources. Users can upload their tracking data in CSV format, and the platform is equipped to handle both 2D and 3D tracking data, with certain limitations in 3D analysis, as currently noted. GPT-4 (OpenAI) was used to accelerate the development of CellTracksColab. All code was carefully tested and validated by the author.

Data and software availability. Multiple test datasets are available on Zenodo. They include the two test datasets that can be directly downloaded from within the CellTracksColab notebooks[26,27]. In addition, the three datasets showcased in this study, their tracking files, and the CellTracksColab dataframes are also available on Zenodo[28–30].

The code for CellTracksColab is publicly available under the MIT license, encouraging broad utilization and adaptation. CellTracksColab's GitHub repository serves as a dynamic platform for tracking the evolution of the code across various versions. Users are encouraged to report issues and suggest features directly through the GitHub interface. A stable version of the code and associated documentation is also archived on Zenodo[31].

The T cell dataset. The T cell dataset used is available on Zenodo[28] and has been detailed in previous

publications[1,13,14]. In summary, Lab-Tek 8 chamber slides (ThermoFisher) were prepared by overnight coating with either 2 μg/mL ICAM-1 or VCAM-1 at a temperature of 4°C. Subsequently, activated primary mouse CD4+ T cells were cleansed and suspended in L-15 media, enriched with 2 mg/mL D-glucose. These T cells were then placed into the chamber slides and incubated for 20 minutes. Post-incubation, a gentle wash was performed to eliminate all unattached cells. The imaging process was conducted using a 10x phase contrast objective at 37°C, utilizing a Zeiss Axiovert 200M microscope equipped with an automated X-Y stage and a Roper EMCCD camera. Time-lapse imaging was executed at intervals of 1 minute over 10 minutes, employing SlideBook 6 software from Intelligent Imaging Innovations.

Cells were automatically tracked using StarDist, directly implemented within TrackMate[1,32,33]. The StarDist model was trained using ZeroCostDL4Mic[12] and is publicly available on Zenodo[34]. This model generated excellent segmentation results on our test dataset (F1 score > 0.99). In TrackMate, the StarDist detector custom model (score threshold = 0.41 and overlap threshold = 0.5) and the Simple LAP tracker (linking max distance = 30 μm; gap closing max distance = 15 μm, gap closing max frame gap = 2 frames) were used.

In CellTracksColab, we conducted a dimensionality reduction analysis employing Uniform Manifold Approximation and Projection (UMAP). The UMAP settings were as follows: number of neighbors (n_neighbors) set to 20, minimum distance (min_dist) to 0, and number of dimensions (n_dimension) to 2. This analysis utilized an array of track metrics, including:

NUMBER_SPOTS, NUMBER_GAPS, NUMBER_SPLITS, NUMBER_MERGES, NUMBER_COMPLEX, LONGEST_GAP, TRACK_DURATION, TRACK_DISPLACEMENT, TRACK_MEAN_SPEED, TRACK_MAX_SPEED, TRACK_MIN_SPEED, TRACK_MEDIAN_SPEED, TRACK_STD_SPEED, TRACK_MEAN_QUALITY, TOTAL_DISTANCE_TRAVELED, MAX_DISTANCE_TRAVELED, CONFINEMENT_RATIO, MEAN_STRAIGHT_LINE_SPEED, LINEARITY_OF_FORWARD_PROGRESSION, MEAN_DIRECTIONAL_CHANGE_RATE, Directionality, Tortuosity, Total_Turning_Angle, Spatial_Coverage, MEAN_CIRCULARITY, MEAN_SOLIDITY, MEAN_SHAPE_INDEX, MEDIAN_CIRCULARITY, MEDIAN_SOLIDITY, MEDIAN_SHAPE_INDEX, STD_CIRCULARITY, STD_SOLIDITY, STD_SHAPE_INDEX, MIN_CIRCULARITY, MIN_SOLIDITY, MIN_SHAPE_INDEX, MAX_CIRCULARITY, MAX_SOLIDITY, MAX_SHAPE_INDEX

Subsequently, clustering analysis was performed using Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN). The parameters included clustering_data_source set to UMAP, min_samples at 20, min_cluster_size at 200, and the metric employed was Euclidean.

The breast cancer cell dataset. The tracked breast cancer cell dataset is available on Zenodo[30]. In this experiment, approximately 50,000 shCTRL or shMYO10 lifeact-RFP DCIS.COM cells[15] were seeded into one well of an ibidi culture-insert 2 well pre-placed in a μ-Slide 8 well. The cells were cultured for 24 hours, after which the culture insert was removed to create a wound-healing assay setup. When appropriate, a fibrillar collagen gel (PureCol EZ Gel) was applied over the cells and allowed to polymerize for 30 minutes at 37°C. Standard culture media was added to all wells, and the cells were left to migrate/invade for two days[35]. Before live cell imaging, the cells were treated with 0.5 μM SiR-DNA (SiR-Hoechst, Tetu-bio) for two hours. Imaging was performed over 14 hours using a Marianas spinning-disk confocal microscope system. This system included a Yokogawa CSU-W1 scanning unit mounted on an inverted Zeiss Axio Observer Z1 microscope (Intelligent Imaging Innovations, Inc.). Imaging was conducted using a 20x (NA 0.8) air Plan Apochromat objective (Zeiss), and images were captured at 10-minute intervals.

Cell tracking was conducted using Fiji[32] and TrackMate[1]. The Stardist detector was employed to detect nuclei using the Stardist versatile model[33]. Tracks were created using the Kalman tracker (a maximum frame gap of 1, a Kalman search radius of 20 μm, and a linking maximum distance of 15 μm). Post-tracking, tracks were filtered so that each track had to contain more than six spots, ensuring a significant amount of data per track, and the total distance traveled by cells had to be greater than 89 μm.

In CellTracksColab, we conducted a dimensionality reduction analysis employing Uniform Manifold Approximation and Projection (UMAP). The UMAP settings were as follows: number of neighbors (n_neighbors) set to 10, minimum distance (min_dist) to 0.5, and number of dimensions (n_dimension) to 2. This analysis utilized an array of track metrics, including:

NUMBER_SPOTS, NUMBER_GAPS, NUMBER_SPLITS, NUMBER_MERGES, NUMBER_COMPLEX, LONGEST_GAP, TRACK_DURATION, TRACK_DISPLACEMENT, TRACK_MEAN_SPEED, TRACK_MAX_SPEED, TRACK_MIN_SPEED, TRACK_MEDIAN_SPEED, TRACK_STD_SPEED, TRACK_MEAN_QUALITY, TOTAL_DISTANCE_TRAVELED, MAX_DISTANCE_TRAVELED, CONFINEMENT_RATIO, MEAN_STRAIGHT_LINE_SPEED, LINEARITY_OF_FORWARD_PROGRESSION, MEAN_DIRECTIONAL_CHANGE_RATE, Directionality, Tortuosity, Total_Turning_Angle, Spatial_Coverage, MEAN_CIRCULARITY, MEAN_SOLIDITY, MEAN_SHAPE_INDEX, MEDIAN_CIRCULARITY, MEDIAN_SOLIDITY, MEDIAN_SHAPE_INDEX, STD_CIRCULARITY, STD_SOLIDITY, STD_SHAPE_INDEX, MIN_CIRCULARITY, MIN_SOLIDITY, MIN_SHAPE_INDEX, MAX_CIRCULARITY,

MAX_SOLIDITY, MAX_SHAPE_INDEX, MaxDistance_edge, MinDistance_edge, StartDistance_edge, EndDistance_edge, MedianDistance_edge, StdDevDistance_edge, DirectionMovement_edge, AvgRateChange_edge, PercentageChange_edge, TrendSlope_edge

Subsequently, clustering analysis was performed using Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN). The parameters included clustering_data_source set to UMAP, min_samples at 20, min_cluster_size at 200, and the metric employed was Canberra.

The filopodia dataset. The tracked filopodia dataset is available on Zenodo[29] and was previously described[17]. U2-OS cells expressing MYO10-GFP, a MYO10 MyTH/FERM deletion construct (EGFP-MYO10$^{\Delta FERM}$), or an MYO10/TLN1 chimera construct (EGFP-MYO10TH) were plated for at least 2 h on fibronectin before the start of live imaging. Images were taken every 5 s at 37 C on an Airyscan microscope, using a 40x objective. MYO10 puncta were tracked using TrackMate using the custom Stardist detector and the simple LAP tracker (Linking max distance=1 micron, Gap-closing max distance=2 microns, Gap-closing max frame gap=2 micron). The StarDist 2D model used was previously described[36]. Briefly, this model was trained for 200 epochs on 11 paired image patches [image dimensions: (512, 512), patch size: (512,512)] with a batch size of 2 and a mean absolute error (MAE) loss function, using the StarDist 2D ZeroCostDL4Mic notebook[12].

In CellTracksColab, we conducted a dimensionality reduction analysis employing Uniform Manifold Approximation and Projection (UMAP). The UMAP settings were as follows: number of neighbors (n_neighbors) set to 10, minimum distance (min_dist) to 0, and number of dimensions (n_dimension) to 2. This analysis utilized an array of track metrics, including:

NUMBER_SPOTS, NUMBER_GAPS, NUMBER_SPLITS, NUMBER_MERGES, NUMBER_COMPLEX, LONGEST_GAP, TRACK_DURATION, TRACK_DISPLACEMENT, TRACK_MEAN_SPEED, TRACK_MAX_SPEED, TRACK_MIN_SPEED, TRACK_MEDIAN_SPEED, TRACK_STD_SPEED, TRACK_MEAN_QUALITY, TOTAL_DISTANCE_TRAVELED, MAX_DISTANCE_TRAVELED, CONFINEMENT_RATIO, MEAN_STRAIGHT_LINE_SPEED, LINEARITY_OF_FORWARD_PROGRESSION, MEAN_DIRECTIONAL_CHANGE_RATE, Directionality, Tortuosity, Total_Turning_Angle, Spatial_Coverage, MEAN_MEAN_INTENSITY_CH1, MEAN_MEDIAN_INTENSITY_CH1, MEAN_MIN_INTENSITY_CH1, MEAN_MAX_INTENSITY_CH1, MEAN_TOTAL_INTENSITY_CH1, MEAN_STD_INTENSITY_CH1, MEAN_CONTRAST_CH1, MEAN_SNR_CH1, MEAN_ELLIPSE_X0, MEAN_ELLIPSE_Y0, MEAN_ELLIPSE_MAJOR, MEAN_ELLIPSE_MINOR, MEAN_ELLIPSE_THETA, MEAN_ELLIPSE_ASPECTRATIO, MEAN_AREA, MEAN_PERIMETER, MEAN_CIRCULARITY, MEAN_SOLIDITY, MEAN_SHAPE_INDEX, MEDIAN_MEAN_INTENSITY_CH1, MEDIAN_MEDIAN_INTENSITY_CH1, MEDIAN_MIN_INTENSITY_CH1, MEDIAN_MAX_INTENSITY_CH1, MEDIAN_TOTAL_INTENSITY_CH1, MEDIAN_STD_INTENSITY_CH1, MEDIAN_CONTRAST_CH1, MEDIAN_SNR_CH1, MEDIAN_ELLIPSE_X0, MEDIAN_ELLIPSE_Y0, MEDIAN_ELLIPSE_MAJOR, MEDIAN_ELLIPSE_MINOR, MEDIAN_ELLIPSE_THETA, MEDIAN_ELLIPSE_ASPECTRATIO, MEDIAN_AREA, MEDIAN_PERIMETER, MEDIAN_CIRCULARITY, MEDIAN_SOLIDITY, MEDIAN_SHAPE_INDEX, STD_MEAN_INTENSITY_CH1, STD_MEDIAN_INTENSITY_CH1, STD_MIN_INTENSITY_CH1, STD_MAX_INTENSITY_CH1, STD_TOTAL_INTENSITY_CH1, STD_STD_INTENSITY_CH1, STD_CONTRAST_CH1, STD_SNR_CH1, STD_ELLIPSE_X0, STD_ELLIPSE_Y0, STD_ELLIPSE_MAJOR, STD_ELLIPSE_MINOR, STD_ELLIPSE_THETA, STD_ELLIPSE_ASPECTRATIO, STD_AREA, STD_PERIMETER, STD_CIRCULARITY, STD_SOLIDITY, STD_SHAPE_INDEX, MIN_MEAN_INTENSITY_CH1, MIN_MEDIAN_INTENSITY_CH1, MIN_MIN_INTENSITY_CH1, MIN_MAX_INTENSITY_CH1, MIN_TOTAL_INTENSITY_CH1, MIN_STD_INTENSITY_CH1, MIN_CONTRAST_CH1, MIN_SNR_CH1, MIN_ELLIPSE_X0, MIN_ELLIPSE_Y0, MIN_ELLIPSE_MAJOR, MIN_ELLIPSE_MINOR, MIN_ELLIPSE_THETA, MIN_ELLIPSE_ASPECTRATIO, MIN_AREA, MIN_PERIMETER, MIN_CIRCULARITY, MIN_SOLIDITY, MIN_SHAPE_INDEX, MAX_MEAN_INTENSITY_CH1, MAX_MEDIAN_INTENSITY_CH1, MAX_MIN_INTENSITY_CH1, MAX_MAX_INTENSITY_CH1, MAX_TOTAL_INTENSITY_CH1, MAX_STD_INTENSITY_CH1, MAX_CONTRAST_CH1, MAX_SNR_CH1, MAX_ELLIPSE_X0, MAX_ELLIPSE_Y0, MAX_ELLIPSE_MAJOR, MAX_ELLIPSE_MINOR, MAX_ELLIPSE_THETA, MAX_ELLIPSE_ASPECTRATIO, MAX_AREA, MAX_PERIMETER, MAX_CIRCULARITY, MAX_SOLIDITY, MAX_SHAPE_INDEX

Subsequently, clustering analysis was performed using Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN). The parameters included clustering_data_source set to UMAP, min_samples at 20, min_cluster_size at 300, and the metric employed was Canberra.

Manuscript preparation. GPT-4 (OpenAI) and Grammarly (Grammarly, Inc.) were used as writing aids while preparing this manuscript. All text sections generated were further edited and validated by the author. No references were provided by GPT-4. All figure panels (except Fig. 2A, 3A, and 4A) and all statistical analyses were generated directly within CellTracksColab and edited in Inkscape (Inkscape Project).

Acknowledgements

CellTracksColab is inspired by several key projects in cell tracking and analysis. We acknowledge the influential contributions of Traject3d, CellPhe, CelltrackR, the Motility-Lab website, and Cellplato on Zenodo. The innovative use of UMAP and HDBSCAN for analyzing tracking data, as featured in CellTracksColab, was first introduced in CellPlato.

I thank Hellyeh Hamidi and Estibaliz Gómez-de-Mariscal for providing feedback on this manuscript. This study was supported by the Research Council of Finland (338537 to G.J.), the Sigrid Juselius Foundation (to G.J.), the Cancer Society of Finland (Syöpäjärjestöt; to G.J.), and the Solutions for Health strategic funding to Åbo Akademi University (to G.J.). This research was supported by the In-FLAMES Flagship Programme of the Academy of Finland (decision numbers: 337530, 337531, 357910, and 357911). The Cell Imaging and Cytometry Core facility (Turku Bioscience, University of Turku, Åbo Akademi University, and Biocenter Finland) and Turku Bioimaging are acknowledged for services, instrumentation, and expertise.

Bibliography

1. Ershov, D. et al. TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines. Nat. Methods 19, 829–832 (2022).

2. Aragaki, H., Ogoh, K., Kondo, Y. & Aoki, K. LIM Tracker: a software package for cell tracking and analysis with advanced interactivity. Sci. Rep. 12, 2702 (2022).

3. Ulicna, K., Vallardi, G., Charras, G. & Lowe, A. R. Automated Deep Lineage Tree Analysis Using a Bayesian Single Cell Tracking Approach. Front. Comput. Sci. 3, (2021).

4. Sugawara, K., Çevrim, Ç. & Averof, M. Tracking cell lineages in 3D by incremental deep learning. eLife 11, e69380 (2022).

5. Wortel, I. M. N., Dannenberg, K., Berry, J. C., Miller, M. J. & Textor, J. CelltrackR: an R package for fast and flexible analysis of immune cell migration data. http://biorxiv.org/lookup/doi/10.1101/670505 (2019) doi:10.1101/670505.

6. Royle, S. quantixed/TrackMateR. (2024).

7. Wiggins, L. et al. The CellPhe toolkit for cell phenotyping using time-lapse imaging and pattern recognition. Nat. Commun. 14, 1854 (2023).

8. Freckmann, E. C. et al. Traject3d allows label-free identification of distinct co-occurring phenotypes within 3D culture by live imaging. Nat. Commun. 13, 5317 (2022).

9. Michael-shannon/cellPLATO: cellPLATO. doi:10.5281/zenodo.8096717.

10. Shannon, M. J., Eisman, S. E., Lowe, A. R., Sloan, T. & Mace, E. M. cellPLATO: an unsupervised method for identifying cell behaviour in heterogeneous cell trajectory data. 2023.10.28.564355 Preprint at https://doi.org/10.1101/2023.10.28.564355 (2023).

11. Mirdita, M. et al. ColabFold: making protein folding accessible to all. Nat. Methods 19, 679–682 (2022).

12. von Chamier, L. et al. Democratising deep learning for microscopy with ZeroCostDL4Mic. Nat. Commun. 12, 2276 (2021).

13. Roy, N. H. et al. LFA-1 signals to promote actin polymerization and upstream migration in T cells. J. Cell Sci. 133, (2020).

14. Fazeli, E. et al. Automated cell tracking using StarDist and TrackMate. http://biorxiv.org/lookup/doi/10.1101/2020.09.22.306233 (2020) doi:10.1101/2020.09.22.306233.

15. Peuhu, E. et al. MYO10-filopodia support basement membranes at pre-invasive tumor boundaries. Dev. Cell 57, 2350-2364.e7 (2022).

16. Campanale, J. P. & Montell, D. J. Who's really in charge: Diverse follower cell behaviors in collective cell migration. Curr. Opin. Cell Biol. 81, 102160 (2023).

17. Miihkinen, M. et al. Myosin-X and talin modulate integrin activity at filopodia tips. Cell Rep. 36, 109716 (2021).

18. Berg, J. S. & Cheney, R. E. Myosin-X is an unconventional myosin that undergoes intrafilopodial motility. Nat. Cell Biol. 4, 246–250 (2002).

19. Hidalgo-Cenalmor, I. et al. DL4MicEverywhere: Deep learning for microscopy made flexible, shareable, and reproducible. 2023.11.19.567606 Preprint at https://doi.org/10.1101/2023.11.19.567606 (2023).

20. Hartley, M. et al. The BioImage Archive - Building a Home for Life-Sciences Microscopy Data. J. Mol. Biol. 434, 167505 (2022).

21. Williams, E. et al. Image Data Resource: a bioimage data integration and publication platform. Nat. Methods 14, 775–781 (2017).

22. Hu, J. et al. Multisite assessment of reproducibility in high-content cell migration imaging data. Mol. Syst. Biol. 19, e11490 (2023).

23. Shim, C. et al. CellTrackVis: interactive browser-based visualization for analyzing cell trajectories and lineages. BMC Bioinformatics 24, 124 (2023).

24. Regot, S., Hughey, J. J., Bajar, B. T., Carrasco, S. & Covert, M. W. High-Sensitivity Measurements of Multiple Kinase Activities in Live Single Cells. Cell 157, 1724–1734 (2014).

25. Sakaue-Sawano, A. et al. Visualizing Spatiotemporal Dynamics of Multicellular Cell-Cycle Progression. Cell 132, 487–498 (2008).

26. Jacquemet, G. T cell dataset for CellTracksColab. (2023) doi:10.5281/zenodo.8413510.

27. Jacquemet, G. T cell dataset for CellTracksColab - 2. (2023) doi:10.5281/zenodo.8420011.

28. Jacquemet, G. CellTracksColab - T cell dataset (full). (2024) doi:10.5281/zenodo.10539720.

29. Jacquemet, G. CellTracksColab - Filopodia dataset. (2024) doi:10.5281/zenodo.10539196.

30. Jacquemet, G. CellTracksColab - breast cancer cell dataset. (2024) doi:10.5281/zenodo.10539020.

31. guijacquemet. guijacquemet/CellTracksColab: v.02. (2023) doi:10.5281/zenodo.10078057.

32. Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. Nat. Methods 9, 676–682 (2012).

33. Schmidt, U., Weigert, M., Broaddus, C. & Myers, G. Cell Detection with Star-Convex Polygons. in Medical Image Computing and Computer Assisted Intervention – MICCAI 2018 (eds. Frangi, A. F., Schnabel, J. A., Davatzikos, C., Alberola-López, C. & Fichtinger, G.) vol. 11071 265–273 (Springer International Publishing, 2018).

34. Roy, N. H. & Jacquemet, G. Combining StarDist and TrackMate example 2 - T cell dataset. (2020) doi:10.5281/zenodo.4034929.

35. Jacquemet, G. et al. FiloQuant reveals increased filopodia density during breast cancer progression. J. Cell Biol. 216, 3387–3403 (2017).

36. Popović, A. et al. Myosin-X recruits lamellipodin to filopodia tips. J. Cell Sci. 136, jcs260574 (2023).

Fig. S1. Track visualization and filtering. Example of track display before (**A**) and after (**B**) smoothing and filtering. The tracks originate from a video of migrating breast cancer cells tracked from their nuclei using TrackMate.

Fig. S2. Evaluating the experimental variability in the T-cell dataset with CellTracksColab. (**A**) This panel presents a stacked histogram showcasing the number of tracks for each biological repeat under different conditions, aiding in evaluating the dataset's balance. Each biological repeat is color-coded, and each histogram segment's specific number of tracks is annotated. (**B**, **C**) Hierarchical Clustering: Utilizing the cosine similarity metric and a complete linkage method, these dendrograms reveal the hierarchical clustering within the dataset. (**B**) FOV-based Clustering Analysis: This dendrogram illustrates the clustering across the ten available Fields of View (FOVs). (**C**) Condition and Repeat-based Clustering: This dendrogram delves deeper by segregating the dataset based on conditions and biological repeats

Fig. S3. Evaluating the experimental variability in the breast cancer cell dataset with CellTracksColab. (**A**) Utilizing the cosine similarity metric and a complete linkage method, this dendrogram assesses the similarity in the dataset between conditions and biological repeats. (**B**) p-value heatmap comparing the differences between the data distribution before and after resampling for each condition and repeats (selected number of track metrics). (**C**) This panel presents a stacked histogram showcasing the number of tracks for each biological repeat under different conditions, aiding in evaluating the dataset's balance. Each biological repeat is color-coded, and each histogram segment's specific number of tracks is annotated.

Fig. S4. Exploring the breast cancer migration dataset using CellTracksColab. (**A**, **B**) Boneferri-corrected p-value and Cohen's d-value mirrored heatmaps for the 'track mean speed' (**A**) and track 'directionality' (**B**) metrics (see Fig. 3C). (**C**) 2D UMAP projection of the entire breast cancer migration dataset, using all available track metrics for dimensionality reduction. Resultant clusters from the HDBSCAN analysis on the 2D UMAP projection. The Canberra method served as the metric for clustering. Each identified cluster is color-coded. (**D**) Heatmap representation, normalized using Z scores, displaying variations in selected track metrics among the clusters. Full heatmaps are available in the Zenodo archive of this dataset. (**E**, **F**) The 'track 'directionality,' (**E**), and 'track mean speed, (**F**)' metrics for each condition for Cluster 2 are summarised in a Tukey boxplot format. For all box plots, the vertical whiskers extend to data points within 1.5× the interquartile range, and the values for each track are shown as dots where each biological replicate is displayed next to each other from R1 to R3 (left to right). Boneferri-corrected p-value and Cohen's d-value mirrored heatmaps are displayed on the right. (**G**) Boneferri-corrected p-value and Cohen's d-value mirrored heatmaps for the 'Direction Movement' metric (see Fig. 3G).

Fig. S5. Exploring filopodia dynamics using CellTracksColab. (**A**) 2D UMAP projection of the entire filopodia dataset, using all available track metrics for dimensionality reduction. Resultant clusters from the HDBSCAN analysis on the 2D UMAP projection. The Canberra method served as the metric for clustering. Each identified cluster is color-coded. (**B**) Heatmap representation, normalized using Z scores, displaying variations in selected track metrics among the clusters. Full heatmaps are available in the Zenodo archive of this dataset. (**C**) Utilizing the cosine similarity metric and a complete linkage method, this dendrogram assesses the similarity in the filopodia dataset between conditions and biological repeats. (**D**) This panel presents a stacked histogram showcasing the number of tracks for each biological repeat under different conditions. Each biological repeat is color-coded, and each histogram segment's specific number of tracks is annotated.