ZNF91 is an endogenous repressor of the molecular phenotype associated with X-linked dystonia-parkinsonism (XDP)

Jimi L. Rosenkrantz[1], Sanaz Raghib[1], J. Elias Brandorff[1], Ashni Kapadia[1], Christine A. Vaine[2], D. Cristopher Bragg[2], Grace Farmiloe[1], Frank M.J. Jacobs[1,3]

Abstract

Background: X-linked dystonia-parkinsonism (XDP) is a severe neurodegenerative disorder resulting from the insertion of an intronic SINE-Alu-VNTR (SVA) retrotransposon in the *TAF1* gene. Recent research has revealed that the pathogenic XDP-SVA insertion leads to dysregulation of *TAF1* transcription, including increased intron retention and decreased expression of exons surrounding the insertion. The Krüppel-associated box (KRAB) zinc finger protein, ZNF91, is a critical repressor of SVA retrotransposons. However, it remains unclear whether ZNF91 is able to repress the XDP-SVA insertion and how this influences the XDP-associated molecular phenotype. In this study, we investigate the role of ZNF91 in repressing the XDP-SVA insertion and its impact on the molecular phenotype associated with XDP.

Methods: Here, we used CRISPR/Cas9 to genetically delete ZNF91 in induced pluripotent stem cell (iPSC) lines derived from XDP patients, as well as isogenic control iPSC lines that lack the XDP-SVA insertion. Total RNA sequencing and capture RNA-sequencing were used to confirm ZNF91 deletion and to assess *TAF1* transcriptional changes between conditions. Furthermore, publicly available transcriptomic data from whole blood and different brain regions were used to assess ZNF91 expression levels across ages.

Results: We found that genetic deletion of ZNF91 exacerbates the molecular phenotype

associated with the XDP-SVA insertion in patient cells, while no difference was observed

when ZNF91 was deleted from isogenic control cells. Additionally, we observed a significant

age-related reduction in ZNF91 expression in whole blood and brain, indicating a potential

role of ZNF91 in the age-dependent onset of XDP.

Conclusions: These findings indicate that ZNF91 plays a crucial role in controlling the

molecular phenotype associated with XDP. Since ZNF91 is a critical epigenetic repressor of

SVAs, this suggests that epigenetic silencing of the XDP-SVA minimizes the severity of the

molecular phenotype. Our results showing that ZNF91 expression levels significantly

decrease with age provide a potential explanation for the age-related progressive

neurodegenerative character of XDP. Collectively, our study provides important insights into

the protective role of ZNF91 in XDP pathogenesis and suggests that modulating ZNF91

levels or targeted repression of the XDP-SVA could be novel therapeutic strategies worth

exploring.

Author affiliations:

1 Evolutionary Neurogenomics, Swammerdam Institute for Life Sciences, University of

Amsterdam, 1098 XH Amsterdam, The Netherlands.

2 The Collaborative Center for X-linked Dystonia-Parkinsonism, Massachusetts General

Hospital, MA 02129, USA

3 Amsterdam Neuroscience, Complex Trait Genetics, University of Amsterdam, 1098 XH

Amsterdam, The Netherlands

Correspondence to: Frank M. J. Jacobs

Science Park 904, 1098 XH Amsterdam, The Netherlands.

f.m.j.jacobs@uva.nl

Running title: ZNF91 represses XDP molecular phenotype.

Keywords:

1. X-linked dystonia-parkinsonism (XDP);

2. SINE-Alu-VNTR (SVA);

3. Krüppel-associated zinc finger protein 91 (ZNF91);

4. TATA-binding protein-associated factor-1 gene (*TAF1*);

5. Neurodegeneration;

6. Retrotransposons;

7. CRISPR/Cas9;

8. induced pluripotent stem cells (iPSCs);

9. Capture RNA-sequencing;

Background

X-linked dystonia-parkinsonism (XDP; MIM:314250), also known as DYT3 or Lubag syndrome, is a rare, progressive neurodegenerative disorder that primarily affects adult males with maternal ancestry from the Philippine Island of Panay. In the local Filipino dialect, XDP is referred to as "lubag," meaning "twisted," due to its characteristic dystonia (abnormal muscle tone and movements) and Parkinsonism (tremors, rigidity, and difficulty with movement) [1,2]. XDP was first described in the clinic nearly 50 years ago [3]. Since then, extensive research has revealed that it is caused by a 2.6 kb SINE-Alu-VNTR (SVA) retrotransposon insertion in intron 32 of the TATA-binding protein-associated factor-1 gene (*TAF1*) on chromosome Xq13 [4,5]. *TAF1* encodes a central protein of the transcription factor II D (TFIID) complex, which plays a crucial role in the transcription of genes involved in neurodevelopmental processes [6–11]. Recent studies using XDP patient cells with and without the disease-specific SVA insertion (XDP-SVA) have shown that XDP-SVA leads to dysregulation of *TAF1* transcription, including increased intron 32 retention and decreased expression of exons surrounding the XDP-SVA insertion [5,12]. *TAF1* dysregulation is thought to underlie the development of XDP, however the exact mechanism by which the XDP-SVA insertion contributes to *TAF1* dysregulation remains poorly understood.

SVA elements are a class of retrotransposons found exclusively in primate genomes [13] and belong to one of three active transposable element (TE) families in the human genome [14–16]. SVAs are a composite of several different repeat elements and consist of a (CCCTCT)n hexamer repeat, an Alu-like element, a variable number tandem repeat (VNTR), SINE-R, and Poly-A region. There are six subfamilies of SVAs (SVA_A to SVA_F). The two youngest subfamilies of SVAs, SVA_E and SVA_F, are human-specific and include many insertions that are polymorphic among human populations [13,17,18]. Polymorphic SVAs contribute to

human genetic diversity and, in some cases, can lead to variation in traits and disease

susceptibility [19–21]. This is in part due to the nature of SVA sequences, which possess

robust gene regulatory potential and can exert a significant influence on nearby gene

expression [22–24]. The epigenetic repression of SVAs is crucial in preventing their

reinsertion into the genome but also to minimize the potential gene regulatory effects that

may arise from their insertion. Many families of TEs, including SVAs, are silenced by

Krüppel-associated box (KRAB) zinc finger (KZNF) proteins, which have been shown to

recruit repressive epigenetic modifiers [22,25,26]. Recent findings have highlighted the

critical role of the primate-specific zinc finger 91 (ZNF91) protein in repressing SVA

elements [22,24]. These previous findings strongly suggest that ZNF91 binds and represses

the pathogenic XDP-SVA insertion. However, the influence of ZNF91-mediated epigenetic

repression on the molecular phenotype of XDP remains unclear.

In this study, we aimed to investigate how ZNF91 influences the XDP-SVA insertion and the

associated molecular phenotype. Using CRISPR/Cas9 we genetically deleted ZNF91 from

XDP induced pluripotent stem cell (iPSC) lines to assess the impact of ZNF91 on the XDP

molecular phenotype. Our findings contribute important insights into the functional role of

ZNF91 in the context of XDP and offer promising prospects for novel therapeutic approaches

in combating XDP.

Methods

XDP iPSC culture

Parental XDP (MIN15i-33363.D and MIN04i-33109.2B) and dSVA (MIN31i-33363.D.3C2

and MIN32i-33109.2B.3A12) iPSC lines were obtained from the WiCell (**Supplemental**

Table 1). XDP probands and respective dSVA iPSCs were cultured at 37 °C with 5% CO_2

on Growth Factor Reduced Matrigel-coated (Corning) plates in mTeSR1 (STEMCELL).

Media was changed daily, and the cells were split when they reached 80-90% confluency

with Accutase. After each passage, 10 μM Y-27632 Rock inhibitor (Tocris) was added to the

media and removed after 24 hours by refreshing the media.

CRISPR☐Cas9 genetic deletion of ZNF91

For CRISPR/Cas9 deletion of ZNF91, two gRNAs designed to target exon 1 of ZNF91 [24]

were cloned and inserted into pX330-SpCas9-HF1 (Addgene #108301). In addition to the

gRNAs, pCAG. GFP (Addgene #11150) was cotransfected for FACS sorting of iPSCs. A day

before transfection, iPSCs were seeded in 60 mm dishes coated with GFR Matrigel and

incubated in mTESR1 with 10 μM Y-27632 Rock inhibitor (Tocris). At the time of

transfection, the cells were 40-60% confluent, and their media was replaced one hour before

transfection with prewarmed mTeSR1 and ROCK inhibitor. For each of the 60 mm dishes,

1.4 μg of each ZNF91-gRNA and 0.2 mg of pCAG. GFP plasmids were transfected. First,

125 µl optiMEM (Thermo Fisher) and 12.5 µl Lipofectamine stem (Thermo Fisher) were

mixed, the plasmids were combined, and optiMEM was added to adjust the volume with the

Lipofectamine mixture. The plasmids and Lipofectamine were mixed and incubated for 10

minutes at room temperature before adding the lipid-DNA mix to the cultures. After

transfection, the cells were incubated for 24 hours. The media was changed the next day with

prewarmed mTeSR1, and the cells were maintained for an additional 24 hours in the same

cell culture condition. For sorting GFP-positive cells, the cells were supplemented with 10

μM Rock inhibitor an hour before Fluorescence-activated cell sorting (FACS). For collecting

single cells, the plates were washed twice with pre-warmed PBS without Ca2+ and Mg2+

(Phosphate-buffered saline), subsequently Versene Solution was added to the plates and

incubated at 37 °C for 7 minutes. The cells were filtered through a 35-mm mesh cell strainer

and sorted on a FACS Aria III with a 100 μm nozzle. Sorted cells were seeded at clonal

density of 6-8x10³ cells per 9.5cm² dishes coated with GFR Matrigel and supplemented by

the Rock inhibitor (10 μM) and mTeSR1. In addition, extra single cells (1-3x10³ cells per

9.5cm² dishes) were collected in MEF (Mouse Embryonic Fibroblasts) supplemented with the

Rock inhibitor (10 μM), Fibroblast Growth Factors (FGF, 8 ng/ml) and ESC medium. The

media was changed the next day and adding the 10 μM Y-27632 Rock inhibitor (Tocris) was

continued for two more days. Single colonies were manually scooped into separated wells of

96-well plates. Propagation of the cells was done by diluting Versene Solution with PBS

(1:3) to use in genotyping, culturing, and freezing.

Genotyping of the expanded cells was done when the cells reached 90-95% confluency. The

cells were treated with a lysis buffer (10 mM Tris-Cl pH 8.0, 10 mM disodium-EDTA, 10

mM NaCl, 0.5% w/v sarcosyl, 40 μg/ml proteinase K) overnight and incubated at 60 °C. The

following day, after precipitating DNA with cold 95% ethanol supplemented with 75 mM

NaCl and washing with 70% ethanol, genomic DNA (gDNA) was extracted and used for

PCR analysis. Four primer sets, targeted internal and external deletion sites of ZNF91, were

designed and used for the detection of successful excision. After selecting ZNF91 knock-out

and Wild-type colonies based on the PCR amplicon sizes, the selected colonies were sent to

Sanger sequencing for confirmation of the interested gene deletion.

Mapping of ZNF91 ChIP-seq data to the XDP-SVA insertion

Publicly available ZNF91 ChIP-seq data from hESCs [24] was mapped to the XDP BAC

clone (AB191243.1). First, raw fastq sequencing reads were trimmed of poor quality and

adapter sequences using trimmomatic [27] (trimmomatic PE -phred33

ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:40). Next, the trimmed reads were aligned to the XDP

BAC clone using BWA-MEM aligner using default settings [28]. Resulting SAM files was

converted into sorted BAM file using samtools [29]. For visualization, BAM files were

converted into bigwig files using deeptools bamCoverage [30] (bamCoverage --

outFileFormat bigwig --binSize 1 --normalizeUsing None --minMappingQuality 1). Bigwig

ZNF91 ChIP-seq coverage results were visualized along the XDP BAC clone using the

Integrative Genomics Viewer (IGV) desktop application [31].

RNA-sequencing

Total RNA sequencing

Three Sanger-validated ZNF91 KO and WT clones per condition were sent for RNA

sequencing. For this, cells were washed once with PBS before collection in TRIzol Reagent.

Total RNA was extracted using TRIzol Reagent (Thermo Fisher), followed by gDNA

removal and RNA purification using an RNA clean and concentrator kit (Zymo Research).

Ribosomal RNA was depleted from total RNA with the rRNA depletion kit (NEB# E6310)

and subsequently prepared for RNA-seq with the NEBNext Ultra Directional RNA Library

Prep Kit (NEB #E7420) at GenomeScan. Samples were sequenced at 150 bp paired-end at an

Illumina Hiseq 4000 device.

Capture RNA-sequencing (Cap-seq)

Two different sets of custom designed MyBaits probes (Arbor Bioscience) were used

throughout this project. The first set of probes (MyBaits1) was designed to target and capture

the *TAF1* cDNA sequence. The second set of probes (MyBaits2, Design ID: D10165*TAF1*)

was designed to target and capture both cDNA and intron sequences of the *TAF1* gene. The

first Cap-seq experiment (using the MyBaits1 probe set) was performed on a total of 12 iPSC

samples from patient 1, including three replicates of a single clone from each cell line. The

second Cap-seq experiment (using the MyBaits2 probe set) was performed on a total of 24

iPSC samples from patient 1 and patient 2, including three replicates of a single clone from

each cell line.

TRIzol RNA isolation was performed as described above. Ribosomal depleted RNA-seq

libraries were prepared using the TruSeq® Stranded Total RNA Library Prep Kit

Human/Mouse/Rat kit (Illumina, 20020596) following manufacturer's instructions. Libraries

were quantified using DNA HS Bioanalyzer chip, and equal molar amounts were pooled

together. Custom MyBaits probes kit was used according to the manufacturer's instructions to

capture and enrich pooled sequencing libraries. Briefly, libraries were blocked using Block

O, C, and X buffers and then hybridized with capture probes for 16-24 hours at 65C. Samples

were washed and cleaned, and KAPA HiFi HotStartReadMix was used to PCR amplify (14

cycles) the libraries. The final amplified and enriched libraries were cleaned using

AmPureXP beads and sequenced on Illumina NextSeq500 using Mid Output flow cell 2x75

cycles.

Analysis of RNA-sequencing data

Raw fastq sequencing reads were trimmed of poor quality and adapter sequences using

FASTP [32] (fastp --qualified_quality_phred 30). Trimmed fastq files were then mapped to

UCSC hg38 human reference genome using STAR [33] (STAR –outFilterMultimapNmax 20

--outFilterMismatchNmax 5 --alignMatesGapMax 0 --alignIntronMax 0 --

outFilterMatchNminOverLread 0.66 --alignEndsType EndToEnd). The human gene

annotation file (Homo_sapiens.GRCh38.104.gtf) was downloaded from ENSEMBL and

converted to UCSC hg38 compatible annotations using the chromToUcsc python script. The

final gene annotation file used also included an annotation for the intronic region upstream of

the XDP-SVA insertion (*TAF1*_partial*TAF1*_intron32, chrX:71424239-71440488). The

STAR-mapped BAM files and our custom gene annotation file were used to generate both

transcript-level and exon-level count tables using FeatureCounts [34] (featureCounts -M -t

exon). Count tables were filtered to only include transcript and exon IDs targeted by MyBaits

probes. Normalization and differential expression analysis were performed using DEseq2

[35]. For both Cap-seq experiments expression was normalized to *TAF1* exons 1-28, and

differential expression was performed using default DEseq2 parameters. A transcript of exon

was considered differentially expressed if padj<0.05 and change was greater than 25%

(L2FC>0.3219 or L2FC<-0.4150).

Intron 32 splicing analysis

Data from both Cap-seq experiments (MyBaits1 and MyBaits2) were used for intron

retention analysis. The number of reads mapping to the last bp of *TAF1* exon 32

(chrX:71424238), and the first bp of intron 32 (chrX:71424239) were extracted from indexed

bam files using the igvtool command line tool [31] (igvtools count -w 1 –query). The number

of reads spliced from exon 32 to exon 33 were extracted from bam files using samtools [29]

by searching the CIGAR string for 29931bp splicing ('29931N') using the following

command (samtools view -F 0x4 $BAM chrX:71424229-71424248 | cut -f 6 | grep '29931N' |

wc -l). The proportion of *TAF1* transcripts with canonical splicing from exon 32 to exon 33

was calculated by dividing the number of reads spliced from exon 32 to exon 33 by the

number of reads mapping to the last bp of *TAF1* exon 32 (chrX:71424238). The proportion of

TAF1 transcripts with intron 32 retention was calculated by dividing the number of reads

mapping to the first bp of intron 32 (chrX:71424239) by the number of reads mapping to the

last bp of *TAF1* exon 32 (chrX:71424238).

Analysis of ZNF91 expression across age

The whole blood microarray expression data used for the analyses described in this manuscript were obtained from The 10,000 Immunomes Project (10Kimmunomes) [36] (https://comphealth.ucsf.edu/app_direct_i/10kimmunomes/) on 06/09/23. The brain RNA-seq expression data, from thirteen distinct regions of the brain, were obtained from V8 of The Genotype-Tissue Expression (GTEx) project [37] (gtex_analysis_v8, rna_seq_data, GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_gene_tpm.gct) on 06/09/23. For the GTEx data, preprocessing involved merging tissue files with subject phenotypes allowing us to match gene expression levels with age for each subject. It's important to note that the GTEx database provides subjects' ages in a ten-year range; for this analysis, we chose to use the lower limit of each range as the representative age. For instance, a subject classified within the 50-59 years old range by GTEx would be considered 50 years old for the analysis. For the gene expression analysis, we investigated the ZNF91 gene along with *TAF1* and a random selection of 18 other ZNF genes from the GTEx database.

Our analysis of gene expression relied on the following Python libraries: pandas, statsmodels, and matplotlib for data processing, statistical modeling, and visualization, respectively. First, the dataset was imported using pandas and two subsets were defined: 'Age' (the independent variable, x) and 'Expression' (the dependent variable, y, indicating gene expression). Next, the statsmodels function 'add_constant' was applied to the 'Age' variable, introducing a constant to the independent variable. This step is important for the ordinary least squares (OLS) regression model (https://www.statsmodels.org/stable/index.html) since it estimates the intercept of the regression line. Following the creation of the model via the sm.OLS function, we fit the model to our data and generated a comprehensive statistical summary. This summary included two critical metrics: the adjusted R-squared value and the p-value. A p-value less than 0.05 was considered statistically significant. Matplotlib was employed to

249 generate   scatterplots   with   fitted   regression   lines   and   heatmaps

250 (https://matplotlib.org/stable/users/index.html).

Results

Generation of stable ZNF91 knockout XDP iPSC lines.

The pathogenic XDP-SVA insertion is situated within intron 32 of the *TAF1* gene. It is a full-

length SVA element comprising the characteristic (CCCTCT)n hexamer repeat, an Alu-like

region, a VNTR, SINE-R, and a Poly-A region [13,38] (**Fig. 1A**). SVAs, like many TE

families, are silenced through the recruitment of members of the KRAB zinc finger (KZNF)

family. The KZNF recruits KAP1 [39,40], along with repressive epigenetic modifiers, which

repress the TE [41–43]. Our research group has recently demonstrated that the ZNF91 KZNF

protein serves as a major repressor of SVA retrotransposons within the human genome,

accomplishing repression by binding to both the Alu-VNTR and VNTR-SINE border regions

within SVA elements [24]. Given that the XDP-SVA insertion contains intact Alu, VNTR,

and SINE regions, it is expected that ZNF91 will bind and repress this specific SVA element.

To visualize the predicted binding sites of ZNF91 at the XDP-SVA element, we mapped

ZNF91 ChIP-seq data from human embryonic stem cells (hESCs) [24] to the XDP-SVA

insertion. The analysis revealed a prominent peak at the Alu-VNTR border region, indicating

that this region is likely bound and repressed by ZNF91 within the pathogenic XDP-SVA

insertion (**Fig. 1A**). This finding is in accordance with prior research by Haring et al., which

similarly demonstrated ZNF91 binding at this specific region within SVA elements in the

human reference genome [24].

It was recently established that the presence of the XDP-SVA insertion leads to dysregulation

of *TAF1* transcription [5,12]. However, it remains unclear how binding of a repressive KZNF

protein to this element influences the XDP molecular phenotype. To assess the functional

implications of ZNF91 binding and repression on the XDP-SVA insertion and associated

molecular phenotype we used CRISPR/Cas9 to genetically delete *ZNF91* in iPSCs derived

from two XDP patients (XDP1 and XDP2) [44] and corresponding isogenic control iPSCs

with XDP-SVA excised (dSVA1 and dSVA2) [5] (**Supplemental Table 1**). For each cell

line, three *ZNF91* knockout (KO) and wild-type (WT) clones were established (**Fig. 1B,**

Supplemental Table 1). To confirm the successful editing of *ZNF91* expression, total RNA-

seq analysis was performed, revealing the absence of *ZNF91* expression in all *ZNF91* KO

(dZNF91) clones and its presence in all *ZNF91* WT clones (**Fig. 1C-D, Supplemental Table**

2). The successful genetic deletion of the *ZNF91* gene in XDP iPSCs provides a valuable

model to investigate the functional consequences of ZNF91 binding and repression on the

XDP-SVA insertion and its associated molecular phenotype.

Knockout of ZNF91 exacerbates the XDP molecular phenotype.

Deletion of ZNF91 increases retention of the *TAF1* intron that harbors the XDP-SVA

insertion.

The presence of the XDP-SVA insertion leads to an elevated level of *TAF1* intron 32

expression [5]. This phenomenon was further validated in our study through analysis of total

RNA-seq data from patient 1 and 2 iPSCs, which showed a marked increase in *TAF1* intron

32 expression in XDP compared to dSVA isogenic control samples (**Fig. 2A, Supplemental**

Table 2). To investigate whether KO of ZNF91 affects this molecular phenotype, we

developed a capture probe set with baits covering the canonical *TAF1* mRNA sequence and

all introns of the *TAF1* gene and performed capture RNA sequencing (Cap-seq) on patient 1

and 2 cell lines. Differential expression (DE) analysis of ZNF91 KO compared with WT

patient cells revealed a significant 130-250% upregulation of intron 32 expression in both

patient cell lines (XDP1-dZNF91 vs. XDP1-WT, padj=1.37E-5, L2FC=1.22; XDP2-dZNF91

vs. XDP2-WT, padj=7.04E-13, L2FC=1.81) (**Fig. 2B-C, Supplemental Table 3**).

Furthermore, analysis of intron 32 splicing showed that canonical splicing was decreased by

~13% and ~32%, while intron 32 retention was increased by ~10% and ~29% in ZNF91 KO

cells compared with WT cells for patients 1 and 2, respectively (**Fig. 2D**). DE analysis

revealed no significant change in intron 32 expression level when ZNF91 was deleted from

the dSVA isogenic control cell lines (dSVA1-dZNF91 vs. dSVA1-WT, padj=0.9670,

L2FC=0.06; dSVA2-dZNF91 vs. dSVA2-WT, padj=0.8754, L2FC=-0.07) (**Fig. 2B-C,**

Supplemental Table 3) and negligible differences in canonical intron 32 splicing and

retention levels were observed in ZNF91 KO compared with WT isogenic control cells (**Fig.**

2D). Similar results were obtained for patient 1 from a pilot Cap-seq experiment (excluding

intron probes) (**Supplemental Fig. S1A-C, Supplemental Table 4**), providing further

support that ZNF91 KO exacerbates the intron 32 molecular phenotype only in the presence

of the XDP-SVA insertion and that increased expression of intron 32 is due in part to failed

splicing and intron retention.

Deletion of ZNF91 decreases transcription of *TAF1* exons located downstream of the

XDP-SVA insertion.

Previous studies have reported a slight decrease in the overall expression level of the *TAF1*

gene due to the presence of the XDP-SVA insertion [5,12]. Subsequent analysis at the exon

level further elucidated that the exons located in proximity to the XDP-SVA insertion site are

primarily responsible for this observed decrease [5]. To investigate the impact of ZNF91 KO

on this molecular phenotype, we utilized Cap-seq data and conducted exon-level differential

expression analysis of the *TAF1* gene. In the case of patient 1, a significant reduction in the

expression of downstream exons (exons 34-36 and 38) was observed upon deletion of ZNF91

in XDP1 iPSCs (XDP1-dZNF91 vs. XDP1-WT). However, no change in downstream exon

expression was observed when ZNF91 was deleted in dSVA isogenic control cells (dSVA1-
dZNF91 vs. dSVA1-WT) (**Fig. 3A-C, Supplemental Fig. S2, Supplemental Table 3-4**).
Similar results were obtained for patient 2, where a significant reduction in the expression of
all downstream exons (exons 33-38) was observed upon deletion of ZNF91 in XDP2 iPSCs
(XDP2-dZNF91 vs. XDP2-WT). Again, no change in downstream exon expression was
observed when ZNF91 was deleted in isogenic control dSVA2 iPSCs (dSVA2-dZNF91 vs.
dSVA2-WT) (**Fig. 3A-C, Supplemental Table 3**). These findings provide compelling
evidence that the loss of ZNF91 exacerbates the *TAF1* exonic molecular phenotype
associated with XDP. Moreover, the absence of the effect in the dSVA isogenic control cells
without XDP-SVA insertions underscores the specific regulatory impact of ZNF91 on the
disease-causing XDP-SVA insertion.

ZNF91 expression decreases with age.

Despite the presence of the causative XDP-SVA mutation in all cells from birth, the onset of
the XDP disease occurs later in life and specifically impacts certain regions of the brain [45–
47]. The factors initiating the late onset and the brain-specific manifestations remain elusive.
In our study, we provide compelling evidence that downregulation of ZNF91 expression
exacerbates the molecular phenotype of XDP. Based on these findings, we hypothesize that
an age-related reduction in *ZNF91* expression in the brain contributes to the disease onset. To
investigate whether a general age-related reduction in ZNF91 expression exists in humans,
we performed linear regression analysis on whole blood transcriptomic data from the 10K
Immunomes Project [36]. To provide a single value summarizing both the magnitude and
direction of the correlation, we devised a metric we call the Directional Coefficient of
Determination (dR^2). This metric is computed by multiplying the coefficient of determination
(R^2) by the sign of the correlation coefficient (r). It ranges from -1 to 1, with values closer to -

1 or 1 indicating stronger negative or positive relationships, respectively. Our analysis

revealed a significant inverse correlation between *ZNF91* expression level and age (dR^2 = -

0.062, p-value = 8.27E-06) (**Fig. 4A**). This result implies that approximately 6.2% of the

variation in *ZNF91* expression can be explained by age in the regression model. This finding

suggests that age plays a critical role in the regulation of *ZNF91* expression in blood, with

older individuals exhibiting lower levels of *ZNF91* than younger individuals. To investigate

the uniqueness of this observation for ZNF91, we extended our analysis to include 10

randomly selected ZNF genes. Intriguingly, none of these genes exhibited a significant

inverse correlation with age (**Fig. 4B**), underscoring the distinct regulatory relationship

between age and ZNF91 expression in the blood.

To investigate if the correlation between ZNF91 expression and age is also observed in brain

regions implicated in XDP, we analyzed RNA-seq data from the GTEx database [37]. This

encompassed 49 different tissues, of which 13 were distinct brain regions. A total of 32% of

the tissues analyzed (16 out of 49) exhibited a significant inverse relationship between

ZNF91 expression and age (**Supplemental Fig. S3**). This correlation was predominantly

observed in brain tissues (12 out of 16), underscoring the specificity of the age-related

reduction in *ZNF91* expression in the brain. Notably, a significant inverse correlation was

observed in the striatal regions impacted by XDP, including the Caudate Basal Ganglia (dR^2

= -0.02145, p-value = 0.02157) and the Putamen Basal Ganglia (dR^2= -0.02153, p-value =

0.03576) [48,49] (**Fig. 4C-D**). The correlation was even more pronounced in the frontal

cortex (dR^2= -0.10354, p-value = 2.02E-06) and cerebellum (dR^2= -0.0484, p-value = 5.79E-

04) - extrastriatal regions recently linked with structural thinning and volume loss in XDP

patients [49,50] (**Fig. 4E-F**). These results indicate that age-related reductions in ZNF91

expression, especially within brain regions affected by XDP, may influence the disease's

onset or progression.

Discussion

In this study, we investigated the impact of genetic deletion of ZNF91, a potent repressor of SVA elements, on the molecular phenotype associated with XDP. We demonstrated that knockout of ZNF91 aggravates the molecular phenotype, including an increase in *TAF1* intron 32 retention and a decrease in the expression of *TAF1* exons downstream of the XDP-SVA insertion. Additionally, our data show that there is a significant inverse correlation between age and ZNF91 expression levels in both whole blood and the brain. The results of this study provide important insights into the functional role of ZNF91 in the context of XDP. By observing the aggravated molecular phenotype in the absence of ZNF91, we can conclude that ZNF91 is an endogenous repressor of the molecular phenotype associated with XDP. Further, the observation that ZNF91 expression levels significantly decrease with age supports its potential involvement in disease onset and/or progression, which is known to occur later in life [45]. In summary, the findings from our study support the following proposed model: the age-related decline in ZNF91 expression unleashes the full pathogenic potential of the XDP-SVA insertion, thereby driving the onset of the disease (**Fig. 5**).

This study represents a significant milestone as it is the first to successfully modulate *TAF1* dysregulation without altering any of the XDP-specific genetic variations [1,5,38]. Previous work utilizing CRISPR/Cas9 deletion of the XDP-SVA has been instrumental in dissecting the causal mechanism of XDP, revealing that the XDP-SVA insertion drives the XDP-specific *TAF1* transcriptional dysregulation [5,12]. The results of this study align with these previous findings and provide novel insights into how ZNF91 impacts the severity of the molecular phenotype. Given ZNF91's established role as a repressor of SVAs, our results strongly suggest that epigenetic repression of the XDP-SVA by ZNF91 serves as a critical regulatory mechanism that keeps the molecular phenotype under control. This indicates a

potential direct relationship between ZNF91-mediated epigenetic control and disease

severity. These new findings provide valuable knowledge for understanding the underlying

mechanisms of XDP and may contribute to the development of epigenetic-based therapies for

this disorder.

Previous research has established the role of ZNF91 in binding and repressing the majority of

SVAs in the human reference genome [22,24]. Our study builds upon this work and

demonstrates the direct involvement of ZNF91 with a polymorphic disease-associated SVA.

Moreover, our results highlight the influence of this interaction on the molecular phenotype

of the disease. This underscores the importance of ZNF91 as a key regulator of SVA

elements and its potential implications in the pathogenesis of XDP and other SVA-associated

diseases. In addition to XDP, polymorphic SVA insertions have previously been associated

with various other human diseases, including Lynch syndrome [51], hemophilia B [52],

Fukuyama-type congenital muscular dystrophy (FCMD) [53], X-linked agammaglobulinemia

(XLA) [54,55], autosomal recessive hypercholesterolemia (ARH) [56], and

neurofibromatosis type I (NF1) [57]. Our findings may open new avenues of investigation

into the mechanisms underlying these other disease-associated SVAs and their impact on

disease susceptibility and progression.

The age of onset of XDP is highly variable (12-64 years old) [58], studies looking at the

hexamer repeat length and nearby single nucleotide polymorphisms (SNPs) have succeeded

in partially explaining this age-at-onset variability (50% and 13%, respectively) [4,58,59].

However, a large proportion (37%) of the age-at-onset variability is still unexplained. Given

the findings of this study, it is reasonable to speculate that ZNF91 expression levels could

contribute to the age-at-onset variability in XDP patients. Those with lower ZNF91

expression levels might experience more severe *TAF1* dysregulation, leading to an earlier

onset of the disease. These epigenetic and transcriptomic differences go undetected by

traditional DNA-based genome wide association studies (GWAS).

It is important to acknowledge that all experiments in this study were conducted using iPSCs,

which primarily mimic early embryonic development—a stage seemingly unaffected by

XDP. Consequently, the findings obtained from these cell types may not fully represent the

events occurring in striatal medium spiny neurons (MSNs), which are predominantly affected

by the disease [46]. Recent studies have reported the successful generation of MSNs from

iPSCs [60–62]. The employment of iPSC-derived MSNs would provide more comprehensive

insights into how ZNF91 deletion impacts the molecular phenotype in the specific neuronal

cell types affected by XDP. However, fully recapitulating the affected cell types using any *in*

vitro model poses challenges, considering that cell lines are limited to several months of

culture, whereas the average age of onset for XDP is approximately 40 years old [45]. This

discrepancy in "age" between *in vitro* models and the actual age of onset poses a difficulty in

fully capturing the late-onset component of XDP *in vitro*. Despite these challenges, the use of

iPSCs in this study provides valuable initial data and lays the groundwork for further research

in this area. The combination of *in vitro* models with other approaches, such as animal

models or postmortem human brain tissue analysis, may offer complementary insights into

the complex disease mechanisms underlying XDP.

Previous investigators have proposed that guanine-rich sequences within the XDP-SVA

might be susceptible to forming non-canonical DNA structures, called G-quadruplexes (G4)

[4,63]. An intriguing hypothesis arising from our ZNF91 KO findings is that in the absence

of ZNF91 binding and/or repression, the intronic XDP-SVA might be susceptible to the

formation of G4 structures. As previous studies suggest, the presence of G4 structures within

a gene can hinder RNA polymerase during transcription [64,65]. Blockage of RNA

polymerase, and subsequent disruption of splicing, could explain the observed increase in

intron 32 retention and subsequent decrease in downstream exon expression. While this

hypothesis offers a compelling explanation for our results, additional investigations are

essential to test this.

Several unanswered questions and intriguing directions for future research have emerged

from this study, which hold the potential to further enhance our understanding of how ZNF91

expression may contribute to XDP pathogenesis. First, it remains unclear whether reduction

of ZNF91 expression levels can trigger similar effects as its complete knockout and whether

there is a "threshold" level of ZNF91 expression required for maximal repression of the

phenotype. Investigating the consequences of varying ZNF91 expression levels, ideally in

MSNs, would shed light on its dosage-dependent effects on the XDP-SVA insertion and the

associated molecular phenotype. Additionally, it remains unclear whether the aggravated

molecular phenotype observed upon loss of ZNF91 contributes to the onset or progression of

this disease and whether the aggravation is caused simply by loss of ZNF91 binding or

epigenetic derepression of the XDP-SVA insertion. Addressing these questions is important

for determining whether modulation of ZNF91 expression and/or repression of the XDP-SVA

are viable therapeutic approaches.

In conclusion, this study provides important insights into the functional role of ZNF91 in the

context of XDP. The results demonstrate that ZNF91 deletion aggravates TAF1

dysregulation, indicating that ZNF91 acts as an endogenous repressor of the XDP-SVA and

the associated molecular phenotype. Importantly, this study is the first to successfully

modulate the XDP molecular phenotype without altering any XDP-specific genetic variants

in the *TAF1* region. Our findings highlight the significance of ZNF91 as a key regulator of

SVA elements and its potential implications in disease pathogenesis, not only in XDP but

also in other SVA-associated disorders. In the broader context, our research contributes to the

growing knowledge of intronic gene regulation in disease pathogenesis. Ultimately, the

identification of ZNF91 as a potential therapeutic target in XDP and other SVA-associated

diseases offers promising prospects for innovative approaches in disease treatment.

List of abbreviations

ARH	Autosomal recessive hypercholesterolemia
Cap-seq	Capture RNA-sequencing
Cas9	CRISPR□associated protein 9
cDNA	Complementary DNA
ChIP-seq	Chromatin Immunoprecipitation Sequencing
CRISPR	Clustered regularly interspaced short palindromic repeats
DE	Differential expression
dSVA	XDP-SVA excised
DYT3	X-linked dystonia-parkinsonism (Alternative name)
ESC	Embryonic Stem Cell
FACS	Fluorescence-activated cell sorting
FCMD	Fukuyama-type congenital muscular dystrophy
FGF	Fibroblast Growth Factor
G4	G-quadruplexes
gDNA	Genomic DNA
gRNA	Guide RNA
hESCs	Human Embryonic Stem Cells
iPSC	Induced pluripotent stem cell
KAP1	KRAB-associated protein 1
KO	Knockout
KRAB	Krüppel-associated box
KZNF	KRAB zinc finger
L2FC	Log2 fold change
NF1	Neurofibromatosis type I
OLS	Ordinary least squares
padj	Adjusted P-value
PBS	Phosphate-Buffered Saline
PEI	Polyethylenimine
Poly-A	Poly-adenosine
RNA-seq	RNA-sequencing
SINE	Short Interspersed Nuclear Element
SVA	SINE-VNTR-Alu
TAF1	TATA-binding protein-associated factor-1
TE	Transposable Element
TFIID	Transcription Factor II D
VCM	Virus Containing Media
VNTR	Variable Number Tandem Repeat
WT	Wild type
XDP	X-linked dystonia-parkinsonism
XDP-SVA	XDP-specific SVA
XLA	X-linked agammaglobulinemia
ZNF91	Zinc Finger Protein 91

Declarations

Ethics approval and consent to participate

The present study involves the use of deidentified XDP patient iPSCs that were derived from a previous study [44]. In the primary study generating the iPSC lines used in this research [44], patients provided informed written consent for their biological samples to be used for scientific investigation.

Consent for publication

All authors have given their consent for publication.

Availability of data and materials

The datasets generated and/or analyzed during the current study are available in the NCBI Sequencing read archive (SRA; https://www.ncbi.nlm.nih.gov/sra/) repository under accession number PRJNA1004173. The ZNF91 ChIP-seq data was downloaded from NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) accession number GSE162571. The XDP BAC clone "Homo sapiens DNA, transposon SVA sequence" was downloaded from the NCBI Nucleotide database (https://www.ncbi.nlm.nih.gov/nuccore/AB191243.1/) accession number AB191243.1.

Competing interests

The authors declare that they have no competing interests.

Funding

Funding for this study was provided by the Massachusetts General Hospital Collaborative Center for X-linked Dystonia-Parkinsonism.

Authors' contributions

Conceptualization: FMJ and JLR; Methodology: JLR, SR, JEB, and AK; Validation: JLR, SR, and JEB; Formal Analysis: JLR and AK; Investigation: JLR, SR, and JEB; Resources: SR, CAV, and DCB; Writing – Original Draft: JLR; Writing – Review & Editing: JLR, DCB, GF, and FMJ; Visualization: JLR and AK; Supervision: FMJ and JLR; Project Administration: FMJ and JLR; Funding Acquisition: FMJ

Acknowledgments

We would like to acknowledge the research subjects and clinical research staff of the Dystonia Partners Research Bank (Massachusetts General Hospital, Boston, USA) for their effort, foundational contributions to this work and commitment to improving the lives of people and families with XDP. Funding for this study was provided by the MGH Collaborative Center for X-Linked Dystonia-Parkinsonism (CCXDP). We would also like to thank Christopher Bragg, Amy Alessi, Johan Jakobsson, and all members of CCXDP for their valuable discussions and guidance on the project; all members of the Jacobs lab for their engaging discussions, and constructive feedback; Gonzalo Congrains Sotomayor for technical assistance with FACS; UvA sequencing core for technical assistance with Illumina sequencing; the Evolutionary Neurogenomics Group, the Molecular Neuroscience group, and others at the Swammerdam Institute for Life Sciences (SILS) for helpful discussions.

Figures

File name: **Figure 1**
File format: .pdf
Title: Generation of stable ZNF91 knockout XDP iPSC lines
Legend: Fig 1. Generation of stable ZNF91 knockout XDP iPSC lines. (A) Schematic of TAF1, XDP-SVA insertion, and ZNF91 ChIP-seq signal. (Top) Ideogram of human Chromosome X, highlighting the genomic location of the TAF1 gene. (Middle) Detailed view of the TAF1 gene structure, the XDP-SVA insertion within intron 32 is highlighted in cyan. (Bottom) Schematic depiction of the XDP-SVA insertion, showing hESC ZNF91 ChIP-seq

signal over the distinct subregions of the XDP-SVA element. (B) Schematic of CRISPR/Cas9
ZNF91 knockout in XDP and dSVA iPSCs. (C) Total RNA-seq coverage tracks at the ZNF91
locus highlighting loss of ZNF91 gene expression in ZNF91 knockout cell lines. Tracks are
scaled to total library size. *Denotes samples used for capture RNA-seq analysis. (D)
Quantification of ZNF91 gene expression. Each bar represents the mean ± SEM of N=3
replicate clones of each condition. DEseq2 default parameters were used for normalization.
525
File name: **Figure 2**
File format: .pdf
Title: Deletion of ZNF91 increases retention of the TAF1 intron that harbors the XDP-SVA
insertion
Legend: Fig 2. Deletion of ZNF91 increases retention of the TAF1 intron that harbors the
XDP-SVA insertion. (A) Total RNA-seq coverage tracks at the TAF1 locus highlighting
increase of intron 32 (i32) expression in XDP cells (red) compared to dSVA isogenic controls
(blue). Tracks are scaled to TAF1 exons 1-28. (B) Capture RNA-seq coverage tracks at TAF1
intron 32, highlighting increased i32 expression in ZNF91 knockout XDP (purple) compared
to WT XDP (red) cells. (C) Quantification of TAF1 i32 expression. A significant
upregulation in ZNF91 KO XDP (purple) compared to WT XDP (red) cells is observed. Each
bar represents the mean ± SEM of N=3 replicates (three replicates of single clones) and the
DE analysis was performed using DESeq2 with default parameters (Wald test, corrected for
multiple testing using the Benjamini and Hochberg method), except for normalization to
TAF1 exons 1-28. Genes were considered differentially expressed if they had an adjusted p-
value (padj) < 0.05 and exhibited greater than 25% change in expression (fold-change > 1.25
or <0.75). *padj<0.05, **padj<0.01, ***padj<0.001, ns = not significant. (D) TAF1 intron 32
splicing results. The percentage of reads with canonical splicing of intron 32 are displayed on
the right, and the percentage of exon 32 reads with retained intron are displayed next on the
left. Percentages are expressed as the mean ± SEM of three replicate samples. Statistical
significance was assessed using a two-sided t-test, *p-value<0.05, ns = not significant.
547
File name: **Figure 3**
File format: .pdf
Title: Deletion of ZNF91 decreases transcription of TAF1 exons located downstream of the
XDP-SVA insertion
Legend: Fig 3. Deletion of ZNF91 decreases transcription of TAF1 exons located
downstream of the XDP-SVA insertion. (A) Capture RNA-seq coverage tracks at TAF1
exons. A reduction of exon 33-38 expression is observed in ZNF91 knockout XDP (purple)
compared to WT XDP (red) cells. Tracks are scaled to TAF1 exons 1-28. (B) Quantification
of TAF1 exon 1-38 expression level in ZNF91 knockout XDP (purple) and WT XDP (red)
cells. (C) Quantification of TAF1 exon 1-38 expression level in ZNF91 knockout dSVA
(green) and WT dSVA (blue) cells. For panels B and C, each bar represents the mean ± SEM
of three independent replicates (three replicates of single clones) and the DE analysis was
performed using DESeq2 with default parameters (Wald test, corrected for multiple testing
using the Benjamini and Hochberg method), except for normalization to TAF1 exons 1-28.
Genes were considered differentially expressed if they had an adjusted p-value (padj) < 0.05
and exhibited greater than 25% change in expression (fold-change > 1.25 or < 0.75).
*padj<0.05. For all panels, exons 33-38, downstream of the XDP-SVA insertion are
highlighted with grey background.
566
File name: **Figure 4**
File format: .pdf

Title: ZNF91 expression decreases with age

Legend: Fig 4. ZNF91 expression decreases with age. (A) Scatter plot illustrating the
relationship between age (x-axis) and ZNF91 TPM expression level (y-axis) in whole blood
samples from 311 individuals. Data sourced from the publicly available 10K Immunomes
database. Linear regression $R2$ and significance level are included below plot. (B) Heatmap
illustrating the linear regression results of gene expression level across age for 10 random
ZNF genes in whole blood (10K immunomes data). Each box represents the directional R2
(dR2) value derived from the regression analysis, with the color intensity indicating the
strength of the relationship. Red boxes signify an inverse correlation between ZNF91
expression and age, while blue boxes indicate a direct correlation. (C-F). Scatter plots
illustrating the relationship between age and ZNF91 expression level in (C-D) striatal and (E-
F) extra-striatal brain regions impacted in XDP. Data sourced from the publicly available
GTEx database. Linear regression R^2 and significance level are included below plot.

File name: **Figure 5**
File format: .pdf
Title: Proposed model: age-related reduction of ZNF91 unleashes full pathogenic potential of
XDP-SVA
Legend: Figure 5. Proposed model: age-related reduction of ZNF91 unleashes full pathogenic
potential of XDP-SVA. ZNF91 expression significantly decreases with age. Further, ZNF91
interacts specifically with the disease-specific XDP-SVA insertion and is an endogenous
repressor of the associated molecular phenotypes. Thus, we propose that the age-related loss
of ZNF91 leads to increased TAF1 intron 32 retention and decreased levels of full-length
TAF1 transcripts in XDP patients. In contrast, the age-related loss of ZNF91 has no effect on
TAF1 transcription in healthy individuals since they lack the XDP-SVA insertion.

Additional files
595

File name: Supplemental Table 1
File format: .xlsx
Title of data: XDP and dSVA iPSC line details
Description of data: Details of XDP and dSVA iPSC cell lines used throughout study.

File name: Supplemental Table 2
File format: .xlsx
Title of data: Total RNA-seq DE results
Description of data: Differential expression (DE) results from total RNA-seq data. N=3
replicate clones of each condition. TAF1 exons and intron 32 (i32) were considered
differentially expressed if they had an adjusted p-value (padj) < 0.05 and exhibited greater
than 25% change in expression (fold-change > 1.25 or <0.75). (*padj<0.05, **padj<0.01,
***padj<0.001). L2FC = Log2 fold-change. DEseq2 default parameters were used for
normalization and statistical analysis (Wald test, corrected for multiple testing using the
Benjamini and Hochberg method).

File name: Supplemental Table 3
File format: .xlsx
Title of data: Capture RNA-seq DE results (including intron capture probes)

Description of data: Differential expression (DE) results from capture RNA-seq experiment (using capture probe set including TAF1 intron sequences). N=3 replicates (three replicates of single clones). TAF1 exons and intron 32 (i32) were considered differentially expressed if they had an adjusted p-value (padj) < 0.05 and exhibited greater than 25% change in expression (fold-change > 1.25 or < 0.75). (*padj<0.05, **padj<0.01, ***padj<0.001). L2FC = Log2 fold-change. The DE analysis was performed using DESeq2 with default parameters (Wald test, corrected for multiple testing using the Benjamini and Hochberg method), except for normalization to TAF1 exons 1-28.

File name: Supplemental Table 4
File format: .xlsx
Title of data: Capture RNA-seq DE results (excluding intron capture probes)
Description of data: Differential expression (DE) results from capture RNA-seq experiment (using capture probe set excluding TAF1 intron sequences). N=3 replicates (three replicates of single clones). TAF1 exons and intron 32 (i32) were considered differentially expressed if they had an adjusted p-value (padj) < 0.05 and exhibited greater than 25% change in expression (fold-change > 1.25 or <0.75). (*padj<0.05, **padj<0.01, ***padj<0.001). L2FC = Log2 fold-change, i32= intron 32. The DE analysis was performed using DESeq2 with default parameters (Wald test, corrected for multiple testing using the Benjamini and Hochberg method), except for normalization to TAF1 exons 1-28.

File name: Supplemental Figures
File format: .pdf
Title of data: Supplemental Figures
Description of data: ***Supplemental Figure S1***. Deletion of ZNF91 increases retention of the TAF1 intron that harbors the XDP-SVA insertion (Pilot Capture RNA-seq results excluding intron probes). (A) Capture RNA-seq coverage tracks at TAF1 exon32/intron32 boundary, highlighting increased i32 expression in ZNF91 knockout XDP (purple) compared to WT XDP (red) cells. (B) Quantification of TAF1 i32 expression. A marked, but not statistically significant, increase of i32 in ZNF91 knockout XDP (purple) compared to WT XDP (red) cells is observed. Each bar represents the mean ± SEM of N=3 independent replicates (three replicates of single clones) and the DE analysis was performed using DESeq2 with default parameters (Wald test, corrected for multiple testing using the Benjamini and Hochberg method), except for normalization to TAF1 exons 1-28. Genes were considered differentially expressed if they had an adjusted p-value (padj) < 0.05 and exhibited greater than 25% change in expression (fold-change > 1.25 or <0.75). *padj<0.05, **padj<0.01, ***padj<0.001, ns = not significant. (C) TAF1 intron 32 splicing results. The percentage of reads with canonical splicing of intron 32 are displayed on the right, and the percentage of exon 32 reads with retained intron are displayed next on the left. Percentages are expressed as the mean ± SEM of three replicate samples. Statistical significance was assessed using a two-sided t-test, *p-value<0.05, ns = not significant. ***Supplemental Figure S2***. Deletion of ZNF91 decreases transcription of TAF1 exons located downstream of the XDP-SVA insertion (Pilot capture RNA-seq results excluding intron probes). (A) Capture RNA-seq coverage tracks at TAF1 exons. A reduction of exon 33-38 expression is observed in ZNF91 knockout XDP (purple) compared to WT XDP (red) cells. Tracks are scaled to TAF1 exons 1-28. (B) Quantification of TAF1 exon 1-38 expression level in ZNF91 knockout XDP (purple) and WT XDP (red) cells. (C) Quantification of TAF1 exon 1-38 expression level in ZNF91 knockout dSVA (green) and WT dSVA (blue) cells. ***Supplemental Figure S3***. ZNF91 expression decreases with age specifically in the brain. Heatmap illustrating the linear regression results of ZNF91 expression across age for various GTEx tissues. Each box

665 represents the directional R^2 (dR^2) value derived from the regression analysis, with the color
666 intensity indicating the strength of the relationship. Red boxes signify an inverse correlation
667 between ZNF91 expression and age, while blue boxes indicate a direct correlation. Only
668 GTEx tissues with more than 30 samples were included in this analysis.
669

References

1. Nolte D, Niemann S, Müller U. Specific sequence changes in multiple transcript system DYT3 are associated with X-linked dystonia parkinsonism. Proc National Acad Sci. 2003;100:10347–52.

2. Evidente VGH, Advincula J, Esteban R, Pasco P, Alfon JA, Natividad FF, et al. Phenomenology of "Lubag" or X☐linked dystonia–parkinsonism. Movement Disord. 2002;17:1271–7.

3. Lee LV, Pascasio FM, Fuentes FD, Viterbo GH. Torsion dystonia in Panay, Philippines. Adv Neurol. 1976;14:137–51.

4. Bragg DC, Mangkalaphiban K, Vaine CA, Kulkarni NJ, Shin D, Yadav R, et al. Disease onset in X-linked dystonia-parkinsonism correlates with expansion of a hexameric repeat within an SVA retrotransposon in TAF1. Proc Natl Acad Sci. 2017;114:E11020–8.

5. Aneichyk T, Hendriks WT, Yadav R, Shin D, Gao D, Vaine CA, et al. Dissecting the Causal Mechanism of X-Linked Dystonia-Parkinsonism by Integrating Genome and Transcriptome Assembly. Cell. 2018;172:897-909.e21.

6. Thomas MC, Chiang C-M. The General Transcription Machinery and General Cofactors. Crit Rev Biochem Mol. 2006;41:105–78.

7. Papai G, Weil PA, Schultz P. New insights into the function of transcription factor TFIID from recent structural studies. Curr Opin Genet Dev. 2011;21:219–24.

8. Anandapadamanaban M, Andresen C, Helander S, Ohyama Y, Siponen MI, Lundström P, et al. High-resolution structure of TBP with TAF1 reveals anchoring patterns in transcriptional regulation. Nat Struct Mol Biol. 2013;20:1008–14.

9. Grünberg S, Hahn S. Structural insights into transcription initiation by RNA polymerase II. Trends Biochem Sci. 2013;38:603–11.

10. Malkowska M, Kokoszynska K, Rychlewski L, Wyrwicz L. Structural bioinformatics of

the general transcription factor TFIID. Biochimie. 2013;95:680–91.

11. Gudmundsson S, Wilbe M, Filipek-Górniok B, Molin A-M, Ekvall S, Johansson J, et al.

TAF1, associated with intellectual disability in humans, is essential for embryogenesis and

regulates neurodevelopmental processes in zebrafish. Sci Rep-uk. 2019;9:10730.

12. Rakovic A, Domingo A, Grütz K, Kulikovskaja L, Capetian P, Cowley SA, et al. Genome

editing in induced pluripotent stem cells rescues TAF1 levels in X☐linked

dystonia☐parkinsonism. Movement Disord. 2018;33:1108–18.

13. Wang H, Xing J, Grover D, Hedges DJ, Han K, Walker JA, et al. SVA elements: a

hominid-specific retroposon family. J Mol Biol. 2005;354:994–1007.

14. Kazazian HH, Wong C, Youssoufian H, Scott AF, Phillips DG, Antonarakis SE.

Haemophilia A resulting from de novo insertion of L1 sequences represents a novel

mechanism for mutation in man. Nature. 1988;332:164–6.

15. Batzer MA, Gudi VA, Mena JC, Foltz DW, Herrera RJ, Deininger PL. Amplification

dynamics of human-specific (HS) alu family members. Nucleic Acids Res. 1991;19:3619–23.

16. Ostertag EM, Goodier JL, Zhang Y, Kazazian HH. SVA elements are nonautonomous

retrotransposons that cause disease in humans. Am J Hum Genet. 2003;73:1444–51.

17. Stewart C, Kural D, Stromberg MP, Walker JA, Konkel MK, Stutz AM, et al. A

comprehensive map of mobile element insertion polymorphisms in humans. PLoS Genet.

2011;7:e1002236.

18. Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A

global reference for human genetic variation. Nature. 2015;526:68–74.

19. Kaer K, Speek M. Retroelements in human disease. Gene. 2013;518:231–41.

20. Hancks DC, Kazazian HH. Roles for retrotransposon insertions in human disease. Mobile

DNA. 2016;7:9–28.

21. Hancks DC, Jr. HHK. Active human retrotransposons: variation and disease. Curr Opin

Genet Dev. 2012;22:191–203.

22. Jacobs FMJ, Greenberg D, Nguyen N, Haeussler M, Ewing AD, Katzman S, et al. An

evolutionary arms race between KRAB zinc-finger genes ZNF91/93 and SVA/L1

retrotransposons. Nature. 2014;516:1–18.

23. Pontis J, Planet E, Offner S, Turelli P, Duc J, Coudray A, et al. Hominoid-Specific

Transposable Elements and KZFPs Facilitate Human Embryonic Genome Activation and

Control Transcription in Naive Human ESCs. Cell Stem Cell. 2019;24:724-735.e5.

24. Haring NL, Bree EJ van, Jordaan WS, Roels JRE, Sotomayor GC, Hey TM, et al. ZNF91

deletion in human embryonic stem cells leads to ectopic activation of SVA retrotransposons

and up-regulation of KRAB zinc finger gene clusters. Genome Res. 2021;31:551–63.

25. Lippman Z, Gendrel A-V, Black M, Vaughn MW, Dedhia N, McCombie WR, et al. Role

of transposable elements in heterochromatin and epigenetic control. Nature. 2004;430:471–6.

26. Rebollo R, Karimi MM, Bilenky M, Gagnier L, Miceli-Royer K, Zhang Y, et al.

Retrotransposon-Induced Heterochromatin Spreading in the Mouse Revealed by Insertional

Polymorphisms. Plos Genet. 2011;7:e1002301.

27. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data.   Bioinformatics   [Internet].   2014;30:2114–20.   Available   from:

https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btu170

28. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv. 2013;

29. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

30. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a

next generation web server for deep-sequencing data analysis. Nucleic Acids Res.

2016;44:W160–5.

31. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.

Integrative genomics viewer. Nat Biotechnol. 2011;29:24–6.

32. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;34:i884–90.

33. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

34. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30:923–30.

35. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

36. Zalocusky KA, Kan MJ, Hu Z, Dunn P, Thomson E, Wiser J, et al. The 10,000

Immunomes Project: Building a Resource for Human Immunology. Cell Rep. 2018;25:513-

522.e3.

37. Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, et al. The Genotype-Tissue

Expression (GTEx) project. Nat Genet. 2013;45:580–5.

38. Makino S, Kaji R, Ando S, Tomizawa M, Yasuno K, Goto S, et al. Reduced neuron-

specific expression of the TAF1 gene is associated with X-linked dystonia-parkinsonism. Am

J Hum Genet. 2007;80:393–406.

39. Nielsen AL, Ortiz JA, You J, Oulad Abdelghani M, Khechumian R, Gansmuller A, et al.

Interaction with members of the heterochromatin protein 1 (HP1) family and histone

deacetylation are differentially involved in transcriptional silencing by members of the TIF1

family. EMBO J. 1999;18:6385–95.

40. Sripathy SP, Stevens J, Schultz DC. The KAP1 Corepressor Functions To Coordinate the

Assembly of De Novo HP1-Demarcated Microenvironments of Heterochromatin Required

for KRAB Zinc Finger Protein-Mediated Transcriptional Repression. Mol Cell Biol.

2006;26:8623–38.

41. Schultz DC, Friedman JR, Rauscher FJ. Targeting histone deacetylase complexes via

KRAB-zinc finger proteins: the PHD and bromodomains of KAP-1 form a cooperative unit

that recruits a novel isoform of the Mi-2α subunit of NuRD. Genes Dev. 2001;15:428–43.

42. Schultz DC, Ayyanathan K, Negorev D, Maul GG, Rauscher FJ. SETDB1: a novel KAP-

1-associated histone H3, lysine 9-specific methyltransferase that contributes to HP1-mediated

silencing of euchromatic genes by KRAB zinc-finger proteins. Genes Dev. 2002;16:919–32.

43. Turelli P, Castro-Diaz N, Marzetta F, Kapopoulou A, Raclot C, Duc J, et al. Interplay of

TRIM28 and DNA methylation in controlling human endogenous retroelements. Genome

Res. 2014;24:1260–70.

44. Ito N, Hendriks WT, Dhakal J, Vaine CA, Liu C, Shin D, et al. Decreased N-TAF1

expression in X-linked dystonia-parkinsonism patient-specific neural stem cells. Dis Model

Mech. 2016;9:451–62.

45. Lee LV, Maranon E, Demaisip C, Peralta O, Borres-Icasiano R, Arancillo J, et al. The

natural history of sex-linked recessive dystonia parkinsonism of Panay, Philippines (XDP).

Parkinsonism Relat D. 2002;9:29–38.

46. Goto S, Lee LV, Munoz EL, Tooyama I, Tamiya G, Makino S, et al. Functional anatomy

of the basal ganglia in X□linked recessive dystonia□parkinsonism. Ann Neurol. 2005;58:7–

17.

47. Acuna P, Supnet-Wells ML, Spencer NA, Guzman JK de, Russo M, Hunt A, et al.

Establishing a natural history of X-linked dystonia parkinsonism. Brain Commun.

2023;5:fcad106.

48. Pasco PMD, Ison CV, Muňoz EL, Magpusao NS, Cheng AE, Tan KT, et al.

Understanding XDP Through Imaging, Pathology, and Genetics. Int J Neurosci.

2011;121:12–7.

49. Arasaratnam CJ, Singh-Bains MK, Waldvogel HJ, Faull RLM. Neuroimaging and

neuropathology studies of X-linked dystonia parkinsonism. Neurobiol Dis. 2021;148:105186.

50. Brüggemann N, Heldmann M, Klein C, Domingo A, Rasche D, Tronnier V, et al.

Neuroanatomical changes extend beyond striatal atrophy in X-linked dystonia parkinsonism.

Parkinsonism Relat D. 2016;31:91–7.

51. Klift HM van der, Tops CM, Hes FJ, Devilee P, Wijnen JT. Insertion of an SVA element,

a nonautonomous retrotransposon, in PMS2 intron 7 as a novel cause of lynch syndrome.

Hum Mutat. 2012;33:1051–5.

52. Nakamura Y, Murata M, Takagi Y, Kozuka T, Nakata Y, Hasebe R, et al. SVA

retrotransposition in exon 6 of the coagulation factor IX gene causing severe hemophilia B.

Int J Hematol. 2015;102:134–9.

53. Kobayashi K, Nakahori Y, Miyake M, Matsumura K, Kondo-Iida E, Nomura Y, et al. An

ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy.

Nature. 1998;394:388–92.

54. Rohrer J, Minegishi Y, Richter D, Eguiguren J, Conley ME. Unusual Mutations in Btk:

An Insertion, a Duplication, an Inversion, and Four Large Deletions. Clin Immunol.

1999;90:28–37.

55. Conley ME, Partain JD, Norland SM, Shurtleff SA, Kazazian HH. Two independent

retrotransposon insertions at the same site within the coding region of BTK. Hum Mutat.

2005;25:324–5.

56. Wilund KR, Yi M, Campagna F, Arca M, Zuliani G, Fellin R, et al. Molecular

mechanisms of autosomal recessive hypercholesterolemia. Hum Mol Genet. 2002;11:3019–

30.

57. Vogt J, Bengesser K, Claes KB, Wimmer K, Mautner V-F, Minkelen R van, et al. SVA

retrotransposon insertion-associated deletion represents a novel mutational mechanism

underlying large genomic copy number changes with non-recurrent breakpoints. Genome

Biol. 2014;15:R80.

58. Laabs B-H, Klein C, Pozojevic J, Domingo A, Brüggemann N, Grütz K, et al. Identifying

genetic modifiers of age-associated penetrance in X-linked dystonia-parkinsonism. Nat

Commun. 2021;12:3216.

59. Westenberger A, Reyes CJ, Saranza G, Dobricic V, Hanssen H, Domingo A, et al. A

hexanucleotide repeat modifies expressivity of X□linked dystonia parkinsonism. Ann

Neurol. 2019;85:812–22.

60. Smith-Geater C, Hernandez SJ, Lim RG, Adam M, Wu J, Stocksdale JT, et al. Aberrant

Development Corrected in Adult-Onset Huntington's Disease iPSC-Derived Neuronal

Cultures via WNT Signaling Modulation. Stem Cell Rep. 2020;14:406–19.

61. Conforti P, Bocchi VD, Campus I, Scaramuzza L, Galimberti M, Lischetti T, et al. In

vitro-derived medium spiny neurons recapitulate human striatal development and complexity

at single-cell resolution. Cell Rep Methods. 2022;2:100367.

62. Henkel LM, Kankowski S, Moellenkamp TM, Smandzich NJ, Schwarz S, Fonzo AD, et

al. iPSC-Derived Striatal Medium Spiny Neurons from Patients with Multiple System

Atrophy Show Hypoexcitability and Elevated α-Synuclein Release. Cells. 2023;12:223.

63. Lexa M, Steflova P, Martinek T, Vorlickova M, Vyskot B, Kejnovsky E. Guanine

quadruplexes are formed by specific regions of human transposable elements. Bmc

Genomics. 2014;15:1032.

64. Broxson C, Beckett J, Tornaletti S. Transcription Arrest by a G Quadruplex Forming-

Trinucleotide Repeat Sequence from the Human c-myb Gene. Biochemistry. 2011;50:4162–

72.

65. Han H, Hurley LH, Salazar M. A DNA polymerase stop assay for G-quadruplex-

interactive compounds. Nucleic Acids Res. 1999;27:537–42.

A

B

TAF1 Exon Expression

C

TAF1 Exon Expression

A

Whole Blood (10K immunomes)

R-squared = 0.05936
p-value = 0.00001

B

Whole Blood (10K immunomes)

dR^2

−0.10   0.00   0.10

ZNF91 | ZNF93 | ZNF184 | ZNF212 | ZNF313 | ZNF488 | ZNF593 | ZNF600 | ZNF675 | ZNF680 | ZNF747 | TAF1

C

ZNF91 Expression Caudate Basal Ganglia

R-squared = 0.02145
p-value = 0.02157

D

ZNF91 Expression Putamen Basal Ganglia

R-squared = 0.02153
p-value = 0.03576

E

ZNF91 Expression Frontal Cortex

R-squared = 0.10354
p-value = 0.00000

F

ZNF91 Expression Cerebellum

R-squared = 0.04844
p-value = 0.00058