Diversification and recurrent adaptation of the synaptonemal complex in *Drosophila*

Rana Zakerzade[1,2,†], Ching-Ho Chang[3,†], Kamalakar Chatla[4], Ananya Krishnapura[4], Samuel P Appiah[4],

Jacki Zhang[4], Robert L Unckless[5], Justin P Blumenstiel[6], and Doris Bachtrog[4], Kevin H-C Wei[1,2,4,*]

[1] Department of Zoology, University of British Columbia, Vancouver, BC, Canada

[2] Life Sciences Institute, University of British Columbia, Vancouver BC, Canada

[3] Basic Sciences Division, Fred Hutch Cancer Center, Seattle, WA, USA

[4] Department of Integrative Biology, University of California Berkeley, Berkeley, CA, USA

[5] Department of Molecular Biosciences, University of Kansas, Lawrence, KS, USA

[6] Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA

* Corresponding author (wei.kevin@ubc.ca)

† Equal contributions

Summary

The synaptonemal complex (SC) is essential for meiotic recombination and faithful chromosome

segregation across eukaryotes, yet components of the SC are often poorly conserved. Here we show

that across the *Drosophila* phylogeny several SC genes have evolved under recurrent positive selection

resulting in orthologs that are barely recognizable. This is partly driven duplications repeatedly

generating paralogs that may have adopted novel germline functions, often in the testes. Unexpectedly,

while most SC genes are thought to be dispensable in the male germline where recombination is

absent in *Drosophila*, elevated testes expression appears to be the norm across the genus and likely

the ancestral state. The evolutionary lability of SC genes in *Drosophila* is likely a repeated source of

adaptive innovations in the germline.

ABSTRACT

The synaptonemal complex (SC) is a protein-rich structure essential for meiotic recombination and faithful chromosome segregation. Acting like a zipper to paired chromosomes during early prophase, the complex consists of central elements bilaterally tethered by the transverse filaments to the lateral elements anchored on either side to the homologous chromosome axes. Despite being found in most major eukaryotic taxa implying a deeply conserved evolutionary origin, several components of the complex exhibit unusually high rates of sequence turnover. This is puzzlingly exemplified by the SC of Drosophila, where the central elements and transverse filaments display no identifiable homologs outside of the genus. Here, we exhaustively examine the evolutionary history of the SC in *Drosophila* taking a comparative phylogenomic approach with high species density to circumvent obscured homology due to rapid sequence evolution. Contrasting starkly against other genes involved in meiotic chromosome pairing, SC significantly shows elevated rates of coding evolution due to a combination of relaxed constraint and recurrent, widespread positive selection. In particular, the central element *cona* and transverse filament *c(3)G* have diversified through tandem and retro-duplications, repeatedly generating paralogs that likely have novel germline functions. In a striking case of molecular convergence, *c(3)G* paralogs that independently arose in distant lineages evolved under positive selection to have convergent truncations to the protein termini and elevated testes expression. Surprisingly, the expression of SC genes in the germline is exceedingly prone to change suggesting recurrent regulatory evolution which, in many species, resulted in high testes expression even though *Drosophila* males are achiasmic. Overall, our study recapitulates the poor conservation of SC components, and further uncovers that the lack of conservation extends to other modalities including copy number, genomic locale, and germline regulation. Considering the elevated testes expression in many Drosophila species and the common ancestor, we suggest that the function of SC genes in the male germline, while still poorly understood, may be a prime target of constant evolutionary pressures driving repeated adaptations and innovations.

INTRODUCTION

Meiotic recombination, the exchange of non-sister, homologous chromosomes through physical crossovers, is an essential genetic mechanism universal to sexually reproducing eukaryotes. It allows for the shuffling of homologous alleles generating novel allelic combinations. This is necessary for maintaining nucleotide diversity and efficacy of selection; without it, chromosomes (like on the non-recombining, degenerate Y or W chromosomes) will irreversibly accumulate deleterious mutations ultimately leading populations to go extinct. At the cellular level, meiotic pairing, synapsis, and resolution of double strand breaks into crossovers are critical for stabilizing meiotic bivalents as failure is typically associated with skyrocketing aneuploidy rates. Therefore, recombination is a crucial genetic process that is necessary for reproductive fitness and long-term species survival.

Despite the critical functionality of recombination and the deep conservation across eukaryotes, aspects of this fundamental genetic mechanism are surprisingly prone to change. Recombination rate has been repeatedly shown to vary drastically between closely related species [1]. Adaptive explanations typically invoke changing environmental (e.g. temperature [2]) or genomic conditions (e.g. repeat content [3]) requiring commensurate shifts in recombination rate to maintain fitness optima [4,5]. Others have suggested intragenomic conflicts with selfish elements [6] or sexual conflict creating unstable equilibria for optimal fitness [7,8]. However, some have argued that changes in recombination rate have little impact on fitness and rate changes are the byproduct of selection on other aspects of the meiotic processes [9]. Several key findings supporting the adaptive interpretation come from *Drosophila* as multiple genes in the pathways necessary for recombination show signatures of rapid evolution due to positive selection [10–13]. Moreover, because recombination is absent in *Drosophila* males and the SC does not assemble during spermatogenesis [14], sexual antagonism due to sex-specific optima of crossover rates is unlikely the underlying driver of adaptive recombination evolution, at least in species with sex-specific achiasmy. Why recombination, an essential genetic mechanism, is prone to change and whether such changes are adaptive remain central questions in evolutionary genetics [15–17].

The paradox of poor conservation but crucial function is exemplified by the synaptonemal

complex (SC), a crucial machinery necessary for meiotic recombination in plants, animals, and major

lineages of fungi [18]. It is a protein complex that acts as zippers to tether homologs together along the

chromosome axes during meiotic prophase I and forms train track-like structures which have been

visualized under electron microscopy across eukaryotic taxa [19,20]. The SC is mirrored along a central

axis composed of central element proteins that are tethered by the transverse filaments to lateral

elements on two sides anchoring into chromatin (Figure 1A) [21]. This highly stereotypical configuration

is found in baker's yeast, mice, flies, and plants, indicative of an evolutionary ancient structure. Yet,

despite the deep evolutionary origin and functional necessity across wide eukaryotic domains, there are

many examples of unexpected exceptions. At the extreme are recombining species such as the fission

yeast that entirely forego the SC [22]. In another instance, the SC of *Caenorhabditis* has been

reconfigured such that the transverse filament – typically a single gene in most SCs – is composed of at

least four genes [23]. Therefore, parts of the SC appear to be curiously flexible in composition whereby

different analogous but perhaps non-homologous pieces can be recruited and replaced [24].

Consistent with this flexibility, sequences of SC components are often poorly conserved at

shorter evolutionary time scales [12,25]. In *Drosophila*, positive selection appears to repeatedly drive

the sequence evolution of the SC, which is composed of the central elements *corona* (*cona*) [26] and

corolla [27], the transverse filament *c(3)G* [28], and the lateral elements *orientation disruptor* (*ord*) [29]

and *c(2)M* [30]. Previously, orthologs of the central region components, *corolla*, *cona,* and *c(3)G,* could

not be found outside of the *Drosophila* genus [12] either reflecting divergence so extensive that

orthology is no longer recognizable, or novel acquisitions of SC components. Flexibility in SC

composition may explain how these molecular transitions are possible without major fitness impacts,

but cannot account for why SC genes appear to be evolving under recurrent adaptation. The recent

explosion of high quality *Drosophila* species genome assemblies [31–37] offer a unique opportunity to

understand the genetic and evolutionary mechanisms driving the strikingly rapid divergence of SC

genes. Here, we systematically revisit the evolution of the SC in *Drosophila* by examining the genomes

and transcriptomes of 48 species scattered across the entire *Drosophila* phylogeny, with dense

representation from three key species groups (*melanogaster*, *obscura*, and *immigrans*). In our

exhaustive analyses, we uncovered frequent duplications of several SC components generating

paralogs with likely novel functions, in addition to repeated sequence evolution under positive selection.

Further, we revealed unexpectedly high rates of expression divergence and regulatory turnover in not

just the ovary but also the male germline, where SC genes are thought to have no function. In fact,

testes-biased expression of SC genes appears to be the norm, and likely the ancestral state,

suggesting SC components have crucial function in male germline, despite the absence of male

recombination. Altogether our study revealed a highly dynamic evolutionary history with repeated bouts

of copy number, sequence, and regulatory evolution that contribute to the overall poor conservation of

SC genes. Further, the surprising transcriptional activity of SC genes in the male germline raises new

possibilities for functions of SC genes unrelated to recombination under repeated directional selection

in addition to their roles in chiasmate meiosis in the female germline.

RESULTS

Poor sequence conservation and frequent duplications of components of the SC

To identify *Drosophila* SC homologs we elected to focus on only species with high quality genome

assemblies with either available annotations and/or RNA-seq data (Supplementary table 1). In addition,

we strategically generated highly contiguous assemblies of two additional species (*D. hypocausta* and

D. niveifrons, belonging to the immigrans group; Supplementary table 2), and testes and ovaries RNA-

seq of eight species (*D. subobscura*, *D. arawakana*, *D. dunni*, *D. innubila*, *D. funebris*, *D. immigrans*, *D.*

hypocausta, and *D. niveifrons*) to either annotate previously unannotated genomes or to refine previous

annotations (Supplementary table 1). Altogether, we compiled a total of 47 species spanning the two

major arms of the *Drosophila* genus (the *Sophophora* and *Drosophila* subgenera), with three species

groups particularly well-represented (*melanogaster*, *obscura*, and *immigrans* groups) (Figure1B) and

the outgroup species *Scaptopdrosophila lebanonensis*.

Figure 1. Sequence conservation, or the lack thereof, of synaptonemal complex components across the *Drosophila* genus. A. Cartoon diagram of the Drosophila SC and its primary constituents. B. Phylogenetic relationships of the 48 species used in this study. Bold characters in the species name denote species shorthands. Species dense groups are labeled and boxed. C. Presence, copy number, and absence of SC components across the phylogeny. The number of blue circles indicated the copy number. D-F. Pairwise blast sequence alignments between orthologs from representative species across the genus. Alignments above the diagonal are from nucleotide blasts of the CDS sequences using blastn. Alignments below the diagonal are from protein blasts of the amino acid sequence using blastp. % blast identity is the length of blast alignments multiplied by the % sequence identity, summed across the gene. G. Distribution of pairwise blastp % identity across the genus for SC (colored) and genes (gray) involved in meiotic pairing during early prophase.

Using a multi-step reciprocal best blast hit approach, we sought to identify orthologs and

paralogs across the genus (Figure 1B and C; Material and methods). However, we found that the gene

structures are frequently malformed regardless of the source of the annotation (publicly available, or

our ones we generated). SC componets are often mis-annotated as truncated or chimeric gene

products or entirely missing in the annotation (Supplementary table 3; for examples see Supplementary

figure 1), likely due to the combination of exacerbating factors such as poor sequence conservation,

frequent presence of tandem duplicates, low RNA-seq reads, and in some cases assembly errors. To

ensure proper sequence alignments, we therefore curated all SC genes and manually re-annotated all

erroneous ones ensuring at a minimum, well-formed CDSs and intact full ORFs (Supplementary figure

1 and 2; see Materials and Methods). Note, because *cona* is a short gene with few exons, we hand-

annotated its orthologs in 8 additional species (see below).

For the lateral elements *c(2)M* and *ord*, sequence homology is somewhat preserved across the

genus, although homology is not always detectable between distant species using Blast (Figure 1D and

Supplementary Figure 3A). However, for the central region genes (*c3G*, *cona*, *corolla*), DNA sequence

homology quickly becomes unrecognizable outside of species groups, while weak protein homology is

only occasionally detectable (Figure 1E, Supplementary Figure 3A). Previously, Hemmer and

Blumentiel 2018 identified SC orthologs in a subset of fly species [12]. Increased species and better

annotations enabled us to identify orthologs previously missed (*cona* in *D. willistoni* and *corolla* in the

outgroup) and resolved discrepant homology relationships (*cona* in the *Drosophila* subgenus, see

below). To determine whether this lack of conservation is common to other genes involved in early

meiotic progression, we curated and identified the homologs of 8 genes necessary for meiotic

chromosome pairing (Figure 1F and Supplementary Figure 3B) across the *Drosophila* genus. The poor

sequence conservation of the SC starkly contrasts from these (Figure 1G): even the most conserved

component of the SC, *ord,* shows significantly poorer sequence homology compared to the least

conserved meiotic pairing gene, *sunn* (Figure 1G; Wilcoxon's Rank sum test p = 1.813e-05).

Furthermore, we revealed multiple independent duplication events, with *c(2)M* being the only

SC gene that remained single-copy across the genus. All SC paralogs were previously unaccounted

with the only exception being *ord* duplicates in *D. miranda*, which was identified to have rampantly

amplified creating over 20 copies (Supplementary figure 2) on the specie's unique neo-sex

chromosomes [38]. Of the poorly conserved components, *c(3)G* and *cona* in particular have recurrent

copy number changes, having more than two copies in eight and thirteen species, respectively, as the

result of independent duplications in at least seven lineages each. This propensity to duplicate is

epitomized by the five *c(3)G* in *D. arawakana* and six *cona* copies in *D. obscura*.

History of repeated duplication and loss of paralogs

Based on protein trees of the SC components, we find that the current copy number distributions reflect

at least 3, 7, 1, and 7 independent duplications of *ord*, *c(3)G*, *corolla*, and *cona*, respectively. The

majority of paralogs are recent species-specific duplications resulting in short branch lengths (Figure 2A

and 3A, and supplementary figures 3). The genomic locations of the copies further reveal that tandem

duplications account for the majority of the observed copies. For the transverse filament *c(3)G*, four of

the seven duplication events are tandems (Figure 2A,C-D), three of which (including the five copies in

D. arawakana) are recent and species-specific. In the lineage leading up to *D. athabasca* and *affinis*, an

older tandem duplication generated duplicates of *c(3)G* and the neighboring gene *pon*, (Figure 2D); one

of the copies which we designated *c(3)G2* is shorter, while showing poorer conservation and longer

branch lengths between the orthologs, suggestive of functional divergence (see below). Similarly, for

cona, three of the seven duplication events are tandems, including the 6 copies in *D. obscura* (Figure

2B, Supplementary Figure 5).

Both *c(3)G* and *cona* experienced several instances of likely retroduplications. *cona* offers two

clear examples of old events in the common ancestor of the *serrata* and *nasuta* subgroups leading to

paralogs shared across all species in the subgroups (Figure 2B). *c(3)G*'s duplication history appears

more convoluted but offers unique insight into its dynamic evolution. There are three non-tandem

duplicates of *c(3)G* found in *D. kikkawai*, *triauraria*, and *innubila* (Figure 2A); the resulting paralogs

found on different chromosomal regions (Figure 2A, Supplementary figure 6A) have minimal

Figure 2. Complex evolution history of synaptonemal complex. A and B. Gene trees of *c(3)G* and *cona* orthologs and paralogs, respectively; nodes with poor bootstrap support are demarcated by circles. Duplicates are labeled in blue and tandem duplicates have asterisks. For *cona*, some branches were adjusted (red) to align with the species tree. For other SC components see Supplementary Figure 4. Color blocks to the right represent different Muller elements on which the genes reside. Unless connected by lines, separate blocks of the same color represent different locations on the same element. C-E. Dotplots showing synteny of genomic regions surrounding *c(3)G* between sister species and/or paralogs. The color of the dots represent the % sequence identity from blastn alignments with darker red reflecting higher identity. In the gene tracks of the displayed regions, *c(3)G* is in purple and other neighboring genes in blue. For E, dotplots depict the local homology of *c(3)G1* and retroduplicate *c(3)G2* (bottom), and additional regions with truncated *c(3)G* (top) in *D. kikkawai*.

neighboring homology to the original (Figure 2E, Supplementary figure 6B) suggesting retroduplication

events instead of larger scale duplications. In the latter two species, the duplications are older events

evidenced by their phylogenetic placements with long branches separating the paralogs, compared to

the species-specific duplication in *D. kikkawai*. For *D. triauraria*, the duplication creating *c(3)G2*

predated the split of the serrata species subgroup, but is no longer found in the derived lineages,

indicating subsequent loss of the paralog. For *D. innubila,* the paralogs are found 180kb apart on the X

chromosome (Supplementary Figure 5B), and the phylogeny suggests that the duplication occurred

after the split from *D. funebris*, although with low bootstrap support (Figure 2A). Interestingly, synteny

information suggests this is not the true relationship; while one of the *c(3)G* copies is found in the same

syntenic block shared with *D. funebris, arawakana*, and *dunni*, the other is in a separate synteny block

shared will nearly all other species in the *Drosophila* subgenus and thus the ancestral copy (Figure 2A,

Supplemental Figure 6C). This synteny pattern is therefore more parsimonious with an old duplication

in the last common ancestor of the dunni, quinaria, and funebris species groups with the original copy

lost in species other than *D. innubila*. Non-allelic gene conversion subsequently homogenized the

duplicates in *D. innubila*, obscuring the true phylogenetic relationship.

In addition to the one *c(3)G* retroduplicate in *D. kikkawai*, we curiously identified numerous loci

across the genome as 5' truncated homologs, none of which were annotated or have RNA-seq reads

mapping (Figure 2E). These truncated and nonfunctional duplicates, along with the two loss events

mentioned, raise the possibility that *c(3)G* experienced not only repeated duplications through

transpositions, but also repeated pseudogenization events. A similar pattern of nonfunctional duplicates

is also observed with *corolla* in *D. arawakana*; despite only one full length *corolla*, there are four

adjacent tandem copies that lack the 5' exon and therefore likely non-functioning (Supplementary

figures 5). Further examining the syntenic relationships of the SC homologs, we find that while the

lateral elements have maintained the same local microsynteny showing a lack of gene movement, the

central region genes have repeatedly relocated to different chromosomes, or different locations on the

same chromosome (Figure 2A, Supplementary figure 6A). The X is likely the ancestral home for all

three but we documented eleven, sex, and five inter- and intrachromosomal movements for *c(3)G*,

corolla, and *cona*, respectively. Such recurrent movements likely through transpositions are unusual for

flies as chromosomal gene content and microsyteny are largely stable while broad chromosome-scale

synteny is scrambled by large scale rearrangements [39]. Curiously, one relatively recent relocation

occurred in the common ancestor of the *pseudoobscura* species, moving *c(3)G* from Muller B to an

euchromatic repeat block on Muller E (Supplementary figure 6), suggesting that such movements may

be mediated by the instability of repetitive sequences, perhaps piggybacking off of transpositions of

transposable elements. Since most of these movements no longer have extant paralogs, corresponding

pseudogenization events were likely common. Therefore, even though most observable paralogs are

young tandem duplicates, retroduplications and pseudogenization events have frequently occurred for

c(3)G, cona, and even *corolla* which has few remnants of duplicates, thus accounting for the existence

of many recent and species-specific copies in different locations but fewer old, shared duplicates.

***c(3)G* and *cona* paralogs likely have novel germline functions**

Much like other transverse filaments, *C(3)G* has an extensive coiled-coil domain flanked by globular

domains at the N- and C- termini connecting the central and lateral elements, respectively [18,40].

Despite the poor sequence conservation, we find that this canonical structure is conserved across the

genus based on AlphaFold protein prediction (Figure 3A) [41,42] and coiled-coil predictions

(Supplementary figure 9) [43]. This unique evolutionary property of structural but not sequence

conservation is also observed by Kursel et al. in *Caenorhabditis*, whereby central element genes have

conserved coiled-coil domains and near invariant protein lengths, but neutrally evolving sequences [25].

In *D. athabasca*, *affinis*, and *triauraria*, while *c(3)G1*s produce longer proteins (690, 692, 830 AAs

respectively) predicted to have the canonical structure, the paralogs *c(3)G2s* all produce

notably shorter proteins (361, 395, and 319 AAs, respectively) with the flanking globular domains

truncated, if not entirely absent, suggesting they no longer function as transverse filaments that can

tether the SC (Figure 3B). Curiously, these paralogs are highly expressed in the testes but lowly

expressed in the ovaries (Figure 2C-E), incongruent with the expectation of female meiotic function.

Despite being independent duplications in lineages separated by over 25 million years, *c(3)G2* in *D.*

triauraria and *D. athabasca/affinis* display remarkably similar structural and regulatory evolution,

revealing molecular convergence for likely male germline function. In a single nuclei RNA-seq dataset

Figure 3. Functional and structural evolution of *c(3)G* and *cona*. A-B. AlphaFold structural prediction of full length c(3)G (A) and diverged paralogs (B). Color represents the confidence of structural prediction. C-E. Gene structure and gonad expression of *c(3)G paralogs* (purple genes) in *D. triauraria* (C), *affinis* (D) and *athabasca* (E). F. Alphafold of distant *cona* orthologs in Drosophila and outgroup species. G. Gene structure and gonad expression of *cona* paralogs in *D. simulans* and *sechelia*. Annotated lncRNA of gene with homology to cona is in orange in the gene tracks. G. Same as F but for D. *serrata*.

of *D. affinis* testes, we further find that while *c(3)G1* and *c(3)G2* are both testis-expressed, they are

most active in different cell populations (Supplementary Figure 10), evidence that the two have

functionally diverged.

Similar to *c(3)G*, *cona* has maintained the same conserved tri-coil structure (Figure 3F), despite

poor protein homology. Curiously, several *cona* duplicates similarly show properties that deviate from

its characterized function in SC formation during female meiosis. At least two recent duplication events

occurred within the *simulans* clade generating two upstream paralogs, one ancestral to the three

simulans species while another found only in *D. sechelia* (Figure 2B and 3G). The *sechelia*-specific

duplicate generates a complete but short ORF and likely protein coding, but the shared paralog only

shows homology at the 3', lacks a complete ORF, and is annotated as a long non-coding RNA (Figure

3G). To evaluate whether this paralog, which we named *cona-like*, is transcriptionally active or

pseudogenized, we examined RNA-seq data, and found high expression in the testes and males but

low-to-no expression in ovaries or females across all *simulans* species (Figure 3G), strongly suggesting

testes function as a lncRNA. Adding to the intrigue, this is not the only instance of a *cona* paralog

generating lncRNA. In the *serrata* group, the retroduplicate, *cona2*, is shared across the species

(Figure 2B), but only in *D. serrata* does it generate a lncRNA (Figure 2H). Unlike *cona-like* in the

simulans clade, this paralog has a well-formed ORF and is expressed in both sexes, suggesting high

protein coding potential. However, the lncRNA is anti-sense as confirmed with strand-specific RNA-seq

(Supplementary figure 11) and includes additional flanking sequences that only show expression in

males. *cona2* likely generates a functional protein in females and ovaries but was incorporated in the

anti-sense direction into lncRNA production in the testes of *D. serrata*. Altogether, these results suggest

that both *c(3)G* and *cona* paralogs have repeatedly adopted germline functions in the testes unrelated

to SC formation.

coronetta (_conta_) is an ancient testes-expressed paralog of _cona_

Previously, Hemmer and Blumenstiel identified *cona* homologs in the *Drosophila* subgenus that were

highly diverged from *Sophophora cona* [12]. One of the proteins they identified through reciprocal best

blastp hit was *GJ20698* in *D. virilis,* a gene producing a short peptide of 109 AA. We were able to find

the orthologs of *GJ20698* across the *Drosophila* subgenus as well as the outgroup, but none of them

were reciprocal best hits to *sophophora cona* – in fact, they have no identifiable *sophophara* homologs

at all. Instead, increased density of species enabled us to correctly identify *D. virilis's GJ16397* (the *D.*

virilis's 2nd best hit to *sophophora cona*) which has orthologs across the *Drosophila* subgenus and *S.*

lebanonensis that are reciprocals best hits with *sophophora cona* (Figure 1F, 2B). The gene tree affirms

Figure 4. Expression and structure of _conta_, an ancient duplicate of _cona_. A. Unrooted protein tree of *cona* and its old duplicate *conta*; major lineages are labeled. B. Annotation and germline expression of *conta* (purple) in *D. virilis* and the outgroup *S. lebanonensis*. C. Alphafold prediction of conta in representative species.

that *sophophora cona* is more closely related to *GJ16397,* which we conclude to be the true *cona*

ortholog (and used for all analyses). *GJ20698*, which we named *coronetta* (*conta*), appears to be a

distant paralog, and, given its presence in the outgroup, emerged prior to the last common ancestor of

Drosophila and *Scaptodrososphila*. Unlike *cona*, *conta* sequence is conserved (Figure 4A), found in the

same syntenic region (in the intron of the gene *teiresias;* Supplementary figure 12), and highly

expressed in the testes but not ovaries (Figure 4B). Structural prediction of *CONTA* reveals distinctly

shorter coiled structures (Figure 4C). Given the absence of *conta*'s ortholog in *sophophora* – not even

in the same syntenic location (Supplementary figure 10) – it has likely been lost, further underscoring

the propensity for SC paralogs to participate in germline function that may be evolutionarily fleeting.

SC proteins are evolving under recurrent positive selection and accelerated by repeated

duplications

Poor sequence homology can result from relaxed constraint due to reduced negative selection or

adaptive protein evolution due to positive selection. Previously, Kursel et al. reported that the elevated

rate of SC protein evolution in *Caenohabditis* reflect relaxed sequence constraint while the coiled-coil

domains and protein lengths are both highly conserved [25]. However, Hemmer and Blumenstiel

identified both elevated rates of protein evolution and signatures of positive selection for *Drosophila* SC

genes [12]. We reassessed the rates of protein evolution by estimating the branch-specific ratio of

nonsynonymous and synonymous rate of protein evolution, represented by omega. Values approaching

0 indicate negative selection while values close to or greater than 1 indicate relaxed constraint and

positive selection, respectively [44]. Notedly, gene-wide omega values, which are predominantly

negative, are typically the composite of several modes of evolution as different residues and/or

domains of the protein can be under different forms and levels of selection [45,46]. Our curated,

species dense SC orthologs and paralogs enabled not only branch-specific, gene-wide estimates

(Figure 5A-B and Supplementary figure 12), but also detection of significant positive selection occurring

only at portions of the protein coding sequence with the Hyphy package [47].

Figure 5. Rate of protein evolution and signatures of positive selection of SC components. A-B. Along the gene trees, branches are colored by their branch-specific rates of protein evolution (omega) with warmer colors representing higher omega. Branches inferred to have significant positive selection in part of the protein are labeled with asterisks (* = p < 0.05 and ** = p < 0.001). See supplementary figure 12 for the remaining SC genes. C. Distribution of branch-specific omega values for the different SC components and genes required for pairing during early prophase I. P-values are from pairwise Wilcoxon's rank sum tests comparing between the lateral and central region genes. For non-SC genes, ** indicates significant differences (p < 0.005) when compared to all SC genes; * indicates significance comparisons except for *ord*. Ratios below SC genes tallies the number of branches showing omega > 1 or significance over total number of branches. D. D_N/D_S of the SC genes compared to the genome-wide distribution estimated of by PAML. E, F. Comparison of the distribution of omega values on branches following duplications (red) versus unduplicated branches (gray). Vertical dotted lines mark the median omega values for duplicated (red) and unduplicated branches (black). P-values were inferred from 1-tailed Wilcoxon's rank sum test.

295   For the more conserved, lateral elements, despite the better preserved sequence homology and

296   predominantly negative gene-wide omega (Figure 5A and Supplementary figure 13), multiple branches

297   still show signatures of positive selection; 28 out of 85 and 38 out of 99 branches display either positive

298   gene-wide omega or significant site-specific selection for *c(2)M* and *ord*, respectively. For the poorly

299   conserved central elements (Figure 5B & Supplementary figure 12), not only do they have significantly

higher omega than the lateral elements (p < 0.00005, pairwise Wilcoxon's rank sum tests; Figure 5C)

indicative of higher rate of protein evolution, over 40% of the branches show either positive omega or

significant signatures of positive selection at parts of the protein. Note, we suspect the omega values

for *c(3)G*, *corolla*, and *cona* may be underestimated due to poor amino acid alignments across much of

the protein. Collectively, the SC shows significantly elevated rates of protein evolution compared to the

other genes involved in meiotic pairing; *sunn* is the only exception (Figure 5C) with omega values

similar to *ord*, but nonetheless significantly lower than the other SC genes. We further used PAML to

infer the rate of protein evolution within the three well represented species groups, and we, again,

consistently find evidence of positive selection. Restricting to a subset of 6 *melanogaster* group species

with decent alignments, we compared the omega values of SC genes to the genome-wide distribution

and found that the SC genes fall between 86.9 and 99.8 percentiles, and are significantly

overrepresented with elevated values despite only 5 genes (Figure 5D; p = 0.000318 Wilcoxon's Rank

Sum Test). Altogether these results demonstrate that all components of the SC have a history of

recurrent adaptive evolution with the central region genes under frequent and repeated positive

selection.

Copy number expansions can allow genes to diversify leading to new functions or subdivision of

existing functions among the paralogs. Both of these scenarios are associated with elevated omega,

either from relaxed functional constraint or positive selection for novel function. To test whether the

recurrent duplications of SC components lead to elevated rates following such functional diversification,

we examined branches after duplications for *c(3)G* and *cona.* We found significantly elevated rates of

protein evolution on such branches with median omega of 0.925 and 0.914, respectively, both

significantly higher than single copy branches (Figure 5E and F; p = 0.000031 and 0.03788,

respectively, one-tailed Wilcoxon's rank sum test). In particular, the *c(3)G2* paralogs with high testes

expression in *D. triauraria*, *athabasca*, and *affinis* (Figure 3C-E) all show clear signatures of adaptive

evolution post duplication (Figure 5B). Although the terminal branches of the latter two species appear

neutrally evolving, the parent branch displays highly elevated and significant omega, indicating strong

positive selection in the common ancestor post duplication. Notably, the original copy, *c(3)G1,* in these

cases also display signatures of positive selection suggesting that the duplications were followed by

functional diversification to both. Moreover, other *c(3)G* duplicates, even the recent ones, show

signatures of adaptive evolution including those in *D. erecta* and *kikkawai*. Similar for the *cona*

duplicates, nearly all the branches of *cona2* in the serrata group show signatures of positive selection,

which accounts for the overall longer branch lengths than those of *cona1* within the same group

(Supplementary figure 12). These repeated signatures of positive selection after duplications strongly

argue for adaptive functional diversification, acting to further accelerate the already rapid evolution of

the SC.

Poor regulatory conservation of SC genes in both female and male germlines

With the exception of *ord* which maintains sister chromatid cohesion in the male germline [48,49] the

Drosophila SC proteins are thought to primarily function in the ovaries and dispensable in the testes

because males are achiasmic and mutant males do not show obvious meiotic or fertility defects [50–

53]. However, a recent elegant study of the male germline by Rubin et al. found that the progression of

pre-meiotic chromosome pairing is slower in *cona* and *c(3)G* mutants, revealing putative male germline

function despite the absence of SC assembly [54]. Inspired by these results, we examined the

expression of SC genes in ovaries and testes RNA-seq datasets from a subset of 38 species (Figure

6A), 14 of which we generated and 26 curated from publicly available datasets (Supplementary table

3). Other than *ord* which is testes-biased across all species, the expression of SC genes – even single

copy ones – is highly unstable contrary to the naive expectation of high ovary and low testes

expression. testes-biased expression appears to be the norm rather than the exception as SC

components are more highly expressed in the testes in over 70% of the species. We further examined

available testes single cell RNA-seq data of *D. miranda* [55]*,* a species with high testes expression of

c(3)G, and find expression concentrated in the pre-meiotic cell types such as germline stem cells and

spermatogonia (Supplementary Figure 14), reminiscent of their reported pre-meiotic activity in *D.*

melanogaster. Thus, even though they are primarily known for their role in female meiosis, SC genes

Figure 6. Variable expression of SC components in the testes and ovaries. Ovaries (red) and testes (blue) expression (transcript per million in log scale) of SC genes across 38 species. Tandem duplicates with similar expression are collapsed into one. Other duplicates are labeled. Gray and white vertical bars differentiate between species.

can also be highly active in testes, arguing for critical male germline function. This is further supported

by the heavily testes-biased expression of *corolla*, *c(2)M*, and *ord*, in the outgroup *S. lebanonensis*,

suggesting that testes function is ancestral.

The striking lability of germline SC expression is particularly evident from several closely related

species pairs whereby expression rapidly switches between testes- and ovaries-bias. For instance,

c(2)M is ovaries-biased in *D. simulans* but testes-biased in the sister species *D. sechelia and*

mauritania (Figure 6). Similar rapid expression change of *c(2)M* is also observed in *D.*

pseudoobscura/*D. persimilis* and *D. athatbasca*/*D. affinis* sister pairs, suggesting such regulatory

evolution occurs regularly. As the RNA-seq datasets came from different studies, we tested the

possibility that the expression differences resulted from external factors like rearing conditions and

examined SC expression from RNA-seq of studies where flies were reared in different environmental

conditions. In the datasets examined [56–58], SC expression shows minimal change in different rearing

temperature (Supplementary Figure 15). Additionally, drastic expression divergence of the germlines

can be observed in RNA-seq datasets we generated of closely related species reared in the same

controlled laboratory condition, including those in the *immigrans* group and the sister species *D.*

arawakana and *D. dunni*. Therefore, highly variable expression of the SC across the genus likely

reflects bona-fide regulatory divergence. Other curious patterns include elevated testes but low-to-no

ovary expression such as *c(3)G* in *D. obscura* and *D. hypocausta*. Most puzzling, there are multiple

lineages where components of the SC show little-to-no expression in both gonads, such as *D. kikkawai*

with little gonad expression of all components. The most extensive regulatory stability appears to be the

nasuta subgroup where the expression of SC components are consistently low across the species,

especially for *c(3)G* and *corolla*. These drastic differences between species cannot be simply driven by

differential tissue contribution in the dissections, as the expression of the SC components are poorly

correlated, with many instances of high expressions of one component and low expression of others

within the same sample. While it is tempting to infer function outside of the germline based on the

absence of expression, even *D. melanogaster* ovaries show low expression of some of the SC

components like *ord* and *cona* arguing that normal SC function may not require large transcript counts.

Nevertheless, these expression patterns reveal that despite the essential roles in regulating crossovers,

germline transcriptional regulation of many SC genes is highly labile and constantly evolving, echoing

their protein and copy number evolution.

DISCUSSION

Trials and tribulations of gene ortholog and CDS searches from genome releases

The species-dense examination of SC evolution was made possible by the large amounts of *Drosophila*

genomes that have been recently published. One of the promises of the ever-increasing number of

genomes is to enable deep and broad investigation of the molecular evolution of genes and pathways.

Naturally, analyses of genic evolution typically require alignments of full length CDSs, which are

distilled from genome annotations. Since good gene structure inference requires additional data such

as RNA-seq, only ~1/3 of the available *Drosophila* genomes have available annotations (including

those we generated here), which were the focus of this study. Notably, recent studies were able to take

advantage of more genomes by focusing on small genes such as protamines [59], sex peptide, and sex

peptide receptor [60], which either have no intron or have simple gene structure, enabling protein

construction without transcript information from RNA-seq. Even when we relied on annotations with

RNA-seq support to identify gene orthologs from annotated genomes using reciprocal best blast hits,

we were flummoxed by the regularity of annotation errors (Supplementary table 3). Rapidly evolving

genes may be particularly prone to such errors due to obscured sequence homology and, in the case of

the SC genes, propensity to duplicate. Without careful manual curation, cross referencing with

homology and RNA-seq data, annotations errors taken at face value would be falsely construed as

evolutionary changes. Below, we discuss some of the pitfalls of identifying orthologs and paralogs from

annotated genomes.

We find that NCBI annotations are generally more reliable than our in-house annotations with

maker but has its own sets of issues, the biggest of which is the lack of access to the proprietary

pipeline. Problems common to all pipelines include improper splice junctions producing truncated ORFs

with premature stops and fused genes creating genes chimeras d – tandem duplicates were particularly

problematic for this as they cause extensive mapping errors of the RNA-seq reads between

neighboring copies. These errors will cause protein sequences with excess length differences and can

be detected as long stretches lacking homology between orthologs in multiple sequence alignments.

The more insidious issues stem from genome assembly errors most often occurring as indels at

homopolymers tracks, an issue common to Pacbio and Nanopore reads [61,62]. While released

genomes should have been polished with Illumina reads to reduce such errors [61], we repeatedly

identified annotation errors caused by frameshift indels at homopolymer tracks, the worst offender of

which is a Refseq genome, released on and annotated by NCBI, rife with such indels (Supplementary

Figure 16). In our in-house annotations with maker, such indels can cause the expected frameshifts

and truncated ORFs or unexpected splice junctions with no RNA-seq support. Puzzlingly, NCBI

annotations add nonsensically short (<5 bp) introns around such frameshifts to maintain the ORF

(Supplementary Figure 15). Because this creates near complete proteins but with small numbers of

missing internal amino acids, it is only detectable by looking at variant calls in mapped Illumina reads.

Careful examination of CDSs and protein sequences extracted from genome assemblies is therefore

necessary to avoid erroneously calling species-specific substitutions, indels, nonsynonmous changes,

psuedogenes, and splice variants.

Mechanisms underlying recurrent duplications and rapid evolution of the SC

Our exhaustive survey for SC orthologs resulted in the surprising identification of many

duplicates across *Drosophila*. Duplication is an important mechanism to diversify gene function as the

resulting paralogs can either evolve novelty or compartmentalize existing functions, with reduced

selective constraint on the gene as a single copy. Indeed, we find elevated rates of protein evolution

following duplications indicative of both relaxed constraint and adaptive evolution. Many of the

duplicates are young and species-specific showing no evidence of expression differences, and

therefore unlikely to have diverged in function. For some of the older *c(3)G* and *cona* paralogs, we

observed repeated and independent acquisition of distinct activities surprisingly in the male germline,

such as testes-specific expressions and incorporation into lncRNA production. In the case of the

tandem duplication of c(3)G in the ancestor of *D. affinis* and *D. athabasca* (Figure 3D-E), the two copies

show clear evidence of positive selection driving diversified function evidenced by important structural

differences (i.e. loss of terminal globular dominas) in the protein and expression in different testes cell

types. While divergence in expression domains is consistent with subfunctionalization, the adaptive

protein evolution strongly argues functional novelty. Notably, a parallel duplication *of c(3)G* occurred in

D. triauraria, which also evolved into a testes-expressed paralog lacking the globular domains that

interact with lateral elements. This surprising molecular convergence raises the intriguing possibility of

a common testes process acting to repeatedly drive the rapid evolution of SC proteins.

For reasons still unclear, the testes appear to be a unique regulatory environment, producing

the largest repertoire of lncRNAs [63,64] and *de novo* genes [65–68], both of which can have critical

roles in spermatogenesis. Considering our unexpected finding of frequent testes expression, we

speculate that testes activity of SC genes – even single copy ones – may provide the opportunities to

generate paralogs that can diversify. This path to functional novelty or diversification in the testes may

be further facilitated by the elevated retrotransposon activity in spermatocytes [69] which can

conceivably provide the necessary machineries for both retro- and tandem duplications. Our

observation that *c(3)G* moved into a satellite-block in the lineages leading up to the pseudoobscura

subpseices (Supplementary figure 8) lends support to this possibility.

However, despite many paralogs, some of which with likely novel functions, pseudogenization

of duplicates also appears common. We have identified multiple instances where extant duplicates in

one species have been lost in neighboring lineages (e.g. *c(3)G2* in *D. triauraria*, and *D. innubila*, and

conta). We also found several examples of remnants of SC duplicates, including truncated copies of

c(3)G and *corolla*. Such dynamic copy number changes raise a perplexing conundrum: why are *cona*

and *c(3)G* prone to produce duplicates under positive selection only for the duplicates to end up as

pseudogenes. One clue may come from the exceptional duplication of *ord* in *D. miranda*, where neo-X-

and neo-Y-linked gametologs, along with other meiosis-related genes, have massively amplified in

tandem [38]. The amplification is hypothesized to be the result of dosage-sensitive sex-ratio meiotic

drivers precipitating an arms race for gene copy numbers on sex chromosomes [38,55]. Similar

dynamics of repeated copy number evolution is also observed for sex ratio drivers and suppressors that

manipulate DNA packaging in X- and Y- bearing sperms [59,70,71]. In such models of meiotic conflicts,

temporary/young duplications may act to increase the gene dosage to either induce selfish

transmission (such as biased sex ratio) or to act as suppressors of drive that restore fitness reduction

associated with non-mendelian transmission. Once the conflict is resolved, drivers and suppressors

may pseudogenize and degenerate as they no longer impart fitness benefits.

Since the intricate orchestration of chromosome movement is necessary for recombination and

faithful disjunction, we find the regulatory variability of SC genes in the ovaries to be perplexing,

particularly in species where SC genes have little-to-no ovary expression. The repeated relocation to of

c(3)G, *cona*, and *corolla* to different chromosomal regions may confer some degree of regulatory

changes, but even c(2)M, which has remained in the same syntenic location, show variable expression.

It is tempting to speculate that the low ovary expression in multiples species correspond to repeated

loss of meiotic function, but we find other possibilities more likely. If SC proteins have long half-lives,

minimal transcript production may be sufficient to support robust SC assembly. However, this possibility

cannot address why species evolved to have drastically different expression profiles. While we

examined available germline expression datasets under different environmental conditions and found

minimal changes in SC expression, we cannot fully rule out extrinsic factors driving to the expression

lability of SC genes. Indeed, recombination rate is sensitive to environmental conditions and life history

such as nutrition [72,73], temperature [2,74,75] and stresses [76], and age [77,78], and species can

differ in their physiological responses that subsequently regulate SC activity. Such mechanisms can be

beneficial in ensuring optimal recombination rates to modulate the amount of genotype diversity in the

offspring [76] or proper progression of meiosis in suboptimal cellular conditions like extreme

temperatures [9]. However, we note that the species we generated germline RNA-seq for were raised

under common laboratory conditions and were at standard, reproductively active age, we still observed

drastic differences in SC regulation likely reflecting true regulatory divergence in both the male and

female germlines.

Our analyses of the protein coding evolution demonstrate that SC genes have an extensive

history of recurrent adaptation with the central region genes being frequent targets of positive selection.

This appears unique to SC genes compared to others necessary for chromosomal progression during

early prophase which are all conserved, other than *sunn*. While the elevated rates of evolution are

partly driven by paralogs diversifying in germline function, orthologs without duplicates also show

signatures of positive selection across the gene trees. This contrasts from the SC genes in

Caenorhabditis, the protein sequence of which are evolving neutrally [25]. Further, while *C(3)G* appears

structurally conserved, the lengths of the proteins are far more variable with a coefficient of length

variation 5 times higher than that of worms (0.17 vs. ~0.03 [25]). While this could reflect repeated

adaptation in *Drosophila* female meiosis and meiotic recombination, our findings that SC genes

frequently function in testes where it is also ancestrally highly expressed compel us to consider

additional avenues under recurrent positive selection, especially since spermatogenesis is fruitful

grounds for meiotic conflicts, sexual selection, and molecular innovations. Pleiotropy tends to increase

molecular constraint [79]. However, given the sequence tolerance of the SC, dual function of

Drosophila SC genes in both oogenesis and spermatogensis may instead predicate a unique scenario

where positive selection in the latter has little pleiotropic impact on the former. Dissecting the function of

SC genes in the *Drosophila* male germline, which is ironically achiasmate, will therefore be critical to

understanding the diversity and evolution of meiotic recombination.

MATERIALS AND METHODS

High molecular weight DNA extraction and genome assembly

To assemble the genomes of *D. hypocausta* and *D. neivefrons*, we followed the Nanopore long read

sequencing pipeline from [33,80]. In short, high molecular weight DNA was extracted using the Qiagen

Blood & Cell Culture DNA Midi Kit from ~100 males of *D. hypocausta* strain 15115-1871.04 from the

National Drosophila Species Stock Center and ~50 females of *D. niveifrons* strain LAE-276 from the

Kyorin Drosophila Species Stock Center. DNA strands were hand spooled after precipitation, followed

by gentle washing with supplied buffers.

RNA-seq preparation and analyses

5 pairs of ovaries and testes were dissected from adult females and males and stored in Trizol at -80

degrees, followed by standard RNA extraction. RNA-seq libraries were generated using either the

NEBNext RNA Library Prep Kit for Illumina with the Stranded and mRNA isolation Modules or the

Illumina Truseq Stranded mRNA Library Prep kit. After quality check with the Fragment Analyzer at

QB3-Berkeley, the libraries were sequenced by Novogene. We aligned the reads (both ones we

generated and downloaded from SRA) using hisat2 (v2.2.1) [81] on either pair-end or single-end mode

to their respective genomes with the –dta flag to allow for downstream transcriptome assembly. After

sorting the aligned reads with samtools (v1.5) [82], we used the featureCount (v2.0.3) in the Subread

package [83] for read-counting over genes, allowing for non-uniquely mapped reads (-M flag). Read

count tables were processed and analyzed in R (v4.2.2) and Rstudio (v2022.12.0). For gene

expression analyses, we normalized the read counts across samples by converting them to transcript

per million (TPM) [84]. For species where we needed to do de novo gene annotation, we used stringtie

v2.1.6 [85] on default for genome-guided transcript assembly.

Gene annotation and manual curation of gene structures

For species that required gene annotation, we ran three rounds of maker [86]. For evidence-based ab

initio gene prediction in the first round, we supplied the transcript assembly from stringtie, de novo

repeat index from RepeatModeler2 [87], transcript sequence from closely related-species and protein

sequence data from D. melanogaster and D. virilis downloaded from FlyBase. The maker results from

round one were used to train the species specific gene model using SNAP [88]. The resulting

snap.hmm file was fed back into maker for round 2. We iterated this process again, refining the gene

models for a 3rd round of maker.

For malformed or missing annotations, we first visualized the gene structures and RNA-seq reads

mapping around them using IGV (v2.16.0) [89]. Additionally, we manually defined the region of the

genome showing gene homology by blastn-ing the well-formed ortholog from a closely related species

to the genome. The combination of RNA-seq reads mapping the blast-hit boundaries provided evidence

to correct erroneous exon-intron injunctions, truncated annotations, chimeric gene structures, and

absent annotations. To update the annotations file (.gff file), we used the genome browser

GenomeView (v2250) [90] to manually edit or add the gene structures including mRNAs, exons, CDSs.

All edited genes have at least full open reading frames, although 5' and 3' UTRs are missing. All

manually annotated features were marked by the flag "hand" in the gffs. The updated .gff is then

exported and sorted using GFF3sort [91], and transcript sequences are retrieved using gffread (0.9.12)

[92]. In several instances, we noticed assembly errors leading to malformed genes. One was *ord* in D.

nasuta which had a stretch of N's within the gene body indicating scaffolding points. The other was in

D. *neivifrons* where c(3)G was annotated as two fragments. This was due to a deletion of a single

nucleotide in the genome causing a shifted reading frame which led to malformed annotations. The

deletion was revealed by RNA-seq read mapping, whereby all reads showed a one basepair insertion.

We fused the fragmented annotations into one, and corrected the transcript sequence to rectify the

erroneous deletion. Lastly, we initially could not identify *corolla* in the primary NCBI genome assembly

of *D. funebris* strongly suggesting gene is loss; however, we were subsequently able to identify it in an

unplaced repeat-rich contig in a separate assembly. For species with malformed annotations that had a

closely related species with well-formed orthologs, we used the program Liftoff [93] to convert the

annotation from one genome to the other.

Homolog search with reciprocal best blast hits of transcripts and/or coding sequences

We first used blastn to identify homologous transcripts between species pairs, and in cases where no

clear best hit was identified, we then used tblastn to identify homologous transcript with protein

sequences. The absence of protein hits with tblastn is then followed by blasting to the genome, to

ensure the absence of an ortholog in the transcript sequences is not merely the result of absent

annotation.

To identify orthologs and paralogs using a reciprocal best blast hit strategy, we reciprocally blastn-ed

transcript sequences from species pairs using the commands:

blastn -task blastn -query species1.transcripts -db species2.transcripts -outfmt "6 qseqid sseqid pident

length qlen slen mismatch gapopen qstart qend sstart send evalue bitscore" -evalue 1

blastn -task blastn -query species2.transcripts -db species1.transcripts -outfmt "6 qseqid sseqid pident

length qlen slen mismatch gapopen qstart qend sstart send evalue bitscore" -evalue 1.

For publicly available genomes with annotation files, we generated the transcript sequences using

gffread, otherwise we used transcript sequences generated by maker. We then used grep to identify

the blast hits and checked whether they are reciprocal best hits of each other. For the *Sophophora* and

Drosophila sub-genera, we used *D. melanogaster* and *D. virilis* sequences downloaded from flybase as

the focal species and blasted them first to their close relatives. When one species yields no blast hit for

a gene, we then use other closely related species where the orthologs was successfully identified. If no

hits can be identified for a species or a clade, we then repeat the same procedure using tblastn to

identify translated protein sequences, as amino acid can be more conserved than nucleotide sequence.

If tblastn fails to identify a homologous transcript, we then blastn-ed to the genome sequence. True

absences/loss of a gene will yield poor or no blast hits, while missing annotation will result in clear

noncontiguous hits with gaps corresponding to introns.

Microsynteny surrounding homolgs and chromosome placement

We extracted the sequences of the homologs including 50kb up and downstream using bedtools slop

and bedtools getfasta. We then pairwise blastn-ed the sequences to each other and filtered out

alignments with E-values of < 0.01 or shorter than 100 bp. To infer the extent of homology in the

flanking sequences, we calculated the proportion of sequence aligned, excluding the positions of the

homolog. Genes are deemed to be in non-syntenic regions if they share < 5% flanking homology. For

species without Muller element designation of chromosomes, we assigned Muller elements by blastn-

ing to the genome of a closely related species where the Muller elements have been determined.

Phylogeny construction

We retrieved the CDS for all genes, removed the stop codon, and converted them first to protein

sequences using EMBOSS Transeq [94]. We then aligned the protein sequences using three aligners

with the commands: prank (v.170427) -protein -showtree [95], mafft (v7.505) --localpair --maxiterate

1000 [96], and muscle (v5.1) [97]. The resulting multi-sequence alignment fasta file were used as the

input for iqtree (v1.6.12) [98] with the flags -AA and -bb 2000 for 2000 iterations of ultrafast boot-

strapping [99]. These trees were manually rooted with *S. lebanonensis* as the outgroup species in

FigTree (v1.4.4) [100], and then Node labels added with phytools (v1.5.1) [101] to the trees to facilitate

downstream rate of evolution analyses with Hyphy. We then selected the resulting trees with the best

bootstrap support and concordance with species tree. *cona* trees were highly inconsistent across

alignment methods with many poorly supported nodes.

Branch-specific rate of protein evolution and positive selection with HyPhy

We used TranslatorX [102] to align the CDS sequence based on the protein alignments. Providing the

CDS alignments and the protein trees, we used the ABSREL module in HyPhy (v2.5.51) [47,103] to

infer the branch-specific rate of protein evolution (Omega) and significant signatures of positive

selection. We wrote a custom script (github.com/weikevinhc/phyloparse) to parse the HyPhy .json

output in R where the trees were reoriented with phytools and visualized with colors representing gene-

wide omega values. The nominal p-values were used for significance. In addition, we used PAML [104]

for the species-group specific test of recurrent positive selection. For comparisons with the rate of

regulatory evolution, we reran HyPhy after removing sequences from species with no RNA-seq data.

Rate of protein evolution with PAML analyses

From the genus wide alignment of SC genes, we extracted six species, *D. melanogaster, D. simulans,*

D. sechelia, D. yakuba, D. erecta, and D. ananassae, the same six *Drosophila* species used to

calculate dN/dS previously in Clark et al 2007 [105]. We then constructed maximum-likelihood trees

using iqtree using parameters '-m MFP -nt AUTO -alrt 1000 -bb 1000 -bnni'. We then used the PAML

parameter (model = 0 and CodonFreq = 2) to estimate the rate of protein evolution across the six

species.

Protein structure prediction with AlphaFold

Structures of proteins previously annotated in NCBI were retrieved from the AlphaFold Protein

Structure Database [41]. For genes we annotated, we used ColabFold (v1.5.2), an implementation of

AlphaFold on the Google Colab platform [42] and selected num_recycles 24, producing structure

predictions that were visualized in UCSF ChimeraX [106].

Data availability:

All generated raw sequences have been deposited onto the SRA under PRJNA1030345 (to be

released upon publication). Relevant intermediate datasets and files have been deposited on dryad

currently available to reviewers at:

https://datadryad.org/stash/share/V4wnkUUIEi_RkeK1XMF9qREOcynzDvA2QjmvjrrAKgU

References

1. True JR, Mercer JM, Laurie CC. Differences in crossover frequency and distribution among three
sibling species of Drosophila. Genetics. 1996;142: 507–523.

2. Stern C. An Effect of Temperature and Age on Crossing-Over in the First Chromosome of
Drosophila Melanogaster. Proc Natl Acad Sci U S A. 1926;12: 530–532.

3. Kent TV, Uzunović J, Wright SI. Coevolution between transposable elements and recombination.
Philos Trans R Soc Lond B Biol Sci. 2017;372. doi:10.1098/rstb.2016.0458

4. Feldman MW, Otto SP, Christiansen FB. Population genetic perspectives on the evolution of
recombination. Annu Rev Genet. 1996;30: 261–295.

5. Bomblies K, Higgins JD, Yant L. Meiosis evolves: adaptation to external and internal environments.
New Phytol. 2015;208: 306–323.

6. Brandvain Y, Coop G. Scrambling eggs: meiotic drive and the evolution of female recombination
rates. Genetics. 2012;190: 709–723.

7. Lenormand T. The evolution of sex dimorphism in recombination. Genetics. 2003;163: 811–822.

8. Sardell JM, Kirkpatrick M. Sex Differences in the Recombination Landscape. Am Nat. 2020;195:
361–379.

9. Morgan CH, Zhang H, Bomblies K. Are the effects of elevated temperature on meiotic
recombination and thermotolerance linked via the axis and synaptonemal complex? Philos Trans R
Soc Lond B Biol Sci. 2017;372. doi:10.1098/rstb.2016.0470

10. Anderson JA, Gilliland WD, Langley CH. Molecular population genetics and evolution of Drosophila
meiosis genes. Genetics. 2009;181: 177–185.

11. Brand CL, Cattani MV, Kingan SB, Landeen EL, Presgraves DC. Molecular Evolution at a Meiosis
Gene Mediates Species Differences in the Rate and Patterning of Recombination. Curr Biol.
2018;28: 1289-1295.e4.

12. Hemmer LW, Blumenstiel JP. Holding it together: rapid evolution and positive selection in the
synaptonemal complex of Drosophila. BMC Evol Biol. 2016;16: 91.

13. Samuk K, Manzano-Winkler B, Ritz KR, Noor MAF. Natural Selection Shapes Variation in
Genome-wide Recombination Rate in Drosophila pseudoobscura. Curr Biol. 2020;30: 1517-
1528.e6.

14. Ault JG, Lin HP, Church K. Meiosis in Drosophila melanogaster. IV. The conjunctive mechanism of
the XY bivalent. Chromosoma. 1982;86: 309–317.

15. Ritz KR, Noor MAF, Singh ND. Variation in Recombination Rate: Adaptive or Not? Trends Genet.
2017;33: 364–374.

16. Zelkowski M, Olson MA, Wang M, Pawlowski W. Diversity and Determinants of Meiotic
Recombination Landscapes. Trends Genet. 2019;35: 359–370.

17. Henderson IR, Bomblies K. Evolution and Plasticity of Genome-Wide Meiotic Recombination
Rates. Annu Rev Genet. 2021;55: 23–43.

18. Page SL, Hawley RS. The genetics and molecular biology of the synaptonemal complex. Annu

665    Rev Cell Dev Biol. 2004;20: 525–558.

666  19. Schmekel K, Skoglund U, Daneholt B. The three-dimensional structure of the central region in a
667     synaptonemal complex: a comparison between rat and two insect species, Drosophila
668     melanogaster and Blaps cribrosa. Chromosoma. 1993;102: 682–692.

669  20. Carpenter AT. Electron microscopy of meiosis in Drosophila melanogaster females. I. Structure,
670     arrangement, and temporal change of the synaptonemal complex in wild-type. Chromosoma.
671     1975;51: 157–182.

672  21. Hughes SE, Miller DE, Miller AL, Hawley RS. Female Meiosis: Synapsis, Recombination, and
673     Segregation in Drosophila melanogaster. Genetics. 2018;208: 875–908.

674  22. Loidl J. S. pombe linear elements: the modest cousins of synaptonemal complexes. Chromosoma.
675     2006;115: 260–271.

676  23. Schild-Prüfert K, Saito TT, Smolikov S, Gu Y, Hincapie M, Hill DE, et al. Organization of the
677     synaptonemal complex during meiosis in Caenorhabditis elegans. Genetics. 2011;189: 411–421.

678  24. Hawley RS. Solving a meiotic LEGO puzzle: transverse filaments and the assembly of the
679     synaptonemal complex in Caenorhabditis elegans. Genetics. 2011. pp. 405–409.

680  25. Kursel LE, Cope HD, Rog O. Unconventional conservation reveals structure-function relationships
681     in the synaptonemal complex. Elife. 2021;10. doi:10.7554/eLife.72061

682  26. Page SL, Khetani RS, Lake CM, Nielsen RJ, Jeffress JK, Warren WD, et al. Corona is required for
683     higher-order assembly of transverse filaments into full-length synaptonemal complex in Drosophila
684     oocytes. PLoS Genet. 2008;4: e1000194.

685  27. Collins KA, Unruh JR, Slaughter BD, Yu Z, Lake CM, Nielsen RJ, et al. Corolla is a novel protein
686     that contributes to the architecture of the synaptonemal complex of Drosophila. Genetics.
687     2014;198: 219–228.

688  28. Page SL, Hawley RS. c(3)G encodes a Drosophila synaptonemal complex protein. Genes Dev.
689     2001;15: 3130–3143.

690  29. Webber HA, Howard L, Bickel SE. The cohesion protein ORD is required for homologue bias
691     during meiotic recombination. J Cell Biol. 2004;164: 819–829.

692  30. Manheim EA, McKim KS. The Synaptonemal complex component C(2)M regulates meiotic
693     crossing over in Drosophila. Curr Biol. 2003;13: 276–285.

694  31. Mahajan S, Wei KH-C, Nalley MJ, Gibilisco L, Bachtrog D. De novo assembly of a young
695     Drosophila Y chromosome using single-molecule sequencing and chromatin conformation capture.
696     PLoS Biol. 2018;16: e2006348.

697  32. Miller DE, Staber C, Zeitlinger J, Hawley RS. Highly Contiguous Genome Assemblies of 15
698     Drosophila Species Generated Using Nanopore Sequencing. G3 . 2018;8: 3131–3141.

699  33. Bracewell R, Chatla K, Nalley MJ, Bachtrog D. Dynamic turnover of centromeres drives karyotype
700     evolution in Drosophila. Elife. 2019;8. doi:10.7554/eLife.49002

701  34. Hill T, Rosales-Stephens H-L, Unckless RL. Rapid divergence of the male reproductive proteins in
702     the Drosophila dunni group and implications for postmating incompatibilities between species. G3 .
703     2021;11. doi:10.1093/g3journal/jkab050

35. Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, et al. Highly contiguous
assemblies of 101 drosophilid genomes. Elife. 2021;10. doi:10.7554/eLife.66405

36. Chakraborty M, Chang C-H, Khost DE, Vedanayagam J, Adrion JR, Liao Y, et al. Evolution of
genome structure in the Drosophila simulans species complex. Genome Res. 2021;31: 380–396.

37. Wei KH-C, Mai D, Chatla K, Bachtrog D. Dynamics and Impacts of Transposable Element
Proliferation in the Drosophila nasuta Species Group Radiation. Mol Biol Evol. 2022;39.
doi:10.1093/molbev/msac080

38. Bachtrog D, Mahajan S, Bracewell R. Massive gene amplification on a recently formed Drosophila
Y chromosome. Nat Ecol Evol. 2019;3: 1587–1597.

39. Schaeffer SW, Bhutkar A, McAllister BF, Matsuda M, Matzkin LM, O'Grady PM, et al. Polytene
chromosomal maps of 11 Drosophila species: the order of genomic scaffolds inferred from genetic
and physical maps. Genetics. 2008;179: 1601–1655.

40. Anderson LK, Royer SM, Page SL, McKim KS, Lai A, Lilly MA, et al. Juxtaposition of C(2)M and
the transverse filament protein C(3)G within the central region of Drosophila synaptonemal
complex. Proc Natl Acad Sci U S A. 2005;102: 4482–4487.

41. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-sequence space with
high-accuracy models. Nucleic Acids Res. 2022;50: D439–D444.

42. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making protein
folding accessible to all. Nat Methods. 2022;19: 679–682.

43. Combet C, Blanchet C, Geourjon C, Deléage G. NPS@: network protein sequence analysis.
Trends Biochem Sci. 2000;25: 147–150.

44. Kimura M. Preponderance of synonymous changes as evidence for the neutral theory of molecular
evolution. Nature. 1977;267: 275–276.

45. Yang Z, Bielawski JP. Statistical methods for detecting molecular adaptation. Trends Ecol Evol.
2000;15: 496–503.

46. Yang Z, Nielsen R. Estimating synonymous and nonsynonymous substitution rates under realistic
evolutionary models. Mol Biol Evol. 2000;17: 32–43.

47. Kosakovsky Pond SL, Poon AFY, Velazquez R, Weaver S, Hepler NL, Murrell B, et al. HyPhy 2.5-
A Customizable Platform for Evolutionary Hypothesis Testing Using Phylogenies. Mol Biol Evol.
2020;37: 295–299.

48. Mason JM. Orientation disruptor (ord): a recombination-defective and disjunction-defective meiotic
mutant in Drosophila melanogaster. Genetics. 1976;84: 545–572.

49. Goldstein LS. Mechanisms of chromosome orientation revealed by two meiotic mutants in
Drosophila melanogaster. Chromosoma. 1980;78: 79–111.

50. Sandler L, Lindsley DL, Nicoletti B, Trippa G. Mutants affecting meiosis in natural populations of
Drosophila melanogaster. Genetics. 1968;60: 525–558.

51. Baker BS, Carpenter AT. Genetic analysis of sex chromosomal meiotic mutants in Drosophilia
melanogaster. Genetics. 1972;71: 255–286.

52. Baker BS, Hall JC. Meiotic mutants: genic control of meiotic recombination and chromosome
segregation. In: Ashburner M NE, editor. The Genetics and Biology of Drosophila. New York:
Academic Press; 1976. pp. 351–434.

53. Lindsley DL, Sandler L, Counce SJ, Chandley AC, Lewis KR. The Genetic Analysis of Meiosis in
Female Drosophila melanogaster [and Discussion]. Philos Trans R Soc Lond B Biol Sci. 1977;277:
295–312.

54. Rubin T, Macaisne N, Vallés AM, Guilleman C, Gaugué I, Dal Toe L, et al. Premeiotic pairing of
homologous chromosomes during *Drosophila* male meiosis. Proc Natl Acad Sci U S A. 2022;119:
e2207660119.

55. Wei KH-C, Chatla K, Bachtrog D. Single cell RNA-seq in Drosophila testis reveals evolutionary
trajectory of sex chromosome regulation. bioRxiv. 2022. p. 2022.12.07.519494.
doi:10.1101/2022.12.07.519494

56. Fast I, Hewel C, Wester L, Schumacher J, Gebert D, Zischler H, et al. Temperature-responsive
miRNAs in Drosophila orchestrate adaptation to different ambient temperatures. RNA. 2017;23:
1352–1364.

57. Marin P, Jaquet A, Picarle J, Fablet M, Merel V, Delignette-Muller M-L, et al. Phenotypic and
Transcriptomic Responses to Stress Differ According to Population Geography in an Invasive
Species. Genome Biol Evol. 2021;13. doi:10.1093/gbe/evab208

58. Voigt S, Froschauer C. Genome-wide temperature-sensitivity of Polycomb group regulation and
reduction thereof in temperate Drosophila melanogaster. Genetics. 2023;224.
doi:10.1093/genetics/iyad075

59. Chang C-H, Mejia Natividad I, Malik HS. Expansion and loss of sperm nuclear basic protein genes
in Drosophila correspond with genetic conflicts between sex chromosomes. Elife. 2023;12.
doi:10.7554/eLife.85249

60. Hopkins BR, Angus-Henry A, Kim BY, Carlisle JA, Thompson A, Kopp A. Decoupled evolution of
the *Sex Peptide* gene family and *Sex Peptide Receptor* in Drosophilidae. Proc Natl Acad Sci U S
A. 2024;121: e2312380120.

61. Dohm JC, Peters P, Stralis-Pavese N, Himmelbauer H. Benchmarking of long-read correction
methods. NAR Genom Bioinform. 2020;2: lqaa037.

62. Huang Y-T, Liu P-Y, Shih P-W. Homopolish: a method for the removal of systematic errors in
nanopore sequencing by homologous polishing. Genome Biol. 2021;22: 95.

63. Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. Diversity and dynamics of the
Drosophila transcriptome. Nature. 2014;512: 393–399.

64. Wen K, Yang L, Xiong T, Di C, Ma D, Wu M, et al. Critical roles of long noncoding RNAs in
Drosophila spermatogenesis. Genome Res. 2016;26: 1233–1244.

65. Begun DJ, Lindfors HA, Kern AD, Jones CD. Evidence for de novo evolution of testis-expressed
genes in the Drosophila yakuba/Drosophila erecta clade. Genetics. 2007;176: 1131–1137.

66. Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. Novel genes derived from noncoding
DNA in *Drosophila melanogaster* are frequently X-linked and exhibit testis-biased expression. Proc
Natl Acad Sci U S A. 2006;103: 9935–9939.

67. Witt E, Benjamin S, Svetec N, Zhao L. Testis single-cell RNA-seq reveals the dynamics of de novo

gene transcription and germline mutational bias in Drosophila. Elife. 2019;8.
doi:10.7554/eLife.47138

68. Van Oss SB, Carvunis A-R. De novo gene birth. PLoS Genet. 2019;15: e1008160.

69. Lawlor MA, Cao W, Ellison CE. A transposon expression burst accompanies the activation of Y-
chromosome fertility genes during Drosophila spermatogenesis. Nat Commun. 2021;12: 6854.

70. Muirhead CA, Presgraves DC. Satellite DNA-mediated diversification of a sex-ratio meiotic drive
gene family in Drosophila. Nat Ecol Evol. 2021;5: 1604–1612.

71. Vedanayagam J, Lin C-J, Lai EC. Rapid evolutionary dynamics of an expanding family of meiotic
drive factors and their hpRNA suppressors. Nat Ecol Evol. 2021;5: 1613–1623.

72. Neel JV. A Relation between Larval Nutrition and the Frequency of Crossing over in the Third
Chromosome of Drosophila Melanogaster. Genetics. 1941;26: 506–516.

73. Belmonte-Tebar A, San Martin Perez E, Nam Cha S, Soler Valls AJ, Singh ND, de la Casa-
Esperon E. Diet effects on mouse meiotic recombination: a warning for recombination studies.
Genetics. 2022;220. doi:10.1093/genetics/iyab190

74. Plough HH. The effect of temperature on crossing-over in Drosophila. Columbia University; 1917.

75. Lloyd A, Morgan C, H Franklin FC, Bomblies K. Plasticity of Meiotic Recombination Rates in
Response to Temperature in Arabidopsis. Genetics. 2018;208: 1409–1420.

76. Singh ND, Criscoe DR, Skolfield S, Kohl KP, Keebaugh ES, Schlenke TA. Fruit flies diversify their
offspring in response to parasite infection. Science. 2015;349: 747–750.

77. Hussin J, Roy-Gagnon M-H, Gendron R, Andelfinger G, Awadalla P. Age-dependent
recombination rates in human pedigrees. PLoS Genet. 2011;7: e1002251.

78. Hunter CM, Robinson MC, Aylor DL, Singh ND. Genetic Background, Maternal Age, and
Interaction Effects Mediate Rates of Crossing Over in Drosophila melanogaster Females. G3 .
2016;6: 1409–1416.

79. Pavličev M, Cheverud JM. Constraints Evolve: Context Dependency of Gene Effects Allows
Evolution of Pleiotropy. Annu Rev Ecol Evol Syst. 2015;46: 413–434.

80. Bracewell R, Tran A, Chatla K, Bachtrog D. Chromosome-Level Assembly of Drosophila bifasciata
Reveals Important Karyotypic Transition of the X Chromosome. G3 . 2020;10: 891–897.

81. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37: 907–915.

82. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

83. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-
and-vote. Nucleic Acids Res. 2013;41: e108.

84. Pachter L. Models for transcript quantification from RNA-Seq. arXiv [q-bio.GN]. 2011. Available:
http://arxiv.org/abs/1104.3889

85. Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol. 2015;33: 290–

295.

86.   Holt C, Yandell M. MAKER2: an annotation pipeline and genome-database management tool for
824   second-generation genome projects. BMC Bioinformatics. 2011;12: 491.

825   87.   Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for
826   automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A.
827   2020;117: 9451–9457.

828   88.   Korf I. Gene finding in novel genomes. BMC Bioinformatics. 2004;5: 59.

829   89.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative
830   genomics viewer. Nat Biotechnol. 2011;29: 24–26.

831   90.   Abeel T, Van Parys T, Saeys Y, Galagan J, Van de Peer Y. GenomeView: a next-generation
832   genome browser. Nucleic Acids Res. 2012;40: e12.

833   91.   Zhu T, Liang C, Meng Z, Guo S, Zhang R. GFF3sort: a novel tool to sort GFF3 files for tabix
834   indexing. BMC Bioinformatics. 2017;18: 482.

835   92.   Pertea G, Pertea M. GFF Utilities: GffRead and GffCompare. F1000Res. 2020;9: 304.

836   93.   Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Bioinformatics. 2021;37:
837   1639–1643.

838   94.   Madeira F, Pearce M, Tivey ARN, Basutkar P, Lee J, Edbali O, et al. Search and sequence
839   analysis tools services from EMBL-EBI in 2022. Nucleic Acids Res. 2022;50: W276–W279.

840   95.   Löytynoja A. Phylogeny-aware alignment with PRANK. Methods Mol Biol. 2014;1079: 155–170.

841   96.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in
842   performance and usability. Mol Biol Evol. 2013;30: 772–780.

843   97.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
844   Nucleic Acids Res. 2004;32: 1792–1797.

845   98.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-
846   TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol
847   Biol Evol. 2020;37: 1530–1534.

848   99.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the Ultrafast
849   Bootstrap Approximation. Mol Biol Evol. 2018;35: 518–522.

850   100.   FigTree. [cited 13 Jul 2023]. Available: http://tree.bio.ed.ac.uk/software/figtree/

851   101.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things):
852   phytools: R package. Methods Ecol Evol. 2012;3: 217–223.

853   102.   Abascal F, Zardoya R, Telford MJ. TranslatorX: multiple alignment of nucleotide sequences
854   guided by amino acid translations. Nucleic Acids Res. 2010;38: W7-13.

855   103.   Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Kosakovsky Pond SL. Less is more:
856   an adaptive branch-site random effects model for efficient detection of episodic diversifying
857   selection. Mol Biol Evol. 2015;32: 1342–1353.

104.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007;24: 1586–

1591.

105.   Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, et al. Evolution of genes
and genomes on the Drosophila phylogeny. Nature. 2007;450: 203–218.

106.   Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF ChimeraX:
Structure visualization for researchers, educators, and developers. Protein Sci. 2021;30: 70–82.