Title: Critical reproductive behaviors in Scaled Quail and Northern Bobwhite are affected by
thermal variability and mean temperature

Authors: William Kirkpatrick[1], Erin Sauer[1], Rachel Carroll[2], Jeremy Cohen[3,4], Craig Davis[5],
Samuel Fuhlendorf[5], Sarah DuRant[1]

Affiliation: [1]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[2]Jones College of Science, Engineering, and Technology, Murray State University, Murray, KY,
USA

[3]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA

[4]Center for Biodiversity and Global Change, Yale University, New Haven, CT, USA

[5]Department of Natural Resource Ecology and Management, Oklahoma State University,
Stillwater, OK, USA

Keywords: climate change, environmental temperature, thermal variation, behavior, incubation,

quail, *Colinus virginianus, Callipepla squamata*

Abstract

Environmental conditions can influence parental care behaviors, which in turn can
greatly affect the microhabitat conditions experienced by developing offspring. It is critical to
uncover relationships between ambient weather, parental behaviors, and offspring thermal
environments to understand the ability of organisms to mitigate environmental conditions and
predict outcomes under climate change. In warm conditions, avian parents may spend longer
periods away from the nest, but we know little of how increased thermal variation drives these
changes or what temporal scale influences the changes. We examined how parents mediate
environmental conditions for developing offspring in two quail species, Scaled Quail (*Callipepla
squamata*) and Northern Bobwhite (*Colinus virginiatus*), by examining changes in the duration
and timing of incubation off-bouts (recesses away from the nest) in relation to mean temperature
and temperature variability experienced over either a 24-hour period (low resolution) or in the
two hours immediately preceding an off-bout (high resolution). We found species-specific
differences in off-bout duration, with Scaled Quail spending a significantly shorter time off the
nest daily. In both species, we detected a positive relationship between average thermal
conditions and the start time of afternoon off-bouts. Alternatively, morning off-bouts occur
earlier in response to increased thermal variation from the previous day. Warm conditions
immediately preceding off-bouts led to increased off-bout duration with the opposite trend in
response to increased variation. Our findings suggest that the thermal environment, in contrasting
ways over varying thermal and temporal scales, can have a significant impact on parental care
behaviors. These alterations can lead to altered thermal environments for offspring across
multiple scales. Therefore, determining how the thermal environment induces behavioral
changes requires high-resolution environmental data.

Introduction

Moderate projections of anthropogenic climate warming estimate an average global temperature increase of 1.1 – 2.6°C by 2100 (1) with projected losses of 7.9% of all species due to climate change (2). However, mean increases in temperature alone will not reflect the shifts in thermal conditions that organisms will experience. The fallacy of the average, known as Jensen's Inequality, highlights the importance of exploring thermal variability in natural systems, as relationships between external conditions and biological responses are often nonlinear and can change drastically under unusual or extreme conditions (ex. the relationship between metabolic rates and average thermal conditions described in Denny 2017; 3–5). Increases in thermal variability are increasing alongside warming average temperatures (6,7), and organisms can have differing responses to increasing thermal averages or variability (8,9).

Anthropogenic climate change is projected to cause phenotypic changes (10) which can be driven or mitigated by plastic behaviors (11). For wildlife, plastic behaviors are necessary to avoid reductions in fitness in response to extreme or novel thermal conditions (12). Thus, behavioral plasticity is an important driver of organismal responses to climate warming. Behaviors employed by parents to care for their offspring (i.e., parental care behaviors) are known to be plastic (13), play an important role in shaping the thermal environment of developing offspring, and are often directly linked to offspring phenotypic expression (14). Ambient thermal conditions can alter parental care behaviors and subsequently modify the thermal environment of the developing offspring to directly influence offspring phenotypes (15,16). Examining shifts in parental care behaviors of wild organisms in response to the natural thermal environment can reveal the ability of parents to modify their behaviors to meet the competing demands of their offspring's thermal needs and self-maintenance, which becomes critical in suboptimal thermal conditions (17,18). Multiple species have expressed these trade-offs (19,20), including large bodied homeotherms like Eastern Wild Turkeys (*Meleagris gallopavo silvestris*; 21).

Environmental temperature may be an important instigator of behavioral and phenotypic change (22,23) over distinct temporal scales that convey different information about how animals respond to the thermal environment. Understanding how short-term thermal fluctuations influence parental behavior in real time is critical for understanding drivers of the developmental environment (24). Low-resolution, temporally averaged thermal data is useful in many contexts, especially when exploring seasonal trends in behavioral patterns (25). For instance, greater than average temperature over the course of a reproductive season in birds (i.e., low-resolution thermal and behavioral data) is associated with shorter incubation periods (26,27), possibly indicating a reduced cost of reproduction for parents. However, assessing only average temperature over the entire incubation stage obscures the thermal variability occurring during this period. During an acute period of hot temperatures, parents may increase nest attendance to shield eggs or seek refuge, leaving eggs exposed to a suboptimal thermal environment. Thermal biologists have explored the relationship between phenological outcomes and their variation alongside long-term thermal trends (28), but less is known about how real-time or recently experienced temperatures shape isolated behavioral responses. Exploring the effects of both low-

and high-resolution thermal data on specific behavioral changes can inform the mechanisms
behind large scale changes in species' ranges, phenotypic changes, and fitness (29).

Birds present a unique opportunity to investigate the relationship between high-frequency
thermal measurements and behavioral decisions due to the role parents play in shaping the
thermal environment for their developing offspring (30). Birds in hot environments require self-
maintenance behaviors such as gular fluttering or retreating to cooler microclimates to avoid
physiological stress (31,32). However, coping behaviors that cause the parent to leave the nest
can come at the cost of offspring care, as the nest is left exposed to external thermal conditions
(14,15,30). The timing of off-bouts (the time a parent spends off the nest) and their duration play
important roles in the development of avian offspring (27). Lengthening and, in some cases,
delaying off-bouts can lead to higher rates of predation (28) or thermal stress, reducing adult and
offspring fitness (33).We know little of the scale at which these tradeoff decisions are being
made and their relationship to natural thermal variation (34,35), but the importance of parental
care on offspring development suggests that reproductive success would benefit from the ability
of parents to delay off-bouts until temperatures are outside lethal levels for embryos (27).
Acclimation to thermal variation and short-term thermal conditions has been evidenced in
ectotherms (36–38), though evidence in endotherms is limited to animals living in the tropics due
to their narrow ranges of thermal tolerance (39). Largely, studies focusing on temperate avian
responses to thermal change focus on phenological adjustments and seasonal shifts in the thermal
environment (40,41). There is little work focusing on avian responses to rapid changes in
temperature, with the majority limited to individual thermoregulatory responses (20,21,42). The
ability of parents to alter their behavior will depend on their ability to withstand physiologically
demanding temperatures and to respond to thermal conditions as they are occurring, thus studies
examining high resolution thermal and behavioral data are necessary.

The primary goal of this study is to determine the influence of average environmental
temperature and thermal variability on incubation behaviors and resulting offspring microhabitat
in Scaled Quail (*Callipepla squamata*) and Northern Bobwhite (*Colinus virginianus*) across two
temporal scales: immediately or over a full day prior to parents leaving the nest. Using data
collected in 2015 and 2016 and described in Carroll *et al*. (30), we compare differences in
behavioral modifications of these species as it relates to temperature variability, average
temperature, and the resolution of thermal data. Scaled and Northern Bobwhite overlap in the
Southwestern US; Northern Bobwhite typically occupy a temperate climate across their range
while Scaled Quail largely occupy arid, desert-like environments like the study locale. Nest
construction is similar in both species, though Scaled Quail select nesting locations with more
favorable vegetation for thermal buffering (30). We hypothesize that: 1) Thermal variability is an
important driver of total daily off-bouts, with greater thermal variation delaying off-bout start
time and shortening individual off-bout duration to avoid inhospitable nest conditions. 2) The
temporal scale of thermal measurements will influence the resulting relationships between
behavior and temperature, with short term thermal trends and behavior exhibiting a stronger
relationship than extended, temporally averaged thermal trends.

Methods

Monitoring Temperature and Incubation Patterns

All nests were monitored in the Beaver River Wildlife Management Area in Oklahoma, USA (36°50′21″N, 100°42′15″W). Environmental temperature data was collected every 30 seconds within and outside the nest using modified Onset Computer HOBOware (Onset Corporation, Borne, MA) thermal sensors. External temperatures were taken with HOBO U23 Pro v2 External Temperature Loggers (30). The internal temperature was taken using egg shaped clay models inserted into the nest that matched the dimensions of quail eggs. Each model egg was inserted into the center of the nest and external sensor was placed within 2m of the nest for standardization of thermal measurements. More details of thermal probes and data collection can be found in Carroll *et al.* (30).

The machine learning software NestIQ was used to determine off-bout length and timing from the raw nest temperature data (43). To assess the impact the rate of thermal change has on off-bout duration, we calculated the average and standard deviation of environmental temperature measurements during the two hours before each off-bout. To assess the impact the daily thermal environment has on total daily off-bout, we calculated average and standard deviation of temperatures from the morning (sunrise to noon), afternoon (noon to sunset), and night (previous day sunset to sunrise) as well as the temperatures from the previous day and previous afternoon. All sunrise and sunset times were recorded for each day of the summers of 2015 and 2016 at the field site.

Statistical Methodology

All statistical analyses were conducted with R version 4.1.2 (44). To test the effect of the mean and standard deviation of temperature during the two-hour period before each off-bout (i.e., high-resolution environmental data) on off-bout duration, we developed a linear mixed model (45). The universal model included mean and standard deviation of the two-hour period preceding the off-bout as predictors and off-bout length (in minutes) as the response variable. We included a two-level categorical predictor for whether the off-bout took place in the morning or afternoon, as each species takes a maximum of two off-bouts per day, in the early morning or late afternoon (referred to as AMPM in models). We also included species (Scaled Quail or Northern Bobwhite). To control for non-independence of the measurements at a single nest and year, we included individual nest ID and year as random intercepts. All possible interactions between predictors were included in the universal model. To test for relationships between temperature and off-bout start time, we conducted a separate linear mixed model with the same predictors, interactions, and random intercepts described above but with off-bout timing as the response variable (Table 1).

To address the effect of temperatures experienced over a longer time period (i.e., low-resolution environmental data) on total daily off-bout, we developed a linear mixed effects model which included the sum of all off-bouts in 24 hours as the response variable. We selected total daily off-bouts as our response variable because the predictors are date-specific thermal measurements instead of off-bout specific thermal measurements. The predictive variables

included the mean and standard deviation of external temperature from the following timeframes
relative to the date: 24hr (temperatures from 12AM-11:59PM of the current day), Morning
(sunrise to 12PM), Afternoon (12PM to sunset), Previous Night (sunset to sunrise of previous
night), Previous Day (12AM to 11:59PM of the previous day), and Previous Afternoon (12PM to
sunset of the previous day). We chose these time periods to allow for comparison of longer-term
trends in temperature. While they are likely highly correlated with one another, our model
selection strategy (see below – *Model Selection*) corrected for this issue. We included a two-level
categorical predictor for species (Scaled or Northern Bobwhite). To control for non-
independence of the nest location and year, we included individual Nest ID and year as random
intercepts. To test for relationships between mean and standard deviation of temperature on the
average daily time of each off-bout, we conducted two separate linear mixed models with the
same predictors, interactions, and random intercepts described above, but the response variables
were the timing of the morning or afternoon off-bout respectively, given that each species takes
at most, two off-bouts during the day in the morning, evening, or both.

Model Selection

For each universal mixed-effects model (high and low-resolution), we used the dredge
function from the MuMIn package for model selection (46). We chose the best fit model by
finding the model with the lowest relative AICc score (47). All values represented in our figures
are partial residual plots generated using the predict function in R.

The best fit model for the high-resolution approach to analyze individual off-bout
duration (referred to in Results as High Duration) included 2-hour mean temperature, 2-hour
standard deviation of temperature, morning or afternoon (AMPM), and species as main effects
with three interactive effects: 2-hour mean*species, 2-hour standard deviation of
temperature*species, and AMPM*species (Table 1).

Two models were used to predict individual off-bout start time, one that included all
morning off-bouts (referred to in Results as High AM Start Time) and one that included all
afternoon off-bouts (referred to in Results as High PM Start Time). Both models included 2-hour
mean temperature and 2-hour standard deviation of temperature as main effects, but only High
AM Start Time included species (Table 1).

The best fit model for predicting total daily off-bout (referred to in Results as Low
Duration) included 24hr standard deviation of temperature, morning standard deviation of
temperature, and previous night standard deviation of temperature; Species and two interactive
effects were included: species*morning standard deviation of temperature and species*previous
night standard deviation of temperature (Table 1).

Two models were used to predict daily off-bout start time, one that included morning
off-bouts (referred to in Results as Low AM Start Time) and one that included afternoon off-
bouts (referred to in Results as Low PM Start Time) since each species take a maximum of two
off-bouts per day in either the morning or evening. The best fit model for the morning off-bouts
(Low AM Start Time) included standard deviation of temperature in the previous day and species
as main effects with no interactive effects (Table 1). The best fit model for the afternoon off-

bouts (Low PM Start Time) included mean temperature during the afternoon and species with no
interactive effects (Table 1).

High-Resolution Best Fit Models							
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AIC$_c$**	**df**	**AIC$_c$ w$_i$**	
High Duration	Off-bout Duration	2hr Mean, 2 hr SD, AMPM, Species	2hr Mean:Species, 2hr SD:Species, AMPM:Species	8952.7	11	0.742	
Null	Off-bout Duration	~	~	9154.8	4	< 0.0001	
High AM Start Time	Off-bout Timing - AM	2hr Mean, 2hr SD, Species	~	1375.2	7	0.429	
Null	Off-bout Timing - AM	~	~	1417.8	4	< 0.001	
High PM Start Time	Off-bout Timing - PM	2hr Mean, 2hr SD	~	1580.9	6	0.425	
Null	Off-bout Timing - PM	~	~	1654.8	4	< 0.001	
Low-Resolution Best Fit Models							
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AIC$_c$**	**df**	**AIC$_c$ w$_i$**	
Low Duration	Daily Off-bout Duration	24hr SD, Morning SD, Previous Night SD, Species	24hr SD:Species, Previous Night SD:Species	5666.7	10	0.02	
Null	Daily Off-bout Duration	~	~	5706	4	< 0.001	
Low AM Start Time	Off-bout Timing - AM	Previous Day SD, Species	~	1378.9	6	0.075	
Null	Off-bout Timing - AM	~	~	1395.6	4	< 0.0001	
Low PM Start Time	Off-bout Timing - PM	Afternoon Mean, Species	~	1349.7	6	0.22	
Null	Off-bout Timing - PM	~	~	1374.4	4	< 0.0001	

202
Table 1. The best fit models for each suite of thermal predictors as selected by AICc using the
package MuMIn. High-resolution: The universal model included mean and standard deviation of
the two-hour period preceding the off-bout as predictors and off-bout length (in minutes) as the
response variable. We included a two-level categorical predictor for whether the off-bout took
place in the morning or afternoon (referred to as AMPM) and species (Scaled Quail or Northern
Bobwhite). The universal model for morning and afternoon start time included all above (except
for AMPM) with individual off-bout start time as the response. Low-resolution: All models were
structured as above except with averaged thermal data over 6 deterministic timeframes as
predictors (24hr (temperatures from 12AM-11:59PM of the current day), Morning (sunrise to
12PM), Afternoon (12PM to sunset), Previous Night (sunset to sunrise of previous night),
Previous Day (12AM to 11:59PM of the previous day), and Previous Afternoon (12PM to sunset
of the previous day)). We included individual nest ID and year as random intercepts in all
models. All possible interactions between predictors were included in each universal model.

Results

High-Resolution Thermal Data – Duration

As described by model High Duration, off-bout duration increased by ~4.4 ± 0.48
minutes per degree increase of 2-hour mean temperature ($\beta = 4.37 \pm 0.48$, t = 9.1, p < 0.001;
Figure 1A). Off-bout duration decreased by ~ 4.2 ± 2.2 minutes per unit increase of 2-hour
standard deviation of temperature, though the association was only moderate ($\beta = -4.16 \pm 2.16$, t
= -1.9, p = 0.055; Figure 1B).

Northern Bobwhite averaged 140 ± 68 minutes per off-bout while Scaled Quail averaged
110 ± 57 minutes ($\beta = 62 \pm 14.7$, t = 4.2, p < 0.001). Northern Bobwhite exhibited longer
afternoon off-bouts (159 ± 61 minutes) than Scaled Quail (Afternoon =108 ± 48 minutes; β = -
43.1 ± 10.5, t = -4.1, p > 0.001; Supplementary Figure 1A) even when mean temperatures 2
hours preceding the off-bout were high ($\beta = -1.8 \pm 0.7$, t = -2.7, p = 0.007, Supplementary Figure
1B). For both species morning off-bouts were not significantly longer than afternoon off-bouts,

though they were different (β = 1.5 ± 7.4, t = 0.2, p = 0.8; Morning = 110 ± 66 minutes;
Afternoon = 138 ± 61 minutes).

High-Resolution Thermal Data – Start Time

As described by model High AM Start Time, per degree increase of 2-hour mean
temperature, morning off-bouts began 9 ± 3 minutes later (β = 0.09 ± 0.03, t = 3.2, p = 0.002;
Figure 1C. Similarly, off-bouts began 37 ± 10 minutes later per unit increase of 2-hour standard
deviation (β = 0.37 ± 0.10, t = 3.7, p < 0.001; Figure 1D). Scaled Quail began their off-bouts 55
± 28 minutes earlier in the morning than Northern Bobwhite, though the association was only
moderate (β = -55 ± 0.28, t = -1.9, p = 0.06).

As described by model High PM Start Time, per degree increase of 2-hour mean
temperature, afternoon off-bouts began 21 ± 3 minutes earlier (β = -0.21 ± 0.03, t = -8,1, p <
0.001; Figure 1E. Alternatively, off-bouts began 33 ± 7 minutes later per unit increase of 2-hour
standard deviation (β = 0.33 ± 0.07, t = 4.5, p < 0.001; Figure 1F).

Figure 1. Partial residual plots showing the relationships between the thermal environment immediately preceding individual off-bouts and off-bout duration (A, B) as well as off-bout start time (C-F). Blue lines exhibit the association across both years while black and gray lines represent 2015 and 2016, respectively. Open circles represent quail nests in 2015, and closed

circles are 2016 nests. X-values represent thermal averages 2 hours prior to the off-bout for A, C, and E. Whereas x-values for B, D, and F represent standard deviation (thermal variation) two hours prior to the off-bout. A) As mean temperature in the 2 hours preceding each off-bout increased, Northern Bobwhite (green) and Scaled Quail (blue) off-bout duration increased. Nest sites in 2015 were cooler on average than 2016 (2015 = 21.5 C, 2016 = 31.2 C). B) As standard deviation of temperature increased, individual off-bout duration decreased. C) Both species delayed taking off-bouts until later in the morning as mean temperature two hours preceding the off-bout increased, and D) a similar relationship exists with standard deviation of temperature. E) Afternoon off-bout start time began earlier as average temperature 2 hours prior to the off-bout increased while F) start time was delayed as thermal variation increased. All points are predicted model values.

Low-Resolution Thermal Data – Total Daily Off-Bout

As described by model Low Duration, per unit increase of 24hr standard deviation of temperature, total daily off-bouts lengthened by 11.9 ± 3.7 minutes ($\beta = 11.9 \pm 3.7$, $t = 3.2$, $p = 0.001$; Figure 2A). Alternatively, per unit increase of morning standard deviation of temperature, total daily off-bout decreased by 4.2 ± 2.3 minutes, though the association is only moderate ($\beta = -4.2 \pm 2.3$, $t = -1.8$, $p = 0.07$; Figure 2B). Scaled Quail spent less total time off the nest daily (156 ± 74 minutes) than Northern Bobwhite (197.2 ± 81 minutes; $\beta = 71 \pm 34$, $t = 2.1$, $p = 0.03$). Scaled Quail had significantly shorter off-bouts than Northern Bobwhite in more variable environments ($\beta = -9.9 \pm 3.8$, $t = -2.5$, $p = 0.01$; Supplementary Figure 1C). There were no other significant effects.

Low-Resolution Thermal Data – Start Time

As described by model Low AM Start Time, per unit increase of previous day standard deviation, off-bout start time in the morning began 15 ± 3 minutes earlier ($\beta = -0.15 \pm 0.03$, $t = -4.4$, $p < 0.001$; Figure 2C).

As described by model Low PM Start Time, as afternoon average temperature increased, afternoon off-bouts started 7 ± 2 minutes later ($\beta = 0.07 \pm 0.02$, $t = 4.1$, $p < 0.001$; Figure 2D).

Figure 2. Partial residual plots showing the relationships between the thermal environment
averaged over defined deterministic timeframes and off-bout duration (A, B), morning off-bout
start time (C), and afternoon start time (D). A) X-values represent the standard deviation of

temperature over a 24 hour period. As thermal variation increased, both species reduced their
total daily off-bout duration. B) X-values represent the standard deviation of temperature over
the morning (sunrise-noon). As thermal variation in the morning increased, both species reduced
their total daily off-bout. C) X-values represent the standard deviation of temperature of the
previous day. Morning off-bouts began earlier as the standard deviation of the temperature of the
previous day increased. D) X-values represent the average afternoon temperature. As afternoons
became warmer on average, both species delayed their off-bouts until later in the evening. All
points are predicted model values.

Discussion

We investigated the relationship between thermal means, thermal variability, and
incubation behaviors across multiple timescales in Scaled Quail and Northern Bobwhite. We
provide evidence that both species are modifying behavior to avoid even short periods of higher
thermal variability, a novel finding among studies of avian parental behavior. Further
relationships between thermal endpoints (mean vs. variation) and behavior differed depending on
the resolution of the data. These results provide evidence that thermal-behavioral relationships
depend on the scale at which endpoints are measured, and scale is important to consider when
contextualizing behavioral outcomes.

Short-term increases to the mean and variation of temperature delay the beginning of the
morning off-bout by ~9 and ~37 minutes, respectively (Figure 1C, D). However, when we
examine similar responses on a low-resolution timescale, thermal variation of the previous day
advanced the start of morning off-bout by 15 minutes (Figure 2C). On the surface, this seems
like the opposite of what was observed on the high-resolution scale, but the interpretation of
these relationships requires contextualization beyond direct comparison, as the high-resolution
measurements examine the relationship between the rate of environmental thermal change and a
direct measurement of decision-making (i.e., individual timing of off-bouts). However, the low-
resolution result examines the relationship between single measurement predictors averaged over
time and their impacts on an averaged response, providing context for periods in the day that
may be influential to general trends in behavior, but these measurements say little about
moment-to-moment decision-making. Similarly, while afternoon average temperatures pushed
each species to delay the start of an off-bout by 7 minutes (Figure 2D), the opposite was
observed on the high-resolution scale (off-bouts began 21 minutes earlier, Figure 1E). These
seemingly alternating relationships provide more evidence that sole characteristic predictors
require careful contextualization. The low-resolution predictors (model name – Low Duration,
0.02 AIC_C w_i) may be significant, but the model weights suggest that the high-resolution
predictors make for a better predictive model (Table 1), likely due to the loss of resolution that
comes with averaging behavioral data.

Northern Bobwhite and Scaled Quail increased the duration of their individual off-bouts
in response to the immediate thermal environment at a rate of 4.4 minutes/mean degree increase
(Figure 1A) and decreased off-bouts in more variable environments at a rate of 4.2 minutes/unit
increase of variation (Figure 1B). Both species are more comfortable leaving eggs in warm,
stable conditions which are less likely to expose eggs to temperatures outside of critical ranges.

Yet, when we examined these relationships at a lower resolution, total daily off-bout was not
influenced by average temperatures, only thermal variation (with significantly reduced model
weights – Table 1; Figure 2A, B). These findings further suggest that the timescales we use to
measure thermal change and behavioral modification alter how we view the plasticity of
organismal behavior to the thermal environment. Multiple studies have shown that the
relationship between temperature and the average lengths of off-bouts predicts avian
reproductive success (48–51). Approaches considering averaged conditions can contextualize
general trends in behavior (52,53), but as we show above, only examining behavior in this way
risks missing the relationship between the rate of thermal change and organismal behavior. Our
findings are supported by studies of other species nesting in hot conditions. Least Terns
(*Sternula antillarum*; 52,53) and Piping Plovers (*Charadrius melodus*; 53) exhibit some
thermoregulatory behaviors but only within only certain breadths of experienced temperature.
Using low-resolution thermal data alone to predict behavior would limit our understanding of
these responses, as the breadth of behavior may be washed out in a sole-characteristic thermal
predictor (56).

Here, both species expressed similar trends in their behavioral modifications but differed
in individual off-bout duration and environmental temperature of their nests (Supplemental
Figure 1). Scaled Quail spent significantly less time off the nest on average (110 minutes) than
Northern Bobwhite (140 minutes). The interactive effects of each model may be influenced by
natural history differences of the two species (57,58). The typical breeding habitat of Northern
Bobwhite is temperate, consisting of habitat exposed to a wide range of temperatures, but
extreme temperatures rarely reach those experienced by Scaled Quail across their range (57).
Scaled Quail are a grassland desert-dwelling species, adapted to the extreme highs and lows
present in the Western Oklahoma field site (58), reflected in superior thermal buffering observed
in nestbuilding compared to Northern Bobwhite (30). Behavioral adaptation to thermal extremes
can be limited by range-specific adaptations (59). Range edges are dynamic, and while high
turnover species adapt quickly to shifts in the environment on the edge of their range (60),
Northern Bobwhite have a limited ability to adapt to novel climates due to high temporal
investment in offspring, long generation time, and slow physical development (57). It should be
noted that direct comparisons between each species based on results found here need to be made
with caution, as thermal and behavioral measurements across each species' range would be
required to make species level comparisons in behavioral plasticity.

Behavioral plasticity during breeding generally benefits both the parent and dependent
offspring by keeping the thermal environment stable over time (61–63), but shifts in behavior
alone cannot completely negate the impacts of poor or unexpected environmental conditions
(64), as self-maintenance may take precedence over offspring care (35). Our results indicate that
the rate of environmental temperature change can have a direct influence on parental behavior.
We also demonstrate that thermal variability can be lost in averages, as evidenced by the lack of
a significant relationship in daily off-bout duration and temperature (Figure 2A) but a significant
relationship between acute thermal environment and off-bout duration (Figure 1A). Therefore,
averaged conditions may provide inaccurate assessments of organismal vulnerability to thermal

change if characterized incorrectly (8,65). Models with averaged, lone sole-characteristic
predictors should not be avoided but contextualized as we do here (56).

Acknowledgements

We would like to thank Rachel Carroll, Craig Davis, and Sam Fuhlendorf for permission to use
their thermal data from Scaled and Northern Bobwhite Quail nests. Thanks to Wayne Hawkins
for assistance with NestIQ.

Funding

This work was supported by the Oklahoma Department of Wildlife Conservation through
funding administered through the Oklahoma Cooperative Fish and Wildlife Research Unit (grant
#F11AF00069), the Oklahoma Agricultural Experiment Station at Oklahoma State University,
and the Bollenbach and Groendyke Endowments.

Ethics

Animal welfare approval was obtained from IACUC at Oklahoma State University.

Data, Code, and Materials

All supporting data and annotated R scripts are available in GitHub or through contact with the
corresponding author.

Competing Interests

We declare we have no competing interests.

Literature Cited

1. Calvin K, Dasgupta D, Krinner G, Mukherji A, Thorne PW, Trisos C, et al. IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. [Internet]. First. Intergovernmental Panel on Climate Change (IPCC); 2023 Jul. Available from: https://www.ipcc.ch/report/ar6/syr/

2. Urban MC. Accelerating extinction risk from climate change. Science. 2015 May;348(6234):571–3.

3. Denny M. The fallacy of the average: on the ubiquity, utility and continuing novelty of Jensen's inequality. Journal of Experimental Biology. 2017 Jan 15;220(2):139–46.

4. Jensen J. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. Acta mathematica. 1906;30(1):175–93.

5. Ruel JJ, Ayres MP. Jensen's inequality predicts effects of environmental variation. Trends in Ecology & Evolution. 1999;14(9):361–6.

6. Bathiany S, Dakos V, Scheffer M, Lenton TM. Climate models predict increasing temperature variability in poor countries. Sci Adv. 2018 May 4;4(5):eaar5809.

7. Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO. Climate Extremes: Observations, Modeling, and Impacts. Science. 2000 Sep 22;289(5487):2068–74.

8. Sheldon KS, Dillon ME. Beyond the Mean: Biological Impacts of Cryptic Temperature Change. Integrative and Comparative Biology. 2016 Jul;56(1):110–9.

9. Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS, et al. Increased temperature variation poses a greater risk to species than climate warming. Proceedings of the Royal Society B: Biological Sciences. 2014 Mar 22;281(1779):20132612.

10. Iler AM, CaraDonna PJ, Forrest JRK, Post E. Demographic Consequences of Phenological Shifts in Response to Climate Change. Annu Rev Ecol Evol Syst. 2021 Nov 2;52(1):221–45.

11. Muñoz MM, Stimola MA, Algar AC, Conover A, Rodriguez AJ, Landestoy MA, et al. Evolutionary stasis and lability in thermal physiology in a group of tropical lizards. Proc R Soc B. 2014 Mar 7;281(1778):20132433.

12. Sinclair BJ, Marshall KE, Sewell MA, Levesque DL, Willett CS, Slotsbo S, et al. Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? Vasseur D, editor. Ecology Letters. 2016 Sep 25;19(11):1372–85.

13. Dorset EE, Sakaluk SK, Thompson CF. Behavioral Plasticity in Response to Perceived Predation Risk in Breeding House Wrens. Evol Biol. 2017 Jun;44(2):227–39.

14. DuRant SE, Willson JD, Carroll RB. Parental Effects and Climate Change: Will Avian Incubation Behavior Shield Embryos from Increasing Environmental Temperatures? Integrative and Comparative Biology [Internet]. 2019 Jun 5 [cited 2019 Oct 13]; Available from: https://academic.oup.com/icb/advance-article/doi/10.1093/icb/icz083/5511833

15. DuRant SE, Hopkins WA, Hepp GR, Walters JR. Ecological, evolutionary, and conservation
implications of incubation temperature-dependent phenotypes in birds: Incubation temperature
and avian offspring phenotype. Biological Reviews. 2013 May;88(2):499–509.

16. Hepp GR, Kennamer RA, Johnson MH. Maternal effects in Wood Ducks: incubation temperature
influences incubation period and neonate phenotype. Funct Ecology. 2006 Apr;20(2):308–14.

17. Nord A, Sandell MI, Nilsson JÅ. Female zebra finches compromise clutch temperature in
energetically demanding incubation conditions: Incubation metabolic rate and clutch temperature.
Functional Ecology. 2010 Oct;24(5):1031–6.

18. Sultan SE. Organism and environment: ecological development, niche construction, and adaptation.
USA: Oxford University Press; 2015.

19. Simeone A, Luna-Jorquera G, Wilson R. Seasonal variations in the behavioural thermoregulation of
roosting Humboldt penguins (Spheniscus humboldti) in north-central Chile. JOURNAL OF
ORNITHOLOGY. 2004 Jan;145(1):35–40.

20. McDiarmid CS, Naguib M, Griffith SC. Calling in the heat: the zebra finch "incubation call" depends
on heat but not reproductive stage. BEHAVIORAL ECOLOGY. 2018 Dec;29(6):1245–54.

21. Nelson SD, Keever AC, Wightman PH, Bakner NW, Argabright CM, Byrne ME, et al. Fine-scale
resource selection and behavioral tradeoffs of eastern wild turkey broods. JOURNAL OF WILDLIFE
MANAGEMENT. 2022 Jul;86(5).

22. Bonamour S, Chevin LM, Reale D, Teplitsky C, Charmantier A. Age-dependent phenological plasticity
in a wild bird. JOURNAL OF ANIMAL ECOLOGY. 2020 Nov;89(11):2733–41.

23. Simmonds EG, Sheldon BC, Coulson T, Cole EF. Incubation behavior adjustments, driven by ambient
temperature variation, improve synchrony between hatch dates and caterpillar peak in a wild bird
population. ECOLOGY AND EVOLUTION. 2017 Nov;7(22):9415–25.

24. Kearney MR, Matzelle A, Helmuth B. Biomechanics meets the ecological niche: the importance of
temporal data resolution. Journal of Experimental Biology. 2012 Apr 15;215(8):1422–4.

25. Cones AG, Crowley PH. Optimal maternal incubation strategies for altricial and precocial birds.
Ecological Modelling. 2020 Nov;436:109290.

26. Diez-Mendez D, Cooper CB, Jose Sanz J, Verdejo J, Barba E. Deconstructing incubation behaviour in
response to ambient temperature over different timescales. JOURNAL OF AVIAN BIOLOGY. 2021
Jul;52(7).

27. Smith MG, Kaiser SA, Sillett TS, Webster MS. Variation in nest characteristics and brooding patterns
of female Black-throated Blue Warblers is associated with thermal cues. AUK. 2018 Jul;135(3):733–
47.

28. Conway CJ. Effects of ambient temperature on avian incubation behavior. Behavioral Ecology. 2000
Mar 1;11(2):178–88.

29. Nowakowski AJ, Frishkoff LO, Agha M, Todd BD, Scheffers BR. Changing Thermal Landscapes:
Merging Climate Science and Landscape Ecology through Thermal Biology. Curr Landscape Ecol Rep.
2018 Dec;3(4):57–72.

30. Carroll RL, Davis CA, Fuhlendorf SD, Elmore RD, DuRant SE, Carroll JM. Avian parental behavior and
nest success influenced by temperature fluctuations. Journal of Thermal Biology. 2018 May;74:140–
8.

31. AlRashidi M, Kosztolányi A, Küpper C, Cuthill IC, Javed S, Székely T. The influence of a hot
environment on parental cooperation of a ground-nesting shorebird, the Kentish plover Charadrius
alexandrinus. Front Zool. 2010;7(1):1.

32. van de Ven TMFN, McKechnie AE, Cunningham SJ. The costs of keeping cool: behavioural trade-offs
between foraging and thermoregulation are associated with significant mass losses in an arid-zone
bird. Oecologia. 2019 Sep 1;191(1):205–15.

33. Nord A, Nilsson JÅ. Long-term consequences of high incubation temperature in a wild bird
population. Biol Lett. 2016 Apr;12(4):20160087.

34. Boulton RL, Richard Y, Armstrong DP. The Effect of Male Incubation Feeding, Food and Temperature
on the Incubation Behaviour of New Zealand Robins. ETHOLOGY. 2010 Jun;116(6):490–7.

35. Deeming C. Avian incubation: behaviour, environment and evolution. Oxford University Press; 2002.

36. Fischer K, Karl I. Exploring plastic and genetic responses to temperature variation using copper
butterflies. Clim Res. 2010 Aug 5;43(1):17–30.

37. Huey RB, Kearney MR, Krockenberger A, Holtum JAM, Jess M, Williams SE. Predicting organismal
vulnerability to climate warming: roles of behaviour, physiology and adaptation. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2012 Jun 19;367(1596):1665–79.

38. Turriago JL, Tejedo M, Hoyos JM, Bernal MH. The effect of thermal microenvironment in upper
thermal tolerance plasticity in tropical tadpoles. Implications for vulnerability to climate warming. J
Exp Zool Pt A. 2022 Aug;337(7):746–59.

39. Sheldon KS. Climate Change in the Tropics: Ecological and Evolutionary Responses at Low Latitudes.
Annu Rev Ecol Evol Syst. 2019 Nov 2;50(1):303–33.

40. Cohen JM, Fink D, Zuckerberg B. Extreme winter weather disrupts bird occurrence and abundance
patterns at geographic scales. Ecography. 2021 Aug;44(8):1143–55.

41. Harrod SE, Rolland V. Demographic responses of eastern bluebirds to climatic variability in
northeastern Arkansas. Population Ecology. 2020 Jul;62(3):317–31.

42. WALSBERG G. THERMAL CONSEQUENCES OF DIURNAL MICROHABITAT SELECTION IN A SMALL BIRD.
ORNIS SCANDINAVICA. 1993 Sep;24(3):174–82.

43. Hawkins WD, DuRant SE. Applications of machine learning in behavioral ecology: Quantifying avian
incubation behavior and nest conditions in relation to environmental temperature. Bouffanais R,
editor. PLOS ONE. 2020 Aug 28;15(8):e0236925.

44. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R
Foundation for Statistical Computing; 2021. Available from: https://www.R-project.org/

45. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using lme4 [Internet].
arXiv; 2014 [cited 2022 Dec 14]. Available from: http://arxiv.org/abs/1406.5823

46. Bartoń K. Package 'MuMIn.' 2022 Oct 12;1(18):79.

47. Burnham KP, Anderson DR, Huyvaert KP. AIC model selection and multimodel inference in
behavioral ecology: some background, observations, and comparisons. Behav Ecol Sociobiol. 2011
Jan;65(1):23–35.

48. Grubbauer P, Hoi H. Female penduline tits (Remiz pendulinus) choosing high quality nests benefit by
decreased incubation effort and increased hatching success. ECOSCIENCE. 1996;3(3):274–9.

49. Taff CC, Freeman-Gallant CR. Female ornamentation, incubation behavior, and reproductive success
in a wild bird. BEHAVIORAL ECOLOGY AND SOCIOBIOLOGY. 2021 Jun;75(6).

50. MacDonald EC, Camfield AF, Jankowski JE, Martin K. An alpine-breeding songbird can adjust dawn
incubation rhythms to annual thermal regimes. AUK. 2014 Oct;131(4):495–506.

51. Zhang L, An B, Shu M, Zhao C, Yang X, Suo Y, et al. Incubation strategies of the Black-necked Crane
(Grus nigricollis) in relation to ambient temperature and time of day. AVIAN RESEARCH. 2017 Jul
20;8.

52. Husby A, Nussey DH, Visser ME, Wilson AJ, Sheldon BC, Kruuk LEB. CONTRASTING PATTERNS OF
PHENOTYPIC PLASTICITY IN REPRODUCTIVE TRAITS IN TWO GREAT TIT (PARUS MAJOR)
POPULATIONS. EVOLUTION. 2010 Aug;64(8):2221–37.

53. Le Vaillant J, Potti J, Camacho C, Canal D, Martinez-Padilla J. Fluctuating selection driven by global
and local climatic conditions leads to stasis in breeding time in a migratory bird. JOURNAL OF
EVOLUTIONARY BIOLOGY. 2021 Oct;34(10):1541–53.

54. Snowden R. Least Tern Nest Thermoregulation in Response to Environmental Temperature and
Human Disturbance [Doctoral]. University of North Carolina Wilmington; 2018.

55. Andes AK, Sherfy MH, Shaffer TL, Ellis-Felege SN. Plasticity of Least Tern and Piping Plover nesting
behaviors in response to sand temperature. JOURNAL OF THERMAL BIOLOGY. 2020 Jul;91.

56. Garcia RA, Allen JL, Clusella-Trullas S. Rethinking the scale and formulation of indices assessing
organism vulnerability to warmer habitats. Ecography. 2019 May;42(5):1024–36.

57. Brennan LA, Hernandez F, Williford D. Northern Bobwhite (Colinus virginianus). In: Billerman SM,
Keeney BK, Rodewald PG, Schulenberg TS, editors. Birds of the World [Internet]. Cornell Lab of

Ornithology; 2020 [cited 2022 Dec 14]. Available from:
https://birdsoftheworld.org/bow/species/norbob/1.0/introduction

58. Dabbert CB, Pleasant G, Schemnitz SD. Scaled Quail (Callipepla squamata). In: Billerman SM, Keeney
BK, Rodewald PG, Schulenberg TS, editors. Birds of the World [Internet]. Cornell Lab of Ornithology;
2020 [cited 2022 Dec 14]. Available from:
https://birdsoftheworld.org/bow/species/scaqua/1.0/introduction

59. Rohwer VG, Purcell JR. Geographic variation in incubation behavior of a widely distributed passerine
bird. Naya DE, editor. PLoS ONE. 2019 Aug 14;14(8):e0219907.

60. Grayson KL, Johnson DM. Novel insights on population and range edge dynamics using an
unparalleled spatiotemporal record of species invasion. Woodcock B, editor. J Anim Ecol. 2018
May;87(3):581–93.

61. Kirkpatrick WH, Sheldon KS. Experimental increases in temperature mean and variance alter
reproductive behaviours in the dung beetle Phanaeus vindex. Biology Letters. 2022;18:20220109.

62. Mamantov MA, Sheldon KS. Behavioural responses to warming differentially impact survival in
introduced and native dung beetles. Behmer S, editor. Journal of Animal Ecology. 2021
Jan;90(1):273–81.

63. Telemeco RS, Elphick MJ, Shine R. Nesting lizards (*Bassiana duperreyi*) compensate partly, but not
completely, for climate change. Ecology. 2009 Jan;90(1):17–22.

64. Sudnick M, Brodie BS, Williams KA. Nature versus nurture: Structural equation modeling indicates
that parental care does not mitigate consequences of poor environmental conditions in Eastern
Bluebirds (*Sialia sialis*). Ecol Evol. 2021 Nov;11(21):15237–48.

65. Clusella-Trullas S, Garcia RA, Terblanche JS, Hoffmann AA. How useful are thermal vulnerability
indices? Trends in Ecology & Evolution. 2021 Nov;36(11):1000–10.

High-Resolution Best Fit Models						
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AIC$_c$**	**df**	**AIC$_c$ w$_i$**
High Duration	Off-bout Duration	2hr Mean, 2 hr SD, AMPM, Species	2hr Mean:Species, 2hr SD:Species, AMPM:Species	8952.7	11	0.742
Null	Off-bout Duration	~	~	9154.8	4	< 0.0001
High AM Start Time	Off-bout Timing - AM	2hr Mean, 2hr SD, Species	~	1375.2	7	0.429
Null	Off-bout Timing - AM	~	~	1417.8	4	< 0.001
High PM Start Time	Off-bout Timing - PM	2hr Mean, 2hr SD	~	1580.9	6	0.425
Null	Off-bout Timing - PM	~	~	1654.8	4	< 0.001
Low-Resolution Best Fit Models						
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AIC$_c$**	**df**	**AIC$_c$ w$_i$**
Low Duration	Daily Off-bout Duration	24hr SD, Morning SD, Previous Night SD, Species	24hr SD:Species, Previous Night SD:Species	5666.7	10	0.02
Null	Daily Off-bout Duration	~	~	5706	4	< 0.001
Low AM Start Time	Off-bout Timing - AM	Previous Day SD, Species	~	1378.9	6	0.075
Null	Off-bout Timing - AM	~	~	1395.6	4	< 0.0001
Low PM Start Time	Off-bout Timing - PM	Afternoon Mean, Species	~	1349.7	6	0.22
Null	Off-bout Timing - PM	~	~	1374.4	4	< 0.0001