Title: Critical reproductive behaviors in Scaled Quail and Northern Bobwhite are affected by
thermal variability and mean temperature

Authors: William Kirkpatrick[1], Erin Sauer[1], Rachel Carroll[2], Jeremy Cohen[3,4], Craig Davis[5],
Samuel Fuhlendorf[5], Sarah DuRant[1]

Affiliation: [1]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[2]Jones College of Science, Engineering, and Technology, Murray State University, Murray, KY,
USA

[3]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA

[4]Center for Biodiversity and Global Change, Yale University, New Haven, CT, USA

[5]Department of Natural Resource Ecology and Management, Oklahoma State University,
Stillwater, OK, USA

Corresponding Author: William Kirkpatrick[1], whkirk97@gmail.com,

2560 Iron Ave, Farmington, AR 72730

Abstract

Animals can respond differently to shifting thermal variability versus thermal averages,
both of which are changing due to climate warming. How these thermal variables affect parental
care behaviors can reveal the ability of parents to modify their behaviors to meet the competing
demands of their offspring's thermal needs and self-maintenance, which becomes critical in
suboptimal thermal conditions. Further, the time frame used to examine the interplay between
temperature and behavioral shifts (e.g., seasonal patterns in care vs. drivers of individual care
decisions) can provide different information about the plasticity of parental care behavior. We
investigated the relationship between thermal means, thermal variability, and incubation
behaviors across multiple timescales in Scaled Quail and Northern Bobwhite. Both species
decreased off-bout length during periods of high thermal variability, a novel finding among
studies of avian parental behavior. Further relationships between thermal endpoints (mean vs.
variation) and behavior differed depending on the temporal scale. For instance, total daily time
spent off the nest was not influenced by daily average temperature, yet individual off-bout
duration increased with increasing average temperature in the two hours prior to the off-bout.
These results provide evidence that thermal-behavioral relationships differ across scales and
likely represent a bird's ability to modify their incubation strategy to rapidly respond to the
immediate thermal environment (altering individual off-bout length based on temperature) to
meet self-maintenance needs while resulting in a similar outcome for their nest (total daily off-
bout time). However, longer off-bout durations during high temperature events can come with
reproductive costs, sometimes resulting in acute offspring mortality when eggs or chicks
experience lethal temperatures.

Keywords: environmental temperature, thermal variation, parental behavior, *Colinus virginianus,*
Callipepla squamata, quail

Introduction

Moderate projections of anthropogenic climate warming estimate an average global temperature increase of 1.1 – 2.6°C by 2100 (IPCC, 2023) with projected losses of 7.9% of all species due to climate change (Urban, 2015). However, mean increases in temperature alone will not reflect the shifts in thermal conditions that organisms will experience (Denny, 2017; Jensen, 1906; Ruel and Ayres, 1999). Increases in thermal variability are occurring alongside warming average temperatures (Bathiany et al., 2018; Easterling et al., 2000), and evidence in ectotherms shows that animals modify their behavior when exposed to increased environmental variation but not average conditions (Kirkpatrick and Sheldon, 2022; Sheldon and Dillon, 2016; Vasseur et al., 2014). The nature of the relationship between organismal behavior and environmental temperature across genera is complex and often nonlinear (Conway and Martin, 2000), requiring careful contextualization with consideration of multiple aspects of the thermal environment.

When wildlife are exposed to novel thermal conditions, plastic behaviors improve fitness outcomes. This plasticity can have multiple layers of fitness benefits or consequences when the responding behavior is related to parental care, because parental care behaviors have implications for parental and offspring fitness, as well as offspring phenotypic development and plasticity. Behaviors employed by parents to care for their offspring (i.e., parental care behaviors) are known to be plastic (Andes et al., 2020; Dorset et al., 2017) and play an important role in shaping the thermal environment of developing offspring (DuRant et al., 2019). Ambient thermal conditions can alter parental care behaviors (Grimaudo et al., 2020; Masero et al., 2012), subsequently modifying the thermal environment of the developing offspring to directly influence offspring phenotypes (DuRant et al., 2013; Hepp et al., 2006). Examining shifts in parental care behaviors of wild organisms in response to the natural thermal environment can reveal the ability of parents to modify their behaviors to meet the competing demands of offspring care and self-maintenance (McDiarmid et al., 2018; Nelson et al., 2022; Nord et al., 2010; Sharpe et al., 2021; Simeone et al., 2004; Sultan, 2015).

Birds in hot environments must invest in self-maintenance behaviors such as gular fluttering or retreating to cooler microclimates to avoid physiological stress (AlRashidi et al., 2010; van de Ven et al., 2019). However, coping behaviors that cause the parent to leave the nest can come at the cost of offspring care, as the nest is left exposed to external thermal conditions (Carroll et al., 2018; DuRant et al., 2019, 2013; Sharpe et al., 2019). The timing of off-bouts (the time a parent spends off the nest) and their duration play important roles in the development of avian offspring (Cooper and Voss, 2013; Smith et al., 2018). Lengthening and, in some cases, delaying off-bouts can lead to higher rates of predation (Conway and Martin, 2000) or thermal stress, reducing adult and offspring fitness (Nord and Nilsson, 2016; Sharpe et al., 2022).We know little of the scale at which these decisions are being made and their relationship to natural thermal variation (Boulton et al., 2010; Deeming, 2002), but the importance of parental care on offspring development suggests that reproductive success would benefit from the ability of parents to delay off-bouts until temperatures are outside lethal levels for embryos (Smith et al., 2018). Largely, studies focusing on avian behavioral responses to thermal change in temperate climates focus on discrete thermoregulatory behaviors (McDiarmid et al., 2018; Nelson et al.,

2022; Walsberg, 1993) or phenological adjustments to rising average temperatures associated
with seasonal shifts in the thermal environment (Cohen et al., 2021; Harrod and Rolland, 2020)
(McDiarmid et al., 2018; Nelson et al., 2022; Walsberg, 1993). Published data typically show
that parents shorten off-bout duration when they are experiencing hot temperatures (Matysioková
and Remeš, 2018; Rohwer and Purcell, 2019). The ability of parents to alter their behavior will
depend on their ability to withstand physiologically demanding temperatures and to respond to
thermal conditions as they are occurring, thus studies examining thermal and behavioral
relationships at varying temporal scales are necessary, e.g. acute responses to immediate
conditions and broad patterns of behavior across seasonal, monthly or daily scales.

Understanding how the real-time thermal environment (temperatures experienced
immediately before exhibition of the observed behavior) influences avian parental behavior is
critical for understanding what drives parents to initiate biologically relevant behaviors (Kearney
et al., 2012). Long-term (> 24 hours or a single diel cycle) thermal data is useful for exploring
seasonal or daily patterns in behavior (Bonamour et al., 2020; Cones and Crowley, 2020; Martin
et al., 2018; Simmonds et al., 2017). For instance, greater than average temperature over the
course of a reproductive season in birds (i.e., low-resolution thermal and behavioral data) is
associated with shorter incubation periods (Capp et al., 2018; Diez-Mendez et al., 2021; Smith et
al., 2018), possibly indicating a reduced cost of reproduction for parents. However, assessing
only average temperature over the entire incubation stage obscures variance in the thermal
environment occurring during this period. During an acute period of hot temperatures, parents
may increase nest attendance to shield eggs or seek refuge, leaving eggs exposed to a suboptimal
thermal environment. The relationship between thermal variance and incubation behaviors may
also vary with scale because thermal variance experienced just before an off-bout may be
indicative of thermal stability or rapidly changing temperatures (i.e., rate of temperature change;
Sharpe et al., 2021). Temperature variation experienced over a day or entire incubation period,
however, may represent the breadth of thermal challenges parents navigated (i.e. the repertoire of
behavioral and physiological mechanisms needed to thermoregulate above and below the thermal
neutral zone). It is not known how thermal variation shapes incubation behaviors in birds, but
acclimation to novel thermal variation and short-term thermal conditions has been evidenced in
ectotherms (Fischer and Karl, 2010; Huey et al., 2012; Turriago et al., 2022). What is known
about how real-time or recently experienced temperatures shape isolated behavioral responses in
birds is mixed, with some species exhibiting reduced off-bout duration in response to increased
ambient temperature (Skrade and Dinsmore, 2012; Zhang et al., 2017) and others showing the
opposite relationship (Bourne et al., 2021; Caldwell and Cornwell, 1975; Croston et al., 2020;
Ringelman et al., 1982). Exploring the effects of acute thermal experiences on variation in avian
behavior can inform the mechanisms behind large scale changes in species' ranges, phenotypic
changes, and fitness (Nowakowski et al., 2018).
Here, we describe the variation in the length and timing of off-bout behaviors in response
to local thermal conditions experienced throughout the day or immediately prior to the off-bout
for two species: Scaled and Northern Bobwhite Quail. We opportunistically used existing data
collected in 2015 and 2016 and described in Carroll *et al*. 2018, to begin exploring if both
average temperatures and thermal variability experienced by parents shape incubation behaviors.
Scaled Quail occupy arid shrublands and prairies from northern Mexico to southern Colorado,

USA (Dabbert et al., 2020) while Bobwhite Quail naturally occur in temperate open or early-
succession forests and prairies in the eastern USA and Mexico (Brennan et al., 2020). Both of
these precocial New World Quail species (family – Ontophoridae) are female-only incubators,
but males have been observed in multiple species of the family assisting late in the incubation
period (Carroll, 1994). Scaled Quail (*Callipepla squamata*) and Bobwhite Quail (*Colinus
virginianus*) overlap in their ranges in western Oklahoma, USA. At range edges, climatic
patterns will determine the future of the species in response to climate warming (Rehm et al.,
2015), and while the limited overlap in range edges in our species suggests evolutionary
distinction, both species have exhibited reduced breeding in response to climatic trends (Carroll
et al., 2018; Errington, 1945; Lusk et al., 2006; Pleasant et al., 2006). Nest construction is similar
in both species, as both are ground nesters that occupy low-growing trees and shrubs in their
shared habitat, though observations suggest that Scaled Quail select nesting locations with more
favorable vegetation for thermal buffering than Northern Bobwhite (8.2℃ versus 5.7℃ cooler
nests on average than random locales near the nest, respectively – Carroll et al., 2018).

We hypothesize that: 1) Increased average temperature will lead to increased off-bout
duration regardless of temporal scale and species, and 2) thermal variability will operate as a
distinct predictor of avian behavior. As mentioned above, little is known about environmental
thermal variability as a predictor of behavior, but we expect that parents will likely avoid leaving
the nest as often or as long in highly variable conditions to reduce the risk of exposing the nest to
temperatures that could quickly become too hot or cold for optimal embryo development.
Because parental incubation behaviors can have serious consequences for eggs, we wanted to
explore whether temperatures eggs experienced when parents were away from the nest were
associated with nest success. Therefore, we also examined nest success (at least 1 egg hatching)
in relation to exposure to lethal temperatures (temperatures above 40 ℃ . We predicted that
failed nests would be more likely to experience lethal temperatures and for longer periods of
time.

Methods

Monitoring Temperature and Incubation Patterns

All nests (37 Bobwhite and 24 Scaled Quail) were monitored in the Beaver River
Wildlife Management Area in Oklahoma, USA (36°50′21″N, 100°42′15″W). Environmental
temperature data was collected every 30 seconds within and outside the nest using modified
Onset Computer HOBOware (Onset Corporation, Borne, MA) thermal sensors. External
temperatures were taken with HOBO U23 Pro v2 External Temperature Loggers (Carroll et al.,
2018). The internal temperature was taken using egg shaped clay models inserted into the nest
that matched the dimensions of quail eggs. Each model egg was inserted into the center of the
nest and external sensor was placed within 2m of the nest for standardization of thermal
measurements. More details of thermal probes and data collection can be found in Carroll *et al.*
2018. Though thermal-behavioral relationships do change depending on humidity, our field site
humidity was not measured at the same frequency or microscale as temperature, so we cannot
incorporate it relevantly into our study. According to weather station data collected at a nearby

meteorological station (NOAA Station US1OKBV0005 – Forgan, OK, USA), precipitation
during the summer was exceedingly rare (and only 9 cm average precipitation on those days).

The machine learning software NestIQ was used to determine off-bout length and timing
from the raw nest temperature data (Hawkins and DuRant, 2020). To assess the impact the rate
of thermal change has on off-bout duration, we calculated the average and standard deviation of
environmental temperature measurements during the two hours before each off-bout. We chose
temperatures 2 hours prior to the off-bout to capture a window of thermal temperature long
enough to assess a meaningful measure of thermal conditions and stability that could drive an
incubation decision without overlapping with longer term diel patterns. To assess the impact the
daily thermal environment has on total daily off-bout, we calculated average and standard
deviation of temperatures from the morning (sunrise to noon), afternoon (noon to sunset), and
night (previous day sunset to sunrise) as well as the temperatures from the previous day and
previous afternoon. We included analyses of the temperatures experienced in the previous day,
because those conditions could affect incubation decisions by contributing to the parent's
physiological ability to withstand current conditions. We chose these deterministic timeframes to
represent a longer-term window of ambient thermal conditions relative to the local diel cycle.
We considered including maximum and minimum temperatures from each timeframe, but they
both correlated highly with thermal variation (>0.9 and -0.9 respectively) and were excluded. All
sunrise and sunset times were recorded for each day of the summers of 2015 and 2016 at the
field site and verified through local weather records.

Statistical Methodology

All statistical analyses were conducted with R version 4.1.2 (R Core Team, 2023). To test
the effect of the mean and standard deviation of temperature during the two-hour period before
each off-bout on off-bout duration, we developed a linear mixed model (Bates et al., 2014). The
universal model included mean and standard deviation of the two-hour period preceding the off-
bout as predictors and off-bout length (in minutes) as the response variable. We included a two-
level categorical predictor for whether the off-bout took place in the morning or afternoon, as
each species typically takes two off-bouts per day, in the early morning or late afternoon
(referred to as AMPM in models). We also included species (Scaled Quail or Northern
Bobwhite). To control for non-independence of the measurements at a single nest and year, we
included individual nest ID, year, and date as random intercepts. All possible interactions
between predictors were included in the universal model. To test for relationships between
temperature and off-bout start time, we conducted a separate linear mixed model with the same
predictors, interactions, and random intercepts described above but with off-bout timing as the
response variable (Table 1).

To address the effect of temperatures experienced over a longer time period on total daily
off-bout, we developed a linear mixed effects model which included the sum duration of all off-
bouts in 24 hours as the response variable. We selected total daily off-bouts as our response
variable because the predictors are date-specific thermal measurements instead of off-bout
specific thermal measurements. The predictive variables included the mean and standard
deviation of external temperature from the following timeframes relative to the date: 24hr

(temperatures from 12AM-11:59PM of the current day), Morning (sunrise to 12PM), Afternoon
(12PM to sunset), Previous Night (sunset to sunrise of previous night), Previous Day (12AM to
11:59PM of the previous day), and Previous Afternoon (12PM to sunset of the previous day).
We chose these time periods to allow for comparison of longer-term trends in temperature.
While they are likely highly correlated with one another, our model selection strategy (see below
– *Model Selection*) corrected for this issue. We included a two-level categorical predictor for
species (Scaled or Northern Bobwhite). To control for non-independence of the nest location and
year, we included individual Nest ID and year as random intercepts. To test for relationships
between mean and standard deviation of temperature on the average daily time of each off-bout,
we conducted two separate linear mixed models with the same predictors, interactions, and
random intercepts described above, but the response variables were the timing of the morning or
afternoon off-bout respectively, given that each species takes at most, two off-bouts during the
day in the morning, evening, or both.

Model Selection

For each universal mixed-effects model (high and low-resolution), we used the dredge
function from the MuMIn package for model selection (Bartoń, 2022). We chose the best fit
model by finding the model with the lowest relative AICc score (Burnham et al., 2011). All
values represented in our figures are partial residual plots generated using the predict function in
R.

The best fit model for the high-resolution approach to analyze individual off-bout
duration (referred to in Results as Individual Duration) included 2-hour mean temperature, 2-
hour standard deviation of temperature, morning or afternoon (AMPM), and species as main
effects with three interactive effects: 2-hour mean*species, 2-hour standard deviation of
temperature*species, and AMPM*species (Table 1).

The best fit model for predicting total daily off-bout (referred to in Results as Total
Duration) included 24 hour and morning standard deviation of temperature as main effects;
standard deviation of morning temperatures was removed due to a high correlation with 24 hour
variation ($r^2 > 0.6$) and lack of a significant effect in the general model. Species and two
interactive effects were also included: species*morning standard deviation of temperature and
species*previous night standard deviation of temperature (Table 1).

Two models were used to predict daily off-bout start time, one that included morning off-
bouts (referred to in Results as Morning Start Time) and one that included afternoon off-bouts
since each species takes two off-bouts per day in either the morning or evening. The best fit
model for the afternoon off-bouts (Total PM Start Time) included mean temperature during the
afternoon and species with no interactive effects (Table 1).

Conditional Model Residuals

Behavioral responses to temperature in wildlife are often nonlinear, especially in extreme
thermal scenarios (Denny, 2017). By examining the conditional residuals of our fitted models,
we can determine linearity and heteroscedasticity of the relationship observed (Santos Nobre and
da Motta Singer, 2007). For the model Individual Duration, the residuals were largely linear,

with a slight deviation above expected observations as residual values increased (Supplementary
Figure 2). However, non-linear and/or heteroscedastic relationships were noted when examining
timing of off-bouts on local temporal scales. Models predicting high-resolution off-bout start
time and morning off-bouts did not meet the necessary assumptions. Therefore, we only
examined models that met the requirements for heteroscedasticity and non-linearity (see
Supplementary Materials to reference conditional residual plots).

Hatching Success

We compared cumulative minutes exposed to temperatures above 40°C in hatched vs.
failed nests. We also constructed a similar model with days exposed to temperatures above 40°C
to see whether failed nests were exposed more frequently to lethal temperatures. To determine
the effect of time exposed to extremely warm conditions, we constructed two one-way
ANOVA's with either total minutes above 40°C experienced or days above 40°C. The models
included nest hatching or failure and species as main effects. Interactive effects of species and
nest failure were also included.

High-Resolution Best Fit Models						
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AICc**	**df**	**AICc wi**
Individual Duration	Off-bout Duration	2hr Mean, 2 hr SD, AMPM, Species	2hr Mean: Species, 2hr SD: Species, AMPM: Species	8940	12	0.80
Null	Off-bout Duration	~	~	9156	5	1.2e-47

Low-Resolution Best Fit Models						
Model Name	**Response Variable**	**Main Effect(s)**	**Interactive Effects**	**AICc**	**df**	**AICc wi**
Total Duration	Daily Off-bout Duration	24hr SD, Morning SD, Species	24hr SD: Species, Previous Night SD: Species	5666	10	0.02
Null	Daily Off-bout Duration	~	~	5706	4	6.2e-11
Morning Start Time	Off-bout Timing - AM	Previous Day SD, Species	~	1367	7	0.12
Null	Off-bout Timing - AM	~	~	1387	5	5.8e-0.6

Table 1. The best fit models for each suite of thermal predictors as selected by AICc using the
package MuMIn. High-resolution: The universal model included mean and standard deviation of
the two-hour period preceding the off-bout as predictors and off-bout length (in minutes) as the
response variable. We included a two-level categorical predictor for whether the off-bout took
place in the morning or afternoon (referred to as AMPM) and species (Scaled Quail or Northern
Bobwhite). Low-resolution: All models were structured as above except with averaged thermal
data over 6 deterministic timeframes as predictors: 24hr (temperatures from 12AM-11:59PM of
the current day), Morning (sunrise to 12PM), Afternoon (12PM to sunset), Previous Night
(sunset to sunrise of previous night), Previous Day (12AM to 11:59PM of the previous day), and
Previous Afternoon (12PM to sunset of the previous day)). We included individual nest ID, year,

and date as random intercepts in all models. All possible interactions between predictors were
included in each universal model.

Results

High-Resolution Thermal Data – Duration

As described by model Individual Duration, off-bout duration increased as mean
temperature increased (β = 4.64 ± 0.46, t = 10, p < 0.001; Figure 1A). However, in response to
increased thermal variation, lengths of individual off-bouts decreased (β = -4.68 ± 2.11, t = -2.2,
p = 0.03; Figure 1B).

Northern Bobwhite averaged 140 ± 68 minutes per off-bout while Scaled Quail averaged
110 ± 57 minutes (β = 62 ± 14.7, t = 4.2, p < 0.001). Northern Bobwhite exhibited longer
afternoon off-bouts (159 ± 61 minutes) than Scaled Quail (Afternoon =108 ± 48 minutes; β = -
43.1 ± 10.5, t = -4.1, p > 0.001) even when mean temperatures 2 hours preceding the off-bout
were high (β = -1.8 ± 0.7, t = -2.7, p = 0.007, Supplementary Figure 2B). For both species,
morning off-bouts were not significantly longer than afternoon off-bouts, though they were
different (β = 1.5 ± 7.4, t = 0.2, p = 0.8; Morning = 110 ± 66 minutes; Afternoon = 138 ± 61
minutes).

Figure 1. Partial residual plots showing the relationships between the thermal environment
immediately preceding individual off-bouts and off-bout duration. Blue lines exhibit the
association across both years while black and gray lines represent 2015 and 2016, respectively.
Open circles represent quail nests in 2015, and closed circles represent 2016 nests. A) As mean
temperature in the 2 hours preceding each off-bout increased, Northern Bobwhite (green) and
Scaled Quail (blue) off-bout duration increased. Nest sites in 2015 were cooler on average than
2016 (2015 = 21.5 C, 2016 = 31.2 C). B) As standard deviation of temperature increased,
individual off-bout duration decreased. All points are predicted model values.

Low-Resolution Thermal Data – Total Daily Off-Bout

As described by model Total Duration, as thermal variability of the total day (24 hour
standard deviation) increased, daily off-bouts lengthened ($\beta = 11.9 \pm 3.7$, $t = 3.2$, $p = 0.001$;
Figure 2A). Alternatively, thermal variation of the previous night was selected for the model but
was not a significant predictor of total daily off-bout ($\beta = 5.6 \pm 6.4$, $t = 0.9$, $p = 0.38$). Scaled
Quail spent less total time off the nest daily (156 ± 74 minutes) than Northern Bobwhite ($197.2 \pm$
81 minutes; $\beta = 71 \pm 34$, $t = 2.1$, $p = 0.03$).

Low-Resolution Thermal Data – Start Time

As described by model Morning Start Time, increasing thermal variation of the previous
day advanced the initiation of the following day's off-bouts into the cooler early morning ($\beta = -$
0.16 ± 0.03, $t = -4.9$, $p < 0.001$; Figure 2B)

Figure 2. Partial residual plots showing the relationships between the thermal environment
measured over selected timeframes, total daily off-bout duration (A), and morning off-bout
initiation (start time) (B). A) As thermal variation increased, both species reduced their total
daily off-bout duration. B) Thermal variation of air temperature of the day preceding the
morning off-bout was a significant negative predictor of start time, advancing off-bout initiation.
Open circles represent quail nests in 2015, and closed circles represent 2016 nests.

Hatching Success

Both species constructed nests in locations that experienced temperatures above the
critical thermal maxima for eggs (~40°C and above). Our analysis revealed a marginal difference
in total minutes exposed to 40°C temperatures in failed vs. hatched nests (one-way ANOVA; *df*
$= 1,31$, $F = 4.1$, $p = 0.051$). For both species, failed nests experienced 40°C+ for 98 ± 133 (mean
$\pm$ SD) minutes compared to 30 ± 56 minutes in successful nests. There was not a significant
effect of the number of days exposed to temperatures > 40°C or species, though failed Northern
Bobwhite nests were typically exposed to less time above the threshold than Scaled Quail nests
(88 minutes ± 140 and 111 ± 135 minutes respectively).

Discussion

We investigated the relationship between average thermal conditions, thermal variability, and incubation behaviors across multiple timescales in Scaled Quail and Northern Bobwhite. We provide evidence that both species reduced time on the nest following short periods of increased thermal variability (i.e. at high rates of temperature change), a novel finding among studies of avian parental behavior. Further, relationships between thermal endpoints (mean vs. variation) and behavior differed depending on scale, in other words, whether we examined environmental temperature just prior to an individual off-bout or thermal conditions across the course of a day and daily incubation patterns. These results provide evidence that the thermal factors shaping discrete behavioral decisions may be washed out when assessing general patterns of behavior over longer time scales (Garcia et al. 2019) and both are important to consider when estimating effects of climate change on reproductive patterns.

Northern Bobwhite and Scaled Quail exhibited plastic behavioral responses to recently experienced temperatures. Both species shortened off-bouts in response to increasing thermal variability (i.e. higher rate of temperature change; Figure 1B). Similarly, total time away from the nest during a 24 hour period decreased with increasing daily thermal variability (Figure 2A, B). Alternatively, increases in recently experienced average temperatures caused parents to increase the length of an individual off-bout (Figure 1A), but daily average temperatures were not important to the total time spent away from the nest in a 24 hour period. These findings demonstrate that thermal variability can act independently of average temperature to shape an important reproductive behavior in birds and the directionality of these relationships can vary with scale. Presumably, birds shortened the length of an individual off-bout with increasing thermal variability because temperature was rapidly changing or unpredictable and could quickly reach lethal temperatures or temperatures that would slow embryonic development. Our study cannot distinguish between decisions driven by parental self-maintenance needs and offspring requirements, so shorter off-bouts during high rates of temperature change could also benefit parents. It is less clear why daily thermal variability would shorten total time away from the nest that day, but it may be that temperatures were changing most dramatically at the times of day that the quail would typically take an off-bout, shortly after sunrise and late afternoon. Alternatively, high daily thermal variability could generally signal that temperatures are unpredictable, causing parents to shorten all incubation breaks that day. The increased care we detected in our study is similar to findings in smallmouth bass (*Micropterus dolomieu*). Bass nesting in more variable habitats increased parental care activities compared to parents nesting in less variable but warmer habitats (Cooke et al., 2003). It was surprising that relationships between daily patterns of incubation and temperature did not match the relationship between off-bout length and the average temperature experienced just before the off-bout. This suggests that temperatures experienced at the time of an off-bout do inform incubation decisions, and these patterns will not always scale up to be reflected in daily or seasonal patterns in incubation constancy (Grubbauer and Hoi, 1996; Husby et al., 2010; Le Vaillant et al., 2021; MacDonald et al., 2014; Taff and Freeman-Gallant, 2021; Zhang et al., 2017). This finding is supported by studies of other species nesting in hot conditions. Least Tern and Piping Plover (*Charadrius melodus*; Andes et al., 2020) incubating parents exhibit some behaviors during incubation that

aid in their thermoregulation but only within certain ranges of environmental temperature and
these relationships would not be obvious when summing time away from the nest across a longer
time period. These individual incubation decisions based on current thermal conditions can have
important implications for nest success. We found that most failed nests were exposed to
temperatures exceeding 40°C on a single day in a single extreme event, this finding suggests that
a single exposure to extreme heat during an off-bout can lead to reproductive failure (and see
Cunningham et al., 2013) and may not be detectable when comparing general patterns of
incubation in relation to average temperature. In some species, there is evidence of population
collapse caused by a short exposure to extreme heat regardless of corresponding daily mean
temperature or parental responses to temperature (Ma et al., 2015).

When comparing behaviors of the two quail species, we found that both species
expressed similar trends in their behavioral modifications to thermal conditions but differed in
individual off-bout duration and temperature of their nests. Scaled Quail spent significantly less
time off the nest on average (156 minutes) than Northern Bobwhite (197 minutes). Species
specific differences in these data were explored elsewhere (Carroll et al. 2018), including this
particular result, so we discuss it here as it relates to the hypotheses we tested. It is interesting
that the two quail species differ in off-bout duration, but not in the relationship between the
responsiveness of their incubation behaviors to thermal conditions, both thermal averages and
variation. Off-bout duration and nest temperatures are likely influenced by natural history
differences of the two species (Brennan et al., 2020; Dabbert et al., 2020) and determined
primarily as a result of Scaled Quail selecting nest sites or substrates with superior thermal
buffering characteristics compared to Northern Bobwhite nests (Carroll et al., 2018). The typical
breeding habitat of Northern Bobwhite is temperate and extreme temperatures rarely reach those
experienced by Scaled Quail (Brennan et al., 2020), which are a grassland desert-dwelling
species adapted to the extreme highs and lows present in the Western Oklahoma field site
(Dabbert et al., 2020). The two species may have experienced different selection pressures for
nest site selection, but plasticity of incubation behaviors to changing thermal conditions indicate
the importance for both species to prevent exposure of nests to lethal temperatures and reduce
care costs to the parent. It should be noted that direct comparisons between each species based
on results found here need to be made with caution, as thermal and behavioral measurements
across each species' range would be required to make definitive species level comparisons in
behavioral plasticity.

Behavioral plasticity during breeding generally benefits both the parent and dependent
offspring by keeping the thermal environment stable over time (Kirkpatrick and Sheldon, 2022;
Mamantov and Sheldon, 2021; Telemeco et al., 2009), but shifts in behavior alone cannot
completely negate the impacts of poor or unexpected environmental conditions (Sharpe et al.,
2019, 2022; Sudnick et al., 2021), as time allotted for self-maintenance may take precedence
over offspring care (Deeming, 2002). Our results indicate that the rate of environmental
temperature change can directly influence parental behavior, highlighting that temperature
variability may be a key player in shaping avian reproductive behaviors. Our analysis also
indicated that discrete behavioral decisions (timing or duration of an off-bout) are made using
thermal information about current conditions, but more general patterns in incubation behaviors

(total daily time spent off the nest) were less responsive to the daily average temperature. In
other words, in these two species, parents are spending a similar amount of time off the nest each
day regardless of average daily temperature, but how that time is metered out is flexible based on
recently experienced thermal conditions and temperature stability. This finding underscores the
importance of testing for relationships between thermal conditions and behavior across multiple
timescales. A single thermal measurement taken over a defined time may provide inaccurate
assessments of organismal vulnerability to thermal change if characterized incorrectly or
incompletely (Clusella-Trullas et al., 2021; Sheldon and Dillon, 2016). Models with averaged,
lone sole-characteristic predictors should not be avoided but contextualized temporally as we do
here (Garcia et al., 2019).

CRediT Authorship Statement

William Kirkpatrick: Conceptualization, Writing - Original Draft, Writing – Editing and Review Methodology, Data Curation, Formal Analysis. Erin Sauer: Formal Analysis, Writing – Editing and Review. Rachel Carroll: Investigation, Resources, Methodology, Data Curation. Jeremy Cohen: Formal Analysis, Writing – Editing and Review. Craig Davis: Investigation, Resources, Methodology, Data Curation, Funding Acquisition. Sam Fuhlendorf: Investigation, Resources, Methodology, Data Curation, Funding Acquisition. Sarah DuRant: Conceptualization, Investigation, Resources, Methodology, Data Curation, Writing - Original Draft, Writing – Editing and Review, Supervision.

Acknowledgements

We would like to thank Rachel Carroll, Craig Davis, and Sam Fuhlendorf for permission to use their thermal data from Scaled and Northern Bobwhite Quail nests. Thanks to Wayne Hawkins for assistance with NestIQ.

Funding

This work was supported by the Oklahoma Department of Wildlife Conservation through funding administered through the Oklahoma Cooperative Fish and Wildlife Research Unit (grant #F11AF00069), the Oklahoma Agricultural Experiment Station at Oklahoma State University, and the Bollenbach and Groendyke Endowments.

Ethical Committee/Institutional Review Board (IRB) Approval

Animal welfare approval was obtained from IACUC at Oklahoma State University.

Data Accessibility Statement

All supporting data and annotated R scripts are available in GitHub or through contact with the corresponding author.

Competing Interests

We declare we have no competing interests.

Literature Cited

AlRashidi, M., Kosztolányi, A., Küpper, C., Cuthill, I.C., Javed, S., Székely, T., 2010. The influence of a hot environment on parental cooperation of a ground-nesting shorebird, the Kentish plover Charadrius alexandrinus. Front Zool 7, 1. https://doi.org/10.1186/1742-9994-7-1

Andes, A.K., Sherfy, M.H., Shaffer, T.L., Ellis-Felege, S.N., 2020. Plasticity of Least Tern and Piping Plover nesting behaviors in response to sand temperature. JOURNAL OF THERMAL BIOLOGY 91. https://doi.org/10.1016/j.jtherbio.2020.102579

Bartoń, K., 2022. Package 'MuMIn' 1, 79.

Bates, D., Mächler, M., Bolker, B., Walker, S., 2014. Fitting Linear Mixed-Effects Models using lme4.

Bathiany, S., Dakos, V., Scheffer, M., Lenton, T.M., 2018. Climate models predict increasing temperature variability in poor countries. Sci. Adv. 4, eaar5809. https://doi.org/10.1126/sciadv.aar5809

Bonamour, S., Chevin, L.-M., Reale, D., Teplitsky, C., Charmantier, A., 2020. Age-dependent phenological plasticity in a wild bird. JOURNAL OF ANIMAL ECOLOGY 89, 2733–2741. https://doi.org/10.1111/1365-2656.13337

Boulton, R.L., Richard, Y., Armstrong, D.P., 2010. The Effect of Male Incubation Feeding, Food and Temperature on the Incubation Behaviour of New Zealand Robins. ETHOLOGY 116, 490–497. https://doi.org/10.1111/j.1439-0310.2010.01759.x

Bourne, A.R., Ridley, A.R., McKechnie, A.E., Spottiswoode, C.N., Cunningham, S.J., Cooke, S., 2021. Dehydration risk is associated with reduced nest attendance and hatching success in a cooperatively breeding bird, the southern pied babbler Turdoides bicolor. CONSERVATION PHYSIOLOGY 9. https://doi.org/10.1093/conphys/coab043

Brennan, L.A., Hernandez, F., Williford, D., 2020. Northern Bobwhite (Colinus virginianus), in: Billerman, S.M., Keeney, B.K., Rodewald, P.G., Schulenberg, T.S. (Eds.), Birds of the World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.norbob.01

Burnham, K.P., Anderson, D.R., Huyvaert, K.P., 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. Behav Ecol Sociobiol 65, 23–35. https://doi.org/10.1007/s00265-010-1029-6

Caldwell, P.J., Cornwell, G.W., 1975. Incubation Behavior and Temperatures of the Mallard Duck. The Auk 92, 706–731. https://doi.org/10.2307/4084783

Capp, E., Liebl, A.L., Cones, A.G., Russell, A.F., 2018. Advancing breeding phenology does not affect incubation schedules in chestnut-crowned babblers: Opposing effects of temperature and wind. Ecol Evol 8, 696–705. https://doi.org/10.1002/ece3.3524

Carroll, J.P., 1994. Family Odontophoridae (New World Quails).

Carroll, R.L., Davis, C.A., Fuhlendorf, S.D., Elmore, R.D., DuRant, S.E., Carroll, J.M., 2018. Avian parental behavior and nest success influenced by temperature fluctuations. Journal of Thermal Biology 74, 140–148. https://doi.org/10.1016/j.jtherbio.2018.03.020

Clusella-Trullas, S., Garcia, R.A., Terblanche, J.S., Hoffmann, A.A., 2021. How useful are thermal vulnerability indices? Trends in Ecology & Evolution 36, 1000–1010. https://doi.org/10.1016/j.tree.2021.07.001

Cohen, J.M., Fink, D., Zuckerberg, B., 2021. Extreme winter weather disrupts bird occurrence and abundance patterns at geographic scales. Ecography 44, 1143–1155. https://doi.org/10.1111/ecog.05495

Cones, A.G., Crowley, P.H., 2020. Optimal maternal incubation strategies for altricial and
precocial birds. Ecological Modelling 436, 109290.
https://doi.org/10.1016/j.ecolmodel.2020.109290
Conway, C.J., Martin, T.E., 2000. Effects of ambient temperature on avian incubation behavior.
Behavioral Ecology 11, 178–188. https://doi.org/10.1093/beheco/11.2.178
Cooke, S.J., Schreer, Jason.F., Philipp, D.P., Weatherhead, P.J., 2003. Nesting activity, parental
care behavior, and reproductive success of smallmouth bass, Micropterus dolomieu, in an
unstable thermal environment. Journal of Thermal Biology 28, 445–456.
https://doi.org/10.1016/S0306-4565(03)00038-X
Cooper, C.B., Voss, M.A., 2013. Avian Incubation Patterns Reflect Temporal Changes in
Developing Clutches. PLoS ONE 8, e65521.
https://doi.org/10.1371/journal.pone.0065521
Croston, R., Hartman, C.A., Herzog, M.P., Casazza, M.L., Feldheim, C.L., Ackerman, J.T.,
2020. Timing, frequency, and duration of incubation recesses in dabbling ducks. Ecology
and Evolution 10, 2513–2529. https://doi.org/10.1002/ece3.6078
Cunningham, S.J., Martin, R.O., Hojem, C.L., Hockey, P.A.R., 2013. Temperatures in Excess of
Critical Thresholds Threaten Nestling Growth and Survival in A Rapidly-Warming Arid
Savanna: A Study of Common Fiscals. PLoS ONE 8, e74613.
https://doi.org/10.1371/journal.pone.0074613
Dabbert, C.B., Pleasant, G., Schemnitz, S.D., 2020. Scaled Quail (Callipepla squamata), in:
Billerman, S.M., Keeney, B.K., Rodewald, P.G., Schulenberg, T.S. (Eds.), Birds of the
World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.scaqua.01
Deeming, C., 2002. Avian incubation: behaviour, environment and evolution. Oxford University
Press.
Denny, M., 2017. The fallacy of the average: on the ubiquity, utility and continuing novelty of
Jensen's inequality. Journal of Experimental Biology 220, 139–146.
https://doi.org/10.1242/jeb.140368
Diez-Mendez, D., Cooper, C.B., Jose Sanz, J., Verdejo, J., Barba, E., 2021. Deconstructing
incubation behaviour in response to ambient temperature over different timescales.
JOURNAL OF AVIAN BIOLOGY 52. https://doi.org/10.1111/jav.02781
Dorset, E.E., Sakaluk, S.K., Thompson, C.F., 2017. Behavioral Plasticity in Response to
Perceived Predation Risk in Breeding House Wrens. Evol Biol 44, 227–239.
https://doi.org/10.1007/s11692-016-9402-7
DuRant, S.E., Hopkins, W.A., Hepp, G.R., Walters, J.R., 2013. Ecological, evolutionary, and
conservation implications of incubation temperature-dependent phenotypes in birds:
Incubation temperature and avian offspring phenotype. Biological Reviews 88, 499–509.
https://doi.org/10.1111/brv.12015
DuRant, S.E., Willson, J.D., Carroll, R.B., 2019. Parental Effects and Climate Change: Will
Avian Incubation Behavior Shield Embryos from Increasing Environmental
Temperatures? Integrative and Comparative Biology. https://doi.org/10.1093/icb/icz083
Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R., Mearns, L.O., 2000.
Climate Extremes: Observations, Modeling, and Impacts. Science 289, 2068–2074.
https://doi.org/10.1126/science.289.5487.2068
Eldridge, W.H., Sweeney, B.W., Law, J.M., 2015. Fish growth, physiological stress, and tissue
condition in response to rate of temperature change during cool or warm diel thermal
cycles. Can. J. Fish. Aquat. Sci. 72, 1527–1537. https://doi.org/10.1139/cjfas-2014-0350

Errington, P.L., 1945. Some Contributions of a Fifteen-Year Local Study of the Northern
Bobwhite to a Knowledge of Population Phenomena. Ecological Monographs 15, 1–34.
https://doi.org/10.2307/1943293
Fischer, K., Karl, I., 2010. Exploring plastic and genetic responses to temperature variation using
copper butterflies. Clim. Res. 43, 17–30. https://doi.org/10.3354/cr00892
Garcia, R.A., Allen, J.L., Clusella-Trullas, S., 2019. Rethinking the scale and formulation of
indices assessing organism vulnerability to warmer habitats. Ecography 42, 1024–1036.
https://doi.org/10.1111/ecog.04226
Grimaudo, A.T., Hope, S.F., DuRant, S.E., Kennamer, R.A., Hallagan, J.J., Hopkins, W.A.,
2020. Ambient temperature and female body condition are related to night incubation
behavior in wood ducks Aix sponsa. JOURNAL OF AVIAN BIOLOGY 51.
https://doi.org/10.1111/jav.02379
Grubbauer, P., Hoi, H., 1996. Female penduline tits (Remiz pendulinus) choosing high quality
nests benefit by decreased incubation effort and increased hatching success.
ECOSCIENCE 3, 274–279. https://doi.org/10.1080/11956860.1996.11682342
Harrod, S.E., Rolland, V., 2020. Demographic responses of eastern bluebirds to climatic
variability in northeastern Arkansas. Population Ecology 62, 317–331.
https://doi.org/10.1002/1438-390X.12048
Hawkins, W.D., DuRant, S.E., 2020. Applications of machine learning in behavioral ecology:
Quantifying avian incubation behavior and nest conditions in relation to environmental
temperature. PLOS ONE 15, e0236925. https://doi.org/10.1371/journal.pone.0236925
Hepp, G.R., Kennamer, R.A., Johnson, M.H., 2006. Maternal effects in Wood Ducks: incubation
temperature influences incubation period and neonate phenotype. Funct Ecology 20, 308–
314. https://doi.org/10.1111/j.1365-2435.2006.01108.x
Huey, R.B., Kearney, M.R., Krockenberger, A., Holtum, J.A.M., Jess, M., Williams, S.E., 2012.
Predicting organismal vulnerability to climate warming: roles of behaviour, physiology
and adaptation. Philosophical Transactions of the Royal Society B: Biological Sciences
367, 1665–1679. https://doi.org/10.1098/rstb.2012.0005
Husby, A., Nussey, D.H., Visser, M.E., Wilson, A.J., Sheldon, B.C., Kruuk, L.E.B., 2010.
CONTRASTING PATTERNS OF PHENOTYPIC PLASTICITY IN REPRODUCTIVE
TRAITS IN TWO GREAT TIT (PARUS MAJOR) POPULATIONS. EVOLUTION 64,
2221–2237. https://doi.org/10.1111/j.1558-5646.2010.00991.x
IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and
III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change
[Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland.
Intergovernmental Panel on Climate Change (IPCC).
https://doi.org/10.59327/IPCC/AR6-9789291691647
Jensen, J., 1906. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. Acta
mathematica 30, 175–193.
Kearney, M.R., Matzelle, A., Helmuth, B., 2012. Biomechanics meets the ecological niche: the
importance of temporal data resolution. Journal of Experimental Biology 215, 1422–
1424. https://doi.org/10.1242/jeb.072249
Kirkpatrick, W.H., Sheldon, K.S., 2022. Experimental increases in temperature mean and
variance alter reproductive behaviours in the dung beetle Phanaeus vindex. Biology
Letters 18, 20220109.

Le Vaillant, J., Potti, J., Camacho, C., Canal, D., Martinez-Padilla, J., 2021. Fluctuating selection
driven by global and local climatic conditions leads to stasis in breeding time in a
migratory bird. JOURNAL OF EVOLUTIONARY BIOLOGY 34, 1541–1553.
https://doi.org/10.1111/jeb.13916
Lusk, J.J., Smith, S.G., Fuhlendorf, S.D., Guthery, F.S., 2006. Factors Influencing Northern
Bobwhite Nest-Site Selection and Fate. Journal of Wildlife Management 70, 564–571.
https://doi.org/10.2193/0022-541X(2006)70[564:FINBNS]2.0.CO;2
Ma, G., Hoffmann, A.A., Ma, C.-S., 2015. Daily temperature extremes play an important role in
predicting thermal effects. Journal of Experimental Biology jeb.122127.
https://doi.org/10.1242/jeb.122127
MacDonald, E.C., Camfield, A.F., Jankowski, J.E., Martin, K., 2014. An alpine-breeding
songbird can adjust dawn incubation rhythms to annual thermal regimes. AUK 131, 495–
506. https://doi.org/10.1642/AUK-13-234.1
Mamantov, M.A., Sheldon, K.S., 2021. Behavioural responses to warming differentially impact
survival in introduced and native dung beetles. Journal of Animal Ecology 90, 273–281.
https://doi.org/10.1111/1365-2656.13366
Martin, T.E., Ton, R., Oteyza, J.C., 2018. Adaptive influence of extrinsic and intrinsic factors on
variation of incubation periods among tropical and temperate passerines. The Auk.
Masero, J.A., Monsa, R., Amat, J.A., 2012. Dual function of egg-covering in the Kentish plover
Charadrius alexandrinus. Behav 149, 881–895. https://doi.org/10.1163/1568539X-
00003008
Matysioková, B., Remeš, V., 2018. Evolution of parental activity at the nest is shaped by the risk
of nest predation and ambient temperature across bird species. Evolution 72, 2214–2224.
https://doi.org/10.1111/evo.13580
McDiarmid, C.S., Naguib, M., Griffith, S.C., 2018. Calling in the heat: the zebra finch
"incubation call" depends on heat but not reproductive stage. BEHAVIORAL
ECOLOGY 29, 1245–1254. https://doi.org/10.1093/beheco/ary123
Nelson, S.D., Keever, A.C., Wightman, P.H., Bakner, N.W., Argabright, C.M., Byrne, M.E.,
Collier, B.A., Chamberlain, M.J., Cohen, B.S., 2022. Fine-scale resource selection and
behavioral tradeoffs of eastern wild turkey broods. JOURNAL OF WILDLIFE
MANAGEMENT 86. https://doi.org/10.1002/jwmg.22222
Nord, A., Nilsson, J.-Å., 2016. Long-term consequences of high incubation temperature in a wild
bird population. Biol. Lett. 12, 20160087. https://doi.org/10.1098/rsbl.2016.0087
Nord, A., Sandell, M.I., Nilsson, J.-Å., 2010. Female zebra finches compromise clutch
temperature in energetically demanding incubation conditions: Incubation metabolic rate
and clutch temperature. Functional Ecology 24, 1031–1036.
https://doi.org/10.1111/j.1365-2435.2010.01719.x
Nowakowski, A.J., Frishkoff, L.O., Agha, M., Todd, B.D., Scheffers, B.R., 2018. Changing
Thermal Landscapes: Merging Climate Science and Landscape Ecology through Thermal
Biology. Curr Landscape Ecol Rep 3, 57–72. https://doi.org/10.1007/s40823-018-0034-8
Pleasant, G.D., Dabbert, C.B., Mitchell, R.B., 2006. Nesting Ecology and Survival of Scaled
Quail in the Southern High Plains of Texas. Journal of Wildlife Management 70, 632–
640. https://doi.org/10.2193/0022-541X(2006)70[632:NEASOS]2.0.CO;2
R Core Team, 2023. R: A language and environment for statistical computing.

Rehm, E.M., Olivas, P., Stroud, J., Feeley, K.J., 2015. Losing your edge: climate change and the
conservation value of range-edge populations. Ecology and Evolution 5, 4315–4326.
https://doi.org/10.1002/ece3.1645
Ringelman, J.K., Longcore, J.R., Owen, R.B., 1982. Nest and Brood Attentiveness in Female
Black Ducks. The Condor 84, 110. https://doi.org/10.2307/1367831
Rodgers, E.M., Cocherell, D.E., Nguyen, T.X., Todgham, A.E., Fangue, N.A., 2018. Plastic
responses to diel thermal variation in juvenile green sturgeon, Acipenser medirostris.
JOURNAL OF THERMAL BIOLOGY 76, 147–155.
https://doi.org/10.1016/j.jtherbio.2018.07.015
Rohwer, V.G., Purcell, J.R., 2019. Geographic variation in incubation behavior of a widely
distributed passerine bird. PLoS ONE 14, e0219907.
https://doi.org/10.1371/journal.pone.0219907
Ruel, J.J., Ayres, M.P., 1999. Jensen's inequality predicts effects of environmental variation.
Trends in Ecology & Evolution 14, 361–366. https://doi.org/10.1016/S0169-
5347(99)01664-X
Santos Nobre, J., da Motta Singer, J., 2007. Residual Analysis for Linear Mixed Models.
Biometrical J 49, 863–875. https://doi.org/10.1002/bimj.200610341
Sharpe, L., Cale, B., Gardner, J.L., 2019. Weighing the cost: the impact of serial heatwaves on
body mass in a small Australian passerine. JOURNAL OF AVIAN BIOLOGY 50.
https://doi.org/10.1111/jav.02355
Sharpe, L.L., Bayter, C., Gardner, J.L., 2021. Too hot to handle? Behavioural plasticity during
incubation in a small, Australian passerine. Journal of Thermal Biology 98, 102921.
https://doi.org/10.1016/j.jtherbio.2021.102921
Sharpe, L.L., Prober, S.M., Gardner, J.L., 2022. In the Hot Seat: Behavioral Change and Old-
Growth Trees Underpin an Australian Songbird's Response to Extreme Heat.
FRONTIERS IN ECOLOGY AND EVOLUTION 10.
https://doi.org/10.3389/fevo.2022.813567
Sheldon, K.S., Dillon, M.E., 2016. Beyond the Mean: Biological Impacts of Cryptic
Temperature Change. Integrative and Comparative Biology 56, 110–119.
https://doi.org/10.1093/icb/icw005
Simeone, A., Luna-Jorquera, G., Wilson, R., 2004. Seasonal variations in the behavioural
thermoregulation of roosting Humboldt penguins (Spheniscus humboldti) in north-central
Chile. JOURNAL OF ORNITHOLOGY 145, 35–40. https://doi.org/10.1007/s10336-
003-0005-6
Simmonds, E.G., Sheldon, B.C., Coulson, T., Cole, E.F., 2017. Incubation behavior adjustments,
driven by ambient temperature variation, improve synchrony between hatch dates and
caterpillar peak in a wild bird population. ECOLOGY AND EVOLUTION 7, 9415–
9425. https://doi.org/10.1002/ece3.3446
Skrade, P.D.B., Dinsmore, S.J., 2012. Incubation patterns of a shorebird with rapid multiple
clutches, the Mountain Plover (Charadrius montanus). CANADIAN JOURNAL OF
ZOOLOGY 90, 257–266. https://doi.org/10.1139/Z11-134
Smith, M.G., Kaiser, S.A., Sillett, T.S., Webster, M.S., 2018. Variation in nest characteristics
and brooding patterns of female Black-throated Blue Warblers is associated with thermal
cues. AUK 135, 733–747. https://doi.org/10.1642/AUK-17-195.1
Sudnick, M., Brodie, B.S., Williams, K.A., 2021. Nature versus nurture: Structural equation
modeling indicates that parental care does not mitigate consequences of poor

environmental conditions in Eastern Bluebirds (*Sialia sialis*). Ecol Evol 11, 15237–
15248. https://doi.org/10.1002/ece3.8207
Sultan, S.E., 2015. Organism and environment: ecological development, niche construction, and
adaptation. Oxford University Press, USA.
Taff, C.C., Freeman-Gallant, C.R., 2021. Female ornamentation, incubation behavior, and
reproductive success in a wild bird. BEHAVIORAL ECOLOGY AND
SOCIOBIOLOGY 75. https://doi.org/10.1007/s00265-021-03033-0
Telemeco, R.S., Elphick, M.J., Shine, R., 2009. Nesting lizards (*Bassiana duperreyi*)
compensate partly, but not completely, for climate change. Ecology 90, 17–22.
https://doi.org/10.1890/08-1452.1
Turriago, J.L., Tejedo, M., Hoyos, J.M., Bernal, M.H., 2022. The effect of thermal
microenvironment in upper thermal tolerance plasticity in tropical tadpoles. Implications
for vulnerability to climate warming. J Exp Zool Pt A 337, 746–759.
https://doi.org/10.1002/jez.2632
Urban, M.C., 2015. Accelerating extinction risk from climate change. Science 348, 571–573.
https://doi.org/10.1126/science.aaa4984
van de Ven, T.M.F.N., McKechnie, A.E., Cunningham, S.J., 2019. The costs of keeping cool:
behavioural trade-offs between foraging and thermoregulation are associated with
significant mass losses in an arid-zone bird. Oecologia 191, 205–215.
https://doi.org/10.1007/s00442-019-04486-x
Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S., Savage,
V., Tunney, T.D., O'Connor, M.I., 2014. Increased temperature variation poses a greater
risk to species than climate warming. Proceedings of the Royal Society B: Biological
Sciences 281, 20132612. https://doi.org/10.1098/rspb.2013.2612
Walsberg, G.E., 1993. Thermal Consequences of Diurnal Microhabitat Selection in a Small Bird.
Ornis Scandinavica 24, 174. https://doi.org/10.2307/3676733
Zhang, L., An, B., Shu, M., Zhao, C., Yang, X., Suo, Y., Se, Y., Dabu, X., 2017. Incubation
strategies of the Black-necked Crane (Grus nigricollis) in relation to ambient temperature
and time of day. AVIAN RESEARCH 8. https://doi.org/10.1186/s40657-017-0076-3
696