ISWI chromatin remodeling complexes recruit NSD2 and H3K36me2

in pericentromeric heterochromatin

Running title: H3K36me2 in heterochromatin

Naoki Goto[1], Kazuma Suke[2], Nao Yonezawa[2], Hidenori Nishihara[1,3], Yuko Sato[1,4], Tomoya

Kujirai[5], Hitoshi Kurumizaka[5], Kazuo Yamagata[2], and Hiroshi Kimura[1,4,*]

[1]School of Life Science and Technology, Tokyo Institute of Technology

[2]Faculty of Biology-Oriented Science and Technology, Kindai University

[3]Department of Advanced Bioscience, Graduate School of Agriculture, Kindai University

[4]Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology

[5]Instite of Quantitative Biology, University of Tokyo

*Correspondence should be addressed:

Hiroshi Kimura

Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology

4259 Nagatsuta-cho, Midori-ku, Yokohama, 226-8503, Japan

Phone: +81-45-924-5742

ORCID: 0000-0003-0854-083X

E-mail: hkimura@bio.titech.ac.jp

Summary

The authors discovered histone H3K36me2, which is believed to be enriched in potentially

active genomic regions, is also located in transcriptionally inactive regions called

heterochromatin in some cell types. The detailed molecular mechanism of its

heterochromatin targeting is now revealed.

Abstract

Histone H3 lysine36 dimethylation (H3K36me2) is generally distributed in the gene body and euchromatic intergenic regions. However, we found that H3K36me2 is enriched in pericentromeric heterochromatin in some mouse cell lines. We here revealed the mechanism of heterochromatin targeting of H3K36me2. Among several H3K36 methyltransferases, NSD2 was responsible for inducing heterochromatic H3K36me2. Depletion and overexpression analyses of NSD2-associating proteins revealed that NSD2 recruitment to heterochromatin was mediated through the imitation switch (ISWI) chromatin remodeling complexes, such as BAZ1B-SMARCA5 (WICH), which directly binds to AT-rich DNA via a BAZ1B domain containing AT-hook-like motifs. The abundance and stoichiometry of NSD2, SMARCA5, and BAZ1B or BAZ2A could determine the localization of H3K36me2 in different cell types. To explore the physiological role of heterochromatic H3K36me2, we analyzed mouse tissues and embryos. As a result, H3K36me2 was found in heterochromatin at the 2- to 4-cell stages of mouse preimplantation embryos, suggesting its involvement in developmental regulation.

Introduction

In eukaryotic cells, DNA is packed into chromatin, which could be categorized into highly condensed heterochromatin and loosely decondensed euchromatin (Woodcock and Ghosh, 2010). Pericentromeric heterochromatin (PCH) is composed of repeat elements, including α-satellites in humans (Altemose et al., 2022) and major satellites in mice (Saksouk et al., 2015). Except during specific stages of development and under some cellular stresses, transcription from PCH is mostly repressed (Smurova and Wulf, 2018). In mouse interphase cell nuclei, PCH forms microscopically visible clusters termed chromocenters. PCH in mice is epigenetically characterized by a high level of histone H3 lysine 9 trimethylation (H3K9me3), H4 lysine 20 trimethylation (H4K20me3), cytosine methylation on CpG sites, and low histone acetylation (Probst and Almouzni, 2008). Other histone modifications, such as H3 lysine 27 monomethylation (H3K27me1) (Peters et al., 2003), H3 lysine 56 trimethylation (H3K56me3) (Jack et al., 2013), H3 lysine 64 trimethylation (H3K64me3) (Lange et al., 2013), and H3 lysine 79 trimethylation (H3K79me3) (Ooga et al., 2008), have been reported to localize to PCH, but the detailed molecular mechanism for specialized localization and physiological significance, are not well understood.

In mammals, histone H3 lysine36 di- and tri-methylation (H3K36me2 and H3K36me3) are

known to function in euchromatin, methylated by histone methyltransferases such as nuclear

receptor SET domain-containing 1/2/3 (NSD1/2/3) and SETD2 (Wagner and Carpenter,

2012) and demethylated by lysine demethylases such as KDM2A/JHDM1A (Blackledge et

al., 2010). H3K36me2 serves as a marker to attract DNMT3A to the intergenic region and

contributes to the establishment of DNA methylation patterns (Weinberg et al., 2019).

H3K36me3 accumulates in transcriptionally active gene bodies by Set2 (an ortholog of

SETD2) in yeast and by SETD2 in mouse ES cells, serving as a scaffold for the histone

deacetylation complex RPD3 (Carrozza et al., 2005; Guan et al., 2023) and DNA

methyltransferase DNMT3B (Neri et al., 2017) to repress aberrant transcription. Such

regulation of DNA methylation via H3K36me2/3 has been shown to play an important role in

germline development (Xu et al., 2019; Shirane et al., 2020). H3K36me2 also shows

antagonistic distribution with H3K27me3. In mouse ES cells, H3K36me2 protects the

genome from excessive accumulation of PRC2-dependent H3K27me3 (Streubel et al., 2018;

Chen et al., 2022). Similar antagonism has been observed in multiple myeloma, which highly

express NSD2 (Popovic et al., 2014).

In addition to the function of H3K36me2/3 in euchromatin, several studies have reported

their roles in heterochromatin. H3K36me2 is enriched in the pericentromere regions of

Drosophila eye discs and represses transcription of transposons such as long interspersed

nuclear element (LINE) and long-terminal repeat (LTR) (Chaouch et al., 2021). KDM2A

depletion causes H3K36me2 enhancement and increased transcription of satellite repeats in

the PCH of mouse and human cells (Frescas et al., 2008). H3K36me3 accumulation in PCH

has also been observed in mouse ES cells (Chantalat et al., 2011). In addition, H3K9me3

and H3K36me3 co-localize in some intergenic regions and gene bodies in mouse ES cells

and can function as potential enhancers (Barral et al., 2022). In fission yeast, H3K36me3 by

Set2 is required for histone deacetylation for RNAi-dependent heterochromatin

establishment (Chen et al., 2008), maintenance of condensed chromatin near subtelomeres

(Matsuda et al., 2015), and transcriptional repression of pericentromeres and subtelomeres

(Georgescu et al., 2020).

ATP-dependent chromatin remodeling complexes are widely conserved in eukaryotes.

Higher eukaryotes have four subfamilies: switch/sucrose non-fermenting (SWI/SNF),

chromodomain-helicase DNA-binding (CHD), imitation switch (ISWI), and inositol-requiring

80 (INO80) (Hasan and Ahuja, 2019). These complexes utilize ATP hydrolysis-derived

energy to drive nucleosome assembly, nucleosome sliding, and histone exchange to

regulate transcription, replication, and repair (Clapier et al., 2017). The CHD and ISWI

complexes play a particularly crucial role in heterochromatin. The NuRD complex recognizes

H3K9me3 via the PHD domain of CHD4 (Musselman et al., 2012) and methylated DNA via

the MBD2 subunit (Zhang et al., 1999), suggesting that the complex is involved in

heterochromatin regulation and transcriptional repression. ISWI subfamily containing

SMARCA5/SNF2H as an ATPase consists of five members in higher eukaryotes; CHRAC

(with BAZ1A/ACF1, CHRAC15, and CHRAC17), ACF (with BAZ1A), WICH (with

BAZ1B/WSTF), NoRC (with BAZ2A/TIP5), and RSF (with and RSF1) (Bartholomew, 2014).

ACF accumulates in PCH during the S phase of mouse cells, and depletion of BAZ1A delays

late S phase progression, suggesting that ACF seems to contribute to PCH replication

(Collins et al., 2002). WICH is associated with DNA replication foci during the S phase in

mouse cells in a PCNA-dependent manner and concentrated on PCH in the late S phase.

Depletion of each subunit of WICH increases H3K9me3 and HP1α/β levels and reduces

chromatin accessibility after replication (Poot et al., 2004). It has also been reported that

WICH localizes to inactive X chromosomes in the late S phase in human cells (Culver-

Cochran and Chadwick, 2012). NoRC suppresses rRNA transcription by inducing DNA

methylation and histone deacetylation on the rDNA promoter in the nucleolus (Zhou et al.,

2002; Santoro et al., 2002). In addition, BAZ2A depletion in mouse cells causes genomic

instability by decreasing H4K20me3 and H3K9me3 in major satellite repeats and rDNA

(Guetg et al., 2010). ISWI complexes are thought to be differentially recruited to their target

regions by SMARCA5 partner factors that bind to different histone modifications (Ito et al.,

1999; Ribeyre et al., 2016; Strohner et al., 2001; Li et al., 2006; Tallant et al., 2015);

however, the mechanism of target recognition on PCH is not well understood.

In a series of immunofluorescence studies using mouse cells, we noticed that some cells

exhibit H3K36me2 enrichment in PCH while others show typical euchromatic distributions.

This intriguing observation led us to investigate the mechanism regulating H3K36me2

localization. Here we show that H3K36me2 could be localized to PCH via the ISWI-NSD2

axis.

Results

H3K36me2 is enriched in pericentromeric heterochromatin in several mouse cell lines

To investigate the nuclear distribution of H3K36me2 in various mouse cell types, we

immunostained H3K36me2 with a specific mouse monoclonal antibody (clone 2C3/CMA332;

Rechtsteiner et al., 2010) and detected it using a confocal microscope. H3K9me3 was co-

stained as a marker of constitutive heterochromatin, or chromocenter. In Pmi28 and C2C12

myoblasts and MC12 embryonic carcinoma cells, H3K36me2 showed typical euchromatin

distribution (Fig. 1A and S1A). In contrast, in immortalized mouse embryonic fibroblasts

(iMEFs), NIH3T3 fibroblasts, and embryonic stem cells (ESCs), H3K36me2 was

concentrated on PCH where H3K9me3 is enriched (Fig. 1B and S1A). Intensity

measurements confirmed that the enrichment of H3K36me2 in PCH was higher in iMEFs,

NIH3T3, and ESCs, compared to the other three cell lines (Fig. 1C). Using

immunofluorescence with another H3K36me2-specific antibody (rabbit monoclonal clone

EPR16994(2); Bressan et al., 2021), we observed the same localization patterns (Fig. S1B).

This indicates that the variations in H3K36me2 distribution are not due to antibody staining

artifacts. In iMEFs that exhibit H3K36me2 PCH localization, other histone marks such as

H3K27me3, H3K9me2, and H3K36me3, were not concentrated in PCH (Fig. S1C-E),

suggesting a unique feature of H3K36me2 independently of these histone marks.

To validate the immunofluorescence data, we analyzed the genome-wide distribution of

H3K36me2 in different mouse cell types using the public ChIP-seq datasets (Barral et al.,

2022; Vian et al., 2018; Yuan et al., 2020). The correlation analysis showed that H3K36me2

and H3K9me3 were highly correlated in ESCs, but not in B cells and pancreatic cancer cells

(Fig. 1D). H3K36me2 in B cells and pancreatic cancer cells exhibited anti-correlation with

H3K9me3 in B cells. Upon analyzing enrichments in repeat elements with transcription units,

H3K9me3 was enriched in LINE and LTR, particularly at the transcription start site (TSS), in

both ESCs and B cells (Fig. S1F and S1G). This is consistent with the silencing role of this

modification in those repeats. H3K36me2 in ESCs, but not in B cells, was also enriched in

LINE and LTR, reflecting its genome-wide correlation with H3K9me3. H3K36me2 in both

ESCs and B cells was enriched at TSS of SINE, which is rich in CpG that can be methylated

(Varshney et al., 2015), in line with the role of H3K36me2 in establishing DNA methylation in

euchromatin (Weinberg et al., 2019). Thus, the enrichment of H3K36me2 in H3K9me3-rich

genomic regions in ESCs observed in public datasets well agreed with immunofluorescence

results.

H3K9me3 and DNA methylation are not required for H3K36me2 in pericentromeric

heterochromatin

H3K9me3 and DNA methylation are the major epigenetic marks concentrated in PCH in

mouse cells. To examine if these marks are required for PCH H3K36me2 localization, we

used knockout (KO) cell lines lacking H3K9me3 and DNA methylation. 5KO iMEFs lack

H3K9me3 because five H3K9me2/3-methyltransferases, i.e., Suv39h1, Suv39h2, Setdb1,

G9a, and Glp, are deleted (Fukuda et al., 2023). TKO ESCs lack DNA methylation because

three DNA methyltransferases, i.e., Dnmt1, Dnmt3a, and Dnmt3b, are deleted (Tsumura et

al., 2006). In both of these cell lines, H3K36me2 remained enriched in PCH (Fig. S2A and

S2B), although the enrichment was slightly reduced in 5KO iMEFs. Therefore, the

localization of H3K36me2 in PCH does not require H3K9me3 and DNA methylation, and the

related methyltransferases.

NSD2 is responsible for H3K36me2 in pericentromeric heterochromatin

To identify which H3K36 methyltransferase targets PCH and mediates H3K36me2, we

analyzed the localization of transiently expressed Halo-tagged enzymes in NIH3T3 cells

(Fig. S2C). Among several methylation and demethylation enzymes, NSD2/WHSC1 and

KDM2A/JHDM1A localized to PCH. KDM2A is indeed known to localize to PCH in an HP1-

dependent manner (Borgel et al., 2017). Although H3K9me3 is required for HP1 localization

to the PCH (Lachner et al., 2001), H3K36me2 was still concentrated in H3K9me3-deficient

cells (Fig. S2A), suggesting that KDM2A per se does not play a primary role in H3K36me2

enrichment in PCH. Therefore, we focused on NSD2-mediated H3K36me2 regulation.

Hereafter, we used iMEFs and Pmi28 cells for cells exhibiting H3K36me2 in PCH and

euchromatin, respectively, for overexpression and iMEFs for KO studies.

Overexpressed Halo-NSD2 localized to PCH in iMEFs but not obvious in Pmi28 cells (Fig.

2A and 2B). Measuring PCH-to-nucleoplasm intensity ratios in individual nuclei revealed that

Halo-NSD2 was enriched in PCH in all iMEF nuclei but its enrichment was lower in Pmi28

nuclei (Fig. 2C). In iMEFs, Halo-NSD2 expression did not affect the enrichment of

H3K36me2 in PCH (Fig. 2D), suggesting that increased levels of NSD2 had little effect. In

Pmi28 cells, when Halo-NSD2 was expressed, PCH H3K36me2 was induced albeit at lower

levels compared to iMEFs, suggesting that NSD2 alone is not sufficient for inducing PCH

H3K36me2.

To validate the role of NSD2 in PCH H3K36me2 in iMEFs, we generated NSD2 KO cell lines

using the CRISPR/Cas9 system (Fig. 2E). NSD2 depletion reduced H3K36me2 levels in

PCH, and this phenotype was rescued by Halo-NSD2 expression (Fig. 2F and 2G). Thus,

this result indicates that NSD2 is responsible for inducing H3K36me2 in PCH.

SMARCA5 is required for NSD2 localization and H3K36me2 in PCH

As the PWWP domain of NSD2 binds to H3K36me2 and stabilizes NSD2 on chromatin

(Sankaran et al., 2016), it is possible that once H3K36me2 is introduced into PCH by ectopic

NSD2 expression, the modification could be maintained through positive feedback. However,

the mechanism of how NSD2 initially targets remains uncertain. Therefore, we searched for

a potential NSD2 interactor on heterochromatin. Among the NSD2-interacting factors

comprehensively analyzed in a previous report (Huang et al., 2019), there were several

candidate proteins associated with PCH, including SMARCA5, PARP1, ORC2, MKI67,

MECP2, KDM2A, and HMGB1 (Fig. 3A) (Malla et al., 2023; Quenet et al., 2008; Prasanth et

al., 2010; Sobecki et al., 2016; Nan et al., 2007; Frescas et al., 2008; Pallier et al., 2003).

These candidate factors and NSD2 were knocked down in iMEFs using the lentivirus shRNA

expression system, and the number of cells clearly showing PCH H3K36me2 was counted

(Fig. 3B). The knockdown efficiency was verified by qPCR. Knockdown of NSD2 had the

most drastic effect, reassuring the critical role of NSD2 in PCH H3K36me2. Among the

candidate proteins, knockdown of the chromatin remodeling factor SMARCA5/SNF2H

substantially reduced the number of cells exhibiting PCH H3K36me2. Therefore, we

performed a deeper analysis of SMARCA5.

Firstly, we overexpressed Halo-SMARCA5 in iMEFs and Pmi28 cells to examine the

localization of the expressed protein and H3K36me2. Halo-SMARCA5 was concentrated in

PCH in iMEFs but not much in Pmi28 cells (Fig. 3C and 3D). SMARCA5 expression did not

affect PCH H3K36me2 levels in both iMEFs and Pmi28 cells (Fig. 3E). These data suggest

that SMARCA5 alone is not sufficient to target PCH and induce H3K36me2 there.

Secondly, we established SMARCA5 KO iMEFs to investigate the requirement of SMARCA5

on H3K36me2 and NSD2 localization in PCH (Fig. 3F). The loss of SMARCA5 led to a

decrease in PCH H3K36me2 levels, and this phenotype was rescued to some extent by the

expression of Halo-SMARCA5 (Fig. 3G and 3H). Furthermore, in SMARCA5 KO cells, PCH

localization of transfected Halo-NSD2 was lower, and the expression of SMARCA5-sfGFP,

but not EGFP alone, also partially restored PCH H3K36me2 enrichment (Fig. 3I). Taken

together, we concluded that SMARCA5 is required, but not sufficient, for PCH H3K36me2 by

recruiting NSD2.

BAZ1B regulates PCH localization of SMARCA5, NSD2, and H3K36me2

In mammalian cells, SMARCA5 forms ISWI complexes with various partner factors, such as

ACF with BAZ1A, WICH with BAZ1B, and NoRC with BAZ2A (Bartholomew, 2014).

Chromatin remodeling is driven by SMARCA5 through its ATPase activity, while the partner

factors are thought to modulate the activity and recruitment of the ISWI complexes to target

sites (Ito et al., 1999; Ribeyre et al., 2016; Strohner et al., 2001). Both BAZ1A and BAZ1B

have been reported to accumulate at the PCH in the S phase and to promote proper

heterochromatin replication (Collins et al., 2002; Poot et al., 2004; Bozhhenok et al., 2002).

BAZ2A also associates with pericentromeric repeats and rDNA in the nucleolus and

contributes to its genome stabilization (Guetg et al., 2010). Therefore, we hypothesized that

BAZ1A, BAZ1B, and/or BAZ2A in the ISWI complexes might bind to PCH to recruit NSD2

(Fig. 4A). To test whether these BAZ proteins target PCH, we expressed their Emerald-
green fluorescent protein (emGFP)-tagged versions in iMEFs and Pmi28 cells. Whereas
BAZ1A diffused throughout the nucleoplasm in most cells of both cell lines (Fig. S3A),
BAZ1B and BAZ2A were concentrated in PCH in both cell types (Fig. 4B, 4C, S3B, and
S3C). Overexpression of BAZ1B and BAZ2A however did not induce PCH H3K36me2 in
Pmi28 cells (Fig. 4D and S3D), probably because other proteins that are critical for PCH
H3K36me2, such as SMARCA5 and NSD2, are not abundantly present.
259
To understand the role of BAZ1B and BAZ2A in PCH H3K36me2, we attempted to establish
single and double KO iMEFs; however, only BAZ1B KO lines were obtained (Fig. 4E)
leading us to focus on the function of BAZ1B. PCH H3K36me2 was indeed diminished in
BAZ1B KO cells and this phenotype was partially restored by the introduction of emGFP-
BAZ1B (Fig. 4F and 4G). The levels of co-expressed Halo-NSD2 and Halo-SMARCA5 at
PCH were both decreased in BAZ1B KO cells (Fig. 4H and 4I). The expression of emGFP-
BAZ1B, but not EGFP, in KO cells rescued and increased the levels of NSD2 (Fig. 4H) and
SMARCA5 (Fig. 4I) in PCH. These results indicate that BAZ1B facilitates PCH localization of
NSD2, SMARCA5, and H3K36me2 (Fig. 4J).
269
BAZ1B binds to AT-rich DNA through its AT-hook-like motifs
To determine how BAZ1B can target to PCH, we expressed truncated BAZ1B mutants fused
with EGFP in iMEFs (Fig. S4A). The C-terminal region containing a bromodomain (BrD) was
found to be essential for BAZ1B localization to the PCH (Fig. 5A); however, the regions both
upstream and downstream of the BrD, rather than the BrD as such, appeared to be critical
for PCH targeting (Fig. S4A). Since neither H3K9me3 nor DNA methylation is required for
PCH H3K36me2 (Fig. S2A and S2B), we hypothesized that BAZ1B directly binds to DNA
elements. Then, we found that three AT-hook-like motifs, conserved across vertebrates (Fig.
5A), are located in the C-terminal region of BAZ1B (Fig. 5A; BrD+AT1-3). These motifs
resemble the typical AT-hook motifs known to bind to AT-rich DNA (Arg-Gly-Arg-Pro;
Aravind and Landsman, 1998). A BAZ1B mutant, which harbors amino acid substitutions in
these AT-hook-like motifs (BrD+AT1-3mut) no longer localized to PCH (Fig. 5A). In addition,
a truncated BAZ1B mutant (AT1-2) containing two AT-hook-like motifs but lacking BrD still
targeted the PCH, although with a more diffuse background (Fig. S4A). These data suggest
that three AT-hook-like motifs are required for DNA binding.
285
We therefore examined whether the BAZ1B domain containing the three AT-hook-like motifs
directly bind to AT-rich DNA in vitro (Fig. 5B). Both EGFP-BrD+AT1-3 and EGFP-BrD+AT1-
3mut were expressed in HEK293T cells and then purified using anti-GFP magnetic beads

(Fig. S4B). These beads were mixed with input DNA, either AT-rich major satellite repeats,

GC-rich EF1α promoter, or 601 sequences (Lowary and Widom, 1998) (Fig. S4C-E).

Following the removal of unbound DNA, the beads were washed, and the bound DNA was

eluted (Fig. 5B). DNA from each fraction was then analyzed by agarose gel electrophoresis

(Fig. 5C). Major satellite repeats bound to BrD+AT1-3 but not to BrD+AT1-3mut (Fig. 5C),

whereas the binding of EF1α promoter to BrD+AT1-3 was much weaker. This result

indicates that the AT-hook-like motifs have a preference for AT-rich DNA. We next examined

whether BrD+AT1-3 could bind to nucleosomal DNA using a 145-bp 601 sequence.

BrD+AT1-3 indeed bound to nucleosomes, although the binding affinity appeared to be

weaker compared to the naked DNA (Fig. 5C). Thus, BAZ1B directly binds to AT-rich DNA

via AT-hook-like motifs, which could recruit SMARCA5 and then NSD2 to induce H3K36me2

in AT-rich PCH.

302　Coordinated expression of NSD2, SMARCA5, and BAZ1B promotes PCH localization

303　of H3K36me2

To determine the molecular differences between cells with and without PCH H3K36me2, we

investigated the protein expression levels of NSD2, SMARCA5, and BAZ1B by

immunoblotting (Fig. 6A). Compared to cells without PCH H3K36me2 (i.e., MC12, C2C12,

and Pmi28), NSD2 and SMARCA5 were highly expressed in NIH3T3 and iMEFs, both of

which display high levels of H3K36me2 in PCH (Fig. 1C). BAZ1B was also highly expressed

in iMEFs. In ESCs, SMARCA5 was relatively highly expressed, whereas levels of NSD2 and

BAZ1b were moderate, possibly explaining the lower PCH enrichment of H3K36me2 in

ESCs compared to iMEFs and NIH3T3 (Fig. 1C).

Using Pmi28 cells, which exhibit the lowest levels of NSD2 and BAZ1B without PCH

H3K36me2, we investigated which proteins or their combinations, among NSD2, SMARCA5,

and BAZ1B or BAZ2A, could induce PCH H3K36me2 by transient expression (Fig. 6B).

When a single protein was expressed, only NSD2 induced a low level of PCH H3K36me2,

compared to a control (EGFP alone), as shown before (Fig. 2D, 3E, 4D, and S3D).

Combining SMARCA5 with either BAZ1B or BAZ2A did not have any effect. This result is

reasonable because a methyltransferase is essential to increase H3K36me2. Co-expression

of NSD2 with SMARCA5 did not markedly differ from the effects of NSD2 alone. By contrast,

drastic increases of PCH H3K36me2 were observed when NSD2 was paired with either

BAZ1B or BAZ2A. In addition, the concurrent expression of NSD2, SMARCA5, and BAZ1B

further slightly increased PCH H3K36me2 levels. Since SMARCA5 levels are not very low in

Pmi28, its overexpression appears to have a minimal impact on PCH H3K36me2 compared

to the combined effects of NSD2 and BAZ1B, or BAZ2A. Taken together with KO analysis

using iMEFs, these results indicate that PCH H3K36me2 is induced by the cooperation of

NSD2, SMARCA5, and BAZ1B or BAZ2A.

H3K36me2 is enriched in heterochromatin in mouse preimplantation embryos at the 2-

to 4-cell stages

Finally, we investigated whether H3K36me2 is also localized to the PCH in mice. We first

chose the spleen, lung, testis, liver, and brain for immunostaining, based on the high

expression of at least two of NSD2, SMARCA5, BAZ1B, and BAZ2A, searching the public

database (Palasca et al., 2018). However, none of these tissues showed H3K36me2 in PCH

(Fig. S5A), suggesting that H3K36me2 is located in euchromatin in adult mouse tissues as

generally observed. We also stained 13.5-day mouse embryo sections with anti-H3K36me2

but did not find PCH localization in any areas closely looked at (Fig. S5B).

We next analyzed totipotent cells in mouse preimplantation embryos since PCH H3K36me2

was found in undifferentiated ESCs. From the zygote to the 2-cell stage, the PCH is

distributed in a ring around the nucleolus precursor body (NPB), and thereafter, the PCH

gradually forms chromocenters until the blastocyst stage (Aguirre-Lavin et al., 2012). In

paternal and maternal pronuclei (PN), H3K36me2 was not found in heterochromatin around

NPBs, although H3K36me2 levels were higher in maternal PN (Fig. 7A and S5C). At both

the 2- and 4-cell stages, however, H3K36me2 was in part concentrated in heterochromatin

around NPBs and chromocenters (Fig. 7A and S5C). Indeed, colocalization analysis of

individual nuclei revealed that H3K36me2 at the 2- and 4-cell stages exhibited a higher

correlation with DNA compared to other stages (Fig. 7B). Although H3K36me2 was not

concentrated on PCH in most cells during the morula and the blastocyst stages, a small

number of nuclei in blastocyst inner cell mass exhibited some enrichments in PCH (Fig. 7A

and 7B). These results indicate that H3K36me2 can be enriched in heterochromatin at

limited developmental stages in vivo. As the heterochromatin in mammalian preimplantation

embryos is transcribed and dynamically reorganized, the deposition of H3K36me2 might be

involved in these processes.

Discussion

Here we showed that H3K36me2, typically associated with euchromatin, is enriched in

heterochromatin in some mouse cell lines and preimplantation embryos at the 2- to 4-cell

stages. Overexpression and KO experiments revealed that H3K36me2 localization at PCH is

regulated through NSD2, SMARCA5, and BAZ1B or BAZ2A.

Mechanisms guiding H3K36me2 to heterochromatin

PCH histone modifications, such as H4K20me3, typically occur in an H3K9me3-dependent

manner (Schotta et al., 2004). The PCH localization of H3K36me2 was, however,

independent of either H3K9me2/3 or DNA methylation. We found that NSD2, the enzyme

responsible for H3K36me2, targets PCH through the ISWI complexes that harbor subunits

with AT-rich DNA binding motifs. We showed that BAZ1B, which forms the WICH with

SMARCA5, has three AT-hook-like motifs intervened by a BrD at its C-terminus. Although

these motifs are not typical AT-hook, they are essential for DNA binding, in agreement with

previous studies showing that variations in the AT-hook core and surrounding amino acids

are somewhat permissive (Aravind and Landsman, 1998; Filarsky et al., 2015). Additionally,

BAZ2A, which forms the NoRC with SMARCA5, was also localized to heterochromatin and

assists in PCH H3K36me2 localization. BAZ2A has been reported to bind to DNA through its

Tip5/ARBP/MBD (TAM) domain with AT-hook motifs (Chen et al., 2021; Filarsky et al.,

2015). Hence, these ISWI complexes may redundantly function in recruiting NSD2 for PCH

H3K36me2 localization. Even though BAZ1A could target PCH via its N-terminal WAC

domain (Tate et al., 1998), we did not observe PCH localization of emGFP-BAZ1A.

In Pmi28 cells, which display euchromatic H3K36me2 localization and express relatively low

levels of BAZ1B and NSD2, the ectopically expressed emGFP-BAZ1B localized to PCH

perhaps via its AT-hook-like motifs that bind to DNA. However, as BAZ family proteins

possess various chromatin binding domains such as PHD and BrD, their binding affinity and

specificity to DNA and histones could be modulated by chromatin modification context

(Tallant et al., 2015; Oppikofer et al., 2017), which may in turn result in diverse localization in

different cell types. The efficiency of ISWI complexes targeting PCH may also depend on the

cell cycle since both BAZ1B and SMARCA5 exhibit heterochromatin localization from mid to

late S phase (Collins et al., 2002; Bozhenok et al., 2002; Poot et al., 2004). The association

of BAZ1B with newly replicated chromatin could be mediated through specific histone

modifications and/or increased DNA accessibility during replication since the AT-hook-like

motifs bind to naked DNA more strongly than to nucleosomes.

Of note, a recent report suggests that the histone methyltransferase DOT1L introduces

H3K79me3 to PCH in a SMARCA5-dependent manner in mouse ESCs (Malla et al., 2023).

Although the SMARCA5's partner in this process remains unidentified, the ISWI complexes

might decorate PCH by attracting various modifying enzymes.

Physiological relevance of H3K36me2 on heterochromatin

The relatively higher expression of NSD2, SMARCA5, and BAZ1B appears to be associated

with H3K36me2 PCH localization in mouse cell lines, but there was no obvious common

feature among cells showing PCH H3K36me2; for example, whereas PCH H3K36me2 in ESCs suggests its relevance to pluripotency, differentiated fibroblasts also showed the same localization. To explore the physiological relevance of PCH H3K36me2, we examined adult mouse tissues and embryos. So far, heterochromatic H3K36me2 was exclusively observed in preimplantation embryos, particularly at the 2- and 4-cell stages. A minor fraction of cells in blastocysts also exhibited PCH H3K36me2 might be related to ESCs. In mice, the major zygotic genome activation undergoes at the 2-cell stage, including transient transcription from major satellite repeat (Probst et al., 2010), rRNA (Yu et al., 2021), retrotransposons LINE (Jachowicz et al., 2017), and MERVL (Sakashita et al., 2023). Appropriate transcription from these repeats is important for normal development. PCH H3K36me2 at these stages may fine-tune heterochromatin transcription as previously suggested (Frescas et al., 2008; Suzuki et al., 2016; Chaouch et al., 2021). H3K36me2 may also play a role in heterochromatin reorganization in embryogenesis mediated through the perinucleolar ring-shaped structures at the 2-cell stage to more typical chromocenters at the later stages. A very recent study has shown a model that H3K9me3 and H3K36 methylation tandemly promote DNA methylation (Sinha et al., 2023). DNA methylation in oocytes is also regulated by H3K36me2 and H3K36me3 (Yano et al., 2022). As both H3K9me3 and DNA methylation are dynamically changed during these stages, H3K36me2, which is targeted to heterochromatin independently of H3K9me3, might play a critical role in DNA methylation on heterochromatin repeat sequences. To support this notion, H3K36me2 localization in ring-shaped heterochromatin was also observed in porcine 2-cell stage embryos, although the zygotic genome activation occurs at the later stages (Diao et al., 2014). H3K36me2/3 is also known to cross-talk with other chromatin modifications, such as antagonistically distributing H3K27me3 (Streubel et al., 2018; Chen et al., 2022), during oocyte maturation and zygotes (Xu et al., Nat Genet, 2019). Transient accumulation of H3K36me2 might prevent H3K27me3 invasion into constitutive heterochromatin before H3K9me3 distribution is established. Further analysis will be required to elucidate the role of H3K36me2 in embryonic development.

Materials and Methods

Plasmid construction
The following Kazusa HaloTag-tagged human cDNA expression vectors were used: NSD2 (FHC00177), NSD3 (FHC00329), SETD2 (FHC00272), SETD3 (FHC12875), SETD7 (FHC00270), SETD8 (FHC06789), ASH2L (FHC01483), SMYD2 (FHC27765), SMYD3

(FHC03496), SETMAR (FHC05461), KDM2A (FHC 00712), KDM3C (FHC36094E), KDM4A

(FHC00602), KDM4B (FHC13945E), KDM4C (FHC00635), and SMARCA5 (FHC01216).

For expressing human BAZ1A, BAZ1B, and BAZ2A, we used the following emerald-green

GFP (emGFP)-tagged vectors from Addgene: BAZ1A (#65371), BAZ1B (#65372), and

BAZ2A (#65373). To construct the super-folder GFP (sfGFP)-tagged SMARCA5 expression

vector, the ORF was amplified by PCR using Halo-SMARCA5 (#FHC01216; Kazusa) as a

template and inserted into sfGFP-N1 vector (#54737; Addgene) digested with EcoRI using

the In-Fusion HD Cloning Kit (Takara Bio). Then, SMARCA5-sfGFP was cloned into the

PB533A-2 vector (Systems Biosciences). To identify the BAZ1B domain required for PCH

localization, enhanced GFP (EGFP)-tagged truncated BAZ1B expression vectors were

constructed. The full-length and domain of interest were amplified by PCR using emGFP-

BAZ1B (#65372; Addgene) as a template and inserted into pEGFP-C2 vector (Takara Bio)

digested with EcoRI using the In-Fusion HD Cloning Kit. One truncated vector without the

PHD finger and bromodomain (BrD) (emGFP-ΔPHD, BrD) was produced by inverse PCR

using emGFP-BAZ1B as a template. To generate a truncated vector with BrD and three

mutated AT-hook-like motifs (EGFP-BrD+AT1-3mut), site-directed mutagenesis was

performed by inverse PCR using EGFP-BrD+AT1-3 as a template. The primers used are

listed in Table S1.

Animal care

Mouse care and experimental procedures were approved by the Institutional Animal

Experiment Committee of the Tokyo Institute of Technology (Application number: D2023004)

and the Animal Care and Use Committee of Kindai University (Application number: KAAT-

22-001). All animal experiments were performed in accordance with institutional and

governmental guidelines. Mice were maintained on a 12:12 dark: light cycle at a constant

temperature of 22–23°C and were fed freely with food and water. For preparing adult mouse

tissue specimens, we used 26-month-old C57BL/6 male mice that were housed in ventilated

cages and kept in a pathogen-free, temperature-controlled facility at the Tokyo Institute of

Technology.

Cell culture

NIH3T3 mouse embryonic fibroblasts (obtained from N. Takagi at Hokkaido University),

MC12 mouse embryonic carcinoma cells (obtained from N. Takagi at Hokkaido University;

Abe et al., 1988), C2C12 myoblasts (obtained from JCRB), and HEK293T cells (obtained

from K. Fujinaga at Sapporo Medical School) were grown in high-glucose Dulbecco's

modified Eagle's medium (DMEM; Nacalai Tesque) with supplements (1%

penicillin/streptomycin [Nacalai Tesque] and 10% fetal bovine serum [FBS; Thermo Fisher

Scientific, Gibco]). Immortalized mouse embryonic fibroblasts (iMEFs; obtained from Y.

Shinkai at RIKEN), H3K9me-related enzymes knockout (5KO) iMEFs (obtained from Y.

Shinkai; Fukuda et al., 2023) were grown in high-glucose DMEM with supplements (1%

penicillin/streptomycin, 10% FBS, 1% Non-essential Amino Acids [NEAA; Wako], 55 µM 2-

Mercaptoethanol [Gibco]). Mouse embryonic stem (ES) cells (TT2; obtained from Y. Shinkai)

were grown in high-glucose DMEM with supplements (1% penicillin/streptomycin, 10% FBS,

1% NEAA, 1% L-glutamine [Gibco], 10^3 units/ml leukemia inhibitory factor [Nacalai Tesque],

100 µM 2-Mercaptoethanol). Triple DNA methyltransferase knockout (TKO) ES cells

(obtained from M. Okano; Tsumura et al., 2006) were grown in high-glucose DMEM with

supplements (1% penicillin/streptomycin, 15% FBS, 1% NEAA, 1 mM sodium pyruvate

[Gibco], 10^3 units/ml leukemia inhibitory factor, 100 µM 2-Mercaptoethanol). Pmi28

myoblasts were grown in Ham's F-10 medium with supplements (1% penicillin/streptomycin,

20% FBS). All of the cells were grown at 37°C under a 5% CO_2 atmosphere.

Generation of Knockout (KO) cells

The CRISPR/Cas9 system was used for KO cell establishment. All guide RNAs (gRNAs)

were designed by CRISPRdirect (https://crispr.dbcls.jp/). The DNA sequences for gRNAs

are listed in Table S1. Annealed oligonucleotides were inserted into pSpCas9(BB)-2A-Puro

(PX459) V2.0 vector (#62988; Addgene) digested with BbsI. iMEFs were transfected with

gRNA vectors using Lipofectamine 3000 (Thermo Fisher Scientific). For the NSD2 knockout,

two gRNAs targeting the PWWP and SET domains were used. The next and 3 days later,

the medium was replaced with the one containing 2 µg/mL puromycin. For single-cell

cloning, puromycin-selected cells were seeded at a density of 500 cells/20 mL in a 15-cm

dish to form colonies. Then, single colonies were picked and transferred to a glass-bottom

(for immunofluorescence) or plastic-bottom (for stock) 96-well plate. To screen KO-candidate

cells for each gene, immunostaining was performed using the following antibodies: anti-

H3K36me2 for NSD2-KO, anti-SMARCA5 for SMARCA5-KO, and anti-BAZ1B for BAZ1B-

KO cells. The depletion of each protein in KO cells was then validated by Western blotting.

Generation of Knockdown (KD) cells

The shRNA sequences were designed using the GPP Web Portal

(https://portals.broadinstitute.org/gpp/public/). shRNA sequences are listed in Table S1. Two

shRNAs were used for each gene knockdown. Annealed oligonucleotides were inserted into

the pLKO.1-Hygro vector (#24150, Addgene) digested with AgeI and EcoRI. The day after

seeding HEK293T cells on 6-cm dishes, pLKO.1-Hygro-based vector, psPAX2 (#12260,

Addgene), and pVSV-G (#PT3343-5, Clontech) vectors were transfected using

Lipofectamine 3000. The following day, the medium was replaced with fresh medium, which

was then collected 48 h post-transfection. Then, iMEFs were incubated with culture

supernatant containing the virus and 4 µg/mL polybrene (Nacalai Tesque) for 24 h. Infected

cells were selected in a medium containing 400 µg/mL hygromycin B Gold (InvivoGen) for 3

days.

Transient transfection and live-cell imaging

For Fig. 2C, S2C, 3C, 3D, 3I, 4H, and 4I, cells were plated on a glass bottom 24-well plate

(AGC Techno Glass). The following day, HaloTag-, sfGFP-, emGFP-, or EGFP-tagged

expression vectors were transfected using Lipofectamine 3000. After 24 h of regrowth, cells

were incubated in a medium containing 2 nM Janelia Fluor 646 (JF646) HaloTag ligand

(Promega) and 0.5 µg/mL Hoechst33342 (Nacalai Tesque) for 30 min. For Fig. S3C, 4C,

S4A, and 5A, cells were only stained with Hoechst33342. After washing twice with

FluoroBrite medium (Thermo Fisher Scientific) containing the supplements, the plate was

positioned on a heated stage (Tokai Hit) at 37°C under 5% CO_2 on a point-scan confocal

microscope (Ti2 with A1 system; Nikon) operated by the built-in software NIS-elements AR

version 5.21.00 with a Plan Apo VC 60x WI DIC N2 (NA 1.2) objective, water immersion lens

(Nikon). Images were acquired using a 405-nm, 488-nm and a 640-nm laser lines (LU-N4;

Nikon), GAsP detectors, a 405/488/561/640 dichroic mirror, and 450/50, 525/50, 595/50,

700/75 emission filters, with the following parameters; Z range, 5 µm; Z step, 1 µm; Pinhole

size, 43.42 µm; Scanner zoom, 1.5x; Scan size, 512 x 512 pixels; and Scan speed, 1

frame/sec.

Immunofluorescence for cultured cells

For immunostaining of histone marks, cells were fixed with 4% formaldehyde in 250 mM

HEPES–NaOH (pH 7.4) and 0.1% Triton X-100 for 5 min at room temperature. After washing

with PBS, cells were permeabilized with 1% Triton X-100 in PBS for 20 min at room

temperature. After blocking with Blocking One-P (Nacalai Tesque) for 20 min at room

temperature, cells were incubated with 2 µg/mL mouse monoclonal antibody directly

conjugated with a fluorescence dye (Alexa Fluor 488, Cy3, or Cy5) and 1 µg/mL

Hoechst33342 overnight at 4°C. After washing with PBS three times, fluorescence images

were collected. Antibodies used are listed in Table S2. For Fig. S1B, cells were stained with

2 µg/mL rabbit anti-H3K36me2 monoclonal antibody, Cy3-conjugated mouse anti-H3K9me3

monoclonal antibody, and 1 µg/mL Hoechst33342 overnight at 4°C, and then incubated with

1 µg/mL Alexa Fluor 488-conjugated anti-rabbit antibody for 1 h at room temperature.

For Fig. 2A, 2B, 2D, 2F, 2G, 3E, 3G, and 3H, 24 h after transfection with HaloTag

expression vectors, cells were fixed with 4% formaldehyde in PBS for 5 min at room

temperature. After washing with PBS, cells were permeabilized with 1% Triton X-100 in PBS

for 20 min at room temperature. After blocking with Blocking One-P (Nacalai Tesque) for 20

min at room temperature, cells were incubated with 2 µg/mL fluorescence dye-conjugated

mouse monoclonal antibody, 2 nM JF646 HaloTag ligand, and 0.5 µg/mL Hoechst33342

overnight at 4°C. For Fig. 4B, 4D, 4F, 4G, S3A, S3B, and S3D, cells were stained without

the HaloTag ligand 24 h after transfection with emGFP-tag expression vectors. For Fig. 6B,

up to three different plasmids were mixed; in the case of using only one or two expression

vectors, the total amount of DNA was equalized using EGFP or Halo vectors. Cells were

fixed 48 h after transfection for immunostaining.

For Fig. 1A, 1B, S1C-E, fluorescence images were collected using a high-resolution

spinning-disk confocal microscope (Ixplore SpinSR; Olympus) operated by the built-in

software CellSens Dimension 3.2, equipped with a UplanApoN 60× OSC2 (NA 1.4)

objective, oil immersion lens at room temperature. Images were acquired using 405-, 488-,

561-, and 640-nm laser lines (OBIS), a 405/488/561/640 dichroic mirror, and 447/60, 525/50,

617/73, and 685/40 emission filters, a scientific complementary metal oxide semiconductor

(sCMOS) camera (ORCA Flash 4.0; Hamamatsu Photonics) with the following parameters; Z

range, 8 µm; Z step, 0.5 µm; Scan size, 2048 x 2048 pixels; Camera adapter magnification,

1x.

For other figures, fluorescence images were collected using a point-scan confocal

microscope (Ti2 with A1 system; Nikon) operated by the built-in software NIS-elements AR

with a Plan Apo VC 60x WI DIC N2 (NA 1.2), water immersion objective lens at room

temperature. Images were acquired using 405-, 488-, 561-, and 640-nm laser lines (LU-N4;

Nikon). Other setting parameters were the same as described in "Transient transfection and

live-cell imaging."

Immunofluorescence for mouse tissue and E13.5 embryo sections

Mice were euthanized by cervical dislocation. Tissues of interest and E13.5 embryos were

isolated and washed with PBS. Tissues and embryos were fixed in 4% formaldehyde in PBS

overnight at 4°C. After washing with PBS, tissues were soaked in the optimal cutting

temperature compound (SAKURA) and frozen in liquid nitrogen. Frozen tissues were

sectioned at 10 µm thick using a microtome (CM3050S; Leica). The sections were attached

to glass slides, washed with PBS-T (0.05% Tween20 in PBS), and heated at 95°C for 20 min

in 10mM citric acid (pH2.0) for antigen retrieval using a decloaking chamber NxGen

(FUNAKOSHI). After washing with PBS-T, the sections were permeabilized with 1% Triton

X-100 in PBS for 20 min, washed with PBS-T, and incubated with Blocking One-P for 20 min

at room temperature. Then, the sections were stained with anti-H3K36me2 Fab conjugated

with Alexa Fluor 488 (20 µg/mL) and the anti-H3K9me3 antibody conjugated with Cy3 (for

adult mouse sections) or with Cy5 (for embryo sections) (10 µg/mL), and 1 µg/mL

Hoechst33342 for three days at room temperature. After washing with PBS-T, the samples

were mounted with Prolong Diamond (Thermo Fisher Scientific) and covered with a coverslip

(MICRO COVER GLASS 24x32mm, MATSUNAMI).

Fluorescence images for adult tissue sections were acquired using a point-scan confocal

microscope (Ti2 with A1 system; Nikon) with a Plan Apo VC 60x WI DIC N2 (NA 1.2)

objective lens as described above except Zoom factor (1x). Images were acquired using

405- 488-, 561-, and 640-nm laser lines (LU-N4; Nikon). Other setting parameters were the

same as described in "Transient transfection and live-cell imaging."

Fluorescence images for embryo sections were acquired using a spinning disk confocal

microscope (CSU-W1; Yokogawa) operated by the built-in software NIS-elements v5.11.03,

equipped with an inverted microscope (Ti-E; Nikon), EM-CCD (iXon 3; Andor), a laser

illumination system (LDI-NIR; Chroma Technology Japan), and a Plan Apo 40x/0.95 DIC

M/N2 dry objective lens at room temperature. Images were acquired using 405-, 470-, and

640-nm laser lines, a 405/470/555/640 dichroic mirror, and 440/40, 520/60, and 690/50

emission filters, with the following parameters; Z range, 5 µm; Z step, 1 µm; Scan size, 1024

x 1024 pixels.

In vitro fertilization and immunofluorescence of mouse preimplantation embryos

For preparing preimplantation embryos, mice were euthanized with cervical dislocation

immediately before oocyte and sperm collection. Cumulus-intact oocytes were collected in

0.2 mL of TYH medium and inseminated with capacitated sperm at a final concentration of

75 sperm/µL. After 1.5 h incubation at 37 °C under 5% CO_2 in the air, cumulus cells were

dispersed by a brief treatment with hyaluronidase (Type-IS, 150 units/mL, Sigma). The

denuded fertilized oocytes were transferred to KSOMaa medium and incubated until they

reached the appropriate stages.

The embryos were fixed with 3.7% paraformaldehyde containing 0.2% Triton X-100 for 20

min at room temperature. After fixation, the embryos were washed twice with PBS containing

3% bovine serum albumin (BSA) and placed for 60 min at room temperature for blocking.

The embryos were then incubated with anti-H3K9me3 antibody conjugated with Alexa Fluor

488 (10 µg/mL) and anti-H3K36me2 antibody conjugated with Cy3 (5 µg/mL) dissolved in

the above blocking buffer at 4 °C for at least 8 h. After washing with PBS, the embryos were

transferred to PBS containing 0.5 µg/mL of DAPI (Nacalai Tesque) in a glass-bottomed dish

(Mat Tek, P35G-1.5-14-C).

The fluorescent observations were performed using a spinning disk based-super resolution

confocal microscopic system (CSU-W1 SoRa; Yokogawa Electric Corp.; Hatano et al., 2022)

equipped with an inverted microscopy (IX73; Olympus) with a 40× silicone immersion

objective lens (UPLSAPO40XS: NA 1.25; Olympus) and an intermediate 4× tube lens.

Three-color fluorescence images in 100 different focal planes at 0.5 µm intervals were

captured with 405-, 488-, and 561-nm laser lines. The images were passed through

emission filters (FF02-460/80 for DAPI, FF01-520/35 for Alexa488, and FF02-617/73 for

Cy3; Yokogawa) and captured by a Prime95B sCMOS camera (Teledyne Photometrics) with

de-noised switch. The system was controlled and images were taken by µ-Manager

microscopy software (Edelstein et al., 2014).

Image analysis

Line profiles were drawn using NIS-Elements AR Analysis version 5.01.00 (Nikon) and

Image J Fiji version 1.54f (https://fiji.sc/). A merged image from single Z-slice images of

different fluorescence channels was created using NIS-Elements. Using the "Plot Profile"

function in Fiji, a straight line was drawn on the image, allowing for the measurement of both

position and intensities. A line profile plot was then generated using R version 4.2.1

(https://www.r-project.org/), with the x-axis representing the distance (µm) and the y-axis

representing the intensity.

Intensity measurements were performed using NIS-Elements (Nikon) and Aivia version

10.5.1 (DRVision). After subtracting the background with NIS-Elements, the Aivia "Cell

Analysis" recipe was run to define regions for chromocenters and cell nuclei based on

Hoechst signals, and then mean pixel intensities were measured. Fold enrichment in PCH

(the mean intensity in the chromocenter region divided by that of the nucleus excluding the

chromocenter) was calculated using R.

Pearson correlation coefficients between two fluorescence intensities in single nuclei were

calculated using the Image J Fiji "Coloc2" function. Plots and significance tests were

performed using R.

Quantitative RT-PCR (RT-qPCR)

Total RNA was isolated from cells using TRIzol (Thermo Fisher Scientific) and further
purified using an RNeasy MinElute Cleanup Kit (Qiagen) according to the manufacturer's
instructions. cDNA was synthesized by reverse transcription using Superscript III Reverse
Transcriptase (Thermo Fisher Scientific) and random hexamer primers. cDNA was mixed
with PowerUP SYBR Green Master Mix (Applied BioSystems) and forward and reverse
primers (Table S1) in 96-well format tubes, before setting to a real-time PCR system
(Agilent) with the following parameters: Initial hold, 50ºC for 2 min and 95ºC for 2 min; PCR,
(95ºC for 15 sec and 60ºC for 60 sec) x 40 cycles; Dissociation curve, 95ºC for 60 sec, 55ºC
for 30 sec, and 95°C for 30 sec. For relative quantification, standard curves were prepared
for the target genes and the reference gene (Gapdh) using cDNA from NIH3T3 cells. The
initial template amounts of the target and reference genes were calculated based on the Ct
values, and the expression levels of the target genes were relative to the reference gene.
Two reaction tubes were prepared for each of the two biological replicates.
669

Protein extraction and immunoprecipitation
HEK293T cells grown on two 10-cm dishes were transfected with an expression vector for
EGFP-BrD+AT1-3 or EGFP-BrD+AT1-3mut using Lipofectamine 2000 (Thermo Fisher
Scientific). The next day, cells were washed with PBS, trypsinized, and collected in a 15-mL
tube by centrifugation (1,200 rpm for 2 min). The cell pellet was suspended in ice-cold PBS,
transferred to a 1.5 mL tube, and washed with ice-cold PBS three times, before lysis in Lysis
buffer (300 mM NaCl, 10 mM HEPES-NaOH [pH7.4], 0.1% TritonX-100, 1× Protease
inhibitor cocktail [Nacalai Tesque]). After centrifugation (20,000 x g for 20 min at 4°C), the
supernatant was transferred to a new 1.5 mL tube. GFP-Trap Magnetic Agarose beads
(Proteintech; 80 µL) were prewashed with wash buffer (10 mM Tris-HCl [ph7.5], 150 mM
NaCl, 0.5 mM EDTA) in a 1.5 mL tube. After collecting beads using a magnetic rack (Thermo
Fisher Scientific), the wash buffer was removed. This washing step was repeated a total of
three times. The cell lysate was then added to the GFP-Trap beads and incubated at 4°C
overnight with gentle mixing. After washing the beads three times with wash buffer, they
were resuspended in the same buffer. Small aliquots were collected from the input cell
lysate, washed supernatant, and bead suspension for SDS-polyacrylamide gel
electrophoresis to evaluate purity and quantity. Aliquots were mixed with an equal volume of
2×SDS-gel loading buffer (4% SDS, 20% glycerol, 120 mM Tris-HCl [pH 6.8], 100 mM DTT,
0.01% bromophenol blue) and heated at 95°C for 5 min. Proteins were then separated on a
15% polyacrylamide gel (Supersep Ace, Fujifilm) alongside a dilution series of BSA for
quantitation. The concentrations of the purified EGFP-BrD+AT1-3 or EGFP-BrD+AT1-3mut
proteins were measured using a standard curve based on known amounts of BSA.
692

DNA-binding assay

AT-rich Major Satellite repeats were amplified by PCR using specific primers (Table S2) and NIH3T3-derived DNA as a template. PCR products were electrophoresed on an agarose gel, and ~250 bp fragments were gel extracted for cloning using a PCR Cloning Kit (NEB). The resulting plasmid was purified by Midiprep (Thermo Fisher Scientific), and the insert sequence was determined by Sanger sequencing. After EcoRI digestion and agarose gel electrophoresis, the insert Major Satellite repeats (289 bp) were gel extracted and purified using a Gel-extraction kit (Qiagen). For preparing GC-rich DNA fragments, the EF1α promoter flanking sequence (78 bp, 67% GC contents) from the PB533A-Halo vector was utilized by AsiSI and MfeI digestion and gel extraction. The preparation of Widom 601 DNA and nucleosome was described previously (Arimura et al., 2012; Vasudevan et al., 2010; Kujirai et al., 2018). Nucleotide sequences used for binding assay are shown in Fig. S4C-E.

For the DNA binding assay, GFP-Trap Magnetic Agarose beads that captured EGFP-BrD+AT1-3 and EGFP-BrD+AT1-3mut proteins were prewashed with High-salt buffer (1 M NaCl, 20 mM Tris-HCl [pH 7.5], 1 mM DTT) to eliminate any residual genomic DNA, before washing twice with Reaction buffer (20 mM Tris-HCl [pH7.5], 1 mM DTT). After removing the Reaction buffer, DNA or nucleosomes (0.28 pmol) in 10 μL of Reaction buffer were added to the beads (0.14 pmol), and the mixture was incubated for 30 min at room temperature with gentle rotation. The tubes were placed on a magnetic rack, and the supernatant was collected. Beads were washed with Wash buffer (50 mM NaCl, 20 mM Tris-HCl [pH7.5], 1 mM DTT) for 5 min at room temperature before the supernatant was collected. Then, Elution buffer (350 mM NaCl, 20 mM Tris-HCl [pH 7.5], 1 mM DTT) was added, incubated at room temperature for 5 min, and the supernatant was collected. All supernatants were electrophoresed on a 2% agarose gel and stained with Gel Red (Biotium).

Western blotting

The day after cells were plated on a 6-cm dish, cells were washed three times with ice-cold PBS and 500 μL of 2×Laemmli buffer (4% SDS, 20% glycerol, 120 mM Tris-HCl [pH 6.8]) was added. Cell lysate was harvested using a cell lifter (Corning) and transferred to a 1.5 mL tube. After mixing with DTT (0.1 M) and bromophenol blue (0.01%), tubes were heated at 95°C for 10 min, proteins were separated in 7.5% SDS-polyacrylamide gels (SuperSep™ Ace; Fujifilm) and transferred to polyvinylidene fluoride membranes (FluoroTrans W; Pall) with EzFastBlot transfer buffer (ATTO) using a transfer system (Trans-Blot Turbo; Bio-Rad). After washing with TBS-T (10 mM Tris-HCl [pH 8.0], 0.1% Tween-20, 150 mM NaCl), membranes were incubated with Blocking One (Nacalai Tesque) for 20 min and then incubated with primary antibodies diluted in Can-Get-Signal Solution I (TOYOBO) for 2 h at

room temperature. After washing in TBS-T for 10 min three times, membranes were

incubated with horse radish peroxidase-labeled secondary antibody (Jackson

ImmunoResearch) diluted in Can-Get-Signal Solution II (TOYOBO) for 1 h at room

temperature. After washing membranes with TBS-T for 10 min three times,

chemiluminescence signals were developed using ImmunoStar LD (Wako) and detected

using a LuminoGraph II Chemiluminescent Imaging System (ATTO). Antibodies used are

listed in Table S2.

ChIP-seq data analysis

ChIP-seq dataset of H3K36me2 and H3K9me3 were obtained from the National Center for

Biotechnology Information Sequence Read Archive (NCBI SRA) for ESCs (PRJNA720715),

B cells (PRJNA324130), and pancreatic cancer cells (PRJNA565773). The fastq data were

preprocessed with fastp ver. 0.23.2 and mapped onto the mouse mm39 genome using

bowtie2 ver. 2.4.5 with the "very-sensitive-local" option. After removing unmapped read data

using sambamba ver. 0.6.6, duplicated reads were removed using the markdup function in

samtools ver. 1.13. The mapped data were utilized to generate bigwig files using

bamCoverage function of deepTools ver. 3.2.1 with binSize of 20 and RPKM normalization.

The correlations among the ChIP-seq data were calculated for consecutive genome bins

using multiBigwigSummary bins function with 1,000 binSize, and the Spearman correlation

was visualized with deepTools plotCorrelation.

Regarding transposable elements, RepeatMasker output for the mouse mm39 genome

(accessed on May 14, 2023) was obtained from the UCSC Genome Browser. For each class

of SINE, LINE, LTR retrotransposons, and DNA transposons, relatively young copies (<10%

divergence from consensus) that were nearly full-length (>90% in length of consensus) were

selected for subsequent analysis. The deepTools computeMatrix tool was used to calculate

the accumulation of ChIP-seq signals within ±3 kbp regions of transcription start sites (TSS)

or the TSS-TES (transcription end sites) of the transposable elements. The resulting data

were visualized using the deepTools plotHeatmap tool.

Amino acid alignments

BAZ1B amino acid sequences of H. sapiens (NP_115784.1), M. musculus (NP_035844.2),

G. gallus (XP_001233717.3), X. tropicalis (XP_004911823.1), and D. rerio

(XP_003200529.2) were aligned using GENETYX version 11 software (Genetyx).

Statistical analysis

Statistical significance tests were performed using R. For comparisons between two groups, the Mann-Whitney U test was applied. For comparisons of three or more groups, the Mann-Whitney U test with Hommel correction was used.

Artificial Intelligence

For English proofreading, Grammarly and ChatGPT-4 (Open AI), with a prompt "The following document is a part of a scientific paper. Please proofread and correct any grammatical errors.", were in part used. No original sentences were generated using AI chatbots.

Online supplemental material

Fig. S1 shows H3K36me2 distribution in various mouse cells, related to Fig. 1. Fig. S2 shows the results of screening of factors that are involved in PCH H3K36me2 localization, including the localization of H3K36me2 in cells lacking H3K9me3 and DNA methylation and the localization of Halo-tagged methyltransferases and demethylase, related to Fig. 2. Fig. S3 shows the localization of BAZ1A and BAZ2A, related to Fig. 4. Fig. S4 shows the details for BAZ1B-DNA interaction, including the localization of various truncated proteins, purification of the DNA binding domain, and DNA sequences used for the binding assay, related to Fig. 5. Fig. S5 shows immunofluorescence data for mouse adult tissue and embryo sections, and line plots for zygote and 2-cell embryos. Tables S1 and S2 list oligonucleotides and antibodies with the conditions used in this study, respectively.

Acknowledgments

We thank Y. Shinkai, M. Okano, N. Takagi, and K. Fujinaga for providing cell lines and members of the Kimura and Yamagata Labs for help and discussion, especially H. Oda for assisting with an imaging analysis and Y. Kono and T. Shimi for instructing the lentivirus system. We also thank the Center for Integrative Biosciences and the Biomaterials Analysis Division, Open Facility Center at Tokyo Institute of Technology for DNA sequencing.

This work was supported by Japan Society for the Promotion of Science KAKENHI (JP17H01417 and JP21H04764 to H. Kimura, JP18H05528 to K. Yamagata, JP20H05690 to T. Kujirai, JP23H05475 to H. Kurumizaka, and JP22K06338 to H. Nishihara), Japan Science and Technology Agency CREST (JPMJCR20S6 to Y. Sato and JPMJCR16G1 to H. Kimura) and ERATO (JPMJER1901 to H. Kurumizaka), and Japan Agency for Medical Research and Development (AMED) Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS) (JP22ama121020 to H. Kimura and JP23ama121009 to H. Kurumizaka).

The authors declare no competing financial interests.

Author contributions: Conceptualization, N. Goto, H. Kimura; investigation, N. Goto, K. Suke,

N. Yonezawa, H. Nishihara; formal analysis, N. Goto; methodology, Y. Sato; resources, T.

Kujirai, H. Kurumizaka; writing – original draft, N. Goto; writing – review and editing, K. Suke,

N. Yonezawa, Y. Sato, H. Nishihara, K. Yamagata, H. Kimura; funding acquisition, Y. Sato,

T. Kujirai, H. Kurumizaka, K. Yamagata, H. Kimura; supervision, H. Kurumizaka, K.

Yamagata, H. Kimura.

References

Abe, K., N. Takagi, and M. Sasaki. 1988. Nonhistone nuclear proteins specific to certain mouse embryonal carcinoma clones having an inactive X chromosome. *Exp. Cell Res.* 179:590–594. doi:10.1016/0014-4827(88)90297-2.

Aguirre-Lavin, T., P. Adenot, A. Bonnet-Garnier, G. Lehmann, R. Fleurot, C. Boulesteix, P. Debey, and N. Beaujean. 2012. 3D-FISH analysis of embryonic nuclei in mouse highlights several abrupt changes of nuclear organization during preimplantation development. *BMC Dev. Biol.* 12:30. doi:10.1186/1471-213X-12-30.

Altemose, N., G.A. Logsdon, A. V. Bzikadze, P. Sidhwani, S.A. Langley, G. V. Caldas, S.J. Hoyt, L. Uralsky, F.D. Ryabov, C.J. Shew, M.E.G. Sauria, M. Borchers, A. Gershman, A. Mikheenko, V.A. Shepelev, T. Dvorkina, O. Kunyavskaya, M.R. Vollger, A. Rhie, A.M. McCartney, M. Asri, R. Lorig-Roach, K. Shafin, J.K. Lucas, S. Aganezov, D. Olson, L.G. de Lima, T. Potapova, G.A. Hartley, M. Haukness, P. Kerpedjiev, F. Gusev, K. Tigyi, S. Brooks, A. Young, S. Nurk, S. Koren, S.R. Salama, B. Paten, E.I. Rogaev, A. Streets, G.H. Karpen, A.F. Dernburg, B.A. Sullivan, A.F. Straight, T.J. Wheeler, J.L. Gerton, E.E. Eichler, A.M. Phillippy, W. Timp, M.Y. Dennis, R.J. O'Neill, J.M. Zook, M.C. Schatz, P.A. Pevzner, M. Diekhans, C.H. Langley, I.A. Alexandrov, and K.H. Miga. 2022. Complete genomic and epigenetic maps of human centromeres. *Science.* 376:eabl4178. doi:10.1126/science.abl4178.

Aravind, L., and D. Landsman. 1998. AT-hook motifs identified in a wide variety of DNA-binding proteins. *Nucleic Acids Res.* 26:4413–4421. doi:10.1093/nar/26.19.4413.

Arimura, Y., H. Tachiwana, T. Oda, M. Sato, and H. Kurumizaka. 2012. Structural analysis of the hexasome, lacking one histone H2A/H2B dimer from the conventional nucleosome. *Biochemistry.* 51:3302–3309. doi:10.1021/bi300129b.

Barral, A., G. Pozo, L. Ducrot, G.L. Papadopoulos, S. Sauzet, A.J. Oldfield, G. Cavalli, and J. Déjardin. 2022. SETDB1/NSD-dependent H3K9me3/H3K36me3 dual heterochromatin maintains gene expression profiles by bookmarking poised enhancers. *Mol. Cell.* 82:816-832.e12. doi:10.1016/j.molcel.2021.12.037.

Bartholomew, B. 2014. ISWI chromatin remodeling: One primary actor or a coordinated effort? *Curr. Opin. Struct. Biol.* 24:150–155. doi:10.1016/j.sbi.2014.01.010.

Blackledge, N.P., J.C. Zhou, M.Y. Tolstorukov, A.M. Farcas, P.J. Park, and R.J. Klose. 2010. CpG Islands Recruit a Histone H3 Lysine 36 Demethylase. *Mol. Cell.* 38:179–190. doi:10.1016/j.molcel.2010.04.009.

Borgel, J., M. Tyl, K. Schiller, Z. Pusztai, C.M. Dooley, W. Deng, C. Wooding, R.J. White, T. Warnecke, H. Leonhardt, E.M. Busch-Nentwich, and T. Bartke. 2017. KDM2A integrates DNA and histone modification signals through a CXXC/PHD module and direct interaction with HP1. *Nucleic Acids Res.* 45:1114–1129. doi:10.1093/nar/gkw979.

Bozhenok, L., P.A. Wade, and P. Varga-Weisz. 2002. WSTF-ISWI chromatin remodeling complex targets heterochromatic replication foci. *EMBO J.* 21:2231–2241. doi:10.1093/emboj/21.9.2231.

Bressan, R.B., B. Southgate, K.M. Ferguson, C. Blin, V. Grant, N. Alfazema, J.C. Wills, M.A.
Marques-Torrejon, G.M. Morrison, J. Ashmore, F. Robertson, C.A.C. Williams, L.
Bradley, A. von Kriegsheim, R.A. Anderson, S.R. Tomlinson, and S.M. Pollard. 2021.
Regional identity of human neural stem cells determines oncogenic responses to
histone H3.3 mutants. *Cell Stem Cell.* 28:877-893.e9. doi:10.1016/j.stem.2021.01.016.

Carrozza, M.J., B. Li, L. Florens, T. Suganuma, S.K. Swanson, K.K. Lee, W.J. Shia, S.
Anderson, J. Yates, M.P. Washburn, and J.L. Workman. 2005. Histone H3 methylation
by Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious
intragenic transcription. *Cell.* 123:581–592. doi:10.1016/j.cell.2005.10.023.

Chantalat, S., A. Depaux, P. Héry, S. Barral, J.Y. Thuret, S. Dimitrov, and M. Gérard. 2011.
Histone H3 trimethylation at lysine 36 is associated with constitutive and facultative
heterochromatin. *Genome Res.* 21:1426–1437. doi:10.1101/gr.118091.110.

Chaouch, A., J. Berlandi, C.C.L. Chen, F. Frey, S. Badini, A.S. Harutyunyan, X. Chen, B.
Krug, S. Hébert, A. Jeibmann, C. Lu, C.L. Kleinman, M. Hasselblatt, P. Lasko, M.
Shirinian, and N. Jabado. 2021. Histone H3.3 K27M and K36M mutations de-repress
transposable elements through perturbation of antagonistic chromatin marks. *Mol. Cell.*
81:4876-4890.e7. doi:10.1016/j.molcel.2021.10.008.

Chen, E.S., K. Zhang, E. Nicolas, H.P. Cam, M. Zofall, and S.I.S. Grewal. 2008. Cell cycle
control of centromeric repeat transcription and heterochromatin assembly. *Nature.*
451:734–737. doi:10.1038/nature06561.

Chen, H., B. Hu, C. Horth, E. Bareke, P. Rosenbaum, S.Y. Kwon, J. Sirois, D.N. Weinberg,
F.M. Robison, B.A. Garcia, C. Lu, W.A. Pastor, and J. Majewski. 2022. H3K36
dimethylation shapes the epigenetic interaction landscape by directing repressive
chromatin modifications in embryonic stem cells. *Genome Res.* 32:825–837.
doi:10.1101/gr.276383.121.

Chen, S., M. Zhou, A. Dong, P. Loppnau, M. Wang, J. Min, and K. Liu. 2021. Structural basis
of the TAM domain of BAZ2A in binding to DNA or RNA independent of methylation
status. *J. Biol. Chem.* 297:101351. doi:10.1016/j.jbc.2021.101351.

Chen, Y., Y. Zhang, Y. Wang, L. Zhang, E.K. Brinkman, S.A. Adam, R. Goldman, B. Van
Steensel, J. Ma, and A.S. Belmont. 2018. Mapping 3D genome organization relative to
nuclear compartments using TSA-Seq as a cytological ruler. *J. Cell Biol.* 217:4025–
4048. doi:10.1083/jcb.201807108.

Clapier, C.R., J. Iwasa, B.R. Cairns, and C.L. Peterson. 2017. Mechanisms of action and
regulation of ATP-dependent chromatin-remodelling complexes. *Nat. Rev. Mol. Cell*
Biol. 18:407–422. doi:10.1038/nrm.2017.26.

Collins, N., R.A. Poot, I. Kukimoto, C. García-Jiménez, G. Dellaire, and P.D. Varga-Weisz.
2002. An ACF1-ISWI chromatin-remodeling complex is required for DNA replication
through heterochromatin. *Nat. Genet.* 32:627–632. doi:10.1038/ng1046.

Culver-Cochran, A.E., and B.P. Chadwick. 2012. The WSTF-ISWI Chromatin Remodeling
Complex Transiently Associates with the Human Inactive X Chromosome during Late
S-Phase Prior to BRCA1 and γ-H2AX. *PLoS One.* 7:1–15.
doi:10.1371/journal.pone.0050023.

Diao, Y.F., R.K. Oqani, X.X. Li, T. Lin, J.W. Kang, and D. Il Jin. 2014. Changes in histone H3
lysine 36 methylation in porcine oocytes and preimplantation embryos. *PLoS One*. 9.
doi:10.1371/journal.pone.0100205.

Edelstein, A.D., M.A. Tsuchida, N. Amodaj, H. Pinkard, R.D. Vale, and N. Stuurman. 2014.
Advanced methods of microscope control using μManager software. *J. Biol. Methods*.
1:e10. doi:10.14440/jbm.2014.36.

Filarsky, M., K. Zillner, I. Araya, A. Villar-Garea, R. Merkl, G. Längst, and A. Németh. 2015.
The extended AT-hook is a novel RNA binding motif. *RNA Biol*. 12:864–876.
doi:10.1080/15476286.2015.1060394.

Frescas, D., D. Guardavaccaro, S.M. Kuchay, H. Kato, A. Poleshko, V. Basrur, K.S.
Elenitoba-Johnson, R.A. Katz, and M. Pagano. 2008. KDM2A represses transcription of
centromeric satellite repeats and maintains the heterochromatic state. *Cell Cycle*.
7:3539–3547. doi:10.4161/cc.7.22.7062.

Fukuda, K., T. Shimi, C. Shimura, T. Ono, T. Suzuki, K. Onoue, S. Okayama, H. Miura, I.
Hiratani, K. Ikeda, Y. Okada, N. Dohmae, S. Yonemura, A. Inoue, H. Kimura, and Y.
Shinkai. 2023. Epigenetic plasticity safeguards heterochromatin configuration in
mammals. *Nucleic Acids Res*. 51:6190–6207. doi:10.1093/nar/gkad387.

Georgescu, P.R., M. Capella, S. Fischer-Burkart, and S. Braun. 2020. The euchromatic
histone mark H3K36me3 preserves heterochromatin through sequestration of an
acetyltransferase complex in fission yeast. *Microb. Cell*. 7:80–92.
doi:10.15698/mic2020.03.711.

Guan, H., P. Wang, P. Zhang, C. Ruan, Y. Ou, B. Peng, X. Zheng, J. Lei, B. Li, C. Yan, and
H. Li. 2023. Diverse modes of H3K36me3-guided nucleosomal deacetylation by Rpd3S.
Nature. 620. doi:10.1038/s41586-023-06349-1.

Guetg, C., P. Lienemann, V. Sirri, I. Grummt, D. Hernandez-Verdun, M.O. Hottiger, M.
Fussenegger, and R. Santoro. 2010. The NoRC complex mediates the heterochromatin
formation and stability of silent rRNA genes and centromeric repeats. *EMBO J.*
29:2135–2146. doi:10.1038/emboj.2010.17.

Hasan, N., and N. Ahuja. 2019. The emerging roles of atp-dependent chromatin remodeling
complexes in pancreatic cancer. 11.

Hatano, Y., D. Mashiko, M. Tokoro, T. Yao, and K. Yamagata. 2022. Chromosome counting
in the mouse zygote using low-invasive super-resolution live-cell imaging. *Genes to
Cells*. 27:214–228. doi:10.1111/gtc.12925.

Hayashi-Takanaka, Y., K. Yamagata, T. Wakayama, T.J. Stasevich, T. Kainuma, T.
Tsurimoto, M. Tachibana, Y. Shinkai, H. Kurumizaka, N. Nozaki, and H. Kimura. 2011.
Tracking epigenetic histone modifications in single cells using Fab-based live
endogenous modification labeling. *Nucleic Acids Res*. 39:6475–6488.
doi:10.1093/nar/gkr343.

Huang, X., R.D. LeDuc, L. Fornelli, A.J. Schunter, R.L. Bennett, N.L. Kelleher, and J.D.
Licht. 2019. Defining the NSD2 interactome: PARP1 PARylation reduces NSD2 histone
methyltransferase activity and impedes chromatin binding. *J. Biol. Chem.* 294:12459–
12471. doi:10.1074/jbc.RA118.006159.

Ito, T., M.E. Levenstein, D. V. Fyodorov, A.K. Kutach, R. Kobayashi, and J.T. Kadonaga.
1999. ACF consists of two subunits, Acf1 and ISWI, that function cooperatively in the
ATP-dependent catalysis of chromatin assembly. *Genes Dev.* 13:1529–1539.
doi:10.1101/gad.13.12.1529.

Jachowicz, J.W., X. Bing, J. Pontabry, A. Bošković, O.J. Rando, and M.E. Torres-Padilla.
2017. LINE-1 activation after fertilization regulates global chromatin accessibility in the
early mouse embryo. *Nat. Genet.* 49:1502–1510. doi:10.1038/ng.3945.

Jack, A.P.M., S. Bussemer, M. Hahn, S. Pünzeler, M. Snyder, M. Wells, G. Csankovszki, I.
Solovei, G. Schotta, and S.B. Hake. 2013. H3K56me3 Is a Novel, Conserved
Heterochromatic Mark That Largely but Not Completely Overlaps with H3K9me3 in
Both Regulation and Localization. *PLoS One.* 8. doi:10.1371/journal.pone.0051765.

Kujirai, T., Y. Arimura, R. Fujita, N. Horikoshi, S. Machida, and H. Kurumizaka. 2018.
Methods for Preparing Nucleosomes Containing Histone Variants BT   - Histone
Variants: Methods and Protocols. G.A. Orsi and G. Almouzni, editors. Springer New
York, New York, NY. 3–20.

Lachner Monika, O'Carroll Do¬nal, Rea Stephen, Mechtler Karl, and Jenuwein Thomas.
2001. Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature.*
410.

Lange, U.C., S. Siebert, M. Wossidlo, T. Weiss, C. Ziegler-Birling, J. Walter, M.E. Torres-
Padilla, S. Daujat, and R. Schneider. 2013. Dissecting the role of H3K64me3 in mouse
pericentromeric heterochromatin. *Nat. Commun.* 4. doi:10.1038/ncomms3233.

Li, H., S. Ilin, W. Wang, E.M. Duncan, J. Wysocka, C.D. Allis, and D.J. Patel. 2006.
Molecular basis for site-specific read-out of histone H3K4me3 by the BPTF PHD finger
of NURF. *Nature.* 442:91–95. doi:10.1038/nature04802.

Lowary, P.T., and J. Widom. 1998. New DNA sequence rules for high affinity binding to
histone octamer and sequence-directed nucleosome positioning. *J. Mol. Biol.* 276:19–
42. doi:10.1006/jmbi.1997.1494.

Malla, A.B., H. Yu, D. Farris, S. Kadimi, T.T. Lam, A.L. Cox, Z.D. Smith, and B.J. Lesch.
2023. DOT1L bridges transcription and heterochromatin formation at mammalian
pericentromeres . *EMBO Rep.* 1–25. doi:10.15252/embr.202256492.

Matsuda, A., Y. Chikashige, D.Q. Ding, C. Ohtsuki, C. Mori, H. Asakawa, H. Kimura, T.
Haraguchi, and Y. Hiraoka. 2015. Highly condensed chromatins are formed adjacent to
subtelomeric and decondensed silent chromatin in fission yeast. *Nat. Commun.* 6:1–3.
doi:10.1038/ncomms8753.

Musselman, C.A., J. Ramírez, J.K. Sims, R.E. Mansfield, S.S. Oliver, J.M. Denu, J.P.
Mackay, P.A. Wade, J. Hagman, and T.G. Kutateladze. 2012. Bivalent recognition of
nucleosomes by the tandem PHD fingers of the CHD4 ATPase is required for CHD4-
mediated repression. *Proc. Natl. Acad. Sci. U. S. A.* 109:787–792.
doi:10.1073/pnas.1113655109.

Nan, X., J. Hou, A. Maclean, J. Nasir, M.J. Lafuente, X. Shu, S. Kriaucionis, and A. Bird.
2007. Interaction between chromatin proteins MECP2 and ATRX is disrupted by

mutations that cause inherited mental retardation. *Proc. Natl. Acad. Sci. U. S. A.*
104:2709–2714. doi:10.1073/pnas.0608056104.

Neri, F., S. Rapelli, A. Krepelova, D. Incarnato, C. Parlato, G. Basile, M. Maldotti, F. Anselmi,
and S. Oliviero. 2017. Intragenic DNA methylation prevents spurious transcription
initiation. *Nature.* 543:72–77. doi:10.1038/nature21373.

Ooga, M., A. Inoue, S.I. Kageyama, T. Akiyama, M. Nagata, and F. Aoki. 2008. Changes in
H3K79 methylation during preimplantation development in mice. *Biol. Reprod.* 78:413–
424. doi:10.1095/biolreprod.107.063453.

Oppikofer, M., M. Sagolla, B. Haley, H.M. Zhang, S.K. Kummerfeld, J. Sudhamsu, E.M.
Flynn, T. Bai, J. Zhang, C. Ciferri, and A.G. Cochran. 2017. Non-canonical reader
modules of BAZ1A promote recovery from DNA damage. *Nat. Commun.* 8.
doi:10.1038/s41467-017-00866-0.

Palasca, O., A. Santos, C. Stolte, J. Gorodkin, and L.J. Jensen. 2018. TISSUES 2.0: An
integrative web resource on mammalian tissue expression. *Database.* 2018:1–12.
doi:10.1093/database/bay003.

Pallier, C., P. Scaffidi, S. Chopineau-Proust, A. Agresti, P. Nordmann, M.E. Bianchi, and V.
Marechal. 2003. Association of Chromatin Proteins High Mobility Group Box (HMGB) 1
and HMGB2 with Mitotic Chromosomes. *Mol. Biol. Cell.* 14:3414–3426.
doi:10.1091/mbc.E02.

Peters, A.H.F.M., S. Kubicek, K. Mechtler, R.J. O'Sullivan, A.A.H.A. Derijck, L. Perez-
Burgos, A. Kohlmaier, S. Opravil, M. Tachibana, Y. Shinkai, J.H.A. Martens, and T.
Jenuwein. 2003. Partitioning and Plasticity of Repressive Histone Methylation States in
Mammalian Chromatin. *Mol. Cell.* 12:1577–1589. doi:10.1016/S1097-2765(03)00477-5.

Poot, R.A., L. Bozhenok, D.L.C. van den Berg, S. Steffensen, F. Ferreira, M. Grimaldi, N.
Gilbert, J. Ferreira, and P.D. Varga-Weisz. 2004. The Williams syndrome transcription
factor interacts with PCNA to target chromatic remodelling by ISWI to replication foci.
Nat. Cell Biol. 6:1236–1244. doi:10.1038/ncb1196.

Popovic, R., E. Martinez-Garcia, E.G. Giannopoulou, Q. Zhang, Q. Zhang, T. Ezponda, M.Y.
Shah, Y. Zheng, C.M. Will, E.C. Small, Y. Hua, M. Bulic, Y. Jiang, M. Carrara, R.A.
Calogero, W.L. Kath, N.L. Kelleher, J.P. Wang, O. Elemento, and J.D. Licht. 2014.
Histone Methyltransferase MMSET/NSD2 Alters EZH2 Binding and Reprograms the
Myeloma Epigenome through Global and Focal Changes in H3K36 and H3K27
Methylation. *PLoS Genet.* 10. doi:10.1371/journal.pgen.1004566.

Prasanth, S.G., Z. Shen, K. V. Prasanth, and B. Stillman. 2010. Human origin recognition
complex is essential for HP1 binding to chromatin and heterochromatin organization.
Proc. Natl. Acad. Sci. U. S. A. 107:15093–15098. doi:10.1073/pnas.1009945107.

Probst, A. V., and G. Almouzni. 2008. Pericentric heterochromatin: Dynamic organization
during early development in mammals. *Differentiation.* 76:15–23. doi:10.1111/j.1432-
0436.2007.00220.x.

Probst, A. V., I. Okamoto, M. Casanova, F. El Marjou, P. Le Baccon, and G. Almouzni. 2010.
A Strand-specific burst in transcription of pericentric satellites is required for

chromocenter formation and early mouse development. *Dev. Cell.* 19:625–638.
doi:10.1016/j.devcel.2010.09.002.

Quénet, D., V. Gasser, L. Fouillen, F. Cammas, S. Sanglier‑Cianferani, R. Losson, and F.
Dantzer. 2008. The histone subcode: poly(ADP‑ribose) polymerase‑1 (Parp‑1) and
Parp‑2 control cell differentiation by regulating the transcriptional intermediary factor
TIF1β and the heterochromatin protein HPIα. *FASEB J.* 22:3853–3865.
doi:10.1096/fj.08-113464.

Rechtsteiner, A., S. Ercan, T. Takasaki, T.M. Phippen, T.A. Egelhofer, W. Wang, H. Kimura,
J.D. Lieb, and S. Strome. 2010. The Histone H3K36 Methyltransferase MES-4 acts
epigenetically to transmit the memory of germline gene expression to progeny. *PLoS*
Genet. 6. doi:10.1371/journal.pgen.1001091.

Ribeyre, C., R. Zellweger, M. Chauvin, N. Bec, C. Larroque, M. Lopes, and A. Constantinou.
2016. Nascent DNA Proteomics Reveals a Chromatin Remodeler Required for
Topoisomerase I Loading at Replication Forks. *Cell Rep.* 15:300–309.
doi:10.1016/j.celrep.2016.03.027.

Sakashita, A., T. Kitano, H. Ishizu, Y. Guo, H. Masuda, M. Ariura, K. Murano, and H. Siomi.
2023. Transcription of MERVL retrotransposons is required for preimplantation embryo
development. *Nat. Genet.* 55:484–495. doi:10.1038/s41588-023-01324-y.

Saksouk, N., E. Simboeck, and J. Déjardin. 2015. Constitutive heterochromatin formation
and transcription in mammals. *Epigenetics and Chromatin.* 8:1–17. doi:10.1186/1756-
8935-8-3.

Sankaran, S.M., A.W. Wilkinson, J.E. Elias, and O. Gozani. 2016. A PWWP Domain of
histone-lysine N-methyltransferase NSD2 binds to dimethylated lys-36 of histone H3
and regulates NSD2 function at chromatin. *J. Biol. Chem.* 291:8465–8474.
doi:10.1074/jbc.M116.720748.

Santoro, R., J. Li, and I. Grummt. 2002. The nucleolar remodeling complex NoRC mediates
heterochromatin formation and silencing of ribosomal gene transcription. *Nat. Genet.*
32:393–396. doi:10.1038/ng1010.

Schotta, G., M. Lachner, K. Sarma, A. Ebert, R. Sengupta, G. Reuter, D. Reinberg, and T.
Jenuwein. 2004. A silencing pathway to induce H3-K9 and H4-K20 trimethylation at
constitutive heterochromatin. *Genes Dev.* 18:1251–1262. doi:10.1101/gad.300704.

Shirane, K., F. Miura, T. Ito, and M.C. Lorincz. 2020. NSD1-deposited H3K36me2 directs de
novo methylation in the mouse male germline and counteracts Polycomb-associated
silencing. *Nat. Genet.* 52:1088–1098. doi:10.1038/s41588-020-0689-z.

Sinha, J., J.F. Nickels, A.R. Thurm, C.H. Ludwig, B.N. Archibald, M.M. Hinks, J. Wan, D.
Fang, and L. Bintu. 2023. The H3.3 K36M oncohistone disrupts the establishment of
epigenetic memory through loss of DNA methylation. bioRxiv. doi:
10.1101/2023.10.13.562147.

Smurova, K., and P. De Wulf. 2018. Centromere and Pericentromere Transcription: Roles
and Regulation … in Sickness and in Health. *Front. Genet.* 9:1–26.
doi:10.3389/fgene.2018.00674.

Sobecki, M., K. Mrouj, A. Camasses, N. Parisis, E. Nicolas, D. Llères, F. Gerbe, S. Prieto, L.
Krasinska, A. David, M. Eguren, M.C. Birling, S. Urbach, S. Hem, J. Déjardin, M.
Malumbres, P. Jay, V. Dulic, D.L.J. Lafontaine, R. Feil, and D. Fisher. 2016. The cell
proliferation antigen Ki-67 organises heterochromatin. *Elife*. 5:1–33.
doi:10.7554/eLife.13722.

Streubel, G., A. Watson, S.G. Jammula, A. Scelfo, D.J. Fitzpatrick, G. Oliviero, R. McCole,
E. Conway, E. Glancy, G.L. Negri, E. Dillon, K. Wynne, D. Pasini, N.J. Krogan, A.P.
Bracken, and G. Cagney. 2018. The H3K36me2 Methyltransferase Nsd1 Demarcates
PRC2-Mediated H3K27me2 and H3K27me3 Domains in Embryonic Stem Cells. *Mol.*
Cell. 70:371-379.e5. doi:10.1016/j.molcel.2018.02.027.

Strohner, R., A. Nemeth, P. Jansa, U. Hofmann-rohrer, R. Santoro, G. La, and I. Grummt.
2001. NoRC - a novel member of mammalian ISWI-containing chromatin remodeling
machines, Strohner et al, 2001. *EMBO J.* 20:4892–4900.

Suzuki, S., H. Kato, Y. Suzuki, Y. Chikashige, Y. Hiraoka, H. Kimura, K. Nagao, C. Obuse,
S. Takahata, and Y. Murakami. 2016. Histone H3K36 trimethylation is essential for
multiple silencing mechanisms in fission yeast. *Nucleic Acids Res.* 44:4147–4162.
doi:10.1093/nar/gkw008.

Tallant, C., E. Valentini, O. Fedorov, L. Overvoorde, F.M. Ferguson, P. Filippakopoulos, D.I.
Svergun, S. Knapp, and A. Ciulli. 2015. Molecular basis of histone tail recognition by
human TIP5 PHD finger and bromodomain of the chromatin remodeling complex
NoRC. *Structure*. 23:80–92. doi:10.1016/j.str.2014.10.017.

Tate, P., M. Lee, S. Tweedie, W.C. Skarnes, and W.A. Bickmore. 1998. Capturing novel
mouse genes encoding chromosomal and other nuclear proteins. *J. Cell Sci.* 111:2575–
2585. doi:10.1242/jcs.111.17.2575.

Tsumura, A., T. Hayakawa, Y. Kumaki, S.I. Takebayashi, M. Sakaue, C. Matsuoka, K.
Shimotohno, F. Ishikawa, E. Li, H.R. Ueda, J.I. Nakayama, and M. Okano. 2006.
Maintenance of self-renewal ability of mouse embryonic stem cells in the absence of
DNA methyltransferases Dnmt1, Dnmt3a and Dnmt3b. *Genes to Cells*. 11:805–814.
doi:10.1111/j.1365-2443.2006.00984.x.

Varshney, D., J. Vavrova-Anderson, A.J. Oler, V.H. Cowling, B.R. Cairns, and R.J. White.
2015. SINE transcription by RNA polymerase III is suppressed by histone methylation
but not by DNA methylation. *Nat. Commun.* 6. doi:10.1038/ncomms7569.

Vasudevan, D., E.Y.D. Chua, and C.A. Davey. 2010. Crystal Structures of Nucleosome Core
Particles Containing the "601" Strong Positioning Sequence. *J. Mol. Biol.* 403:1–10.
doi:10.1016/j.jmb.2010.08.039.

Vian, L., A. Pękowska, S.S.P. Rao, K.R. Kieffer-Kwon, S. Jung, L. Baranello, S.C. Huang, L.
El Khattabi, M. Dose, N. Pruett, A.L. Sanborn, A. Canela, Y. Maman, A. Oksanen, W.
Resch, X. Li, B. Lee, A.L. Kovalchuk, Z. Tang, S. Nelson, M. Di Pierro, R.R. Cheng, I.
Machol, B.G. St Hilaire, N.C. Durand, M.S. Shamim, E.K. Stamenova, J.N. Onuchic, Y.
Ruan, A. Nussenzweig, D. Levens, E.L. Aiden, and R. Casellas. 2018. The Energetics
and Physiological Impact of Cohesin Extrusion. *Cell*. 173:1165-1178.e20.
doi:10.1016/j.cell.2018.03.072.

Wagner, E.J., and P.B. Carpenter. 2012. Understanding the language of Lys36 methylation
at histone H3. *Nat. Rev. Mol. Cell Biol.* 13:115–126. doi:10.1038/nrm3274.

Weinberg, D.N., S. Papillon-Cavanagh, H. Chen, Y. Yue, X. Chen, K.N. Rajagopalan, C.
Horth, J.T. McGuire, X. Xu, H. Nikbakht, A.E. Lemiesz, D.M. Marchione, M.R. Marunde,
M.J. Meiners, M.A. Cheek, M.C. Keogh, E. Bareke, A. Djedid, A.S. Harutyunyan, N.
Jabado, B.A. Garcia, H. Li, C.D. Allis, J. Majewski, and C. Lu. 2019. The histone mark
H3K36me2 recruits DNMT3A and shapes the intergenic DNA methylation landscape.
Nature. 573:281–286. doi:10.1038/s41586-019-1534-3.

Woodcock, C.L., and R.P. Ghosh. 2010. Chromatin Higher-order Structure and Dynamics.
Cold Spring Harb Perspect Biol. 2.

Xu, Q., Y. Xiang, Q. Wang, L. Wang, J. Brind'Amour, A.B. Bogutz, Y. Zhang, B. Zhang, G.
Yu, W. Xia, Z. Du, C. Huang, J. Ma, H. Zheng, Y. Li, C. Liu, C.L. Walker, E. Jonasch, L.
Lefebvre, M. Wu, M.C. Lorincz, W. Li, L. Li, and W. Xie. 2019. SETD2 regulates the
maternal epigenome, genomic imprinting and embryonic development. *Nat. Genet.*
51:844–856. doi:10.1038/s41588-019-0398-7.

Yano, S., T. Ishiuchi, S. Abe, S.H. Namekawa, G. Huang, Y. Ogawa, and H. Sasaki. 2022.
Histone H3K36me2 and H3K36me3 form a chromatin platform essential for DNMT3A-
dependent DNA methylation in mouse oocytes. *Nat. Commun.* 13:4440.
https://doi.org/10.1038/s41467-022-32141-2

Yu, H., Z. Sun, T. Tan, H. Pan, J. Zhao, L. Zhang, J. Chen, A. Lei, Y. Zhu, L. Chen, Y. Xu, Y.
Liu, M. Chen, J. Sheng, Z. Xu, P. Qian, C. Li, S. Gao, G.Q. Daley, and J. Zhang. 2021.
rRNA biogenesis regulates mouse 2C-like state by 3D structure reorganization of peri-
nucleolar heterochromatin. *Nat. Commun.* 12:1–21. doi:10.1038/s41467-021-26576-2.

Yuan, S., R. Natesan, F.J. Sanchez-Rivera, J. Li, N. V. Bhanu, T. Yamazoe, J.H. Lin, A.J.
Merrell, Y. Sela, S.K. Thomas, Y. Jiang, J.B. Plesset, E.M. Miller, J. Shi, B.A. Garcia,
S.W. Lowe, I.A. Asangani, and B.Z. Stanger. 2020. Global regulation of the histone
mark H3K36ME2 underlies epithelial plasticity and metastatic progression. *Cancer*
Discov. 10:854–871. doi:10.1158/2159-8290.CD-19-1299.

Zhang, Y., H.H. Ng, H. Erdjument-Bromage, P. Tempst, A. Bird, and D. Reinberg. 1999.
Analysis of the NuRD subunits reveals a histone deacetylase core complex and a
connection with DNA methylation. *Genes Dev.* 13:1924–1935.
doi:10.1101/gad.13.15.1924.

Zhou, Y., R. Santoro, and I. Grummt. 2002. The chromatin remodeling complex NoRC
targets HDAC1 to the ribosomal gene promoter and represses RNA polymerase I
transcription. *EMBO J.* 21:4632–4640. doi:10.1093/emboj/cdf460.

1138

1139
1140

 Figure Legends

 Figure 1. H3K36me2 is enriched in pericentromeric heterochromatin in some mouse

 cells

  (A-C) Localization of H3K36me2 by Immunofluorescence. Pmi28 cells (A) and iMEFs (B)

 were fixed and stained with anti-H3K36me2 (green), anti-H3K9me3 (red), and Hoechst

 33342 (blue). (A and B) Single optical sections of confocal microscope images are shown.

 Insets show magnified views of indicated areas. Line plots indicate intensity profiles of

 H3K36me2, H3K9me3, and DNA along the yellow line in the merged images. (C)

 Quantification of the intensity ratio of H3K36me2 in PCH to that in the nucleoplasm in

 various mouse cell lines. Two box plots display data from two biological replicates with dots

 representing individual cells (round for the first experiment and triangular for the second).

 Center lines represent medians; box limits indicate the 25th and 75th percentiles; whiskers

 extend 1.5 times the interquartile range from these percentiles. The numbers of total cells

 analyzed (n) and p-value comparing ESCs (H3K36me2 in PCH) to MC12 cells (H3K36me2

 in euchromatin), calculated using the Mann-Whitney U test, are indicated. (D) Genome-wide

 correlation heatmap of H3K36me2 and H3K9me3 in mouse ESCs, B cells, and pancreatic

 cancer cells. Spearman's correlation coefficients are indicated inside cells. Scale bars,

 10 µm.

 Figure 2. NSD2 is the enzyme responsible for H3K36me2 at pericentromeric

 heterochromatin

 (A-D) Effect of Halo-NSD2 expression on H3K36me2 localization. iMEFs and Pmi28 cells

 were transfected with the Halo-NSD2 expression vector, and treated with JF646 HaloTag

 ligand and Hoechst33342. After live imaging, cells were fixed and stained with anti-

 H3K36me2. (A and B) Examples of single optical sections for Halo-NSD2 (red), H3K36me2

 (green), and Hoechst33342 (blue). Halo-NSD2 expressed (NSD2+) and unexpressed

 (NSD2-) iMEFs (A) and Pmi28 cells (B) are shown. (C and D) PCH-to-nucleus intensity

 ratios for Halo-NSD2 in live cells (C) and H3K36me2 by immunofluorescence (D). (E-G)

 Effect of NSD2 KO on H3K36me2 localization. (E) The protein levels of NSD2 evaluated by

 Western blotting in WT and NSD2 KO#1 and KO#2 iMEFs. Lamin B served as a loading

 control. Positions of size standards are indicated on the left. (F and G) Immunofluorescence

 and quantification. (F) WT, NSD2 KO#2, and KO#1 without and with Halo-NSD2 were

 stained with anti-H3K36me2 (green) and Hoechst33342 (red). Single optical section images

 are displayed. (G) PCH-to-nucleus intensity ratios for H3K36me2. See Fig. 1 legend for the

 details of box plots. Scale bars, 10 µm.

Figure 3. SMARCA5 modulates NSD2 localization and H3K36me2 levels in PCH

1178
(A) Candidates of NSD2-interacting proteins that mediate heterochromatic H3K36me2.
Among the NSD2-interacting factors previously identified (Huang et al., 2019), some are
known to be enriched in PCH. (B) Percentages of cells clearly showing PCH H3K36me2
when the indicated proteins are depleted by shRNA-mediated knockdown. The knockdown
efficiencies are indicated on the right as relative expression levels to the control (scramble)
shRNA, measured by RT-qPCR. (C-E) Localization of Halo-SMARCA5 and H3K36me2.
iMEFs and Pmi28 cells, transfected with the Halo-SMARCA5 expression vector, were
incubated in JF646 HaloTag ligand and Hoechst33342. After live imaging, cells were fixed
and stained with anti-H3K36me2. (C) Single optical sections for Halo-SMARCA5 (green) and
Hoechst33342 (red) in living cells. (D and E) PCH-to-nucleus intensity ratios of Halo-
SMARCA5 in live cells (D) and H3K36me2 by immunofluorescence (E). (F-I) Effects of
SMARCA5 KO on the localization of H3K36me2 and NSD2. (F) SMARCA5 protein levels
evaluated by Western blotting in WT and SMARCA5 KO#3 iMEFs. Lamin B served as a
loading control. The positions of size standards are indicated on the left. (G and H)
Immunofluorescence of H3K36me2 and quantification. (G) WT and SMARCA5 KO#3 without
and with Halo-SMARCA5 were stained with anti-H3K36me2 (green) and Hoechst33342
(red). Single optical section images are shown. (H) PCH-to-nucleus intensity ratios of
H3K36me2. (I) Intensity ratios of Halo-NSD2 in WT and SMARCA5 KO#3 cells without and
with SMARCA5-sfGFP or EGFP. See Fig. 1 legend for the details of box plots. Scale bars,
10 µm.

1199

Figure 4. BAZ1B regulates the PCH localization of SMARCA5 and NSD2

1200
(A) Candidates of ISWI complexes that bind to chromatin and recruit NSD2 to PCH. (B-D)
Localization of emGFP-BAZ1B and H3K36me2. iMEFs and Pmi28 cells transfected with
emGFP-BAZ1B expression vector were stained with Hoechst33342 for live cell imaging. For
immunofluorescence, cells were fixed and stained with anti-H3K36me2. (B) Examples of
single optical sections for emGFP-BAZ1B (red), H3K36me2 (green), and Hoechst33342
(blue). (C and D) PCH-to-nucleus intensity ratios of emGFP-BAZ1B in living cells (C) and
H3K36me2 by immunofluorescence (D). (E-I) Effects of BAZ1B KO on the localization of
H3K36me2, NSD2, and SMARCA5. (E) BAZ1B protein levels evaluated by Western blotting
in WT and BAZ1B KO#9 and KO#10 iMEFs. Lamin B served as a loading control. The
positions of size standards are indicated on the left. (F and G) Immunofluorescence of
H3K36me2 and quantification. (F) WT, BAZ1B KO#10, and KO#9 without and with emGFP-
BAZ1B were stained with anti-H3K36me2 (green) and Hoechst33342 (red). Single optical
section images are shown. (G) PCH-to-nucleus intensity ratios of H3K36me2. (H and I)
Intensity ratios of Halo-NSD2 (H) and Halo-SMARCA5 (I) in WT and BAZ1B KO#9 cells

without and with emGFP-BAZ1B or EGFP. See Fig. 1 legend for the details of box plots. (J)

Schematic drawing of NSD2 recruitment to PCH through the BAZ1B-SMARCA5 complex

(WICH). Scale bars, 10 µm.

1218

Figure 5. BAZ1B binds to AT-rich DNA through its AT-hook-like motif

(A) BAZ1B domain that targets PCH. Schematic drawing of the full length (FL), truncated

BAZ1B consisting of bromodomain (BrD) and three AT-hook-like motifs (BrD+AT1-3), and

those motifs mutant (BrD+AT1-3mut) is shown, with their representative live-cell images on

the right. Amino acid sequence alignments of BAZ1B AT-hook-like motifs in different species

and the substituted amino acids in AT1-3mut are shown. (B and C) DNA binding assay. (B)

Experimental design. EGFP-tagged BAZ1B BrD+AT1-3 and BrD+AT1-3mut were expressed

in HEK293T cells and purified using anti-GFP nanobody magnetic beads. After incubating

BAZ1B domain-bound anti-GFP beads with naked DNA or nucleosomes for 30 min in 20 mM

Tris-HCl [pH7.5] buffer, unbound DNA/nucleosomes were collected. After washing in 20 mM

Tris-HCl [pH7.5] buffer containing 50 mM NaCl, bound-DNA/nucleosomes were eluted using

20 mM Tris-HCl [pH7.5] containing 350 mM NaCl. (C) Detection of DNA in different fractions

by agarose-gel electrophoresis. The positions of size standards are indicated on the right.

Scale bar, 10 µm.

1233

Figure 6. Cooperative high expression of NSD2, SMARCA5, and BAZ1B promotes PCH

localization of H3K36me2

(A) Relative protein levels of NSD2, SMARCA5, and BAZ1B in mouse cell lines evaluated by

Western blotting. Lamin B was detected as a loading control. The positions of size standards

are indicated on the left. (B) PCH-to-nucleus intensity ratio of H3K36me2. Pmi28 cells were

transfected with the expression vectors of indicated proteins, fixed, and stained with anti-

H3K36me2, JF646 HaloTag ligand, and Hoechst33342. When expressing three proteins, an

equal amount of each expression vector was mixed. When expressing one or two proteins,

the same amount of each vector was used as was the case of triple expression, and the total

DNA amount was adjusted using the EGFP- or Halo-expression vector. See Fig. 1 legend for

the details of box plots. Note that p-values were calculated using the Mann-Whitney U test

and Hommel correction with the EGFP control.

1246

Figure 7. H3K36me2 is distributed in a ring-shaped PCH around the NPB in mouse

preimplantation embryos

Localization of H3K36me2 in mouse preimplantation embryos at various stages. Embryos

were fixed and stained with anti-H3K36me2 (red), anti-H3K9me3 (green), and Hoechst

33342 (blue). (A) Images of single optical sections. Low-power views of merged images are

shown on the left. Magnified views of indicated nuclei are shown on the right. PN,

pronucleus; TE, trophectoderm; and ICM, inner cell mass. (B) Pearson correlation

coefficients of DNA and H3K36me2 (top) or H3K9me3 (bottom). The number of nuclei (n)

and p-values, calculated using the Mann-Whitney U test and Hommel correction to the 4-cell

stage, are shown. See Fig. 1 legend for the details of box plots. Scale bars, 10 µm

1257

1258

1259

1260

Figure 1. H3K36me2 is enriched in pericentromeric heterochromatin in some mouse cells

Figure 2. NSD2 is the enzyme responsible for H3K36me2 at pericentromeric heterochromatin

Figure 3. SMARCA5 modulates NSD2 localization and H3K36me2 level in PCH

Figure 4. BAZ1B regulates the PCH localization of SMARCA5 and NSD2

A

BAZ1B domain structure

	WAC	DDT	WSD	PHD	BrD		PCH localization
FL							+
				AT1	AT2	AT3	
BrD+AT1-3							+
BrD+AT1-3mut							-

	AT1		AT2		AT3
X.tropicalis	R S S R G R N Y A	A R R G R P P K N	P G R G R K R K K
G.gallus	R S S R G R N Y T	S R Q G R H Q R K	H S R G R R R K K
M.musculus	R N S R G R N Y T	A R P G R P P G K	Q S R G R R Q K K
H.sapiens	R N S R G R N Y T	A R S G R R P G K	Q S R G R R Q K K
AT1-3mut	R N S D G D N Y T	A R S G D D P G K	Q S D G D D Q K K

B

anti-GFP magnetic beads

BrD+AT1-3 or BrD+AT1-3mut

N — C

EGFP

Naked DNA or Nucleosome

Incubate (30 min)

unbound DNA

Wash

N — C

bound DNA

C

	BrD+AT1-3				BrD+AT1-3mut			
	input	unbound DNA	wash	bound DNA	unbound DNA	wash	bound DNA	(bp)
Major Satellite Repeat								300
EF1 promoter								100
601 DNA for Nucleosome								100
Nucleosome								100

Figure 5. BAZ1B binds to AT-rich DNA through its AT-hook-like motif

Figure 6. Cooperative high expression of NSD2, SMARCA5, and BAZ1B promotes PCH localization of H3K36me2

Figure 7. H3K36me2 is distributed in a ring-shaped PCH around the NPB in mouse preimplantation embryos

Figure S1.

Figure S1. H3K36me2 is enriched in pericentromeric heterochromatin and LINE repeats in some mouse cell lines

(A-E) Immunofluorescence. Cells were fixed and stained with antibodies directed against histone modifications and Hoechst 33342. Single optical sections of confocal microscope images are shown. (A) H3K36me2 (green) and Hoechst33342 (red) in various mouse cell lines. (B) Pmi28 cells and iMEFs stained with another H3K36me2-specific antibody (green; rabbit monoclonal) with H3K9me3 (red) and Hoechst33342 (blue). (C-E) Localization of various modifications (blue), such as H3K27me3 (C), H3K9me2 (D), and H3K36me3 (E), respect to H3K36me2 (green) and H3K9me3 (red) in iMEFs. Insets show magnified views of the indicated PCH areas. (F and G) ChIP-seq signal enrichments of H3K36me2 and H3K9me3 on repeat elements in ESCs and B cells from the transcription start site (TSS) to the transcription end site (TES) (F) and around TSS (G). Scale bars, 10 µm.

Figure S2.

Figure S2. H3K36me2 PCH localization does not depend on H3K9me3 or DNA methylation and is induced by NSD2

(A and B) PCH localization of H3K36me2 in cells lacking H3K9me2/3 (A) or DNA methylation (B). 5KO iMEFs (lacking G9a, Glp, Setdb1, Suv39h1, and Suv39h2) (A) and TKO ESCs (lacking Dnmt1, Dnmt3a, and Dnmt3b) (B) were fixed and stained with anti-H3K36me2 (green) and Hoechst33342 (red). Single optical sections are shown on the left. Box plots display the intensity ratios of H3K36me2 in PCH to the nucleus. See Fig. 1 legend for the details of box plots. (C) Representative live-cell images of HaloTag-tagged H3K36 methylases and demethylases. NIH3T3 cells transfected with expression vectors for indicated Halo-tagged enzymes were stained with JF646 HaloTag ligand (green) and Hoechst33342 (red). Scale bars, 10 μm.

Figure S3. Effect of BAZ1A and BAZ2A on H3K36me2 localization
iMEFs and Pmi28 cells transfected with emGFP-BAZ1A (A) or emGFP-BAZ2A (B-D) expression vectors were stained with Hoechst 33342 for live-cell imaging (C). Cells were fixed and stained with anti-H3K36me2. (A and B) Examples of single optical sections of cells expressing emGFP-BAZ1A (A) and emGFP-BAZ2A (B) (red), H3K36me2 (green), and Hoechst33342 (blue). (C and D) PCH-to-nucleus intensity ratios of emGFP-BAZ2A in live cells (C) and H3K36me2 by immunofluorescence (D). See Fig. 1 legend for the details of box plots. Scale bars, 10 μm.

Figure S4. BAZ1B domain targeting PCH

(A) Schematic drawing of various truncated BAZ1B and their localizations. (B) Expression and purification of EGFP-tagged BAZ1B BrD+AT1-3 and BrD+AT1-3mut were evaluated by SDS-polyacrylamide gel electrophoresis and Coomassie Brilliant Blue staining. HEK293T cells transfected with the expression vectors were lysed and the supernatant (input) was incubated with GFP Trap anti-GFP nanobody magnetic beads at 4°C overnight. After removing the unbound fraction (Flowthrough), beads were washed, and aliquots were mixed with SDS-gel loading buffer (GFP-IP). Serial dilution series of bovine serum albumin (BSA) were loaded to estimate the amounts of recovered protein bound to beads. (C-E) Nucleotide sequences of DNA used for binding assay. (C) Major Satellite Repeats. (D) EF1 promoter. (E) 601 DNA. Sequences with more than three consecutive AT-bases are highlighted in red. Scale bar, 10 μm.

A

	Merge	H3K36me2	H3K9me3	DNA	Merge

Spleen

Lung

B

E13.5 male

#1 Brain (pons area)
#2 Cartilage primordium
#3 Stomach

Merge	H3K36me2	H3K9me3	DNA	Merge

#1

#2

#3

C

H3K36me2	DNA	Merge

Maternal

2cell

Maternal

2cell

Normalized Intensity

μm

— H3K36me2
— DNA

Figure S5.

Figure S5. H3K36me2 in mouse tissues, E13.5, and preimplantation embryos

(A and B) Immunofluorescence of tissue sections (A) and 13.5-day embryo sections (B) stained with antibodies against H3K36me2 (red), H3K9me3 (green), and Hoechst 33342 (blue). Single optical sections of confocal microscope images are shown. (A) Low-power views of merged images are shown on the left (Scale bars, 50 μm). Individual and merged images of the indicated area are shown on the right (Scale bars, 10 μm). (B) A tiled image covering the whole embryo is shown on the left (Scale bar, 500 μm). Single images for the brain (pons area; #1), cartilage primordium (#2), and stomach (#3) (Scale bars, 50 μm) and magnified views of the indicated areas are shown from the middle to the right (Scale bars, 10 μm). (C) Different localization of H3K36me2 in maternal PN in zygotes and the 2-cell nucleus in preimplantation embryos analyzed by line plots. Intensity profiles of H3K36me2 and DNA along the yellow lines are shown. Scale bar, 10 μm.

Table S1. Oligonucleotides

Primers	Sequence (5'-3')
Primers for generating Halo-SMARCA5	
InF_EcoR1_SM5_F	CTCAAGCTTCGAATTATGTCGTCCGCGGCCGAG
InF_EcoR1_SM5_R2	CGGTACCGTCGACTGCAGAATTAGTTTCAGCTTCTTTTTTCTTCC
Primers for generating truncated BAZ1B	
FL_F	ATCTCGAGCTCAAGCTTCGAAATGGCGCCGCTCCTGGGC
FL_R	GTCGACTGCAGAATTCTTCTTCTGTCTTCGTCCCC
ΔPHD,BrD_F	CATAAACACCTTCCTGGCC
ΔPHD,BrD_R	GTACCTCTTCTCTAGTCTCTCTTTAAG
PHD+BrD+AT1-3_F	ATCTCGAGCTCAAGCTTCGAACAGGACATTATTCACTCTATTC
PHD+BrD+AT1-3_R	GTCGACTGCAGAATTCTTCTTCTGTCTTCGTCCCC
PHD_F	ATCTCGAGCTCAAGCTTCGAACAGGACATTATTCACTCTATTC
PHD_R	GTCGACTGCAGAATTGCAAGCTGGGCACTGCCAC
BrD+AT1-3_F	ATCTCGAGCTCAAGCTTCGAACAGCCCGCTACTGCCAGG
BrD+AT1-3_R	GTCGACTGCAGAATTCTTCTTCTGTCTTCGTCCCC
BrD+AT1-2_F	ATCTCGAGCTCAAGCTTCGAACAGCCCGCTACTGCCAGG
BrD+AT1-2_R	GTCGACTGCAGAATTCAACAGAGCCACTAGACACTG
BrD+AT3_F	ATCTCGAGCTCAAGCTTCGAAGAGCTGCAGAAGTGTGAAGAG
BrD+AT3_R	GTCGACTGCAGAATTCTTCTTCTGTCTTCGTCCCC
AT1-2_F	ATCTCGAGCTCAAGCTTCGAACAGCCCGCTACTGCCAGG
AT1-2_R	GTCGACTGCAGAATTCAGGCTTTGCCTCCGGGAG
AT3_F	ATCTCGAGCTCAAGCTTCGAACATAAACACCTTCCTGGCCAC
AT3_R	GTCGACTGCAGAATTCTTCTTCTGTCTTCGTCCCC
gRNAs for generating knockout cells	
Nsd2_gRNA_1_s (PWWP1 domain)	CACCGAAAATACAACGTTGGTGATT
Nsd2_gRNA_1_as (PWWP1 domain)	AAACAATCACCAACGTTGTATTTTC
Nsd2_gRNA_2_s (SET domain)	CACCGTGGGGCCAGCATCAATTATG
Nsd2_gRNA_2_as (SET domain)	AAACCATAATTGATGCTGGCCCCAC
Smarca5_gRNA_1_s	CACCGCTTGCTCCGTTCTACGGTGT
Smarca5_gRNA_1_as	AAACACACCGTAGAACGGAGCAAGC
Baz1b_gRNA_1_s	CACCGTTCCCTGGCTTCTTGGCGGC
Baz1b_gRNA_1_as	AAACGCCGCCAAGAAGCCAGGGAAC
shRNAs for generating knockdown cells	
Smarca5_shRNA_1_s	CCGGGTCCGAGTGTTTCGCTTTATACTCGAGTATAAAGCGAAACACTCGGACTTTTTG
Smarca5_shRNA_1_as	AATTCAAAAAGTCCGAGTGTTTCGCTTTATACTCGAGTATAAAGCGAAACACTCGGAC
Smarca5_shRNA_2_s	CCGGCGTCGAATTAAAGCTGATGTTCTCGAGAACATCAGCTTTAATTCGACGTTTTTG
Smarca5_shRNA_2_as	AATTCAAAAACGTCGAATTAAAGCTGATGTTCTCGAGAACATCAGCTTTAATTCGACG

Parp1_shRNA_1_s	CCGGGCCCTTGGAAACATGTATGAACTCGAGTTCATACATGTTTCCAAGGGCTTTTTG
Parp1_shRNA_1_as	AATTCAAAAAGCCCTTGGAAACATGTATGAACTCGAGTTCATACATGTTTCCAAGGGC
Parp1_shRNA_2_s	CCGGTCGACGTCAACTACGAGAAACCTCGAGGTTTCTCGTAGTTGACGTCGATTTTTG
Parp1_shRNA_2_as	AATTCAAAAATCGACGTCAACTACGAGAAACCTCGAGGTTTCTCGTAGTTGACGTCGA
Orc2_shRNA_1_s	CCGGATCAATCCTGAACTCTATAACCTCGAGGTTATAGAGTTCAGGATTGATTTTTTG
Orc2_shRNA_1_as	AATTCAAAAAATCAATCCTGAACTCTATAACCTCGAGGTTATAGAGTTCAGGATTGAT
Orc2_shRNA_2_s	CCGGGCATCTATTGACCATCTCAATCTCGAGATTGAGATGGTCAATAGATGCTTTTTG
Orc2_shRNA_2_as	AATTCAAAAAGCATCTATTGACCATCTCAATCTCGAGATTGAGATGGTCAATAGATGC
Mki67_shRNA_1_s	CCGGGACAGTCTAGGGCAAGTATAACTCGAGTTATACTTGCCCTAGACTGTCTTTTTG
Mki67_shRNA_1_as	AATTCAAAAAGACAGTCTAGGGCAAGTATAACTCGAGTTATACTTGCCCTAGACTGTC
Mki67_shRNA_2_s	CCGGCTACTGTTCTGATCCATTAATCTCGAGATTAATGGATCAGAACAGTAGTTTTTG
Mki67_shRNA_2_as	AATTCAAAAACTACTGTTCTGATCCATTAATCTCGAGATTAATGGATCAGAACAGTAG
Mecp2_shRNA_1_s	CCGGCTGACAAAGCTTCCCGATTAACTCGAGTTAATCGGGAAGCTTTGTCAGTTTTTG
Mecp2_shRNA_1_as	AATTCAAAAACTGACAAAGCTTCCCGATTAACTCGAGTTAATCGGGAAGCTTTGTCAG
Mecp2_shRNA_2_s	CCGGGCTGGAAAGTATGATGTATATCTCGAGATATACATCATACTTTCCAGCTTTTTG
Mecp2_shRNA_2_as	AATTCAAAAAGCTGGAAAGTATGATGTATATCTCGAGATATACATCATACTTTCCAGC
Kdm2a_shRNA_1_s	CCGGGCTCCAAACCAACAAATATAACTCGAGTTATATTTGTTGGTTTGGAGCTTTTTG
Kdm2a_shRNA_1_as	AATTCAAAAAGCTCCAAACCAACAAATATAACTCGAGTTATATTTGTTGGTTTGGAGC
Kdm2a_shRNA_2_s	CCGGCGATGTCTTGTAGATAAGTTACTCGAGTAACTTATCTACAAGACATCGTTTTTG
Kdm2a_shRNA_2_as	AATTCAAAAACGATGTCTTGTAGATAAGTTACTCGAGTAACTTATCTACAAGACATCG
Hmgb1_shRNA_1_s	CCGGTGACAAGGCTCGTTATGAAAGCTCGAGCTTTCATAACGAGCCTTGTCATTTTTG
Hmgb1_shRNA_1_as	AATTCAAAAATGACAAGGCTCGTTATGAAAGCTCGAGCTTTCATAACGAGCCTTGTCA
Hmgb1_shRNA_2_s	CCGGGATGACAAGCAGCCCTATGAGCTCGAGCTCATAGGGCTGCTTGTCATCTTTTTG
Hmgb1_shRNA_2_as	AATTCAAAAAGATGACAAGCAGCCCTATGAGCTCGAGCTCATAGGGCTGCTTGTCATC
Nsd2_shRNA_1_s	CCGGCCTGGTGCTCATGATACTAAACTCGAGTTTAGTATCATGAGCACCAGGTTTTTG
Nsd2_shRNA_1_as	AATTCAAAAACCTGGTGCTCATGATACTAAACTCGAGTTTAGTATCATGAGCACCAGG
Nsd2_shRNA_2_s	CCGGCCCACTCCTTCACAATCATACCTCGAGGTATGATTGTGAAGGAGTGGGTTTTTG
Nsd2_shRNA_2_as	AATTCAAAAACCCACTCCTTCACAATCATACCTCGAGGTATGATTGTGAAGGAGTGGG
Scramble_s	CCGGCAACAAGATGAAGAGCACCAACTCGAGTTGGTGCTCTTCATCTTGTTGTTTTTG
Scramble_as	AATTCAAAAACAACAAGATGAAGAGCACCAACTCGAGTTGGTGCTCTTCATCTTGTTG

Primers for RT-qPCR

Smarca5_RT_F	TGCCTTGGGGATCAAAAGCT
Smarca5_RT_R	TGGAGGCAAACTCTTTTCAACA
Parp1_RT_F	GTCAAGAAGACGGCCGAGG
Parp1_RT_R	CCCAATGTCTTCTCCGCCTT
Orc2_RT_F	ACATCTGATCGAACGCTGCA
Orc2_RT_R	TCAGCAGAAAAGGAAGGCACA
Mki67_RT_F	ACCGAAAAGTGGACTTCAGTGA

Mki67_RT_R	TCTGCTGCTGCTTCTCCTTC
Mecp2_RT_F	TGTATATTTGATCAATCCCCAGGG
Mecp2_RT_R	TCATTAGGGTCCAAGGAGGTG
Kdm2a_F	GTGCATCTCACCCATTTTGAAC
Kdm2a_R	ATCGCTACTGGCAAGTTCTTCC
Hmgb1_RT_F	GGCTGACAAGGCTCGTTATG
Hmgb1_RT_R	TGCATTGGGGTCCTTGAACT
Nsd2_RT_F	GTGGCCTTGCATGGTTTCTG
Nsd2_RT_R	ATCCAAGCTCTTTCTGGGGC

Primers for cloning major satellites

MajSat_2_Fw	TATGGCGAGGAAAACTGAAAAAG
MajSat_2_Rv	TTTCACGTCCTAAAGTGTGTATTTC

Table S2. Antibodies

Primary antibody			Immunofluorescence				Western blotting
Antigen	Animal	Clone/Name (Source)	for cultured cells	for tissues	for 13.5-day embryo	for preimplantation embryos	
H3K36me2	Mouse	CMA332/2C3 (Rechtsteiner et al., 2010)	2 µg/mL (Alexa488 or Cy3-conjugated)	-	-	2.5 µg/mL (Cy3-conjugated)	-
H3K36me2	Rabbit	ab176921 (Abcam)	2 µg/mL	-	-	-	-
H3K36me3	Mouse	CMA333/13C9 (Rechtsteiner et al., 2010)	2 µg/mL (Cy5-conjugated)	-	-	-	-
H3K9me2	Mouse	CMA317/6D11 (Hayashi-Takanaka et al., 2011)	2 µg/mL (Cy5-conjugated)	-	-	-	-
H3K9me3	Mouse	CMA318/2F3 (Hayashi-Takanaka et al., 2011)	2 µg/mL (Cy3-conjugated)	10 µg/mL (Cy3-conjugated)	10 µg/mL (Cy5-conjugated)	5 µg/mL (Alexa488-conjugated)	-
H3K27me3	Mouse	CMA323/1E7 (Hayashi-Takanaka et al., 2011)	2 µg/mL (Cy5-conjugated)	-	-	-	-
NSD2	Rabbit	HPA015801 (Atlas Antibodies)	-	-	-	-	1:2000
SMARCA5	Rabbit	ab72499 (Abcam)	0.5 µg/mL	-	-	-	1:2000
BAZ1B	Mouse	sc-514287 (Santa Cruz)	0.5 µg/mL	-	-	-	1:500
LAMIN B1/2	Mouse	2D8 (Chen et al., JCB, 2018)	-	-	-	-	1:2000
Secondary antibody							
Mouse IgG	Donkey	715-005-150 (Jackson ImmunoResearch)	1 µg/mL (Alexa488-conjugated)	-	-	-	-
Rabbit IgG	Donkey	711-005-152 (Jackson ImmunoResearch)	1 µg/mL (Alexa488-conjugated)	-	-	-	-
Mouse IgG	Goat	Affinipure (Jackson ImmunoResearch)	-	-	-	-	1:10000 (HRP-conjugated)
Rabbit IgG	Goat	Affinipure (Jackson ImmunoResearch)	-	-	-	-	1:10000 (HRP-conjugated)

Fab

H3K36me2	Mouse	CMA332/2C3 (Rechtsteiner et al., 2010)	-	20 µg/mL (Alexa488-conjugated)	20 µg/mL (Alexa488-conjugated)