Evaluation of the neutralising antibody response in human and hamster sera against SARS-CoV-2

variants up to and including BA.2.86 using an authentic virus neutralisation assay

Naomi S. Coombes[1], Kevin R. Bewley[1], Yann Le Duff[2], Nassim Alami-Rahmouni[2], Kathryn A. Ryan[1],

Sarah Kempster[2], Deborah Ferguson[2], Elizabeth R. Davies[1], Thomas M. Weldon[1], Eleanor S. Cross[1],

Lauren Smith[1], Conner Norris[1], Karly Rai Rogers-Broadway[1], Kuiama Lewandowski[1], Samantha

Treagus[1], Steven T. Pullan[1], Bassam Hallis[1], Sue Charlton[1], Yper Hall[1], Simon G. P. Funnell[1,3,4]

1 UK Health Security Agency, Porton Down, Salisbury, Wiltshire, United Kingdom

2 Scientific Research and Innovation, Medicines and Healthcare products Regulatory Agency, South

Mimms, United Kingdom.

3 Quadram Institute Bioscience, Rosalind Franklin Road, Norwich Research Park, Norfolk, NR4 7UQ,

UK

4 World Health Organisation, Avenue Appia 20, 1211, Geneva, Switzerland

Abstract

New vaccines, therapeutics and immunity elicited by natural infection create evolutionary pressure

on SARS-CoV-2 to evolve and adapt to evade vaccine-induced and infection-elicited immunity.

Vaccine and therapeutics developers thus find themselves in an "arms race" with the virus. The

ongoing assessment of emerging SARS-CoV-2 variants remains essential as the global community

transitions from an emergency response to a long-term management plan. Here, we describe how

an authentic virus neutralisation assay using low passage clinical virus isolates has been employed to

monitor resistance of emerging virus variants to neutralising antibodies from humans and

experimentally infected hamsters. Sera and plasma from people who received three doses of a

vaccine as well as those who received a bivalent booster were assessed against SARS-CoV-2 variants,

up to and including BA.2.86. Contemporary or recent virus variants showed substantial resistance to

neutralisation by antibodies from those who had received three vaccines but were still effectively

neutralised by antibodies from individuals who had received a bivalent booster (ancestral/BA.1).

Convalescent sera from hamsters that had been experimentally infected with one of seven virus

variants (ancestral, BA.1, BA.4, BA.5.2.1, XBB.1.5, XBB.1.16, XBB.2.3) were also tested here. The most

contemporary variant, BA.2.86, was effectively neutralised by sera from hamsters infected with

XBB.1.5 and XBB.1.16 but it was not neutralised by sera from those infected with BA.5.2.1. These

data support the recommendations given by the WHO that a new vaccine was required and should

consist of an XBB sub-lineage antigen.

<u>Introduction</u>

Coronavirus 2019 disease (COVID-19) caused by SARS-CoV-2 continues to cause significant

disruption to healthcare systems and economies worldwide having caused over 700 million cases

and 6.7 million deaths since its emergence in December 2019 [1]. The WHO downgraded the COVID-

19 pandemic from a public health emergency of international concern (PHEIC) on 5 May 2023, this

marked a transition from emergency response to long-term management [2]. However, the virus

continues to evolve in response to changing population immunity leading to the regular

identification of new viral variants. Sequences of emerging variants are continuously monitored and

assigned as new sub-lineages by the Phylogenetic Assignment of Named Global Outbreak Lineages

(PANGOLIN) tool. At the time of writing there are 3,569 designated SARS-CoV-2 lineages with five of

these having been classified as variants of concern (VOC) by the WHO Technical Advisory Group on

Virus Evolution: Alpha (B.1.1.7), Beta (B.1.351), Delta (B.1.617.2), Gamma (P.1) and Omicron

(B.1.1.529) [3-5].

A global effort to design and manufacture COVID-19 vaccines began as soon as the viral genome was

published and the first COVID-19 vaccine was deployed in the UK on 8[th] December 2020 (Comirnaty)

[6]. This was followed by Vaxzevria on 30[th] December 2020 and Spikevax on 8[th] January 2021 [7, 8].

These vaccines were the most widely used worldwide for the initial course of three immunisations

and all were based on the ancestral (clade A) virus spike glycoprotein. In the first year of vaccines

being available it has been predicted that vaccination prevented 14.4 million deaths due to COVID-

19 [9]. The emergence of novel SARS-CoV-2 variants brought with it the concern that ancestral-

based vaccines would no longer provide sufficient protection to vulnerable people in at-risk groups.

Indeed, our data utilising the gold-standard authentic virus neutralisation assay indicated that the

latest variants were becoming increasingly resistant to ancestral-based immune responses [10-12].

Bivalent vaccines were developed in response to this concern and have been designed to not only

target the ancestral strain but also Omicron sub-lineage spike glycoproteins. These were approved

for use in the UK (BA.1/ancestral bivalent) for at-risk groups only in the Autumn of 2022, and the US

(BA.5/ancestral bivalent) for anyone aged 18 or over in August 2022 [13, 14]. Restricting vaccination

to only at-risk groups in the UK and low vaccine uptake in the US, where only 34% adults received a

fourth booster, means that most of the population who have received a vaccination are still

immunised with an ancestral spike-based vaccine [15]. However, since the relaxing of restrictions

worldwide, many people will also have received some degree of natural immunity due to infection

with one or more of the circulating variants. The most recent vaccines to be approved elicit

immunity against contemporary XBB recombinants, the latest to gain authorisation consists of an

XBB.1.5 component [16-18].

The recent emergence of the BA.2.86 variant has highlighted the value of sustained sequence

monitoring due to the sudden appearance of a virus with vastly different spike sequence compared

to other circulating variants or those represented in the latest vaccines. As part of a CEPI-funded

"Agility" project, we have been monitoring and performing risk assessments of SARS-CoV-2 variants

since November 2020. We previously presented our analysis of 20 different variants using early

pandemic human convalescent and triple-vaccinated plasma and serum panels as well as

international standards [12]. Here we describe our latest data generated using serum or plasma

collected from a panel of individuals who have received up to four doses of a COVID-19 vaccine using

an authentic virus neutralisation assay. These assays have been performed using well characterised,

low passage SARS-CoV-2 viruses derived from clinical isolates including the BA.2.86 variant. We also

present neutralisation data generated using the sera of experimentally infected hamsters which

were convalescent from either the ancestral virus (pre-D614G), BA.1, BA.4, BA.5.2.1, XBB.1.5,

XBB.1.16 or XBB.2.3 (mono-variant convalescent). We have also compared the antigenic maps of

these two data sets.

Methods

Human vaccinee sera and antibody standards

SARS-CoV-2 variants were tested against two panels of vaccinee samples collected from staff

volunteers. The first panel, consisting of sera from individuals who had received three doses of an

ancestral spike-based COVID-19 vaccine, was collected as part of the ESCAPE study as previously

described [12, 19]. The second panel was assembled from staff members at MHRA (n=8 plasma

samples) and UKHSA (n=14 sera samples) who had received a fourth vaccination consisting of

bivalent (BA.1/ancestral) vaccine. Samples were collected at a median of 78 days after the fourth

vaccination. When tested for the presence of anti-N antibodies, 19 of the 22 individuals had

evidence of convalescence for naturally acquired SARS-CoV-2 infection. Dates of previous infections

were provided by 14 individuals which identified an infection range of February 2021 to December

2022. Two reference materials were included in this study: a working reagent for anti-SARS-CoV-2

immunoglobulin (NIBSC 21/234), which is a pool of several pre-Alpha convalescent plasma, and the

International Standard for antibodies to SARS-CoV-2 VOC (NIBSC 21/338), which is a pool of 265

individuals who are both vaccinated and convalescent following Alpha, Beta or Delta infection [20,

21].

Virus bank preparation and quality control checks

Unless otherwise stated in Table 1, viruses were isolated from sequence-confirmed clinical swabs at

UKHSA. SARS-CoV-2 variants were isolated and propagated on Vero/hSLAM cells (ECACC 04091501)

before being subjected to quality control checks as previously described [12]. Briefly, titrations were

performed using a focus-forming assay with immunostaining for nucleocapsid to visualise foci [22].

Sighting for each variant into the Focus Reduction Neutralisation Test (FRNT) to calculate dilution at

which a median of 130 foci could be counted in non-neutralised control wells was performed. Whole

genome sequencing (SISPA-Illumina) was performed on final virus assay banks, as described

previously, to confirm presence of FCS S1/S2 boundary, check for mixed bases as well as confirm

variant identity [12, 23-25].

FRNT and statistics

Median neutralising antibody titres (ND_{50}) were measured with FRNT and calculated with a Probit

regression analysis using R v4.1.2 (R Core Team, Vienna, Austria) as described previously [11, 12, 22].

Log_{10}-transformed ND_{50} titres were analysed using a mixed-effects linear model in R, as described

previously [12, 26].

Antigenic cartography

Antigenic maps were generated using the Racmacs (https://acorg.github.io/Racmacs/) package in R

as described previously [27, 28]. The number of dimensions was set to 2, number of optimisations to

500 and minimum column basis to none. Neutralisation titres from human and hamster sera were

used to generate two separate maps.

Hamster sera

All *in vivo* experimental work was conducted under the authority of a UK Home Office approved

project licence that has been subject to local ethical review at either UK Health Security Agency

Porton Down or Medicines and Healthcare products Regulatory Agency, South Mimms by their

respective Animal Welfare and Ethical Review Body (AWERB) as required by the *Home Office*

Animals (Scientific Procedures) Act 1986.

Sera collected at day 27 post-infection from hamsters exposed to ancestral virus (n=6) and BA.1

(n=5) or day 28 post-infection with BA.4 (n=6) and BA.5.2.1 (n=6). These hamster sera were

generated as previously described [29, 30]. Sera was collected from hamsters infected with XBB1.16

and XBB.2.3 (n=6 per variant) with a target dose of $1x10^4$ FFU/200 µL via intranasal instillation

(method previously described [29, 30]). Blood samples were collected during necropsy at 28 days

post infection.

Sera from XBB.1.5 convalescent hamsters were collected as follows. Four hamsters (approximately

10 weeks old) were administered $1x10^5$ FFU/50µL intranasally of XBB.1.5. After infection, oral swabs

were taken to confirm infection by lateral flow and RT-qPCR. Hamsters were terminated at 28 days

post-infection and clotted whole blood was centrifuged (1,000RPM, 5 mins) to obtain serum.

Results

Through monitoring of the currently circulating SARS-CoV-2 variants, we identified and sourced

nasopharyngeal swabs from sequence-confirmed patients through our clinical networks and Pillar 1

testing (primarily positive tests conducted in hospital) in England. We then utilised an authentic virus

FRNT using low passage SARS-CoV-2 variants of clinical origin to perform risk assessments with

regards to cross-reactivity of humoral immunity. SISPA-Illumina sequencing was performed on stocks

prior to use in the FRNT to confirm the lineage and check for cell culture-induced mutations. We

note that one of the BA.2.86 variants (BA.2.86 – A) described here contains an unexpected amino

acid change within the furin cleavage site (FCS) (S:R682W). However, work published by others

indicates that changes in this position are unlikely to alter the neutralisation profile, though this

mutation makes this stock unsuitable for *in vivo* studies [31-34]. Neutralisation assays were

therefore performed with this virus whilst efforts continued to generate a bank of BA.2.86 with a

wild-type FCS (BA.2.86 – B). The variants used in this study, and their sources, are detailed in Table 1.

Table 1 – SARS-CoV-2 variants used in this study

WHO Designation	Pangolin Lineage	GISAID Clade/Lineage	Nextstrain Clade	Date Assigned	GISAID ID of Isolation Swab (Where Known) and/or Source
Ancestral virus	B	O	19A	Jan 2020	EPI_ISL_406844
Alpha	B.1.1.7	GRY	20I (V1)	Dec 2020	EPI_ISL_683466
Beta	B.1.351	GH/501Y.V2	20H (V2)	Dec 2020	EPI_ISL_770441
Delta	B.1.617.2	G/452R.V3	21A	May 2021	EPI_ISL_2742236
Zeta–FioCruz *	P.2	GR/484K.V2	20B/S.484K	Mar 2021	FioCruz, Brazil
Zeta–BEI *	P.2	GR/484K.V2	20B/S.484K	Mar 2021	BEI (NR-55439)
XE	XE	GRA	Recombinant	Mar 2022	EPI_ISL_11586931
XF	XF	GRA	Recombinant	Mar 2022	EPI_ISL_10458256
Omicron	BA.1	GRA	21K	Dec 2021	EPI_ISL_7400555
Omicron	BA.1.1	GRA	21K	Jan 2022	EPI_ISL_8165999
Omicron	BA.2	GRA	21L	Dec 2021	Not available
Omicron	BA.2.12.1	GRA	22C	Apr 2022	Gavin Screaton (University of Oxford)
Omicron	BA.2.75.3	GRA	22D	Jun 2022	EPI_ISL_13882158
Omicron	BA.2.75.2	GRA	22D	Aug 2022	Not available
Omicron	BA.4	GRA	22A	Apr 2022	EPI_ISL_13157810
Omicron	BA.4.6	GRA	22A	Jul 2022	Not available
Omicron	BA.5.2.1	GRA	22B	Apr 2022	EPI_ISL_12810908
Omicron	BQ.1.22	GRA	22E	Sep 2022	EPI_ISL_15581064
XBB.1.1	XBB.1.1	GRA	22F	Oct 2022	EPI_ISL_15682231
Omicron	CH.1.1	GRA	23C	Oct 2022	EPI-ISL-16615644
XBB.1.5	XBB.1.5	GRA	23A	Jan 2023	EPI-ISL-16587654
XBB.1.16	XBB.1.16	GRA	23B	Mar 2023	EPI_ISL_17666853
XBB.2.3	XBB.2.3	GRA	23E	Jan 2023	Not available
EG.5.1.1	EG.5.1.1	GRA	23F	Jun 2023	Not available
Omicron*	BA.2.86 (A)	GRA	21L	Aug 2023	EPI_ISL_18111770
Omicron*	BA.2.86 (B)	GRA	21L	Aug 2023	Not available

* For Zeta and BA.2.86, assays were conducted with stocks derived from viruses of different sources.

Our previously published data indicated that immunity derived from neutralising antibodies was

becoming less effective for those who had received three doses of a first-generation COVID-19

vaccine against the currently circulating variants [12]. Whilst a newer panel of sera was acquired, we

continued our assessment of triple vaccinated individuals who had received an original ancestral

based vaccine. This panel had been tested against ten SARS-CoV-2 variants (including ancestral) at

the point of publishing our previous paper. We have subsequently tested this panel against a further

six variants (Zeta – FC, Zeta – BEI, BA.4.6, BA.2.75.2, BQ.1.22 and XBB.1.1), as shown in Figure 1.

Our previous report demonstrated a significant difference in ND_{50} titres of our pre-Alpha

convalescent panel when tested against two different Zeta isolates. This was notable due to most

other published data for this variant demonstrating comparable neutralisation titres to those against

Delta. However, another group demonstrated a similarly large resistance to neutralisation of Zeta

with a panel of early convalescent sera but not a panel of vaccinee sera [35]. We therefore sought to

investigate whether the same pattern would be observed with our triple vaccinee panel (Figure 1).

Geometric mean (GM) neutralisation titres of this panel against Zeta – FC and Zeta – BEI were 199

and 671 corresponding to a 26.2-fold and 7.8-fold reduction relative to ancestral, respectively. The

difference between the titres measured against the two Zetas was significant (fold-change = 3.4; ME-

ANOVA; $p < 0.05$). Of the four remaining additional variants that were tested against this panel since

our previous paper, BA.4.6 and BA.2.75.2 resulted in GM titres above the lower limit of detection of

the assay (LLOD), thus permitting more accurate estimates of fold-reduction relative to ancestral

virus. The fold-reduction for BA.4.6 relative to ancestral was statistically significant at 105. When

compared to its close parent lineage, BA.4, there was also a significant fold-reduction of 2.1. Against

BA.2.75.2 the triple-vaccinated panel gave a GM titre of 46.1 which equated to a 113-fold reduction

relative to ancestral and 3.7-fold reduction relative to a closely related variant, BA.2.75.3. Two of the

most recent variants (BQ.1.22 and XBB.1.1) were substantially resistant to neutralisation to sera

from triple-vaccinated individuals with most titres falling below the LLOD. Sera which fall below the

LLOD are assigned the value of the lowest dilution used in the assay (1/20), thus fold-reduction

calculations do not provide accurate estimates. The lower limit of quantification (LLOQ) for the

ancestral virus assay (previously determined as 58) is also displayed [11]. Although the LLOQ was not

determined for subsequent variants, the precision of the assay is likely to be lower for titres falling

below this value. However, the estimated fold-reductions relative to ancestral virus for BQ.1.22 and

XBB.1.1 were estimated at 197 and 226-fold.

In addition, to assessing the sera from individuals of the panel, the neutralisation titres of the NIBSC

working reagents 21/234 and/or International Standard 21/338 were also assessed in every assay.

These are displayed in Figure 1 as green squares or blue circles, respectively. As described

previously, these are pools of pre-Alpha convalescent plasma (21/234) or individuals who were both

vaccinated and convalescent following an Alpha, Beta or Delta infection (21/338). These standards

are typically more potent at neutralising the virus compared to the panel. However, it was noted

that for BA.2.75.2, BA.4, BA.4.6 and BA.5.2.1, the 21/234 standard was yielding ND_{50} values

comparable to those from the lowest responders of our panel and was completely unable to

neutralise BQ.1.22. The newest 21/338 standard retains good reactivity and ability to neutralise all

variants assessed herein.

To continue to assess humoral immunity to currently circulating SARS-CoV-2 variants, a selection of

participants who had received a fourth (booster) vaccine were sought. A panel of 22 individuals was

assembled from donations given by eight MHRA staff members and 14 UKHSA staff members who

had received three doses of a first-generation COVID-19 vaccine followed by a booster with a

bivalent (BA.1/ancestral) vaccine. Samples were collected a median of 78 days after the most recent

vaccination. Among the 22 individuals, 19 had a known previous SARS-CoV-2 infection between

February 2021 and December 2022.

This four-dose panel was first assessed against the ancestral virus which resulted in the highest

neutralisation titres with a GM of 12,400 (Figure 2). Next, we assessed the titres of this panel against

one of the earlier Omicron sub lineages, BA.4. The GM neutralisation titres of the panel against BA.4

was 1,060 which equated to a 11.7-fold reduction relative to ancestral virus. To date this panel has

been tested against a further eight variants with the largest resistance to neutralisation seen for the

Omicron sub lineage, CH.1.1, with a GM titre for the panel of 62.9 and fold-reduction of 180.0.

Another Omicron sub lineage, BQ.1.22, also displayed notable resistance to neutralisation by the

four-dose panel which generated a GM titre of 92.9 corresponding to a 121.9-fold-reduction relative

to ancestral virus. The recombinants, XBB.1.1, XBB.1.5, XBB.1.16, XBB.2.3 and EG.5.1.1 all gave

comparable GM titres between 102 and 146. Similarly, the most recently emerged Omicron sub-

lineage at the time of writing, BA.2.86, was neutralised equally well by those who have had received

four doses of a COVID-19 vaccine. Here, we tested two BA.2.86 isolates with or without an FCS

substitution (S:R682W), BA.2.86 – A and BA.2.86 – B, respectively. The GM titres of the panel against

each virus was 99.3 and 143, equating to a 124.7 and 89.4-fold reduction relative to ancestral,

respectively. The difference between BA.2.86 – A and BA.2.86 – B was a 1.4-fold increase and was

not-significant (p=0.32). The numerical breakdown of these data and the model estimates of

geometric means are presented in Table S2.

Antigenic cartography is a powerful method for visualising the evolution of a pathogen by plotting

the neutralisation titres of different panels of sera against successive variants or strains. It was

originally developed as a means of monitoring changes in seasonal influenza virus and has since

been utilised by numerous groups to compare SARS-CoV-2 variants [28, 36, 37]. Here, we utilise this

technique to visualise the antigenic relationships of the SARS-CoV-2 variants tested against the

panels of human sera or plasma described in this and our previous publication (Figure 3). The data

plotted here include all titres from a panel of pre-Alpha convalescent plasma, triple-vaccinated

individuals, boosted (BA.1/ancestral bivalent) individuals and the three standards (20/136, 21/334,

21/338). Each square of the map represents one antigenic unit which equates to a two-fold change

in neutralisation titres. The ancestral virus is presented as a red circle and sits in the centre of the

serum panels. Pre-Omicron variants were coloured as dark green circles and are situated within an

inner circle around the serum panels. In the influenza field, vaccines are updated when circulating

strains are at least two units away from the vaccine strain, thus marking it as genetically different

[36]. Of the pre-Omicron variants, Beta and Zeta-FC are the furthest from ancestral virus with a gap

of over 4 antigenic units, though in perpendicular directions from ancestral. Substantial diversity can

be seen in post-Omicron variants which are grouped and coloured as Omicron-BA.1x (light green),

Omicron-BA.2.x (sky blue), Omicron-BA.4/5.x (yellow), late Omicron (lilac), XE/XF recombinants

(pink) and XBB recombinants (neon blue). However, some patterns can be seen, for example the

recombinants XF and XE cluster with the virus from which their spike was derived. The recombinant

XF is located with BA.1/BA.1.1 at approximately 5.5 units from ancestral and XE with BA.2/BA.2.12.1

at 2.5 units from ancestral. Also, the BA.4 and BA.5.2.1 cluster closely at approximately 5 units away

from ancestral and, while the BA.4.6 sub lineage is a similar distance from ancestral, it is also

approximately 4.5 units away from its parent, BA.4. The XBB family of variants cluster together at

approximately six units from ancestral. However, within this cluster there are three antigenic units

between the two most distant variants, XBB.1.16 and EG.5.1.1. The late Omicron variants (BQ.1.22,

CH.1.1 and BA.2.86) form another separate cluster approximately seven antigenic units from

ancestral. Interestingly, BA.2.75.2 and BA.2.75.3 appear on the opposite side of the map to BA.2

with BA.2.75.2 located closer to the late Omicrons than to BA.2.75.3.

One of the known limitations of generating antigenic maps with human neutralisation data is the

diversity of pre-existing immunity that is associated with multi-exposure sera (vaccine and naturally

acquired). We therefore utilised single-exposure convalescent sera collected during our *in vivo*

hamster infection studies to generate more antigenically defined maps. In Figure 4, we have plotted

neutralisation titres of sera from hamsters collected on day 27 (ancestral and BA.1) or day 28 (BA.4,

BA.5.2.1, XBB.1.5, XBB.1.16, XBB.2.3) after a single virus infection by the intranasal route. Each panel

of sera was tested against the variant used as the challenge virus. Fold-changes against subsequent

variants tested are presented as relative to the virus used for infection (Figure 4). Data for hamsters

infected with ancestral, BA.1, BA.4 and BA.5.2.1 are described previously but also presented here

[29, 30]. As our results from the human data indicated that the FCS substitution in the BA.2.86 – A

virus did not affect neutralisation titres, repeat assays with these volume-limited sera were not

performed with the second virus.

Briefly, sera taken from hamsters infected with ancestral virus gave significantly lower GM titres

with a 3.3- and 3.5-fold reduction for Alpha and Delta relative to ancestral, respectively. Sera taken

from BA.1 convalescent hamsters showed a similar ability to neutralise both BA.1 and ancestral but

were unable to neutralise BQ.1.22 and XBB.1.1 at the lowest dilution used here (1/30). For BA.4

convalescent hamster sera, BQ.1.22 and XBB.1.5 were the only variants which were significantly

more resistant to neutralisation by these sera, with the latter resulting in GM titres all below the

LLOD. In hamsters that were convalescent for BA.5.2.1, GM titres against ancestral virus relative to

BA.5.2.1 were not significantly different. Titres against all other variants tested against BA.5.2.1

convalescent sera were significantly lower relative to BA.5.2.1. Against BQ.1.22, XBB.1.1, XBB.1.5

and EG.5.1.1 titres were above the LLOD meaning fold changes could be determined accurately with

7.5-, 94.4-, 80.1-, 73.8-fold reductions, respectively. Notably, when tested against BA.2.86 – A, none

of the sera from BA.5.2.1 infected hamsters were able to neutralise the virus with all titres falling

below the LLOD. GM titres of convalescent sera from hamsters infected with XBB.1.5 tested against

EG.5.1.1 were significantly reduced relative to XBB.1.5 with a 1.8-fold reduction. When these sera

were tested against BA.2.86 there was no significant difference in GM neutralising titres.

Convalescent sera from hamsters experimentally infected with XBB.1.16 were tested against

ancestral virus, Beta, BA.1, BA.5.2.1, XBB.1.5, EG.5.1.1, XBB.2.3 and BA.2.86 – A. GM titres for this

panel against all variants were significantly reduced relative to XBB.1.16 with fold-reductions of 6.0,

77.5, 55.0, 17.1, 6.8, 10.4, 12.2 and 33.1, respectively. Finally, convalescent sera from hamsters

experimentally infected with XBB.2.3 were able to neutralise ancestral, XBB.1.16 and EG.5.1.1

equally well to XBB.2.3. Against all other variants (Beta, BA.1, BA.5.2.1, XBB.1.5, BA.2.86 – A), the

neutralisation titres for these sera were significantly lower with Beta displaying the most resistance

to neutralisation by this panel. The numerical breakdown of these data and the model estimates of

geometric means are presented in Tables S3-S9.

When these data were used to generate an antigenic map (Figure 5) the points corresponding to

individual sera were more widely distributed compared to the human map. As expected, each

antigenic variant clusters with sera from animals infected with homologous virus. Of the pre-

Omicron variants, Beta is again the most antigenically distant at approximately three units from

ancestral. BQ.1.22 is located in the same area of the map as Beta but another antigenic unit further

from ancestral. The most antigenically distinct variant with the hamster map was XBB.1.1 at 5.5 units

from ancestral and three units away from the nearest variant on the map, BA.1. The most

contemporary variant tested here, BA.2.86 – A, is approximately five units from ancestral but

clusters closely to XBB.1.5 with just over one antigenic units' distance.

Discussion

SARS-CoV-2 continues to cause worldwide disruption, despite the downgrading of the COVID-19

pandemic marking the end of the public health emergency. Since its emergence, SARS-CoV-2 has

been acquiring fitness-enhancing mutations in response to changing population immunity driven by

both infection and vaccination [38]. It is thought that additional pressures on the virus such as the

clinical use of antiviral drugs, long-term infections in immunocompromised individuals and spillback

events from zoonotic infections are also likely to contribute to the increased diversity in viral

sequences [39, 40]. Here, we present our updated assessment of emerging SARS-CoV-2 variants in

an authentic virus neutralisation assay against panels of sera or plasma from individuals who have

received three doses of a COVID-19 vaccine as well as those who received a bivalent (BA.1/ancestral)

booster. We also assessed neutralisation titres from experimentally infected hamsters that were

convalescent for seven SARS-CoV-2 variants (ancestral, BA.1, BA.4, BA.5.2.1, XBB.1.5, XBB.1.16,

XBB.2.3) and map both these and the human data antigenically.

In our previous publication we described a distinct difference in neutralisation titres against two

different isolates of the Zeta variant when tested against our pre-Alpha convalescent panel. Another

group from the University of Geneva showed a similar large reduction in neutralisation titres with an

early pandemic convalescent panel of sera against Zeta that was not seen against triple vaccinated

individuals [41]. We therefore tested our two Zeta isolates against our three-dose vaccinee panel,

however these viruses continued to show substantial resistance to neutralisation by this panel with a

significant difference between the two isolates. A sequence comparison with data kindly supplied by

the University of Geneva revealed identical spike sequences for all three viruses but with a three

amino acid deletion in the N gene of the virus discussed in the Bekliz paper (208-210). Zeta – BEI and

the University of Geneva virus also have three additional non-synonymous mutations in Orf1a

leading to amino acid changes: V1071A, P1810L and S3149F. With no differences in the main viral

proteins exposed to neutralising antibodies (spike, membrane, envelope), it appears that the

sequence is not responsible for the variation in titres seen here. Other potential explanations were

described in our previous paper and include the potential impact of secretion of free spike protein,

defective interfering particles and differences in spike glycosylation [12, 42-46].

Our data from individuals who had received three doses of an ancestral spike-based vaccine

indicated that antibodies against the two variants, BA.2.75.2 and BA.4.6, were less able to neutralise

these viruses with titres nearing the LLOD. BA.2.75.2 has three additional amino acid changes

relative to BA.2.75.3 (identical spike to BA.2.75) which are thought to increase immune evasion

(S:R346T, S:F486S, S:D1199N) [47, 48]. Similarly, BA.4.6 also contains S:R346T relative to its parent,

BA.4. Our neutralisation data supports this prediction with a higher immune escape of BA.2.75.2

compared to BA.4.6 and agrees with others who show similar trends [49, 50].

Further testing of this population demonstrated that the sera in this panel had no detectable

neutralising activity against the later variants, BQ.1.22 and XBB.1.1. The BQ.1.22 variant used in this

study has the same spike sequence as the BQ.1.1 variant, which was dominant in December 2022,

with three additional changes relative to its parent BA.5 (S:R346T, S:K444T, S:N460K). Similarly, the

recombinant XBB.1.1, circulating around the same time, had changes in several of the same

locations including S:R346T, S:L368I, S:N460K, S:F486S, S:R493Q and S:F490S. These two variants

were also resistant to neutralisation by panels of triple vaccinated volunteers in other studies as well

as in those with breakthrough infections with BA.1 and BA.2 [51, 52].

In the UK, as boosters containing bivalent antigens were deployed in the Autumn of 2022, we

continued our assessment of emerging variants with samples from individuals who had received a

BA.1/ancestral booster. The majority also had at least one breakthrough infection between February

2021 and December 2022 (Alpha to XBB.1.5). Results from this panel showed that neutralisation

titres for the two variants which were no longer neutralised by the three-dose panel (BQ.1.22 and

XBB.1.1) were effectively neutralised by this four-dose panel. Indeed, all the subsequent

recombinants and Omicron sub-lineages tested here showed similar neutralisation titres for all

variants between CH.1.1 to BA.2.86. This includes the two BA.2.86 isolates tested here, with and

without the presence of an amino acid change in the FCS (S:R682W), with no significant difference

seen for sera tested against the two viruses. These data are in alignment with others who have

demonstrated that changes in the FCS are not important in relation to virus neutralisation *in vitro*

but alter how the virus behaves *in vivo* [31-34]. All variants tested (CH.1.1, BQ.1.22, XBB.1.1, XBB.1.5,

XBB.1.16, XBB.2.3, EG.5.1.1, BA.2.86) displayed increased resistance to neutralisation to this four-

dose panel relative to both ancestral and BA.4, with similar GM titres for all. This aligns with results

presented by other groups using both pseudovirus (PSV) and authentic virus assays against panels of

vaccinated (three- and four-dose) with additional break-through infection in a globally diverse

population [53-60]. It is also of interest that the GM titres are comparable despite the genetic

diversity in the variants tested against our four-dose panels. This is likely due to the range of variants

to which these individuals have acquired natural immunity via break-through infections. We note

however that, due to vaccination campaigns targeting at-risk subgroups within the population, the

sera or plasma from individuals in our panel of boosted vaccinees are part of a limited subset of the

population considered to be at risk. Whilst this is informative in terms of the effects of vaccines on

immunity it may not be representative of levels of vaccine-induced immunity in the general

population.

Antigenic mapping of the human data shows co-localisation of the pre-Omicron variants in the

centre of the map with a larger antigenic spread following the emergence of Omicron and

recombinants. This is comparable to maps presented by others using human data, though direct

comparisons are challenging given the diversity in different panels of human sera used between

studies [28, 61-64]. For example, many will have had different vaccination schedules/ formulations

and exposure to unidentified viral variants in addition to human genetic diversity leading to a range

of humoral immune responses. However, while distances between variants in different maps may

vary, the general pattern is the same with pre-Omicron variants localised to the centre and clusters

of related variants moving out from the centre in a chronological manner. As others have shown, the

XBB family of variants cluster further away from ancestral than BA.4/5 and late Omicron variants

such as BQ.1.22, CH.1.1 and BA.2.86 are more distant again [60, 65].

One limitation of ours and others human antigenic maps is the clustering of sera in one location on

the map, corresponding to the ancestral spike that drove the pandemic and was used as the basis of

the first-generation vaccines. This effect is less prominent when using the neutralisation titres from

sera taken from hamsters 27- or 28-days post-infection with a single, known SARS-CoV-2 variant at a

standard infectious dose. Antigenic mapping of these data highlighted two antigenically distinct

viruses: BQ.1.22 and XBB.1.1. Although the most contemporary variant to date, BA.2.86, is one of

the furthest antigenically from ancestral, it clustered closely with XBB.1.5. This suggests that

antibodies raised against the XBB.1.5 spike antigen have good cross-reactivity against BA.2.86, thus

supporting the transition to XBB.1.5 based vaccines. Others have also shown XBB.1.1, BQ.1.1 and

BA.2.86 to be distinct in antigenic maps generated with hamster or mouse data [60, 65]. Using single

exposure sera from hamsters has the added benefits of being generated by a known priming

antigen, given at the same dose with the duration since infection being known and fixed. These are

therefore more predictive of antigenic relationships when mapped. Our hamster data also indicate

that a BA.5 spike alone does not generate a detectable neutralising antibody response to one of the

most recent variants, BA.2.86. Taken together, these data suggested a turning point in the virus'

evolution and signalled to vaccine developers that an updated vaccine was required. On 18 May

2023 the WHO Technical Advisory Group on COVID-19 Vaccine Composition (TAG-CO-VAC) published

their recommendations to develop vaccines to target contemporary XBB-descendent lineages [66].

Numbers of sequenced SARS-CoV-2 genomes are decreasing due to reduced global surveillance

activity since the end of the pandemic. As a result, identifying variants of concern in a timely manner

is increasingly challenging. This was highlighted by the apparent sudden emergence of BA.2.86 in

August 2023 despite some models retrospectively predicting it had been circulating since May 2023

[56, 67]. Continued monitoring of emerging variants and their assessment in programmes such as

Agility is essential to build on existing knowledge of immune evasion of contemporary and newly

emerging variants and to better advise policymakers and vaccine manufacturers on future strategies.

Acknowledgements

This work was funded by the Coalition for Epidemic Preparedness Innovations' (CEPI) Agility

Program. The authors gratefully acknowledge the support from the Porton Sequencing Laboratory

and the Biological Services Porton at the UK Health Security Agency, Porton Down, United Kingdom.

The authors thank our staff volunteers for providing serum and plasma samples so that we can

assemble our testing panels. We also acknowledge Samuel Wilks and Derek Smith for their help and

advice with using the Racmacs package. The authors are grateful to Professor Isabella Eckerle and Dr

Meriem Bekliz for sharing sequence data for their Zeta variant and useful discussions.

Ethics

The ESCAPE study was conducted in accordance with the Declaration of Helsinki and the Governance

Arrangements for Research Ethics Committees (GAfREC). It was approved by the PHE Research Ethics

and Governance Group (R&D REGG Ref NR0190, March 2020). Informed consent was obtained from

all subjects involved. Donations from MHRA staff members were collected with written informed

consent from donors, study was approved by the MHRA local ethics committee (HuMAC ref:

20/05/SK).

References

1.   Organization, W.H. *WHO Coronavirus (COVID-19) Dashboard*. 2023 [retrieved 2023 27th

407   September]; Available from: https://covid19.who.int/.

408   2.   Organization, W.H. *Statement on the fifteenth meeting of the IHR (2005) Emergency*

409   *Committee on the COVID-19 pandemic*. 2023 [retrieved 2023 27th September]; Available

410   from: https://www.who.int/news/item/05-05-2023-statement-on-the-fifteenth-meeting-of-

411   the-international-health-regulations-(2005)-emergency-committee-regarding-the-

412   coronavirus-disease-(covid-19)-pandemic.

413   3.   O'Toole, Á., E. Scher, A. Underwood, B. Jackson, V. Hill, J.T. McCrone, R. Colquhoun, C. Ruis,

414   K. Abu-Dahab, B. Taylor*, et al.*, *Assignment of epidemiological lineages in an emerging*

415   *pandemic using the pangolin tool.* Virus Evolution, 2021. **7**(2).

416   4.   Rambaut, A., E.C. Holmes, Á. O'Toole, V. Hill, J.T. Mccrone, C. Ruis, L. Du Plessis, and O.G.

417   Pybus, *A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic*

418   *epidemiology.* Nature Microbiology, 2020. **5**(11): p. 1403-1407.

419   5.   WHO. *Tracking SARS-CoV-2 variants*. 2023 [retrieved 2023 27th September]; Available from:

420   https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/.

421   6.   England, N. *Landmark moment as first NHS patient receives COVID-19 vaccination*. 2020

422   [retrieved 2023 10th October]; Available from:

423   https://www.england.nhs.uk/2020/12/landmark-moment-as-first-nhs-patient-receives-

424   covid-19-

425   vaccination/#:~:text=Landmark%20moment%20as%20first%20NHS%20patient%20receives%

426   20COVID%2D19%20vaccination,-

8%20December%202020&text=The%20biggest%20vaccine%20campaign%20in,jab%20follo

wing%20its%20clinical%20approval.

7.   Care, D.o.H.a.S. *Second COVID-19 vaccine authorised by medicines regulator*. 2020

[retrieved 2023 10th October]; Available from:

https://www.gov.uk/government/news/second-covid-19-vaccine-authorised-by-medicines-

regulator.

8.   Agency, M.a.H.p.R. *Moderna vaccine becomes third COVID-19 vaccine approved by UK*

regulator. 2021 [retrieved 2023 10th October]; Available from:

https://www.gov.uk/government/news/moderna-vaccine-becomes-third-covid-19-vaccine-

approved-by-uk-regulator.

9.   Watson, O.J., G. Barnsley, J. Toor, A.B. Hogan, P. Winskill, and A.C. Ghani, *Global impact of*

the first year of COVID-19 vaccination: a mathematical modelling study. Lancet Infect Dis,

2022. **22**(9): p. 1293-1302.

10.   TGHN. *The Global Health Network - Covax Epidemic Preparedness Innovations - Agility*. 2023

[retrieved 2023 1st October]; Available from: https://epi.tghn.org/covax-overview/enabling-

sciences/agility_epi/.

11.   Bewley, K.R., N.S. Coombes, L. Gagnon, L. McInroy, N. Baker, I. Shaik, J.R. St-Jean, N. St-

Amant, K.R. Buttigieg, H.E. Humphries*, et al.*, *Quantification of SARS-CoV-2 neutralizing*

antibody by wild-type plaque reduction neutralization, microneutralization and pseudotyped

virus neutralization assays. Nat Protoc, 2021. **16**(6): p. 3114-3140.

12.   Coombes, N.S., K.R. Bewley, Y. Le Duff, M. Hurley, L.J. Smith, T.M. Weldon, K. Osman, S.

Pullan, N. Berry, B. Hallis*, et al.*, *Assessment of the Biological Impact of SARS-CoV-2 Genetic*

Variation Using an Authentic Virus Neutralisation Assay with Convalescent Plasma, Vaccinee

Sera, and Standard Reagents. Viruses, 2023. **15**(3).

13.   MHRA. *First bivalent COVID-19 booster vaccine approved by UK medicines regulator*. 2022

[retrieved 2022 5th December]; Available from:

https://www.gov.uk/government/news/first-bivalent-covid-19-booster-vaccine-approved-

by-uk-medicines-regulator.

14.   FDA. *Coronavirus (COVID-19) Update: FDA Authorizes Moderna, Pfizer-BioNTech Bivalent*

COVID-19 Vaccines for Use as a Booster Dose. 2022 [retrieved 2022 5th December];

Available from: https://www.fda.gov/news-events/press-announcements/coronavirus-

covid-19-update-fda-authorizes-moderna-pfizer-biontech-bivalent-covid-19-vaccines-use.

15.   Prevention, C.f.D.C.a. *COVID-19 Vaccination Coverage and Vaccine Confidence Among*

Adults. 2023 [retrieved 2023 16th October]; Available from:

https://www.cdc.gov/vaccines/imz-

managers/coverage/covidvaxview/interactive/adults.html.

16.   Agency, M.a.H.p.R. *MHRA approves Moderna's adapted COVID-19 vaccine (Spikevax) that*

targets Omicron XBB.1.5. 2023 [retrieved 2023 10th October]; Available from:

https://www.gov.uk/government/news/mhra-approves-modernas-adapted-covid-19-

vaccine-spikevax-that-targets-omicron-xbb15.

17.   Agency, M.a.H.p.R. *MHRA approves Pfizer/BioNTech's adapted COVID-19 vaccine*

(Comirnaty) that targets Omicron XBB.1.5. 2023 [retrieved 2023 10th October]; Available

from: https://www.gov.uk/government/news/mhra-approves-pfizerbiontechs-adapted-

covid-19-vaccine-comirnaty-that-targets-omicron-xbb15.

18.   FDA. *FDA Takes Action on Updated mRNA COVID-19 Vaccines to Better Protect Against*

Currently Circulating Variants. 2023 [retrieved 2023 10th October]; Available from:

https://www.fda.gov/news-events/press-announcements/fda-takes-action-updated-mrna-

covid-19-vaccines-better-protect-against-currently-circulating.

19.   Harris, R.J., H.J. Whitaker, N.J. Andrews, F. Aiano, Z. Amin-Chowdhury, J. Flood, R. Borrow, E.

Linley, S. Ahmad, L. Stapley*, et al.*, *Serological surveillance of SARS-CoV-2: Six-month trends*

and antibody response in a cohort of public health workers. J Infect, 2021. **82**(5): p. 162-169.

20.   NIBSC. *Working Standard Working reagent for anti-SARS-CoV-2 immunoglobulin NIBSC code:*

479   *21/234*. 2022 [retrieved 2022 12th December]; Available from:

480   https://www.nibsc.org/documents/ifu/21-234.pdf.

481   21.   NIBSC. *WHO International Standard 1st International Standard for antibodies to SARS-CoV-2*

482   *variants of concern NIBSC code: 21/338*. 2022 [retrieved 2022 12th December]; Available

483   from: https://www.nibsc.org/documents/ifu/21-338.pdf.

484   22.   Ryan, K.A., R.J. Watson, K.R. Bewley, C. Burton, O. Carnell, B.E. Cavell, A. Challis, N.S.

485   Coombes, K. Emery, R. Fell*, et al.*, *Convalescence from prototype SARS-CoV-2 protects Syrian*

486   *hamsters from disease caused by the Omicron variant*. 2021, Cold Spring Harbor Laboratory.

487   23.   Lewandowski, K., Y. Xu, S.T. Pullan, S.F. Lumley, D. Foster, N. Sanderson, A. Vaughan, M.

488   Morgan, N. Bright, J. Kavanagh*, et al.*, *Metagenomic Nanopore Sequencing of Influenza Virus*

489   *Direct from Clinical Respiratory Samples.* J Clin Microbiol, 2019. **58**(1).

490   24.   Li, H., *Aligning sequence reads, clone sequences and assemblycontigs with BWA-MEM.*

491   arXiv:1303.3997v2 [q-bio.GN], 2013.

492   25.   Danecek, P., J.K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M.O. Pollard, A. Whitwham, T.

493   Keane, S.A. McCarthy, R.M. Davies*, et al.*, *Twelve years of SAMtools and BCFtools.*

494   Gigascience, 2021. **10**(2).

495   26.   R. *R Core Team: A language and environment for statistical computing. R Foundation for*

496   *Statistical Computing*. 2014 [retrieved 2023 24th February]; Available from: http://www.R-

497   project.org/.

498   27.   Wilks, S. *Racmacs: Antigenic Cartography Macros*. 2023 [retrieved 2023 1st October];

499   Available from: https://acorg.github.io/Racmacs/, https://github.com/acorg/Racmacs/.

500   28.   Wilks, S.H., B. Muhlemann, X. Shen, S. Tureli, E.B. LeGresley, A. Netzl, M.A. Caniza, J.N.

501   Chacaltana-Huarcaya, V.M. Corman, X. Daniell*, et al.*, *Mapping SARS-CoV-2 antigenic*

502   *relationships and serological responses.* bioRxiv, 2023.

29.   Davies, E.R., K.A. Ryan, K.R. Bewley, N.S. Coombes, F.J. Salguero, O.T. Carnell, S.

Biddlecombe, M. Charlton, A. Challis, E.S. Cross*, et al.*, *The Omicron Sub-Variant BA.4*

Displays a Remarkable Lack of Clinical Signs in a Golden Syrian Hamster Model of SARS-CoV-2

Infection. Viruses, 2023. **15**(5).

30.   Ryan, K.A., K.R. Bewley, R.J. Watson, C. Burton, O. Carnell, B.E. Cavell, A. Challis, N.S.

Coombes, E.R. Davies, J. Edun-Huges*, et al.*, *Syrian hamster convalescence from prototype*

SARS-CoV-2 confers measurable protection against the attenuated disease caused by the

Omicron variant. PLoS Pathog, 2023. **19**(4): p. e1011293.

31.   Chan, Y.A. and S.H. Zhan, *The Emergence of the Spike Furin Cleavage Site in SARS-CoV-2.* Mol

Biol Evol, 2022. **39**(1).

32.   Valleriani, F., L. Jurisic, C. Di Pancrazio, R. Irelli, E. Ciarrocchi, M. Martino, A. Cocco, E. Di

Felice, M.L. Colaianni, N. Decaro*, et al.*, *A Deletion Encompassing the Furin Cleavage Site in*

the Spike Encoding Gene Does Not Alter SARS-CoV-2 Replication in Lung Tissues of Mink and

Neutralization by Convalescent Human Serum Samples. Pathogens, 2022. **11**(10).

33.   Klimstra, W.B., N.L. Tilston-Lunel, S. Nambulli, J. Boslett, C.M. McMillen, T. Gilliland, M.D.

Dunn, C. Sun, S.E. Wheeler, A. Wells*, et al.*, *SARS-CoV-2 growth, furin-cleavage-site*

adaptation and neutralization using serum from acutely infected hospitalized COVID-19

patients. J Gen Virol, 2020. **101**(11): p. 1156-1169.

34.   Peacock, T.P., D.H. Goldhill, J. Zhou, L. Baillon, R. Frise, O.C. Swann, R. Kugathasan, R. Penn,

J.C. Brown, R.Y. Sanchez-David*, et al.*, *The furin cleavage site in the SARS-CoV-2 spike protein*

is required for transmission in ferrets. Nature Microbiology, 2021. **6**(7): p. 899-909.

35.   Bekliz, M., K. Adea, P. Vetter, C.S. Eberhardt, K. Hosszu-Fellous, D.-L. Vu, O. Puhach, M.

Essaidi-Laziosi, S. Waldvogel-Abramowski, C. Stephan*, et al.*, *Neutralization of ancestral*

SARS-CoV-2 and variants Alpha, Beta, Gamma, Delta, Zeta and Omicron by mRNA

vaccination and infection-derived immunity through homologous and heterologous variants.

2021, Cold Spring Harbor Laboratory.

36.   Smith, D.J., A.S. Lapedes, J.C. de Jong, T.M. Bestebroer, G.F. Rimmelzwaan, A.D.M.E.

Osterhaus, and R.A.M. Fouchier, *Mapping the Antigenic and Genetic Evolution of Influenza*

Virus. Science, 2004. **305**(5682): p. 371-376.

37.   Fonville, J.M., S.H. Wilks, S.L. James, A. Fox, M. Ventresca, M. Aban, L. Xue, T.C. Jones, Y.

Wong, A. Mosterin*, et al.*, *Antibody landscapes after influenza virus infection or vaccination.*

Science, 2014. **346**(6212): p. 996-1000.

38.   Basile, K., J.E. Agius, W. Fong, K. McPhie, M. Fennel, D. Ko, L. Heuston, C. Lam, A. Arnott,

S.C.-A. Chen*, et al.*, *Genomic evolution of SARS-CoV-2 variants of concern under in*

vitro neutralising selection pressure following two doses of the Pfizer-BioNTech

BNT162b2 COVID-19 vaccine. bioRxiv, 2023: p. 2023.09.24.558921.

39.   Sanderson, T., R. Hisner, I.a. Donovan-Banfield, H. Hartman, A. Løchen, T.P. Peacock, and C.

Ruis, *A molnupiravir-associated mutational signature in global SARS-CoV-2 genomes.* Nature,

2023.

40.   Markov, P.V., M. Ghafari, M. Beer, K. Lythgoe, P. Simmonds, N.I. Stilianakis, and A.

Katzourakis, *The evolution of SARS-CoV-2.* Nature Reviews Microbiology, 2023. **21**(6): p. 361-

379.

41.   Bekliz, M., K. Adea, P. Vetter, C.S. Eberhardt, K. Hosszu-Fellous, D.L. Vu, O. Puhach, M.

Essaidi-Laziosi, S. Waldvogel-Abramowski, C. Stephan*, et al.*, *Neutralization capacity of*

antibodies elicited through homologous or heterologous infection or vaccination against

SARS-CoV-2 VOCs. Nat Commun, 2022. **13**(1): p. 3840.

42.   Brun, J., S. Vasiljevic, B. Gangadharan, M. Hensen, A. V. Chandran, M.L. Hill, J.L. Kiappes, R.A.

Dwek, D.S. Alonzi, W.B. Struwe*, et al.*, *Assessing Antigen Structural Integrity through*

Glycosylation Analysis of the SARS-CoV-2 Viral Spike. ACS Central Science, 2021. **7**(4): p. 586-

593.

43.   Kwilas, S., R.M. Liesman, L. Zhang, E. Walsh, R.J. Pickles, and M.E. Peeples, *Respiratory*

Syncytial Virus Grown in Vero Cells Contains a Truncated Attachment Protein That Alters Its

Infectivity and Dependence on Glycosaminoglycans. Journal of Virology, 2009. **83**(20): p.

10710-10718.

44.   Routhu, N.K., S.D. Lehoux, E.A. Rouse, M.R.M. Bidokhti, L.B. Giron, A. Anzurez, S.P. Reid, M.

Abdel-Mohsen, R.D. Cummings, and S.N. Byrareddy, *Glycosylation of Zika Virus is Important*

in Host–Virus Interaction and Pathogenic Potential. International Journal of Molecular

Sciences, 2019. **20**(20): p. 5206.

45.   Letarov, A.V., V.V. Babenko, and E.E. Kulikov, *Free SARS-CoV-2 Spike Protein S1 Particles May*

Play a Role in the Pathogenesis of COVID-19 Infection. Biochemistry (Mosc), 2021. **86**(3): p.

257-261.

46.   Girgis, S., Z. Xu, S. Oikonomopoulos, A.D. Fedorova, E.P. Tchesnokov, C.J. Gordon, T.M.

Schmeing, M. Gotte, N. Sonenberg, P.V. Baranov*, et al.*, *Evolution of naturally arising SARS-*

CoV-2 defective interfering particles. Commun Biol, 2022. **5**(1): p. 1140.

47.   Miller, N.L., T. Clark, R. Raman, and R. Sasisekharan, *Insights on the mutational landscape of*

the SARS-CoV-2 Omicron variant receptor-binding domain. Cell Rep Med, 2022. **3**(2): p.

100527.

48.   Greaney, A.J., T.N. Starr, and J.D. Bloom, *An antibody-escape estimator for mutations to the*

SARS-CoV-2 receptor-binding domain. Virus Evolution, 2022. **8**(1).

49.   Planas, D., T. Bruel, I. Staropoli, F. Guivel-Benhassine, F. Porrot, P. Maes, L. Grzelak, M. Prot,

S. Mougari, C. Planchais*, et al.*, *Resistance of Omicron subvariants BA.2.75.2, BA.4.6, and*

BQ.1.1 to neutralizing antibodies. Nat Commun, 2023. **14**(1): p. 824.

50.   Dijokaite-Guraliuc, A., R. Das, R. Nutalai, D. Zhou, A.J. Mentzer, C. Liu, P. Supasa, S.J.

Dunachie, T. Lambe, E.E. Fry*, et al.*, *Antigenic characterization of SARS-CoV-2 Omicron*

subvariant BA.4.6. Cell Discov, 2022. **8**(1): p. 127.

51.   Wang, X., S. Jiang, S. Jiang, X. Li, J. Ai, K. Lin, S. Lv, S. Zhang, M. Li, J. Li*, et al.*, *Neutralization of*

SARS-CoV-2 BQ.1.1, CH.1.1, and XBB.1.5 by breakthrough infection sera from previous and

recent waves in China. Cell Discov, 2023. **9**(1): p. 64.

52.   Jiang, X.L., K.L. Zhu, X.J. Wang, G.L. Wang, Y.K. Li, X.J. He, W.K. Sun, P.X. Huang, J.Z. Zhang,

H.X. Gao*, et al.*, *Omicron BQ.1 and BQ.1.1 escape neutralisation by omicron subvariant*

breakthrough infection. Lancet Infect Dis, 2023. **23**(1): p. 28-30.

53.   Wang, X., S. Jiang, W. Ma, X. Li, K. Wei, F. Xie, C. Zhao, X. Zhao, C. Li, R. Qiao*, et al.*, *Enhanced*

neutralization of SARS-CoV-2 XBB sub-lineages and BA.2.86 by a tetravalent COVID-19

vaccine booster. bioRxiv, 2023: p. 2023.09.14.557682.

54.   Qu, P., K. Xu, J.N. Faraone, N. Goodarzi, Y.-M. Zheng, C. Carlin, J.S. Bednash, J.C. Horowitz,

R.K. Mallampalli, L.J. Saif*, et al.*, *Immune Evasion, Infectivity, and Fusogenicity of SARS-CoV-2*

Omicron BA.2.86 and FLip Variants. bioRxiv, 2023: p. 2023.09.11.557206.

55.   Hu, Y., J. Zou, C. Kurhade, X. Deng, H.C. Chang, D.K. Kim, P.-Y. Shi, P. Ren, and X. Xie, *Less*

neutralization evasion of SARS-CoV-2 BA.2.86 than XBB sublineages and CH.1.1. bioRxiv,

2023: p. 2023.09.10.557047.

56.   Khan, K., G. Lustig, K. Reedoy, Z. Jule, C. Römer, F. Karim, Y. Ganga, M. Bernstein, Z. Baig, B.

Mahlangu*, et al.*, *Evolution and neutralization escape of the SARS-CoV-2 BA.2.86 subvariant.*

medRxiv, 2023: p. 2023.09.08.23295250.

57.   Uriu, K., J. Ito, Y. Kosugi, Y.L. Tanaka, Y. Mugita, Z. Guo, J. Alfredo A Hinay, O. Putri, Y. Kim, R.

Shimizu*, et al.*, *Transmissibility, infectivity, and immune resistance of the SARS-CoV-2 BA.2.86*

variant. bioRxiv, 2023: p. 2023.09.07.556636.

58.   Lasrado, N., A.-r.Y. Collier, N.P. Hachmann, J. Miller, M. Rowe, E.D. Schonberg, S.L.

Rodrigues, A. LaPiana, R.C. Patio, T. Anand*, et al.*, *Neutralization Escape by SARS-CoV-2*

Omicron Subvariant BA.2.86. bioRxiv, 2023: p. 2023.09.04.556272.

59.   Sheward, D.J., Y. Yang, M. Westerberg, S. Öling, S. Muschiol, K. Sato, T.P. Peacock, G.B.K.

Hedestam, J. Albert, and B. Murrell, *Sensitivity of BA.2.86 to prevailing neutralising antibody*

responses. bioRxiv, 2023: p. 2023.09.02.556033.

60.   Yang, S., Y. Yu, F. Jian, W. Song, A. Yisimayi, X. Chen, Y. Xu, P. Wang, J. Wang, L. Yu*, et al.*,

Antigenicity and infectivity characterisation of SARS-CoV-2 BA.2.86. Lancet Infect Dis, 2023.

61.   Wang, W., S. Lusvarghi, R. Subramanian, N.J. Epsi, R. Wang, E. Goguet, A.C. Fries, F.

608   Echegaray, R. Vassell, S.A. Coggins*, et al.*, *Antigenic cartography of well-characterized human*

609   *sera shows SARS-CoV-2 neutralization differences based on infection and vaccination history.*

610   Cell Host Microbe, 2022. **30**(12): p. 1745-1758 e7.

611   62.   Straten, K.v.d., D. Guerra, M.J.v. Gils, I. Bontjer, T.G. Caniels, H.D.G.v. Willigen, E. Wynberg,

612   M. Poniman, J.A. Burger, J.H. Bouhuijs*, et al.*, *Mapping the antigenic diversification of SARS-*

613   *CoV-2.* medRxiv, 2022: p. 2022.01.03.21268582.

614   63.   van der Straten, K., D. Guerra, M.J. van Gils, I. Bontjer, T.G. Caniels, H.D.G. van Willigen, E.

615   Wynberg, M. Poniman, J.A. Burger, J.H. Bouhuijs*, et al.*, *Antigenic cartography using sera*

616   *from sequence-confirmed SARS-CoV-2 variants of concern infections reveals antigenic*

617   *divergence of Omicron.* Immunity, 2022. **55**(9): p. 1725-1731 e4.

618   64.   Astakhova, E.A., A.A. Morozov, M.G. Byazrova, M.M. Sukhova, A.A. Mikhailov, A.R.

619   Minnegalieva, A.A. Gorchakov, and A.V. Filatov, *Antigenic Cartography Indicates That the*

620   *Omicron BA.1 and BA.4/BA.5 Variants Remain Antigenically Distant to Ancestral SARS-CoV-2*

621   *after Sputnik V Vaccination Followed by Homologous (Sputnik V) or Heterologous*

622   *(Comirnaty) Revaccination.* International Journal of Molecular Sciences, 2023. **24**(13): p.

623   10493.

624   65.   Mykytyn, A.Z., M.E. Rosu, A. Kok, M. Rissmann, G. van Amerongen, C. Geurtsvankessel, R.D.

625   de Vries, B.B.O. Munnink, D.J. Smith, M.P.G. Koopmans*, et al.*, *Antigenic mapping of*

626   *emerging SARS-CoV-2 omicron variants BM.1.1.1, BQ.1.1, and XBB.1.* Lancet Microbe, 2023.

627   **4**(5): p. e294-e295.

628   66.   Organization, W.H. *Statement on the antigen composition of COVID-19 vaccines*. 2023

629   [retrieved 2023 27th September]; Available from: https://www.who.int/news/item/18-05-

630   2023-statement-on-the-antigen-composition-of-covid-19-vaccines.

631   67.   Rothstein, A.P., X. Qiu, K. Robison, S. Collins, G. Muir, B. Lu, A.M. Plocik, B.B. Simen, and C.W.

632   Philipson, *Bayesian phylogenetics on globally emerging SARS-CoV-2 variant BA.2.86 suggest*

633   *global distribution and rapid evolution.* bioRxiv, 2023: p. 2023.09.08.556912.

Figure 1. Neutralising antibody titres of a panel of triple vaccinated human participant sera determined by Focus Reduction Neutralisation Test (FRNT) against authentic SARS-CoV-2 Variants.

A panel of ten sera from triple vaccinated human volunteers were assessed by FRNT with 16 authentic SARS-CoV-2 variants. Data are presented as median and interquartile ranges of titre ND_{50}, titres of panel members falling outside of these ranges are plotted as open small circles. Fold-changes relative to ancestral virus were calculated by regression analysis using a mixed-effect model and are presented where the difference is statistically significant ($p<0.05$, two-way ANOVA with Tukey's *post hoc* test). The Lower Limit of Detection (LLOD) of 1/20 is indicated by the blue dotted line. The Lower Limit of Quantification (LLOQ) of 1/58 as determined for the ancestral assay, is indicated by the grey dotted line. Neutralisation titres of the standards; NIBSC WR 21/234 and 21/338, are presented as a green box and blue circle, respectively.

Figure 2. Neutralising antibody titres of a panel of sera/plasma from human volunteers who received three doses of an ancestral based vaccine as well as a BA.1 bivalent booster as determined by Focus Reduction Neutralisation Test (FRNT) against authentic SARS-CoV-2 Variants.

Sera/plasma from participants who received four doses of a vaccine were assessed by FRNT with 11 authentic SARS-CoV-2 variants. The panel consists of sera (n=14) and plasma (n=8) samples from human participants who received three doses of an ancestral based vaccine as well as a BA.1 bivalent booster. Of these, 19 out of 22 had at least one confirmed breakthrough infection with SARS-CoV-2. Data are presented as median and interquartile ranges of titre (ND$_{50}$). Fold-changes relative to ancestral virus were calculated by regression analysis using a linear mixed effect model and are presented where the difference is statistically significant (p<0.05, two-way ANOVA with Tukey's *post hoc* test). The Lower Limit of Detection (LLOD) of 1/20 is indicated by the blue dotted line. The Lower Limit of Quantification (LLOQ) of 1/58 as determined for the ancestral assay, is indicated by the grey dotted line. Neutralisation titres of the NIBSC WR 21/338 is presented as a blue circle.

Figure 3. Antigenic map of SARS-CoV-2 variants assessed against human pre-Alpha convalescent, vaccinee sera or plasma and antibody standards

Median neutralisation titres (ND_{50}) for human sera/plasma were generated against 34 authentic SARS-CoV-2 variants and plotted as antigenic maps. Serum/Plasma are plotted as open-coloured boxes and virus variants (antigens) as filled circles. Convalescent plasma (n=11) collected early in the pandemic (pre-Alpha), sera (n=10) from triple-vaccinated individuals and sera/plasma (n=22) from those who had received a BA.1 bivalent booster are shown as black, green, and blue boxes, respectively. The standards, WHO IS 20/136, NIBSC WR 21/334 and IS 21/338 are also plotted as red boxes. Virus variants are shown as coloured circles.

Figure 4. Neutralising antibody titres of panels of convalescent sera from hamsters experimentally infected with SARS-CoV-2 variants determined using an authentic virus Focus Reduction Neutralisation Test (FRNT)

Sera was collected from Golden Syrian hamsters at day 27 or day 28 post-infection with ancestral (A), BA.1 (B), BA.4 (C), BA.5.2.1 (D), XBB.1.5 (E), XBB.1.16 (F) or XBB.2.3 (G) SARS-CoV-2 variants. Median neutralisation titres (ND_{50}) were generated by FRNT using 14 authentic SARS-CoV-2 viruses. Data are presented as median and interquartile ranges of titre (ND_{50}). Fold-changes relative to the homologous virus were calculated by regression analysis using a mixed effect model and are presented where the difference is statistically significant ($p < 0.05$, two-way ANOVA with Tukey's *post hoc* test). The Lower Limit of Detection (LLOD) of 1/30 is indicated by the blue dotted line. The Lower Limit of Quantification (LLOQ) of 1/58 as determined for the ancestral assay, is indicated by the grey dotted line. NS = not significant at the $p < 0.05$ level.

Figure 5. Antigenic map of SARS-CoV-2 variants assessed against convalescent sera collected from experimentally infected hamsters

Sera was collected from Golden Syrian hamsters at day 27 or day 28 post-infection with ancestral, BA.1, BA.4, BA.5.2.1, XBB.1.5, XBB.1.16 or XBB.2.3 SARS-CoV-2 variants. Median neutralisation titres (ND$_{50}$) were generated by Focus Reduction Neutralisation Test (FRNT) using 14 authentic SARS-CoV-2 viruses and plotted using antigenic cartography. Serum/Plasma are plotted as open-coloured boxes and virus variants (antigens) as filled circles.