Title

Synaptic plasticity in the medial preoptic area of male mice encodes social experiences with female

and regulates behavior toward young

Kazuki Ito[1], Keiichiro Sato[1], Yousuke Tsuneoka[2,3], Takashi Maejima[4], Hiroyuki Okuno[5], Yumi
Hamasaki[1], Shunsaku Murakawa[1], Chihiro Yoshihara[2,6], Sayaka Shindo[2], Haruka Uki[1], Stefan
Herlitze[7], Masahide Seki[8], Yutaka Suzuki[8], Takeshi Sakurai[9], Kumi O Kuroda[2,6], Masabumi
Minami[1], Taiju Amano[1,2]*****

9
[1] Department of Pharmacology, Graduate School of Pharmaceutical Sciences, Hokkaido
University, 060-0812, Japan
[2] Laboratory for Affiliative Social Behavior, RIKEN Center for Brain Science, 351-0198, Japan
[3] Department of Anatomy, School of Medicine, Toho University, 143-8540, Japan
[4] Department of Integrative Neurophysiology, Graduate School of Medical Sciences, Kanazawa
University, 920-8640, Japan
[5] Laboratory of Biochemistry and Molecular Biology, Graduate School of Medical and Dental
Sciences, Kagoshima University, Kagoshima, 890-8520, Japan
[6] Dept. of Life Science and Technology, Tokyo Institute of Technology
[7] Department of General Zoology and Neurobiology, ND7/31, Ruhr-University Bochum, 44780,
Germany

[8] Department of Computational Biology and Medical Sciences, Graduate School of Frontier
Sciences, The University of Tokyo, 277-8562, Japan

[9] Faculty of Medicine/International Institute for Integrative Sleep Medicine (IIIS), University
of Tsukuba, 305-8575, Japan
* Correspondence to: Taiju Amano, Department of Pharmacology, Graduate School of

Pharmaceutical Sciences, Hokkaido University, Kita 12, Nishi 6, Kita-ku, Sapporo Hokkaido

060-0812, Japan Tel: +81-11-706-3246, Fax: +81-11-706-4987, E-mail:

tamano@pharm.hokudai.ac.jp

Keywords:

paternal behavior, infanticide, aggression, social behavior, medial preoptic area, bed nucleus of the

stria terminalis, electrophysiology

Summary

A dramatic shift from aggressive infanticidal to paternal social behaviors is an essential event for males after mating. While the central part of the medial preoptic area (cMPOA) has been shown to critically mediate the paternal behaviors in mice, how this brain region becomes activated by mating and subsequent interaction with pups has not been investigated. Here, we demonstrate that the reduction in inhibitory synaptic strength towards the cMPOA provided by posteriordorsal medial amygdala (MePD) neurons is a key event for the post-mating behavioral shift in males. Consistent with this, we found optogenetic disinhibition of Me^{Cartpt} to the cMPOA synapses reduces male aggression towards pups. We further provide evidence that the cMPOA of paternal mice mediated pup-induced neural plastic changes in the bed nucleus of the stria terminalis. These findings provide possible function of cMPOA neural circuits required for the reception to young in male mice.

Introduction

Reproductive behavior in male includes courting, mating and parental behavior. Polygynous male mammals, which form reproductive groups with one male and multiple females, often show aggressive behavior toward un-familiar infant conspecifics[1-5]. Infanticide of non-offspring young is commonly believed to improve reproductive success by achieving a shorter inter-birth interval for

females, a situation more favorable to breeding lower competition for resources, and the avoidance

of misdirected paternal investment[3,6]. After mating, however, the same male gradually stops

committing infanticide and becomes paternal, even toward non-biological offspring[5]. This behavioral

transition from infanticide to paternal care in C57BL/6 male mice is shown to require not only the

experience of ejaculation but also cohabitation with the pregnant female[7,8]. The exact neural

mechanisms governing this behavioral transition through social experiences with female and infant

remained unknown.

The medial preoptic area (MPOA) has been identified as one of the most important brain regions

regulating parental behavior[9]. Within the MPOA, the central part (cMPOA), located ventral to the

cluster of magnocellular oxytocin neurons (the anterior commissural nucleus), appears to play a pivotal

role in parental behavior because bilateral cMPOA lesions abolished this behavior and promoted

infanticide in both maternal and paternal mice[10,11]. Likewise, optogenetic activation of the MPOA in

virgin male mice delayed infanticidal behavior[11,12]. However, the neural circuit mechanisms that control

the MPOA neuronal activities was not clear. Previously, we reported that behavioral mode transition

from infanticide to parental was initiated by lesions of vomeronasal organ (VNO), which transmits

olfactory information to the accessory olfactory bulb[8]. Similar results were obtained in the experiment

using genetically mutated mice with impaired vomeronasal signaling[12]. Tracing studies using classical

retrograde tracer or rabies virus indicate that sensory information from accessary olfactory system is

mediated by the bed nucleus of the stria terminalis (BST) and the medial amygdala (Me) and is

transmitted to the MPOA[13-15]. It is reported that aggressive behavior toward pup is elicited by activation

of GABAergic Me[16, 17].

The present study evaluates the efficacy of synaptic transmission from the Me to putative

Galanin positive cMPOA neurons from three groups: (i) males that experienced mating and pup

exposure, (ii) males that experienced mating but had not pup exposure, and (iii) virgin males. We

observed that inhibitory inputs from the Me to cMPOA were suppressed ahead of pup delivery,

leading to cMPOA disinhibition. Optogenetic inhibition of Me–cMPOA synapses significantly

reduced the proportion of mice showing aggressive behavior toward pups. Furthermore, we found

paternal experiences increased synaptic inhibition in the rhomboid nucleus of the bed nucleus of the

stria terminalis (BSTrh), one of a downstream structure of the MPOA[11]. Together, findings indicate

that alterations in plasticity at Me–cMPOA synapses occur as a result of social experience with a

female partner, which may function to prime a shift in behavioral response toward pups. This is

followed by paternal experience with pups inducing changes in BSTrh.

Results

Sensitivity to GABAergic synaptic inputs from the Me to the cMPOA are suppressed by mating

experience and cohabitation with a late gestational female

To map the projections from the Me to cMPOA, we took advantage of the high expression of the

peptide, cocaine and amphetamine regulated transcript (Cart), in the subpopulation of posterior-

dorsal Me (MePD) neurons[18,19]. Within the Me, clusters of Cart-positive neurons were clearly

visualized in the MePD by immunostaining with an anti-Cart antibody (Figure S1A). We infused

biotin dextran amine (BDA), an anterograde neuron tracer, into the Cart-positive area of the Me

(Figure 1A and Figure B) and observed a projection to the cMPOA (Figure 1C and Figure S1B). The

cMPOA is located ventral to the anterior commissural nucleus, which contains magnocellular

oxytocin neurons and thus could be visualized using an antibody for the oxytocin-associated protein

neurophysin I (NPI)[10]. In contrast to the Cart-positive area of the Me, the Cart-negative area did

not send a dense projection to the cMPOA regardless of tiny BDA leak in Cart-positive area of the

Me (Figure S1C-F).

To address the cell-type in the cMPOA, the cMPOA neurons were manually collected using

a glass pipette and RNA-seq analysis was performed to evaluate mRNA expression levels. A six cell-

sample was combined and represented as data for one mouse. Here, we established three age-

matched experimental groups of male mice with different levels of paternal experience: 1) males

each co-housed with one female for mating, gestation, delivery, and remaining with the pups for the

first 3 postnatal days (paternal group), 2) males each co-housed with one female mating partner

until late gestation without exposure to pups (fathers in gestation experience; FGE group[20]), and 3)

virgin male mice. Table S1 lists the differentially expressed genes (DEGs) in cMPOA neurons among

virgin (V), FGE (F), and paternal (P) mice. Further, we evaluated the expression levels of 21

representative genes levels previously reported in activated MPOA neurons during social behavior[21]

to explore the recorded neuron's cell-type in this study and found no difference in their expression

levels among the groups (Figure 1D). Among all samples, expression level of *Galanin* (*Gal*) was the

highest. Previous studies have shown that the activation of Gal-Cre labeled MPOA neurons can

block infanticide in virgin male mice[12,14]. We visualized *Galanin* mRNA-positive neurons with

Alexa647 using *in situ* hybridization chain reaction (HCR). We observed a banded *Galanin*-

expressing region extending ventrally from the posterior part of the anterior commissure (AC) and

covering the cMPOA region (Figure 1E). Biocytin injected neurons during patch clamp recording

were visualized by reacting it with streptavidin conjugated to fluorescein isothiocyanate (FITC). We

observed that 85.0% (34 in 40 cells) of the recorded neurons were positive for *Galanin* mRNA co-

staining (Figure 1F and 1G). Meanwhile, Esr1, another marker gene in the MPOA region related to

parental behavior[21-23], was moderately detected among the groups (Figure 1D). Histological analysis

revealed that ERα-positive neurons (protein from the Esr1 translation) covered the cMPOA region,

which is locaged ventrally to NPI-positive neuronal clusters (Figure 1H). Compared with *Galanin*,

the ERα-positive neurons were broadly expressed in MPOA (Figure 1I). Further, 50% of patch clamp

recorded neurons (7 of 14 cells) were ERα positive (Figure 1J and 1K).

Next, to examine if social experience-dependent plastic changes in the cMPOA underlie the

shift from infanticidal to paternal behavior, we performed whole-cell patch clamp recordings of

cMPOA neurons from paternal, FGE, and virgin group mice to compare the synaptic responses.

There were no significant differences in passive membrane properties of cMPOA neurons among the

groups (Table S2 and S3). The projections including those from Cart-positive Me to the cMPOA were

electrically stimulated (Figure 1L), and the cell types of the recorded neurons were confirmed by post hoc *in situ*

hybridization staining. The average evoked inhibitory postsynaptic potential amplitude was

significantly smaller in the FGE group *Galanin*-positive cMPOA neurons compared with the virgin

group *Galanin*-positive cMPOA neurons (Figure 1M). A similar trend was observed in the

experiments in the blind targeted cMPOA neurons from the virgin and FGE groups (Figure S2A).

Considering that a high percentage of *Galanin*-positive neurons were targeted in the cMPOA

(Figure 1G), we considered that we were, in most cases, recording from *Galanin*-positive neurons.

Consequently, cell types are not discussed for each of the following experiments. Similar to FGE

mice, paternal mice showed a significant level of decreased eIPSP amplitude in the cMPOA

compared with the virgin mice (Figure 1N). We also addressed the effects of pregnant female

exposure alone (i.e., without the experience of copulation and delivery) on virgin males (virgin male

with late gestation; VMLG group, Figure S2B) or sexual experience without staying until late

gestation of female partner (male with mating experience; MME, Figure S2D). As a result, eIPSP

amplitude in the cMPOA in VMLG and MME mice were not changed (Figure S2C and S2E),

suggesting that either mating experience or cohabitation with a late gestational female was not

enough to reduce eIPSP amplitude in the cMPOA. To test the postsynaptic factors contribution to

plastic changes in paternal mice, a G-protein signaling blocker GDPßS was intracellularly infused.

The decreased eIPSP amplitude in the paternal group mice's cMPOA was reversed by GDPßS

intracellular infusion (Figure 1O). Collectively, these data suggested that mating and cohabitation

with a female, but not experience with pup, activates intracellular signals inducing disinhibition in

the cMPOA, which densely include *Galanin*-positive neurons.

Average evoked excitatory postsynaptic potentials (eEPSPs) amplitudes did not differ

between the groups (Figure 1P).

Me^{Cartpt} neuron projections to cMPOA neurons are modified by female exposure alone

To directly test if the eIPSP decrease observed in the cMPOA of FGE group mice is mediated by

input from Me neurons, we utilized Cartpt (which encodes Cart)-Cre mice. Cartpt-Cre × Ai9

reporter mice showed that 21.2% of NeuN-positive neurons in MePD were Cartpt (tdTomato)

positive (Figure 2A). We first infused an AAV vector encoding Cre-dependent eYFP into the Me of

Cartpt-Cre mice for pathway identification (Figure 2B). Indeed, we observed a clear Cre-dependent

eYFP expression in Me^{Cartpt} neurons and eYFP-labeled projections into the cMPOA (Figure 2C and

2D), consistent with the tracer study (Figure 1B and 1C). Next, we infused an AAV vector to express

blue light-sensitive channelrhodopsin, ChR2(H134R) in Me^{Cartpt} neurons and performed whole-cell

patch clamp recordings from cMPOA neurons in slices prepared from these mice. Application of blue

light to brain slices including the cMPOA depolarized the synaptic terminals derived from Me^{Cartpt}

neurons and induced both excitatory and inhibitory synaptic currents in cMPOA neurons (Figure 2E

and 2F), so we evaluated social experience-dependent synaptic changes at both excitatory and

inhibitory pathways. At the recording potential (−60 mV), glutamatergic and GABAergic ionotropic

receptor currents should flow in the opposite direction. Thus, glutamatergic and GABAergic

predominance can be distinguished by the net change in membrane current. Because of the

abundance of GABAergic neurons in the MePD[13], including retrogradely labeled cMPOA projecting

MePD neurons (Figure S3), we chose an experimental protocol that emphasized the EPSC

amplitude. The driving force for EPSC and IPSC, calculated by subtraction of the recording

potential (−60 mV) and reversal potential (−1 mV for EPSC, −87 mV for IPSC calculated by Nernst

equation), were about 59 mV and 27 mV, respectively. Under this experimental protocol, we counted

cMPOA neurons expressing predominantly blue light-induced EPSCs or IPSCs by the net

membrane current change. Whereas 83.3% of cMPOA neurons were IPSC-dominant in virgin male

mice, only 21.4% were IPSC-dominant in FGE males (Figure 2G). In accordance with the electrically

evoked responses (Figure 1O), intracellular perfusion of a G-protein signaling blocker GDPβS into

the cMPOA neurons reversed the ratio of IPSC- dominant neurons in FGE males to the level of

virgin males. As expected, control group mice expressing only eYFP did not show light-evoked

synaptic responses (Figure 2G). The data presented so far suggest that experiencing mating and

cohousing with a pregnant female impaired GABAergic synaptic transmission from Me^{Cartpt} into the

cMPOA.

Optogenetic inhibition of Me-to-cMPOA inputs change the spontaneous synaptic activity

To examine whether inhibition of Me^{Cartpt} terminals in the cMPOA modulate neural and/or

synaptic activity, we attempted to specifically silence these synapses. For this, we infused an AAV

vector encoding vertebrate long-wavelength opsin (vLWO)[24,25] bilaterally into the Me of virgin

Cartpt-Cre mice to suppress Me^{Cartpt} inputs (Figure 3A). To confirm the inhibitory effects of vLWO,

we performed whole-cell patch clamp recordings from Me^{Cartpt} neurons expressing vLWO-eGFP.

Application of green light hyperpolarized Me^{Cartpt} neurons (Figure 3B and 3C) but did not in the

control group (Figure 3D and 3E).

Next, we infused an AAV vector encoding ChR2 and vLWO into the Me of virgin Cartpt-Cre

mice to confirm the inhibitory effects of vLWO at the synaptic terminal (Figure 3F and 3G). After

more than 4 weeks, brain slices were prepared and recorded from cMPOA neurons. Green light

could affect ChR2 and therefore instead of using green light as above, red light was used to

stimulate the vLWO. The amplitude of blue light induced IPSC was significantly reduced by vLWO

activation (Figure 3H and 3I). Furthermore, the suppressed synaptic inputs from the Me^{Cartpt} to the

MPOA could result not only in reduced direct synaptic transmission but also in more frequent

inhibitory synaptic currents by disinhibition of surrounding interneurons. To address this

hypothesis, we observed spontaneous IPSCs (sIPSC) in each cMPOA neuron before and after the

suppression of inputs from the Me^{Cartpt} via vLWO activation (Figure 3J-L). Within 25 recorded

neurons, sIPSC frequency was increased more than 10% in 13 neurons and reduced more than 10%

in 1 neuron. The averaged frequency of the sIPSC in cMPOA was significantly increased (Figure 3L).

The application of tetorodotoxin (TTX) blocked the effects of vLWO activation (Figure 3M and 3N).

These data suggest that inhibition of the Me^{Cartpt} inputs to the MPOA could modify the activity of

each neuron in the cMPOA by acting on inter-neuronal network.

In vivo optogenetic inhibition of Me-cMPOA inputs suppresses infanticidal behavior

We previously demonstrated that Me neurons in infanticidal virgin males are highly activated and

express c-Fos protein following exposure to donor pups contained in wire mesh ball[8]. Consistent

with those results, in this study we observed that exposure of infanticidal virgin males to a pup in

wire mesh ball resulted in a significantly increased number of c-Fos- and eYFP-positive Me^{Cartpt}

neurons as compared with a wire mesh ball exposure alone (control) (Figure 4A-C).

Next, we aimed to address the causal relationship between synapse plastic changes in the

cMPOA and behavioral changes. To address the behavioral causality of the impaired Me-cMPOA

inputs, we infused an AAV vector encoding vLWO into the Me of virgin Cartpt-Cre mice and two

optic fibers were bilaterally implanted above the cMPOA for optogenetic stimulation. Green light

was delivered to the cMPOA during behaviors toward pups being observed (Figure 4D). The latency

to first sniffing did not differ significantly between the vLWO group and control group (Figure 4E).

However, the proportion of vLWO-expressing mice that did not show infanticidal behavior was

significantly larger than in the control group on 4 successive days (Figure 4F and 4G). After the 4-

day experiments with green light delivery, we examined the behavioral pattern toward pups

without green light delivery for some individuals. The proportion of vLWO group mice that showed

infanticidal behavior did not differ between day 4 (with light) and post-test (without light) (Figure

S4). Similar results were observed in the experiments, in which Me input to cMPOA was inhibited

in vGAT-IRES-Cre mice (Figure 5). Taken together, these data suggest that Me[Cartpt] neurons

normally serve to suppress cMPOA neurons and promote infanticidal behavior.

Paternal experience enhances inhibitory transmission in the BSTrh through a postsynaptic

mechanism

We have demonstrated that the decreased inhibition of MPOA by Me input upon experiences with

female is a key event for suppressing the infanticide in the FGE. However, about half of FGE mice

still show the infanticidal behavior, whereas almost all paternal mice do not show infanticidal

behavior[8]. Next, we examined the synaptic change in paternal mice upon experiences with pup.

Downstream structure of the MPOA, the rhomboid nucleus of the bed nucleus of the stria

terminalis (BSTrh), has been shown to be activated by infanticidal behavior and negatively

regulated by cMPOA[11]. To address the impact of social experiences on the BSTrh synaptic property,

we measured the changes in the excitatory－inhibitory (E/I) ratio of the virgin, FGE and paternal

groups, in a manner similar to our previous studies[26]. The average amplitude of IPSPs evoked by

electrical stimulation of the stria terminalis (Figure 6A) was enhanced in the BSTrh of the paternal

group compared to FGE and virgin groups (Figure 6B). However, the average eEPSP amplitudes did

not differ significantly between these groups (Figure 6C). These data suggest that, in the paternal

mice, increased inhibition in the BSTrh critically takes place following the disinhibition in the Me–

cMPOA pathway as was seen in FGE mice.

We hypothesized that the cMPOA plays a critical role in the increase of inhibition in the

BSTrh. To evaluate this hypothesis, we specifically lesioned the MPOA. To this end, we bilaterally

injected NMDA into the cMPOA in virgin and paternal mice (Figure 6D and S5) and, 3 to 9 days

later, performed whole-cell recordings from BSTrh. In the BSTrh of ipsilaterally lesioned paternal

mice, the average eIPSP amplitude was significantly smaller than unlesioned paternal mice (Figure

6E). These data strongly suggest that the increased inhibitory input to the BSTrh is mediated by

the cMPOA in mice showing paternal behavior. It should be noted that cMPOA fiber inputs into

BSTrh were not likely stimulated to evoke synaptic responses in the current protocol because

cMPOA lesion unchanged eIPSP amplitude in virgin mice (Figure 6F).

We next addressed if this increased inhibition onto the BSTrh takes place through a pre- or

post-synaptic mechanisms. There were no significant differences in passive membrane properties

among the three groups (Table S4). We carried out patch-clamp analysis on BSTrh of paternal group

mice and found out that while an inclusion of the GDPβS in the pipette does reduce the amplitude of

eIPSPs to the level of virgin group (Figure 6G), eEPSP did not show obvious changes (Figure 6H).

These data suggest that the increased eIPSP amplitude in the BSTrh occurs mostly by postsynaptic

mechanisms. We also analyzed the paired-pulse ratios to compare the release probability from

presynaptic terminal and found out that the ratio of eEPSCs and eIPSCs did not differ between

paternal and virgin groups (Figure 6I). Altogether, we conclude that the cMPOA in the paternal

mice mediates plastic changes at GABA synapses in the BSTrh mainly by postsynaptic mechanisms,

and these sequential events are centrally involved in the fundamental switch in the paternal

behaviors of the father animals.

Discussion

The results showed that multi-step plastic changes in synapses correlate with the transition from

infanticidal to parental behavior in male mice. The inhibitory synapses from Me[Cartpt] neurons in the

cMPOA were impaired by interactions with females. This plastic change in synapses is the potential

trigger for the behavioral transition from virgin to FGE. Our targeted cMPOA area are composed by

Galanin-positive neurons (Fig 1D-G), consistent with the previous reports that the activation of

Galanin-positive MPOA neuron cause inhibition of aggressive behavior toward pup[12,14]. In addition,

the shift from FGE to paternal was accompanied by an increased BSTrh inhibition. These plastic

changes caused the change of balance of excitatory–inhibitory synaptic transmission (E/I balance),

which correlated positively with the neuronal activity of the cMPOA and BSTrh, when virgin or

paternal males contacted with pups.

It was previously reported that the number of males showing infanticide decreased about

two weeks after ejaculation without cohabitation with the female partner[7,27]. This discrepancy may

be due to strain differences or by the different experimental/breeding environments. The behavior of

males toward pups appears to be controlled by factors triggered by ejaculation, which remain

effective for several subsequent weeks, thus preventing infanticide of the male's own offspring.

Followingly, additional mechanisms, such as activation of oxytocinergic neuron in paraventricular

nucleus of hypothalamus to initiate parental care[28], would complete a series of behavioral changes

of male, from aggression to parenting onset.

The Me has been implicated in various social behaviors, including social investigation,

mating, and aggression[29-33]. In males, the functional association between the Me and the MPOA is

critical for mating behavior[34,35]. The Me comprises several subdivisions with distinct projections and

cell types[36,37], including the MePD, which is a sexually dimorphic nucleus crucial for aggressive

behaviors[29,31,38,39]. We found the level of pup exposure-induced c-Fos expression in the VNO and

MePD was higher in infanticidal virgins than in paternal male mice[8]. On the other hand, the eIPSP

amplitude in cMPOA neurons of paternal mice elicited by the stimulation of axonal fibers derived

from Me was reversed by intracellular GDP-ßS infusion (Figure 1O). These data suggest that both

decreased efficacy in synaptic transmission in cMPOA neurons, which we report, and hypoactivity of

MePD neurons likely contribute to behavioral selection in male mice. However, the notion that the

decrease of Me-to-cMPOA inputs in paternal mice causes a behavioral transition from infanticide to

parenting can be overly simplistic, since other brain areas should also provide inputs into the

cMPOA and can be modified by mating experiences. Further research is needed to elucidate this

phenomenon of behavioral transition in male mice.

Although we utilized Cartpt-Cre mice to map the Me−cMPOA pathway, Cart may function

as a neurotransmitter[40] and so influence Me to MPOA transmission. However, agonists and

antagonists specific for Cart transmission have not been identified, so we could not address the

influence of Cart released from the Me. In this study, we confirmed that manipulation of Me inputs to

the cMPOA in vGAT-IRES-Cre mice led to behavioral changes like those observed in Cartpt-Cre mice

(Figure 5). By considering data showing disinhibition in the cMPOA (Figure 1 and 2), we consider that

the plastic changes of GABAergic transmission at Me synapses strongly contribute to the shift from

infanticide to paternal behavior. On the other hand, we detected EPSCs by optical stimulation of Me

inputs into the cMPOA of Cartpt-Cre mice. Therefore, it is possible that subpopulation of Me neurons

might specifically control some cMPOA neurons and the associated behavioral pattern. The MPOA

contributes to several types of intrinsic behaviors, which in turn activate the MPOA according to

different patterns[11,23,32,33,41-44]. However, the mechanisms underlying the expression of one

appropriate behavior remains unknown. Suppressing the accessory olfactory signals mediating the

Me, presumably changes neural activity patterns in the cMPOA. This is supported by the results

that temporal disinhibition of the Me−cMPOA pathway modifies behavioral patterns and increases

cMPOA inter-neuronal network activity, as reflected by the increased frequency of postsynaptic

currents (Figure 3). In addition to the Me, other inputs into the cMPOA would drive several inter-

neuronal network activity patterns to achieve the multi-functional MPOA and behavioral choices.

Several studies report that activation of MPOA neurons in virgin males disturbs infanticide,

although parental behavior is not initiated[11,12]. Other studies report that activation of MPOA

neurons expressing Esr1 or afferents to ventral tegmental area (VTA) promotes parental behavior,

as measured by the latency to retrieve pups to the nest[22,23]. Here, the basal motivation for parenting

should be carefully distinguished from the behavioral choice toward the pups.

Uncovering the primary trigger of cMPOA plastic changes would promote the

understanding of how mammalians become parental. Neither mating nor cohabitation with a

pregnant female caused sufficient cMPOA disinhibition (Figure S2B and S2C). Because no

significant cMPOA disinhibition was observed in virgin males co-housed with a late-gestation

pregnant female (the VMLG group), the sensory inputs from pregnant females were unlikely to

induce synaptic changes in the cMPOA (Figure S2D and S2E). However, our cohabitation protocol

lasted only 3 days; it is possible that longer cohabitation is required for cMPOA disinhibition,

although longer cohabitation may be confused by the Bruce effect (i.e., spontaneous pregnancy

termination triggered by the scent of an unfamiliar male). Moreover, mating experience without

cohabitation with the female partner until late gestation did not change the GABAergic

transmission in the cMPOA. We are focusing on the previous reports that mating experiences

increase the dopamine turnover[45] and that plastic changes of the expression level of dopamine D_2

receptor in the MPOA were reported[46]. Because the function of dopamine receptors can change over

time[47,48], this may help explain the behavioral transition triggered by the social experience with

female partners. We observed that intracellular GDPßS infusion into cMPOA neurons reversed

eIPSP amplitude and the ratio of inhibitory- to excitatory-dominant neurons (Figure 1O, and 2G).

These data suggest that postsynaptic GABA$_A$ receptor activity on cMPOA neurons is downregulated

(i.e., neuronal disinhibition) by intracellular signals triggered in response to social experience with

females. Multiple signaling molecules are reported to be involved in the function of GABA$_A$

receptors[49]. Moreover, altered gene expression levels among virgin, FGE, and paternal mice were

recently reported[50]. Among them, the intracellular signaling molecules controlled by GPCR or small

G proteins are one of the possible targets to address the mechanisms of plastic changes in the

cMPOA.

Interestingly, unlike the synaptic changes in the Me^{Cartpt} - cMPOA pathways of FGE mice,

an inhibitory shift was observed in the BSTrh E/I balance, which was induced by the caring

experience (Figure 6). What is the biological significance of the increased BSTrh inhibition? Like

previous report[11], we observed that BSTrh inactivation by hM4Di significantly delayed infanticidal

behavior in virgin male mice without parental behavior initiation (Figure S6). This suggests that

BSTrh does not block parental behavior directly. Considering the time course for increased

inhibition between FGE and paternal mice, BSTrh plastic changes could contribute to pup

sensitization, which underlie the onset of pup caring. The cMPOA lesions reversed the increased

BSTrh inhibition in paternal mice to the level of virgin mice, suggesting that the inhibitory

transmission of BSTrh was controlled by cMPOA (Figure 6D and 6E). This is supported by the

following facts (1) cMPOA neurons project into the BSTrh and (2) unilateral lesion of the cMPOA

increases the number of c-Fos positive neurons in the ipsilateral BSTrh of mice exposed to pups[11].

Because of increased BSTrh inhibition in paternal mice is sensitive to intracellular GDPβS infusion

(Figure 6G), some metabotropic receptors of neurotransmitter or peptide may be influencing BSTrh

during and/or after pup sensitization. The cMPOA neurons activated during parental behavior

contain many kinds of neurotransmitters, including Galanin, neurotensin, and tachykinin2[10,51].

Moreover, the anterior commissural nucleus, adjacent to the cMPOA, contains the third largest

population of magnocellular oxytocin neurons. Therefore, cMPOA neurons or adjacent neurons may

directly and persistently control $GABA_A$ receptor activity in the BSTrh of paternal mice via

metabotropic receptors.

Converging data suggest that the cMPOA neurons contribute to both infanticide inhibition

and parental behavior by modifying the inter-neuronal network and projecting to the downstream

including BSTrh. There is no doubt that uncovering the relevant MPOA subgroup for each function

requires the analysis of neural circuits. Moreover, understanding the modulatory mechanism of

synaptic plasticity in the cMPOA and BSTrh may provide evidence for possible interventions in

parental stress and subsequent child maltreatment.

Author contributions

TA conceived the study with support from KOK. TA, TS and MM designed the experiments. TA, KI,

and KS conducted the behavioral experiments. TA and KI performed stereotaxic surgeries. TA, YT,

KI, KS, YH, SM and SS performed histological experiments. TA and HU performed whole-cell patch

clamp recordings. TM, TS and SH developed the AAV virus expressing vLWO. TA and HO prepared

sample for RNA sequence and YS, MS and KS analyzed data. TA wrote the manuscript with support

from KOK and YT. All authors contributed to editing the manuscript.

Acknowledgements

This work was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant

Number 25713044, 16K19747, 16H06279 (PAGS), 18K07584, 21K07494 (to TA), 26282220 (to KOK),

15K01832 (to TM), 19H03328, 20H05068 (to HO), Suzuken Memorial Foundation (to TA), Narishige

Neuroscience Research Foundation (to TA), Nishinomiya Basic Research Fund, Japan (to TA), and

the fellowship of Special Postdoctoral Researchers Program at RIKEN (to TA), Hokkaido University,

Global Facility Center (GFC), Pharma Science Open Unit (PSOU), funded by MEXT under "Support

Program for Implementation of New Equipment Sharing System," RIKEN Brain Science Institute

(2011-2015 to KOK), Deutsche Forschungsgemeinschaft He2471/18 Priority Program (SPP1926),

SFB874, SFB1280, DFG2471/21, DFG2471/23 project B10 (to SH). We appreciate Dr. Goichi Miyoshi

for critically reading this manuscript prior to submission, Dr. Shigeyoshi Fujisawa and Kazunari

Miyamichi for supporting genetically modified mice transportation, RIKEN Research Resource

Center for maintenance of animals, Kiyomi Imamura, Terumi Horiuchi, Dr. R. Jude Samulski and

the UNC Vector Core for helpful supports.

STAR Methods

Animals

The Animal Experiment Committee of the RIKEN Brain Science Institute and Institutional Animal

Care and Use Committee at Hokkaido University approved all animal experiments, which were

conducted in compliance with the National Institute of Health guidelines for the care and use of

laboratory animals.

Animal maintenance and male exposure to different levels of mating and paternal experience

Most C57BL/6J male mice were bred at the RIKEN Brain Science Institute and Graduate School of

Pharmaceutical Sciences, Hokkaido University unless specially described. For the BDA (Thermo

Fisher Scientific) tracer injection into Me (Figure 1A-C and S1), we utilized C57BL/6J male mice

from Japan SLC. Cartpt-Cre mice (Jackson Laboratory, stock number 009615), vGAT-IRES-Cre

mice (Jackson Laboratory, stock number 016962) and Ai9 (Jackson Laboratory, stock number

007909) were bred at the Graduate School of Pharmaceutical Sciences, Hokkaido University. We

backcrossed vGAT-IRES-Cre mice with C57BL/6J mice at least five generations after arrival from

Jackson Laboratory. Mice were housed in individually ventilated cages and provided ad libitum

access to water and food and maintained under a 12-h light/dark cycle in cages lined with TEK

Fresh Standard bedding (Envigo). Mice were weaned at postnatal day 28 (P28). All mutant

heterogenic mice were co-housed with wild-type mice until the experiments. To produce paternal

group mice, a virgin male mouse was housed with a female for mating. After delivery, the male

stayed with pups for 3 days. All paternal group males experienced one delivery and were used for

experiments within 20 days after the birth of pups. If pups did not survive for more than 3 days, the

male was not included in the paternal group for experiments. Other virgin male mice were mated

and co-housed with a female only until late gestation but not delivery, termed 'fathers in gestation

experience' (FGE) mice.

Stereotaxic surgery

BDA (1.25 µg/ dissolved in 12.5 nL 0.1 M phosphate buffer), N-methyl-D-aspartic acid (NMDA,

Sigma-Aldrich, 20 mg/ml in saline), and AAV5-EF1α-DIO-hChR2(H134R)-eYFP (6.6×10^{12} genome

copies/ml, provided by Vector Core at the University of North Carolina (UNC) at Chapel Hill, UNC

vector core), AAV5-EF1α-DIO-eYFP (5.6×10^{12} genome copies/ml, UNC vector core), AAV10-EF1α-

DIO-vLWO-eGFP-5HT1A (3.7×10^{13} or 1.8×10^{13} genome copies/ml, UNC vector core), AAV10-EF1α-

DIO-eYFP (1.2×10^{14} or 3.7×10^{13} genome copies/ml, UNC vector core), rAAV2-EF1a-DO_DIO-

tdTomato_EGFP-WPRE (more than 7.0×10^{12} vg/ml, Addgene), AAV2-hSyn-HA-IRES-eGFP ($3.3 \times$

10^{12} genome copies/ml, UNC vector core), and AAV2-hSyn-HA-hM4Di(Gi)-IRES-mCitrine (5.6×10^{12}

genome copies/ml, UNC vector core) (100−400 nL/hemisphere), were administered under anesthesia.

Briefly, 2−3-month-old male mice were anesthetized by intraperitoneal (i.p.) sodium pentobarbital

(30 mg/kg) or the mixture of medetomidine hydrochloride (0.3mg/kg), midazolam (4mg/kg),

butorphanol tartrate (5 mg/kg)[52], and local subcutaneous (s.c.) lidocaine hydrochloride. The skull

was exposed and holes drilled for stereotaxic injection. Glass capillaries of ~50 μm tip diameter were

filled with oil and backfilled with the test drug or AAV vector. Injection was targeted by referring to

the Mouse Brain Atlas of Flanklin & Paxinos (2007) coordinates for the Me (AP −1.5 mm, ML 2.1

mm, DV −5.4 mm and AP −1.2 mm, ML 1.9 mm, DV −5.6 mm), cMPOA (AP 0.1 mm, ML 0.55 mm,

DV −5.1 mm), and BSTrh (AP −0.1 mm, ML 1.2 mm, DV −4.2 mm). After injection, the skin was

closed with a nylon suture. Mice were kept in single housing 3−4 days for recovery. Mice injected

with AAV vectors were then maintained under group housing for more than 4 weeks to allow for

expression of vector genes. One week before behavioral testing, optic fibers were implanted into the

targeted area as described below.

Histological analysis

Mice were anesthetized by sodium pentobarbital (50 mg/kg, i.p.) and transcardially perfused with

4% paraformaldehyde (PFA) dissolved in phosphate buffered saline (PBS, pH 7.4). Brains were

removed from the skull and post-fixed overnight in PFA at 4 °C, followed by incubation in PBS/20%

sucrose for 1 day and PBS/30% sucrose for 1−2 days. Brains were then embedded in O.C.T.

Compound (Sakura Finetek) at −80 °C and sectioned at 40 μm using a Cryostat (Leica Biosystems)

unless otherwise indicated. Some brains were used to prepare sections for electrophysiological

analysis. In the case of BSTrh recording after MPOA lesion, serial slices including MPOA were

sectioned at 80 μm in cutting solution (see below) using a Leica VT1200 Semiautomatic Vibrating

Blade Microtome (Leica). These slices and the slices including BSTrh used for electrophysiological

analysis were fixed in 4% PFA and immunostained.

For fluorescent immunohistochemical staining, brain slices were incubated in PBS

containing 0.1% Triton X-100 (PBST) for more than 15 min and then in 0.4% Block Ace (Dainihon-

Seiyaku) for an hour. Slices were then incubated at 4 °C overnight in Block Ace solution containing

primary antibody. After several rinses in PBST, slices were incubated in PBST containing secondary

antibody at room temperature. Table S5 shows antibody list.

To stain the BDA anterograde tracer, sections were incubated with Alexa 568-conjugated

streptavidin (1:500, Thermo Fisher Scientific) for 2 hours. After washout of the secondary antibody

or streptavidin, slices were mounted on glass slides with VECTASHIELD Mounting Medium (Vector

Laboratories). For 3, 3-diaminobenzidine (DAB) staining, immunolabeled slices were incubated in

0.1 M glycine for 10 min, 0.5% H_2O_2 for 30 min, and VECTASTAIN ABC reagent (Vector

Laboratories) overnight at 4 °C. After rinsing several times with PBST, the slices were incubated

with Vector DAB Substrate (Vector Laboratories) including nickel chloride for about 5 min. The

stained slices were mounted on gelatin-coated glass slides and dried. After nissl staining, slides

were treated with Softmount (Wako). Labeling of *Galanin* positive neurons followed modified

protocol of *in situ* hybridization chain reaction (HCR) using short hairpin DNAs[53,54] using probes

listed in Table S6. Fixed brain section re-sliced at 75 μm after patch clamp recording with internal

solution containing 1% biocytin (Sigma) and 100 nU RNase inhibitor (Promega), or 30 μm brain

section fixed by 4% PFA were soaked in methanol for 10 minutes. After washing with PBST, the

sections were prehybridized for 10 min at 37°C in a hybridization buffer containing 10% dextran

sulfate, 1× standard saline citrate (SSC), 0.1% Tween 20, 50 μg/ml heparin, 1× Denhardt's solution.

The sections were moved to another hybridization solution containing a mixture of 20 nM split-

initiator probes, and incubated overnight at 37°C. For *in situ* HCR, 3 μM hairpin DNA solutions

were separately snap-cooled before use. The sections were incubated in amplification buffer (10%

dextran sulfate in 5× SSCT) with 60 nM hairpin DNA pairs for 2 hours at 25°C. Then, the samples

were washed with PBST three times at room temperature.

Immunofluorescence images were captured using an incident-light fluorescence microscope

(Leica DM6000B, Leica), confocal laser-scanning microscope (FV-10, Olympus), or fluorescence

microscope (BZ-X700, Keyence). To allow for comparison among trials, the contrast and brightness

of all photographs were adjusted linearly and uniformly using Adobe Photoshop CS5 (Adobe

Systems) unless specially described. For the lesion study, sections were immunostained with anti-

NeuN antibody to confirm cell death in the target area. The loss of NeuN-immunoreactive cells in

the NMDA-injected group was compared with the saline group to judge the lesioned area.

Electrophysiology

Electrophysiological recordings were performed as described previously[26]. The artificial cerebral

spinal fluid (ACSF) contained 126 mM NaCl, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 1 mM $MgCl_2$, 2 mM

CaCl$_2$, 26 mM NaHCO$_3$, and 10 mM glucose (pH 7.3). For a cutting solution, NaCl in ACSF was

replaced with the same concentration of choline chloride. Mice were injected with pentobarbital (30

mg/kg, i.p.) followed by transcardiac perfusion of ice-cold cutting solution. The brains were removed,

and 230-μm brain slices including the target area prepared in ice-cold choline chloride-based cutting

solution using a Leica VT1200 Semiautomatic Vibrating Blade Microtome. Brain slices were stored

in warmed ACSF at 32 °C for 20–30 min and then kept at room temperature until recording.

For electrophysiological recordings, slices were superfused with 32 °C–34 °C ACSF at 2–4

mL/min in a chamber mounted on a microscope. Neurons were identified using a 40× or 60× lens

and an infrared camera (IR-1000, DEGE-MTI). Glass electrodes (World Precision Instruments) of 4–

9 MΩ resistance were used for whole-cell patch clamp recordings in response to electrical

stimulation using ~0.5 MΩ glass electrodes containing ACSF. Recording electrodes were filled with

a potassium-based internal solution (132 mM K-gluconate, 3 mM KCl, 10 mM HEPES, 0.5 mM

EGTA, 1 mM MgCl$_2$, 12 mM Na-phosphocreatine, 4 mM Mg-ATP, 0.5 mM Na-GTP, 0.2-0.3 %

biocytin, pH 7.25) and signals were recorded using an Axopatch 700B amplifier (Molecular Devices).

Only cells with access resistance <30 MΩ and exhibiting action potentials with >60 mV amplitude

evoked by positive current injection were included in the analysis. Whole-cell currents were filtered

at 3 kHz. The liquid junction potential (11 mV) was compensated. Stimulating electrodes were

placed at the dorsal BSTrh or cMPOA to stimulate stria terminalis. To activate the ChR2-expressing

axon terminals, we utilized a LED (465 nm, LEX2-LZ4-B, Brainvision Inc.). To activate the vLWO-

expressing neurons, green light generated by a 100 W mercury lamp (U-RFL-T) was applied

through a fluorescence band pass filter (520-550 nm) (Olympus). Red light was provided by Fiber-

Coupled LED (625 nm, Thorlabs). Evoked synaptic currents and potentials were measured at least

three times at the same stimulus intensity and averaged. Responses with action potentials were

omitted from the analysis. We recorded evoked excitatory postsynaptic potentials (eEPSPs) and

eEPSCs at a holding potential of −85 mV in the presence of 100 μM picrotoxin. eIPSPs were

recorded at −65 mV and eIPSCs at −60 mV in the presence of 20 μM CNQX and 20 μM MK-801. In

experiments using optogenetic suppression using vLWO, slices were also perfused with 25 μM 9-cis-

retinal, 0.1% dimethyl sulfoxide, 0.025% (±)- α-tocopherol and 0.2% bovine serum albumin.

Behavioral test

The tests of paternal behavior toward pups were performed as described previously[26]. Mice were

housed individually in cages containing new purified paper bedding (Alpha-Dri, Shepherd Specialty

Papers) and a cotton square (Nestlet, Ancare). After 1–2 days, three pups aged 1–6 days were placed

into the cage corner avoiding the nest. Tests were performed once each day for 30 min on 4

successive days during the daytime unless otherwise indicated. The endpoint of the aggressive

behavior toward pups to terminate the experiments was 2-s screams or visible wounds on the pup

skin. Wounded pups were immediately euthanized. The paternal behavioral score was evaluated

according to previous reports as follows: 4 = all pups were retrieved, 3 = 1 or 2 pups were retrieved,

2 = no pup was retrieved, 1 = at least one pup was attacked >3 min after placement in the test cage,

and 0 = at least one pup was attacked within 3 min after placement in the test cage.

For c-Fos mapping, the procedure was modified from a previous study[8]. Briefly, mice were

exposed to a pup protected inside wire-mesh ball (tea balls, 45 mm diameter; Minex Metal, Tsubame,

Japan) with about 10 holes (5 mm diameter) placed in a cage corner avoiding the nest for 30 min.

Ninety minutes later, the adult male was anesthetized with pentobarbital (50 mg/kg, i.p.) and

transcardially perfused with 4% PFA/PBS for brain isolation as described above. The c-Fos

immunoreactive cells were counted automatically by Image J software (National Institutes of

Health, Bethesda, MD, USA).

Optogenetic and chemogenetic manipulation of neural activity in vivo

Zirconia ferrules optical fibers (Thorlabs) of 200 μm diameter core (NA = 0.37) were cut and polished

for >70% coefficient of transmission. These constructs were bilaterally implanted into the mouse

brain area 1-mm above the target area and fixed with dental cement. After the mouse had recovered

from the surgery (one week later), the optic fibers were connected via Ceramic Split Mating with

1×2 Branching Fiber-optic Patch Cords-Glass (0.37 NA, Doric Lenses) to a diode laser (450 nm or

532 nm, Changchun New Industries Optoelectronics Tec.) through a FC/PC connector. Averaged

light power at the tips between the bilateral fiber tips was set to 30−40 mW (450 nm) or 10−12 mW

(532 nm). Light pulses were controlled by TTL Pulse Generators (Doric Lenses). After connecting

the fibers, experimental mice were allowed free actiTon for 15 min. During the last minute of the

behavioral test, mice received constant light on for 20 s without light pulses followed by a 10-s off

period. This photostimulation protocol was continued throughout the 30-min behavioral experiment.

After the behavioral experiments, animals were perfused with 4% PFA to prepare the brain slices.

Only animals with bilateral expression of fluorescence labeled fiber and the bilateral optic fibers

above targeted area between 0.3-1.0 mm from the targeted area were included for the data analysis.

For the chemogentic silencing, we infused CNO (2.0 mg / kg, i.p.) and observed the behavioral

pattern 30-35 min later. After the behavioral experiments, we immunostained the brain slices with

anti-GFP. Only animals with bilateral expression of fluorescent signal in the targeted area were

included for the data analysis.

RNA-seq analysis

Mouse brain slices were prepared using essentially the same procedures for electrophysiological

experiments. Six cMPOA neurons were manually collected by using $1 \sim 2$ MΩ glass electrodes

containing ACSF and suspended in cell lysis buffer (0.5% Triton-X100, 0.5 U/μL RNase inhibitor, 5

ng/μL yeast rRNA). The cell samples were immediately frozen and stored at −80°C until starting

reverse transcription. Reverse transcription and cDNA library preparation were performed using

the Smart-seq2 method[55] with some modifications. Briefly, 12 μM dT$_{30}$VN primer (5'-

AAGCAGTGGTATCAACGCAGAGTACT$_{30}$VN-3) were added into microtubes containing cell lysates

(final concentration, 1 μM); then, samples were heat-denatured at 70 °C for 3 min, followed by

immediate incubation on ice. Next, samples were mixed with a reverse transcription reaction

solution containing locked nucleic acid (LNA)-template switching oligo primer (5'-

AAGCAGTGGTATCAACGCAGAGTACrGrG+G-3', rG; riboguanosines, +G; LNA-modified guanosine,

final concentration of 0.6 μM) and reverse transcriptase (SMARTscribe, Clontech, final

concentration of 5 U/μL), and incubated at 42°C for 120 min followed by at 70°C for 10 min to

terminate the reaction. The resultant cDNA samples were purified using solid phase reversible

immobilization (SPRI) beads (AMPure XP, Beckman Coulter). The purified cDNA samples were

amplified by PCR using the IS primer (5'-NH$_2$-AAGCAGTGGTATCAACGCAGAGT-3', final

concentration of 0.24 μM) and Tks Gflex DNA polymerase (Takara Bio, final concentration of 0.025

U/μL). The amplified cDNA samples were purified by SPRI beads. The quality and quantity of cDNA

sample was confirmed by an Agilent 2100 Bioanalyzer using the High sensitivity DNA kit (Agilent

Technology, CA, USA).

The cDNA samples were then processed to generate indexed libraries for sequencing using a

NEBNext Ultra II DNA Library Prep Kit for Illumina (#E7645, New England Biolabs Inc., Ipswich,

MA, USA), according to manufacturer's instructions. Then, libraries were sequenced for 50-bp

single-end reads using HiSeq 3000 (Illumina, San Diego, CA, USA). Output fastq files were aligned

to the mouse genome (GRCm38/mm10) using STAR aligner[56]. Library size normalization was

performed using R (v.4.1.0) by calculating transcripts per million (TPM). Genes with <5 TPM in all

samples were excluded from the analysis. We selected 22 genes for neuronal cluster-mediated social

behavior in the MPOA based on previous reports[21]. However, *Oprd1* was excluded because of its low

expression. Heatmaps were drawn using gplots (v.3.1.1). Differentially expressed genes (DEGs)

between virgin, FGE, and paternal mice were analyzed using DESeq2 (v.1.32.0). Genes with a false

discovery rate-adjusted $p \leq 0.05$ and a $\log_2$ fold change ≥ 2 in either direction were considered DEGs.

Statistical analysis

Group means were compared by Fisher's exact test, the two-tailed t-test, one-way or two-way

repeated measures ANOVA followed by a post hoc test as indicated. All statistical tests were two-

tailed and conducted with GraphPad Prism software 6 and 9 (GraphPad Software, Inc., La Jolla, CA,

USA). A P-value < 0.05 was deemed statistically significant. Statical data was written in the figure

legends and Table S7.

Data availability

All data that support the findings presented in this study are available from the corresponding

author upon request. The raw sequences have been deposited in the DNA Data Bank of Japan

(DDBJ) under the DDBJ BioProject umbrella with accession number PRJDB8470, and to the DDBJ

Read Archive DRA008585 (BioSample ID: SAMD00175991- SAMD00176002).

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Taiju Amano (tamano@pharm.hokudai.ac.jp).

Figure Legends

Figure 1 The medial amygdala sends projections to the cMPOA

(A) Medial amygdala in the parasagittal section. HC; hippocampus, cp; cerebral peduncle, opt; optic

tract, EA; extended amygdala, AHi; amygdalohippocampal area, MeP; medial amygdaloid nucleus,

posterior part, MeA; medial amygdaloid nucleus, anterior part, LOT; nucleus of the lateral olfactory

tract, AA; anterior amygdaloid area.

(B) Representative images of the Me stained with anti-Cart visualized with Alexa 488 and BDA

visualized with Alexa 568-conjugated streptavidin (magenta). Scale bar = 200 μm.

(C) Representative images of the Me injected with the tracer BDA in the MPOA. Parasagittal

sections are shown from lateral to medial. Fixed sections were visualized with Alexa 568-conjugated

streptavidin (magenta). In parallel, fixed sectioned were immunostained with anti-NPI and

visualized with Alexa 488 (green). Main targeted area as the cMPOA (right, lateral 0.5 mm) was

represented as the area of yellow line by reference to our previous study[11]. Scale bar = 200 μm.

(D) left, summary of experimental procedure. middle, RNA-seq identification of candidate gene

expressed in the cMPOA neurons. Heatmap showing a *z*-scored TPM of selected genes for neuronal

cluster-mediated social behavior in the MPOA[21]. n = 4 (virgin), 4 (FGE), and 4 (paternal). right,

Social experiences of virgin, FGE, and paternal group mice.

(E) Representative images of the MPOA. *Galanin* mRNA Alexa 647 -conjugated hairpin DNA (#S23)

(magenta), streptavidin conjugated with FITC (green) and DAPI (blue). (e1-e3) magnified images

corresponding area in (e), Scale bar = 200 μm for (e, left) and 20 μm for (e1-e3)

(F) Representative images of cMPOA neurons after whole-cell patch-clamp recording. Biocytin was

infused from recording pipette. Fixed sections visualized with Alexa 488-conjugated streptavidin

(green) and *in situ* HCR for *Galanin* mRNA Alexa 647 -conjugated hairpin DNA (#S23) (magenta).

Scale bar = 20 μm.

(G) Ratio of *Galanin* positive cMPOA neurons targeted for whole-cell patch-clamp recording.

(H) Fixed sections were immunostained with anti-NPI visualized with a secondary antibody

conjugated to Alexa 488 (green) and immunostained with anti-ERα visualized with a secondary

antibody conjugated to Alexa 568 (magenta). Scale bar = 300 μm.

(I) Fixed sections were visualized with *in situ* HCR for *Galanin* mRNA Alexa 647 -conjugated hairpin

DNA (#S23) (magenta), immunostained with anti-ERα visualized with a secondary antibody

conjugated to Alexa 488 (green), and DAPI (blue). Scale bar = 200 μm. f, fornix, ac, anterior

commissure.

(J) Representative images of cMPOA neurons after whole-cell patch-clamp recording. Biocytin was

infused from recording pipette. Fixed sections visualized with Alexa 568-conjugated streptavidin

(magenta) and immunostained with anti-ERα and secondary antibody conjugated to Alexa 488

(green). Scale bar = 20 μm.

(K) Ratio of ERα positive cMPOA neurons targeted for whole-cell patch-clamp recording.

(L) Schematic image of the recording and stimulating electrodes on a parasagittal section including

the cMPOA. BSTv; bed nucleus of the stria terminalis, ventral part,

(M) Input−output curves of stimulus-evoked IPSPs from the *Galanin* positive cMPOA of virgin mice

(n = 12 cells; 7 animals) and FGE mice (n = 7 cells; 5 animals). Values are presented as the mean ±

standard error. (bottom) Representative traces used to construct input−output curves (40, 80, 120,

160, and 200 μA stimuli). Two-way RM ANOVA, $F_{(1, 17)} = 4.964$, $P = 0.0397$ followed by Holm-

Sidak post hoc test, $*P < 0.05$, $**P < 0.01$ vs. virgin group.

(N) Input−output curves of stimulus-evoked IPSPs from the cMPOA of virgin mice (n = 11 cells; 4

animals) and paternal mice (n = 11 cells; 4 animals). Values are presented as the mean ± standard

error. (bottom) Representative traces used to construct input−output curves (25, 75, 125, 175 and

225 μA stimuli, performed in a different setup and at different stimulation intensity ranges from

Figure 1E, 1G and 1H). Two-way RM ANOVA, $F_{(1, 20)} = 10.72$, $P = 0.0038$ followed by Holm-Sidak

post hoc test, $***P < 0.001$, $****P < 0.0001$ vs. virgin group.

(O) Input−output curves of stimulus-evoked IPSPs from paternal mice recorded without (n = 7 cells;

2 animals) or with GDPβS (1 mM) in the pipette solution (n = 16 cells; 4 animals). Values are

presented as the mean ± standard error. (bottom) Representative traces used to construct

input—output curves (40, 80, 120, 160, and 200 μA stimuli). Two-way RM ANOVA, F (1, 21) = 6.212,

P = 0.0211 followed by Holm-Sidak post hoc test, *P < 0.05, **P < 0.01 vs. paternal group without

GDPβS.

(P) Input—output curves of stimulus-evoked EPSPs from the cMPOA of virgin mice (n = 21 cells; 3

animals) and paternal mice (n = 14 cells; 3 animals). Values are presented as mean ± standard error.

(bottom) Representative traces used to construct input—output curves (40, 80, 120, 160, and 200 μA

stimuli). Two-way RM ANOVA, F (1, 33) = 0.7974, P > 0.05.

Figure 2 Cre-positive neuron in the medial amygdala of Cartpt-Cre mice project to cMPOA neuron

(A) Representative images of the MePD in Cartpt-Cre × Ai9 reporter mice. Among MePD neurons

immunostained with anti-NeuN antibody visualized with a secondary antibody conjugated to Alexa

647 (blue), 21.2% Cartpt neurons (magenta) were tdTomato positive (1 male, 1 female). Scale bar =

300 μm. opt; optic tract, LV; lateral ventricle, AHi; amygdalohippocampal area, MePD; posterior-

dorsal medial amygdala, MePV; posterior-ventral medial amygdala

(B) AAV5-EF1α-DIO-eYFP or AAV5-EF1α-DIO-ChR2-eYFP was injected into the Me of Cartpt-Cre

mice. After the expression of eYFP with/without ChR2, whole-cell patch clamp recordings were

performed from cMPOA neuron in parasagittal sections of virgin and FGE mouse brain.

(C, D) Representative images of eYFP emission from Me (C) and MPOA (D). Me^{Cartpt} neurons sent

projection fibers into the MPOA. Scale bar = 200 μm.

(E) (left)Schematic image of the recording electrodes on a parasagittal section including the cMPOA.

Input fiber from Me^{Cartpt} expressed ChR2 were stimulated by blue light application. BSTv; bed

nucleus of the stria terminalis, ventral part, (right) Diagram of dominant responses of IPSC and

EPSC with opposite current direction as that of GABAergic and glutamatergic ionotropic receptors.

(F) Representative traces of blue light-evoked IPSCs and EPSCs in IPSC- and EPSC-dominant

cMPOA neurons. When Me^{Cartpt} neurons were injected with control AAV virus, no responses to

optical activation were observed in cMPOA neurons.

(G) Relative frequencies of excitatory- and inhibitory-dominant cMPOA neurons in virgin mice (n =

12 cells; 4 animals), FGE mice (n = 14 cells; 3 animals), and FGE mice with GDPβS (1 mM) in the

pipette solution (n = 9 cells; 2 animals). A significant difference in the ratio of excitatory- to

inhibitory-dominant cMPOA neurons between virgin and FGE group mice was observed (Fisher's

exact probability test, **P = 0.0048). A significant difference in the ratio of excitatory- and

inhibitory-dominant cMPOA neurons between FGE mice and FGE mice with GDPβS (1 mM) in the

pipette solution (FGE + GDPβS group) was also observed (Fisher's exact probability test, #P =

0.0131).

Figure 3 The modulation of Me inputs into the cMPOA

(A) AAV10-EF1α-DIO-vLWO-eGFP or AAV10-EF1α-DIO-eYFP were injected into the Me of Cartpt-

Cre mice and whole-cell recordings were obtained from fluorescent-labeled Me neurons.

(B) Representative resting membrane potential traces recorded from a Me^{Cartpt} neuron expressing

vLWO. Green light was applied for 20 s (green bar).

(C) Effects of green light application on the resting membrane potential in the Me^{Cartpt} neurons

expressing vLWO. The data are shown as the averaged potentials between −6 s and −1 s before the

onset of green light application (pre-green light) and between 5 s and 10 s after the onset of green

light application (green light). Lines represent the data obtained from individual neurons (n = 5),

and bars represent the averaged data obtained from five neurons. **P = 0.0017, two-tailed paired t-

test (t = 7.466, df = 4), compared with pre-green light application.

(D) Representative resting membrane potential traces recorded from a Me^{Cartpt} neuron expressing

control eYFP. Green light was applied for 20 s (green bar). Five animals were excluded due to

exclusion criteria.

(E) Effects of green light application on the resting membrane potential in the Me^{Cartpt} neurons

expressing control eYFP. Lines represent the data obtained from individual neurons (n = 10), and

bars represent the averaged data obtained from nine neurons. P = 0.6308, two-tailed paired t-test

(t=0.4975 df=9), compared with pre-green light application.

(F) AAV5-DIO-ChR2(H134R)-mCherry and AAV10-EF1α-DIO-vLWO-eGFP-5HT1A was injected into

the Me of Cartpt-Cre mice and whole-cell recordings were obtained from cMPOA neurons.

(G) Representative images of the Me expressing eGFP and mCherry. Scale bar = 200 μm.

(H) Representative traces of blue light-evoked IPSCs in the cMPOA neurons. Red bar indicates red

light application (2 s).

(I) Effects of red-light application on the blue light-evoked postsynaptic currents in the cMPOA

neurons. Lines represent the data obtained from individual neurons (n = 4), and bars represent the

averaged data obtained. ***P = 0.0007, two-tailed paired t-test (t=14.25 df=3), compared with pre-

red light application.

(J) AAV10-EF1α-DIO-vLWO-eGFP was injected into the Me of Cartpt-Cre mice and whole-cell

recordings were obtained from cMPOA neurons.

(K) Representative trace of green light-induced modulation of sIPSCs in the cMPOA neurons. Data

were obtained in the presence of CNQX (20 μM) and MK-801 (20 μM). Green light was applied for

20 s.

(L) Effects of green light application on the frequency of sIPSC in the cMPOA neurons. Lines

represent the data obtained from individual neurons (n = 25), and bars represent the averaged data.

Repeated measures ANOVA, F (1.468, 35.23) = 3.962, P = 0.0396, followed by Dunnett's multiple

comparisons post hoc test *P = 0.0332 compared with pre-green light application. One data showed

large sIPSC frequency. It was judged as an outlier in Grubbs' test. Without this data point, the same

repeated measure ANOVA analysis resulted in F (1.733, 39.87) = 4.149, P=0.0279, indicating that

the conclusion remains when leaving out this outlier.

(M) Representative trace of sIPSCs in the cMPOA neurons in the presence of TTX (1 μM).

(N) Application of TTX blocked the green light induced changes of IPSCs frequency. Lines represent

the data obtained from individual neurons (n = 12), and bars represent the averaged data. Repeated

measures ANOVA, F (1.236, 13.60) = 0.2769, P = 0.6561.

Figure 4 Optogenetic manipulation of the Me^{Cartpt}–cMPOA input suppressed infanticidal behavior of

virgin male mice

(A, B) Representative images of c-Fos induction in the Me following pup exposure (A) and in the

control Me (B). Fixed sections were stained with anti-c-Fos and visualized with Alexa 647 (magenta).

eYFP was expressed by injection of AAV10-EF1α-DIO-eYFP. Scale bar = 200 μm for 10× images.

Scale bar = 50 μm for 60× images.

(C) Percentage of c-Fos-positive Me^{Cartpt} neurons out of all eYFP-positive Me^{Cartpt} neurons following

pup exposure. A pup kept in a metal net to avoid biting attacks was placed in the test cage for 30

min (n = 3). An empty metal net was added as a control (n = 3). ****P < 0.0001, two-tailed unpaired

t-test (t = 23.41, df = 4).

(D) AAV10-EF1α-DIO-vLWO-eGFP-5HT1A or AAV10-EF1α-DIO-eYFP was injected into the Me of

Cartpt-Cre mice. Five weeks later, glass fibers were implanted above the cMPOA for pathway

stimulation. During the behavioral test, 532 nm green light was applied following a repeated

protocol of 20 s on and 10 s off.

(E) Latencies to first sniffing were compared between vLWO and control group mice. Two-way RM

ANOVA, F (1, 16) = 1.642, P = 0.2183.

(F, G) Relative frequencies of paternal/infanticidal behaviors toward pups during manipulation of

the Me^{Cartpt}-to-cMPOA input by vLWO activation. Significant difference in the ratio showing

infanticidal behavior between mice expressing vLWO-eGFP (n = 8, F) or eYFP only (n = 10, G) *P =

0.0229, **P = 0.0065, Fisher's exact probability test. Eight animals of vLWO group were excluded

due to exclusion criteria.

Figure 5

Optogenetic inhibition of the Me−cMPOA input in vGAT-IRES-Cre mice suppressed infanticidal

behavior of virgin male mice

(A) AAV10- EF1α-DIO-vLWO-eGFP-5HT1A or AAV10-EF1α-DIO-eYFP was injected into the Me of

vGAT-IRES-Cre mice. Five weeks later, glass fibers were implanted above the cMPOA. During the

behavioral test, 532 nm green light was repeatedly applied at 20 s on and 10 s off.

(B) Representative images of eYFP emission from the MPOA of vGAT-IRES-Cre mice (green).

Sections including the Me were immunostained with anti-NPI and visualized with a secondary

antibody conjugated to Alexa 594 (magenta).

(C) Representative images of eYFP expressed in the Me of vGAT-IRES-Cre mice (green). Sections

were immunostained with anti-Cart and visualized with a secondary antibody conjugated to Alexa

594 (magenta). All scale bar = 200 μm.

(D, E) Relative frequencies of paternal/infanticidal behaviors during vLWO activation at the Me^{vGAT}

input into cMPOA. Significant difference in the ratio of mice showing infanticide between vLWO (n

= 7, D) and control (n = 10, E) groups (Fisher's exact probability test, *P = 0.0345).

Figure 6 Inhibitory synaptic inputs into the BSTrh were potentiated in paternal group mice

(A) Whole-cell patch clamp recordings were performed on BSTrh neurons (Rec). To observe the

evoked synaptic responses, electrical stimulation was delivered through a stimulating electrode

(Stim). rh, rhomboid nucleus; pr, principal nucleus; LV, lateral ventricle; ic, internal capsule; f,

fornix; D3V, dorsal third ventricle; GP, globus pallidus; ac, anterior commissure; ST, stria terminalis

(B) Input−output curves of stimulus-evoked IPSPs from virgin mice (n = 16 cells; 10 animals),

paternal mice (n = 17 cells; 11 animals), and FGE mice (n = 10 cells; 5 animals). Values are

presented as the mean ± standard error. (inset) Representative traces used to construct

input−output curves (50, 100, 150, and 200 µA stimuli). Two-way RM ANOVA, $F_{(2, 40)}$ = 5.416, P =

0.0083, followed by Holm-Sidak post hoc test, *P < 0.05, **P < 0.01 vs. virgin group.

(C) Input−output curves of stimulus-evoked EPSPs from virgin mice (n = 9 cells; 6 animals),

paternal mice (n = 15 cells; 6 animals), and FGE mice (n = 6 cells; 3 animals). Values are presented

as the mean ± standard error. (inset) Representative traces used to construct input−output curves

(25 and 100 µA stimuli). Two-way RM ANOVA, $F_{(2, 27)}$ = 0.4838, P > 0.05.

(D) Photographs of the MPOA from sham-operated and NMDA-lesioned mice. Sections were

immunostained with anti-NeuN and visualized with DAB. Arrowhead indicates the ventral edge of

lesioned area. Scale bar = 500 µm.

(E) Input–output curves of stimulus-evoked IPSPs from paternal sham-operated mice (sham; n = 26

from 7 mice, NMDA; n = 15 from 6 mice). Values are presented as the mean ± standard error. (inset)

Representative traces used to construct input–output curves (25, 75, 125, and 175 µA stimuli). Two-

way RM ANOVA, $F_{(1, 29)} = 10.30$, $P = 0.0032$ followed by Holm-Sidak post hoc test, $*P < 0.05$, $**P <$

0.01, $***P < 0.001$ vs. virgin group.

(F) Input–output curves of stimulus-evoked IPSPs from sham-operated virgin mice (sham; n = 7

cells; 4 animals) and cMPOA-lesioned virgin mice (NMDA; n = 9 cells; 4 animals). Values are

presented as the mean ± standard error. (inset) Representative traces used to construct

input–output curves (25, 75, 125, and 175 µA stimuli). Two-way RM ANOVA, $F_{(1, 14)} = 0.9177$, $P >$

0.05.

(G) Input–output curves of stimulus-evoked IPSPs from virgin mice (n = 10 cells; 4 animals) and

paternal mice (n = 9 cells; 4 animals) recorded with GDPßS (1 mM) in the pipette solution. Values

are presented as mean ± standard error. (inset) Representative traces used to construct

input–output curves (50, 100, 150, and 200 µA stimuli). Two-way RM ANOVA, $F_{(1, 17)} = 0.1907$, P

> 0.05.

(H) Input–output curves of stimulus-evoked EPSPs from virgin mice (n = 9 cells; 5 animals) and

paternal mice (n = 10 cells; 4 animals) recorded with GDPßS (1 mM) in the pipette solution. Values

are presented as mean ± standard error. (inset) Representative traces used to construct

input−output curves (25, 75 and 125 μA stimuli). Two-way RM ANOVA, F (1, 17) = 0.4811, P > 0.05.

(I) (top) Paired-pulse ratio of eEPSCs (inter-stimulation interval = 50 ms) from virgin mice (n = 6

cells; 5 animals) and paternal mice (n = 7 cells; 6 animals). Two-tailed unpaired t-test: P > 0.05.

(bottom) Paired-pulse ratio of eIPSCs (inter-stimulation interval = 100 ms) from virgin mice (n = 10

cells; 9 animals) and paternal mice (n = 15 cells; 10 animals). Two-tailed unpaired t-test, P > 0.05.

References

1.   Elwood, R.W. (1977). Changes in Responses of Male and Female Gerbils (Meriones-
Unguiculatus) Towards Test Pups during Pregnancy of Female. Anim Behav *25*, 46−51. Doi
10.1016/0003-3472(77)90066-5.

2.   Hausfater, G., and Hrdy, S.B. (1984). Infanticide: Comparative and Evolutionary
Perspectives (Routledge).

3.   Hrdy, S.B. (1974). Male-Male Competition and Infanticide among Langurs (Presbytis-
Entellus) of Abu, Rajasthan. Folia Primatol *22*, 19−58.

4.   van Schaik, C.P., and Janson, C.P. (2000). Infanticide by Males and its Implications
(Cambridge University Press).

5.   vom Saal, F.S., and Howard, L.S. (1982). The regulation of infanticide and parental
behavior: implications for reproductive success in male mice. Science *215*, 1270-1272.

6.   Palombit, R.A. (2015). Infanticide as Sexual Conflict: Coevolution of Male Strategies and
Female Counterstrategies. Csh Perspect Biol *7*. 10.1101/cshperspect.a017640.

7.   vom Saal, F.S. (1985). Time-contingent change in infanticide and parental behavior induced
by ejaculation in male mice. Physiol Behav *34*, 7-15.

8.   Tachikawa, K.S., Yoshihara, Y., and Kuroda, K.O. (2013). Behavioral transition from attack
to parenting in male mice: a crucial role of the vomeronasal system. The Journal of
neuroscience : the official journal of the Society for Neuroscience *33*, 5120-5126.
10.1523/JNEUROSCI.2364-12.2013.

9.   Numan, M. (1974). Medial preoptic area and maternal behavior in the female rat. J Comp
Physiol Psychol *87*, 746-759.

10.   Tsuneoka, Y., Maruyama, T., Yoshida, S., Nishimori, K., Kato, T., Numan, M., and Kuroda,
K.O. (2013). Functional, anatomical, and neurochemical differentiation of medial preoptic
area subregions in relation to maternal behavior in the mouse. The Journal of comparative

neurology *521*, 1633-1663. 10.1002/cne.23251.

11. Tsuneoka, Y., Tokita, K., Yoshihara, C., Amano, T., Esposito, G., Huang, A.J., Yu, L.M.,
Odaka, Y., Shinozuka, K., McHugh, T.J., and Kuroda, K.O. (2015). Distinct preoptic-BST
nuclei dissociate paternal and infanticidal behavior in mice. The EMBO journal *34*, 2652-
2670. 10.15252/embj.201591942.

12. Wu, Z., Autry, A.E., Bergan, J.F., Watabe-Uchida, M., and Dulac, C.G. (2014). Galanin
neurons in the medial preoptic area govern parental behaviour. Nature *509*, 325-330.
10.1038/nature13307.

13. Keshavarzi, S., Sullivan, R.K., Ianno, D.J., and Sah, P. (2014). Functional properties and
projections of neurons in the medial amygdala. The Journal of neuroscience : the official
journal of the Society for Neuroscience *34*, 8699-8715. 10.1523/JNEUROSCI.1176-14.2014.

14. Kohl, J., Babayan, B.M., Rubinstein, N.D., Autry, A.E., Marin-Rodriguez, B., Kapoor, V.,
Miyamishi, K., Zweifel, L.S., Luo, L., Uchida, N., and Dulac, C. (2018). Functional
circuit architecture underlying parental behaviour. Nature *556*, 326-331. 10.1038/s41586-
018-0027-0.

15. Simerly, R.B., and Swanson, L.W. (1986). The organization of neural inputs to the medial
preoptic nucleus of the rat. The Journal of comparative neurology *246*, 312-342.
10.1002/cne.902460304.

16. Mei, L., Yan, R., Yin, L., Sullivan, R.M., and Lin, D. (2023). Antagonistic circuits
mediating infanticide and maternal care in female mice. Nature. 10.1038/s41586-023-06147-9.

17. Chen, P.B., Hu, R.K., Wu, Y.E., Pan, L., Huang, S., Micevych, P.E., and Hong, W. (2019).
Sexually Dimorphic Control of Parenting Behavior by the Medial Amygdala. Cell *176*, 1206-
1221.e1218. 10.1016/j.cell.2019.01.024.

18. Broberger, C. (1999). Hypothalamic cocaine- and amphetamine-regulated transcript (CART)
neurons: histochemical relationship to thyrotropin-releasing hormone, melanin-
concentrating hormone, orexin/hypocretin and neuropeptide Y. Brain Res *848*, 101-113.

19. Wu, Y.E., Pan, L., Zuo, Y., Li, X., and Hong, W. (2017). Detecting Activated Cell
Populations Using Single-Cell RNA-Seq. Neuron *96*, 313-329 e316.
10.1016/j.neuron.2017.09.026.

20. Sato, K., Hamasaki, Y., Fukui, K., Ito, K., Miyamichi, K., Minami, M., and Amano, T.
(2020). Amygdalohippocampal Area Neurons That Project to the Preoptic Area Mediate Infant-
Directed Attack in Male Mice. The Journal of neuroscience : the official journal of the
Society for Neuroscience *40*, 3981-3994. 10.1523/JNEUROSCI.0438-19.2020.

21. Moffitt, J.R., Bambah-Mukku, D., Eichhorn, S.W., Vaughn, E., Shekhar, K., Perez, J.D.,
Rubinstein, N.D., Hao, J., Regev, A., Dulac, C., and Zhuang, X. (2018). Molecular, spatial,
and functional single-cell profiling of the hypothalamic preoptic region. Science *362*.
10.1126/science.aau5324.

22. Fang, Y.Y., Yamaguchi, T., Song, S.C., Tritsch, N.X., and Lin, D. (2018). A Hypothalamic

882   Midbrain Pathway Essential for Driving Maternal Behaviors. Neuron *98*, 192-207 e110.
883   10.1016/j.neuron.2018.02.019.

884   23.   Wei, Y.C., Wang, S.R., Jiao, Z.L., Zhang, W., Lin, J.K., Li, X.Y., Li, S.S., Zhang, X.,
885   and Xu, X.H. (2018). Medial preoptic area in mice is capable of mediating sexually
886   dimorphic behaviors regardless of gender. Nat Commun *9*, 279. 10.1038/s41467-017-02648-0.

887   24.   Masseck, O.A., Spoida, K., Dalkara, D., Maejima, T., Rubelowski, J.M., Wallhorn, L.,
888   Deneris, E.S., and Herlitze, S. (2014). Vertebrate cone opsins enable sustained and highly
889   sensitive rapid control of Gi/o signaling in anxiety circuitry. Neuron *81*, 1263-1273.
890   10.1016/j.neuron.2014.01.041.

891   25.   Soya, S., Takahashi, T.M., McHugh, T.J., Maejima, T., Herlitze, S., Abe, M., Sakimura, K.,
892   and Sakurai, T. (2017). Orexin modulates behavioral fear expression through the locus
893   coeruleus. Nat Commun *8*, 1606. 10.1038/s41467-017-01782-z.

894   26.   Amano, T., Shindo, S., Yoshihara, C., Tsuneoka, Y., Uki, H., Minami, M., and Kuroda, K.O.
895   (2017). Development-dependent behavioral change toward pups and synaptic transmission in
896   the rhomboid nucleus of the bed nucleus of the stria terminalis. Behav Brain Res *325*, 131-
897   137. 10.1016/j.bbr.2016.10.029.

898   27.   Elwood, R.W. (1985). Inhibition of Infanticide and Onset of Paternal Care in Male-Mice
899   (Mus-Musculus). J Comp Psychol *99*, 457-467. Doi 10.1037//0735-7036.99.4.457.

900   28.   Inada, K., Hagihara, M., Tsujimoto, K., Abe, T., Konno, A., Hirai, H., Kiyonari, H., and
901   Miyamichi, K. (2022). Plasticity of neural connections underlying oxytocin-mediated
902   parental behaviors of male mice. Neuron. 10.1016/j.neuron.2022.03.033.

903   29.   Hong, W., Kim, D.W., and Anderson, D.J. (2014). Antagonistic control of social versus
904   repetitive self-grooming behaviors by separable amygdala neuronal subsets. Cell *158*, 1348-
905   1361. 10.1016/j.cell.2014.07.049.

906   30.   Kondo, Y. (1992). Lesions of the medial amygdala produce severe impairment of copulatory
907   behavior in sexually inexperienced male rats. Physiol Behav *51*, 939-943.

908   31.   Li, Y., Mathis, A., Grewe, B.F., Osterhout, J.A., Ahanonu, B., Schnitzer, M.J., Murthy,
909   V.N., and Dulac, C. (2017). Neuronal Representation of Social Information in the Medial
910   Amygdala of Awake Behaving Mice. Cell *171*, 1176-1190 e1117. 10.1016/j.cell.2017.10.015.

911   32.   Sano, K., Nakata, M., Musatov, S., Morishita, M., Sakamoto, T., Tsukahara, S., and Ogawa,
912   S. (2016). Pubertal activation of estrogen receptor alpha in the medial amygdala is
913   essential for the full expression of male social behavior in mice. Proc Natl Acad Sci U S
914   A *113*, 7632-7637. 10.1073/pnas.1524907113.

915   33.   Sano, K., Tsuda, M.C., Musatov, S., Sakamoto, T., and Ogawa, S. (2013). Differential
916   effects of site-specific knockdown of estrogen receptor alpha in the medial amygdala,
917   medial pre-optic area, and ventromedial nucleus of the hypothalamus on sexual and
918   aggressive behavior of male mice. Eur J Neurosci *37*, 1308-1319. 10.1111/ejn.12131.

919   34.   Dominguez, J., Riolo, J.V., Xu, Z., and Hull, E.M. (2001). Regulation by the medial

920   amygdala of copulation and medial preoptic dopamine release. The Journal of neuroscience :
921   the official journal of the Society for Neuroscience *21*, 349-355.

35. Kondo, Y., and Arai, Y. (1995). Functional association between the medial amygdala and the medial preoptic area in regulation of mating behavior in the male rat. Physiol Behav *57*, 69-73.

36. Choi, G.B., Dong, H.W., Murphy, A.J., Valenzuela, D.M., Yancopoulos, G.D., Swanson, L.W., and Anderson, D.J. (2005). Lhx6 delineates a pathway mediating innate reproductive behaviors from the amygdala to the hypothalamus. Neuron *46*, 647-660. 10.1016/j.neuron.2005.04.011.

37. Pardo-Bellver, C., Cadiz-Moretti, B., Novejarque, A., Martinez-Garcia, F., and Lanuza, E. (2012). Differential efferent projections of the anterior, posteroventral, and posterodorsal subdivisions of the medial amygdala in mice. Front Neuroanat *6*, 33. 10.3389/fnana.2012.00033.

38. Morris, J.A., Jordan, C.L., and Breedlove, S.M. (2008). Sexual dimorphism in neuronal number of the posterodorsal medial amygdala is independent of circulating androgens and regional volume in adult rats. The Journal of comparative neurology *506*, 851-859. 10.1002/cne.21536.

39. Unger, E.K., Burke, K.J., Jr., Yang, C.F., Bender, K.J., Fuller, P.M., and Shah, N.M. (2015). Medial amygdalar aromatase neurons regulate aggression in both sexes. Cell Rep *10*, 453-462. 10.1016/j.celrep.2014.12.040.

40. Rogge, G., Jones, D., Hubert, G.W., Lin, Y., and Kuhar, M.J. (2008). CART peptides: regulators of body weight, reward and other functions. Nat Rev Neurosci *9*, 747-758. 10.1038/nrn2493.

41. Allen, W.E., DeNardo, L.A., Chen, M.Z., Liu, C.D., Loh, K.M., Fenno, L.E., Ramakrishnan, C., Deisseroth, K., and Luo, L. (2017). Thirst-associated preoptic neurons encode an aversive motivational drive. Science *357*, 1149-1155. 10.1126/science.aan6747.

42. Chung, S., Weber, F., Zhong, P., Tan, C.L., Nguyen, T.N., Beier, K.T., Hormann, N., Chang, W.C., Zhang, Z., Do, J.P., et al. (2017). Identification of preoptic sleep neurons using retrograde labelling and gene profiling. Nature *545*, 477-481. 10.1038/nature22350.

43. McHenry, J.A., Otis, J.M., Rossi, M.A., Robinson, J.E., Kosyk, O., Miller, N.W., McElligott, Z.A., Budygin, E.A., Rubinow, D.R., and Stuber, G.D. (2017). Hormonal gain control of a medial preoptic area social reward circuit. Nat Neurosci *20*, 449-458. 10.1038/nn.4487.

44. Tan, C.L., Cooke, E.K., Leib, D.E., Lin, Y.C., Daly, G.E., Zimmerman, C.A., and Knight, Z.A. (2016). Warm-Sensitive Neurons that Control Body Temperature. Cell *167*, 47-59 e15. 10.1016/j.cell.2016.08.028.

45. Fumero, B., Fernandez-Vera, J.R., Gonzalez-Mora, J.L., and Mas, M. (1994). Changes in monoamine turnover in forebrain areas associated with masculine sexual behavior: a

microdialysis study. Brain Res *662*, 233-239.

46. Nutsch, V.L., Will, R.G., Robison, C.L., Martz, J.R., Tobiansky, D.J., and Dominguez, J.M.
(2016). Colocalization of Mating-Induced Fos and D2-Like Dopamine Receptors in the Medial
Preoptic Area: Influence of Sexual Experience. Front Behav Neurosci *10*, 75.
10.3389/fnbeh.2016.00075.

47. Ren, W.J., Centeno, M.V., Berger, S., Wu, Y., Na, X.D., Liu, X.G., Kondapalli, J.,
Apkarian, A.V., Martina, M., and Surmeier, D.J. (2016). The indirect pathway of the
nucleus accumbens shell amplifies neuropathic pain. Nature Neuroscience *19*, 220-+.
10.1038/nn.4199.

48. Thompson, D., Martini, L., and Whistler, J.L. (2010). Altered ratio of D1 and D2 dopamine
receptors in mouse striatum is associated with behavioral sensitization to cocaine. PLoS
One *5*, e11038. 10.1371/journal.pone.0011038.

49. Jacob, T.C., Moss, S.J., and Jurd, R. (2008). GABA(A) receptor trafficking and its role in
the dynamic modulation of neuronal inhibition. Nat Rev Neurosci *9*, 331-343.
10.1038/nrn2370.

50. Seelke, A.M.H., Bond, J.M., Simmons, T.C., Joshi, N., Settles, M.L., Stolzenberg, D.,
Rhemtulla, M., and Bales, K.L. (2018). Fatherhood alters gene expression within the MPOA.
Environ Epigenet *4*, dvy026. 10.1093/eep/dvy026.

51. Tsuneoka, Y., Yoshida, S., Takase, K., Oda, S., Kuroda, M., and Funato, H. (2017).
Neurotransmitters and neuropeptides in gonadal steroid receptor-expressing cells in medial
preoptic area subregions of the male mouse. Sci Rep *7*, 9809. 10.1038/s41598-017-10213-4.

52. Kawai, S., Takagi, Y., Kaneko, S., and Kurosawa, T. (2011). Effect of three types of mixed
anesthetic agents alternate to ketamine in mice. Exp Anim *60*, 481-487.

53. Tsuneoka, Y., Atsumi, Y., Makanae, A., Yashiro, M., and Funato, H. (2022). Fluorescence
quenching by high-power LEDs for highly sensitive fluorescence in situ hybridization.
Front Mol Neurosci *15*, 976349. 10.3389/fnmol.2022.976349.

54. Tsuneoka, Y., and Funato, H. (2020). Modified in situ Hybridization Chain Reaction Using
Short Hairpin DNAs. Front Mol Neurosci *13*, 75. 10.3389/fnmol.2020.00075.

55. Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R.
(2014). Full-length RNA-seq from single cells using Smart-seq2. Nat Protoc *9*, 171-181.
10.1038/nprot.2014.006.

56. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Graphical Abstract