Graded changes in local functional connectivity of the brain in young people with depression

*Alec J. Jamieson[1], Christopher G. Davey[2], Jesus Pujol[3], Laura Blanco-Hinojo[3], & Ben J. Harrison[1]

[1]Melbourne Neuropsychiatry Centre, Department of Psychiatry, The University of Melbourne, Australia

[2]Department of Psychiatry, The University of Melbourne, Australia

[3]MRI Research Unit, Department of Radiology, Hospital del Mar, Barcelona, Spain

Corresponding author: Dr. Alec Jamieson. Melbourne Neuropsychiatry Centre, Department of Psychiatry, The University of Melbourne. Level 3, 161 Barry St. Carlton, VIC 3053. Phone: (+61) 449 501 556 | Email: alec.jamieson@unimelb.edu.au

Abstract

Major depressive disorder (MDD) is marked by significant changes to the coupling of spontaneous neural activity within various brain regions. However, many methods for assessing this local connectivity use fixed or arbitrary neighborhood sizes, resulting in a decreased capacity to capture smooth changes to the spatial gradient of local correlations. A newly developed method sensitive to classical anatomo-functional boundaries, Iso-Distant Average Correlation (IDAC), was therefore used to examine depression associated alterations to the local functional connectivity of the brain. One-hundred and forty-five adolescents and young adults with MDD and 95 healthy controls underwent a resting-state fMRI scan. Whole-brain functional connectivity maps of intracortical neural activity within iso-distant local areas (5-10mm, 15-20mm, and 25-30mm) were generated to characterize local fMRI signal similarities. Across all spatial distances, MDD participants demonstrated greater local functional connectivity of the bilateral posterior hippocampus, retrosplenial cortex, dorsal insula, fusiform gyrus, and supplementary motor area. Additionally, in the short and medium range connections there were depression associated alterations in the midcingulate (15-20mm and 25-30mm) and subgenual anterior cingulate (15-20mm). Our study identified increased synchrony of the neural activity in several regions commonly implicated in the neurobiology of depression; however, a subset of identified effects was dependent on the spatial distance under consideration. Longitudinal examination of these effects will clarify whether these differences are also found in other age groups and if this synchrony is additionally altered by continued disease progression.

Introduction

Major depressive disorder (MDD) is a highly prevalent and disabling mental health condition characterized by pervasive symptoms of sadness, hopelessness, and anhedonia [1]. Depression often first arises during adolescence [2, 3], with early onset being a strong predictor of recurrent depression [4, 5] as well as greater reductions in quality of life and wellbeing into adulthood [6]. Early adolescence is marked by increased use of maladaptive emotional regulation strategies which in turn are associated with exacerbated mental health issues [7, 8]. One such strategy, known expressive suppression, involves inhibiting the outward expression of emotional behavior [9], has been specifically associated with depressive symptoms severity [10]. However, there remains limited work examining neurobiological underpinnings of this early stage of the disorder.

Previous neuroimaging research has attempted to uncover the neurobiological underpinnings of MDD by focusing on long-range functional connectivity, examining the temporal correlations of large-scale, spatially distinct brain networks. Within this framework, MDD has been associated with alterations across multiple intrinsic brain networks [11, 12], including the default mode (DMN), salience, and central executive networks [13]. However, there has been increasing interest in assessing local functional connectivity profiles [14, 15], as the functional integration of local neuronal groups appears to represent a key organizing principle in the brains of higher vertebrates [16]. Local functional connectivity measures assess the temporal coherence of the functional magnetic resonance imaging (fMRI) signal between a given voxel and neighboring voxels [14]. Measures of this 'localized synchrony' provide a framework for examining functional disruptions at rest without a priori constraints, thereby allowing for data-driven identification of regional abnormalities across a range of mental health disorders [17]. Moreover, regional variation in homogeneity has been associated with the hierarchical organization of information processing across the brain, and therefore alterations to these measures may serve as key markers of human brain function [14, 18] and a sensitive tool for identifying brain alterations across mental health disorders [19, 20].

Meta-analyses examining regional homogeneity in MDD have revealed inconsistent differences in the directionality of effects and regions implicated [21-23]. Altered local connectivity has been noted across regions including the parahippocampal gyrus, insula, and medial prefrontal cortex [21, 22]. Other local functional connectivity measures, including functional connectivity density and dynamic regional phase synchrony, have illustrated reduced local connectivity in the anterior cingulate cortex, precuneus, hippocampus, thalamus, and insula [24-26]. Conversely, recent work examining first-episode MDD observed

increased regional homogeneity of the hippocampus and insula but reduced local connectivity of the orbitofrontal cortex [23]. It is likely that the specific method used, including how neighborhood boundaries are defined, as well as chronicity of the disorder, contributes to the inconsistencies observed between studies. Despite these inconsistencies, the observed alterations appear to be localized to regions implicated in affective processing and regulation [27, 28]. Of particular note is the implication of insula dysfunction across measures of local and remote functional connectivity [29-32], due to its hypothesized role in the integration of autonomic, emotional, and interoceptive stimuli [33] and potential specificity in delineating MDD from bipolar disorder [29].

      We aimed to investigate local functional connectivity alterations in a large sample of adolescents and young adults with MDD, using the recently introduced framework - IsoDistant Average Correlation (IDAC) [34]. Measures of local functional connectivity often use arbitrary and binarized neighborhood boundaries, thereby losing the capacity to describe the rich smooth spatial gradient of local fMRI correlations [35-37]. IDAC assesses the average temporal correlation of one voxel with all neighboring voxels within different spatial lags, thereby overcoming these limitations [34]. Based on previous findings, we hypothesize that MDD participants would illustrate increased local functional connectivity of the insula, anterior cingulate, and hippocampus and reduced local functional connectivity of the orbitofrontal cortex. Given the developmental nature of our cohort, we additionally investigated whether local connectivity was associated with measures of depressive symptom severity, and emotion processing and regulation.

Methods

One hundred and forty-nine unmedicated, help-seeking MDD participants were recruited as part of the Youth Depression Alleviation-Combined Treatment and -Augmentation trials, for full details see Davey and colleagues [38] and Berk and colleagues [39]. MDD participants were between 15 to 25 years of age and were recruited through specialist mental health clinics in the northern and western suburbs of Melbourne, Australia. These participants had a current diagnosis of MDD, as assessed by the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID) [40]. Depressive symptoms were at least of a moderate level of severity, indicated by a Montgomery-Åsberg Depression Rating Scale (MADRS) score greater than or equal to 20. Exclusion criteria included a lifetime or current diagnosis of a psychotic or bipolar disorder, current treatment with antidepressant medication, or MRI contraindications including pregnancy. One hundred age and sex-matched healthy participants were also

recruited through online advertisements. They had no past mental health disorder diagnoses as assessed through SCID criteria. Due to excessive head motion (three MDD participants, four controls; see below) and incidental findings (one MDD participant, one healthy control), a total of four MDD participants and five controls were excluded from further analysis. This resulted in 95 healthy controls and 145 MDD participants in our final sample. This study and consent process was approved by the Melbourne Health Human Research and Ethics Committee. An informed consent form was provided to, and signed by, all participants in this study. For participants under the age of 18, both participant and parental consent were required.

Image Acquisition

A 3T General Electric Signa Excite system with an eight-channel phased-array head coil was used in combination with ASSET parallel imaging. The functional sequence consisted of a single shot gradient-recalled echo-planar imaging sequence in the steady state (repetition time, 2000 ms; echo time, 35 ms; and pulse angle, 90°) in a 23-cm field-of-view, with a 64 x 64-pixel matrix and a slice thickness of 3.5mm (no gap). Thirty-six interleaved slices were acquired parallel to the anterior-posterior commissure line with a 20° anterior tilt to better cover ventral prefrontal brain regions. The total sequence duration was 8 minutes, corresponding to 240 whole-brain echo-planar imaging volumes. Participants were instructed to keep their eyes closed for the duration of the scan. The first 4 volumes from each run were automatically discarded to allow for signal equilibration. A T1-weighted high-resolution anatomical image was acquired for each participant to assist with functional timeseries co-registration (140 contiguous slices; repetition time, 7.9 ms; echo time, 3 ms; flip angle, 13°; in a 25.6 cm field-of-view, with a 256 x 256-pixel matrix and a slice thickness of 1 mm). To assist with noise reduction and head immobility, all participants used earplugs and had their heads supported with foam-padding inserts.

Image Preprocessing

Imaging data was transferred to a Unix-based platform that ran MATLAB Version 9.3 (The MathWorks Inc., Natick, USA) and Statistical Parametric Mapping (SPM) Version 12 v7487 (Wellcome Trust Centre for Neuroimaging, London, UK). Preprocessing followed previously reported steps for IDAC analysis [41]. In brief, for each subject functional MRI images were slice-time corrected, realigned, and co-registered to their corresponding anatomical image with an affine transformation. Spatial normalization occurred through a back-transformation

process, as such the individual 3D anatomical images were segmented and registered to MNI space so that the resulting deformation fields could be applied to the IDAC maps (see below). Images were then re-sliced to 3x3x3 mm resolution and smoothed by convolving the image with a 4x4x4 mm full width at half maximum Gaussian kernel. Motion fingerprint [42] was used to quantify participant head motion. Participants were excluded if movement exceeded a mean total displacement of 3 mm (~1 native voxel).

IDAC analysis

IDAC maps describe the pattern of correlation decay propagating from each voxel across the brain. This graded change in local functional connectivity provides a more comprehensive characterisation of the brain's functional structure and has been shown to differentiate the human cortex into regions consistent with traditional brain atlases as well as being sensitive to different brain functional states [34]. For a detailed overview of IDAC see Macia and colleagues [34]. Whole-cortex IDAC maps were computed using the mean correlation z-score of each voxel with all neighboring voxels placed at increasingly spaced iso-distant intervals. IDAC maps were calculated for three iso-distant intervals: 5-10 mm, 15-20 mm, and 25-30 mm, and conducted separately for the right and left hemisphere in native space. Covariates included in this analysis were the six rigid body realignment parameters, their first-order derivatives, average white matter, cerebrospinal fluid, and global brain signal. Motion scrubbing using the realignment parameters from preprocessing was additionally conducted to discard motion-affected image volumes [43]. For volumes with greater than 0.2 mm of inter-frame motion, the corresponding volume and those immediately preceding and succeeding were discarded. A discrete cosine transform filter was applied for frequencies outside the .01–0.1 Hz interval.

The resulting maps were then normalized to the Montreal Neurological Institute (MNI) space using a back-transformation process to enable group inference. RGB color overlays were used for displaying the values from the three distances simultaneously. For this study, red represented results from 5-10 mm IDAC maps, green from 15-20 mm, and blue from 25-30 mm. Overlaps of these distances are illustrated through the respective secondary colors.

Statistical Analysis

Between groups analysis were conducted using a 2×3 ANOVA (group [healthy control and MDD] by distance [5–10, 15–20, and 25–30 mm]). Following examination of the Effect of Group F contrast, post hoc t-tests were performed to examine group differences in both directions for each of the three IDAC maps. This resulted in six primary contrasts of interest. Accordingly, we adjusted the false-discovery error rate (FDR) corrected threshold to .008 (0.05/6). Thus all results displayed were estimated with a whole-brain, FDR corrected threshold of $p < .008$, KE ≥ 10 voxels. Secondary level covariates included age, gender, and handedness.

To investigate whether these between group differences were also associated with depressive symptom severity we estimated voxel-wise correlations between MADRS scores and whole-brain functional connectivity for the 5–10, 15–20, and 25–30 mm IDAC distances. Given the regions implicated in the between groups comparison, we then examined whether local functional connectivity changes were associated with the scores from the Emotional Regulation Questionnaire (ERQ)[44]. The instrument assesses the two means by which people attempt to regulate their emotions: by expressive suppression ('Suppression') and by cognitive reappraisal ('Reappraisal'). The scores on the subscales were normally distributed across groups (Supplementary Figure S1), and analyses were conducted across both groups and in the MDD group alone. Analyses of between-group differences for clinical and demographic characteristics were calculated in SPSS version 27 (IBMCorp., Armonk, NY). Comparisons were adjusted for multiple comparisons using the Holm-Bonferroni correction [45] to determine significance ($p < .05$).

Results

Group Differences Between Demographic and Clinical Characteristics

Demographic and clinical characteristics of the sample are reported in Table 1. Healthy controls and MDD patients differed significantly on MADRS symptoms ($t(233.5) = -58.80$, $p < .001$), ERQ Suppression ($t(241) = -5.94$, $p < .001$) and Reappraisal scores ($t(216.6) = 10.60$, $p < .001$) as well as in IQ ($t(242) = 4.03$, $p < .001$).

Functional Structure of the Cerebral Cortex

To ensure that our results are consistent with previously illustrated anatomofunctional boundaries [34, 41, 46], whole-brain maps were generated across the three distances for the healthy controls (Figure 1). Consistent with previous effects, the visual association cortex

demonstrated high connectivity across all local distance ranges. The bilateral inferior parietal lobules, including the angular and supramarginal gyri, were connected at short and medium distances. Morevoer

Group Differences in the Functional Structure of the Cerebral Cortex

The Effect of Group contrast demonstrated that MDD participants and healthy controls had significant differences across the three connectivity distances (Figure 2, Supplementary Table S1). These effects were most predominately observed in the bilateral posterior hippocampus extending to the retrosplenial cortex and ventral posterior cingulate as well as to the lingual and fusiform gyri. Additional large effects were observed in the dorsal mid to posterior insular cortices and right dorsal premotor cortex. Subsequent t-test for each distance 5–10, 15–20, and 25–30 mm illustrated that these results were all driven by greater local connectivity in MDD participants compared to controls (Figure 3). As such, across all distances MDD participants demonstrated greater local functional connectivity of the bilateral extended posterior hippocampus, fusiform and lingual gyri, dorsal mid-insula, and premotor cortex (Figure 4; Supplementary Table S2, S3 and S4).

There were no significant group by distance interactions. However, the longer distances (15–20 and 25–30 mm) demonstrated more apparent alterations in the left anterior insula, as well as across the cingulate, including the midcingulate (15–20 and 25–30 mm), ventral anterior (15–20 mm) and subgenual anterior cingulate (25–30 mm; Figure 3). The 25–30 mm distance was also observed to demonstrate differences between groups in the supplementary motor and premotor cortices.

The Associations Between Depressive Symptoms, Emotional Regulation, and Local Functional Connectivity

We observed a significant association between ERQ Suppression scores and local connectivity across all three IDAC distances (Figure 5). Specifically, there was a consistent association between these scores and the local connectivity of the bilateral insula cortex and fusiform gyrus. Distance specific effects were also observed in the rostral anterior cingulate (15–20 mm), amygdala (15-20 and 25-30 mm), hippocampus (15-20 and 25-30 mm), midcingulate (25-30 mm), as well as increased associations with a much larger spatial extent of the insula for the long distance (25-30 mm). The association between ERQ Suppression scores and insula local functional connectivity was also observed when examining the MDD

group alone, however, to a much lesser extent (Supplementary Figure S2; uncorrected, $p <$ .001). No significant correlations were observed between IDAC measures and MADRS scores or ERQ Reappraisal scores (pFDR < .05, KE ≥ 10). To determine whether local connectivity associations with ERQ Suppression scores were due to an indirect association with depressive symptom severity, we examined the correlations between these variables. This identified a significant but small correlation between ERQ Suppression and MADRS scores (τb = .251, p < .001). Notably, however, this was less strong than the correlation between ERQ Reappraisal and MADRS Scores (τb = -.384, p < .001), indicating that these local connectivity associations were unlikely to be simply an indirect assessment of depressive severity.

Discussion

Our investigation of the differences in local functional connectivity patterns present in adolescents and young adults with MDD showed that MDD participants had greater multi-distance local functional connectivity across several brain regions, with the largest effects observed in the retrosplenial cortex, posterior hippocampus, dorsal mid-insula, and the fusiform and lingual gyri. We did not identify reduced local connectivity in the orbitofrontal cortex as hypothesized. Additionally, we observed consistent associations between ERQ Suppression scores and the local connectivity of regions including the insula and fusiform gyrus across participants, which may have contributed to the between group differences observed in these areas.

Posterior Hippocampal and Retrosplenial Alterations

The posterior hippocampus and retrosplenial cortex have consistently shown functional alterations in depression. Meta-analyses have identified increased regional homogeneity of the left hippocampus in drug naïve MDD participants [23]. Similarly, reduced suppression of the hippocampus has been observed in depressed participants during loss reinforcement learning conditions, with the degree of this hyperactivity being associated with severity of symptoms [47]. Depression has also been associated with functional alterations in the retrosplenial cortex [48-50]. Notably, both regions are components of the medial temporal subsystem of the DMN [51], and have been broadly implicated in affective memory and information processing, particularly autobiographical memory [52, 53]. The retrosplenial cortex has also been implicated in mental imagery and projecting oneself in time, both key components in episodic memory [54]. As such, these regions are spatially and functionally

positioned to link DMN subsystems necessary for episodic memory retrieval and core DMN regions, such as the posterior cingulate cortex, necessary for the generation of self-conceptualization [53, 55, 56]. Increased local synchrony of these regions observed in depressed individuals may therefore represent changes to the processing and encoding of emotional memory [57], and specifically contribute to the negatively biased recall of stimuli hypothesized by cognitive models of depression [58].

In rodent work the retrosplenial cortex has been implicated in abnormal long-term metabolic stress responses and vulnerability to depression [59]. Excitatory efferent projections from the dorsal hippocampus, equivalent to the posterior hippocampus in humans [60], to the retrosplenial cortex have been identified as contributors to the generalization of negative memories [61]. The hippocampus demonstrates a high proportion of glutamatergic pyramidal cells [62, 63], with co-activation of these neurons during rest being shown to significantly increase following chronic stress exposure [64]. Thus, due to the sensitivity of these regions to stress [65], hippocampus and retrosplenial cortex dysfunction may represent a potential mechanism through which early life adversity results in changes to the processing and encoding of emotional memory. This in turn, would facilitate the reductions in hippocampal volume observed in older depressed participants [66-68]. It therefore appears that IDAC could be sensitive to stress-related neuroplastic changes impacting extended hippocampal function. Future examination of how these changes in local synchrony progress as a function of the chronicity of the disorder will provide insight about the longitudinal effect these local alterations have on other regions and networks.

Dorsal Mid-Insula Alterations

Compared to healthy controls, MDD participants have shown decreased activity of the mid-insula cortex during interoceptive processing [69, 70] and recall [70]. MDD participants additionally illustrate increased resting-state connectivity between the dorsal mid-insula and affective regions, including the amygdala, subgenual cingulate cortex, and orbitofrontal cortex [69]. It has been proposed that the insula illustrates a posterior-to-anterior progression of interoceptive processing, with the posterior insula mapping interoceptive signals [71, 72], and the anterior insula associated with interoceptive awareness [73, 74]. Within this framework the dorsal mid-insula is well placed to serve as an intermediary between these processes. Primate cytoarchitectonics of the precentral insular gyrus further supports this functioning, as the mid-insula both projects to [75], and receives input from [76], the anterior and posterior insula. Moreover, the mid-insula has previously been associated with interoceptive processing [77],

including both interoceptive attention [78, 79] and interoceptive recall [70]. Together this work indicates that this part of the insula may be responsible for processing interoceptive prediction errors, which occur following a mismatch between expectations concerning physiological states and bodily signals [80]. A recent meta-analysis examining alterations across multiple psychiatric disorders during interoception identified consistent changes to the mid-insula [81]. As such, the integrative role of the mid-insula may increase its sensitivity to dysfunctional signals emerging from disparate bodily and brain systems [33, 81] thereby contributing to abnormal emotional states [82].

Interestingly, we observed that the local functional connectivity of the insula was correlated with habitual use of expressive suppression. Due to expressive suppression involving inhibiting the outward expression of emotional behavior [9], its use may be viewed as an attempt to exert top-down control of interoceptive signals [81, 83]. Maladaptive emotional regulation strategies are well noted in depressive samples [84], including increased use of expressive suppression and decreased use of cognitive reappraisal [85]. While a recent review has highlighted mixed evidence concerning whether expressive suppression is associated with depression [86], longitudinal research suggests that expressive suppression may both predict [10] and be predicted by depressive symptoms [87]. The precise nature of this relationship likely depends on the developmental stage being investigated [86]. Age has been suggested as an important mediator of this relationship given that older adults use expressive suppression more frequently, however, experience less psychological distress from its use [88]. Thus, the young nature of our sample may explain the large difference in expressive suppression observed between the healthy controls and MDD participants.

Limitations

The strength and novelty of this study should be considered in the context of its limitations. Our results were derived from young adults and adolescents. It remains unclear whether these same effects would be present in older populations and how these local connectivity profiles change longitudinally and as a function of illness chronicity. Additionally, while the homogeneity of the current sample is a major strength it also limits the generalizability of our findings. As such, further work in transdiagnostic samples examining the relationship between local functional connectivity and factors such as expressive suppression may be warranted.

Conclusion

We have examined the local functional connectivity of the cerebral cortex using a novel approach to investigate brain alterations associated with MDD. We identified differences in the local synchrony of several regions, including the retrosplenial cortex, posterior hippocampus, and dorsal mid-insula. Across these regions depressed individuals demonstrated increased coupling of intracortical activity. Additionally, across participants, increased local functional connectivity of the insula was correlated with use of expressive suppression as an emotional regulation strategy. Longitudinal examination of these between group local synchrony differences will aid in identifying whether this relationship changes as a function of the chronicity of the disorder.

Declaration of Competing and Financial Interests

The authors declare no biomedical financial interests or potential conflicts of interest.

Funding

This work was supported by National Health and Medical Research Council of Australia (NHMRC) Project Grants (1064643) to BJH and to CGD (1024570).

Acknowledgments

The authors thank Katerina Stephanou, Lisa Incerti and Rebecca Kerestes for contributions to data collection, as well as staff from the Sunshine Hospital Medical Imaging Department (Western Health, Melbourne).

Table 1

Comparison of Characteristics Between Healthy Controls and Major Depressive Disorder Patients

Characteristics	Healthy Controls (N = 95)		MDD (N = 145)		Hedges' g or Cramer's V	p
	Mean or N	*SD* or Percentage	Mean or N	*SD* or Percentage		
Age (years)	20.1	2.9	19.9	2.8	.07	1.00
Female	52	54.7	81	54.4	.01	1.00
Handedness (Right)	78	82.1	139	92.7	.18	.095
Full scale IQ	110.1	8.6	105.3	9.4	.53	<.001
Ethnicity					.23	1.00
Caucasian	70	73.7	122	81.9		
Black or African	1	1.1	1	0.1		
Asian	19	20.0	23	15.4		
South American	3	3.3	1	0.1		
Indian	1	1.1	1	0.1		
Middle Eastern	1	1.1	2	0.1		
MADRS score	2.13	2.8	33.0	5.4	-6.76	<.001
ERQ Suppression score	14.35	4.7	17.79	4.2	-.78	<.001
ERQ Reappraisal score	30.76	6.2	22.0	6.9	1.02	<.001
Age of MDD onset	-	-	15.5	2.7	-	-

Figure 1. Iso-Distant Average Correlation (IDAC) brain maps across all three distances projected onto a cortical surface. Results displayed are from healthy control participants (*N* = 95) with overlay of the IDAC distances (top), 5-10 mm (red), 15-20mm (green), and 25-30mm (blue) with values capped at 95% of their respective maximal *t* values. The overlay of the 3 maps together (bottom) illustrates the consistent and unique effect across these distances through primary RGB colors and their secondary combinations. Left = Left.

Figure 2. Main effects of group across the three local connectivity distances (5-10mm, 15-20mm, and 25-30mm).

Figure 3. Group differences in the local functional connectivity for each of the 5-10mm, 15-20mm and 25-30mm distances (MDD participants > healthy controls).

Figure 4. Red, green, and blue overlays illustrating the overlapping alterations to local functional connectivity present in MDD participants.

Figure 5. Red, green, and blue overlays illustrating associations between ERQ Suppression scores and local functional connectivity across all participants.

References

1.   American Psychiatric Association D, Association AP. *Diagnostic and statistical manual of mental disorders: DSM-5*, vol. 5. American psychiatric association Washington, DC2013.

2.   Gore FM, Bloem PJ, Patton GC, Ferguson J, Joseph V, Coffey C *et al.* Global burden of disease in young people aged 10-24 years: a systematic analysis. *Lancet* 2011; **377**(9783)**:** 2093-2102.

3.   Thapar A, Collishaw S, Pine DS, Thapar AK. Depression in adolescence. *The lancet* 2012; **379**(9820)**:** 1056-1067.

4.   Fergusson DM, Horwood LJ, Ridder EM, Beautrais AL. Subthreshold depression in adolescence and mental health outcomes in adulthood. *Arch Gen Psychiat* 2005; **62**(1)**:** 66-72.

5.   Klein DN, Shankman SA, Lewinsohn PM, Seeley JR. Subthreshold Depressive Disorder in Adolescents: Predictors of Escalation to Full-Syndrome Depressive Disorders. *J Am Acad Child Psy* 2009; **48**(7)**:** 703-710.

6.   Zisook S, Lesser I, Stewart JW, Wisniewski SR, Balasubramani GK, Fava M *et al.* Effect of age at onset on the course of major depressive disorder. *Am J Psychiat* 2007; **164**(10)**:** 1539-1546.

7.   Cracco E, Goossens L, Braet C. Emotion regulation across childhood and adolescence: evidence for a maladaptive shift in adolescence. *Eur Child Adoles Psy* 2017; **26**(8)**:** 909-921.

8.   Silk JS, Steinberg L, Morris AS. Adolescents' emotion regulation in daily life: Links to depressive symptoms and problem behavior. *Child Dev* 2003; **74**(6)**:** 1869-1880.

9.   Gross JJ. Emotion Regulation: Current Status and Future Prospects. *Psychol Inq* 2015; **26**(1)**:** 1-26.

10.   Larsen JK, Vermulst AA, Ha T, Engels RCME, Geenen R, van Middendorp H *et al.* Emotion Regulation in Adolescence: A Prospective Study of Expressive Suppression and Depressive Symptoms. *J Early Adolescence* 2013; **33**(2)**:** 184-200.

11.   Li BJ, Friston K, Mody M, Wang HN, Lu HB, Hu DW. A brain network model for depression: From symptom understanding to disease intervention. *CNS Neurosci Ther* 2018; **24**(11)**:** 1004-1019.

12.   Brakowski J, Spinelli S, Dörig N, Bosch OG, Manoliu A, Holtforth MG *et al.* Resting state brain network function in major depression – Depression symptomatology, antidepressant treatment effects, future research. *Journal of Psychiatric Research* 2017; **92:** 147-159.

13.   Dunlop K, Talishinsky A, Liston C. Intrinsic Brain Network Biomarkers of Antidepressant Response: a Review. *Curr Psychiatry Rep* 2019; **21**(9)**:** 87.

14.   Jiang L, Zuo XN. Regional Homogeneity: A Multimodal, Multiscale Neuroimaging Marker of the Human Connectome. *Neuroscientist* 2016; **22**(5)**:** 486-505.

15.   Zuo XN, Xu T, Jiang L, Yang Z, Cao XY, He Y *et al.* Toward reliable characterization of functional homogeneity in the human brain: preprocessing, scan duration, imaging resolution and computational space. *Neuroimage* 2013; **65:** 374-386.

16.   Tononi G, Sporns O, Edelman GM. A measure for brain complexity: relating functional segregation and integration in the nervous system. *Proc Natl Acad Sci U S A* 1994; **91**(11)**:** 5033-5037.

17.   Guo WB, Sun XL, Liu L, Xu Q, Wu RR, Liu ZN *et al.* Disrupted regional homogeneity in treatment-resistant depression: a resting-state fMRI study. *Prog Neuropsychopharmacol Biol Psychiatry* 2011; **35**(5)**:** 1297-1302.

18.   Jiang L, Xu T, He Y, Hou X-H, Wang J, Cao X-Y *et al.* Toward neurobiological characterization of functional homogeneity in the human cortex: regional variation, morphological association and functional covariance network organization. *Brain Structure and Function* 2015; **220**(5)**:** 2485-2507.

19.   Canario E, Chen D, Biswal B. A review of resting-state fMRI and its use to examine psychiatric disorders. *Psychoradiology* 2021; **1**(1)**:** 42-53.

20.   Wei Y, Chang M, Womer FY, Zhou Q, Yin Z, Wei S *et al.* Local functional connectivity alterations in schizophrenia, bipolar disorder, and major depressive disorder. *J Affect Disord* 2018; **236:** 266-273.

21.   Chen ZQ, Du MY, Zhao YJ, Huang XQ, Li J, Lui S *et al.* Voxel-wise meta-analyses of brain blood flow and local synchrony abnormalities in medication-free patients with major depressive disorder. *J Psychiatry Neurosci* 2015; **40**(6)**:** 401-411.

22.   Iwabuchi SJ, Krishnadas R, Li C, Auer DP, Radua J, Palaniyappan L. Localized connectivity in depression: a meta-analysis of resting state functional imaging studies. *Neurosci Biobehav Rev* 2015; **51:** 77-86.

23.   Hao H, Chen C, Mao W, Zhong J, Dai Z. Aberrant brain regional homogeneity in first-episode drug-naive patients with major depressive disorder: A voxel-wise meta-analysis. *J Affect Disord* 2019; **245:** 63-71.

24.   Ke Z, Qing G, Zhiliang L, Fei X, Xiao S, Huafu C *et al.* Abnormal functional connectivity density in first-episode, drug-naive adult patients with major depressive disorder. *J Affect Disorders* 2016; **194:** 153-158.

25.   Zheng R, Chen Y, Jiang Y, Zhou B, Han S, Wei Y *et al.* Abnormal voxel-wise whole-brain functional connectivity in first-episode, drug-naïve adolescents with major depression disorder. *Eur Child Adolesc Psychiatry* 2022.

26.   Tang Q, Cui Q, Chen Y, Deng J, Sheng W, Yang Y *et al.* Shared and distinct changes in local dynamic functional connectivity patterns in major depressive and bipolar depressive disorders. *J Affect Disord* 2022; **298**(Pt A)**:** 43-50.

27.   Rolls ET, Grabenhorst F. The orbitofrontal cortex and beyond: From affect to decision-making. *Progress in Neurobiology* 2008; **86**(3)**:** 216-244.

28.   Uddin LQ, Nomi JS, Hebert-Seropian B, Ghaziri J, Boucher O. Structure and Function of the Human Insula. *J Clin Neurophysiol* 2017; **34**(4)**:** 300-306.

29.   Pastrnak M, Simkova E, Novak T. Insula activity in resting-state differentiates bipolar from unipolar depression: a systematic review and meta-analysis. *Sci Rep-Uk* 2021; **11**(1).

30.   Manoliu A, Meng C, Brandl F, Doll A, Tahmasian M, Scherr M *et al.* Insular dysfunction within the salience network is associated with severity of symptoms and aberrant inter-network connectivity in major depressive disorder. *Frontiers in Human Neuroscience* 2014; **7**.

31.   Iwabuchi SJ, Peng DH, Fang YR, Jiang KD, Liddle EB, Liddle PF *et al.* Alterations in effective connectivity anchored on the insula in major depressive disorder. *European Neuropsychopharmacology* 2014; **24**(11)**:** 1784-1792.

32.   Jamieson AJ, Harrison BJ, Razi A, Davey CG. Rostral anterior cingulate network effective connectivity in depressed adolescents and associations with treatment response in a randomized controlled trial. *Neuropsychopharmacology* 2022; **47**(6)**:** 1240-1248.

33.   Sliz D, Hayley S. Major depressive disorder and alterations in insular cortical activity: a review of current functional magnetic imaging research. *Front Hum Neurosci* 2012; **6:** 323.

34.   Macia D, Pujol J, Blanco-Hinojo L, Martinez-Vilavella G, Martin-Santos R, Deus J. Characterization of the Spatial Structure of Local Functional Connectivity Using Multidistance Average Correlation Measures. *Brain Connect* 2018; **8**(5)**:** 276-287.

35.   Sepulcre J, Liu H, Talukdar T, Martincorena I, Yeo BT, Buckner RL. The organization of local and distant functional connectivity in the human brain. *PLoS Comput Biol* 2010; **6**(6)**:** e1000808.

36.   Tomasi D, Volkow ND. Functional connectivity density mapping. *Proc Natl Acad Sci U S A* 2010; **107**(21)**:** 9885-9890.

37.   Zang Y, Jiang T, Lu Y, He Y, Tian L. Regional homogeneity approach to fMRI data analysis. *Neuroimage* 2004; **22**(1)**:** 394-400.

38.   Davey CG, Chanen AM, Hetrick SE, Cotton SM, Ratheesh A, Amminger GP *et al.* The addition of fluoxetine to cognitive behavioural therapy for youth depression (YoDA-C): a randomised, double-blind, placebo-controlled, multicentre clinical trial. *Lancet Psychiat* 2019; **6**(9)**:** 735-744.

39. Berk M, Mohebbi M, Dean OM, Cotton SM, Chanen AM, Dodd S *et al.* Youth Depression Alleviation with Anti-inflammatory Agents (YoDA-A): a randomised clinical trial of rosuvastatin and aspirin. *BMC Med* 2020; **18**(1)**:** 16.

40. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV Axis I Disorders: Clinician Version. In: Werner PD, Widiger TA (eds). *SCID-CV*1997.

41. Pujol J, Pujol N, Mané A, Martínez-Vilavella G, Deus J, Pérez-Sola V *et al.* Mapping alterations in the local synchrony of the cerebral cortex in schizophrenia. *Eur Psychiat* 2023.

42. Wilke M. An alternative approach towards assessing and accounting for individual motion in fMRI timeseries. *Neuroimage* 2012; **59**(3)**:** 2062-2072.

43. Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* 2014; **84:** 320-341.

44. Gross JJ, John OP. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol* 2003; **85**(2)**:** 348-362.

45. Holm S. A Simple Sequentially Rejective Multiple Test Procedure. *Scand J Stat* 1979; **6**(2)**:** 65-70.

46. Pujol J, Blanco-Hinojo L, Macia D, Alonso P, Harrison BJ, Martinez-Vilavella G *et al.* Mapping Alterations of the Functional Structure of the Cerebral Cortex in Obsessive-Compulsive Disorder. *Cereb Cortex* 2019; **29**(11)**:** 4753-4762.

47. Johnston BA, Tolomeo S, Gradin V, Christmas D, Matthews K, Steele JD. Failure of hippocampal deactivation during loss events in treatment-resistant depression. *Brain* 2015; **138**(Pt 9)**:** 2766-2776.

48. Kumar P, Waiter G, Ahearn T, Milders M, Reid I, Steele JD. Abnormal temporal difference reward-learning signals in major depression. *Brain* 2008; **131**(Pt 8)**:** 2084-2093.

49. Zhu X, Wang X, Xiao J, Liao J, Zhong M, Wang W *et al.* Evidence of a dissociation pattern in resting-state default mode network connectivity in first-episode, treatment-naive major depression patients. *Biol Psychiatry* 2012; **71**(7)**:** 611-617.

50. Harel EV, Tennyson RL, Fava M, Bar M. Linking major depression and the neural substrates of associative processing. *Cogn Affect Behav Neurosci* 2016; **16**(6)**:** 1017-1026.

51. Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, Buckner RL. Functional-anatomic fractionation of the brain's default network. *Neuron* 2010; **65**(4)**:** 550-562.

52.   Miller AM, Vedder LC, Law LM, Smith DM. Cues, context, and long-term memory: the role of the retrosplenial cortex in spatial cognition. *Front Hum Neurosci* 2014; **8:** 586.

53.   Andrews-Hanna JR, Smallwood J, Spreng RN. The default network and self-generated thought: component processes, dynamic control, and clinical relevance. *Ann N Y Acad Sci* 2014; **1316**(1)**:** 29-52.

54.   Chrastil ER. Heterogeneity in Human Retrosplenial Cortex: A Review of Function and Connectivity. *Behav Neurosci* 2018; **132**(5)**:** 317-338.

55.   Kaboodvand N, Backman L, Nyberg L, Salami A. The retrosplenial cortex: A memory gateway between the cortical default mode network and the medial temporal lobe. *Human Brain Mapping* 2018; **39**(5)**:** 2020-2034.

56.   Davey CG, Harrison BJ. The self on its axis: a framework for understanding depression. *Transl Psychiatry* 2022; **12**(1)**:** 23.

57.   Jaworska N, Yang XR, Knott V, MacQueen G. A review of fMRI studies during visual emotive processing in major depressive disorder. *World J Biol Psychiatry* 2015; **16**(7)**:** 448-471.

58.   Disner SG, Beevers CG, Haigh EA, Beck AT. Neural mechanisms of the cognitive model of depression. *Nat Rev Neurosci* 2011; **12**(8)**:** 467-477.

59.   Harro J, Kanarik M, Kaart T, Matrov D, Koiv K, Mallo T *et al.* Revealing the cerebral regions and networks mediating vulnerability to depression: oxidative metabolism mapping of rat brain. *Behav Brain Res* 2014; **267:** 83-94.

60.   Fanselow MS, Dong HW. Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* 2010; **65**(1)**:** 7-19.

61.   Ren LY, Meyer MAA, Grayson VS, Gao P, Guedea AL, Radulovic J. Stress-induced generalization of negative memories is mediated by an extended hippocampal circuit. *Neuropsychopharmacology* 2022; **47**(6)**:** 1280-1280.

62.   Freund TF, Buzsaki G. Interneurons of the hippocampus. *Hippocampus* 1996; **6**(4)**:** 347-470.

63.   Olbrich HG, Braak H. Ratio of pyramidal cells versus non-pyramidal cells in sector CA1 of the human Ammon's horn. *Anat Embryol (Berl)* 1985; **173**(1)**:** 105-110.

64.   Tomar A, Polygalov D, Chattarji S, McHugh TJ. Stress enhances hippocampal neuronal synchrony and alters ripple-spike interaction. *Neurobiol Stress* 2021; **14:** 100327.

65.   Corcoran KA, Yamawaki N, Leaderbrand K, Radulovic J. Role of retrosplenial cortex in processing stress-related context memories. *Behav Neurosci* 2018; **132**(5)**:** 388-395.

66.   Huang Y, Coupland NJ, Lebel RM, Carter R, Seres P, Wilman AH *et al.* Structural changes in hippocampal subfields in major depressive disorder: a high-field magnetic resonance imaging study. *Biol Psychiatry* 2013; **74**(1)**:** 62-68.

67.   Malykhin NV, Coupland NJ. Hippocampal neuroplasticity in major depressive disorder. *Neuroscience* 2015; **309:** 200-213.

68.   MacQueen G, Frodl T. The hippocampus in major depression: evidence for the convergence of the bench and bedside in psychiatric research? *Mol Psychiatry* 2011; **16**(3)**:** 252-264.

69.   Avery JA, Drevets WC, Moseman SE, Bodurka J, Barcalow JC, Simmons WK. Major Depressive Disorder Is Associated With Abnormal Interoceptive Activity and Functional Connectivity in the Insula. *Biol Psychiat* 2014; **76**(3)**:** 258-266.

70.   DeVille DC, Kerr KL, Avery JA, Burrows K, Bodurka J, Feinstein JS *et al.* The Neural Bases of Interoceptive Encoding and Recall in Healthy Adults and Adults With Depression. *Biol Psychiat-Cogn N* 2018; **3**(6)**:** 546-554.

71.   Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nature reviews neuroscience* 2002; **3**(8)**:** 655-666.

72.   Kuehn E, Mueller K, Lohmann G, Schuetz-Bosbach S. Interoceptive awareness changes the posterior insula functional connectivity profile. *Brain Struct Funct* 2016; **221**(3)**:** 1555-1571.

73.   Gu XS, Hof PR, Friston KJ, Fan J. Anterior insular cortex and emotional awareness. *Journal of Comparative Neurology* 2013; **521**(15)**:** 3371-3388.

74.   Craig AD. How do you feel—now? The anterior insula and human awareness. *Nature reviews neuroscience* 2009; **10**(1)**:** 59-70.

75.   Mesulam MM, Mufson EJ. Insula of the Old-World Monkey .1. Architectonics in the Insulo-Orbito-Temporal Component of the Paralimbic Brain. *Journal of Comparative Neurology* 1982; **212**(1)**:** 1-22.

76.   Mufson EJ, Mesulam MM. Insula of the old world monkey. II: Afferent cortical input and comments on the claustrum. *J Comp Neurol* 1982; **212**(1)**:** 23-37.

77.   Centanni SW, Janes AC, Haggerty DL, Atwood B, Hopf FW. Better living through understanding the insula: Why subregions can make all the difference. *Neuropharmacology* 2021; **198**.

78.   Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB. A link between the systems: functional differentiation and integration within the human insula revealed by meta-analysis. *Brain Structure & Function* 2010; **214**(5-6)**:** 519-534.

79.   Schulz SM. Neural correlates of heart-focused interoception: a functional magnetic resonance imaging meta-analysis. *Philos T R Soc B* 2016; **371**(1708).

80.   Barrett LF, Simmons WK. Interoceptive predictions in the brain. *Nat Rev Neurosci* 2015; **16**(7)**:** 419-429.

81.   Nord CL, Lawson RP, Dalgleish T. Disrupted Dorsal Mid-Insula Activation During Interoception Across Psychiatric Disorders. *Am J Psychiat* 2021; **178**(8)**:** 761-770.

82.   Namkung H, Kim SH, Sawa A. The Insula: An Underestimated Brain Area in Clinical Neuroscience, Psychiatry, and Neurology: (Trends in Neuroscience 40, 200-207, 2017). *Trends Neurosci* 2018; **41**(8)**:** 551-554.

83.   Paulus MP, Stein MB. Interoception in anxiety and depression. *Brain Struct Funct* 2010; **214**(5-6)**:** 451-463.

84.   Schafer JO, Naumann E, Holmes EA, Tuschen-Caffier B, Samson AC. Emotion Regulation Strategies in Depressive and Anxiety Symptoms in Youth: A Meta-Analytic Review. *J Youth Adolesc* 2017; **46**(2)**:** 261-276.

85.   Joormann J, Gotlib IH. Emotion regulation in depression: relation to cognitive inhibition. *Cogn Emot* 2010; **24**(2)**:** 281-298.

86.   Dryman MT, Heimberg RG. Emotion regulation in social anxiety and depression: a systematic review of expressive suppression and cognitive reappraisal. *Clin Psychol Rev* 2018; **65:** 17-42.

87.   Larsen JK, Vermulst AA, Eisinga R, English T, Gross JJ, Hofman E *et al.* Social Coping by Masking? Parental Support and Peer Victimization as Mediators of the Relationship Between Depressive Symptoms and Expressive Suppression in Adolescents. *J Youth Adolescence* 2012; **41**(12)**:** 1628-1642.

88.   Brummer L, Stopa L, Bucks R. The influence of age on emotion regulation strategies and psychological distress. *Behav Cogn Psychother* 2014; **42**(6)**:** 668-681.