13 Graded changes in local functional connectivity of the cerebral cortex in young people with
14 depression
15
16
17

18 *Alec J. Jamieson[1], Christopher G. Davey[1], Jesus Pujol[2], Laura Blanco-Hinojo[2], & Ben J.
19 Harrison[1]
20 [1]Department of Psychiatry, The University of Melbourne, Australia
21 [2]MRI Research Unit, Department of Radiology, Hospital del Mar, Barcelona, Spain

29 **Corresponding author:** Dr. Alec Jamieson. Melbourne Neuropsychiatry Centre,
30 Department of Psychiatry, The University of Melbourne. Level 3, 161 Barry St. Carlton, VIC
31 3053. Phone: (+61) 449 501 556 | Email: alec.jamieson@unimelb.edu.au

32
33
34

Abstract

Major depressive disorder (MDD) is marked by significant changes to the coupling of

spontaneous neural activity within various brain regions. However, many methods for

assessing this local connectivity use fixed or arbitrary neighborhood sizes, resulting in a

decreased capacity to capture smooth changes to the spatial gradient of local correlations. A

newly developed method sensitive to classical anatomo-functional boundaries, Iso-Distant

Average Correlation (IDAC), was therefore used to examine depression associated alterations

to the local functional connectivity of the brain. One-hundred and forty-five adolescents and

young adults with MDD and 95 healthy controls underwent a resting-state fMRI scan.

Whole-brain functional connectivity maps of intracortical neural activity within iso-distant

local areas (5-10mm, 15-20mm, and 25-30mm) were generated to characterize local fMRI

signal similarities. Across all spatial distances, MDD participants demonstrated greater local

functional connectivity of the bilateral posterior hippocampus, retrosplenial cortex, dorsal

insula, fusiform gyrus, and supplementary motor area. Additionally, in the short and medium

range connections there were depression associated alterations in the midcingulate (15-20mm

and 25-30mm) and subgenual anterior cingulate (15-20mm). Our study identified increased

synchrony of the neural activity in several regions commonly implicated in the neurobiology

of depression; however, a subset of identified effects was dependent on the spatial distance

under consideration. Longitudinal examination of these effects will clarify whether these

differences are also found in other age groups and if this synchrony is additionally altered by

continued disease progression.

Introduction

Major depressive disorder (MDD) is a highly prevalent and disabling mental health condition characterized by pervasive symptoms of sadness, hopelessness, and anhedonia [1]. Depression often first arises during adolescence [2,3], with early onset being a strong predictor of recurrent depression [4,5] as well as greater reductions in quality of life and wellbeing into adulthood [6]. Early adolescence is marked by increased use of maladaptive emotional regulation strategies which in turn are associated with exacerbated mental health issues [7,8]. One such strategy, known expressive suppression, involves inhibiting the outward expression of emotional behavior [9], has been specifically associated with depressive symptoms severity [10]. However, there remains limited work examining neurobiological underpinnings of this early stage of the disorder.

Previous neuroimaging research has attempted to uncover the neurobiological underpinnings of MDD by focusing on long-range functional connectivity, examining the temporal correlations of large-scale, spatially distinct brain networks. Within this framework, MDD has been associated with alterations across multiple intrinsic brain networks [11,12], including the default mode (DMN), salience, and central executive networks [13]. However, there has been increasing interest in assessing local functional connectivity profiles [14,15], as the functional integration of local neuronal groups appears to represent a key organizing principle in the brains of higher vertebrates [16]. Local functional connectivity measures assess the temporal coherence of the functional magnetic resonance imaging (fMRI) signal between a given voxel and neighboring voxels [14]. Measures of this 'localized synchrony' provide a framework for examining functional disruptions at rest without a priori constraints, thereby allowing for data-driven identification of regional abnormalities across a range of mental health disorders [17]. Moreover, regional variation in homogeneity has been associated with the hierarchical organization of information processing across the brain, and therefore alterations to these measures may serve as key markers of human brain function [14,18] and a sensitive tool for identifying brain alterations across mental health disorders [19,20].

Meta-analyses examining regional homogeneity in MDD have revealed inconsistent differences in the directionality of effects and regions implicated [21-23]. Altered local connectivity has been noted across regions including the parahippocampal gyrus, insula, and medial prefrontal cortex [21,22]. Other local functional connectivity measures, including functional connectivity density and dynamic regional phase synchrony, have illustrated reduced local connectivity in the anterior cingulate cortex, precuneus, hippocampus, thalamus, and insula [24-26]. Conversely, recent work examining first-episode MDD observed

increased regional homogeneity of the hippocampus and insula but reduced local connectivity

of the orbitofrontal cortex [23]. It is likely that the specific method used, including how

neighborhood boundaries are defined, as well as chronicity of the disorder, contributes to the

inconsistencies observed between studies. Despite these inconsistencies, the observed

alterations appear to be localized to regions implicated in affective processing and regulation

[27, 28]. Of particular note is the implication of insula dysfunction across measures of local and

remote functional connectivity [29-32], due to its hypothesized role in the integration of

autonomic, emotional, and interoceptive stimuli [33] and potential specificity in delineating

MDD from bipolar disorder [29].

We aimed to investigate local functional connectivity alterations in a large sample of

adolescents and young adults with MDD, using the recently introduced framework -

IsoDistant Average Correlation (IDAC) [34]. Measures of local functional connectivity often

use arbitrary and binarized neighborhood boundaries, thereby losing the capacity to describe

the rich smooth spatial gradient of local fMRI correlations [35-37]. IDAC assesses the average

temporal correlation of one voxel with all neighboring voxels within different spatial lags,

thereby overcoming these limitations [34]. Based on previous findings, we hypothesize that

MDD participants would illustrate increased local functional connectivity of the insula,

anterior cingulate, and hippocampus and reduced local functional connectivity of the

orbitofrontal cortex. Given the developmental nature of our cohort, we additionally

investigated whether local connectivity was associated with measures of depressive symptom

severity, and emotion processing and regulation.

Methods

One hundred and forty-nine unmedicated, help-seeking MDD participants were recruited as

part of the Youth Depression Alleviation-Combined Treatment and -Augmentation trials, for

full details see Davey and colleagues [38] and Berk and colleagues [39]. MDD participants were

between 15 to 25 years of age and were recruited through specialist mental health clinics in

the northern and western suburbs of Melbourne, Australia. These participants had a current

diagnosis of MDD, as assessed by the Structured Clinical Interview for DSM-IV Axis I

Disorders (SCID) [40]. Depressive symptoms were at least of a moderate level of severity,

indicated by a Montgomery-Åsberg Depression Rating Scale (MADRS) score greater than or

equal to 20. Exclusion criteria included a lifetime or current diagnosis of a psychotic or

bipolar disorder, current treatment with antidepressant medication, or MRI contraindications

including pregnancy. One hundred age and sex-matched healthy participants were also

recruited through online advertisements. They had no past mental health disorder diagnoses

as assessed through SCID criteria. Due to excessive head motion (three MDD participants,

four controls; see below) and incidental findings (one MDD participant, one healthy control),

a total of four MDD participants and five controls were excluded from further analysis. This

resulted in 95 healthy controls and 145 MDD participants in our final sample. This study and

consent process was approved by the Melbourne Health Human Research and Ethics

Committee. An informed consent form was provided to, and signed by, all participants in this

study. For participants under the age of 18, both participant and parental consent were

required.

Image Acquisition

A 3T General Electric Signa Excite system with an eight-channel phased-array head coil was

used in combination with ASSET parallel imaging. The functional sequence consisted of a

single shot gradient-recalled echo-planar imaging sequence in the steady state (repetition

time, 2000 ms; echo time, 35 ms; and pulse angle, 90°) in a 23-cm field-of-view, with a 64 x

64-pixel matrix and a slice thickness of 3.5mm (no gap). Thirty-six interleaved slices were

acquired parallel to the anterior-posterior commissure line with a 20° anterior tilt to better

cover ventral prefrontal brain regions. The total sequence duration was 8 minutes,

corresponding to 240 whole-brain echo-planar imaging volumes. Participants were instructed

to keep their eyes closed for the duration of the scan. The first 4 volumes from each run were

automatically discarded to allow for signal equilibration. A T1-weighted high-resolution

anatomical image was acquired for each participant to assist with functional timeseries co-

registration (140 contiguous slices; repetition time, 7.9 ms; echo time, 3 ms; flip angle, 13°;

in a 25.6 cm field-of-view, with a 256 x 256-pixel matrix and a slice thickness of 1 mm). To

assist with noise reduction and head immobility, all participants used earplugs and had their

heads supported with foam-padding inserts.

Image Preprocessing

Imaging data was transferred to a Unix-based platform that ran MATLAB Version 9.3 (The

MathWorks Inc., Natick, USA) and Statistical Parametric Mapping (SPM) Version 12 v7487

(Wellcome Trust Centre for Neuroimaging, London, UK). Preprocessing followed previously

reported steps for IDAC analysis [41]. In brief, for each subject functional MRI images were

slice-time corrected, realigned, and co-registered to their corresponding anatomical image

with an affine transformation. Spatial normalization occurred through a back-transformation

process, as such the individual 3D anatomical images were segmented and registered to MNI

space so that the resulting deformation fields could be applied to the IDAC maps (see below).

Images were then re-sliced to 3x3x3 mm resolution and smoothed by convolving the image

with a 4x4x4 mm full width at half maximum Gaussian kernel. Motion fingerprint [42] was

used to quantify participant head motion. Participants were excluded if movement exceeded a

mean total displacement of 3 mm (~1 native voxel).

IDAC analysis

IDAC maps describe the pattern of correlation decay propagating from each voxel across the

brain. This graded change in local functional connectivity provides a more comprehensive

characterisation of the brain's functional structure and has been shown to differentiate the

human cortex into regions consistent with traditional brain atlases as well as being sensitive

to different brain functional states [34]. For a detailed overview of IDAC see Macia and

colleagues [34]. Whole-cortex IDAC maps were computed using the mean correlation z-score

of each voxel with all neighboring voxels placed at increasingly spaced iso-distant intervals.

IDAC maps were calculated for three iso-distant intervals: 5-10 mm, 15-20 mm, and 25-30

mm, and conducted separately for the right and left hemisphere in native space. Covariates

included in this analysis were the six rigid body realignment parameters, their first-order

derivatives, average white matter, cerebrospinal fluid, and global brain signal. Motion

scrubbing using the realignment parameters from preprocessing was additionally conducted

to discard motion-affected image volumes [43]. For volumes with greater than 0.2 mm of inter-

frame motion, the corresponding volume and those immediately preceding and succeeding

were discarded. A discrete cosine transform filter was applied for frequencies outside the

.01–0.1 Hz interval.

The resulting maps were then normalized to the Montreal Neurological Institute

(MNI) space using a back-transformation process to enable group inference. RGB color

overlays were used for displaying the values from the three distances simultaneously. For this

study, red represented results from 5-10 mm IDAC maps, green from 15-20 mm, and blue

from 25-30 mm. Overlaps of these distances are illustrated through the respective secondary

colors.

Statistical Analysis

Between groups analysis were conducted using a 2×3 ANOVA (group [healthy control and

MDD] by distance [5–10, 15–20, and 25–30 mm]). Following examination of the Effect of

Group F contrast, post hoc t-tests were performed to examine group differences in both

directions for each of the three IDAC maps. This resulted in six primary contrasts of interest.

Accordingly, we adjusted the false-discovery error rate (FDR) corrected threshold to .008

(0.05/6). Thus all results displayed were estimated with a whole-brain, FDR corrected

threshold of $p < .008$, KE ≥ 10 voxels. Secondary level covariates included age, gender, and

handedness.

207   To investigate whether these between group differences were also associated with

depressive symptom severity we estimated voxel-wise correlations between MADRS scores

and whole-brain functional connectivity for the 5–10, 15–20, and 25–30 mm IDAC distances.

Given the regions implicated in the between groups comparison, we then examined whether

local functional connectivity changes were associated with the scores from the Emotional

Regulation Questionnaire (ERQ) [44]. The instrument assesses the two means by which people

attempt to regulate their emotions: by expressive suppression ('Suppression') and by

cognitive reappraisal ('Reappraisal'). The scores on the subscales were normally distributed

across groups (Supplementary Figure S1), and analyses were conducted across both groups

and in the MDD group alone. Analyses of between-group differences for clinical and

demographic characteristics were calculated in SPSS version 27 (IBMCorp., Armonk, NY).

Comparisons were adjusted for multiple comparisons using the Holm-Bonferroni correction

[45] to determine significance ($p < .05$).

221   **Results**

Group Differences Between Demographic and Clinical Characteristics

Demographic and clinical characteristics of the sample are reported in Table 1. Healthy

controls and MDD patients differed significantly on MADRS symptoms ($t(233.5) = - 58.80$,

$p < .001$), ERQ Suppression ($t(241) = -5.94$, $p < .001$) and Reappraisal scores ($t(216.6) =$

10.60, $p < .001$) as well as in IQ ($t(242) = 4.03$, $p < .001$).

Functional Structure of the Cerebral Cortex

To ensure that our results are consistent with previously illustrated anatomo-functional

boundaries [34, 41, 46], whole-brain maps were generated across the three distances for the

healthy controls (Figure 1). Consistent with previous effects, the visual association cortex

demonstrated high connectivity across all local distance ranges. The bilateral inferior parietal

lobules, including the angular and supramarginal gyri, were connected at short and medium

distances.

Group Differences in the Functional Structure of the Cerebral Cortex

The Effect of Group contrast demonstrated that MDD participants and healthy controls had

significant differences across the three connectivity distances (Figure 2, Supplementary Table

S1). These effects were most predominately observed in the bilateral posterior hippocampus

extending to the retrosplenial cortex and ventral posterior cingulate as well as to the lingual

and fusiform gyri. Additional large effects were observed in the dorsal mid to posterior

insular cortices and right dorsal premotor cortex. Subsequent t-test for each distance 5–10,

15–20, and 25–30 mm illustrated that these results were all driven by greater local

connectivity in MDD participants compared to controls (Figure 3). As such, across all

distances MDD participants demonstrated greater local functional connectivity of the

bilateral extended posterior hippocampus, fusiform and lingual gyri, dorsal mid-insula, and

premotor cortex (Figure 4; Supplementary Table S2, S3 and S4).

There were no significant group by distance interactions. However, the longer distances (15–

20 and 25–30 mm) demonstrated more apparent alterations in the left anterior insula, as well

as across the cingulate, including the midcingulate (15–20 and 25–30 mm), ventral anterior

(15–20 mm) and subgenual anterior cingulate (25–30 mm; Figure 3). The 25–30 mm distance

was also observed to demonstrate differences between groups in the supplementary motor

and premotor cortices.

The Associations Between Depressive Symptoms, Emotional Regulation, and Local

Functional Connectivity

We observed a significant association between ERQ Suppression scores and local

connectivity across all three IDAC distances (Figure 5). Specifically, there was a consistent

association between these scores and the local connectivity of the bilateral insula cortex and

fusiform gyrus. Distance specific effects were also observed in the rostral anterior cingulate

(15–20 mm), amygdala (15-20 and 25-30 mm), hippocampus (15-20 and 25-30 mm),

midcingulate (25-30 mm), as well as increased associations with a much larger spatial extent

of the insula for the long distance (25-30 mm). The association between ERQ Suppression

scores and insula local functional connectivity was also observed when examining the MDD

group alone, however, to a much lesser extent (Supplementary Figure S2; uncorrected, $p <$

.001). No significant correlations were observed between IDAC measures and MADRS

scores or ERQ Reappraisal scores (pFDR $< .05$, KE ≥ 10). To determine whether local

connectivity associations with ERQ Suppression scores were due to an indirect association

with depressive symptom severity, we examined the correlations between these variables.

This identified a significant but small correlation between ERQ Suppression and MADRS

scores ($\tau b = .251$, $p < .001$). Notably, however, this was less strong than the correlation

between ERQ Reappraisal and MADRS Scores ($\tau b = -.384$, $p < .001$), indicating that these

local connectivity associations were unlikely to be simply an indirect assessment of

depressive severity.

277   **Discussion**

Our investigation of the differences in local functional connectivity patterns present in

adolescents and young adults with MDD showed that MDD participants had greater multi-

distance local functional connectivity across several brain regions, with the largest effects

observed in the retrosplenial cortex**,** posterior hippocampus, dorsal mid-insula, and the

fusiform and lingual gyri. We did not identify reduced local connectivity in the orbitofrontal

cortex as hypothesized. Additionally, we observed consistent associations between ERQ

Suppression scores and the local connectivity of regions including the insula and fusiform

gyrus across participants, which may have contributed to the between group differences

observed in these areas.

Posterior Hippocampal and Retrosplenial Alterations

The posterior hippocampus and retrosplenial cortex have consistently shown functional

alterations in depression. Meta-analyses have identified increased regional homogeneity of

the left hippocampus in drug naïve MDD participants [23]. Similarly, reduced suppression of

the hippocampus has been observed in depressed participants during loss reinforcement

learning conditions, with the degree of this hyperactivity being associated with severity of

symptoms [47]. Depression has also been associated with functional alterations in the

retrosplenial cortex [48-50]. Notably, both regions are components of the medial temporal

subsystem of the DMN [51], and have been broadly implicated in affective memory and

information processing, particularly autobiographical memory [52, 53]. The retrosplenial cortex

has also been implicated in mental imagery and projecting oneself in time, both key

components in episodic memory [54]. As such, these regions are spatially and functionally

positioned to link DMN subsystems necessary for episodic memory retrieval and core DMN

regions, such as the posterior cingulate cortex, necessary for the generation of self-

conceptualization [53, 55, 56]. Increased local synchrony of these regions observed in depressed

individuals may therefore represent changes to the processing and encoding of emotional

memory [57], and specifically contribute to the negatively biased recall of stimuli hypothesized

by cognitive models of depression [58].

In rodent work the retrosplenial cortex has been implicated in abnormal long-term

metabolic stress responses and vulnerability to depression [59]. Excitatory efferent projections

from the dorsal hippocampus, equivalent to the posterior hippocampus in humans [60], to the

retrosplenial cortex have been identified as contributors to the generalization of negative

memories [61]. The hippocampus demonstrates a high proportion of glutamatergic pyramidal

cells [62, 63], with co-activation of these neurons during rest being shown to significantly

increase following chronic stress exposure [64]. Thus, due to the sensitivity of these regions to

stress [65], hippocampus and retrosplenial cortex dysfunction may represent a potential

mechanism through which early life adversity results in changes to the processing and

encoding of emotional memory. This in turn, would facilitate the reductions in hippocampal

volume observed in older depressed participants [66-68]. It therefore appears that IDAC could be

sensitive to stress-related neuroplastic changes impacting extended hippocampal function.

Future examination of how these changes in local synchrony progress as a function of the

chronicity of the disorder will provide insight about the longitudinal effect these local

alterations have on other regions and networks.

Dorsal Mid-Insula Alterations

Compared to healthy controls, MDD participants have shown decreased activity of the mid-

insula cortex during interoceptive processing [69, 70] and recall [70]. MDD participants

additionally illustrate increased resting-state connectivity between the dorsal mid-insula and

affective regions, including the amygdala, subgenual cingulate cortex, and orbitofrontal

cortex [69]. It has been proposed that the insula illustrates a posterior-to-anterior progression of

interoceptive processing, with the posterior insula mapping interoceptive signals [71, 72], and the

anterior insula associated with interoceptive awareness [73, 74]. Within this framework the

dorsal mid-insula is well placed to serve as an intermediary between these processes. Primate

cytoarchitectonics of the precentral insular gyrus further supports this functioning, as the

mid-insula both projects to [75], and receives input from [76], the anterior and posterior insula.

Moreover, the mid-insula has previously been associated with interoceptive processing [77],

including both interoceptive attention [78, 79] and interoceptive recall [70]. Together this work

indicates that this part of the insula may be responsible for processing interoceptive

prediction errors, which occur following a mismatch between expectations concerning

physiological states and bodily signals [80]. A recent meta-analysis examining alterations

across multiple psychiatric disorders during interoception identified consistent changes to the

mid-insula [81]. As such, the integrative role of the mid-insula may increase its sensitivity to

dysfunctional signals emerging from disparate bodily and brain systems [33, 81] thereby

contributing to abnormal emotional states [82].

Interestingly, we observed that the local functional connectivity of the insula was

correlated with habitual use of expressive suppression. Due to expressive suppression

involving inhibiting the outward expression of emotional behavior [9], its use may be viewed as

an attempt to exert top-down control of interoceptive signals [81, 83]. Maladaptive emotional

regulation strategies are well noted in depressive samples [84], including increased use of

expressive suppression and decreased use of cognitive reappraisal [85]. While a recent review

has highlighted mixed evidence concerning whether expressive suppression is associated

with depression [86], longitudinal research suggests that expressive suppression may both

predict [10] and be predicted by depressive symptoms [87]. The precise nature of this relationship

likely depends on the developmental stage being investigated [86]. Age has been suggested as

an important mediator of this relationship given that older adults use expressive suppression

more frequently, however, experience less psychological distress from its use [88]. Thus, the

young nature of our sample may explain the large difference in expressive suppression

observed between the healthy controls and MDD participants.

Limitations

The strength and novelty of this study should be considered in the context of its limitations.

Our results were derived from young adults and adolescents. It remains unclear whether these

same effects would be present in older populations and how these local connectivity profiles

change longitudinally and as a function of illness chronicity. Additionally, while the

homogeneity of the current sample is a major strength it also limits the generalizability of our

findings. As such, further work in transdiagnostic samples examining the relationship

between local functional connectivity and factors such as expressive suppression may be

warranted.

Conclusion

We have examined the local functional connectivity of the cerebral cortex using a novel

approach to investigate brain alterations associated with MDD. We identified differences in

the local synchrony of several regions, including the retrosplenial cortex, posterior

hippocampus, and dorsal mid-insula. Across these regions depressed individuals

demonstrated increased coupling of intracortical activity. Additionally, across participants,

increased local functional connectivity of the insula was correlated with use of expressive

suppression as an emotional regulation strategy. Longitudinal examination of these between

group local synchrony differences will aid in identifying whether this relationship changes as

a function of the chronicity of the disorder.

Declaration of Competing and Financial Interests

The authors declare no biomedical financial interests or potential conflicts of interest.

Funding

This work was supported by National Health and Medical Research Council of Australia

(NHMRC) Project Grants (1064643) to BJH and to CGD (1024570).

Acknowledgments

The authors thank Katerina Stephanou, Lisa Incerti and Rebecca Kerestes for contributions to

data collection, as well as staff from the Sunshine Hospital Medical Imaging Department

(Western Health, Melbourne).

397   Table 1

398   *Comparison of Characteristics Between Healthy Controls and Major Depressive Disorder*

399   *Patients*

Characteristics	Healthy Controls (N = 95)		MDD (N = 145)		Hedges' g or Cramer's V	p
	Mean or N	SD or Percentage	Mean or N	SD or Percentage		
Age (years)	20.1	2.9	19.9	2.8	.07	1.00
Female	52	54.7	81	54.4	.01	1.00
Handedness (Right)	78	82.1	139	92.7	.18	.095
Full scale IQ	110.1	8.6	105.3	9.4	.53	<.001
Ethnicity					.23	1.00
Caucasian	70	73.7	122	81.9		
Black or African	1	1.1	1	0.1		
Asian	19	20.0	23	15.4		
South American	3	3.3	1	0.1		
Indian	1	1.1	1	0.1		
Middle Eastern	1	1.1	2	0.1		
MADRS score	2.13	2.8	33.0	5.4	-6.76	<.001
ERQ Suppression score	14.35	4.7	17.79	4.2	-.78	<.001
ERQ Reappraisal score	30.76	6.2	22.0	6.9	1.02	<.001
Age of MDD onset	-	-	15.5	2.7	-	-

403
Figure 1. Iso-Distant Average Correlation (IDAC) brain maps across all three distances

projected onto a cortical surface. Results displayed are from healthy control participants ($N =$

95) with overlay of the IDAC distances (top), 5-10 mm (red), 15-20mm (green), and 25-

30mm (blue) with values capped at 95% of their respective maximal t values. The overlay of

the 3 maps together (bottom) illustrates the consistent and unique effect across these

distances through primary RGB colors and their secondary combinations. Left = Left.

Figure 2. Main effects of group across the three local connectivity distances (5-10mm, 15-20mm, and 25-30mm).

445 *Figure 3.* Group differences in the local functional connectivity for each of the 5-10mm, 15-

446 20mm and 25-30mm distances (MDD participants > healthy controls).

Figure 4. Red, green, and blue overlays illustrating the overlapping alterations to local functional connectivity present in MDD participants.

Figure 5. Red, green, and blue overlays illustrating associations between ERQ Suppression scores and local functional connectivity across all participants.

References

1. American Psychiatric Association D, Association AP. *Diagnostic and statistical manual of mental disorders: DSM-5*, vol. 5. American psychiatric association Washington, DC2013.

2. Gore FM, Bloem PJ, Patton GC, Ferguson J, Joseph V, Coffey C *et al.* Global burden of disease in young people aged 10-24 years: a systematic analysis. *Lancet* 2011; **377**(9783)**:** 2093-2102.

3. Thapar A, Collishaw S, Pine DS, Thapar AK. Depression in adolescence. *The lancet* 2012; **379**(9820)**:** 1056-1067.

4. Fergusson DM, Horwood LJ, Ridder EM, Beautrais AL. Subthreshold depression in adolescence and mental health outcomes in adulthood. *Arch Gen Psychiat* 2005; **62**(1)**:** 66-72.

5. Klein DN, Shankman SA, Lewinsohn PM, Seeley JR. Subthreshold Depressive Disorder in Adolescents: Predictors of Escalation to Full-Syndrome Depressive Disorders. *J Am Acad Child Psy* 2009; **48**(7)**:** 703-710.

6. Zisook S, Lesser I, Stewart JW, Wisniewski SR, Balasubramani GK, Fava M *et al.* Effect of age at onset on the course of major depressive disorder. *Am J Psychiat* 2007; **164**(10)**:** 1539-1546.

7. Cracco E, Goossens L, Braet C. Emotion regulation across childhood and adolescence: evidence for a maladaptive shift in adolescence. *Eur Child Adoles Psy* 2017; **26**(8)**:** 909-921.

8. Silk JS, Steinberg L, Morris AS. Adolescents' emotion regulation in daily life: Links to depressive symptoms and problem behavior. *Child Dev* 2003; **74**(6)**:** 1869-1880.

9. Gross JJ. Emotion Regulation: Current Status and Future Prospects. *Psychol Inq* 2015; **26**(1)**:** 1-26.

10. Larsen JK, Vermulst AA, Ha T, Engels RCME, Geenen R, van Middendorp H *et al.* Emotion Regulation in Adolescence: A Prospective Study of Expressive Suppression and Depressive Symptoms. *J Early Adolescence* 2013; **33**(2)**:** 184-200.

11. Li BJ, Friston K, Mody M, Wang HN, Lu HB, Hu DW. A brain network model for depression: From symptom understanding to disease intervention. *CNS Neurosci Ther* 2018; **24**(11)**:** 1004-1019.

12. Brakowski J, Spinelli S, Dörig N, Bosch OG, Manoliu A, Holtforth MG *et al.* Resting state brain network function in major depression – Depression symptomatology, antidepressant treatment effects, future research. *Journal of Psychiatric Research* 2017; **92:** 147-159.

13. Dunlop K, Talishinsky A, Liston C. Intrinsic Brain Network Biomarkers of Antidepressant Response: a Review. *Curr Psychiatry Rep* 2019; **21**(9)**:** 87.

562   14.   Jiang L, Zuo XN. Regional Homogeneity: A Multimodal, Multiscale Neuroimaging
563   Marker of the Human Connectome. *Neuroscientist* 2016; **22**(5)**:** 486-505.
564

565   15.   Zuo XN, Xu T, Jiang L, Yang Z, Cao XY, He Y *et al.* Toward reliable
566   characterization of functional homogeneity in the human brain: preprocessing, scan
567   duration, imaging resolution and computational space. *Neuroimage* 2013; **65:** 374-
568   386.
569

570   16.   Tononi G, Sporns O, Edelman GM. A measure for brain complexity: relating
571   functional segregation and integration in the nervous system. *Proc Natl Acad Sci U S*
572   *A* 1994; **91**(11)**:** 5033-5037.
573

574   17.   Guo WB, Sun XL, Liu L, Xu Q, Wu RR, Liu ZN *et al.* Disrupted regional
575   homogeneity in treatment-resistant depression: a resting-state fMRI study. *Prog*
576   *Neuropsychopharmacol Biol Psychiatry* 2011; **35**(5)**:** 1297-1302.
577

578   18.   Jiang L, Xu T, He Y, Hou X-H, Wang J, Cao X-Y *et al.* Toward neurobiological
579   characterization of functional homogeneity in the human cortex: regional variation,
580   morphological association and functional covariance network organization. *Brain*
581   *Structure and Function* 2015; **220**(5)**:** 2485-2507.
582

583   19.   Canario E, Chen D, Biswal B. A review of resting-state fMRI and its use to examine
584   psychiatric disorders. *Psychoradiology* 2021; **1**(1)**:** 42-53.
585

586   20.   Wei Y, Chang M, Womer FY, Zhou Q, Yin Z, Wei S *et al.* Local functional
587   connectivity alterations in schizophrenia, bipolar disorder, and major depressive
588   disorder. *J Affect Disord* 2018; **236:** 266-273.
589

590   21.   Chen ZQ, Du MY, Zhao YJ, Huang XQ, Li J, Lui S *et al.* Voxel-wise meta-analyses
591   of brain blood flow and local synchrony abnormalities in medication-free patients
592   with major depressive disorder. *J Psychiatry Neurosci* 2015; **40**(6)**:** 401-411.
593

594   22.   Iwabuchi SJ, Krishnadas R, Li C, Auer DP, Radua J, Palaniyappan L. Localized
595   connectivity in depression: a meta-analysis of resting state functional imaging studies.
596   *Neurosci Biobehav Rev* 2015; **51:** 77-86.
597

598   23.   Hao H, Chen C, Mao W, Zhong J, Dai Z. Aberrant brain regional homogeneity in
599   first-episode drug-naive patients with major depressive disorder: A voxel-wise meta-
600   analysis. *J Affect Disord* 2019; **245:** 63-71.
601

602   24.   Ke Z, Qing G, Zhiliang L, Fei X, Xiao S, Huafu C *et al.* Abnormal functional
603   connectivity density in first-episode, drug-naive adult patients with major depressive
604   disorder. *J Affect Disorders* 2016; **194:** 153-158.
605

606   25.   Zheng R, Chen Y, Jiang Y, Zhou B, Han S, Wei Y *et al.* Abnormal voxel-wise whole-
607   brain functional connectivity in first-episode, drug-naïve adolescents with major
608   depression disorder. *Eur Child Adolesc Psychiatry* 2022.
609

610   26.   Tang Q, Cui Q, Chen Y, Deng J, Sheng W, Yang Y *et al.* Shared and distinct changes
611   in local dynamic functional connectivity patterns in major depressive and bipolar
612   depressive disorders. *J Affect Disord* 2022; **298**(Pt A)**:** 43-50.
613

614   27.   Rolls ET, Grabenhorst F. The orbitofrontal cortex and beyond: From affect to
615   decision-making. *Progress in Neurobiology* 2008; **86**(3)**:** 216-244.
616

617   28.   Uddin LQ, Nomi JS, Hebert-Seropian B, Ghaziri J, Boucher O. Structure and
618   Function of the Human Insula. *J Clin Neurophysiol* 2017; **34**(4)**:** 300-306.
619

620   29.   Pastrnak M, Simkova E, Novak T. Insula activity in resting-state differentiates bipolar
621   from unipolar depression: a systematic review and meta-analysis. *Sci Rep-Uk* 2021;
622   **11**(1).
623

624   30.   Manoliu A, Meng C, Brandl F, Doll A, Tahmasian M, Scherr M *et al.* Insular
625   dysfunction within the salience network is associated with severity of symptoms and
626   aberrant inter-network connectivity in major depressive disorder. *Frontiers in Human*
627   *Neuroscience* 2014; **7**.
628

629   31.   Iwabuchi SJ, Peng DH, Fang YR, Jiang KD, Liddle EB, Liddle PF *et al.* Alterations
630   in effective connectivity anchored on the insula in major depressive disorder.
631   *European Neuropsychopharmacology* 2014; **24**(11)**:** 1784-1792.
632

633   32.   Jamieson AJ, Harrison BJ, Razi A, Davey CG. Rostral anterior cingulate network
634   effective connectivity in depressed adolescents and associations with treatment
635   response in a randomized controlled trial. *Neuropsychopharmacology* 2022; **47**(6)**:**
636   1240-1248.
637

638   33.   Sliz D, Hayley S. Major depressive disorder and alterations in insular cortical activity:
639   a review of current functional magnetic imaging research. *Front Hum Neurosci* 2012;
640   **6:** 323.
641

642   34.   Macia D, Pujol J, Blanco-Hinojo L, Martinez-Vilavella G, Martin-Santos R, Deus J.
643   Characterization of the Spatial Structure of Local Functional Connectivity Using
644   Multidistance Average Correlation Measures. *Brain Connect* 2018; **8**(5)**:** 276-287.
645

646   35.   Sepulcre J, Liu H, Talukdar T, Martincorena I, Yeo BT, Buckner RL. The
647   organization of local and distant functional connectivity in the human brain. *PLoS*
648   *Comput Biol* 2010; **6**(6)**:** e1000808.
649

650   36.   Tomasi D, Volkow ND. Functional connectivity density mapping. *Proc Natl Acad Sci*
651   *U S A* 2010; **107**(21)**:** 9885-9890.
652

653   37.   Zang Y, Jiang T, Lu Y, He Y, Tian L. Regional homogeneity approach to fMRI data
654   analysis. *Neuroimage* 2004; **22**(1)**:** 394-400.
655

656   38.   Davey CG, Chanen AM, Hetrick SE, Cotton SM, Ratheesh A, Amminger GP *et al.*
657   The addition of fluoxetine to cognitive behavioural therapy for youth depression
658   (YoDA-C): a randomised, double-blind, placebo-controlled, multicentre clinical trial.
659   *Lancet Psychiat* 2019; **6**(9)**:** 735-744.

661   39.   Berk M, Mohebbi M, Dean OM, Cotton SM, Chanen AM, Dodd S *et al.* Youth
662   Depression Alleviation with Anti-inflammatory Agents (YoDA-A): a randomised
663   clinical trial of rosuvastatin and aspirin. *BMC Med* 2020; **18**(1)**:** 16.
664

665   40.   First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for
666   DSM-IV Axis I Disorders: Clinician Version. In: Werner PD, Widiger TA (eds).
667   *SCID-CV*1997.
668

669   41.   Pujol J, Pujol N, Mané A, Martínez-Vilavella G, Deus J, Pérez-Sola V *et al.* Mapping
670   alterations in the local synchrony of the cerebral cortex in schizophrenia. *Eur Psychiat*
671   2023.
672

673   42.   Wilke M. An alternative approach towards assessing and accounting for individual
674   motion in fMRI timeseries. *Neuroimage* 2012; **59**(3)**:** 2062-2072.
675

676   43.   Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods
677   to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*
678   2014; **84:** 320-341.
679

680   44.   Gross JJ, John OP. Individual differences in two emotion regulation processes:
681   implications for affect, relationships, and well-being. *J Pers Soc Psychol* 2003; **85**(2)**:**
682   348-362.
683

684   45.   Holm S. A Simple Sequentially Rejective Multiple Test Procedure. *Scand J Stat*
685   1979; **6**(2)**:** 65-70.
686

687   46.   Pujol J, Blanco-Hinojo L, Macia D, Alonso P, Harrison BJ, Martinez-Vilavella G *et*
688   *al.* Mapping Alterations of the Functional Structure of the Cerebral Cortex in
689   Obsessive-Compulsive Disorder. *Cereb Cortex* 2019; **29**(11)**:** 4753-4762.
690

691   47.   Johnston BA, Tolomeo S, Gradin V, Christmas D, Matthews K, Steele JD. Failure of
692   hippocampal deactivation during loss events in treatment-resistant depression. *Brain*
693   2015; **138**(Pt 9)**:** 2766-2776.
694

695   48.   Kumar P, Waiter G, Ahearn T, Milders M, Reid I, Steele JD. Abnormal temporal
696   difference reward-learning signals in major depression. *Brain* 2008; **131**(Pt 8)**:** 2084-
697   2093.
698

699   49.   Zhu X, Wang X, Xiao J, Liao J, Zhong M, Wang W *et al.* Evidence of a dissociation
700   pattern in resting-state default mode network connectivity in first-episode, treatment-
701   naive major depression patients. *Biol Psychiatry* 2012; **71**(7)**:** 611-617.
702

703   50.   Harel EV, Tennyson RL, Fava M, Bar M. Linking major depression and the neural
704   substrates of associative processing. *Cogn Affect Behav Neurosci* 2016; **16**(6)**:** 1017-
705   1026.
706

707   51.   Andrews-Hanna JR, Reidler JS, Sepulcre J, Poulin R, Buckner RL. Functional-
708   anatomic fractionation of the brain's default network. *Neuron* 2010; **65**(4)**:** 550-562.
709

710 52. Miller AM, Vedder LC, Law LM, Smith DM. Cues, context, and long-term memory:
711 the role of the retrosplenial cortex in spatial cognition. *Front Hum Neurosci* 2014; **8:**
712 586.
713

714 53. Andrews-Hanna JR, Smallwood J, Spreng RN. The default network and self-
715 generated thought: component processes, dynamic control, and clinical relevance.
716 *Ann N Y Acad Sci* 2014; **1316**(1)**:** 29-52.
717

718 54. Chrastil ER. Heterogeneity in Human Retrosplenial Cortex: A Review of Function
719 and Connectivity. *Behav Neurosci* 2018; **132**(5)**:** 317-338.
720

721 55. Kaboodvand N, Backman L, Nyberg L, Salami A. The retrosplenial cortex: A
722 memory gateway between the cortical default mode network and the medial temporal
723 lobe. *Human Brain Mapping* 2018; **39**(5)**:** 2020-2034.
724

725 56. Davey CG, Harrison BJ. The self on its axis: a framework for understanding
726 depression. *Transl Psychiatry* 2022; **12**(1)**:** 23.
727

728 57. Jaworska N, Yang XR, Knott V, MacQueen G. A review of fMRI studies during
729 visual emotive processing in major depressive disorder. *World J Biol Psychiatry*
730 2015; **16**(7)**:** 448-471.
731

732 58. Disner SG, Beevers CG, Haigh EA, Beck AT. Neural mechanisms of the cognitive
733 model of depression. *Nat Rev Neurosci* 2011; **12**(8)**:** 467-477.
734

735 59. Harro J, Kanarik M, Kaart T, Matrov D, Koiv K, Mallo T *et al.* Revealing the cerebral
736 regions and networks mediating vulnerability to depression: oxidative metabolism
737 mapping of rat brain. *Behav Brain Res* 2014; **267:** 83-94.
738

739 60. Fanselow MS, Dong HW. Are the dorsal and ventral hippocampus functionally
740 distinct structures? *Neuron* 2010; **65**(1)**:** 7-19.
741

742 61. Ren LY, Meyer MAA, Grayson VS, Gao P, Guedea AL, Radulovic J. Stress-induced
743 generalization of negative memories is mediated by an extended hippocampal circuit.
744 *Neuropsychopharmacology* 2022; **47**(6)**:** 1280-1280.
745

746 62. Freund TF, Buzsaki G. Interneurons of the hippocampus. *Hippocampus* 1996; **6**(4)**:**
747 347-470.
748

749 63. Olbrich HG, Braak H. Ratio of pyramidal cells versus non-pyramidal cells in sector
750 CA1 of the human Ammon's horn. *Anat Embryol (Berl)* 1985; **173**(1)**:** 105-110.
751

752 64. Tomar A, Polygalov D, Chattarji S, McHugh TJ. Stress enhances hippocampal
753 neuronal synchrony and alters ripple-spike interaction. *Neurobiol Stress* 2021; **14:**
754 100327.
755

756 65. Corcoran KA, Yamawaki N, Leaderbrand K, Radulovic J. Role of retrosplenial cortex
757 in processing stress-related context memories. *Behav Neurosci* 2018; **132**(5)**:** 388-
758 395.
759

760   66.   Huang Y, Coupland NJ, Lebel RM, Carter R, Seres P, Wilman AH *et al.* Structural
761   changes in hippocampal subfields in major depressive disorder: a high-field magnetic
762   resonance imaging study. *Biol Psychiatry* 2013; **74**(1)**:** 62-68.
763

764   67.   Malykhin NV, Coupland NJ. Hippocampal neuroplasticity in major depressive
765   disorder. *Neuroscience* 2015; **309:** 200-213.
766

767   68.   MacQueen G, Frodl T. The hippocampus in major depression: evidence for the
768   convergence of the bench and bedside in psychiatric research? *Mol Psychiatry* 2011;
769   **16**(3)**:** 252-264.
770

771   69.   Avery JA, Drevets WC, Moseman SE, Bodurka J, Barcalow JC, Simmons WK. Major
772   Depressive Disorder Is Associated With Abnormal Interoceptive Activity and
773   Functional Connectivity in the Insula. *Biol Psychiat* 2014; **76**(3)**:** 258-266.
774

775   70.   DeVille DC, Kerr KL, Avery JA, Burrows K, Bodurka J, Feinstein JS *et al.* The
776   Neural Bases of Interoceptive Encoding and Recall in Healthy Adults and Adults
777   With Depression. *Biol Psychiat-Cogn N* 2018; **3**(6)**:** 546-554.
778

779   71.   Craig AD. How do you feel? Interoception: the sense of the physiological condition
780   of the body. *Nature reviews neuroscience* 2002; **3**(8)**:** 655-666.
781

782   72.   Kuehn E, Mueller K, Lohmann G, Schuetz-Bosbach S. Interoceptive awareness
783   changes the posterior insula functional connectivity profile. *Brain Struct Funct* 2016;
784   **221**(3)**:** 1555-1571.
785

786   73.   Gu XS, Hof PR, Friston KJ, Fan J. Anterior insular cortex and emotional awareness.
787   *Journal of Comparative Neurology* 2013; **521**(15)**:** 3371-3388.
788

789   74.   Craig AD. How do you feel—now? The anterior insula and human awareness. *Nature*
790   *reviews neuroscience* 2009; **10**(1)**:** 59-70.
791

792   75.   Mesulam MM, Mufson EJ. Insula of the Old-World Monkey .1. Architectonics in the
793   Insulo-Orbito-Temporal Component of the Paralimbic Brain. *Journal of Comparative*
794   *Neurology* 1982; **212**(1)**:** 1-22.
795

796   76.   Mufson EJ, Mesulam MM. Insula of the old world monkey. II: Afferent cortical input
797   and comments on the claustrum. *J Comp Neurol* 1982; **212**(1)**:** 23-37.
798

799   77.   Centanni SW, Janes AC, Haggerty DL, Atwood B, Hopf FW. Better living through
800   understanding the insula: Why subregions can make all the difference.
801   *Neuropharmacology* 2021; **198**.
802

803   78.   Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB. A link between the systems:
804   functional differentiation and integration within the human insula revealed by meta-
805   analysis. *Brain Structure & Function* 2010; **214**(5-6)**:** 519-534.
806

807   79.   Schulz SM. Neural correlates of heart-focused interoception: a functional magnetic
808   resonance imaging meta-analysis. *Philos T R Soc B* 2016; **371**(1708).
809

810   80.   Barrett LF, Simmons WK. Interoceptive predictions in the brain. *Nat Rev Neurosci*
811   2015; **16**(7)**:** 419-429.
812
813   81.   Nord CL, Lawson RP, Dalgleish T. Disrupted Dorsal Mid-Insula Activation During
814   Interoception Across Psychiatric Disorders. *Am J Psychiat* 2021; **178**(8)**:** 761-770.
815
816   82.   Namkung H, Kim SH, Sawa A. The Insula: An Underestimated Brain Area in Clinical
817   Neuroscience, Psychiatry, and Neurology: (Trends in Neuroscience 40, 200-207,
818   2017). *Trends Neurosci* 2018; **41**(8)**:** 551-554.
819
820   83.   Paulus MP, Stein MB. Interoception in anxiety and depression. *Brain Struct Funct*
821   2010; **214**(5-6)**:** 451-463.
822
823   84.   Schafer JO, Naumann E, Holmes EA, Tuschen-Caffier B, Samson AC. Emotion
824   Regulation Strategies in Depressive and Anxiety Symptoms in Youth: A Meta-
825   Analytic Review. *J Youth Adolesc* 2017; **46**(2)**:** 261-276.
826
827   85.   Joormann J, Gotlib IH. Emotion regulation in depression: relation to cognitive
828   inhibition. *Cogn Emot* 2010; **24**(2)**:** 281-298.
829
830   86.   Dryman MT, Heimberg RG. Emotion regulation in social anxiety and depression: a
831   systematic review of expressive suppression and cognitive reappraisal. *Clin Psychol*
832   *Rev* 2018; **65:** 17-42.
833
834   87.   Larsen JK, Vermulst AA, Eisinga R, English T, Gross JJ, Hofman E *et al.* Social
835   Coping by Masking? Parental Support and Peer Victimization as Mediators of the
836   Relationship Between Depressive Symptoms and Expressive Suppression in
837   Adolescents. *J Youth Adolescence* 2012; **41**(12)**:** 1628-1642.
838
839   88.   Brummer L, Stopa L, Bucks R. The influence of age on emotion regulation strategies
840   and psychological distress. *Behav Cogn Psychother* 2014; **42**(6)**:** 668-681.
841
842