***De novo* genome assemblies from two Indigenous Americans from Arizona identify new**

polymorphisms in non-reference sequences

Çiğdem Köroğlu[1], Peng Chen[1], Michael Traurig[1], Serdar Altok[2], Clifton Bogardus[1], Leslie J

Baier[1]

[1]Diabetes Molecular Genetics Section, Phoenix Epidemiology and Clinical Research Branch,

National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health,

Phoenix, AZ, USA

[2]Independent Researcher, Phoenix, AZ, USA

Corresponding Author:

Leslie J. Baier, PhD

445 North 5th Street, Phoenix, AZ 85004, USA

Phone: +1 602 440 6589 (office); Fax: +1 602 253 4140

E-mail: lbaier@phx.niddk.nih.gov

Short title: Analysis of non-reference sequences in Indigenous American genomes

Word count: 4,150 (excluding abstract, author summary, and references)

ABSTRACT

There is a collective push to diversify human genetic studies by including underrepresented

populations. However, analyzing DNA sequence reads involves the initial step of aligning the

reads to the GRCh38/hg38 reference genome which is inadequate for non-European ancestries. To

help address this issue, we created a modified hg38 reference map using *de novo* sequence

assemblies from Indigenous Americans living in Arizona (IAZ). Using HiFi SMRT long-read

sequencing technology, we generated *de novo* genome assemblies for one female and one male

IAZ individual. Each assembly included ~17 Mb of DNA sequence not present (non-reference

sequence; NRS) in hg38, which consists mostly of repeat elements. Forty NRSs totaling 240 kb

were uniquely anchored to the hg38 primary assembly generating a modified hg38-NRS reference

genome. DNA sequence alignment and variant calling were then conducted with WGS sequencing

data from 387 IAZ cohorts using both the hg38 and modified hg38-NRS reference maps. Variant

calling with the hg38-NRS map identified ~50,000 single nucleotide variants present in at least

5% of the WGS samples which were not detected with the hg38 reference map. We also directly

assessed the NRSs positioned within genes. Seventeen NRSs anchored to regions including an

identical 187 bp NRS found in both de novo assemblies. The NRS is located in *HCN2* 79 bp

downstream of exon 3 and contains several putative transcriptional regulatory elements.

Genotyping of the *HCN2*-NRS revealed that the insertion is enriched in IAZ (MAF = 0.45)

compared to Caucasians (MAF = 0.15) and African Americans (MAF = 0.03). This study shows

that inclusion of population-specific NRSs can dramatically change the variant profile in an under-

represented ethnic groups and thereby lead to the discovery of previously missed common

variations.

AUTHOR SUMMARY

GRCh38/hg38 reference genome has been the standard reference for large-scale human genetics studies. However, it does not adequately represent sequences of non-European ancestry. In this study, using long-read sequencing technology, we constructed *de novo* sequence assemblies from two Indigenous Americans from Arizona. We then compared the *de novo* assemblies to the hg38 reference genome to identify non-reference sequences (NRSs). We integrated these NRSs into our whole-genome sequencing (WGS) variant calling pipeline to improve read alignment and variant detection. We also directly assessed the NRSs positioned within genes. Inclusion of population-specific NRSs dramatically changed the variant profile of our study group with under-represented ethnicity, revealing common variation not detected by our previous population-level WGS and genotyping studies.

INTRODUCTION

The lack of diversity in human genetic studies is of growing concern, particularly as personalized medicine is becoming a reality. The majority of genomics data, approximately 80 percent, predominantly represents individuals of European ancestry, highlighting a significant underrepresentation of other populations [1]. This lack of diversity has had adverse consequences in genetic testing and variant interpretation, particularly for patients of African and Asian ancestry [2,3]. Recent large-scale genome studies, such as the All of Us initiative [4], the Human Heredity and Health in Africa (H3Africa) project [5], and the GenomeAsia100K project [6], have made significant efforts to address this disparity by actively increasing the representation of diverse populations. However, aside from the challenges associated with sample collection for sequencing, these endeavors face two main limitations when generating genomic data for underrepresented

populations. One limitation is due to short-read sequencing being the main method in large

population studies. While short-reads offers valuable insights through comparisons of allele

frequencies and functional annotations, they struggle to capture rare variants and accurately

characterize structural variants in isolated populations [7-9]. This restricts the comprehensive

understanding of genetic diversity and hampers the identification of variants associated with

diseases. The second limitation stems from the reliance on a human genome reference during DNA

sequence read analysis. The current reference map, the GRCh38/hg38 reference genome, primarily

comprises sequences derived from a single DNA donor of African-European origin, accounting

for approximately 70% of the primary assembly sequences [10]. Consequently, the current

reference map still exhibits limitations in adequately representing the genetic diversity found

within diverse populations. It is important to recognize that individuals may carry unique

sequences that are not represented in the reference map, thereby harboring genomic regions with

potential disease associations and other phenotypic relevance. Therefore, it has been imperative

for the genomic research in isolated populations to move beyond the traditional short-read

sequencing and linear reference genome paradigm.

Long-read sequencing technologies have boosted our power in resolving complex regions of the

genome, detecting structural variants, and assembling complete chromosomes [11-13]. In the

recent years, these technologies have advanced in accuracy and yield, enabling variant detection

at a large scale, with recent studies showcasing their potential in population-scale analyses [13,14].

Significant steps have also been taken to address the issue of the low ancestral diversity in the

human genome reference. The Human Pangenome Reference Consortium has been working on

constructing a reference that encompasses diverse genomes, and they have published the first draft

of the pangenome, consisting of 47 genome assemblies from a diverse cohort [15]. However, there

are still challenges to overcome in growing and refining this reference, both in sampling and in

computational analysis. The current sampling efforts do not include Indigenous American or

Aboriginal peoples (https://humanpangenome.org/samples/). Indigenous American people are

genetically distinct compared to other populations in the United States, with a significant variation

within the group itself [16]. Principal component analysis from a previous study including three

ancestral populations from the 1000 Genomes database (European, East Asian, and African

ancestries) and our Indigenous American population from Arizona (IAZ) showed a distinct cluster

for IAZ on the principal component space [17]. Therefore, exploratory genomics studies using

novel methods are warrented in indigenous populations to uncover previously unrecognized

genetic variations and capture novel disease associations.

In this study, we aimed to improve identification of genetic variations that have not been previously

studied among IAZ population. We utilized *de novo* genome assemblies to identify non-reference

segments (NRSs) in this cohort. These NRSs represent novel genomic segments not present in the

existing reference map. By analyzing these segments, both directly and by incorporating them into

our whole-genome sequencing (WGS) variant calling pipeline, we sought to uncover previously

unrecognized polymorphisms which can be assessed for a potential role in metabolic disease

enriched in   this underrepresented ethnic group.

RESULTS

***De novo* IAZ genome assemblies**

DNA samples from one female (sample 1) and one male (Sample 2) Indigenous American from

Arizona (IAZ) were sequenced using the PacBio HiFi SMRT sequencing platform. Summary

reports for the raw PacBio sequencing data are shown in S1 Table. For both samples, the SMRT

sequencing reactions produced approximately 64 Gb of high-fidelity subreads (>Q20) with a mean

length of ~12 kb, which were used to generate the *de novo* genome assemblies. Primary contigs

with high contiguity were generated using HiFiasm assembler (Table 1). Assembly 1 (female

IAZ) and assembly 2 (male IAZ) yielded a total genome length of 3.05 and 3.07 Gb, respectively,

indicating high assembly quality regarding genome completeness when compared to the 2.96 Gb

GRCh38/hg38 reference genome excluding gaps. Subsequent rounds of MUMmer alignment

identified 19 Mb and 17 Mb of DNA sequence for assembly 1 and assembly 2, respectively, that

could not be aligned to the hg38 reference genome. These non-reference sequences (NRSs) have

a minimum length of 100 bps and a hg38 sequence identity < 80%.

125
126   **Table 1. Summary statistics for the IAZ *de novo* assemblies.**

Contiguity metric	Assembly 1	Assembly 2
N50	48.04 Mb	36.34 Mb
NG50	41.32 Mb	32.19 Mb
Area under the N curve	46.30 Mb	39.89 Mb

Assembly 1, female IAZ individual. Assembly 2, male IAZ individual. N50 is the length
of the shortest assembled contig at 50% of the total assembly length. NG50 is the length
of the shortest assembled contig at 50% of the total genome length. For NG50 and area
under the N curve, 3.3 Gb of genome size is assumed.

NRS size distribution for the IAZ *de novo* assemblies

The two IAZ *de novo* assemblies had similar numbers of NRSs and size distribution patterns with

a total length of 19.3 Mb for assembly 1 (n = 2,177 NRSs) and 16.34 Mb for assembly 2 (n = 1908

NRSs) (Fig 1A-F). For both assemblies, most of the NRSs are 10 kb and shorter but only accounted

for 14.4% and 13.4% of the total lengths for assembly 1 and assembly 2 respectively (Fig 1A and

D). There were 130 NRSs for both assemblies with sizes greater than 10 kb–100 kb with medium

sizes of 20.5 kb for assembly 1 and 22.7 kb for assembly 2 (Fig 1B and E). 93 NRSs are larger

than 100 kb (assembly 1, n = 50; assembly 2 n = 43), including 3 outlier NRSs larger than 500 kb

(Fig 1C and F).

Identification of repeat and transposable elements

Screening the NRSs for repeat and transposable elements using RepeatMasker revealed that

approximately 95% of the NRSs in both assemblies primarily consisted of satellites, simple

repeats, and a small fraction of LINE and SINE transposable elements (S1 Fig). The three largest

NRSs with lengths of 637 kb, 1.88 Mb (assembly 1), and 1.47 Mb (assembly 2) were individually

analyzed for their repeat content. They consisted almost entirely of satellites and simple repeats

(ranging from 99.7% to 99.9%) rendering their alignment to chromosomes challenging.

Mapping the NRSs to the GRCh38/hg38 reference genome

NRSs totaling 241 kb could be placed into 40 distinct locations within the primary scaffold of

GRCh38/hg38. Only 4 of the NRSs totaling 7.6 kb were placed into centromeric regions

suggesting that there was no enrichment of centromeric DNA which is in contrast to long-read

sequencing projects involving other populations [18, 19]. However, when the alignments for all of

the NRSs with at least one anchored flanking site were analyzed, the percentage of NRSs aligning

to centromeric or telomeric regions was ~30%. The 233.4 kb non-centromeric NRSs includes a

large 215 kb insertion aligning to chrY and ~18.4 kb of NRS anchoring to 13 autosomal

chromosomes, chrX, and 2 pseudoautosomal regions on chrY. Among the 36 non-centromeric

NRSs, 17 are located within intronic regions of 13 protein coding and 4 non-protein-coding genes.

Three NRSs were present in both assemblies and only 1, *HCN2*-NRS, was identical in size and

therefore was selected for further characterization (Table 2). The *HCN2*-NRS in not present in the

reference genomes, including the T2T-CHM13 map, suggesting that the *HCN2*-NRS may not exist

in the DNA samples used for reference genome construction. *HCN2*-NRS is located 79 bp

downstream of exon 3 and replaces 228 bp of the reference sequence including a small portion of

a predicted cis-regulatory element (CRE). Individuals who carry the 187 bp HCN2-NRS also have

an adjacent 100 bp deletion (S2 Fig). The *HCN2*-NRS matches (100% identity) with 187 bps

downstream of exon 4 spanning a putative CRE containing various transcription factor binding

sites suggesting that *HCN2*-NRS may have functional importance (Fig 2, S2 Fig).

Table 2. Non-reference sequences anchoring within genes.

Gene	Insertion region[a]	NRS size (bps)	Both assemblies
LINC02270	Chr4:12237666–12237666	139	No
F11-AS1	Chr4:186435038–186435150	4714	No
BTNL9	Chr5:181046016–181046016	386	No
DYNC2I1	Chr7:158915724–158915724	679	No
MROH5	Chr8:141492602–141492602	1722	No
GTPBP4	Chr10:1011004–1011004	341	No
ART1	Chr11:3653879–3653937	877	Yes[b]
PDGFD	Chr11:104078565–104078565	579	No
LOC728715	Chr12:9404585–9404585	836	Yes[b]
GNPTAB	Chr12:101747637–101747644	658	No
MPG	Chr16:81854–81857	344	No
CACNA1H	Chr16:1186044–1186087	466	No
HCN2	Chr19:605301–605528	187	Yes
ADGRG2	ChrX:19074564–19074564	429	No
RGN	ChrX:47089179–47089179	1127	No
CRLF2	ChrY:1198256–1198271[c]	287	No
P2RY8	ChrY:1496850–1496850[c]	136	No

169 NRS, non-reference sequence. [a]Position based on GRCh38/hg38. Indicated intervals are replaced by NRSs.
170 [b]NRS sizes slightly different between the two assemblies. [c]Pseudoautosomal region.
171

Population frequency and association analyses for *HCN2*-NRS

To confirm the presence of the *HCN2*-NRS, the region encompassing the putative NRS was PCR

amplified and sequenced in the two IAZ individuals used for the *de novo* genome assembly along

with some of their family members. Of the 10 individuals sequenced, 1 was heterozygous and 9

were homozygous for *HCN2*-NRS suggesting that the NRS segment was common in the IAZ

population. To determine the frequency of *HCN2*-NRS in IAZ, ~3,400 IAZ individuals were

genotyped using a custom designed TaqMan probe that detects the presence of the NRS. The

HCN2-NRS is the minor allele with a frequency of 0.45. Genotyping was repeated for several

individuals by PCR to verify that the custom probe was working properly (S3 Fig). For

comparison, the *HCN2*-NRS was also genotyped in the African Ancestry in Southwest USA

(ASW) and British from England and Scotland (GBR) DNA panels from the 1000 genomes

project. *HCN2*-NRS was not as common in these two ethnic groups with frequencies of 0.03 and

0.15 in the ASW and GBR populations, respectively. Despite being enriched in IAZ, the *HCN2*-

NRS allele was not significantly associated with any obesity or type 2 diabetes-related traits which

are highly prevalent in this population (data not shown).

Structural variation in the *de novo* assemblies

Structural variants (SV) were analyzed directly from the *de novo* assemblies using Assemblytics,

which distinguishes between insertions/deletions and contractions/expansions of repeat elements

[20]. A total of 12,265 and 12,133 SVs were detected for Sample 1 and Sample 2, respectively,

which is similar to an Assemblytics report for a human reference assembly (11,206 SVs) [20] but

lower than the number of SVs reported in other studies that used assembly-based or long-read

mapping approaches [18, 21]. The lower number of SVs in our study can be attributed to the

stringency of our SV detection method. For both samples, the larger SVs (500bp–10,000bp) are

predominately composed of tandem expansions (S4 Fig). The distribution graphs also show peaks

of SV counts at around 300 bp and 6,000 bp (S4 Fig), a size distribution pattern similar to a

previous report [18].

Variant calling using the amended hg38-NRS map

Variant calling using our short-read WGS data generated for 387 IAZ individuals and the amended

hg38-NRS map identified 11,130,171 variants. Using the new hg38-NRS map identified 161,502

gained single nucleotide variants (SNVs) including 695 missense substitutions that were not

captured when variant calling was performed using the hg38 reference map (Fig 3). Additionally,

180,940 SNVs that were captured using the hg38 map were not detected (lost) with the hg38-NRS

map (Fig 3). Most of the gained SNVs (n = 111,976) were low-frequency variants; while the

remaining gained SNVs (n = 49,526) were detected in $\geq$ 5% of the WGS samples. The accuracy

of our variant calling with the hg38-NRS map was assessed by sequencing 25 gained SNVs (15

reported and 10 not reported in dbSNP). For the 15 dbSNPs, 13 of the gained SNVs were verified

and for the 10 non-dbSNPs, 6 were validated. Most of the non-dbSNPs SNVs reside in repetitive

regions of the genome which may account for the reduced accuracy.

DISCUSSION

In this study, our goal was to enhance the identification of genetic variations that remain largely

unexplored within an Indigenous American cohort from Arizona. We focused on identifying NRSs

in *de novo* genome assemblies constructed for the two participants from the IAZ cohort. The newly

identified NRSs signify novel genomic regions not found in the current GRCh38/hg38 reference

map. By incorporating these NRS segments into our WGS variant calling pipeline, we aimed to

uncover previously undetected polymorphisms.

The *de novo* assemblies were constructed using long-read PacBio HiFi sequencing which produces

highly accurate long-read (>10 kb) sequencing data [22]. The sizes for the *de novo* genome

assemblies were similar to hg38 and exhibited high contiguity with a mean N50 value of 42.2 Mb.

We identified > 10,000 SVs per *de novo* genome assembly which is in line with the expectations

for long-read sequencing [23]. This assembly-based approach for SV detection proved particularly

advantageous for capturing variants of substantial size [24], and the categorization facilitated by

Assemblytics allowed us to differentiate between SVs occurring in repetitive regions, such as

tandem expansions and repeat contractions, and those in non-repetitive regions

(insertions/deletions). It is worth noting that the largest insertions and deletions (> 500 kb) often

resided in regions characterized by high repeat content. We discovered 220 shared deletions (>

500 kb) in both *de novo* assemblies with 90 SVs located within genes (data not shown). The

application of a high-throughput genotyping approach with custom CNV probes could potentially

reveal commonly occurring deletions and uncover potential associations with diseases. However,

probe designs for these regions, predominantly repetitive in content, would pose a challenge.

The detected NRSs in two assemblies were found to be highly enriched in repeat sequences,

especially satellites and simple repeats. This high repeat content was consistent with the previous

literature on non-reference genomic parts [21, 25], and rendered the precise placement of NRSs

into chromosomes challenging. We identified 1.5 Mb of non-repetitive NRS per assembly and 241

kb was anchored to specific chromosomal regions in the primary assembly of hg38. The largest

NRS (215 kb) was anchored to chrY. This large NRS is probably a result of chrY being incomplete

in the current hg38 map due to its complex repeat structure [26].

Several of the NRSs anchored to intronic regions and we focused on a particular NRS in *HCN2*.

HCN2-NRS was detected in both de novo assemblies and genotyping showed that the *HCN2*-NRS

allele is common (MAF = 0.45) in the IAZ population. We also determined that *HCN2*-NRS is

present in Caucasians and African Americans but at lower frequencies (MAF = 0.15 and 0.03,

respectively).

The *HCN2*-NRS matches (100% identity) with a region located downstream of exon 4 that

encompasses a putative *cis*-regulatory element suggesting that individuals carrying the *HCN2*-

NRS may have additional transcription factor binding sites which may potentially influence gene

expression. *HCN2* encodes for a voltage-gated cation channel predominantly expressed in the heart

and the brain and plays a crucial role in migraine, inflammatory, and neuropathic pain including

diabetic neuropathy [27-29]. HCN2 ion channels initiate neuropathic pain by modulating action

potential firing in nociceptor neurons involved in sensing pain intensity [30].

Compared to Caucasians, chronic widespread and localized musculoskeletal pain disorders are

uncommon among IAZ [31], however, IAZ have high rates of diabetic neuropathy and pain from

rheumatoid arthritis [32, 33]. It is possible that the common *HCN2*-NRS allele might modestly

contribute to diabetic neuropathy since there are multiple rare intronic variants with nominal

associations with diabetic neuropathy reported in the T2D Knowledge Portal

(https://t2d.hugeamp.org/). However, the current study has no data on diabetic neuropathy pain to

address this hypothesis.

Our validation assessment for "gained" variants, namely the subset of genomic variants that were

only called using hg38-NRS as the reference, using Sanger sequencing underlines the reliability

of our WGS data analysis with the extended reference map. However, the calls arising from

repetitive regions provided lower rates of true positives. Further research is needed to understand

the functional significance of the gained variants and potential contributions to diseases among

IAZ population.

The overall findings we presented have expanded our understanding of genetic diversity within

Indigenous American populations. The presence of unique genetic elements and variants

unrepresented in existing reference maps demonstrates the importance of delving into the

genomics of underrepresented populations. By combining long-read sequencing and novel NRS

identification methods, we were able to uncover previously elusive genetic variations. This study

not only highlights the value of studying isolated populations but also emphasizes the need for

diverse and comprehensive reference maps to accurately capture the complexity of human genetic

diversity. Our work contributes to the broader goal of enhancing genomic research's inclusivity

and represents a step toward more precise understanding of genetic factors underlying health

disparities in underrepresented populations.

METHODS

Long-read sequencing and *de novo* assembly

Two DNA samples isolated from one female and one male individual were sequenced using the

PacBio HiFi SMRT sequencing platform at the DNA Sequencing Center at Brigham Young

University (Provo, UT). SMRT bell adapted DNA libraries were size selected and ~15 kb fractions

were run on 3 SMRT cells (8M trays) for a duration of 30 hours for each sample. Bioinformatic

analysis for the construction of the IAZ *de novo* assemblies was done by the DNA Sequencing

Center at Brigham Young University. Selected quality reads were assembled using Hifiasm

version 0.11 (r302). Continuity of the reads and other summary statistics were calculated using the

caln50 package (https://github.com/lh3/calN50).

Detection of non-reference sequences and structural variants in the *de novo* assemblies

All computational analyses were run on the high-performance computing environment NIH HPC

Biowulf cluster (http://hpc.nih.gov). Visualization was performed on RStudio using ggplot2

package of R (version 4.3.1).

The *de novo* assemblies were compared to the Genome Reference Consortium Human Build 38

patch release 13 (GRCh38.p13/hg38) using the alignment script of NUCmer from the MUMmer

package (version 4.0.0beta2). Each assembly was aligned to GRCh38.p13/hg38 with the

parameters --maxmatch -l 150 -c 400.   The resulting alignments were filtered to retain only the

single best alignment for each contig using the delta-filter -q command. The show-coords tool was

then used to extract the coordinates and percent identity of the aligned contigs. To identify the

contigs that failed to align to hg38 (< 80% sequence identity), a custom script (available at

https://github.com/cigdemkoroglu/NRS/blob/main/unaligned_contigs.R) was used to parse the

output of the alignment and extract the unaligned contigs and contigs with low identity. Bedtools

version 2.29.2 was employed to generate fasta files containing only the unaligned portions of the

assemblies (NRSs). To improve the alignment sensitivity and capture more contigs, the alignment

workflow was repeated on the initial output of NRSs with relaxed NUCmer settings (--maxmatch

-l 100 -c 200). This allowed for a more permissive alignment approach to increase the number of

contigs that could be aligned to the reference. Final NRSs were analyzed for repetitive sequences

using RepeatMasker and Assemblytics was used to detect structural variants > 50 bp

(http://assemblytics.com/).

310   Anchoring the NRSs to the GRCh38.p13/hg38 primary assembly

For each NRS, 100 bp of flanking sequences either side of the NRSs were aligned to the nucleotide

collection database using the blastn. Contigs from the *de novo* assemblies with both flanking sites

anchoring to the primary assembly chromosomes of hg38 were selected and appended to hg38 to

generate   an   extended   reference   genome   (hg38-NRS)   using   perEditor

(https://systemsbio.ucsd.edu/perEditor/). NRSs spanning a reference region longer than the NRS

itself and those aligning to the same region, indicating repeat expansions, were not incorporated

into the hg38-NRS reference genome. In cases where NRSs overlapped between both assemblies,

only the region from assembly 1 was included.

Variant calling using hg38 and hg38-NRS reference genomes

Whole genomes for 387 IAZ individuals were previously sequenced using Illumina short-read

chemistry (30× average coverage). Variant discovery was performed in parallel on whole genome

sequencing data from 387 IAZ samples using either hg38 or hg38-NRS as the reference. GATK

4.2.6 was used following the best practices workflow, which includes base quality score

recalibration, variant calling and joint genotyping by HaplotypeCaller and GenotypeGVCFs, and

variant quality score recalibration. Databases for the known variation in the GATK resource

bundle were position adjusted based on hg38-NRS using a custom lift-over script (available at

https://github.com/cigdemkoroglu/NRS/blob/main/lift_database.R). Monomorphic variants,

variants with a call rate <95%, variants with quality scores below the truth sensitivity level 99%

threshold, and variants with discordant genotypes between the duplicate pairs were removed from

the final vcf files using VCFtools.

Identification of gained single nucleotide variants (SNVs) in the hg38-NRS reference genome

To compare variant calling between hg38 and hg38-NRS reference genomes, ANNOVAR was

first used to annotate the variants identified with hg38. Next, to annotate the variants identified

with hg38-NRS and to ensure compatibility between the two reference maps, the hg38-NRS

variant dataset was lifted-over using custom scripts (https://github.com/cigdemkoroglu/NRS).

Lastly, to identify lost and gained SNVs, the hg38 and hg38-NRS annotated datasets were

compared using the LostGained tool (https://github.com/ cigdemkoroglu/NRS/blob/main

/LostGained.R). Among the gained variants, 15 missense variants that are reported in dbSNP but

were not detected by our in-house genotyping studies using the Affymetrix SNP Array 6.0 or our

whole-genome and whole-exome sequencing studies along with 10 randomly selected novel SNVs

were verified by sequencing.

Genotyping of the *HCN2*-NRS

Genotyping of the *HCN2*-NRS in the Indigenous Americans from Arizona (IAZ) was performed

using DNA isolated from 3,410 individuals who participated in a community-based longitudinal

study examining type 2 diabetes-related traits conducted by the National Institute of Diabetes and

Digestive and Kidney Diseases (NIDDK) in Phoenix, Arizona between the years 1965 and 2007

[34]. All participants in the study were self-reported full-heritage Indigenous American (all eight

great grandparents identified as IAZ). The *HCN2*-NRS was also genotyped in the African Ancestry

in Southwest USA (ASW) and British from England and Scotland (GBR) DNA panels from the

1000 genomes project. The ASW and GBR DNA samples were obtained from the NHGRI Sample

Repository for Human Genetic Research at the Coriell Institute for Medical Research: Repository

IDs MGP00015 and MGP00003, respectively. Genotyping of the *HCN2*-NRS was done using

custom designed TaqMan probes that detected the presence of either the reference sequence (RS)

or NRS. (Thermo Fisher Scientific, Waltham, MA). Statistical analyses were performed using the

software of the SAS Institute (Cary, NC, USA) as previously described [35].

Acknowledgments

We thank Chris Wiedrich for his help in genotype validation, Sayuko Kobes for running

association tests, and Paolo Piaggi for providing statistical consultation. This work was supported

by the Intramural Research Program of the NIDDK, NIH. We utilized the computational resources

of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

Author contributions:

Conceptualization: Cigdem Koroglu, Clifton Bogardus

Study design & data analysis: Cigdem Koroglu

Consultation regarding data analysis: Peng Chen

Supervision: Leslie J. Baier

Software: Serdar Altok, Peng Chen, Cigdem Koroglu

Computational troubleshooting: Cigdem Koroglu, Serdar Altok

Genotyping: Michael Traurig, Cigdem Koroglu

Visualization: Cigdem Koroglu, Michael Traurig

Writing -original draft: Cigdem Koroglu

Writing -review & editing: Leslie J. Baier, Cigdem Koroglu, Michael Traurig, Peng Chen, Clifton

Bogardus, Serdar Altok

Competing interests: The authors declare no competing interests.

References

1. Gurdasani D, Barroso I, Zeggini E, Sandhu MS. Genomics of disease risk in globally diverse populations. Nat Rev Genet. 2019; 20(9): 520-535.

2. Manrai AK, Funke BH, Rehm HL, Olesen MS, Maron BA, Szolovits P, et al. Genetic Misdiagnoses and the Potential for Health Disparities. N Engl J Med. 2016; 375(7): 655-665.

3. Popejoy AB, Fullerton SM. Genomics is failing on diversity. Nature. 2016; 538(7624): 161-164.

4. All of Us Research Program Investigators; Denny JC, Rutter JL, Goldstein DB, Philippakis
A, Smoller JW, et al. The "All of Us" Research Program. N Engl J Med. 2019; 381(7):
668-676.

5. GenomeAsia100K Consortium. The GenomeAsia 100K Project enables genetic
discoveries across Asia. Nature. 2019; 576(7785): 106-111.

6. Mulder N, Abimiku A, Adebamowo SN, de Vries J, Matimba A, Olowoyo P, et al.
H3Africa: current perspectives. Pharmgenomics Pers Med. 2018; 11: 59-66.

7. Chheda H, Palta P, Pirinen M, McCarthy S, Walter K, Koskinen S, et al. Whole-genome
view of the consequences of a population bottleneck using 2926 genome sequences from
Finland and United Kingdom. Eur J Hum Genet. 2017; 25(4): 477-484.

8. Beyter D, Ingimundardottir H, Oddsson A, Eggertsson HP, Bjornsson E, Jonsson H, et al.
Long-read sequencing of 3,622 Icelanders provides insight into the role of structural
variants in human diseases and other traits. Nat Genet. 2021; 53(6): 779-786.

9. Wu Z, Jiang Z, Li T, Xie C, Zhao L, Yang J, et al. Structural variants in the Chinese
population and their impact on phenotypes, diseases and population adaptation. Nat
Commun. 2021; 12(1): 6501.

10. Schneider VA, Graves-Lindsay T, Howe K, Bouk N, Chen HC, Kitts PA, et al. Evaluation
of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of
the reference assembly. Genome Res. 2017; 27(5): 849-864.

11. Chaisson MJ, Huddleston J, Dennis MY, Sudmant PH, Malig M, Hormozdiari F, et al.
Resolving the complexity of the human genome using single-molecule sequencing. Nature.
2015; 517(7536): 608-611.

12. Logsdon GA, Vollger MR, Eichler EE. Long-read human genome sequencing and its
applications. Nat Rev Genet. 2020; 21(10): 597-614.

13. Marx V. Method of the year: long-read sequencing. Nat Methods. 2023; 20(1): 6-11.

14. De Coster W, Weissensteiner MH, Sedlazeck FJ. Towards population-scale long-read
sequencing. Nat Rev Genet. 2021; 22(9): 572-587.

15. Liao WW, Asri M, Ebler J, Doerr D, Haukness M, Hickey G, et al. A draft human
pangenome reference. Nature. 2023; 617(7960): 312-324.

16. Redd AJ, Chamberlain VF, Kearney VF, Stover D, Karafet T, Calderon K, et al. Genetic
structure among 38 populations from the United States based on 11 U.S. core Y
chromosome STRs. J Forensic Sci. 2006; 51(3): 580-585.

17. Kim HI, Ye B, Gosalia N; Regeneron Genetics Center; Köroğlu Ç, Hanson RL, et al.
Characterization of Exome Variants and Their Metabolic Impact in 6,716 American
Indians from the Southwest US. Am J Hum Genet. 2020; 107(2): 251-264.

18. Ameur A, Che H, Martin M, Bunikis I, Dahlberg J, Höijer I, et al. De Novo Assembly of
Two Swedish Genomes Reveals Missing Segments from the Human GRCh38 Reference
and Improves Variant Calling of Population-Scale Sequencing Data. Genes (Basel). 2018;
9(10): 486.

19. Gao Y, Yang X, Chen H, Tan X, Yang Z, Deng L, et al. A pangenome reference of 36
Chinese populations. Nature. 2023; 619(7968): 112-121.

20. Nattestad M, Schatz MC. Assemblytics: a web analytics tool for the detection of variants
from an assembly. Bioinformatics. 2016; 32(19): 3021-3023.

21. bert P, Audano PA, Zhu Q, Rodriguez-Martin B, Porubsky D, Bonder MJ, et al. Haplotype-

resolved diverse human genomes and integrated analysis of structural variation. Science.

2021; 372(6537): eabf7117.

22. Hon T, Mars K, Young G, Tsai YC, Karalius JW, Landolin JM, et al. Highly accurate long-

read HiFi sequencing data for five complex genomes. Sci Data. 2020; 7(1): 399.

23. Ho SS, Urban AE, Mills RE. Structural variation in the sequencing era. Nat Rev Genet.

2020; 21(3): 171-189.

24. Mahmoud M, Gobet N, Cruz-Dávalos DI, Mounier N, Dessimoz C, Sedlazeck FJ.

Structural variant calling: the long and the short of it. Genome Biol. 2019; 20(1): 246.

25. Li R, Tian X, Yang P, Fan Y, Li M, Zheng H, et al. Recovery of non-reference sequences

missing from the human reference genome. BMC Genomics. 2019; 20(1): 746.

26. Skaletsky H, Kuroda-Kawaguchi T, Minx PJ, Cordum HS, Hillier L, Brown LG, et al. The

male-specific region of the human Y chromosome is a mosaic of discrete sequence classes.

Nature. 2003; 423(6942): 825-837.

27. Young GT, Emery EC, Mooney ER, Tsantoulas C, McNaughton PA. Inflammatory and

neuropathic pain are rapidly suppressed by peripheral block of hyperpolarisation-activated

cyclic nucleotide-gated ion channels. Pain. 2014; 155(9): 1708-1719.

28. Tsantoulas C, Ng A, Pinto L, Andreou AP, McNaughton PA. HCN2 Ion Channels Drive

Pain in Rodent Models of Migraine. J Neurosci. 2022; 42(40): 7513-7529.

29. Tsantoulas C, Laínez S, Wong S, Mehta I, Vilar B, McNaughton PA. Hyperpolarization-

activated cyclic nucleotide-gated 2 (HCN2) ion channels drive pain in mouse models of

diabetic neuropathy. Sci Transl Med. 2017; 9(409): eaam6072.

30. Emery EC, Young GT, Berrocoso EM, Chen L, McNaughton PA. HCN2 ion channels play a central role in inflammatory and neuropathic pain. Science. 2011; 333(6048): 1462-1466.

31. Jacobsson LT, Nagi DK, Pillemer SR, Knowler WC, Hanson RL, Pettitt DJ, et al. Low prevalences of chronic widespread pain and shoulder disorders among the Pima Indians. J Rheumatol. 1996; 23(5): 907-909.

32. Jaiswal M, Fufaa GD, Martin CL, Pop-Busui R, Nelson RG, Feldman EL. Burden of Diabetic Peripheral Neuropathy in Pima Indians With Type 2 Diabetes. Diabetes Care. 2016; 39(4): e63-64.

33. Del Puente A, Knowler WC, Pettitt DJ, Bennett PH. High incidence and prevalence of rheumatoid arthritis in Pima Indians. Am J Epidemiol. 1989; 129(6): 1170-1178.

34. Knowler WC, Bennett PH, Hamman RF, Miller M. Diabetes incidence and prevalence in Pima Indians: a 19-fold greater incidence than in Rochester, Minnesota. Am J Epidemiol. 1978; 108(6): 497-505.

35. Day SE, Traurig M, Kumar P, Piaggi P, Koroglu C, Kobes S, et al. Functional variants in cytochrome b5 type A (CYB5A) are enriched in Southwest American Indian individuals and associate with obesity. Obesity (Silver Spring). 2022; 30(2): 546-552.

Figure legends

Fig 1

Histograms depicting the length distribution of NRSs identified in the indigenous Sample 1 (panels A to C) and Sample 2 (panels D to F). Dashed lines represent the median values for each histogram. **A**. Small NRSs ranging from 100 bp and 10 kb, with a median length of 776 bp. **B**. NRSs sized between 10 kb and 100 kb, with a median size of 20.5 kb. **C**. NRSs longer than 100 kb are shown,

with a median size of 192 kb. Except for two outliers indicated by arrows, all NRSs in this group

are shorter than 500 kb. The arrows in this panel highlight a 637 kb NRS and a 1.88 Mb NRS, both

of which consist of simple repeats and microsatellites, and do not align to any hg38 contigs with

high sequence identity. **D**. Small NRSs with varying lengths between 100 bp and 10 kb and with

a median length of 617 bp. **E**. NRSs sized between 10 kb and 100 kb. The median size is 22.7 kb.

F. NRSs longer than 100 kb, with a median size of 187.8 kb. There is a single NRS, marked with

an arrow, exceeding 500 kb in length. This 1.47 Mb NRS comprises solely of simple repeats and

microsatellites, and does not anchor to any hg38 contigs.

Fig 2

HCN2-NRS and chr19-RS side by side alignment. Source: UCSC Genome Browser

(GRCh38/hg38).

Fig 3

Number of SNVs gained (light gray) and lost (dark gray) per chromosome for the cohort of 387

short-read WGS samples when analyzed with the hg38-NRS map compared to hg38 map.

Figure 1

Figure

Figure 2

Size	Identity	Chromosome	Strand	Start	End
187 bp	100%	19	+	608436	608622

```
HCN2-NRS 000001 caagtggcgcgctcagggcagagaccgcgtgatggggagcaggggaccct 000050
         >>>>>> |||||||||||||||||||||||||||||||||||||||||||||||||| >>>>>>
Chr19-RS 608436 caagtggcgcgctcagggcagagaccgcgtgatggggagcaggggaccct 608485

HCN2-NRS 000051 gggggtctgtggctgtgattaggaggagaaacggccggtggtgagaggtt 000100
         >>>>>> |||||||||||||||||||||||||||||||||||||||||||||||||| >>>>>>
Chr19-RS 608486 gggggtctgtggctgtgattaggaggagaaacggccggtggtgagaggtt 608535

HCN2-NRS 000101 ctctgaccctgtgacagggagggaggtccagagcctggagatggacgtgg 000150
         >>>>>> |||||||||||||||||||||||||||||||||||||||||||||||||| >>>>>>
Chr19-RS 608536 ctctgaccctgtgacagggagggaggtccagagcctggagatggacgtgg 608585

HCN2-NRS 000151 ggcgggggctggggagggggatgcaggctgggcaggaa 000187
         >>>>>> |||||||||||||||||||||||||||||||||||||| >>>>>>
Chr19-RS 608586 ggcgggggctggggagggggatgcaggctgggcaggaa 608622
```

Figure

Figure 3

Lost and Gained SNVs

Figure