Comparison of alternative solvents for *in situ* extraction of hydrocarbons from the colonial green alga *Botryococcus braunii* race B (Showa)

Takehiro A. Ozawa-Uyeda[a], Sebastian Overmans[b], Bárbara Bastos de Freitas[b], Edmundo Lozoya-Gloria[a]** and Kyle J. Lauersen[b]*

[a] Genetic Engineering Department, CINVESTAV-IPN Irapuato Unit, C.P. 36824 Irapuato, Gto., Mexico

[b] Bioengineering Program, Biological and Environmental Sciences and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

*Corresponding author: kyle.lauersen@kaust.edu.sa
**Co-corresponding author: edmundo.lozoya@cinvestav.mx

Abstract

The colony-forming, green microalga *Botryococcus braunii* secretes petroleum-like hydrocarbons, which enables the non-destructive continuous *in situ* extraction, 'milking', of these extracellular products during culture growth without cell lysis. This work compares the suitability of 15 different solvents, including alkanes, halogenated solvents, and green solvents, for *in situ* extraction of *B. braunii* race B (Showa strain) hydrocarbons after acclimation to moderate salinity stress. After 24 h of extraction, bio-based terpene 'green' solvents such as ɣ-terpinene showed the highest hydrocarbon recovery, around 10-fold greater than with conventional alkane solvents. Brominated alkanes and liquid perfluorocarbons (FCs) formed a lower phase to algal cultures rather than an upper phase as with other solvents, but only bromodecane effectively captured extracellular hydrocarbons similar to conventional alkane solvents. However, bromodecane and all green solvents were too toxic for two-phase continuous culture contact hydrocarbon milking, leading to 33–100% chlorophyll content loss. To overcome the biologically adverse effects of these solvents with suitable hydrocarbon recovery, future research should focus on their application in short-term extraction period 'milking' systems to minimize algal-solvent contact and enable continuous extraction.

Keywords:

Botryococcus braunii, microalgae, *in situ* extraction, hydrocarbons, green solvents

Introduction

The colony-forming green microalga *Botryococcus braunii* race B produces and secretes large amounts of liquid triterpene hydrocarbons, botryococcenes, during its exponential growth phase, which are stored in its extracellular matrix (ECM) (Largeau *et al.*, 1980; Suzuki *et al.*, 2013). Botryococcenes can be used to produce petroleum-equivalent fuels suitable for combustion engines (Jin *et al.*, 2016; Lozoya-Gloria *et al.*, 2019). In contrast to most other oleaginous microalgae, which accumulate oils within the cell, extracellular *B. braunii* hydrocarbons can be harvested without requiring cell lysis (Suzuki *et al.*, 2013; Ennaceri *et al.*, 2023). This trait makes it a suitable microalga for the process of non-destructive *in situ* extraction, termed 'milking', of lipophilic compounds. Milking can eliminate the steps of algal biomass downstream processing: harvesting, dewatering and cell disruption (Jackson *et al.*, 2017; Kleinert and Griehl 2022; Ennaceri *et al.*, 2023).

B. braunii typically exhibits slow growth rates compared to other photosynthetic microbes, which has limited its broader biotechnological exploitation (Lozoya-Gloria *et al.*, 2019). Additionally, the colony surface of *B. braunii* race B is covered with an ultrastructure called a retaining wall, which is decorated with a fibrillar sheath composed mainly of arabinose-galactose polysaccharides (Weiss *et al.*, 2012). This outer barrier hinders the entry of many organic solvents, which limits milking efficiencies (Furuhashi *et al.*, 2016b). One pretreatment that was found to improve the extraction rate of botryococcenes is cultivation of the alga in brackish medium (Furuhashi *et al.*, 2013; 2016a). It was proposed that increased salinity in the culture medium decreased the production or integrity of the

polysaccharides that block non-polar solvents and increases colony size and flotation

(Furuhashi *et al*., 2016a; 2016b).

Extraction of hydrocarbons from *B. braunii* has been tested with different types of

solvents, including traditional alkanes solvents such as hexane (C6) and heptane (C7),

partially polar solvents such as 1-octanol, and alternatives solvents such dimethyl ether

(DME) and switchable-polarity solvents (SPS) (Frenz *et al*., 1989; Samori *et al*., 2010;

Kanda *et al*., 2013). Short chain alkanes are reported to have high botryococcene

extraction efficiencies but low biocompatibility, resulting in rapid loss of cell viability after

long contact times (Kleinert and Griehl, 2021). Dodecane (C12) is biocompatible with *B.*

braunii cultures but has reduced hydrocarbon extractability compared to hexane (Mehta

et al., 2019). Alkane solvents have significant drawbacks that hinder scaling their

application to larger culture volumes like flammability and emulsion formation at the

culture-solvent interface (Lauersen, 2019).

It was recently reported that halogenated perfluorocarbon (FC) liquids can enable

milking of sesqui- and di-terpenoids from engineered algal cultures (Overmans and

Lauersen, 2022). These have not yet been tested on *B. braunii*. Green solvents are

another class of chemicals that are bio-based in origin and have gained popularity in

process chemistry due to their biological origins and/or biodegradability (Kumar *et al*.,

2017; De Jesus and Maciel Filho, 2020). Monoterpenes derived from essential oils such

as D-limonene, α-pinene, and p-cymene are considered 'green' solvents because they are

obtained from renewable feedstocks, like plants and crop wastes (Kumar *et al*., 2017).

Lipid extraction of microalgae and oilseeds using terpene green solvents was shown to

have equal or superior quality and yields than with hexane (De Jesus and Maciel Filho,

2020). In this study, we evaluated the suitability of halogenated and green solvents as

alternatives to conventional alkanes for botryococcene extraction from *B. braunii* race B

(Showa strain) cultures grown in brackish media.

Materials and Methods

Reagents

The solvents tested in this work were grouped into straight-chain alkanes,

halogenated solvents, and green solvents. Alkanes included n-hexane, n-decane, n-

dodecane (≥99%), and n-pentadecane, which were obtained from Acros Organics (Geel,

Belgium), VWR International (Fontenay-sous-Bois, France), Sigma-Aldrich (St. Louis,

USA) and Alfa Aesar (Heysham, England), respectively. Halogenated solvents included

brominated alkanes (bromododecane and bromododecane) and chlorinated alkane

(chlorodecane), acquired from Sigma Aldrich (St. Louis, USA). Additionally, two

perfluorocarbon liquids, hereafter referred to as FCs, were used in this work: the

perfluorinated ether FC-770 and the perfluorinated amine FC-3283, obtained from

Fluorochem (Glossop, UK) and Acros Organics (Geel, Belgium), respectively. The green

solvents included three monoterpenes (para-cymene (99%), gamma-terpinene, and

alpha-pinene) and three alcohols (1-heptanol, nerol, and nopol), all purchased from

Sigma-Aldrich (Taufkirchen, Germany). According to GlaxoSmithKline's (GSK) green

solvents guide for assessing green credentials of solvents, 1-heptanol scored highly

compared to other solvents, such as para-cymene and n-hexane (Alder *et al*., 2016). For

chlorophyll quantification, ethanol (96% vol) and methanol were purchased from VWR

International (Fontenay-sous-Bois, France).

Microalgal cultures and growth conditions

Batch cultures of *Botryococcus braunii* race B, Showa strain (Nonomura, 1988), were grown in 1-L Erlenmeyer flasks in Algem photobioreactors (Algenuity©, UK) at 25 °C under 130 µmol photons $m^{-2} s^{-1}$ with a 12h:12h light:dark cycle and sparged with a 7% CO_2 in air mixture. The gas mix was supplied periodically to cultures at 25 cc min^{-1} according to the pH control level set on the photobioreactor, with the online pH measured continuously using a calibrated pH probe (Broadley James, UK) to maintain a constant pH of 7.5 in the culture media. *B. braunii* was cultured in biological triplicates (n = 3) in three different media: modified Chu-13 medium as the standard freshwater medium based on the recipe in (Furuhashi *et al.*, 2013), modified Chu-13 medium with 37.5 mM NaCl, and modified Chu-13 medium with 5% Red Sea water (RSW).

To avoid a severe algal growth inhibition due to relatively high salinity (e.g. 150 mM NaCl in 1/4 seawater medium) as previously reported (Furuhashi *et al.*, 2013), the artificial seawater media were reduced to 3–4 g L^{-1} brackish salinity medium (Furuhashi *et al.*, 2016a; 2022). Culturing *B. braunii* race B (Showa strain) and *B. braunii* race A (Yamanaka strain) in brackish media with 3 and 4 g L^{-1} salinity, respectively, improved their hydrocarbon extractability without inhibiting algal growth compared to the algae cultured in Chu13 freshwater medium (Furuhashi *et al.*, 2016a; 2022). Instead of assuming that all the salt in the seawater is NaCl, the content of NaCl accounted for 57.6% of dissolved salt in the artificial seawater medium used by Furuhashi *et al.* (2016a). Therefore, 3 and 4 g L^{-1} salinity medium contained 30 and 40 mM NaCl, respectively. Brackish medium was prepared by mixing filtered (0.2 µm) and autoclaved RSW with modified Chu-13 medium. The concentration of total dissolved solids in RSW was previously determined to be

around 750 mM NaCl (Abdel-Aal *et al.*, 2015). To get a final NaCl concentration within the

range of 30–40 mM, similar to the most suitable concentrations used by Furuhashi *et al.*

(2016a; 2022), Chu-13 medium was adjusted to a final concentration of 5% RSW (37.5

mM NaCl). All cultures were grown in the Algem photobioreactors with a working volume

of 400 mL for 18 days before being recovered for further analyses.

Determination of growth parameters

Final algal biomass was determined gravimetrically by obtaining a pellet of 5 mL

algal culture after adding an equal volume of 96% ethanol into a pre-weighed glass tube,

mixed manually several times, and centrifuged at 3,000 × *g* for 1 min to sediment all algal

cells. The addition of ethanol helps to precipitate the floating layer of algal colonies, but

the incubation time should be as short as possible to minimize extraction of undesired

pigments. After discarding the ethanol, the remaining algal pellet was dried in an oven at

110 °C for 24 h and cooled for 30 min before weighing. The dry cell weight (DCW; in mg)

of the microalgae was obtained by subtracting the dry weight of the empty glass tube from

the loaded glass tube. The specific growth rate was calculated using the following

equation:

152
$$\mu = \frac{\ln(Xn/X_0)}{(t_n - t_0)}$$

where, μ is the specific growth rate (day^{-1}), Xn is the final biomass concentration (g/L) at

time t_n (days), and X_0 is the initial biomass concentration (g/L) at time t_0 (days).

Total chlorophyll content was obtained by adding 1 mL 96% ethanol to 1 mL culture

samples, mixed by inversion several times, and centrifuged at 11,000 × *g* for 1 min to

sediment all suspended algal cells. After discarding the ethanol, the pellet was

resuspended in 1 mL 100% methanol and incubated for 30 min at 45 ºC in the dark to

extract chlorophyll. The cell debris was pelleted by centrifugation at 11,000 $\times$ g for 1 min

while the supernatant was used to measure the absorbance at 666, 653, and 470 nm with

a spectrophotometer (TECAN Infinite M1000 PRO). Total chlorophyll content was

calculated according to Wellburn (1994). The values of total chlorophyll content were then

expressed on a dry weight basis (mg chlorophyll/ g DCW) using the DCW of the algal

samples.

Determination of hydrocarbon content

Algal pellet samples were collected by centrifugation as described above, and

DCW was determined. Hydrocarbons were extracted from the dried algal sample by

adding 5 mL of n-hexane and incubating for 30 min at room temperature with manual

shaking at 10 min intervals. The extraction solvent containing botryococcene

hydrocarbons, which turned yellowish-orange due to the carotenoids found within the

ECM (Eroglu and Melis, 2010), was separated from the algal residue by decantation and

collected in a separate glass flask. This process was repeated until the solvent phase

became colorless. All n-hexane extracts were combined and evaporated in an oven at

110 ºC overnight. Hydrocarbon amounts were calculated gravimetrically and expressed

as a percentage of biomass DCW. For positive identification of botryococcenes in

samples, the extracts were resuspended in 1 mL of n-hexane or n-dodecane, and then

transferred to gas chromatography (GC) vials (n = 3) and stored at -20 °C until further

analysis as described below.

Preliminary screening of solvents for two-phase 'milking' of *B. braunii* race B in low salinity culture medium

Two-phase living extractions (i.e., 'milking') of extracellular hydrocarbons were performed with alkanes, halogenated-, and green solvents by gently adding 250 µL of each respective solvent (25% v/v) onto 750 µL liquid cultures of *B. braunii* that were previously grown in brackish medium for 18 days. All two-phase cultures were grown for 24 h in duplicates in 2-mL microcentrifuge tubes on an orbital shaker at 120 rpm with a 12h:12h light:dark cycle at room temperature. The solvent layer was then collected following centrifugation at $4,500 \times g$ for 5 min to obtain two separate layers. From each sample, 150 µL of the solvent fraction was transferred into individual GC vials and stored at -20 °C until further analysis, as described below.

Alternative solvents for *in situ* extraction of hydrocarbons produced by *B. braunii* race B

The two most efficient extraction solvents of each group were selected from the initial screening. Two-phase living extractions of extracellular hydrocarbons were performed by gently adding 1 mL of each respective solvent onto 9 mL liquid cultures of *B. braunii* (10% v/v) that were previously grown in a brackish medium for 18 days. The two-phase cultures were cultivated for 24 h in triplicates in 50-mL Erlenmeyer flasks on an orbital shaker at 120 rpm with a 12h:12h light:dark cycle at room temperature. The solvent and algal samples were obtained at two different time points: 2 h and 24 h of cultivation. At each time point, the solvent layer was collected after centrifugation at 4,500 $\times g$ for 5 min. From each sample, 150 µL of the solvent fraction was transferred into separate GC vials and stored at -20 °C until further analysis.

Evaluation of hydrocarbon recovery rate

Hydrocarbon extractability from *B. braunii* culture was expressed as the hydrocarbon recovery rate, defined as the percentage of botryococcene hydrocarbons extracted from the wet algal samples using different types of solvents compared to the amount extracted from the dried algal sample using hexane as described above. The extracted botryococcene content was determined from the sum of two main peak areas detected at retention times of 9.7 min. and 10.2 min., per mg of dry algal cell weight.

Gas chromatography

All solvent extracts were analyzed using a gas chromatograph equipped with a mass spectrometer and a flame ionization detector (GC-MS-FID) with modifications to a previously described protocol (Thapa *et al.*, 2016). Briefly, the GC-MS-FID analyses were performed using an Agilent 7890A gas chromatograph connected to a 5975C inert MSD with a triple-axis detector. The system was equipped with a DB-5MS column (30 m × 0.25 mm i.d., 0.25 μm film thickness) (Agilent J&W, USA). The temperatures were set to: injector (260 °C), interface (250 °C), and ion source (200 °C). One μL of liquid solvent sample was injected in splitless mode with an autosampler (Model G4513A, Agilent). Column flow was kept constant at 2.5 mL min^{-1} with He as a carrier gas. The initial GC oven temperature of 220 °C was held for 1 min, then raised to 280 °C at a rate of 5 °C min^{-1}, followed by 2 °C min^{-1} to 300 °C and held for 5 min. Mass spectra were recorded after a 3 min solvent delay using a scanning range of 50–750 m/z at 20 scans s^{-1}.

Gas chromatograms were manually reviewed for quality control before botryococcene GC peak areas were integrated using MassHunter Workstation software

version B.08.00 (Agilent Technologies, USA). The NIST library (National Institute of

Standards and Technology, Gaithersburg, MD, USA) was used to identify

botryococcenes.

Statistical analysis

Differences between the mean values obtained from each assay were assessed

through one-way analysis of variance (ANOVA) followed by Tukey's HSD post-hoc test

with a confidence level of 95%. All statistical analyses were performed using R software

v3.4.3 (https://www.r-project.org/) through the RStudio interface.

238
239

Results and Discussion

B. braunii growth and hydrocarbon extractability in different media

After 18 days of cultivation, there were no significant differences in specific growth rates of *B. braunii* cultured in modified Chu13 medium (0.077 ± 0.002 day^{-1}; mean $\pm$ SEM) and in 5% RSW (0.064 ± 0.007 day^{-1}), but the algae cultured in 37.5 mM NaCl (0.050 ± 0.005 day^{-1}) showed a significant decrease ($p < 0.01$) in the specific growth rate compared to the other treatments (Fig. 1A). *B. braunii* cultured in 5% RSW and in 37.5 mM NaCl had significant decreases in chlorophyll content (3.6 ± 0.40 mg g^{-1} DCW and 3.4 ± 0.28 mg g^{-1} DCW, respectively) and appeared less healthy (chlorotic phenotype) compared to algae cultured in non-saline medium (5.3 ± 0.13 mg g^{-1} DCW) (Fig. 1B).

The hydrocarbon content in dry biomass of *B. braunii* cultured in the brackish medium was 33 ± 0.5% DCW compared to 40 ± 2.1% DCW in modified Chu13 medium (Fig. 1C). However, the hydrocarbon recovery rate from suspended *B. braunii* cells in 5% RSW (37.3 ± 5.1%) was 2-fold greater than in the suspended algae cultured in 37.5 mM NaCl conditions (18.8 ± 2.4%), while the hydrocarbon extractability in non-saline medium was non-existent (Fig. 1D). The presence of C_{34} and C_{32} botryococcenes from suspended *B. braunii* cells cultured in 5% RSW was confirmed by the appearance of two distinct peaks (9.7 min and 10.2 min, respectively) with appropriate mass spectra in GC-MS chromatograms (Fig. 1E). These results indicate that culturing *B. braunii* in modified Chu-13 medium with 5% RSW is possible without strongly reducing its growth and encourages the production of extracellular botryococcenes. This is in line with previous reports (Furuhashi *et al.*, 2013; 2016; 2022), where *B. braunii* cultivation in brackish medium

reduced the cultures' growth rate but improved hydrocarbon extractability from algal

biomass compared to cultivation in non-saline medium.

Suitability screening of solvents for two-phase cultivation of *B. braunii* race B in

brackish culture medium

A recent study has shown that the Showa strain is one of the most promising for *in*

situ extraction because of its high solvent resistance, hydrocarbon extractability, and lipid

productivity (Kleinert and Griehl 2021). Our results showed that modified Chu-13 medium

with 5% RSW was the best for growing and improving botryococcene recovery rate around

cultured *B. braunii* Showa (Fig. 1). Because *B. braunii* hydrocarbons are mainly (90–95%)

stored in the ECM, most of them can be extracted by thermal pretreatment (i.e. drying or

heating of wet algae) or culturing the algae in brackish medium before soaking the algae

in a non-polar organic solvent such as hexane that does not permeate the cells (Largeau

et al., 1980; Suzuki *et al.*, 2013; Furuhashi *et al.*, 2016b). Hydrocarbons obtained in this

way are typically referred to as extracellular hydrocarbons. After recovering the

extracellular hydrocarbons, a small amount of intracellular hydrocarbons can also be

extracted with a mixture of chloroform and methanol that can reach inside the cells

(Largeau *et al.*, 1980). For the initial solvent screening, cultures grown in 5% RSW were

treated with 14 different non-polar solvents at 25% (v/v), whose physicochemical

properties are shown in Table 1.

After 2 and 24 h extraction, the solvents exhibited a highly variable performance in

extracting botryococcene hydrocarbons from the algal cells (Fig. 2A). This was visually

indicated by the co-extraction of extracellular carotenoids that had previously been

reported to be associated with the botryococcene fraction from the Showa strain (Eroglu

and Melis, 2010). Most of the solvents showed yellowish coloration at different levels in

the upper phase (Fig. 2A). Brominated alkanes and FCs formed a lower phase (Fig. 2A).

Alkane solvents showed a similar but relatively poor hydrocarbon recovery rates (Fig. 2B).

Halogenated solvents were worse than alkanes except for bromodecane, which was the

best of all previously tested solvents (Fig. 2B). However, the hydrocarbon recovery rate

with green solvents (e.g. 34.2 ± 0.5% with α-pinene) was found to be up to ~10-fold and

~3.8-fold greater than with dodecane (3.3 ± 0.1%) and bromodecane (8.9 ± 1.2%),

respectively (Fig. 2B). The exception was nopol which did not extract botryococcenes (Fig.

2B). This indicates that green solvents, such as ɣ-terpinene and α-pinene, were more

efficient at extracting botryococcenes *in situ* compared to conventional alkane solvents

and their halogen derivatives (Fig. 2B).

Although both α-pinene and ɣ-terpinene yielded the highest extraction efficiency of

botryococcenes in this screening (Fig. 2B), we observed a severe degradation of

chlorophyll in all green solvents extracts after 24 h of incubation compared to alkanes and

halogenated solvents (Fig. 2C). The total chlorophyll content in *B. braunii* after two-phase

cultivation with solvents was measured to evaluate solvent biocompatibility. The addition

of solvents to *B. braunii* led to differing extents of chlorosis. Of the different two-phase

cultivations, algal cultures with alkanes showed a slight decrease in the chlorophyll

content compared to cultivations without solvent (Fig. 2C). Algae with brominated alkanes

and chlorododecane demonstrated a stronger reduction in chlorophyll content (1.7–2.1

mg g^{-1} DCW) than those with straight-chain alkanes (3.0–3.6 mg g^{-1} DCW) while cultures

with FCs had chlorophyll contents (3.9–4.2 mg g^{-1} DCW) similar to the no-solvent

treatment (4.4 ± 0.08 mg g^{-1} DCW) (Fig. 2C). Green solvents were the most toxic solvents,

particularly the alcohols, leading to a drastic decrease in chlorophyll content to around 0–

0.4 mg g^{-1} DCW (Fig. 2C).

Our results indicate that green solvents are too toxic for two-phase cultivations,

unlike longer chain alkanes and halogenated solvents. A previous study reported similar

results after comparing the effects of decane, its alcohol derivative decanol, and the green

solvent D-limonene, where decanol was the most toxic solvent, followed by D-limonene

and decane, (Concha *et al.*, 2018). Here, brominated alkanes and FCs formed a lower

phase or 'underlay' with algal cultures, unlike other solvents that formed a distinct layer

on top of the culture (Fig. 2A). Of the heavy solvents, only bromodecane could capture

hydrocarbons from wet algal samples (Fig. 2B). In contrast, previous work reported that

FCs could replace the use of dodecane to extract the sesqui- and diterpenoids produced

by engineered green microalgae (Overmans and Lauersen, 2022).

Selected solvents for *in situ* extraction of hydrocarbons produced by *B. braunii*

race B

We evaluated *in situ* hydrocarbon extraction in larger culture volumes at two

incubation periods, 2 h and 24 h, with six solvents: hexane, dodecane, bromodecane,

bromododecane, p-cymene and ɣ-terpinene. *B. braunii* cultured in 5% RSW medium were

gently overlaid with each solvent at 10% (v/v), which is the standard culture:solvent ratio

used to capture terpenoids or hydrocarbons from engineered algae (Overmans and

Lauersen, 2022; Yunus *et al.*, 2022). Regardless of solvent density, we observed the

formations of emulsions in cultures with all solvents after shaking the mixtures at 120 rpm

even after 2 h (Fig. 3A). This formation was previously identified as one technical limitation

for isoprenoid milking with hydrophobic solvents on algal culture (Lauersen, 2019). After

2 h of extraction, the alkane solvents dodecane and hexane showed a yellow coloration,

indicating hydrocarbon extraction, while the aqueous lower phase contained algae and

culture medium. After 24 h of extraction, algal cultures were only green in dodecane, which

also exhibited a more intense yellow color accumulation than hexane, where cells were

chlorotic or dead (Fig. 3A).

Brominated solvents formed a lower phase to the aqueous algal culture where

colonies clumped, but no hydrocarbon extraction occurred from either solvent. Terpene

green solvents were found to extract the hydrocarbons, with aggregation of colonies

visible at 2 h and severe chlorophyll degradation at 24 h of extraction (Fig. 3A). After a 2

h contact time between solvents and the 5% RSW-acclimated *B. braunii* cells, we detected

the highest hydrocarbon content in hexane, followed by p-cymene and γ-terpinene (Fig.

3B). The hydrocarbon recovery rate in extractions with dodecane and brominated alkanes

were found to be much higher after 24 h than after 2 h (Fig. 3B). Bromodecane showed a

similar hydrocarbon extractability compared to the two alkanes after 24 h of incubation.

Even though the hydrocarbon recovery rate with bromodecane resulted in a 3.8-fold

increase compared to bromododecane (Fig. 3B), both of the tested brominated alkanes

caused a similar decrease in chlorophyll content compared to dodecane from 3.7 ± 0.08

mg g^{-1} DCW to ~2.6 mg g^{-1} DCW (Fig. 3C). *B. braunii* race B cells cultured with hexane

and p-cymene showed a fast decrease in chlorophyll content (3.56 ± 0.16 mg g^{-1} DCW

and 2.62 ± 0.08 mg g^{-1} DCW, respectively), whereas algal cells with γ-terpinene ($4.21 \pm$

0.06 mg g^{-1} DCW) showed a similar chlorophyll content to cells with dodecane after 2 h

of incubation (4.37 ± 0.07 mg g^{-1} DCW) (Fig. 3A and 3C). Longer contact time (24 h) with

γ-terpinene was detrimental to algal cells (0.66 ± 0.06 mg g^{-1} DCW), as could be seen in

the hexane treatment (Fig. 3A and 3C). This result indicates that the bromodecane

underlay was amenable to the extraction of extracellular hydrocarbons and carotenoids

from *B. braunii* cells, but it was more aggressive as a two-phase culture system than the

conventional dodecane overlay.

After 24 h of extraction, *B. braunii* cultured in brackish medium with 10% (v/v) of

solvents showed lower hydrocarbon recovery rates compared to the algae cultured with

25% (v/v) of solvents (Fig. 2B and 3B). Increasing culture salinity to 10% RSW (75 mM

NaCl) could improve the hydrocarbon extraction efficiency from suspended algal cells

without requiring a high solvent content. It was shown that the hydrocarbon recovery rate

from *B. braunii* race B cultured in brackish medium with a salinity of 70 mM NaCl for 18

days was nearly equal to that from the freeze-dried algae (100% hydrocarbon recovery

rate), but the biomass production was reduced (Furuhashi *et al.*, 2016a). Another strategy

could focus on milking hydrocarbons from wet algae in their early growth phase after each

subculture into fresh brackish medium. For example, it was reported that *B. braunii* race

B cultured in brackish media with a salinity of 30 and 50 mM NaCl for 18 days showed a

hydrocarbon recovery rate of around 12% and 36% of dry heptane extraction,

respectively. However, the algae on day 24, which was 6 days after subcultivation in

brackish media with a salinity of 30 and 50 mM NaCl, showed an increase in the

hydrocarbon recovery rate of around 35% and 90% of dry heptane extraction, respectively

(Furuhashi *et al.*, 2016a).

The highest hydrocarbon recovery rates were detected in p-cymene (13.3 ± 0.5%)

and γ-terpinene (13.6 ± 0.8%) extracts after 24 h of incubation, both showing hydrocarbon

extractabilities 13- and 15-fold greater than hexane (1.0 ± 0.4%) and dodecane (0.9 ±

0.1%), respectively (Fig. 3B). As far as we know, this is the first report of using γ-terpinene

in botryococcenes extraction. On the other hand, p-cymene has been used to extract lipids

from different biomass sources, including oleaginous microalgae (De Jesus and Maciel

Filho, 2020). Even though we found a similar hydrocarbon extractability between p-

cymene and ɣ-terpinene when the solvent content was 10% v/v (Fig. 3B), p-cymene was

more aggressive on the cells than ɣ-terpinene (Fig. 3C). It has been previously reported

that p-cymene was more toxic against insects than γ-terpinene, indicating the latter will

likely be more appropriate for future botryococcene milking setups (Gong and Ren, 2020).

The results indicate that ɣ-terpinene could be used as an alternative solvent to

replace hexane for *in situ* extractions of hydrocarbons, but only for short extraction times.

Different bioprocess designs to the *in situ* extraction procedure used in our study could

help to reduce the toxic effects of solvent contact on algal cells. For example, it was

recently shown that *B. braunii* cultures grown in a bubble column bioreactor attached to

an extraction column could achieve a long-term *in situ* extraction of extracellular

hydrocarbons over 30 days by using a daily extraction time of 5 h with hexane (Kleinert

and Griehl 2022). Other *in situ* extraction systems and setups, such as membrane

filtration, have shown to be successful in extracting exopolysaccharides (EPS) from *B.*

braunii CCALA778 by using a microfiltration hollow-fiber membrane without requiring

solvent (García-Cubero *et al.,* 2018). A similar system was used to continuously extract

the heterologous sesquiterpenoid patchoulol with a hydrophobic hollow-fiber membrane

acting as a physical interaction contact matrix between engineered algal cells and the

solvent dodecane (Overmans *et al.*, 2022).

Hexane is a suitable extraction solvent due to its low boiling point (69 °C), low cost,

and high extraction efficiency (Jackson *et al.*, 2017; Ennaceri *et al.*, 2023). Using hexane

facilitates solvent separation from the botryococcenes by distillation, as they have boiling

points well above hexane (> 300 °C; Jackson *et al.*, 2017). Due to its higher volatility,

hexane would be superior to brominated alkanes and the terpene green solvents with

boiling points > 150 °C, but only if the downstream process relies on energy-intensive

distillation (Table 1). Other emerging techniques, including organic solvent nanofiltration

and alternative wetting of membranes, may enable low-energy separations of

botryococcenes, similar to those recently reported for sesquiterpene alcohols (Jackson *et*

al., 2017; Alduraiei *et al.*, 2021; Overmans *et al.*, 2022). Terpene green solvents used in

this work (134–136 g/mol; Table 1) have smaller molecular weights than botryococcenes

(e.g., 466.8 g/mol $C_{34}H_{58}$), which would facilitate their downstream separation by these

techniques. Although p-cymene may be the least expensive alternative solvent when it

compares to hexane for botryococcene extraction (Table 1), solvent recycling of p-cymene

and other green solvents would make it more competitive for the downstream processing

of algae-based biofuels. Furthermore, the improved extraction efficiencies and

sustainable nature of green solvents make them attractive alternatives to hexane.

418
Conclusions

Using brackish cultivation to improve hydrocarbon extractability is a step-change in

milking efficiency potential for *B. braunii* cultures, but the solvent choice was limited to the

use of straight-chain alkanes. Here, we show that terpene green solvents and heavy

solvents like bromodecane could also be used as alternative solvents to hexane for

milking botryococcenes from *B. braunii* race B cultured in brackish medium. However,

they are less biocompatible than dodecane, leading to 33–100% chlorophyll content loss.

Our results warrant follow-up testing of these parameters in short-time scale solvent

contactor systems that may improve the *B. braunii* hydrocarbon milking processes.

Declarations

Acknowledgments

We would like to thank Chandrasekaran Lakshmipathy of the KAUST Lab Equipment Maintenance (LEM) team for assistance in upgrading and initializing the GC-FID-MS unit.

Authors' contributions

TAO-U performed the experiments, contributed to the research planning, experimental design, data collection and visualization, formal analysis, and wrote the original draft manuscript. SO contributed to the research planning, experimental design, GC-MS analysis, and data visualization, and reviewed the manuscript. BBdF contributed to experimental design, assisted in the operation of Algem photobioreactors, and reviewed the manuscript. EL-G contributed to funding acquisition and data visualization, and reviewed the manuscript. The research was mainly conducted in the laboratory of KJL, who was responsible for experimental design, project scope, funding acquisition and reviewing the manuscript. All authors read and approved the final manuscript.

Funding

TAO-U is financially supported by CONAHCYT (Consejo Nacional de Humanidades, Ciencias y Tecnologías) Mexico with a PhD fellowship (CVU 780487) and the KAUST Visiting Student Research Program (VSRP). The research reported in this publication was supported by KAUST baseline funding awarded to KJL.

Availability of data and materials

All data generated or analyzed during this study are included in this article and its

supplementary information file.

Ethics approval and consent to participate

Not applicable.

Consent for publication

All the authors consented to the publication of this work.

Competing Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

References

1. Abdel-Aal, E. A., Farid, M. E., Hassan, F. S., & Mohamed, A. E. (2015). Desalination of Red Sea water using both electrodialysis and reverse osmosis as complementary methods. *Egyptian Journal of Petroleum*, *24*(1), 71-75. https://doi.org/10.1016/j.ejpe.2015.02.007

2. Alder, C. M., Hayler, J. D., Henderson, R. K., Redman, A. M., Shukla, L., Shuster, L. E., & Sneddon, H. F. (2016). Updating and further expanding GSK's solvent sustainability guide. *Green Chemistry*, *18*(13), 3879-3890. https://doi.org/10.1039/C6GC00611F

3. Alduraiei, F., Manchanda, P., Pulido, B., Szekely, G., & Nunes, S. P. (2021). Fluorinated thin-film composite membranes for nonpolar organic solvent nanofiltration. *Separation and Purification Technology*, *279*, 119777. https://doi.org/10.1016/j.seppur.2021.119777

4. Concha, E., Heipieper, H. J., Wick, L. Y., Ciudad, G. A., & Navia, R. (2018). Effects of limonene, n-decane and n-decanol on growth and membrane fatty acid composition of the microalga *Botryococcus braunii*. *AMB Express*, *8*, 189. https://doi.org/10.1186/s13568-018-0718-9

5. De Jesus, S. S., & Maciel Filho, R. (2020). Recent advances in lipid extraction using green solvents. *Renewable and Sustainable Energy Reviews*, *133*, 110289. https://doi.org/10.1016/j.rser.2020.110289

6. Ennaceri, H., Nwoba, E. G., Ogbonna, C. N., Bahri, P. A., & Moheimani, N. R. (2023). Progress of non-destructive hydrocarbon extraction technology of *Botryococcus braunii*. *Algal Research*, *73*, 103156. https://doi.org/10.1016/j.algal.2023.103156

7. Eroglu, E., & Melis, A. (2010). Extracellular terpenoid hydrocarbon extraction and quantitation from the green microalgae *Botryococcus braunii* var. Showa. *Bioresource Technology*, *101*(7), 2359-2366. https://doi.org/10.1016/j.biortech.2009.11.043

8. Frenz, J., Largeau, C., Casadevall, E., Kollerup, F., & Daugulis, A. J. (1989). Hydrocarbon recovery and biocompatibility of solvents for extraction from cultures

of *Botryococcus braunii*. *Biotechnology and bioengineering*, *34*(6), 755-762.
https://doi.org/10.1002/bit.260340605

9. Furuhashi, K., Saga, K., Okada, S., & Imou, K. (2013). Seawater-cultured
Botryococcus braunii for efficient hydrocarbon extraction. *PLoS One*, *8*(6), e66483.
https://doi.org/10.1371/journal.pone.0066483

10. Furuhashi K, Hasegawa F, Saga K, Okada S, Kaizu Y, & Imou K. (2016a). Effects
of culture medium salinity on the hydrocarbon extractability, growth, and
morphology of *Botryococcus braunii*. *Biomass and Bioenergy, 91*, 83–90.
https://doi.org/10.1016/j.biombioe.2016.05.007

11. Furuhashi, K., Noguchi, T., Okada, S., Hasegawa, F., Kaizu, Y., & Imou, K.
(2016b). The surface structure of *Botryococcus braunii* colony prevents the entry
of extraction solvents into the colony interior. *Algal Research*, 16, 160-166.
https://doi.org/10.1016/j.algal.2016.02.021

12. Furuhashi, K., Magota, A., Liu, Y., Hasegawa, F., Okada, S., Kaizu, Y., & Imou, K.
(2022). Thinning of *Botryococcus braunii* Colony Sheath by Pretreatment
Enhances Solvent-Based Hydrocarbon Recovery. *Phycology*, *2*(4), 363-373.
https://doi.org/10.3390/phycology2040020

13. García-Cubero, R., Wang, W., Martín, J., Bermejo, E., Sijtsma, L., Togtema, A., ...
& Kleinegris, D. M. (2018). Milking exopolysaccharides from *Botryococcus braunii*
CCALA778 by membrane filtration. *Algal Research*, *34*, 175-181.
https://doi.org/10.1016/j.algal.2018.07.018

14. Gong, X., & Ren, Y. (2020). Larvicidal and ovicidal activity of carvacrol, p-cymene,
and γ-terpinene from *Origanum vulgare* essential oil against the cotton bollworm,
Helicoverpa armigera (Hübner). *Environmental Science and Pollution*
Research, *27*, 18708-18716. https://doi.org/10.1007/s11356-020-08391-2

15. Jackson, B. A., Bahri, P. A., & Moheimani, N. R. (2017). Repetitive non-destructive
milking of hydrocarbons from *Botryococcus braunii*. *Renewable and Sustainable*
Energy Reviews, *79*, 1229-1240. https://doi.org/10.1016/j.rser.2017.05.130

16. Jin, J., Dupré, C., Yoneda, K., Watanabe, M. M., Legrand, J., & Grizeau, D. (2016).
Characteristics of extracellular hydrocarbon-rich microalga *Botryococcus braunii*

for biofuels production: Recent advances and opportunities. *Process Biochemistry*, *51*(11), 1866-1875. https://doi.org/10.1016/j.procbio.2015.11.026

17. Kanda, H., Li, P., Yoshimura, T., & Okada, S. (2013). Wet extraction of hydrocarbons from *Botryococcus braunii* by dimethyl ether as compared with dry extraction by hexane. *Fuel*, *105*, 535-539. https://doi.org/10.1016/j.fuel.2012.08.032

18. Kleinert, C., & Griehl, C. (2021). Identification of suitable Botryococcus braunii strains for non-destructive in situ hydrocarbon extraction. *Journal of Applied Phycology*, *33*(2), 785-798. https://doi.org/10.1007/s10811-020-02342-7

19. Kleinert, C., & Griehl, C. (2022). In situ extraction (milking) of the two promising *Botryococcus braunii* strains Showa and Bot22 under optimized extraction time. *Journal of Applied Phycology, 34*(1), 269-283. https://doi.org/10.1007/s10811-021-02633-7

20. Kumar, S. J., Kumar, G. V., Dash, A., Scholz, P., & Banerjee, R. (2017). Sustainable green solvents and techniques for lipid extraction from microalgae: A review. *Algal Research*, *21*, 138-147. https://doi.org/10.1016/j.algal.2016.11.014

21. Lauersen, K. J. (2019). Eukaryotic microalgae as hosts for light-driven heterologous isoprenoid production. *Planta*, *249*, 155-180. https://doi.org/10.1007/s00425-018-3048-x

22. Largeau, C., Casadevall, E., Berkaloff, C., & Dhamelincourt, P. (1980). Sites of accumulation and composition of hydrocarbons in *Botryococcus braunii*. *Phytochemistry*, *19*(6), 1043-1051. https://doi.org/10.1016/0031-9422(80)83054-8

23. Lozoya-Gloria, E., Morales-de la Cruz, X. and Ozawa-Uyeda, T. A. (2019). "The Colonial Microalgae *Botryococcus braunii* as Biorefinery". Ch. 7. In: *Microalgae - From Physiology to Application.* Vítová M. (Ed.) IntechOpen, London, UK. Pp. 123-144. https://doi.org/10.5772/intechopen.88206

24. Mehta, P., Jackson, B. A., Nwoba, E. G., Vadiveloo, A., Bahri, P. A., Mathur, A. S., & Moheimani, N. R. (2019). Continuous non-destructive hydrocarbon extraction from *Botryococcus braunii* BOT-22. *Algal Research*, *41*, 101537. https://doi.org/10.1016/j.algal.2019.101537

25. Nonomura, A. M. (1988). *Botryococcus braunii* var. Showa (Chlorophyceae) from

Berkeley, California, United States of America. *Japanese Journal of Phycology*, *36*,

285-291. https://patents.google.com/patent/USPP6169P/en

26. Overmans, S., Ignacz, G., Beke, A. K., Xu, J., Saikaly, P. E., Szekely, G., &

Lauersen, K. J. (2022). Continuous extraction and concentration of secreted

metabolites from engineered microbes using membrane technology. *Green*

Chemistry, *24*(14), 5479-5489. https://doi.org/10.1039/D2GC00938B

27. Overmans, S., & Lauersen, K. J. (2022). Biocompatible fluorocarbon liquid

underlays for *in situ* extraction of isoprenoids from microbial cultures. *RSC*

Advances. *12*(26), 16632-16639. https://doi.org/10.1039/D2RA01112C

28. Samorì, C., Torri, C., Samorì, G., Fabbri, D., Galletti, P., Guerrini, F., ... & Tagliavini,

E. (2010). Extraction of hydrocarbons from microalga *Botryococcus braunii* with

switchable solvents. *Bioresource Technology*, *101*(9), 3274-3279.

https://doi.org/10.1016/j.biortech.2009.12.068

29. Suzuki, R., Ito, N., Uno, Y., Nishii, I., Kagiwada, S., Okada, S., & Noguchi, T.

(2013). Transformation of lipid bodies related to hydrocarbon accumulation in a

green alga, *Botryococcus braunii* (Race B). *PLoS One*, *8*(12), e81626.

https://doi.org/10.1371/journal.pone.0081626

30. Thapa, H. R., Naik, M. T., Okada, S., Takada, K., Molnár, I., Xu, Y., & Devarenne,

T. P. (2016). A squalene synthase-like enzyme initiates production of

tetraterpenoid hydrocarbons in *Botryococcus braunii* Race L. *Nature*

Communications, *7*, 11198. https://doi.org/10.1038/ncomms11198

31. Weiss, T. L., Roth, R., Goodson, C., Vitha, S., Black, I., Azadi, P., ... &

Goodenough, U. (2012). Colony organization in the green alga *Botryococcus*

braunii (Race B) is specified by a complex extracellular matrix. *Eukaryotic*

Cell, *11*(12), 1424-1440. https://doi.org/10.1128/ec.00184-12

32. Wellburn, R. W. (1994). The spectral determination of chlorophylls a and b, as well

as total carotenoids, using various solvents with spectrophotometers of different

resolution. *Journal of Plant Physiology*, *144*(3), 307-313.

https://doi.org/10.1016/S0176-1617(11)81192-2

33. Yunus, I. S., Anfelt, J., Sporre, E., Miao, R., Hudson, E. P., & Jones, P. R. (2022).
Synthetic metabolic pathways for conversion of CO_2 into secreted short-to medium-
chain hydrocarbons using cyanobacteria. *Metabolic engineering*, *72*, 14-23.
https://doi.org/10.1016/j.ymben.2022.01.017

Figure captions

Figure 1. Analysis of *B. braunii* (Showa strain) cultured for 18 days in three different

media. (**A**) Specific growth rate, (**B**) total chlorophyll content, (**C**) extracellular hydrocarbon

content, and (**D**) hydrocarbon recovery rate of *B. braunii* race B cells grown in modified

Chu-13 medium (Chu13), modified Chu-13 medium with 37.5 mM NaCl (37.5 mM NaCl),

and modified Chu-13 medium with 5% Red Sea water (5% RSW). Recovered

hydrocarbons from dry algae by hexane extraction accounted for a 100% hydrocarbon

recovery rate. (**E**) Representative GC-FID chromatograms of botryococcenes extracted

with commonly used alkane solvents obtained from wet *B. braunii* algal samples. Error

bars represent the standard error of the mean of three biological replicates (SEM; n = 3).

Statistical significance was determined by one-way ANOVA with Tukey's HSD post-hoc

test ($p < 0.05$). Within each panel, different letters above bars indicate instances where

means were significantly different ($p < 0.05$).

Figure 2. Extraction and toxicity properties of all different solvents. (**A**) *B. braunii* race B

cells grown in culture medium with 5% RSW after 2 h and 24 h extraction with 25% (v/v)

of solvents. (**B**) Hydrocarbon recovery rates after 24 h of two-phase living *B. braunii*

extraction cultures with different solvents. Recovered hydrocarbons from dry algae by

hexane extraction accounted for a 100% hydrocarbon recovery rate.(**C**) Total chlorophyll

content in *B. braunii* cultured in 5% RSW medium after 24 h extraction with each solvent.

Error bars represent the standard error of the mean of two biological replicates (SEM; n =

2). Statistical significance was determined by one-way ANOVA with post-hoc Tukey's

HSD test. Letters above the bars indicate significantly different mean concentrations ($p <$

0.05).

Figure 3. Extraction and toxicity properties of selected solvents. (**A**) *B. braunii* cells grown in culture medium with 5% RSW after a contact time of 2 h and 24 h with 10% (v/v) of either (1) hexane, (2) dodecane, (3) bromododecane, (4) bromodecane, (5) p-cymene, or (6) ɣ-terpinene. (**B**) Hydrocarbon recovery rates after 2 h and 24 h of two-phase living *B. braunii* extraction cultures with solvents. Recovered hydrocarbons from dry algae by hexane extraction accounted for a 100% hydrocarbon recovery rate. (**C**) Total chlorophyll content of *B. braunii* cultured in medium with 5% RSW after 2 h and 24 h extraction time with solvents. Error bars represent the standard error of the mean of three biological replicates (SEM; n = 3). Statistical significance was determined by one-way ANOVA with post-hoc Tukey's HSD test. Lower- and upper-case letters above the bars indicate the post hoc results of the 2 h and 24 h data sets, respectively. Means were considered significantly different at level $p < 0.05$.

630   **Figure 1:**

Figure 2:

636 **Figure 3:**

639 **Tables**

640 **Table 1**. Physico-chemical properties of the solvents used in this work.

	Solvent	Molecular formula	Molecular weight (g/mol)	Density (g/mL)	Boiling point (ºC)	Vapor pressure at 25 ºC (mm Hg)	Water solubility at 20 ºC (mg/L)	Cost ($ USD)
Alkane	Decane	$C_{10}H_{22}$	142	0.73	174	1.43	0.052	178 L^{-1}
solvents	Dodecane	$C_{12}H_{26}$	170	0.75	216	0.13	0.0037	389 L^{-1}
	Pentadecane	$C_{15}H_{32}$	212	0.77	271	0.005	0.00004	568 kg^{-1}
Halogenated	Bromodecane	$C_{10}H_{21}Br$	221	1.07	241	0.04	Insoluble	129 kg^{-1}
solvents	Bromododecane	$C_{12}H_{25}Br$	249	1.04	276	0	Insoluble	146 kg^{-1}
	Chlorododecane	$C_{12}H_{25}Cl$	205	0.73	< 1000	0	0.052	117 L^{-1}
	FC-770	C_4F_9NO	399	1.79	95	49.3	1.3	375 kg^{-1}
	FC-3283	$C_9F_{21}N$	521	1.82	128	10.8	< 5.0	378 kg^{-1}
Green	α-Pinene	$C_{10}H_{16}$	136	0.86	156	4.5	2.49	403 L^{-1}
solvernts	ɣ-Terpinene	$C_{10}H_{16}$	136	0.85	183	1.1	8.72	318 L^{-1}
	p-Cymene	$C_{10}H_{14}$	134	0.86	177	1.5	23.4	86 L^{-1}
	Heptanol	$C_7H_{16}O$	116	0.82	176	0.22	1670	129 L^{-1}
	Nerol	$C_{10}H_{18}O$	154	0.88	225	0.03	1311	111 kg^{-1}
	Nopol	$C_{13}H_{18}O$	166	0.97	230–240	0.008	333.5	170 kg^{-1}
Control	Hexane	C_6H_{14}	86	0.66	69	153	0.01	84 L^{-1}

641 Hexane is the standard botryococcene extraction solvent.

642　**Graphical Abstract**